

Response to reviewers for manuscript acp-2016-738: **60 years of UK visibility measurements: impact of meteorology and atmospheric pollutants on visibility**

We thank the reviewers for their time and excellent insights which have helped us to improve the manuscript. We now thank them in the acknowledgements.

We respond to all of the reviewers' points below. Responses are given in red.

**Anonymous Referee #1**

This is an interesting study that measures 60 years of UK visibility in different environments (e.g. urban, rural, and marine) and shows the impact of meteorology and atmospheric pollutants on visibility. The authors use horizontal visibility data along with meteorological data from British Atmospheric Data Centre (BADC) to analyse UK visibility trends from 1950-2013. Although the authors extend the work of Doyle and Dorling (2002) to analyse UK visibility trends, but the reviewer find the dissimilarities of visibility values between Singh et al. (2016) and Doyle and Dorling (2002) results for the period of 1950 to 1997. The authors should explicitly describe why the visibility values presented in their study is different than the results from similar study by Doyle and Dorling (2002). In addition, specific descriptions on the explanations and discussions on atmospheric sciences (e.g. reason for the reduction of air pollutants in urban areas) are insufficient. The author also develop a light extinction model for generating predictions of historic aerosol and gas scattering and absorbing properties. But the authors should provide more detailed discussions on the uncertainties which can arise in their modelling study. More specific comments are provided under 'Technical comments'. The manuscript is reasonably well written, but there are a lot of typographical errors throughout, which are noted under 'Editorial comments'. In my opinion, the manuscript is worth publishing, but some reviewer's concerns existed, which need to be addressed, then this should end up being a paper suitable for publication in ACP.

**Technical comments**

1. Generally the reduction of visibility is found with increasing aerosol particles concentrations which has been explained briefly in introduction part of the paper. The results in this paper shows that the visibility of the urban areas has been improved year to year due to the reduction in air pollution for most of the monitoring stations in UK. But very little has been discussed about the possible reasons of the reduction in air pollution. Are they for cleaner fuel usage in the vehicles? Is it for increased deposition at the building surface due to the urbanisation and human population? Or do the authors have any other suggestion? In marine and rural environments, the natural emissions of aerosol precursors (e.g. DMS from ocean and terpenes from rural plants) are dominating which cannot be

controlled. Do the authors think that this could be the reason for decreased visibility over time for marine and rural stations?

Thanks for this useful comment. The primary aim of this paper is not a detailed source of apportionment study; therefore we cannot be certain what caused the visibility changes at the different sites. Previous work in the literature has shown that changes in fuel use that came about after the clean Air Act are likely reasons. We now provide more information on the likely causes determining the visibility on Page 10 Line 6 to Page 10 Line 12 “Improved visibility at most of the sites is due to reduction in air pollution and the likely changes in fuel use and consumption that took place after 1956 Clean Air Act. The Clean Air Act was introduced with the aims of reducing smog, smoke and sulphur dioxide concentrations in the environment. In particular, the policy focused on industrial emission sources and its reduction (Williams, 2004). Recently, Harrison et al. (2015) shown that concentration of sulphur dioxide, coal smoke, nitrogen dioxide, suspended matter (black smoke) and PM were significantly reduced in the UK over last five decades as the result of switching to cleaner fuels after 1956 Clean Air Act.”

**2.** After comparing the visibility results of this paper with Doyle and Dorling (2002) results, the reviewer found the similar variation trend for the period of 1950-1995, but the visibility values are found to be lower for all stations in this study than Doyle and Dorling (2002) study. Why does this paper produce lower visibility values? No explanation/comparison has been shown in the paper.

We agree that there is similar variation trend for the period of 1950-1997, with our work showing slightly lower visibility values compared to Doyle and Dorling (2002). Doyle and Dorling (2002) filtered data for 12 noon, relative humidity <90% and Present weather codes (PR code) of 00-05 in their statistical analysis for the period of 1950-1997. In our study we used mean averaging for statistical analysis, where we filtered data for 12 noon and relative humidity <99%. As discussed already in the manuscript PR codes were not available after 1997 and hence we could not use them (Page 4 Line 16 to Page 4 Line 21):

“Unfortunately the use of present weather codes largely ceased with the introduction of automated meteorological stations and insufficient PR codes were available after the year 1997. It is noted, that if the present weather codes were available they would have been useful to screen the data for rain or other precipitation events. Due to unavailability of present weather codes during required study period (1950-2012), data filtration was done on the bases of RH limits instead of PR codes. Data were removed when the relative humidity reading was > 99 % which is highly suggestive of rain or other precipitation events.”

The differences in our filtering approach, compared to that used in Doyle and Dorling, is the reason why we have lower visibility values compared to Doyle and Dorling (2002). More explanation and comparison are now added in Page 9 Line16 to Page 9 Line 23 “A similar variation in visibility trends is observed for the period of 1950-1997, comparing with Doyle and Dorling (2002). However, this study reports overall lower visibility values when compared to Doyle and Dorling (2002). These differences are due to slightly different data filtering methodologies. Doyle and Dorling (2002) filtered data for 12 noon, relative humidity > 90% and PR codes of 00-05 in their statistical analysis for the period of 1950-1997. However, due to uncertainty and unavailability of PR code after 1997 we did not use these codes. Furthermore we performed mean averaging for statistical analysis, where data is filtered for 12 noon and relative humidity > 99 %. The details of uncertainty and unavailability of PR codes and used data filtration method are given in data and methodology sections”.

3. 12 noon data has been taken as the daily data, however there could be the variation of the visibility throughout the day because of the variation of the meteorological parameters and the concentration of aerosol particles. These need to be discussed as uncertainties of their results. The authors excluded the data for 99% precipitation. How much percent data points of 99% precipitation? How the whole data analysis has been affected after excluding these data points?

Although analysis could be performed for any hour of the day, we chose 12 noon because as stated on Page 4 Line 22 manual observations of visibility can be affected by low light levels. Therefore, we chose a time when light levels were near their maximum.

As shown in Table R1 below the number of data points removed due to the filtering of data with RH >99% is very low. The filtered data accounts for 0.91 – 3.44 % of the total data dependent upon site location. Therefore removing these points does not make any significant difference. We now make the point in the manuscript on Page 4 Line 21 “Removal of data with RH > 99% removes between 0.91 – 3.44 % of the data dependent on site location”.

Table R1

<b>Station</b>	<b>Total data points</b>	<b>Data points of above 90% precipitation</b>	<b>Data points of above 99% precipitation</b>	<b>% (data points of above 90% precipitation)</b>	<b>% (data points of above 99% precipitation)</b>
Aldergrove	23370	4491	619	19.25 %	2.65 %
Heathrow	23322	2460	292	10.53 %	1.25 %
Leuchars	20814	2839	190	13.63 %	0.91 %

Nottingham	20351	3749	620	18.42 %	3.04 %
Plymouth	23183	4719	798	20.35 %	3.44 %
Ringway	20027	2140	195	10.68 %	0.97 %
Tiree	20412	4601	472	22.54 %	2.31 %
Waddington	23163	4014	710	17.33 %	3.07 %

4. Page 1: The reviewer think that some of the sentences (e.g. Moreover, degradation in visibility can be hazardous to . . . . . sea and air accidents, The site locations include . . . . . marine environments, the model incorporates parameterizations . . . . . and particle and gas adsorption) in Abstract are not meaningful. Instead, they can only be placed in introduction and methodology part.

We think a brief description of visibility and the model strengthens the abstract by providing context for the paper. No changes have been made.

5. Page 2 Line 3: The literature review (WMO, 2003) is very old. The authors should consider updating their literature review by using recent report by WMO (2015). There are some other places in the introduction part where the references can be updated. Reference: World Meteorological Organization, 2015: Manual on the Global Observing System. Volume I – Global aspects, (WMO-No. 544), Geneva.

Suggested changes have been implemented in the manuscript.

6. Page 4, Line 32: The reader might be confused in many places of the manuscript as authors used ‘human observation’ and ‘manual observation’ for same meaning. As human observation is more common term for visibility measurement, the reviewer suggest the authors to change ‘manual observation’ to ‘human observation’ throughout the manuscript.

Suggested changes have been implemented in the manuscript; where “manual” has changed to “human” throughout manuscript.

7. Page 5, Line 20-24: The authors claimed that at high visibility the automatic sensors perform sub-optimally at coastal site (e.g. Tiree) due to accumulation of sea salt residue. If this is the case, it will

also be applicable for another coastal site, Leuchars. But the station Leuchars did not show any deviation when the measurement moved from manual to automation. How the authors will explain the different measurement behaviour for similar type of stations, Tiree and Leuchars?

We agree that station Leuchars did not appear to show any deviation when the measurement moved from manual to automation. We suggest two possible reasons, firstly as shown in Figure 1, the average wind speed of Tiree is higher than Leuchars, which causes more sea salt to be generated and transported to the site. Secondly, as mentioned in the main manuscript Page 12 Line 28 that Tiree Island has a very flat landscape, which is not sheltered from the wind in any direction, thereby potentially allowing salt to accumulate more readily on the sensor.

**8.** Page 9, Line 4: What localized sources close to visiometer at the Plymouth site? Are they aerosol particles?

The following sentence has been added in Page 9 Line 10 about localized source at Plymouth site “(ship and traffic emissions from nearby ports and roads)”.

**10.** Page 10, line 9-13: The sentence is contradictory to the reviewer. The reviewer can see from the decadal polar plot that the visibility has been improved decade to decade when the wind comes from the south to east direction, but the reviewer doesn’t understand how this is connected with the part “the higher wind speeds from the direction of Belfast leads to lower visibility over Aldregrove”. Overall the lower visibility in the south to east direction can suit with the above statement.

We have revised the sentence, which now reads “It is clearly seen that visibility has improved the most when wind comes from the south to east direction which covers mainland urban areas such as Belfast, the major regional city.” (Page 10 Line 30)

**11.** Page 12, Lines 10-15: Is there any specific reasons for higher visibilities on Friday for Leuchars and on Wednesday for Ringway?

We checked the data on those two days to investigate any extreme high visibility values, but found no specific reasons. We note that the visibility changes on these days are slight compared to the weekend effect. The main focus of this graph is to show the difference between weekday and weekend visibility. This graph has now been replaced with normalized weekday visibility graph in

response to one of Reviewer 2's questions, where mean weekday visibility normalized to Sunday mean values provides a direct estimate of the percentage differences in weekday visibility values.

**12.** Page 13, Line 5: It would strength the manuscript if the authors can show the relationship of the visibility with temperature. From Fig 5, the reviewer can't see any relationship between visibility and air temperature.

Most sites show clear anti-correlation between temperature and relative humidity as is expected under UK meteorological conditions. Hence a correlation also exists between temperature and visibility. We have now added figure S5 for correlation statistics. The following sentence has been added in Page 13 Line 29 "The correlation statistics between visibility, relative humidity, air temperature and wind speed are provided for all stations in supplementary Fig. S5."

**13.** Figure 2: Measurement methodology Supplementary Table shows that the measurement was performed manually at Ringway from 2004 to onwards, but in the figure-2, there are no data points from 2004. And also the shading should be white as they are manual measurements. And for Heathrow, the red shading and blue shading are the measurements using the same instrument?

We agree, we have mentioned in Supplementary Table S1 that visibility measurement was performed manually at Ringway from 2004 onward; however, very limited numbers of observations at 12 noon are available for statistical analysis and hence we have not able to show annual and seasonal variability after 2004.

There was a mistake in the shading; it has now been replaced. For Heathrow red shading and blue shading are the measurements using the different instrument. More details are now added in Figure 2 caption; Page 24 "Shading indicates changes in measurement methodology, where white is human observation, while blue and red are automated observation using different instruments."

**14.** Figure 5: The reviewer doesn't think this figure is necessary, as most of them already shown in previous figures. Instead, this figure can be placed in the supplementary Information. However, the rose plot for annual average (for full data series) can be placed in the main manuscript which will be easier for reader to see the overall influence of wind speed and wind direction on visibility. The decadal seasonal polar plots can be kept in Supplementary Information.

The suggested change has been implemented, where Figure 5 has been placed in the supplementary Information as Figure S4. Figure 1 shows the wind rose plot for annual average (for full data series) and we think should remain in the main manuscript.

**15.** Figure 7: The green shaded region has been shown from 1990s, but most of the stations start visiometer measurement from 2000. Will it be 2000s instead of 1990s?

The suggested change has been implemented in Figure 7, where green shade has been changed from 1990s to 2000s.

### **Editorial Comments**

Page 1 Line 16: examples of urban areas are preferable.

Suggested change has been implemented in the manuscript

Page 2, Line 23: 'sharp changes' can be replaced by 'sharp decreases'

Suggested change has been implemented in the manuscript

Page 2, Line 25: 'describe' need to be replaced by 'described'

Suggested change has been implemented in the manuscript

Page 3, Line 23: 'decline' should be 'been declined'

Suggested change has been implemented in the manuscript

Page 4, Line 4: 'to' needs to be added in between 'help' and 'explain'

Suggested change has been implemented in the manuscript

Page 4, Line 5: 'They' should be replaced by 'We'.

Thanks for pointing out this. This sentence was unclear. We now state the following “A new model is also presented which can aid in future visibility prediction under different climate and pollution scenarios.”

Page 4, Line 29: Is the term ‘human observation’ or ‘human observer’?

Suggested change has been implemented, where human observation has changed with human observer

Page 5, Line 10, please delete ‘than’

Suggested change has been implemented in the manuscript

Page 5, line 25, please add ‘the’ after assess

Suggested changes have been implemented in the manuscript

## **Anonymous Referee #2**

General comments The study explores UK horizontal visibility, using observations from a number of stations of different characteristics. Actually the study extends the work by Doyle & Dorling (2002) who reported UK visibility improvement from 1950-1997 due to antipollution measures. The extension alone is not so useful as regards estimation of long term trends, since a very strong step change occurred after changes in observational methods. However, authors perform detailed analyses regarding meteorological influence, the role of RH, and develop a light extinction model which make the study interesting. Many points however need to be reconsidered, corrected and clarified.

## **Major Comments**

The study updates the work by Doyle & Dorling 2002, who study UK visibility from 1950-1997. The same stations and the same visibility hour (12Z) have been used in both studies. So, one would expect to see exactly the same values of visibility for the common period which is not true. In contrast, the authors estimate lower values almost in all stations. Is it because of different filters? Or averaging procedure? Following Sloane (1982), Doyle & Dorling exclude visibility values when RH >90%. The authors use another filter (99%) which means that they use more high RH days. I can suppose that this is the reason for the observed differences in the two studies. Some clarifications are required however. How was determined the filter 99%?



We agree that there is similar variation trend for the period of 1950-1997, with our work showing slightly lower visibility values compared to Doyle and Dorling (2002). Doyle and Dorling (2002) filtered data for 12 noon, relative humidity <90% and Present weather codes (PR code) of 00-05 in their statistical analysis for the period of 1950-1997. In our study we used mean averaging for statistical analysis, where we filtered data for 12 noon and relative humidity <99%. As we mentioned in manuscript PR codes were not available after 1997 and hence we could not use them (Page 4 Line 16 to Page 4 Line 21):

“Unfortunately the use of present weather codes largely ceased with the introduction of automated meteorological stations and insufficient PR codes were available after the year 1997. It is noted, that if the present weather codes were available they would have been useful to screen the data for rain or other precipitation events. Due to unavailability of present weather codes during required study period (1950-2012), data filtration was done on the bases of RH limits instead of PR codes. Data were removed when the relative humidity reading was > 99 % which is highly suggestive of rain or other precipitation events.”

The differences in our filtering, compared to that used in Doyle and Dorling, is the reason why we have little lower visibility values from Doyle and Dorling (2002). More explanation and comparison are now added in Page 9 Line16 to Page 9 Line 23 “A similar variation in visibility trends is observed for the period of 1950-1997, comparing with Doyle and Dorling (2002). However, this study reports overall lower visibility values when compared to Doyle and Dorling (2002). These differences are due to slightly different data filtering methodologies. Doyle and Dorling (2002) filtered data for 12 noon, relative humidity > 90% and PR codes of 00-05 in their statistical analysis for the period of 1950-1997. However, due to uncertainty and unavailability of PR code after 1997 we did not use these codes. Furthermore we performed mean averaging for statistical analysis, where data is filtered for 12 noon and relative humidity > 99 %. The details of uncertainty and unavailability of PR codes and used data filtration method are given in data and methodology sections”.

The choice of RH filter (RH<99%) was chosen because this paper was interested in deriving aerosol parameters from the visibility data. In particular, to be able to understand changes in particle hygroscopicity, via the gamma ( $\gamma$ ) parameter, the analysis needs a wide range of RH to fit the model as successfully as possible. If we exclude visibility values when RH >90%, we are losing likely 10-22 % data points (please see Table R1 in Reviewer 1 replied comments). This is the reason we excluded visibility values when RH >99%.

The authors relate visibility with meteorology, however, precipitation is a fundamental parameter which is missing from this analysis. Precipitation increases RH, but also is related to scavenging of

particles in the atmosphere, possibly improving visibility. Precipitation frequency than amount is more important indicator and consequent cleanup of the atmosphere is more important in these cases. So I am wondering if any relevant data are available from nearby stations.

We agree with the referee it would be nice to investigate the role of precipitation upon visibility. Unfortunately in all cases collocated precipitation data are not available for the required study period; therefore, precipitation could not be a focus of this manuscript.

Averaging procedure of visibility is not mentioned. Which code/protocol has been used for human visibility observations? Since uncertainties are much higher in high visibility ranges (as you also mention in Page 5, line 5), visibility follows a rather geometric distribution. Did you use simple mean or a geometric mean for visibility?

The method of measuring visibility changed from human observations to electronic visimeters. This was done at different times for each station. The impact of this change is dramatic as easily seen in Fig. 2. I would say that it is impossible under these circumstances to draw a conclusion for the long term trend of visibility. Are the two methods compared at any station? Is there any parallel period with human + electronic observations? This is a common procedure to evaluate and compare the two methods. If such parallel measurements are available, then authors need to make proper comparisons/calibration and provide a better transition from the first to the second period.

Simple mean method is used for the visibility averaging in the analysis, now has mentioned in Page 5 Line 36. We now also produce boxplots showing the median, interquartile range, outliers etc... in the supplementary material. We note that the median is often close in value to the mean average and the trends remain the same. The following sentences have been added in Page 10 Line 3 to Line 5 “The improvement in median visibility at most of the sites can be seen in supplementary Fig. S1. Boxplots of the decadal visibility are also produced showing the median, interquartile range, outliers etc. (see supplementary Figure S2).”

The details of visibility observations are provided within the UK Met Office guidelines ([https://badc.nerc.ac.uk/data/ukmo-midas/ukmo\\_guide.html](https://badc.nerc.ac.uk/data/ukmo-midas/ukmo_guide.html)).

“Visibility is defined as the greatest distance at which an object in daylight can be seen and recognised, or at night could be seen and recognised if the illumination was raised to daylight levels. Observations should be made at ground level not from observation towers or roof tops. The long standing method of observation has been estimation by the observer using known fixed reference points, such as trees or buildings, which stand out well against the background. Each reference point

should subtend an angle of at least  $0.5^\circ$  at the eye. Estimation of visibility at night is prone to greatest error and should, ideally, be performed with the aid of suitable fixed lights. Visibility estimates on airfields, where accuracy is of particular importance, are often aided in this way. On occasions when the visibility varies in different directions, the minimum value should be reported in the main part of the message and this is the value stored in MIDAS. The guidance to observers at coastal stations states that only visibility over land should be reported; any differing values over the sea being noted in the remarks column of the weather register though it is not clear how closely this practice is followed at voluntary stations.”

For more detail in manuscript, the following line has been added in the manuscript Page 4 Line 29 to Page 4 Line 31 as “More details of visibility observations method are found in UK Met Office Surface Data Users guidelines ([https://badc.nerc.ac.uk/data/ukmo-midas/ukmo\\_guide.html](https://badc.nerc.ac.uk/data/ukmo-midas/ukmo_guide.html)).”

Unfortunately, there are no periods when both human and electronic observations were taken in parallel, therefore we cannot perform a comparison between the two measurement types. To support the conclusion of long term trend in visibility, we have included an additional Figures S1 and S2 in the supplementary material, which show density distribution and decadal box plots of visibility at each station.

The authors use wind roses from surface wind data and perform an extended analysis on visibility variation with respect to wind speed/direction. This is related to air mass origin and associated air pollution or RH sources. Although they perform a reasonable analysis, I think that additional information is required regarding local or long transport pollution from distant sources. Frequently, surface winds reflect very local phenomena (breezes, circulations due to UHI effect, channelling phenomena etc). Although authors refer to long range transport of air pollution from central Europe (for eastern sector) there is no information on long range transport (trajectories, frequencies etc). European emissions increased after the 1950s and decreased after the 1980s. Is UK unaffected from these changes? Is it all local pollution? A discussion on this is necessary. In general, information on local pollution sources and reasons for improving per sector is not adequate. Relative to this, in page 14, line 9, the authors seem to speculate.

Similar to our response to reviewer 1, we note that this paper does not attempt to give a detailed source apportionment of pollution in the UK. An analysis of very local phenomena and long range transport trajectories and frequencies over a multi decade time period represents another major piece of work and is beyond the scope of this study.

Whilst we have not performed trajectory analysis, we have noted previous studies have shown synoptic scale pollution events from Europe affecting UK pollution levels. This information is given in Page 12 Line 19 to 23. We have also added a more recent reference in this section (Crilley et al., 2015) which provides a similar conclusion.

How do you define good or poor visibility? In Fig 2 authors present long-term trends of the annual/seasonal visibility averages and find an overall positive trend in most stations. However, this cannot provide information on the relative improvement in different visibility ranges. Is the improvement higher in low, average or higher visibilities? I would like to see a frequency distribution of different visibility ranges for different sub-periods, which would be much more informative on visibility improvement.

We have now included density distribution graph of decadal visibility for each site in the supplement (Figure S1), where we can clearly see the improvement in median visibility. We have also included boxplot in the supplement to show the decadal changes in visibility over last six decades (Figure S2). The following sentences have been added in Page 10 Line 3 to Line 5 “The improvement in median visibility at most of the sites can be seen in supplementary Fig. S1. Boxplots of the decadal visibility are also produced showing the median, interquartile range, outliers etc. (see supplementary Figure S2).” We have added a definition of poor visibility “(< 2.0 km, (Founda et al., 2016))” in Page 2 Line 5.

In Fig. 5 the authors provide long term records of annual visibility and annual averages of different meteorological parameters. A comparison is attempted between variation of visibility and meteorological variables. I have some questions here. Annual visibility was calculated using daily measurements at 12Z. How other variables were averaged? Do averages refer to 24-hour periods? From the figure it comes out (visually) that visibility is anticorrelated (in low frequencies) with RH. However, RH changes do not refer to 12Z (I think) and also these changes are small enough (in the range of very few units of %, for instance from 75% to 78% or something like that). What mean annual WD refers to? Is it prevailing wind direction? How was calculated? In the same figure, wind speed variability does not seem to be positively correlated (as expected) with visibility. Decreasing trends of wind speed in some stations are accompanied with increasing trends in visibility. Does it mean that wind speed is less influential? Perhaps a running correlation coefficient between visibility and other meteorological variables would be more informative on the influence of such variables and possible temporal changes of this influence. The relationship with air temperature is tentative. At

urban area in particular, air temperature increase could refer to nocturnal increases due to urban heat island effect (but visibility refers to noon). Some clarifications are required.

Yes, annual visibility was measured at 12 noon. The other variables such as RH, air temperature, wind speed and wind direction are also average values for 12 noon. This information has now been included in supplementary Figure S4 caption, which has been removed from the main manuscript (Figure 5) to supplement as per suggestion of Reviewer 1.

Wd refers to the prevailing wind direction. More explanation is now added in Page 6 Line 18 “These calculations were performed using the timePlot function in the openair package for R statistical program, which works on vector functions for wind direction averaging.”

We looked at the correlation between different meteorological variables. This data is now provided in Supplementary Figure S5. Statistically significant correlations are observed between visibility and air temperature, relative humidity and wind speed.

Model: The authors present a model for light extinction, making a number of assumptions and simplifications. Which could be the cost (uncertainty arising from these assumptions)? Despite assumptions, the model has an absolutely perfect performance with observations. Any explanation? What about the other stations?

We understand the excellent agreement between model and observations as evidence that the model is capturing the key features of the underlying physics of atmospheric visibility. We were very pleased by this result! Whilst the model is good – it is not perfect – and there are a number of mismatches between model and observation albeit slight. We state clearly in the manuscript the assumptions we make, in particular, we state the following on Page 8 Line 1 “To reduce the number of parameters within Eq. (5), it is assumed that  $\beta_{abs}(RH) = \beta_{abs}(dry)$ . This assumption always holds for gas absorption; and it is largely true for aerosol particles as well, although it is noted that particle absorption can increase due to lensing effects in mixed phase aerosol, and this lensing effect will be affected by aerosol water content e.g. (Lack and Cappa, 2010).”

Page 4, line 5: The aim of the study is implemented? what do you mean UK projections of meteorology (climate change? it is not clear). And what do you mean with pollution projections? Local or regional? What kind of projections? For which pollutants?

This sentence was unclear. We now state the following “A new model is also presented which can aid in future visibility prediction under different climate and pollution scenarios.” We hope the model parameterization will be used by many future researchers who will find varied uses for it. For example, we can imagine that effect of both local and regional pollution upon visibility could be investigated. The effect of future RH predictions under climate change conditions could also be investigated.

### **Minor Comments**

Abstract, Line 2. It can be removed from abstract

We believe this sentence is useful to provide context to the study and have not removed it.

Page 2, line 16: rearrange using chronological order

Thanks for this comment. We have found that ACP referencing format does not support chronological order. We have used ACP templates in EndNote, hence could not arrange the references in chronological order.

In the analysis of week day variations of visibility , the information provided in Page 12, line 14 is confusing and I also think wrong (regarding the calculations). I do understand the meaning of this analysis.

The calculation was wrong. We have simplified the text to the following - “Lower traffic and industrial emissions at the weekend are the likely reasons for better visibility at the weekend due to less pollutant emissions. The inherent assumption in this analysis is that traffic is higher during week days compared to the weekend. It is noted that visibility tends to peak on Sunday (rather than both Saturday and Sunday) and this may reflect the non-negligible timescale required for pollutant removal by wind driven dispersion, i.e. the build-up of pollution during weekdays is not fully dispersed until Sunday. The same argument explains why visibility is typically higher on Mondays compared to the other weekdays later in the week.”

In Figures 3 (right side), it is better to use normalized values. For instance you can normalize values with the maximum visibility value for a direct estimation of % differences.

Figure 3 has been updated to use normalized values for the day of the week graph, where mean weekday visibility is normalized to the mean Sunday values for direct estimate of the percentage differences in weekday visibility values.

Page 5, line 25: do you mean the sensor was not cleaned? How can you be sure that all other stations are cleaned properly?

Unfortunately, we have no record for when they were cleaned, however we were informed by the UK Met Office that protocols were followed with regards to cleaning after high aerosol sea salt loads at coastal stations. As we described in the main manuscript, Tiree Island has a very flat landscape, which is not sheltered from the wind in any direction, possibly allowing salt to accumulate.

### **Technical Comments**

Although English is in general good, some syntax errors exist in the paper. Missing comma in many cases make the text hard to understand. Figures quality needs to be improved. Use legends in Fig. 3 or use analogous (with variables) colors in the axis Fig3. Indicate in the legends what dashed lines represent (left side) and bars (right side).

Thanks for highlighting this. Suggested changed has been implemented in manuscript, where the typos in the manuscript have now been remedied. The higher resolution figures will be uploaded with the final manuscript. As per your suggestion we have also included analogous colours for variables in the Figure 3.

### **References**

- Crilley, L. R., Bloss, W. J., Yin, J., Beddows, D. C., Harrison, R. M., Allan, J. D., Young, D. E., Flynn, M., Williams, P., and Zotter, P.: Sources and contributions of wood smoke during winter in London: assessing local and regional influences, *Atmospheric Chemistry and Physics*, 15, 3149-3171, 2015.
- Doyle, M., and Dorling, S.: Visibility trends in the UK 1950–1997, *Atmospheric Environment*, 36, 3161-3172, 2002.
- Founda, D., Kazadzis, S., Mihalopoulos, N., Gerasopoulos, E., and Lianou, M.: Long-term visibility variation in Athens (1931–2013): A proxy for local and regional atmospheric aerosol loads, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2015-1025, in review, 2016., 2016.
- Harrison, R. M., Pope, F. D., and Shi, Z.: Trends in Local Air Quality 1970–2014, *Still Only One Earth: Progress in the 40 Years Since the First UN Conference on the Environment*, 2015, 58,
- Lack, D., and Cappa, C.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, *Atmospheric Chemistry and Physics*, 10, 4207-4220, 2010.

Williams, M.: Air pollution and policy—1952–2002, *Science of the total environment*, 334, 15-20, 2004.



# 60 years of UK visibility measurements: impact of meteorology and atmospheric pollutants on visibility

Ajit Singh, William J. Bloss and Francis D. Pope

School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, B15 2TT, UK

*Correspondence to:* Francis D. Pope ([f.pope@bham.ac.uk](mailto:f.pope@bham.ac.uk))

## Abstract

10 Reduced visibility is an indicator of poor air quality. Moreover, degradation in visibility can be hazardous to human safety; for example, low visibility can lead to road, rail, sea and air accidents. In this paper, we explore the combined influence of atmospheric aerosol particle and gas characteristics, and meteorology, on long-term visibility. We use visibility data from eight meteorological stations, situated in the UK, which have been running since the 1950s. The site locations include urban, rural and marine environments.

15 Most stations show a long term trend of increasing visibility which is indicative of reductions in air pollution, **examples of urban areas**. Additionally, the visibility at all sites show a very clear dependence on relative humidity indicating the importance of aerosol hygroscopicity on the ability of aerosol particles to scatter radiation. The dependence of visibility on other meteorological parameters, such as wind speed and wind direction is also investigated. Most stations show long term increases in temperature which can be ascribed to  
20 either climate change, land-use changes (e.g. urban heat island effects) or a combination of both; the observed effect is greatest in urban areas. The impact of this temperature change upon local relative humidity is discussed.

To explain the long term visibility trends and their dependence on meteorological conditions, the measured data were fitted to a newly developed light extinction model to generate predictions of historic aerosol and gas scattering and absorbing properties. In general, an excellent fit was achieved between measured and modelled  
25 visibility for all 8 sites. The model incorporates parameterizations of aerosol hygroscopicity, particle concentration, particle scattering, and particle and gas absorption. This new model should be applicable and is easily transferrable to other data sets worldwide. Hence, historical visibility data can be used to assess trends in aerosol particle properties. This approach may help constrain global model simulations which attempt to generate aerosol fields for time periods when observational data are scarce or non-existent. Both the measured  
30 visibility and the modelled aerosol properties reported in this paper highlight the success of the UK's Clean Air Act, which was passed in 1956, in cleaning the atmosphere of visibility reducing pollutants.

**Keywords:** Visibility, meteorology, pollutants, aerosol, RH and hygroscopicity

## 1 Introduction

The meteorological definition of visibility is the “distance at which the contrast of a given object with respect to its background is just equal to the contrast threshold of an observer” (WMO, 1992, 2015). In general, good visibility is a desirable feature of any geographical location and its importance should not be neglected (Doyle and Dorling, 2002). Poor visibility (< 2.0 km, (Founda et al., 2016)) can affect the transportation of goods and people, whether it is by rail, road, sea or air. Low visibility can lead to accidents and thus is a concern for public safety. Tourism is often dependent on good visibility for appreciation points of interest (Singh and Dey, 2012). For example a study at Grand Canyon Park in USA has shown that visitor frequency in the park has reduced as visibility decreased (NAPAP, 1990).

Typically in cloud free sky, visibility can vary from ca. 5 - 100 kilometres dependent on atmospheric composition and conditions. Visibility is reduced by the interaction of light with atmospheric gases and aerosol particles which can absorb or scatter the light; consequently visibility is greatest within non-polluted pristine atmospheres, other factors (e.g. meteorology) being equal. Many previous studies have investigated the link between atmospheric composition and visibility (Jinhuan and Liquan, 2000;Schichtel et al., 2001;Wu et al., 2005;Park et al., 2003;Yang et al., 2007;Tiwari et al., 2011;Park et al., 2006;Founda et al., 2016;Cao et al., 2012;Watson and Chow, 2006). These investigations demonstrate that visibility is markedly influenced by the size, chemical composition, and concentration of airborne particles. Reduced visibility is attributed mainly to high concentrations of aerosol particles, and in general, scattering effects are the dominant visibility reducing mechanism within the atmosphere. Within heavily polluted atmospheres, visibility can decrease rapidly due to the presence of aerosol particles (Husar et al., 1981). For example, during the 1952 London smog events visibility declined to a few metres due to high air pollution (caused by a rise in smoke and other pollutant concentrations in the atmosphere (Wilkins, 1954)) as discussed in detail by Brimblecombe (1987). More recently, a study by Sati and Mohan (2014) also found sharp decreases in visibility due to increased particulate matter (PM) and NO<sub>2</sub> concentrations during a smog event on November 2012 at Delhi. Similarly, Zhang et al. (2006) described the PM influence upon visibility reduction at Beijing, China. Festivals involving fireworks, which release aerosol particles upon detonation, are a good example of spatially and temporally localised pollution events which may lead to reduced visibility (Singh et al., 2015;Seidel and Birnbaum, 2015;Kong et al., 2015).

In addition to aerosol and gas concentrations and composition, specific meteorological conditions can also affect visibility (Sloane, 1983). There exists a body of literature on urban visibility studies, which attempt to connect visibility with meteorological parameters (e.g. (Hänel, 1972;Clarke et al., 1978;Lee, 1983;Lee, 1990;Haywood and Boucher, 2000)).Whilst temperature (T), relative humidity (RH), wind speed ( $w_s$ ) and wind direction ( $w_d$ ) do not affect clear sky visibility directly, they can influence the sources and sinks of the trace gases and aerosol particles in the atmosphere. For example, high wind speeds can re-suspend dust particles and generate sea spray aerosol particles. Windy conditions can also lead to a cleaning effect by replacing polluted air with cleaner air. Temperature can influence the production of secondary organic aerosol (SOA) particles, for example, via the chemical formation and partitioning between the gas and particle phase. Relative humidity (RH) not only affects the sources and sinks of gases and aerosols, it also directly influences the size and composition of aerosol particles. Nearly all atmospheric aerosol particles are hygroscopic to some degree; hence, their size is dependent

upon the local RH. As RH increases, hygroscopic particles take up water, through absorption and adsorption, and grow in size, volume and weight. The addition of water also changes the overall particle composition. This typically lowers the mean refractive index of the particle since the refractive index of water is lower than other common aerosol components, such as minerals, organics, sulphates and nitrates (Harrison et al., 2004). Under high humidity conditions, a high particle loading in the lower atmosphere can increase fog formation and thus severely reduce visibility (Tiwari et al., 2011). It has previously been shown that monthly variations in visibility are negatively correlated with RH (Singh and Dey, 2012). Other studies have shown how the RH effect on particle hygroscopic growth can influence in visibility change (Liu et al., 2012). Thus, both PM loading and meteorological factors, such as relative humidity, are important for the assessment of the causes of visibility reduction. Other factors may also be important such as vegetation density, industrial development, urbanization and human population since these factors affect surface type and can effect aerosol deposition (Diederer et al., 1985).

In the last few years, worldwide interest in atmospheric visibility has grown, but few studies examine UK visibility. Previously, a long-term trend analysis of visibility was performed at eight UK weather stations between 1950 and 1997 by Doyle and Dorling (2002), where improved visibility was identified at most of the stations, mainly after 1973 due to oil crises and less consumption. Summer visibility trends for five different sites in London and southern England for the period of 1962 to 1979 were analysed by Lee (1983), and it was also found that a rise in visibility was observed at all sites. Gomez and Smith (1984) quantified the seasonal visibility trends at Oxford during 1926 to 1985 and observed a clear reduction in visibility from 1926 to 1944, a notable rise after World War II from 1944 to 1952, and another reduction from 1952 to 1966 (mainly in the summer season); the visibility improved again after 1966 in all seasons due to the reduction in aerosol concentration (Gomez and Smith, 1987). It is also found that after the 1956 Clean Air Act, fog occurrence has been declined at Oxford and nearby rural areas due to drop in smoke concentration, urban heat Island effect and other public activities (Gomez and Smith, 1984). Analyses by Lee (1985) in central Scotland for the period of 1962 to 1982, has mentioned about the effect of 1973 oil crises on visibility and air quality, where a significant increase in visibility was shown primarily in urban areas due to a major reduction in sulphate aerosol concentration. A similar study on historical visibility trends at 22 different UK meteorological stations (includes urban, rural and marine areas) during 1962 to 1990 was performed by Lee (1994). A clear rise in visibility was identified at most of the sites due to reduction in coal and smoke emissions (Lee, 1994). Furthermore, a steady reduction in fog frequency with improved visibility correlated with decreased smoke pollution at Glasgow airport was noted (Harris and Smith, 1982). The correlation between various air pollutants (such as  $\text{NH}_4^+$ , and non-marine pollutants  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$ ) and visibility at northwest England, UK were also performed in the 1980s, where strong negative correlations were found between visibility and these pollutants (Colbeck and Harrison, 1984). At present, most UK urban cities are relatively polluted (Defra, 2011) compared to rural locations, with pollutant sources dominated by vehicular emissions (Colvile et al., 2001). The 1956 Clean Air Act led to general improvements in UK air quality; however, there still exist many negative effects of air quality on the UK population such as impaired human health ((Defra, 2011; Harrison et al., 2015).

The present study investigates visibility in the UK focusing on 8 specific sites. The same sites were previously investigated by Doyle and Dorling (2002) who presented long term UK visibility trends for 1950-1997 and the

dependence of the measured visibility on meteorological conditions. In this paper we build upon the work of Doyle and Dorling to analyse UK visibility trends from 1950-2013. Furthermore, we extend the analysis by investigating causes of the observed visibility trends; in particular we investigate the role of air pollutant concentrations upon UK visibility. The outputs from this work help to explain historic visibility trends in the UK. A new model is also presented which can aid in future visibility prediction under different climate and pollution scenarios.

## 2 Data

Daily archived horizontal visibility data, defined as the visibility distance along a horizontal line at the earth's surface, were obtained from the British Atmospheric Data Centre (BADC) which is run by the UK's Natural Environment Research Council ([www.badc.nerc.ac.uk](http://www.badc.nerc.ac.uk)). The archive contains visibility data, in addition to other relevant meteorological parameters, archived at an hourly time resolution. In addition to visibility, the following meteorological parameters were also utilised: RH, wind speed, wind direction and air temperature, present weather (PR) code which provides further qualitative detail about the weather conditions. A description of the present weather codes is provided in the table ([www.badc.nerc.ac.uk/data/ukmo-midas/WH\\_Table.html](http://www.badc.nerc.ac.uk/data/ukmo-midas/WH_Table.html)) at [www.badc.nerc.ac.uk](http://www.badc.nerc.ac.uk). Unfortunately the use of present weather codes largely ceased with the introduction of automated meteorological stations and insufficient PR codes were available after the year 1997. It is noted, that if the present weather codes were available they would have been useful to screen the data for rain or other precipitation events. Due to unavailability of present weather codes during required study period (1950-2012), data filtration was done on the bases of RH limits instead of PR codes. Data were removed when the relative humidity reading was > 99 % which is highly suggestive of rain or other precipitation events. Removal of data with RH > 99% removes between 0.91 – 3.44 % of the data dependent on site location. Since the ability of visibility observers is affected by light levels, with greater difficulty encountered in night time measurements (Lee, 1990) the daily data used for this study were all measured at 12 noon for all sites.

Meteorological data were collected for the eight UK stations which possess near continuous time series data starting in the 1950s and continuing to present day. The eight stations are Aldergrove, Heathrow, Ringway, Nottingham, Plymouth, Tiree, Leuchars, and Waddington, and details of the stations are given in Table 1 and Fig. 1.

The visibility data sets are based on ground based measurement using a variety of techniques. More details of visibility observations method are found in UK Met Office Surface Data Users guidelines ([https://badc.nerc.ac.uk/data/ukmo-midas/ukmo\\_guide.html](https://badc.nerc.ac.uk/data/ukmo-midas/ukmo_guide.html)). Until the late 1990s all visibility measurements were performed by human observer. Subsequently data collection was automated using visibility sensors (visiometers). See supplementary material Table S1 for detail on measurement type used and dates of service.

There are advantages and disadvantages with both human observation and visiometers. Clearly from a manpower perspective, visiometers are preferred. Human observation provides a true measure of visibility since

the observer is looking for objects located at a known distances away from their location, however, the visibility measurements are imprecise by nature since results can vary according to the contrast and illuminance thresholds (ability to discern and sensitivity to light, respectively) of the observer's eyes (WMO, 2008). Since human observation requires objects to observe the measurement is quantized by the geographical spread of available objects i.e. there is not a continuum of measurement locations. Consequently, human observations provide a lower limit to the actual visibility. Distances between objects to observe can be large especially at the longer distances measured (> 10 km) which leads to reductions in accuracy at high visibility. At high elevation the visibility calculation can be different from that at the surface (Malm et al., 1981). Visiometers automatically measure the extinction of light over a small distance (typically ca. 1 m) and from the measured extinction can estimate visibility. In particular automatic visibility measuring instruments consist of a light transmitter and receiver, the light extinction observed between the transmitter and receiver is then used to estimate the visibility (Jebson, 2008). These automated estimates of visibility are more objective and reproducible compared to human observation. However, since the visiometer only measures air local to the device it can be much more affected by variations in local air quality. This is likely to be a more important consideration at urban meteorological sites where air composition is more heterogeneous, compared to rural sites, due to the greater number of pollutant sources in urban areas.

The change from human to automatic measurement occurred at different times for the different sites (see supplementary Table S1). It is clear for most sites investigated, that the changeover from manually observed to automatically measured data leads to step changes in the visibility reported, see Fig. 2 and further discussion in methodology section. This is unsurprising given the discussion above. In particular, clear deviations away from the long term trend measured under human observation are observed at Aldergrove, Plymouth and Tiree stations once automation was introduced (see supplementary Table S1). After consultation with the UK Met Office it was noted that automated sensors can be unreliable during high visibility events when compared to human readings. In particular automatic sensors perform sub-optimally at coastal sites unless the sensor is cleaned regularly, due to accumulation of sea salt residue. Unfortunately, the Tiree station was reported to fall into this category.

To assess the effects of the gaseous pollutant nitrogen dioxide (NO<sub>2</sub>) on visibility, daily ground based measured data of NO<sub>2</sub> was obtained from the Department of Environment Food and Rural Affairs (Defra) (<https://uk-air.defra.gov.uk/>) for one observing station (Harlington), closely co-located to the Heathrow meteorological station (ca. 1.3 miles distant). NO<sub>2</sub> data were only available for 9 years (2004-2012) of the visibility study period.

### 3 Methodology

#### 3.1 Trend analysis of visibility and other meteorological parameters

60 year trend analyses have been performed on the visibility dataset described in section 2. For long term trend analysis each days value was averaged (simple mean) to determine trends over decadal, annual and seasonal

cycles. The seasonal periods were defined, as is typical, as winter (Dec-Feb), spring (Mar-May), summer (Jun-Aug), and autumn (Sep-Nov). Diurnal, day of the week and monthly averaged trends of visibility and RH were determined at each site using the 60 years of dataset, where weekdays and weekend are categorised as Monday-Friday and Saturday-Sunday respectively.

- 5 To examine the hygroscopic growth effect of aerosol particles upon visibility, the decadal data sets were disaggregated into RH bins. The aerosol hygroscopic growth effect on visibility was examined by using decadal mean visibility within specific relative humidity bins with the following boundaries: 52.5-57.5 %, 57.5-62.5 %, 62.5-67.5 %, 67.5-72.5 %, 72.5-77.5 %, 77.5-82.5 %, 82.5-87.5 %, 87.5-92.5 %, 92.5-97.5 %. We excluded data with  $RH > 97.5$  % due to likely presence of fog and mist at RH greater than this threshold.
- 10 To highlight the daily variation in RH, histograms of daily RH (at 12 noon) were generated using the following boundaries (0-10 %, 10-20 %, 20-30 %, 30-40 %, 40-50 %, 50-60 %, 60-70 %, 70-80 %, 80-90 %, and 90-100 %).

To evaluate the dominant meteorology at each site several meteorological analyses were conducted. Wind rose plots using the complete dataset time series were generated to highlight the dominant wind speed and direction for all sites. Decadal-seasonal bivariate polar plots of visibility using wind direction and wind speed allow for spatial analysis of likely pollution sources (Carslaw and Ropkins, 2012). Finally time series plots of the following meteorological parameters were generated, RH, wind speed, wind direction and air temperature. **These calculations were performed using the timePlot function in the openair package for R statistical program, which works on vector functions for wind direction averaging.**

20

### 3.2 Estimation of aerosol and gas phase properties through analysis of RH dependent visibility

In this section the contribution of aerosol particles and gases upon visibility is estimated via mathematical modelling. In general horizontal visibility ( $V$ ) can be defined via Koschmieder equation Eq. (1), where, horizontal visibility shows an inverse relationship with the extinction coefficient ( $\beta_{ext}$ ). In the Eq. 1, constant ( $k$ ) is equal to 3.912 which assumes a contrast threshold of 2 % (Koschmieder, 1924). The constant ( $k$ ) is a measured by the threshold sensitivity of the observer's eye (Schichtel et al., 2001; Chang et al., 2009), which can vary from 2 to 5 % (Appel et al., 1985).

30

$$V = k/\beta_{ext} \quad (1)$$

The extinction coefficient depends upon ( $\beta_{ext}$ ) is the sum of the scattering ( $\beta_{sca}$ ) and absorption coefficients ( $\beta_{abs}$ ) as shown in Eq. (2).

$$\beta_{ext} = \beta_{sca} + \beta_{abs} \quad (2)$$

In the atmosphere, aerosol particles and gas phase species can both contribute to light scattering and absorption. However, the contribution of gas phase scattering to the total extinction is negligible except in the most pristine environments. Hence under UK conditions, the scattering component of the extinction coefficient can be assumed to be completely dominated by the presence of aerosol particles.

The ability of an individual particle to scatter radiation is dependent on its size, shape, morphology and refractive index (Appel et al., 1985; Liu and Daum, 2000). The particle scattering coefficient ( $\beta_{sca}$ ) can be estimated by Mie theory as shown in Eq. (3) (Tang, 1996);

$$\beta_{sca} = \int_0^{\infty} \pi \left(\frac{D}{2}\right)^2 Q_{sca}(\alpha, \lambda, n) Nf(D) dD \quad (3)$$

Where,  $D$  represents particle diameter, the aerosol size distribution is given by  $Nf(D)$  and  $\alpha$  is the size parameter ( $\alpha = \pi D/\lambda$ ).  $N$  is particle number concentration and  $Q_{sca}(\alpha, \lambda, n)$  is single-particle scattering cross section, which depends upon size parameter ( $\alpha$ ), wavelength ( $\lambda$ ) and refractive index ( $n$ , which is composition dependent). All these particle characteristics can change as the particle undergoes water uptake or loss which is dependent on the local RH. To parameterise the aerosol scattering enhancement due to water uptake an approach, similar to Titos et al. (2014), is taken. The scattering enhancement is parameterised using a single hygroscopicity parameter ( $\gamma$ ) using Eq. (4), where  $\beta_{sca}(RH)$  and  $\beta_{sca}(dry)$  are the aerosol scattering coefficients under a specified RH condition and completely dry conditions, respectively.

$$\frac{\beta_{sca}(RH)}{\beta_{sca}(dry)} = \left(1 - \frac{RH}{100}\right)^{-\gamma} \quad (4)$$

Rearranging Eq. (1), Eq. (2), and Eq. (4) allows for the relationship in Eq. (5) to be derived, where  $\beta_{abs}(RH)$  and  $\beta_{abs}(dry)$  are the combined aerosol and gas absorption coefficients under a specified RH condition and completely dry conditions, respectively.

$$Vis(RH) = \frac{3.912}{\left(1 - \frac{RH}{100}\right)^{-\gamma} \times \left(\frac{3.912}{Vis(dry)} - \beta_{abs}(dry)\right) + \beta_{abs}(RH)} \quad (5)$$

To reduce the number of parameters within Eq. (5), it is assumed that  $\beta_{abs}(RH) = \beta_{abs}(dry)$ . This assumption always holds for gas absorption; and it is largely true for aerosol particles as well, although it is noted that particle absorption can increase due to lensing effects in mixed phase aerosol, and this lensing effect will be affected by aerosol water content e.g. (Lack and Cappa, 2010).

- 5 Equation (5) can be further simplified by assuming that all absorption due to both gases and particles is negligible compared to the RH dependent aerosol scattering, leading to the two parameter Eq.(6).

$$\log [Vis(RH)] = \gamma \log \left[ 1 - \left( \frac{RH}{100} \right) \right] + \log [Vis(dry)] \quad (6)$$

- 10 Equations (5) and (6) can be used to obtain information about aerosol scattering and gas and aerosol absorption, with associated assumptions, through fitting of the measured visibility at a given RH. Equation (6) is linear and so can be fitted using the linear least squares fitting algorithm, whereas Eq. (5) requires non-linear least squares fitting algorithm. The statistical program R was used for all fittings (Version 0.99.489). The ‘lm’ algorithm was used for linear fitting, and the ‘nls’ fitting algorithm was used for the non-linear fitting. The ‘nls’ algorithm was  
 15 always initially run with no lower or upper boundaries for the 3 fitting parameters ( $Vis(dry)$ ,  $\beta_{abs}$  and  $\gamma$ ) specified. However, when fits produced negative values for  $\beta_{abs}$ , which are physically impossible, a lower boundary for  $\beta_{abs}$  was specified to be zero.

### 3.3 Gas absorption

- 20 All gases scatter radiation via Rayleigh scattering but the effect is negligible in all but the most pristine visibility conditions (which are not observed in this study). The only atmospheric gas present at levels that lead to significant absorption of visible light is  $NO_2$  (Ferman et al., 1981; Groblicki et al., 1981). The contribution of  $NO_2$  to visibility can be quantified by its absorption coefficient ( $\beta_{NO_2abs}$ ). The effect of the  $NO_2$  absorption coefficient, at 550 nm wavelength, was calculated using the relationship from Groblicki et al. (1981), shown in  
 25 Eq. (7), where  $[NO_2]$  is the  $NO_2$  in ppm.

$$\beta_{NO_2abs} = 3.3 \times 10^{-4} [NO_2] \quad (7)$$

## 4. Results and Discussion

### 4.1 Historical trend of annual and seasonal visibility



The annual and seasonal mean visibility at 12 noon have been calculated for all eight stations, see Fig. 2. The effect of changing the visibility observation technique from human observation to automatic observation via visimeters (which is highlighted by different shading in Fig. 2) is very clear at some sites. In particular, two stations, Tiree and Aldergrove, do not show realistic values after the changeover from human to automated measurement, with the changeovers coinciding with large and sustained drops in recorded visibility. The effect of human to automated changeovers at Heathrow, Leuchars, Nottingham, Ringway and Waddington sites appears to be minimal, with the pre-changeover long term trends being continued after the changeover. Furthermore the annual data from these sites exhibit similar year to year variance before and after changeover. The long term trend at the Plymouth site is similar before and after changeover but the year to year variance is much reduced once measurement automation is installed. This likely indicates strong localised sources (ship and traffic emissions from nearby ports and roads) close to the visimeter at the Plymouth site. Henceforth it is assumed that all stations, except Aldergrove and Tiree, are performing adequately for both human and automated visibility measurement. Therefore the time series, as shown in Fig. 2, are used in their entirety for the analysis of these six stations. The time series data for the Aldergrove and Tiree stations are used up until automation occurs.

A similar variation in visibility trends is observed for the period of 1950-1997, comparing with Doyle and Dorling (2002). However, this study reports overall lower visibility values when compared to Doyle and Dorling (2002). These differences are due to slightly different data filtering methodologies. Doyle and Dorling (2002) filtered data for 12 noon, relative humidity > 90% and PR codes of 00-05 in their statistical analysis for the period of 1950-1997. However, due to uncertainty and unavailability of PR code after 1997 we did not use these codes. Furthermore we performed mean averaging for statistical analysis, where data is filtered for 12 noon and relative humidity > 99 %. The details of uncertainty and unavailability of PR codes and used data filtration method are given in data and methodology sections

Clear trends of increasing annual visibility are observed for four sites: Ringway, Waddington, Nottingham, and Heathrow with the rate of visibility increase being  $0.339\pm 0.016$  km year<sup>-1</sup>,  $0.293\pm 0.010$  km year<sup>-1</sup>,  $0.235\pm 0.023$  km year<sup>-1</sup> and  $0.201\pm 0.018$  km year<sup>-1</sup>, respectively, where standard errors were determined at the 95 % confidence interval. A, more gradual increasing trend was observed at the Leuchars site ( $0.157\pm 0.019$  km year<sup>-1</sup>). The Plymouth site shows a more variable trend with increases from ca. 1950-1990 followed by decreases from ca. 1990-2006 which is then followed by more increases in the most recent measurements. The long term trend for Plymouth 1950-2013 is near constant ( $0.040\pm 0.021$  km year<sup>-1</sup>). Both the Aldergrove and Tiree sites, with the automated data omitted, show near constant long term visibility with long term rates of visibility change calculated to be  $0.0562\pm 0.021$  km year<sup>-1</sup> and  $-0.0892\pm 0.014$  km year<sup>-1</sup>, respectively.

The seasonal trends for the 8 sites are detailed in Table 2. Poorest visibility was observed in the winter season compared to other seasons mostly due to the seasonal rise in RH (discussed in section 4.3). Another reason is the greater concentration of particles in the environment due to lower mixing layer height in the winter season (Jayamurugan et al., 2013). Furthermore, the long term rate of visibility change in the winter season is significantly higher as compared to spring, summer and autumn seasons for all stations apart from the Ringway station. At Ringway station the rate of change of visibility is higher in spring ( $0.363\pm 0.018$  km year<sup>-1</sup>) as compared to winter ( $0.330\pm 0.020$  km year<sup>-1</sup>). All stations show positive rates of visibility change in winter

season except for Tiree ( $-0.186 \pm 0.012 \text{ km year}^{-1}$ ). It is also observed that Aldergrove station shows negative rate of visibility change in the summer season ( $-0.417 \pm 0.036 \text{ km year}^{-1}$ ).

The improvement in median visibility at most of the sites can be seen in supplementary Fig. S1. Moreover, Boxplots of the decadal visibility are also produced showing the median, interquartile range, outliers etc. (see supplementary Figure S2).

Improved visibility at most of the sites is due to reduction in air pollution and the likely changes in fuel use and consumption that took place after 1956 Clean Air Act. The Clean Air Act was introduced with the aims of reducing smog, smoke and sulphur dioxide concentrations in the environment. In particular, the policy focused on industrial emission sources and its reduction (Williams, 2004). Recently, Harrison et al. (2015) shown that concentration of sulphur dioxide, coal smoke, nitrogen dioxide, suspended matter (black smoke) and PM were significantly reduced in the UK over last five decades as the result of switching to cleaner fuels after 1956 Clean Air Act.

## 4.2 Evaluation of historical wind-data

### 4.2.1 Wind Roses for the 8 stations

A graphical representation of historical wind speed and direction at the eight chosen stations is shown in Fig. 1 using the wind rose polar co-ordinate representation. These graphs describe the most probable wind speeds and directions over the whole time series (Carslaw and Ropkins, 2012). As expected, the graphs show that the predominant wind directions in the UK are from the southwest. However, there are clear variations between the different stations. The range of wind speed varies from  $0\text{-}35 \text{ m s}^{-1}$  dependent upon location, with the more coastal sites experiencing greater average wind speeds.

### 4.2.2 Analysis of influence of wind speed and wind direction on visibility

Decadal-seasonal bivariate polar plots are presented for all eight stations in supplementary Fig. S6; these diagrams provide information on the variation of visibility with wind speed and direction and can suggest locations for visibility degrading sources. The detailed analyses of each site are given below:

**Aldergrove:** Overall, lower values of visibility were observed when the wind was from the south to east, while above average values were collected when the wind was from the north to west direction. Intermediate visibility was generally observed when the wind came from the south to west or north to east quadrants. Distinct differences are observed between the different seasons. In particular, in the summer visibility with wind from the north to west direction was higher compared to other seasons in every decade. It is clearly seen that visibility has improved the most when wind comes from the south to east direction which covers mainland urban areas such as Belfast, the major regional city. It is noted that the seasonal and polar trends are similar between the visiometer (1950s–1990s) and human derived (2000s–2010s) data sets even though the absolute magnitudes are different as noted above.

**Heathrow:** Low visibility was observed whenever wind speeds were lower than  $5 \text{ m s}^{-1}$  in any direction which implies a significant local source of visibility degrading pollutants. Since Heathrow is the site of major international airport, with commensurate road and other transport infrastructure, this is not surprising. Overall, lower visibility is also seen when the wind direction comes from the northeast to southeast direction which is consistent with visibility reducing pollution arriving from the Greater London area. The highest visibilities are typically observed when the wind direction is from the north to southwest which is consistent with less densely populated surrounding areas. In particular during summer visibility in the northwest wind direction was highest compared to other seasons in every decade. It is identified that visibility has improved in all wind directions, but most significantly in the easterly direction which covers the London urban centre. The change in visibility illustrates the dramatic improvement of air quality in London since the introduction of the Clean Air Act in 1950s (Brimblecombe, 2006).

**Leuchars:** Two distinct spatial groupings of visibility are clearly observed. When the wind direction comes from the northeast to southwest (clockwise) visibility is generally lower, and it is generally higher when the wind direction is from the northeast to southwest (anti-clockwise). The lowest visibilities are from the southeast direction in all seasons. The spatial pattern of low visibility suggests a maritime aerosol source as the major source of visibility reduction whilst high visibility was associated with air which had passed over the predominantly rural Scotland. Visibility in the northwesterly wind direction was highest in the summer months, as expected see Fig. 2 and 3, compared to other seasons in every decade.

**Nottingham:** Like Heathrow, the poorest visibility conditions occurred when wind speed was below  $10 \text{ m s}^{-1}$  suggesting local sources of visibility degrading pollutants. Visibility is often lowest when the wind comes from the southeast direction consistent with the relative placement of Nottingham city centre to this direction (the meteorological station is actually located in Watnall just about 5 miles of Nottingham city centre). Visibility is generally highest when the wind comes from the west and southwest directions which is largely consistent with air masses passing over less urban areas compared to the other wind directions. During the summer months, visibility in southwest direction was highest compared to other seasons in every decade. It is clear from Fig. 2 that visibility has increased in all seasons, and the strongest improvement is seen in air from the southeast as seen in [supplementary Fig. S6](#).

**Plymouth:** In general, the lowest visibility was observed when the wind comes from southeast to southwest direction which is consistent with maritime air causing the lowest visibility which suggests a maritime source of aerosol causing visibility degradation. The highest visibilities are observed when wind comes from the northwest to northeast directions, and in particular the northeast, this is consistent with airmasses passing over relatively rural areas. Regardless of the direction of wind, the summer months showed higher visibility than all other seasons. It is identified that visibility has improved over time for all wind directions.

**Ringway:** Overall visibility was poor at low wind speeds and when the wind direction was from the northeast to southeast. Ringway is the location of Manchester International Airport so, like Heathrow, there is likely to be a significant local source of visibility degrading pollutants arising from the airport and its associated infrastructure. The wind directions associated with higher visibility are a lot more variable in time and space when compared to other locations. However, in general, high wind speeds from either the northwest or south

west directions are often associated with higher visibility. Since the 1960s visibility has improved for all wind directions. In particular, visibility associated with air masses coming from the direction of the Greater Manchester Area to the north has shown a marked increase since the 1970s.

5 **Tiree:** The island of Tiree has by far the highest visibility at low wind speeds. Overall low visibility was observed when wind came from the west to southeast, while highest visibility occurred with wind from the northeast. The spatial variation of low visibility is consistent with a maritime source of visibility impairing aerosols. The higher the wind speed typically the lower the visibility which is consistent with greater aerosol production from greater wave activity (Venkataraman et al., 2002). The higher visibility from the northeast is consistent with air masses passing over the larger rural highlands of Scotland. Visibility was relatively stable for  
10 all wind directions for all decades of the **human** observation data series which is consistent with this rural maritime site being largely unperturbed by anthropogenic pollution.

**Waddington:** In general, lower visibility is observed when wind speeds are lower than  $10 \text{ m s}^{-1}$  which is consistent with local pollution sources. Low visibility is also observed when the wind direction is from the east to southeast which potentially indicates a maritime source. Higher visibility is observed from the west at high  
15 wind speeds. Visibility has improved for wind from all directions since the 1970s.

Overall it is clear that visibility has improved at most of the sites for most local wind directions. The most marked improvements in visibility are seen in directions when air masses pass over major metropolitan areas such as Greater London and Greater Manchester. Whilst most of the visibility changes can be ascribed to the location of the meteorological stations with respect to either urban or maritime sources, it is noted that for most  
20 sites the wind direction with the lowest visibility overall is often from the East, i.e. continental Europe and hence synoptic scale pollution events which affect visibility. Poor air quality, in the UK, is often associated with synoptic scale events originating in continental Europe (Charron et al., 2007a;Charron et al., 2007b;Charron et al., 2013;Lee et al., 2006;Crilley et al., 2015)

#### **4.3 Correlation between RH and visibility: seasonal, day of the week and decadal effects**

25 Figure 3 provides monthly values for visibility and RH, averaged over the whole time series, for each station. This figure clearly illustrates that visibility shows a strong seasonal cycle which is anti-correlated with RH at all stations. The relationship at Tiree is less strong compared to the other seven sites. The geographical location of Tiree, which is a maritime island, is the likely reason for the RH trend being different to the other stations. Tiree Island has a very flat landscape, which does not provide shelter from wind in any direction; this directly affects  
30 the local meteorology (Holliday, 2004). Overall, the monthly trends indicate that visibility is lowest in winter and highest in summer with spring and autumn being intermediate in visibility values.

In addition to the seasonal cycle, there is a clear day of the week effect on visibility changes at most sites (Fig. 3), where visibility improves sharply at the weekend with Sunday showing the highest visibility. It is observed that visibility improves at Sunday from 5 % to 12.5 % (depending upon area) as compared to other week days  
35 (Mon-Fri). **Lower traffic and industrial emissions at the weekend are the likely reasons for better visibility at the weekend due to less pollutant emissions. The inherent assumption in this analysis is that traffic is higher during week days compared to the weekend. It is noted that visibility tends to peak on Sunday (rather than both**

Saturday and Sunday) and this may reflect the non-negligible timescale required for pollutant removal by wind driven dispersion, i.e. the build-up of pollution during weekdays is not fully dispersed until Sunday. The same argument explains why visibility is typically higher on Mondays compared to the other weekdays later in the week.

- 5 The long term decadal (1950s–2010s) variation in visibility with RH is shown in Fig. 4, for all 8 stations, where the visibility is averaged within RH bins. A qualitatively similar pattern has been observed for all stations: Visibility is observed to vary strongly with relative humidity, which clearly indicates a significant particle hygroscopicity effect on visibility. It is noted that very high RH can also be indicative of precipitation which also decreases visibility.
- 10 To further highlight the effect of RH on visibility, the mean monthly visibility trend is compared to RH for the 60 years of data recorded at the Waddington station, see [supplementary Fig. S3](#). A scatter plot of visibility versus RH reveals a clear near-linear relationship (linear fit  $R^2 = 0.60$ ) between the variables. Removal of the long term trend in the visibility data was achieved by fitting the visibility to a quadratic function and subtracting the quadratic function from the time series. A scatter plot of the long term detrended visibility data versus RH
- 15 reveals a more linear relationship ( $R^2 = 0.66$ ) where every rise in RH of 10 % results in a reduction of approximately 5 km of visibility.

#### 4.4 Effect of long term changes in meteorological parameters upon visibility

- The long term trends in visibility are compared to the other recorded meteorological parameters: RH, air
- 20 temperature, wind speed and wind direction ([supplementary Fig. S4](#)). It is observed that at most of the stations RH decreases as average air temperature increases. Previous literature observed that the UK mean air temperature and sea surface temperature have increased by about 1°C and 0.7°C respectively between the early 1970s and mid 2000s (Jenkins, 2007). However, overall UK mean RH decreased about 2.7 % between the 1961 and 2006 (Jenkins, 2007). This reduction in RH is also seen more widely in the mid-latitudes (Willett et al.,
- 25 2014). The temperature change is likely due to climate change, land-use (urban heat island) effects or a combination of both. Clearly, urban heat island effects can only affect stations that are located in urban areas ([supplementary Fig. S4](#)). However, as [supplementary Fig. S4](#) shows, visibility is strongly related to relative humidity and hence to the air temperature of a given location, highlighting a possible indirect effect of climate change and urban heat island effects on regional visibility. [The correlation statistics between visibility, relative](#)
- 30 [humidity, air temperature and wind speed are provided for all stations in supplementary Fig. S5.](#)

#### 4.5 Mathematical fitting of measured visibility

- Equations 5 and 6 were fit to the decadal visibility data subset into distinct RH bins, as detailed in section 3.2. It is found that the fitted data is able to match the observed visibility extremely well ( $R^2 > 0.98$ ) for all stations; for
- 35 example see Fig. 5 for Heathrow station. The last decade, starting in 2010, has the poorest fit, albeit still with an  $R^2 = 0.95$ , but only comprises 3 years of data.

We have quantified, in section 4.1, that the decadal observed visibility has improved at most of the stations, which is a direct indicator of change in the combination of aerosol concentration, aerosol composition, gas concentration and RH. To better understand these changes in visibility, the absorption coefficient ( $\beta_{abs}$ ), scattering coefficient ( $\beta_{sca}$ ), particle hygroscopicity parameter ( $\gamma$ ), and dry visibility ( $Vis(dry)$ ) have all been calculated via constructed model (Eq. (4)) described in section 3.2.

The determined model output parameters ( $Vis(dry)$ ,  $\gamma$ ,  $\beta_{sca}$  and  $\beta_{abs}$ ) are presented in Fig. 6, where analysis has been carried out for all sites within each decade; however, the following discussion only considers data that was measured manually, due to the impacts of measurement methodology changes noted above. A clear improvement in calculated dry visibility was observed for Plymouth, Heathrow, Ringway, Nottingham and Waddington, while only minor changes were observed at Aldergrove, Leuchars and Tiree (Fig. 6a and supplementary Table S2). Broadly, the 5 sites in England are similar with all showing an upwards trend in visibility, whereas the Scottish and Northern Irish sites have greater dry visibilities but less discernible trend with time.

The derived value for  $\gamma$  has decreased slightly at Heathrow, Leuchars and Ringway sites over those decades (Fig. 6b and supplementary Table 2), which indicates a decrease in hygroscopicity over the time (and a concomitant improvement in visibility). Tiree is the only station which showed increased hygroscopicity parameter values, implying a rise in aerosol particle hygroscopicity which results in a drop in visibility. The other stations like Aldergrove, Ringway, Plymouth, and Waddington show very little change in hygroscopicity parameter values.

Reductions in scattering coefficient are observed at all sites except Aldergrove. The scattering coefficients calculated at RH = 75 % is shown in Fig. 6d. Larger decreases in the scattering coefficient are observed at the urban sites compared to the rural sites. Reductions are also observed in the absorption coefficient at most sites but there is much more variability compared to the scattering coefficient. It is interesting to note that the two most remote sites, both in Scotland, have increasing absorption coefficients, which is potentially indicative of episodes of long range transport of absorbing aerosol to these pristine sites becoming more frequent. As expected, both the absorption and scattering coefficients show an inverse relationship with the observed visibility (Fig. 6a and 6c).

The change in the fitted values for dry visibility and scattering coefficient are not significantly affected by the change in visibility measurement from human observation to visimeters. Contrastingly, the absorption coefficient and gamma values are much more influenced by measurement technique. This likely indicates that local sources have markedly different absorption and hygroscopicity parameters compared to more regional sources; whereas their local and regional scattering properties are relatively similar.

The modelled scattering coefficient, at 75 % RH, is always higher than the absorption coefficient for all sites and times. However, at lower RH the two values become more comparable, see supplementary Fig. S7 which examines the contribution of the scattering coefficient to the total extinction coefficient at Heathrow. The non-negligible contribution of the absorption coefficient to the total extinction coefficient indicates that the model

shown in Eq. (5) is not appropriate for the data reported in this paper. However, for other locations with lower concentrations of absorbing species, gas or aerosol, the model may be valid and the benefit of a linear fitting algorithm, compared to a non-linear algorithm, could be exploited. It is shown the contribution of aerosol scattering to total extinction has remained relatively constant over time which indicates that the reduction in particulate matter has decreased both the absorbing and scattering fractions in equal measure.

Seasonal decadal changes in aerosol parameters were calculated for the Heathrow station (supplementary Fig. S8). In general, an improved dry visibility with reduced  $\beta_{abs}$  and  $\gamma$  values was observed for all seasons over time. However, during winter months the greatest improvement in dry visibility with a reduction in  $\beta_{abs}$  was noted.

Trends in visibility for those data acquired at a single RH value of 70 % (67.5-72.5 %) during the period of 1950s to 1990s were investigated for the Heathrow site to demonstrate the disaggregation of the RH effect on visibility from aerosol concentration effect upon visibility. At constant RH, a clear improved visibility was determined for the study period (supplementary Fig. S9). The result implies significant changes in aerosol composition/concentration are driving the visibility trend. Hence improving air quality contributes significantly to better visibility.

#### 4.6 Effect of nitrogen dioxide gas upon visibility at Heathrow

The potential influence of NO<sub>2</sub> levels upon visibility was analysed using data from the Harlington station (proximate to the Heathrow site), for the period 2004 - 2012. The annual mean concentration of NO<sub>2</sub> varied from 33.6  $\mu\text{g m}^{-3}$  to 38.5  $\mu\text{g m}^{-3}$ , peaking in 2005 (Table 3). The NO<sub>2</sub> influence on observed visibility (in the RH bin centred at 75 % (72.5-77.5 %)) was greatest in the year of 2005 (where it contributed  $4.7 \pm 1.6$  % in total extinction) and lowest for 2012 ( $3.3 \pm 1.5$  % in total extinction) with the remaining visibility reduction being caused by aerosol extinction. Overall, during 2004 to 2012 NO<sub>2</sub> contributed approximately 4 % to the observed visibility change, while the remaining 96 % contributed arose from aerosol particles and fog. However it is worth considering the contribution of NO<sub>2</sub> towards the total extinction coefficient during the 1970s when visibility was very low (16.5 km) as compared to 2012 (25.24 km) and NO<sub>2</sub> levels higher. Unfortunately NO<sub>2</sub> data is not available before 2004 at nearby Heathrow site, but a recent study shows that, NO<sub>x</sub> emission in UK has almost doubled in the time period 1970 to 2012 (Harrison et al., 2015). Using the UK NO<sub>x</sub> record for 1970 from Harrison et al. (2015), we assumed the annual mean NO<sub>2</sub> concentration in 1970 is double what is measured in the year 2012 (34.6  $\mu\text{g m}^{-3}$ ) as emission estimates are approximately related to concentration. This assumption does not take into account the changing vehicle fleet with corresponding changing emissions of NO and NO<sub>2</sub> (Carslaw and Rhys-Tyler, 2013). Using this data the absorption coefficient for NO<sub>2</sub> was calculated. In particular, a higher absorption coefficient ( $\beta_{NO_2abs}$ ) in 1970 (0.0121  $\text{km}^{-1}$ ) as compared to 2012 (0.00507  $\text{km}^{-1}$ ) was identified. However, the contribution of NO<sub>2</sub> to the total extinction coefficient remained at 5.2 % in 1970, only about 2 % higher than in 2012.

## 4.7 Conclusions

Long term trends in visibility for 8 meteorological stations situated in the UK have been investigated. In general, visibility has improved at most of the stations through time. The improvements are greatest in urban areas, and are attributed to reductions in aerosol particle loadings and decreases in atmospheric RH. Visibility was found to be lowest during winter and highest in the summer due to seasonal variations in RH and likely changes in the mixing layer height. The rate of change of visibility was higher in winter for all stations, with the exception of Ringway. A sharp positive increment (5-12.5 %) in visibility was observed on Sundays, as compared to other days of the week (Mon-Sat), which is most likely due to weekend reductions in traffic and other particulate matter emission sources.

Bivariate polar plots of visibility, which account for both the influence of wind speed and wind direction, explained the influence of wind on likely source areas of visibility reducing aerosols. These bivariate polar plots identified likely locations for visibility reducing pollutants sources and their variation over time. Overall, an improved visibility at most of the stations in almost all directions was observed with notable improvements when the air masses moved over metropolitan areas, for example, Greater Manchester and Greater London Areas. At most sites, low visibility was observed when the winds came from the direction of continental Europe which may indicate an influence of regional pollution events leading to visibility reductions. Significant changes in visibility were observed with changes in relative humidity, which indicates a strong dependency of visibility on aerosol hygroscopicity. The measured RH at all sites was typically in the range of 60-80% and variations of a few percent in this RH range can have significant effects on visibility. Many sites showed long term decreases in RH which correlated with increases in air temperature, and had the effect of improving visibility. If the trend of increasing RH continues, the UK can expect further improvements in visibility for the same pollutant loading.

Calculations indicate that the majority of visibility reduction is caused by PM, however, a non-negligible contribution of light absorption is due to NO<sub>2</sub> gas. For the Heathrow station, over the time period 2004-2012, light absorption by NO<sub>2</sub> was calculated to contribute approximately 4% to the total visibility reduction, with the remainder caused by PM absorption and scattering. The NO<sub>2</sub> contribution was likely to have been significantly higher in prior decades due to the higher NO<sub>x</sub> emissions and hence atmospheric concentrations.

A light extinction model was developed to explain the dependency of visibility upon meteorology and aerosol characteristics. The agreement between the modelled and measured visibility is excellent. The model suggests that there have been significant changes in aerosol concentration over the last 60 years. The model incorporates parameterizations of aerosol hygroscopicity, particle concentration, particle scattering, and particle and gas absorption. The developed model is easily transferrable and applicable to other data sets worldwide.

Visibility can be used as a proxy for aspects of air quality, in particular particulate matter and nitrogen dioxide. Since visibility measurements can extend back for hundreds of years whilst air quality measurements typically only go back decades albeit with a few sparse datasets going back longer in time. The approach demonstrated in this paper has potential for generating historical air quality indications for locations with visibility records.



## *Acknowledgments*

We thank the University of Birmingham for supporting Ajit Singh through the Elite Scholarship Scheme. We are also thankful to UK Met Office and DEFRA for the provision of data used in this research. We also appreciate both reviewers for their valuable comments and suggestions, which has improved the quality of manuscript.

## **References**

- 10 Appel, B., Tokiwa, Y., Hsu, J., Kothny, E., and Hahn, E.: Visibility as related to atmospheric aerosol constituents, *Atmospheric Environment* (1967), 19, 1525-1534, 1985.
- Brimblecombe, P.: *The big smoke*, Methuen London; New York, 1987.
- Brimblecombe, P.: The clean air act after 50 years, *Weather*, 61, 311-314, 2006.
- Cao, J.-j., Wang, Q.-y., Chow, J. C., Watson, J. G., Tie, X.-x., Shen, Z.-x., Wang, P., and An, Z.-s.: Impacts of aerosol compositions on visibility impairment in Xi'an, China, *Atmospheric Environment*, 59, 559-566, 2012.
- 15 Carslaw, D. C., and Ropkins, K.: Openair—an R package for air quality data analysis, *Environmental Modelling & Software*, 27, 52-61, 2012.
- Carslaw, D. C., and Rhys-Tyler, G.: New insights from comprehensive on-road measurements of NO<sub>x</sub>, NO<sub>2</sub> and NH<sub>3</sub> from vehicle emission remote sensing in London, UK, *Atmospheric Environment*, 81, 339-347, 2013.
- 20 Chang, D., Song, Y., and Liu, B.: Visibility trends in six megacities in China 1973–2007, *Atmospheric Research*, 94, 161-167, 2009.
- Charron, A., Birmili, W., and Harrison, R. M.: Factors influencing new particle formation at the rural site, Harwell, United Kingdom, *Journal of Geophysical Research: Atmospheres*, 112, 2007a.
- Charron, A., Harrison, R. M., and Quincey, P.: What are the sources and conditions responsible for exceedences of the 24h PM<sub>10</sub> limit value (50µgm<sup>-3</sup>) at a heavily trafficked London site?, *Atmospheric Environment*, 41, 1960-1975, 2007b.
- Charron, A., Degrendele, C., Laongsri, B., and Harrison, R. M.: Receptor modelling of secondary and carbonaceous particulate matter at a southern UK site, *Atmospheric Chemistry and Physics*, 13, 1879-1894, 2013.
- 30 Clarke, A., Gascoigne, M., Henderson-Sellers, A., and Williams, A.: Modelling air pollution in Leeds (UK), *International Journal of Environmental Studies*, 12, 121-132, 1978.
- Colbeck, I., and Harrison, R. M.: Ozone—secondary aerosol—visibility relationships in North-West England, *Science of the Total Environment*, 34, 87-100, 1984.
- Colville, R., Hutchinson, E., Mindell, J., and Warren, R.: The transport sector as a source of air pollution, 35 *Atmospheric environment*, 35, 1537-1565, 2001.

- Crilley, L. R., Bloss, W. J., Yin, J., Beddows, D. C., Harrison, R. M., Allan, J. D., Young, D. E., Flynn, M., Williams, P., and Zotter, P.: Sources and contributions of wood smoke during winter in London: assessing local and regional influences, *Atmospheric Chemistry and Physics*, 15, 3149-3171, 2015.
- Defra: Air Pollution in the UK 2010, A report prepared by AEA for Defra and the Devolved Administrations  
 5 Department for Environment, Food and Rural Affairs, London, 2011.
- Diederer, H., Guicherit, R., and HolLonder, J.: Visibility reduction by air pollution in the Netherlands, *Atmospheric Environment (1967)*, 19, 377-383, 1985.
- Doyle, M., and Dorling, S.: Visibility trends in the UK 1950–1997, *Atmospheric Environment*, 36, 3161-3172, 2002.
- 10 Ferman, M. A., Wolff, G. T., and Kelly, N. A.: An assessment of the gaseous pollutants and meteorological conditions associated with Denver's brown cloud, *Journal of Environmental Science & Health Part A*, 16, 315-339, 1981.
- Founda, D., Kazadzis, S., Mihalopoulos, N., Gerasopoulos, E., and Lianou, M.: Long-term visibility variation in Athens (1931–2013): A proxy for local and regional atmospheric aerosol loads, *Atmos. Chem. Phys. Discuss.*,  
 15 doi:10.5194/acp-2015-1025, in review, 2016., 2016.
- Gomez, B., and Smith, C.: Atmospheric pollution and fog frequency in Oxford, 1926–1980, *Weather*, 39, 379-384, 1984.
- Gomez, B., and Smith, C.: Visibility at Oxford, 1926–1985, *Weather*, 42, 98-106, 1987.
- Groblicki, P. J., Wolff, G. T., and Countess, R. J.: Visibility-reducing species in the Denver “brown cloud”—I.  
 20 Relationships between extinction and chemical composition, *Atmospheric Environment (1967)*, 15, 2473-2484, 1981.
- Hänel, G.: Computation of the extinction of visible radiation by atmospheric aerosol particles as a function of the relative humidity, based upon measured properties, *Journal of Aerosol Science*, 3, 377-386, 1972.
- Harris, B., and Smith, K.: Cleaner air improves visibility in Glasgow, *Geography*, 67, 137-139, 1982.
- 25 Harrison, R. M., Jones, A. M., and Lawrence, R. G.: Major component composition of PM 10 and PM 2.5 from roadside and urban background sites, *Atmospheric Environment*, 38, 4531-4538, 2004.
- Harrison, R. M., Pope, F. D., and Shi, Z.: Trends in Local Air Quality 1970–2014, *Still Only One Earth: Progress in the 40 Years Since the First UN Conference on the Environment*, 2015, 58,
- Haywood, J., and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric  
 30 aerosols: A review, *Reviews of geophysics*, 38, 513-543, 2000.
- Holliday, J.: Winds of Change "An exhibition about the history of Tiree's weather station and early flights to the island", An Iodhlann Scarinish, Isle of Tiree, 2004.
- Husar, R. B., Holloway, J. M., Patterson, D. E., and Wilson, W. E.: Spatial and temporal pattern of eastern US haziness: a summary, *Atmospheric Environment (1967)*, 15, 1919-1928, 1981.
- 35 Jayamurugan, R., Kumaravel, B., Palanivelraja, S., and Chockalingam, M.: Influence of temperature, relative humidity and seasonal variability on ambient air quality in a coastal urban area, *International Journal of Atmospheric Sciences*, 2013, 2013.
- Jebson, S.: Fact sheet number 17: Weather observations, 2008.
- Jenkins, G. J.: The climate of the United Kingdom and recent trends, Exeter: Met Office Hadley Centre, 2007.

- Jinhuan, Q., and Liqun, Y.: Variation characteristics of atmospheric aerosol optical depths and visibility in North China during 1980–1994, *Atmospheric Environment*, 34, 603-609, 2000.
- Kong, S., Li, L., Li, X., Yin, Y., Chen, K., Liu, D., Yuan, L., Zhang, Y., Shan, Y., and Ji, Y.: The impacts of firework burning at the Chinese Spring Festival on air quality: insights of tracers, source evolution and aging processes, *Atmospheric Chemistry and Physics*, 15, 2167-2184, 2015.
- Koschmieder, H.: Theorie der horizontalen Sichtweite, *Beitr. Phys. Freien. Atmos.*, 12, 33-55, 1924.
- Lack, D., and Cappa, C.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, *Atmospheric Chemistry and Physics*, 10, 4207-4220, 2010.
- Lee, D.: Trends in summer visibility in London and southern England 1962–1979, *Atmospheric Environment* (1967), 17, 151-159, 1983.
- Lee, D.: A preliminary analysis of long-term visibility trends in central scotland, *Journal of climatology*, 5, 673-680, 1985.
- Lee, D.: Regional Variations in Longterm Visibility Trends in the UK, 1962—1990, *Geography*, 108-121, 1994.
- Lee, D. O.: The influence of wind direction, circulation type and air pollution emissions on summer visibility trends in southern England, *Atmospheric Environment. Part A. General Topics*, 24, 195-201, 1990.
- Lee, J. D., Lewis, A. C., Monks, P. S., Jacob, M., Hamilton, J. F., Hopkins, J. R., Watson, N. M., Saxton, J. E., Ennis, C., and Carpenter, L. J.: Ozone photochemistry and elevated isoprene during the UK heatwave of August 2003, *Atmospheric Environment*, 40, 7598-7613, 2006.
- Liu, X., Zhang, Y., Cheng, Y., Hu, M., and Han, T.: Aerosol hygroscopicity and its impact on atmospheric visibility and radiative forcing in Guangzhou during the 2006 PRIDE-PRD campaign, *Atmospheric environment*, 60, 59-67, 2012.
- Liu, Y., and Daum, P. H.: The effect of refractive index on size distributions and light scattering coefficients derived from optical particle counters, *Journal of Aerosol Science*, 31, 945-957, 2000.
- Malm, W. C., Walther, E. G., O'Dell, K., and Kleiner, M.: Visibility in the southwestern United States from summer 1978 to spring 1979, *Atmospheric Environment* (1967), 15, 2031-2042, 1981.
- NAPAP: Visibility Existing and Historical Conditions – Causes and Effects, , National Acid Precipitation Assessment Program, Report 24, 1990.
- Park, R. J., Jacob, D. J., Chin, M., and Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Park, R. J., Jacob, D. J., Kumar, N., and Yantosca, R. M.: Regional visibility statistics in the United States: Natural and transboundary pollution influences, and implications for the Regional Haze Rule, *Atmospheric Environment*, 40, 5405-5423, 2006.
- Sati, A. P., and Mohan, M.: Analysis of air pollution during a severe smog episode of November 2012 and the Diwali Festival over Delhi, India, *International Journal of Remote Sensing*, 35, 6940-6954, 2014.
- Schichtel, B. A., Husar, R. B., Falke, S. R., and Wilson, W. E.: Haze trends over the United States, 1980–1995, *Atmospheric Environment*, 35, 5205-5210, 2001.
- Seidel, D. J., and Birnbaum, A. N.: Effects of Independence Day fireworks on atmospheric concentrations of fine particulate matter in the United States, *Atmospheric Environment*, 115, 192-198, 2015.

- Singh, A., and Dey, S.: Influence of aerosol composition on visibility in megacity Delhi, *Atmospheric Environment*, 62, 367-373, 2012.
- Singh, A., Bloss, W. J., and Pope, F. D.: Remember, remember the 5th of November; gunpowder, particles and smog, *Weather*, 70, 320-324, 2015.
- 5 Sloane, C. S.: Summertime visibility declines: meteorological influences, *Atmospheric Environment* (1967), 17, 763-774, 1983.
- Tang, I. N.: Chemical and size effects of hygroscopic aerosols on light scattering coefficients, *Journal of Geophysical Research: Atmospheres*, 101, 19245-19250, 1996.
- Titos, G., Jefferson, A., Sheridan, P., Andrews, E., Lyamani, H., Alados-Arboledas, L., and Ogren, J.: Aerosol  
 10 light-scattering enhancement due to water uptake during the TCAP campaign, *Atmospheric Chemistry and Physics*, 14, 7031-7043, 2014.
- Tiwari, S., Payra, S., Mohan, M., Verma, S., and Bisht, D. S.: Visibility degradation during foggy period due to anthropogenic urban aerosol at Delhi, India, *Atmospheric Pollution Research*, 2, 116-120, 2011.
- Venkataraman, C., Reddy, C. K., Josson, S., and Reddy, M. S.: Aerosol size and chemical characteristics at  
 15 Mumbai, India, during the INDOEX-IFP (1999), *Atmospheric Environment*, 36, 1979-1991, 2002.
- Watson, J., and Chow, J.: Visibility and air pollution, *WIT Transactions on Ecology and the Environment*, 99, 2006.
- Wilkins, E.: Air pollution and the London fog of December, 1952, *Journal of the Royal Sanitary Institute*, 74, 1-15, 1954.
- 20 Willett, K., Dunn, R., Thorne, P., Bell, S., De Podesta, M., Parker, D., Jones, P., and Williams Jr, C.: HadISDH land surface multi-variable humidity and temperature record for climate monitoring, *Climate of the Past*, 10, 1983-2006, 2014.
- Williams, M.: Air pollution and policy—1952–2002, *Science of the total environment*, 334, 15-20, 2004.
- WMO: International Meteorological Vocabulary, World Meteorological Organization, WMO- NO. 182,  
 25 Geneva, 1992.
- WMO: Guide to Meteorological Instruments and Methods of Observation, 7th edition, World Meteorological Organization, WMO-No. 8, Geneva, 2008.
- WMO: Manual on the Global Observing System, Volume I – Global Aspects, World Meteorological Organization, WMO-No. 544, Geneva, 2015.
- 30 Wu, D., Tie, X., Li, C., Ying, Z., Lau, A. K.-H., Huang, J., Deng, X., and Bi, X.: An extremely low visibility event over the Guangzhou region: A case study, *Atmospheric Environment*, 39, 6568-6577, 2005.
- Yang, L.-x., Wang, D.-c., Cheng, S.-h., Wang, Z., Zhou, Y., Zhou, X.-h., and Wang, W.-x.: Influence of meteorological conditions and particulate matter on visual range impairment in Jinan, China, *Science of the Total Environment*, 383, 164-173, 2007.
- 35 Zhang, Y.-h., Shao, M., and Cheng, C.-l.: Quantitative relationship between visibility and mass concentration of PM<sub>2.5</sub> in Beijing, *Journal of Environmental Sciences*, 18, 475-481, 2006.

## Tables

**Table 1** Study stations with area and length of data description

No.	Station Name	Station code (src id)	Area	Period	Length of Data (in Year)
1	Aldergrove	1450	Urban (Airport)	1950-2012	63
2	Heathrow	708	Urban (Airport)	1950-2012	63
3	Ringway	1135	Urban (Airport)	1950-2004	55
4	Nottingham	556	Urban	1957-2012	56
5	Plymouth	1336	Urban (near coastal area)	1950-2012	63
6	Tiree	18974	Rural (Airport, near Coastal area)	1957-2012	56
7	Leuchars	235	Rural (RAF, near coastal area)	1957-2012	56
8	Waddington	384	Rural (RAF, Airport)	1950-2012	63

\* RAF stands for Royal Air Force

**Table 2** Rate of change of visibility (in  $\text{kmyear}^{-1}$ ) with their standard error at 95% confidence interval


Satiation	Year	Annual	Winter	Spring	Summer	Autumn
<b>Plymouth</b>	1950-2012	$0.040 \pm 0.021$	$0.152 \pm 0.017$	$0.006 \pm 0.025$	$-0.043 \pm 0.031$	$0.049 \pm 0.022$
<b>Aldergrove</b>	1950-2002	$0.056 \pm 0.021$	$0.110 \pm 0.019$	$0.831 \pm 0.030$	$-0.417 \pm 0.036$	$0.074 \pm 0.029$
<b>Heathrow</b>	1950-2011	$0.201 \pm 0.018$	$0.231 \pm 0.021$	$0.181 \pm 0.020$	$0.145 \pm 0.028$	$0.226 \pm 0.020$
<b>Ringway</b>	1950-2004	$0.339 \pm 0.016$	$0.331 \pm 0.020$	$0.363 \pm 0.018$	$0.316 \pm 0.025$	$0.343 \pm 0.018$
<b>Waddington</b>	1950-2012	$0.293 \pm 0.010$	$0.331 \pm 0.019$	$0.245 \pm 0.016$	$0.270 \pm 0.018$	$0.325 \pm 0.016$
<b>Leuchars</b>	1957-2012	$0.157 \pm 0.019$	$0.286 \pm 0.027$	$0.140 \pm 0.030$	$0.030 \pm 0.034$	$0.180 \pm 0.025$
<b>Tiree</b>	1957-2002	$-0.089 \pm 0.014$	$-0.186 \pm 0.014$	$-0.035 \pm 0.015$	$-0.098 \pm 0.015$	$-0.046 \pm 0.015$
<b>Nottingham</b>	1957-2012	$0.235 \pm 0.023$	$0.293 \pm 0.022$	$0.214 \pm 0.024$	$0.149 \pm 0.033$	$0.270 \pm 0.022$

**Table 3** Gases contribution in visibility change over Heathrow airport


<b>Year</b>	<b>NO<sub>2</sub> concentration (µg m<sup>-3</sup>)</b>	<b>NO<sub>2</sub> (ppm)</b>	<b>Total Extinction coefficient (km<sup>-1</sup>) by all effects (using E1)</b>	<b>Absorption coefficient in km<sup>-1</sup> (<math>\beta_{NO_2abs}</math>) by NO<sub>2</sub></b>	<b>% contribution of NO<sub>2</sub> in total extinction coefficient</b>
<b>2004</b>	38.3	0.0203	0.1475	0.00671 ± 0.0023	4.5 ± 1.5
<b>2005</b>	38.5	0.0204	0.1425	0.00675 ± 0.0023	4.7 ± 1.6
<b>2006</b>	36.9	0.0196	0.1978	0.00648 ± 0.0022	3.3 ± 1.1
<b>2007</b>	36.9	0.0197	0.1855	0.00649 ± 0.0029	3.5 ± 1.4
<b>2008</b>	34.7	0.0185	0.1759	0.00600 ± 0.0026	3.4 ± 1.4
<b>2009</b>	36.3	0.0193	0.1681	0.00636 ± 0.0023	3.8 ± 1.2
<b>2010</b>	34.4	0.0183	0.1755	0.00604 ± 0.0023	3.4 ± 1.3
<b>2011</b>	33.6	0.0179	0.1614	0.00589 ± 0.0025	3.6 ± 1.5
<b>2012</b>	34.6	0.0184	0.1550	0.00507 ± 0.0024	3.5 ± 1.5
<b>1970*</b>	69.2	0.0368	0.2370	0.0121	5.12

\*estimated values given for 1970 (see main text for details)

## Figures





**Figure 1** Geographical location of measurement stations used. Location point colours describe location type: red - urban airport; blue - urban; purple - rural/remote and green - rural airport. Also presented are mean wind rose statistics whole time period (approximately 60 years) for all eight stations.




**Figure 2** Historical trend of annual and seasonal visibility derived from daily (12 noon) observations by station: **a)** Aldergrove **b)** Heathrow, **c)** Leuchars, **d)** Nottingham, **e)** Plymouth, **f)** Ringway, **g)** Tiree, **h)** Waddington. **Shading** indicates changes in measurement methodology, where white is human observation, while blue and red are automated observation using different instruments. For further details see the supplementary Table S1










**Figure 3** Mean monthly visibility and RH (Left-side) and average weekday visibility normalized to Sunday mean values (Right-side) at all eight sites: **a)** Aldergrove **b)** Heathrow **c)** Leuchars **d)** Nottingham **e)** Plymouth **f)** Ringway **g)** Tiree **h)** Waddington.








**Figure 4** Decadal visibility at specific range of relative humidity (left side) and number of days in % during different relative humidity (right side)



**Figure 5** Comparisons of modelled and observed visibility at specific range of RH using Eq. (4) at Heathrow station. The observed visibility is presented with standard error bars at 95 % confidence interval.



**Figure 6** Model output parameters **a)** Dry visibility, **b)** gamma and **c)** absorption coefficient and **d)** scattering coefficient at 75 % n.b. from 1950s to 2010s. The green shaded region shows the start of visiometer era at most of the stations (see supplementary Table S1 to see the starting year of visiometer measurement).

\* See Supplementary Table S2 for model output parameter values including their uncertainties

*Supplement of*



**60 years of UK visibility measurements: impact of meteorology and atmospheric pollutants on visibility**



Ajit Singh, William J. Bloss and Francis D. Pope



Correspondence to: Francis D. Pope ([f.pope@bham.ac.uk](mailto:f.pope@bham.ac.uk))





# Supplementary Figures












**Figure S1** Frequency distribution of different visibility range for different sub periods (1950s to 2010s).



**Figure S2** Boxplot of decadal visibility at eight different study sites, where red dot denotes mean value of decadal visibility



**Figure S3** Mean monthly visibility at different RH conditions at Waddington



**Figure S4** Time-series of meteorological components relative humidity (RH), air temperature (T), wind speed ( $w_s$ ), and prevailing wind direction ( $w_d$ ) including visibility (V), where all variables are averaged at 12 noon. Shaded lines show smooth fit line at 95 % confidence interval.

	Visibility	RH	Temp	Wind speed
<b>Aldegrave</b>	1			
Visibility	1			
RH	-.519**	1		
Temp	.199**	-.373**	1	
Wind speed	.095**	-.028**	-.050**	1

	Visibility	RH	Temp	Wind speed
<b>Heathrow</b>	1			
Visibility	1			
RH	-.542**	1		
Temp	.322**	-.540**	1	
Wind speed	.261**	-.084**	-0.008	1

	Visibility	RH	Temp	Wind speed
<b>Leuchars</b>	1			
Visibility	1			
RH	-.688**	1		
Temp	.179**	-.353**	1	
Wind speed	.124**	-.208**	-.084**	1

	Visibility	RH	Temp	Wind speed
<b>Norringham</b>	1			
Visibility	1			
RH	-.583**	1		
Temp	.299**	-.511**	1	
Wind speed	.272**	-.072**	-.054**	1

	Visibility	RH	Temp	Wind speed
<b>Plymouth</b>	1			
Visibility	1			
RH	-.589**	1		
Temp	.185**	-.186**	1	
Wind speed	.220**	-.059**	-.073**	1

	Visibility	RH	Temp	Wind speed
<b>Ringway</b>	1			
Visibility	1			
RH	-.549**	1		
Temp	.342**	-.423**	1	
Wind speed	.269**	-.070**	-.018**	1

	Visibility	RH	Temp	Wind speed
<b>Tiree</b>	1			
Visibility	1			
RH	-.612**	1		
Temp	.041**	-.389**	1	
Wind speed	.331**	-.099**	-.226**	1

	Visibility	RH	Temp	Wind speed
<b>Waddington</b>	1			
Visibility	1			
RH	-.633**	1		
Temp	.340**	-.550**	1	
Wind speed	.232**	-.091**	-.015*	1

\* Statistically significant value (p < 0.05)

\*\* Statistically significant value (p < 0.01)


Temp- Air Temperature

RH- Relative Humidity

**Figure S5** Correlation coefficient (r) values between different variables, where daily data at 12 noon was used for calculation for last six decades




## Aldergrove




# Heathrow


## 1950s




## 1960s




## 1970s




## 1980s





## 1990s



## 2000s \*



## 2010s \*




Mean visibility (km)

Leuchars

1950s

1960s


1970s





1980s

1990s

2000s



2010s \*




# Nottingham

## 1950s

## 1960s


## 1970s





## 1980s

## 1990s


## 2000s \*



## 2010s \*



Mean visibility (km)




# Ringway

## 1950s

## 1960s


## 1970s



## 1980s


## 1990s

## 2000s \*





## 2010s \*

Missing data




Mean visibility (km)






**Figure S6** Decadal seasonal polar plots for all eight stations for 1950s, 1960s, 1970s, 1980s, 1990s, 2000s and 2010s (left to right). \* represents years where visiometer measured data is included.






**Figure S7** (a) Scattering coefficient ( $\beta_{sca}$ ), (b) total extinction coefficient ( $\beta_{ext}$ ) and (c) contribution of scattering coefficient in total extinction coefficient at Heathrow. Estimates of error are not included here to improve clarity.



**Figure S8** Model output parameters **a)** absorption coefficient ( $\beta_{abs}$ ), **b)** Gamma ( $\gamma$ ), and **c)** dry visibility at different seasons for Heathrow site.



**Figure S9** Decadal observed visibility at 70 % RH (range 67.5 -72.5 %) for Heathrow site. Error bars represent standard error at 95 % confidence interval.

## Supplementary Tables

**Table S1** Method of visibility measurement at different station with its used time period, where present indicates the sensor is still installed and being used.

	<b>Method/ Sensor/ Equipment Type Name with their working period</b>					
<b>Station Name</b>	<b>Manually</b>		<b>Sensor 1</b>		<b>Sensor 2</b>	
<b>Aldergrove</b>	01/01/1926	24/01/2003	<b>VISMETER -BELFORT 6230A</b> 24/01/2003   28/08/2012		<b>PRESENT WEATHER SENSOR - FD12P</b> 28/08/2012   Present	
<b>Heathrow</b>	01/01/1947	01/01/2000	<b>VISMETER -BELFORT 6230A</b> 01/01/2000   15/06/2005		<b>VISMETER -BELFORT (Replaced with new one)</b> 15/06/2005   Present	
<b>Ringway</b>	01/01/1941	01/01/2000	<b>VISIBILITY: VISIOMETER</b> 01/01/2000   01/11/2004		<b>Manually</b> 01/11/2004   Present	
<b>Nottingham</b>	01/01/1941	01/01/2000	<b>VISMETER -BELFORT 6230A</b> 01/01/2000   Present		-----	
<b>Plymouth</b>	01/01/1920	23/01/1997	<b>VISMETER -BELFORT 6230A</b> 23/01/1997   16/12/2010		<b>PRESENT WEATHER SENSOR - FD12P</b> 16/12/2010   Present	
<b>Tiree</b>	01/01/1926	16/12/2010	<b>VISMETER -BELFORT 6230A</b> 16/12/2010   present		-----	
<b>Leuchars</b>	01/01/1921	16/12/2010	<b>VISMETER -BELFORT 6230A</b> 16/12/2010   present		-----	
<b>Waddintgon</b>	01/01/1946	01/01/2000	<b>VISMETER -BELFORT 6230A</b> 01/01/2000   present		-----	

**Table S2** Model output parameters (*Vis(dry)*,  $\beta_{abs}$ , Gamma ( $\gamma$ ) and  $\beta_{sca}$ )

Station	Output parameters				
	Decade	<i>Vis(dry)</i> [m]	$\beta_{abs}$ [m <sup>-1</sup> ]	Gamma ( $\gamma$ )	$\beta_{sca}$ [m <sup>-1</sup> ]
Aldergrove	1950s	66168.94 ± 6406.58	1.91E-05 ± 2.07E-05	0.72 ± 0.13	0.000108 ± 0.000104
	1960s	67145.63 ± 6770.0	1.42E-05 ± 1.87E-05	0.78 ± 0.11	0.00013 ± 7.87E-05
	1970s	46450.98 ± 2542.37	5.54E-05 ± 1.25E-05	0.89 ± 0.11	9.93E-05 ± 5.27E-05
	1980s	60581.79 ± 3474.03	3.68E-05 ± 1.04E-05	0.91 ± 0.09	9.80E-05 ± 5.08E-05
	1990s	72645.1 ± 5414.83	4.30E-07 ± 1.77E-05	0.65 ± 0.09	0.000132 ± 4.29E-05
	2000s	33033.3 ± 2175.54	0 ± 6.14E-05	0.38 ± 0.11	0.0002 ± 2.86E-05
	2010s	27816.69 ± 1028.35	0.000134 ± 8.58E-06	1.05 ± 0.19	2.65E-05 ± 4.21E-05
Heathrow	1950s	30712.64 ± 1505.63	9.99E-05 ± 1.42E-05	1.28 ± 0.15	0.000104 ± 0.000161
	1960s	33724.2 ± 1690.38	8.65E-05 ± 1.32E-05	1.26 ± 0.13	7.87E-05 ± 0.000168
	1970s	29053.48 ± 1034.41	0.00011 ± 1.10E-05	1.26 ± 0.12	5.27E-05 ± 0.000127
	1980s	39380.87 ± 1684.78	6.63E-05 ± 1.23E-05	1.03 ± 0.11	5.08E-05 ± 0.000138
	1990s	47969.8 ± 1723.021	6.00E-05 ± 7.77E-06	1.17 ± 0.11	4.29E-05 ± 0.000109
	2000s	39486.93 ± 1182.31	7.14E-05 ± 8.96E-06	0.10 ± 0.10	2.86E-05 ± 0.000111
	2010s	29442.48 ± 782.86	0.000127 ± 5.86E-06	1.40 ± 0.18	4.21E-05 ± 4.22E-05
Ringway	1950s	19009.64 ± 970.33	0.000165 ± 2.34E-05	1.22 ± 0.15	0.000222 ± 0.000104
	1960s	24280.22 ± 1312.52	9.83E-05 ± 2.43E-05	1.10 ± 0.12	0.000287 ± 7.87E-05
	1970s	28983.71 ± 1337.76	8.77E-05 ± 1.69E-05	1.09 ± 0.11	0.000215 ± 5.27E-05
	1980s	38266.72 ± 1971.32	4.22E-05 ± 1.91E-05	0.89 ± 0.10	0.000208 ± 5.08E-05
	1990s	50337.53 ± 2318.8	4.59E-05 ± 1.03E-05	1.06 ± 0.10	0.000139 ± 4.29E-05
	2000s	50433.99 ± 2835.64	5.67E-05 ± 1.07E-05	1.18 ± 0.15	0.000107 ± 2.86E-05
	2010s	-----	-----	-----	-----
Nottingham	1950s	23730.96 ± 458.202	9.08E-05 ± 4.96E-05	1.10 ± 0.22	0.00034 ± 0.000104
	1960s	31058.79 ± 2110.9	7.17E-05 ± 2.07E-05	1.15 ± 0.11	0.000267 ± 7.87E-05
	1970s	35827.23 ± 1938.96	6.68E-05 ± 1.45E-05	1.11 ± 0.10	0.000197 ± 5.27E-05
	1980s	60996.37 ± 4725.79	7.06E-06 ± 1.72E-05	1.0 ± 0.10	0.000204 ± 5.08E-05

	1990s	39257.86 ± 1923.46	7.24E-05 ± 1.22E-05	1.10 ± 0.12	0.000127 ± 4.29E-05
	2000s	29994.36 ± 905.23	0.0001 ± 1.12E-05	0.80 ± 0.084	9.25E-05 ± 2.86E-05
	2010s	28622.94 ± 935.70	0.00013 ± 7.48E-06	1.21 ± 0.18	3.55E-05 ± 4.21E-05
<b>Plymouth</b>	1950s	37567.9 ± 2677.97	3.23E-05 ± 2.92E-05	0.73 ± 0.11	0.000197 ± 0.000104
	1960s	43782.5 ± 3244.77	3.92E-05 ± 1.89E-05	0.84 ± 0.10	0.000161 ± 7.87E-05
	1970s	46748.18 ± 2948.9	5.74E-05 ± 1.22E-05	1.06 ± 0.11	0.000114 ± 5.27E-05
	1980s	49226.96 ± 2903.94	4.95E-05 ± 1.20E-05	1.03 ± 0.11	0.000124 ± 5.08E-05
	1990s	38659.78 ± 1743.28	3.84E-05 ± 2.16E-05	0.56 ± 0.08	0.000137 ± 4.29E-05
	2000s	22827.89 ± 786.67	0.000115 ± 2.33E-05	0.53 ± 0.08	0.000118 ± 2.86E-05
	2010s	36199.28 ± 2242.56	5.56E-05 ± 2.63E-05	0.62 ± 0.12	0.000124 ± 4.21E-05
<b>Tiree</b>	1950s	91524.38 ± 22121.51	1.24E-05 ± 3.19E-05	0.83 ± 0.27	9.62E-05 ± 0.00010
	1960s	95940.8 ± 9045.804	1.43E-05 ± 1.10E-05	0.79 ± 0.10	7.92E-05 ± 7.87E-05
	1970s	108150.7 ± 11993.07	1.69E-06 ± 1.28E-05	0.74 ± 0.09	9.58E-05 ± 5.27E-05
	1980s	79744.05 ± 5324.328	3.21E-05 ± 7.33E-06	1.02 ± 0.10	6.96E-05 ± 5.08E-05
	1990s	72533.1 ± 4859.271	3.85E-05 ± 7.47E-06	1.10 ± 0.11	7.08E-05 ± 4.29E-05
	2000s	40831.35 ± 2358.797	6.26E-05 ± 1.53E-05	0.73 ± 0.09	9.12E-05 ± 2.86E-05
	2010s	35281.66 ± 2462.969	7.74E-05 ± 2.19E-05	0.71 ± 0.13	8.96E-05 ± 4.21E-05
<b>Leuchars</b>	1950s	68678.15 ± 8261.79	2.89E-05 ± 1.70E-05	1.22 ± 0.18	0.000151 ± 0.000104
	1960s	91554.03 ± 5428.33	2.86E-05 ± 5.05E-06	1.43 ± 0.09	0.000103 ± 7.87E-05
	1970s	77735.19 ± 4208.63	2.93E-05 ± 6.50E-06	1.21 ± 0.09	0.000112 ± 5.27E-05
	1980s	66272.27 ± 2493.00	4.19E-05 ± 5.22E-06	1.18 ± 0.08	8.82E-05 ± 5.08E-05
	1990s	78279.2 ± 3279.53	2.56E-05 ± 6.22E-06	1.02 ± 0.08	9.99E-05 ± 4.29E-05
	2000s	62485.8 ± 1901.10	4.97E-05 ± 4.21E-06	1.18 ± 0.08	6.64E-05 ± 2.86E-05
	2010s	56086.63 ± 2125.58	6.11E-05 ± 4.97E-06	1.34 ± 0.13	5.56E-05 ± 4.21E-05
	1950s	36683.17 ± 2690.569	6.47E-05 ± 1.93E-05	1.12 ± 0.13	0.000198 ± 0.000104
	1960s	37699.07 ± 2375.559	6.34E-05 ± 1.62E-05	1.05 ± 0.10	0.000172 ± 7.87E-05
	1970s	39651.07 ± 1685.186	7.30E-05 ± 9.43E-06	1.14 ± 0.09	0.000125 ± 5.27E-05

<b>Waddington</b>	1980s	49441.39 ± 2317.583	5.18E-05 ± 9.29E-06	1.13 ± 0.09	0.00013 ± 5.08E-05
	1990s	45305.49 ± 1503.447	6.99E-05 ± 6.24E-06	1.28 ± 0.10	9.71E-05 ± 4.29E-05
	2000s	46205.36 ± 1299.249	6.84E-05 ± 5.59E-0	1.08 ± 0.09	7.29E-05 ± 2.86E-0
	2010s	44260.81 ± 1370.792	8.24E-05 ± 4.57E-06	1.47 ± 0.15	4.60E-05 ± 4.21E-05