

Supplement to Technical Note: Conversion of Isoprene Hydroxy Hydroperoxides (ISOPOOH) on Metal Environmental Simulation Chamber Walls

5 Anne-Kathrin Bernhammer^{1,2}, Martin Breitenlechner^{1,*}, Frank N. Keutsch³, Armin Hansel^{1,2} and the CLOUD Team⁴

¹Institute for Ion and Applied Physics, University of Innsbruck, 6020 Innsbruck, Austria

²IONICON Analytik GmbH, 6020 Innsbruck, Austria

³John A. Paulson School of Engineering and Applied Sciences and Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA, USA

10 ⁴CERN, Geneva, Switzerland

* now at: John A. Paulson School of Engineering and Applied Sciences and Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA, USA

Correspondence to: Armin Hansel (Armin.Hansel@uibk.ac.at)

Table S1: Structures and names according to the MCM Leeds as used in this paper

MCM Nomenclature	Structure	MCM Nomenclature	Structure
C5H8 (isoprene)		HCHO (formaldehyde)	
MVK (methyl vinyl ketone)		MACR (methacrolein)	
MVKOO		MACROO	
ISOPBO		ISOPDO	
ISOPBO2		ISOPDO2	
ISOPBOOH (1,2-ISOPOOH)		ISOPDOOH (4,3-ISOPOOH)	
IEPOXB (<i>trans</i> - β -IEPOX)			