

Aerosol measurements during COPE: composition, size and sources of CCN and INPs at the interface between marine and terrestrial influences

J. W. Taylor¹, T. W. Choularton¹, A. M. Blyth², M. J. Flynn¹, P. I. Williams^{1,3}, G. Young¹, K. N. Bower¹, J. Crosier^{1,3}, M. W. Gallagher¹, J. R. Dorsey^{1,3}, Z. Liu¹, P. D. Rosenberg⁴

¹ Centre for Atmospheric Science, School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester, UK

² National Centre for Atmospheric Science, University of Leeds, Leeds, UK

³ National Centre for Atmospheric Science, University of Manchester, Manchester, UK

⁴ School of Earth and Environment, University of Leeds, Leeds, UK

Correspondence to: Jonathan W. Taylor (jonathan.taylor@manchester.ac.uk)

Abstract. Heavy rainfall from convective clouds can lead to devastating flash flooding, and observations of aerosols and clouds are required to improve cloud parameterisations used in precipitation forecasts. We present measurements of

boundary layer aerosol concentration, size and composition from a series of research flights performed over the southwest peninsula of the UK during the COPE experiment of summer 2013. We place emphasis on periods of southwesterly winds, which locally are most conducive to convective cloud formation, when marine air from the Atlantic reached the peninsula. Accumulation mode mass loadings were typically $2 - 3 \mu\text{g m}^{-3}$ (corrected to standard m^3 at 1013.25 hPa and 273.15 K), the majority of which was sulphuric acid over the sea, or ammonium sulphate inland, as terrestrial ammonia sources neutralised the aerosol. The cloud condensation nuclei (CCN) concentrations in these conditions were $\sim 150 - 280 \text{ cm}^{-3}$ at 0.1% and $400 - 500 \text{ cm}^{-3}$ at 0.9% supersaturation (SST), which are in good agreement with previous Atlantic measurements, and the cloud drop concentrations at cloud base ranged from $100 - 500 \text{ cm}^{-3}$. The concentration of CCN at 0.1% SST was well correlated with non-sea-salt sulphate, meaning marine sulphate formation was likely the main source of CCN. Marine organic aerosol (OA) had a similar mass spectrum to previous measurements of sea spray OA, and was poorly correlated with CCN.

In one case study that was significantly different to the rest, polluted anthropogenic emissions from the southern and central UK advected to the peninsula, with significant enhancements of OA, ammonium nitrate and sulphate, and black carbon. The CCN concentrations here were around six times higher than in the clean cases, and the cloud drop number concentrations were 3 – 4 times higher.

Sources of ice-nucleating particles (INPs) were assessed by comparing different parameterisations of the nucleation of ice used to predict INP concentrations, using measured aerosol concentrations as input. The parameterisations based on total aerosol produced INP concentrations ~~that~~ agreed within an order of magnitude with measured first ice concentrations at cloud temperatures as low as -12°C . Composition-specific parameterisations for mineral dust, fluorescent particles and sea

spray OA were 3 – 4 orders of magnitude lower, meaning either a source of INPs was present that was not characterised by our measurements, and/or one or more of the composition-specific parameterisations greatly underestimated INPs in this environment.

5 1 Introduction

Flash flooding occurs when rain is heavy and persistent. The southwest peninsula of the UK, (shown in Figs. 1 and 2) is particularly ~~vulnerable-prone~~ to flash flooding because convective clouds form along convergence lines that often persist for many hours, as was the case in the Boscastle flash flood of August 2004 (Golding et al., 2005). This type of event develops when the wind blows from the southwest and aligns with the peninsula. Convergence associated with sea breeze fronts from the north and south coasts generates quasi-stationary convective systems (Warren et al., 2014). In the Boscastle case, individual cells moved with the wind, but new cells were constantly generated at the upwind end, causing the system as a whole to remain in a quasi-stationary position. Each cell developed over a similar timeframe, and they therefore precipitated over the same region, causing highly localised flooding. The potential of convective clouds to generate persistent, heavy rainfall, and consequent flooding, along the peninsula demonstrates the importance of understanding cloud formation and development in the region.

As convection is initiated in the newly developing cells, boundary layer aerosols are lifted up to provide the cloud condensation nuclei (CCN) necessary for cloud drop formation. Increased concentrations of larger and more soluble aerosols enhance CCN numbers and, consequently, enhanced cloud drop number concentration (CDNC). This inhibits the formation of warm precipitation by reducing average drop size (Fan et al., 2007; Huang et al., 2008), which also affects riming efficiency in mixed-phase clouds (Klett and Davis, 1973). At ~~lower-sub-zero~~ temperatures, aerosols provide the ice nucleating particles (INPs) required for primary ice formation, which may then ~~initiate-lead to~~ secondary ice processes such as the Hallett-Mossop rime-splintering processes (e.g. Huang et al., 2008; Phillips et al., 2001). Interactions between the warm rain and secondary ice processes are thought to be key to determining the timing and location of cold precipitation (Taylor et al., 2016). Aerosols may influence these processes by affecting cloud and drizzle drop size, and by determining the concentrations of primary ice forming at different cloud temperatures. Koren et al. (2005) found that convective clouds forming in polluted air may have a higher cloud fraction, lower ice fraction and lower droplet radius than those forming in clean Atlantic air.

In the prevailing west/southwesterly winds, Atlantic air influences the southwest peninsula, whereas in northerly or easterly winds the region receives anthropogenic pollution from the UK and Western Europe. Local emissions from farming and transport may also influence the aerosol over the peninsula.

The COncvective Precipitation Experiment (COPE) (Leon et al., 2015) took place in the southwest peninsula during July-August 2013. Multiple aircraft made in-situ measurements of aerosol and clouds, as well as airborne remote sensing measurements, on days predicted to be favourable for convective cloud formation. Additional ground-based aerosol and X-band radar measurements were performed throughout the duration of the experiment. The main aim of COPE was to improve quantitative precipitation forecasting in numerical weather prediction (NWP) models. Clark et al. (2016) describe state-of-the-art convection-permitting models, which are used operationally for rainfall forecasting both in the UK and elsewhere. The development and application of these NWP models requires the use of parameterisations for processes that would be too small in scale or too costly computationally to explicitly evaluate on an operational basis. A parameterisation may be used, for example, to calculate the autoconversion rate of cloud water to rain, rather than explicitly evaluating processes such as collision-coalescence, condensation and evaporation over sub-grid timescales.

The development of improved parameterisations, and investigating the relative importance of different processes, involves the use of more detailed research models. For example, Connolly et al. (2009) described the aerosol-cloud-precipitation interaction model (ACPIM), one of several research models which investigators will use as part of COPE to study different cloud-aerosol interactions and microphysical processes. ACPIM requires information on aerosol composition and size distribution, as well as environmental variables, to initialise its explicit cloud microphysics scheme. The purpose of this paper is to describe the boundary layer aerosol measured during the aircraft campaign in terms of concentration, size, and composition, and to determine the main sources of CCN and INPs, in order to inform modelling studies aiming to improve forecasts of convective precipitation in the region. Additional measurements of cloud microphysics (e.g. Taylor et al., 2016) may be used to evaluate the importance of different cloud microphysical processes, while the X-band radar measurements provide a dataset to evaluate the accuracy of the distribution and rate of precipitation in a regional model.

2. Experimental

2.1 Aircraft measurements

Most of the data used in this analysis were measured by instrumentation mounted on the UK BAe-146-301 Atmospheric Research Aircraft during six case study flights over the southwest peninsula, which took place between 05 July and 03 August 2013. Each flight had a designated flight number, which are listed in Table 1. Additional airborne measurements were performed with the University of Wyoming King Air 200T (UWKA) and UK Met Office Civil Contingency 208 Aircraft (MOCCA), but these are not discussed here. The BAe-146 was fitted with a variety of aerosol and cloud probes. Here we only discuss instruments relevant to our analysis; Leon et al (2015) provide a full list of instrumentation aboard the aircraft. All concentrations are corrected to standard temperature (273.15 K) and pressure (1013.25 hPa). A typical flight plan involved performing aerosol runs in the boundary layer ~~at an altitude of ~500m~~, before making measurements of clouds at higher levels. The flight plans of the boundary-layer aerosol runs in each flight are shown in Figure 2, and the altitude and

duration of the aerosol runs are listed in Table 1. Vertical profiles of potential temperature showed the boundary layer height was ~ 750 – 1250 m above mean sea level (AMSL; unless otherwise stated, all altitudes henceforth are AMSL), meaning these aerosol runs were conducted in the boundary layer.

The concentration of condensation nuclei (CN, i.e. the aerosol number concentration) larger than 2.5 nm was measured using a model 3786-LP water-filled condensation particle counter (WCPC; Hering et al., 2005) manufactured by Aerosol Dynamics Inc., based on the TSI model 3786 but modified for use at low pressure. The absolute uncertainty in accuracy of the CN concentration measurement is $\pm 12\%$.

A Droplet Measurement Technologies (DMT) dual-column cloud condensation nuclei counter (CCNC; Roberts and Nenes, 2005) measured the concentrations of CCN at nominal supersaturations (SST) of 0.1% and 0.9%, though on 18 July the higher SST was set to 0.6%. The relative uncertainty in the supersaturation measurements is $\pm 10\%$ (Roberts et al., 2010; Trembath, 2013), and the uncertainty associated with the flow calibration and counting efficiency of the particle counter is typically $\sim 640\%$ (Trembath, 2013, p189). The CCNC sampled behind a pressure-controlled inlet set to 650 hPa.

The aerosol size distribution was measured with a custom-built inboard scanning mobility particle sizer (SMPS; Wang and Flagan, 1990), and wing-mounted Particle Measurement Systems passive cavity aerosol spectrometer probe (PCASP) - 100X, with electronics upgraded by DMT, and DMT cloud droplet probe (CDP). The PCASP and CDP were size calibrated as described by Rosenberg et al. (2012). The SMPS measures the mobility diameter of particles 20 – 320 nm, while the PCASP and CDP measure the optical diameter of particles 0.12 – 3.4 μm and 2 – 50 μm respectively. The PCASP inlet has active drying, and the SMPS inlet line was dried to $<45\%$ RH by ram heating and the temperature increase as the sample line entered the cabin. The CDP measures aerosols and cloud drops at ambient humidity.

The composition of nonrefractory submicron aerosol was measured with an Aerodyne Research Inc. compact time-of-flight aerosol mass spectrometer (AMS; Canagaratna et al., 2007; Drewnick et al., 2005), which reports concentrations of organic aerosol (OA), ~~non-sea salt~~ sulphate (~~nss~~- SO_4^{2-}), nitrate (NO_3^-) and ammonium (NH_4^+) ~~and non-sea salt chloride (nss-Chl⁻)~~ as standard. Additionally, we used the fragmentation table given by Langley et al (2010) to calculate concentrations of methanesulphonic acid (MSA), and adjusted the fragmentation tables of OA and ~~nss~~- SO_4 accordingly. The AMS can also be used to report chloride (Chl⁻) concentrations, which would be likely to be influenced by sea salt in a marine environment. Previous studies have attempted to quantify sea salt using the AMS, but the scaling factor used to correct the data for the instrument's poor detection efficiency of sea salt was highly variable (Nuaaman et al., 2015; Ovadnevaite et al., 2012; Schmale et al., 2013; Zorn, 2009). As no calibrations for sea salt were carried out in the field, we are unable to estimate the scaling factor, and thus the Chl⁻ concentration, with any reasonable accuracy, and we therefore do not report Chl⁻ concentrations in this analysis. The SO_4^{2-} measurement may have a contribution from sea-salt sulphate, but this is likely to be of the order of 1% (Schmale et al., 2013).

The AMS was calibrated at the start and end of each flight using monodisperse ammonium nitrate to calculate the ionisation efficiency (IE) for NO_3^- , corrected as described by Morgan et al. (2014, Supplementary material). The ammonium nitrate

calibrations were also used to calculate the relative ionisation efficiency (RIE) of NH_4^+ , and a separate ammonium sulphate calibration was performed to calculate the RIE of SO_4^{2-} . The AMS IE calibration was stable within $\pm 12\%$ throughout the campaign, and the absolute accuracy is $\sim 30\%$ (Bahreini et al., 2009). The collection efficiency was calculated as recommended by Middlebrook et al. (2012), and the ~~calculated~~ aerosol volume, calculated with volume mixing using organic and inorganic densities of 1.27 g cm^{-3} and 1.77 g cm^{-3} respectively (Cross et al., 2007), showed qualitative agreement with the aerosol volume measured by the SMPS and PCASP.

Refractory black carbon (BC) concentrations were measured with a DMT single-particle soot photometer (SP2; Schwarz et al., 2010), which is the same instrument used in previous studies on the BAe-146 (e.g. Taylor et al., 2014), but upgraded to be functionally identical to the current SP2 model D. SP2 calibrations were carried out at the start of each flight, and in the laboratory after the campaign. The incandescence detectors, which measure BC mass on a single-particle basis, were calibrated using 300 nm mobility diameter Aquadag, corrected for the difference in response between the calibration particles and ambient soot (Baumgardner et al., 2012; Laborde et al., 2012b). Using this calibration, the uncertainty in the reported BC mass concentrations are $\pm 10\%$ when sampling diesel soot, and $\sim 20\%$ when sampling biomass burning (Laborde et al., 2012a, 2012b).

The online inboard aerosol instrumentation (i.e. the AMS, SP2, SMPS and CCNC) sampled using Rosemount inlets, which have sample efficiencies of approximately unity for particles smaller than 600 nm in diameter when sampling marine aerosol (Trembath et al., 2012).

Bulk aerosol was collected on two 47-mm Nuclepore filters, with $10 \mu\text{m}$ and $1 \mu\text{m}$ pore sizes, connected in serial. The filter sampling system is described by (Formenti et al., 2008). The size and composition of particles collected on the filters was determined offline after the campaign using scanning electron microscopy (Phillips FEI XL30 Environmental Scanning Electron Microscope with Field-Emission Gun (ESEM-FEG), with an Energy-Dispersive X-Ray Spectroscopy (EDX) system (Hand et al., 2010)). Particles were classified by composition using the scheme described by Young et al. (2016). An example filter composition measurement from the COPE campaign is presented by Leon et al. (2015). In our analysis, we only use the filter measurements to estimate the concentration of mineral dust for use in INP parameterisations.

The CO concentration was measured using an AeroLaser VUV fluorescence monitor model AL5002 (Gerbig et al., 1999), and the 3-D wind vector was measured with a de-iced Aventech aircraft-integrated meteorological measurement system (AIMMS)-20 turbulence probe (Beswick et al., 2008). A Stratton Park Engineering Company SPEC 2DS stereo probe was used to derive ice concentration, using the data processing rules described by Taylor et al. (2016).

2.2 Ground site measurements

The COPE ground site (located at 50.6466°N , 4.63323°W) hosted a suite of aerosol and meteorological instrumentation operating continuously throughout July and August 2013. Leon et al. (2015) provide a full list of instrumentation at the ground site. In this analysis, we only use the size-resolved measurements of fluorescent aerosol made by a ~~DMT~~ University

of Hertfordshire wideband integrated bioaerosol sensor (WIBS) 4M, which we consider to be primary biological aerosol particles (PBAP). Although the WIBS may also detect other fluorescent particles, the fluorescent particle concentration is generally considered to be a lower limit on the PBAP concentration (Huffman et al., 2010).

3. Results and discussion

3.1 Airmass history

Figure 1 shows 5-day back trajectories for each of the case study flights, calculated using the Hybrid Single Particle Lagrangian Integrated Trajectory Model (HYSPLIT) (Draxler and Hess, 1998). Trajectories were initiated every 60 seconds from the aircraft flight track during the boundary layer aerosol runs. The trajectories were performed with 3-hourly 0.5° Global Data Assimilation System (GDAS) reanalysis data, using full vertical dynamics on 23 model levels. Two additional trajectories were released from the ground site (at an altitude of 10m above ground level) on 03 August at 1100 and 1200UTC, the same time period as the aerosol runs, as the ground site measurements are used on this date in Sect. 3.5. These two ground-site trajectories follow a similar path to the ones released from the aircraft flight track on 03 August, meaning the aircraft and ground site were sampling from the same airmass on this date. Although the turbulent mixing in the boundary layer means the accuracy of individual trajectories is uncertain, examining the general trend provides information on the history of the airmass, and possible changes in its cloud nucleating potential, which is determined by synoptic-scale winds.

Figure 2 shows the flight path the aircraft took during the below-cloud runs, and the boundary layer wind speed and direction measured during these runs. The wind data allow us to assess the accuracy and potential influence of the back trajectories' final stages, and examine local wind fields such as sea breeze circulations that are not captured by the meteorology data used to initiate the trajectories.

Figures 1 and 2 panels (c), (e) and (f) show that the 25 and 29 July and 03 August case studies shared the synoptic features of the "Boscastle" type meteorology, i.e. winds blowing marine air from the North Atlantic over the peninsula from the southwest (hereafter referred to as southwesterly winds (SWW)) (Golding, 2005). The back trajectories showed that the airmasses had been over the North Atlantic for at least five days prior to landfall. They also showed sharp changes in direction over the ocean; comparison with ~~the~~ UK Met Office synoptic charts (available at http://www1.wetter3.de/error_adblocker_archiv_ukmet.html) showed these were due to circulation around low-pressure systems as they moved east, and were associated with frontal precipitation, ~~which~~ This precipitation is likely to have washed out ~~any~~ the majority of any long-range transported aerosol as the trajectories show the airmass remained in the lower troposphere, below the likely source of frontal precipitation. On 28 July, the situation was reasonably similar in that the airmass had spent time over the North Atlantic, but reached the peninsula from the south, and may have passed over the

northwest tips of Spain and France. Local sea breeze circulations may still have caused convergence on these four days, but they had only a minor influence on the absolute wind speed and direction, as the synoptic-scale winds were so strong.

The meteorology on 05 and 18 July were markedly different to the other case studies. On these two days, high pressure systems located over the UK meant that synoptic-scale boundary layer winds were much weaker. The measured winds were generally only a few metres per second and the greater variation in wind direction suggest that the local sea-breeze circulations had more of an influence on absolute wind direction on these days. On 05 July, the back trajectories show that for the few days prior to being sampled, the airmass had passed over the Atlantic, though the easternmost trajectories passed over the southern tip of Ireland and the south coast of Wales. It is difficult to assess the accuracy of these parts of the trajectories when local winds were dominant, though there was some anthropogenic influence on this flight, as discussed in the next section. The trajectories for 18 July suggest the air over the peninsula had passed over many parts of England, Wales and Ireland, and are therefore likely to have also had a strong anthropogenic influence.

The colours in Fig. 2 divide sections along the flight track into “Northwest (NW) coast”, “Inland” and “Other” influenced regions. The “NW Coast” regions include areas over the sea, and the regions labelled “Other” are not used in this analysis. Depending on the wind directions, aerosol in these regions may be influenced by marine or terrestrial emissions, or both. The trajectories and wind data in Figs. 1 and 2 show that for flights on 05, 25 and 29 July, and 03 August, the sections on the NW coast were affected by marine air that had not been influenced by land, though this was not the case for the other two flights. Dividing the data into these sections allows us to investigate changes in aerosol chemistry and properties at the land-sea interface. This topic is discussed further in the next section.

3.2 Submicron aerosol composition

3.2.1 Marine aerosol

Figure 3 shows the average aerosol composition measured by the AMS and SP2 in the NW coastal regions during the six case study flights. The data from Figure 3 are also listed in Table 1. In clean Atlantic air (NW coast on 05, 25 and 29 July and 03 August, as identified in the previous section) the composition was fairly consistent. Around 60 – 70% of submicron aerosol mass was sulphate, in the form of a mixture of sulphuric acid and ammonium bisulphate as there was insufficient ammonium for full neutralisation. This is in agreement with measurements of North Atlantic aerosol made by Ovadnevaite et al. (2014). In the marine airmasses, the ratio of $\text{MSA}/\text{H}_2\text{SO}_4$ was ~ 0.05 , which is perhaps lower than average, but within the range typically observed in clean marine air in summer (e.g. Dall’Osto et al., 2010; Huebert et al., 1996; Ovadnevaite et al., 2014; Phinney et al., 2006). MSA and sulphate were well correlated ($R^2 = 0.66$), and the source of both is likely to be oxidation of biogenic dimethyl sulphide (DMS) emissions. OA was the only other major aerosol component, though the ratio of $\text{H}_2\text{SO}_4/\text{OA}$ did show some variation. It is unlikely that anthropogenic pollution, either from shipping or long-range transported, made any significant contribution to the aerosol loadings; in clean Atlantic air, the OA showed no correlation with BC ($R^2 = 0.0$), while H_2SO_4 showed a weak negative correlation ($R^2 = 0.15$). These results are in agreement with a

similar assessment of Atlantic aerosol made at Mace Head, Ireland (O'Dowd et al., 2014). The source of the marine OA is discussed further in Sect. 3.2.3.

The total mass loading measured by the AMS + SP2 varied from $2.4 - 3.6 \mu\text{g m}^{-3}$ in these flights. ~~Non-sea salt chloride and black carbon were~~ as only a minor components, ~~and low~~, but non-zero levels of nitrate were measured, despite the fact that aerosol nitrate is not expected to form in acidic conditions as the sulphate displaces any nitrate to the gas phase. Interpretation of this apparent measured nitrate is discussed in Sect. 3.2.4.

3.2.2 Inland aerosol

Submicron inland aerosol composition on flying days is shown in Figure 4. On flying days between 25 – 29 July and 03 August, inland aerosol had composition similar to the Atlantic marine air, but with the addition of ammonium. On 25 and 28 July, the ammonium was sufficient to fully neutralise the sulphate, but on 29 July and 03 August, the sulphate was a mixture of ammonium sulphate and bisulphate. The total measured mass loadings were within 10% of the marine air for the corresponding flight.

Agriculture is the largest land use category in the southwest peninsula (Morton et al., 2011), including a mix of both arable and pastoral farming, predominantly cattle grazing on managed grass fields. Previous measurements have found elevated levels of ammonia in the southwest peninsula compared to those found in the Atlantic (Quinn et al., 1996; Sutton et al., 2001). The terrestrial ammonia emissions are therefore most likely to be of agricultural origin.

Compared to the last four flights, the inland aerosol on 05 July had a much larger OA fraction. The low wind speeds on this day mean local aerosol sources are likely to have had more of an effect. Inland on 05 July, OA had a correlation with BC of $R^2 = 0.56$, and with the aircraft's CO measurement an $R^2 = 0.69$. Additionally, the BC mass loading was around a factor of 10 higher inland than in the Atlantic air. Taken together, this suggests that local fossil fuel emissions were at least partly responsible for the increased OA inland on 05 July. Some of the easternmost trajectories passed over the southern tips of Wales and Ireland, which are an alternative source of the increased anthropogenic influence inland on 05 July.

18 July was somewhat of an outlier in terms of the COPE case studies in that the aerosol mass loadings were much higher than the other flying days. The back trajectories showed that the air mass had come from England, Wales and/or Ireland, so a strong anthropogenic influence would be expected. in contrast to the clean marine air from the southwest seen in the other cases. There was not much difference between the aerosol measured inland and on the NW coast, as the air over both areas had come from the East, bringing anthropogenic pollution to the region. OA was the largest component, followed by similar amounts of ammonium nitrate and sulphate, which were both fully neutralised by ammonium.

3.2.3 OA composition and origin

Examining the ratios of different fragments in the OA mass spectrum can give an indication of OA sources as well as their level of ageing. AMS factor analysis most-commonly tends to pick out factors with differing levels of oxidation. Fresh fossil

fuel emissions appear as hydrocarbon-like OA (HOA), while more aged aerosol including secondary OA (SOA) appears as oxygenated OA (OOA), which may be semi-volatile (SV-OOA) or more aged low-volatility (LV-OOA) (Jimenez et al., 2009).

OA measured in marine background air may be primary OA from sea spray (Ovadnevaite et al., 2011), SOA from marine VOC emissions (Decesari et al., 2011), or long-range transported pollution. Globally, primary marine OA is thought to dominate over marine SOA formation (Arnold et al., 2009; Gantt and Meskhidze, 2013). As discussed in Sect. 3.2.2, local anthropogenic emissions contribute to OA concentrations inland, while the marine OA would have less of an influence in areas further away from the sea.

The OA mass spectrum measured over the NW coast on 05 July is shown in Fig. 5. The spectrum is dominated by peaks at m/z 28 and 44 from CO^+ and CO_2^+ , which are prescribed to be equal in the default fragmentation table. These peaks are much larger than m/z 43 as is typical in oxidised OA spectra, these peaks are much larger than m/z 43 from C_3H_7^+ or $\text{H}_3\text{C}_2\text{O}^+$, as the CO^+ and CO_2^+ ions are more likely to be formed by fragmentation of more oxidised organic molecules. Table 2 presents correlations between NW coast OA spectra from 05, 25 and 29 July, and 03 August, with several spectra from literature. The literature spectra are HOA and two OOA factors (Morgan et al., 2010), sea spray (Ovadnevaite et al., 2011), marine biogenic SOA (Chang et al., 2011, which was well correlated with ~~non~~-SOA), and the “Organic factor” measured in the Arctic ocean by Chang et al. (2011), which was thought most likely to be sea spray OA and was poorly correlated with sulphate. Though there are many previous measurements of different OA components in Europe, we choose those from Morgan et al (2010) because they were measured with the same instrument aboard the BAe-146, so any inter-instrument variability should be minimised. OOA-1 was more oxidised than OOA-2, but no assessment was made of their volatility. To ensure the comparison was fair, only m/z values where all spectra had data points were used. The marine OA spectra showed the best correlation with sea spray OA, indicating this is a likely source, though the correlations with the other oxidised factors were still reasonably good.

Figure 6 shows the fractions f_{44} and f_{43} (the fraction of OA measured at m/z 44 and 43 respectively) for all six case study flights, divided up into NW coastal and inland data. The literature values from Table 2 are also plotted. Ambient OOA measurements tend to fall within the diagonal dashed lines in Fig. 6 (Ng et al., 2010), and fossil fuel emissions may be expected to fall close to the HOA point from Morgan et al. (2010) upon emission and progress approximately through OOA-2 to OOA-1 with increased oxidation.

The four marine spectra from this study have f_{44} indicative of a moderate level of oxidation but, on average, slightly low f_{43} . Of the selected literature values, the marine OA all fall closest to the sea spray OA, rather than the marine biogenic. If the inland OA was composed of a mix of marine and terrestrial OA, lines drawn from the marine data points through their corresponding inland points should intersect in a region representative of terrestrial OA. For the four flights with clean marine spectra, the lines would intersect in a region around (0.1, 0.1), close to the SV-OOA data point from Morgan et al (2010). This is consistent with the terrestrial aerosol source being similar on each flying day, and being moderately oxidised.

It is not fully clear what fraction of the terrestrial OA is anthropogenic or biogenic. It is difficult to pick out anthropogenic locally produced SOA from biogenic as the mass spectra look similar, and photochemical production is strongest for both during peak daylight hours. Most AMS factor analysis studies are only able to pick out regional background and locally produced OOA (Jimenez et al., 2009 and references therein). Slowik et al. (2010) identified a biogenic SOA factor by correlation with biogenic VOCs, but noted the mass spectrum was very similar to SV-OOA formed from anthropogenic pollution. As discussed in Sect. 3.2.2, the correlation between OA and CO is indicative of a fossil fuel component of the terrestrial OA, but we also cannot rule out a biogenic component.

3.2.4 Interpretation of nitrate measurements

The formation of nitrate aerosol is only possible when there is sufficient ammonium (or other cations) to fully neutralise any sulphate aerosol (Seinfeld and Pandis, 1998). However, our observations in Figure 3 and 4 show the apparent presence of aerosol nitrate under acidic conditions. The measurements are above the detection limit for nitrate ($\sim 0.03 \mu\text{g}/\text{sm}^3$ for a 30s averaging time), and are ubiquitous throughout the runs, meaning they are not an artefact of averaging two air masses of differing composition.

Nitrate appears in the AMS almost entirely in peaks at $m/z = 30$ and 46 (Allan et al., 2003), for ions NO^+ and NO_2^+ respectively. When sampling nitrate species, the ratio of these two peaks is determined by the heater temperature and aerosol volatility. For a fixed heater temperature, less volatile nitrate species decompose further during their slower vaporisation, and have a higher 30/46 ratio when measured (Drewnick et al., 2015). The AMS may detect other nitrate species including salts such as NaNO_3 and KNO_3 , as well as organic nitrates, as long as their boiling point is low enough.

Table 3 shows the ratios of m/z 36/40 measured on each flying day during ammonium nitrate calibrations and ambient sampling periods. For 18 July, the 30/46 ratio was the same during the calibration and coastal/inland aerosol. As the ammonium in these periods was sufficient to fully neutralise the aerosol, this gives us a high degree of confidence that the NO_3 measured on this flight was ammonium nitrate. However, on all other flights the ambient 30/46 ratio was higher than the calibration values, particularly over the sea. Sea salt-based nitrates may be found in marine air, such as NaNO_3 formed by reaction of nitric acid with NaCl . This would be found predominantly in coarse mode aerosol rather than accumulation mode (Harrison and Pio, 1983), but some particles may be detected by the AMS. Alfarra (2004) measured nebulised NaNO_3 and recorded a 30/46 ratio of 29.2, which is significantly higher than those measured during COPE. Presumably, the same argument applies with these nitrate salts as with ammonium nitrate: the nitrate would be displaced by sulphate unless the aerosol was fully neutralised. NaNO_3 would only be possible if the coarse mode aerosol was not in equilibrium with the accumulation mode. We must therefore investigate the possibility of organic compounds giving signals at the same m/z as the nitrate peaks.

Ovadnevaite et al. (2011) measured high-resolution mass spectra of primary marine OA, which allowed them to identify the specific ions measured at particular m/z values. Around 1.7% of primary marine OA was measured at m/z 30, most of which

was CH₂O, and 0.2% was at m/z 46. If the measured nitrate concentrations in COPE were attributable to organic species, they would make up 6 – 12% of organic mass in marine air when corrected for relative ionization efficiency. Rollins et al. (2010) measured 30/46 ratios of 0.99 – 5.30 for various organonitrates, which is more in line with our measurements.

To summarise, we are not able to conclusively explain the reported nitrate measurements in acidic conditions. Several past studies have faced similar problems in various different regions e.g. (Allan et al., 2004, 2006, 2014). It is likely that they are partly due to interference from organic species, but the inland 36/40 ratios, where enhanced ammonium concentrations were measured, were closer to those expected from NaNO₃. The measurements would need to be performed with a high-resolution mass spectrometer to confirm the organic fraction at the nitrate peaks. There may also be a contribution from sea salt-based nitrate species such as NaNO₃, but these are also difficult to quantify with our measurements. On all flights other than 18 July, the total nitrate loadings were a relatively small fraction of the total mass measured by the AMS, as the ~~non~~-SO₄ and OA loadings were both several times higher.

3.3 Aerosol size distributions

Figure 7 shows the submicron aerosol size distributions measured in NW coastal regions during the six case study flights.

The standard deviations ~~of each in~~ are plotted to show the spatial variation over the runs, and were much larger than the calculated standard errors for the size range shown. The SMPS data were normalised to the PCASP in the overlap region. Three submicron aerosol modes are evident from the distributions: nucleation, Aitken, and accumulation modes, with varying magnitudes in the different flights. Coarse-mode aerosols were also measured, but the supermicron distribution is not shown in Fig. 7 as the bin sizing errors and overlap between the PCASP and CDP made the features difficult to distinguish. The mean diameters of the Aitken, accumulation and coarse modes are listed in Table 4. The mean diameter of the nucleation modes was likely to be smaller than the SMPS can reliably measure.

Many of all the measured distributions did not drop to near zero in the lowest bin ~~were open at the lower end, meaning suggesting~~ nucleation mode aerosols were present smaller than the 20nm size of the smallest bin. In all flights, the concentrations of particles in all bins <40 nm in diameter was higher inland than over the sea. Additionally, the total CN concentration (listed in Table 1, and shown in Fig. 8a) was always higher and more variable inland, and the standard deviations of smallest particles measured were higher inland. These measurements suggest there was a source of small particles forming inland, which was not uniform in spatial location.

On most flights, the Aitken mode was centred on ~~340~~ – 60 nm, and the accumulation mode was centred around ~~1150~~ – ~~22190~~ nm. The gap in between was characteristic of the “Hoppel dip”, thought to be caused by cloud processing of marine aerosol (Hoppel et al., 1986). Similar size distributions have been measured in summertime North Atlantic aerosol (Asmi et al., 2011; Dall’Osto et al., 2011). The accumulation mode on 05 July was larger, with modal diameter ~275 nm over the sea, which may be due to increased photochemical aerosol formation in the clear weather. 18 July was also an exception, showing just one wide accumulation mode, and the size distribution was consistent with aged UK urban aerosol (e.g.

Formatted: Don't keep with next

Bohnenstengel et al., 2015). The features of the marine aerosol size distribution seen on other days, such as a prominent Aiken mode and Hoppel dip, were also absent on this day due to the lack of a marine influence. The number concentration of aerosols in the accumulation mode on 18 July was significantly higher than on any of the other days studied, due to the anthropogenic pollution influence. The concentration of particles in the inland nucleation mode on 05 July was also higher than on any of the other flights, and this mode was broader than the other flights, which may be due to the stagnant winds allowing photochemical processing to occur within the airmass with minimal mixing of clean marine air. Other than these

two flights, where fossil fuel emissions had a larger effect than the case studies, the accumulation mode aerosol size distribution was fairly consistent.

By replacing the H^+ ions with NH_4^+ , the neutralisation of the marine sulphuric acid inland should add to the inorganic mass, causing a marginal increase in average particle size. However, the corresponding increase in inorganics was balanced out by similar decrease in ~~ass~~- SO_4 inland, and no consistent differences were observed between the corresponding aerosol size distributions.

One clear feature of the terrestrial influence was the presence of nucleation mode aerosols inland. In all cases, the inland aerosol concentration was significantly higher than those measured over the sea, or typical marine concentrations in the absence of nucleation events (Dall'Osto et al., 2011; Jensen et al., 1996), and the aerosol size distributions showed concentrations of particles <60 nm were always higher inland. Coastal nucleation is unlikely to explain these measurements, as there was no bias in concentrations towards coastal regions.

It is possible that the formation of these particles is also linked to agricultural emissions. Although particle formation rates from H_2SO_4 - NH_3 - H_2O ternary nucleation are lower than those observed in ambient environments, ternary nucleation involving other agricultural emissions (such as amines) may explain our results (Almeida et al., 2013; Pirjola et al., 2000). The high standard deviation/mean ratio of the inland CN concentrations in SWW suggests the process was spatially variable, rather than ubiquitous throughout the boundary layer. Again, however, it is also clear that the generation of these particles had little influence on the CCN concentrations, which were similar over the sea and inland.

It is also of interest to examine the concentrations of giant (dry diameter, $1 \leq D_p \leq 10 \mu m$) and ultragiant particles ($D_p > 10 \mu m$), as these may have an enhanced effect on precipitation formation via the warm rain process (Johnson, 1982), and our observations may be used to inform modelling studies (e.g. Blyth et al., 2013). The number concentrations of giant (N_{GA}) and ultragiant (N_{UGA}) are listed in Table 1. N_{GA} varied by around a factor of two between cases, with values of around one per cubic centimetre, but there was no obvious trend to explain the differences between the different days. For the SWW cases, the inland N_{GA} was fairly proportional to the NW coastal values, but lower by 15 – 40%. These are likely to be sea spray aerosols, and indeed sea salt made up nearly 80% of supermicron aerosol volume on 3rd August (Leon et al., 2015). Additionally, on these days the mean diameter of the coarse aerosol was larger over the NW coast than inland. These measurements are consistent with dilution/deposition of sea salt particles inland being greater than any terrestrial coarse aerosol emissions such as PBAP or dust on these days.

The concentrations of ultragiant aerosol also showed around a factor of two variation between the different cases, and were 3 – 6 L⁻¹, other than the NW coastal measurement on 29 July, which was 11 L⁻¹. Again, it is difficult to pick out any clear trend in the values. On some days, N_{UGA} was higher over the sea, whereas on others it was higher inland. In all cases, N_{GA} and N_{UGA} comprised a very small fraction of the total aerosol number concentration, including the concentration of particles large enough to affect CCN populations.

3.4. Aerosol influence on CCN and cloud drop concentrations

3.4.1 Linking CCN and CDNC

CCN data were only available on 18, 25 and 29 July, and 03 August. Figure 8b shows the average boundary layer CCN concentrations measured on these flights, as well as a comparison to CDNC measured in updrafts near cloud base. Figure 8c shows the in-cloud distribution of vertical wind near cloud base; we use an updraft threshold of 2 m s⁻¹, which is around the 75th percentile of cloud base vertical wind on the four flights shown. By choosing only data in updrafts, as well as excluding data from cloud edges, we aim to minimise the influence of entrainment, though it is unlikely to be eliminated. The higher values of CDNC are those least influenced by entrainment, and those with the strongest updrafts when the cloud formed.

The CN and CCN concentrations showed no correlation, and the fraction of CN active as CCN varied from 2% to 50%. This fraction was largest on 18 July, when the average particle size was the largest of all the flights. On all flights, the CCN fraction was lower inland than over the NW coast, particularly in the SWW cases where it was limited to single-figure percentages even at 0.9% SST. As noted in the previous section, the majority of inland particles were smaller than 60 nm and therefore too small to act as CCN. In these cases, the aerosol size distributions were a more important factor than the CN concentrations for determining CCN concentrations.

On 18 July, the peak CDNC was close to the CCN concentration at 0.1% SST. For the SWW flights, the peak CDNC was more comparable to the CCN concentrations measured at 0.9% SST, though the median CDNC was between the two CCN measurements, which is reasonable based on previous estimates of SST in cumulus clouds. For example, Politovich and Cooper (1988) calculated SST up to ~0.3% in cloud passes with a high fraction of air from near cloud base.

Comparing CCN measured inland and over the sea, the differences were generally <10%, with the largest being 30%. There was no clear systematic difference between land and sea. The addition of ammonium inland did not always increase inorganics, as the change was not always larger than the inherent variability in the ~~ass~~SO₄. On an air mass level, there appears to be an aerosol influence on cloud drop concentrations - the higher aerosol concentrations in the polluted air caused higher CCN concentrations. This difference was also manifested in the CDNC, which was several times higher in the polluted case on 18 July.

Dall'Osto et al. (2010) reported measurements of aerosol composition and CCN at Mace Head, Ireland during late spring and early summer. Mace Head shares similarities with the southwest peninsula of the UK in that it experiences Atlantic air under the prevailing westerly winds but is also a receptor site for aged urban pollution during easterlies. The CCN concentrations

Formatted: Superscript

on 18 July were around double those measured by Dall'Osto et al. (2010) in continental pollution, though the aerosol mass loadings here were also significantly higher. In marine airmasses, the CCN concentrations at Mace Head were $\sim 200 \text{ cm}^{-3}$ at 0.1% SST and $\sim 400 \text{ cm}^{-3}$ at 1.0% SST. These numbers are both in good agreement with our measurements in the SWW cases.

5 3.4.2 Aerosol chemistry and CCN

In marine air, CCN may be produced by sea spray or condensation of secondary aerosols, primarily H_2SO_4 . These processes can interact, for example primary sea spray subsequently coated with H_2SO_4 . Particles produced by sea spray may be composed of sea salt, OA, or a combination of the two, with OA more dominant in submicron particles (Cavalli et al., 2004). Sea salt is an effective CCN; OA from sea spray is likely to be insoluble (Gershey, 1983), but may be internally-mixed with sea salt and/or serve as a condensation nucleus for secondary organic or inorganic aerosol formation. During runs over the southwest peninsula, terrestrial emissions may also have influenced CCN concentrations, such as the addition of ammonia or advected aged urban emissions.

As an air parcel cools as it rises, there may be some co-condensation of semi-volatile species such as ammonium nitrate or SV-OOA, which may further enhance CCN concentrations (Topping et al., 2013). As the aerosol inlets on board the BAe-146 do not reliably sample aerosol during cloud penetrations, we are unable to assess how much of an impact this made, but we note that sulphuric acid / ammonium sulphate (the largest aerosol component on most case studies) is not semi-volatile at the temperatures sampled.

As the source functions of sea spray and secondary organic and inorganic marine and terrestrial aerosols all depend on different factors, they would not be expected to be correlated. Therefore, by comparing the measured CCN concentrations to different chemical species, we may gain information on the sources of the CCN.

Figure 9 shows the relationships between CCN concentrations measured aboard the BAe-146 at 0.1% SST and the PCASP number concentration, AMS total mass, non-sea salt inorganics, and OA. Although the validity of the nitrate measurement is unclear on some flights, the nitrate fraction on these days was low. The data in Figure 9 are split between the three SWW case study flights and the polluted case of 18 July. If we did not make this distinction, all the variation and correlation coefficients would be dominated by the difference between the marine and polluted airmasses, rather than variations in concentration within these airmasses.

The correlation between CCN and aerosol number concentration measured by the PCASP was good, and the data fell close to the 1:1 line, meaning the critical diameter was close to the PCASP's lower cutoff of $\sim 120 \text{ nm}$. Good correlations ($R^2 > 0.66$) were found in both marine and inland environments comparing CCN with total AMS mass and inorganics. The correlations with OA were lower, but non-zero, and the OA correlation in marine air was better than inland. These results are consistent with the main source of CCN being accumulation mode sulphate-containing particles in both marine and inland air. Liu et al. (1996) found a similar correlation with sulphate at 0.06% SST for cloud-processed aerosol.

Figure 10 shows correlations between CCN at 0.9% SST and the same variables as in Figure 9, but only for the SWW flights. The correlations with all four x-variables are lower, meaning CCN at the higher SST are less directly linked to accumulation mode aerosol. In Fig. 10 (a) the CCN are above the 1:1 line with the PCASP number concentration, meaning the critical diameter was below 120 nm. It is therefore not surprising that the correlations at this SST are lower, as the mass measurements are also biased towards larger accumulation mode particles. The species trends in correlation were the same as at 0.1% SST though, in that the correlation with inorganics and AMS total mass were higher than those with OA.

While our data suggest that the average CCN were composed primarily of ~~msm~~-SO₄, it does not provide information on the origin of these particles. For example, marine ~~msm~~-SO₄ may condense onto primary sea-spray OA particles, which are largely insoluble when initially formed (O'Dowd and de Leeuw, 2007). Pierce et al. (2007) showed that this type of process can significantly enhance CCN concentrations globally by providing a surface for sulphuric acid to condense on to. Particle emissions from sea spray are strongly linked to wind speed (O'Dowd et al., 1997). This would mean the marine particle concentrations on 29 July would likely be the highest due to the stronger wind speeds, as shown in Fig. 2. While the marine CN concentration on 29 July was higher than on 25 July, it is difficult to draw firm conclusions from such a limited comparison.

Shipping emissions may also serve as condensation nuclei for secondary aerosol (e.g. Langley et al., 2010), but the low BC concentrations in our case studies suggests this was of little importance. Alternatively, new particles may form through nucleation of gas-phase precursors, and open-ocean nucleation events are thought to be most likely to be related to biological activity (O'Dowd et al., 2010). We gain some insight into particle formation by comparing to the size distributions classified as “clean marine” and “open-ocean nucleation” by Dall'Osto et al. (2011), measured at Mace Head. In particular, the value of dN/dlogDp in our lowest size bin (20 nm) is a useful comparison for the smallest particles, which have experienced the least secondary growth. On 05 and 25 July, the values of the lowest size bin were ~100 cm⁻³, which is consistent with the “clean marine” values (~80 – 150 cm⁻³). The concentration on 29 July was higher at 985 cm⁻³, which is closer to that in “open-ocean nucleation” airmasses (~800 – 2500 cm⁻³), but the value on 03 August of 516 cm⁻³ fell between the two categories. It is therefore not clear what contribution primary and secondary aerosol formation had overall to the aerosol number concentration in marine air, but the majority of CCN mass was composed of ~~msm~~-SO₄, which must be secondary.

This leads us to the conclusions that in the SWW cases, the main factor determining CCN concentrations was the size and composition of marine aerosols reaching the peninsula. The majority of accumulation mode aerosol measured by the AMS in these cases was ~~msm~~-SO₄. The measured CCN also exhibited a high degree of correlation with inorganics, which were predominantly ~~msm~~-SO₄ and its cations. If this correlation were due to the mixing of two homogeneous airmasses, the OA would also show a similar degree of correlation with CCN; however, the R² values for OA were much lower. The data therefore show that whatever process is responsible for the production of sulphate is also largely responsible for determining the concentration of CCN. Our results are consistent with the prevailing theory that the majority of marine CCN are composed of ~~msm~~-SO₄ formed from DMS oxidation (Charlson et al., 1987).

3.5 Sources of INPs

The concentrations of primary ice and the temperatures at which they develop are a large source of uncertainty in the timeline of cloud glaciation. During COPE, the lowest cloud top temperatures were $\sim -15^{\circ}\text{C}$ (Taylor et al., 2016). Reviews by Hoose and Möhler (2012) and Murray et al. (2012) identified the only species that have been shown to behave as ~~INPs~~^{ice} in laboratory experiments at these temperatures as mineral dust; primary biological aerosol particles (PBAP) such as bacteria (Diehl et al., 2006), fungal spores and pollen (Pouleur et al., 1992); and black carbon, although the latter is relatively inefficient. Emissions data for bacteria and fungal spores is limited but global model studies on clouds and precipitation including these species acting as INPs suggest a small but significant zonal influence on precipitation rates (Spracklen and Heald, 2014). ~~but~~ Their concentrations at cloud formation levels are generally much lower than mineral dusts, though there is evidence that ice-active biological material internally-mixed with mineral dust and may enhance the ice nucleating potential of the dust (Augustin-Bauditz et al., 2016). Recently, attention has also focused on marine surface organics, which may be aerosolised as sea spray (Burrows et al., 2013).

Mineral dust concentration and size distribution were measured using the BAe-146 filter system, by combining the ‘silicates’ and ‘mixed silicate’ categories described by Young et al. (2016). Leon et al. (2015) show a more detailed breakdown of the composition of particles from 03 August. BC and OA concentrations were measured on-line using the SP2 and AMS. There was no instrument capable of isolating PBAP aboard the aircraft, but the ground site hosted a WIBS, which made measurements of UV auto-fluorescent particles. The reader should see Sect. 2 for further details of the instrumentation.

In the absence of INP measurements, parameterisations must be used to calculate ~~INP~~^{ice} concentrations. These are empirical relationships derived by comparing measurements of INPs and various chemical species. Figure 11 shows comparisons of several parameterisations, as well as measured first ice concentrations from COPE. The aerosol measurements used for Figure 11 are those from 03 August, as this had the best measurements of ice formation, as presented by Taylor et al. (2016). The measured ice concentrations are from a series of penetrations through a cloud in the early stages of development, when secondary ice processes were thought to have at most a minor influence on ice concentrations (Taylor et al., 2016).

DeMott et al. (2010) based a parameterisation (hereafter D10) for ~~INP~~^{ice} concentrations on the total concentration of aerosol larger than $0.5\ \mu\text{m}$ ($D > 0.5\ \mu\text{m}$), derived from ~~ice-nuclei~~^{INP} measurements with continuous flow diffusion chambers in a variety of environments around the world. Tobo et al. (2013) formed a similar parameterisation (hereafter T13-T) using data measured in a North American pine forest. We used a $D > 0.5\ \mu\text{m}$ concentration of $4.1\ \text{cm}^{-3}$ derived from a combination of the PCASP and CDP measurements during the inland aerosol runs. For comparison, the concentration collected on the filters was $4.35\ \text{cm}^{-3}$, and calculations performed using this number were not discernibly different when plotted on Figure 11. The ~~INP~~^{ice} calculations based on D-10 and T13-T both fall within an order of magnitude of the measured first ice concentrations. The T13-T trace is closer to the measured ice concentrations than D10, meaning the ice active fraction during COPE was more similar to that in a forest ecosystem than in the average of the environments studied in DeMott et al. (2010).

Niemand et al. (2012) and DeMott et al. (2015) provide parameterisations (hereafter referred to as N12 and D15 respectively) base on mineral dusts. N12 uses the integrated mineral dust size distribution to calculate the number of ice surface active sites using measured size distributions, while D15 again simply uses the number concentration of dust particles larger than 0.5 μm , which in our case was 0.24 cm^{-3} based on the filter measurements. ~~Young et al. (2016) found that in a previous study, the concentrations of particles collected on the filters agreed well with in-situ probes for particles larger than 0.5 μm , but the concentrations at filters' collection efficiency was poor for smaller particle sizes were compared poorly.~~ (Young et al., 2016). As D15 uses the dust concentration larger than this size, and N12 is based on the dust surface area distribution, which peaks at supermicron sizes, the filter measurements are expected to be a robust measurement for performing these calculations within the uncertainty of the parameterisations. Both mineral dust parameterisations give ~~ININP~~ concentrations several orders of magnitude below the measured first ice concentrations.

As well as T13-T, Tobo et al. (2013) constructed a parameterisation (hereafter referred to as T13-F) based on measurements of fluorescent particles larger than 0.5 μm , which are likely to be PBAP. The measurements upon which this parameterisation was based were made using an ultraviolet aerodynamic particle sizer (UV-APS), which has a different instrument response to the WIBS. Healy et al. (2014) made concurrent measurements using a UV-APS and a WIBS, and found the WIBS channel 3 (which uses the same fluorophore as the UV-APS) was well correlated with the UV-APS fluorescent concentration, but a factor of 2.7 higher due to the instruments' different size ranges.

Between 1100 – 1500 UTC on 03 August, the total concentration of fluorescent particles measured by the WIBS was $100 \pm 40 \text{ L}^{-1}$. The fluorescent concentration measured using channel 3 was $50 \pm 26 \text{ L}^{-1}$, which corresponds to a UV-APS equivalent concentration of $19 \pm 10 \text{ L}^{-1}$. Using this concentration in the T13-F parameterisation generates ~~ININP~~ concentrations that are also several orders of magnitude lower than the measured ice concentrations. The difference between the predicted ~~ININP~~ concentrations based on T13-T and T13-F are due to the different coarse aerosol composition between the forest environment studied in Tobo et al. (2013) and the marine environment studied here, which is likely to have higher concentrations of sea salt. There may be some difference between the PBAP concentration measured at the ground site that that at the altitude of the aircraft measurements. However, it is unlikely that the PBAP concentration at ~500 m would be three orders of magnitude higher than at ground level, which is what would be required for the INP concentrations calculated using the T13-F parameterisation to agree with the measured ice concentrations.

As an additional estimate of biological INPs, Mohler et al (2008) performed chamber experiments using *pseudomonas syringae* bacteria, and found that the fraction of these bacteria that were active INPs at -10C was 0.5%. Multiplying this fraction by the PBAP concentration measured by the WIBS gives an ~~ININP~~ concentration of 0.5 L^{-1} at this temperature. As *pseudomonas syringae* are amongst the most active biological INPs studied thus far (Lorv et al., 2014), this value may be considered an upper estimate for biological INPs at this temperature. Though it is possible to classify WIBS measurements into different PBAP types, even within simple metaclass discrimination (bacteria, fungal spores and pollen) the uncertainty

in the derived concentrations can be large, e.g. up to a factor of 5 for bacteria, depending on the assumptions used in the analysis approach, as discussed by Crawford et al. (2015).

To date there are no ~~INP~~ parameterisations for soot in the relevant temperature range, and laboratory studies show mixed results, though it is likely ~~to be a worse~~ that soot particles are less efficient INPs than mineral dust (Hoose and Möhler, 2012).

The total mineral dust mass concentration measured was $4 \mu\text{g m}^{-3}$, whereas the total BC concentration was 30 ng m^{-3} , so the ~~INP~~ concentrations from BC were likely to have been several orders of magnitude lower than mineral dust.

Wilson et al. (2015) recently published a mass-based parameterisation for marine surface organics measured in the European Arctic. To our knowledge, this is the only parameterisation published for this type of ~~INP~~. Our results in Figure 11 show that aerosolised marine surface organics are likely to have produced INPs in similar concentrations to mineral dust and PBAP.

In summary, both parameterisations based on total aerosol larger than $0.5 \mu\text{m}$ produced ~~INP~~ concentrations of a similar order of magnitude to the measured first ice concentrations. However, all the composition-specific parameterisations produced ~~INP~~ concentrations several orders of magnitude lower. This therefore leaves us with two possible conclusions: either there was a source of INPs that was not characterised by our measurements (i.e. a species other than PBAP, mineral dust and sea spray OA that ~~were~~ was significantly active INPs at $-5 > T > -15^\circ\text{C}$), and/or one or more of the composition-specific parameterisations underestimate ~~INP~~ concentrations when used in the southwest peninsula of the UK. The measured ice concentrations lie in the temperature range at the high end of most of the parameterisations tested. For example, the data used to generate the fit parameters for T13-F included just seven data points at -10°C , and the majority of points were from temperatures $\leq -20^\circ\text{C}$. Mason et al. (2015) also noted the poor performance of ~~INP~~ parameterisations at a coastal site, and noted the good correlation of PBAP with INPs between -15 and -25°C in air from the Pacific Ocean. It is possible the sources of INPs are similar between the two studies, but without further characterisation of INPs active at temperatures higher than -15°C , we are limited to such speculation.

4. Summary and conclusions

We have presented measurements of boundary layer aerosol concentration, size distribution and composition from a series of research flights performed over the southwest peninsula of the UK during the COPE campaign of summer 2013. Several case studies showed marine air from the Atlantic Ocean advecting to the peninsula from the south/southwest. The CCN and submicron aerosol in these cases were predominantly sulphate-based, and typical for Atlantic marine air. During Easterly winds, polluted air from the UK brought urban pollution to the region; the submicron aerosol was composed of ammonium nitrate and organics, and CCN concentrations were several times higher than the marine cases. In one marine case, ~~INP~~ concentrations were estimated using by inputting field measurements into several size- and composition-specific

parameterisations. Bulk parameterisations predicted ~~ININP~~ concentrations within an order of magnitude to measured first ice concentrations, but attempts to identify the composition of the INPs were largely inconclusive.

The periods of southwesterly winds, which brought marine air from the Atlantic to the peninsula, are of particular interest as they are the most conducive to convective cloud formation in the region. The CCN concentrations in these conditions were ~150 – 280 cm⁻³ at 0.1% SST, and 400 – 500 cm⁻³ at 0.9% SST, which are in good agreement with previous measurements in Atlantic air performed at Mace Head (Dall'Osto et al., 2010). The marine aerosol size distributions were also similar to those measured at Mace Head during “clean marine” conditions (Dall'Osto et al., 2011), though some flights had enhanced concentrations of nucleation mode particles, the source of which is not clear from our measurements.

The majority of accumulation mode mass was composed of ~~ss-SO~~₄ on these days, which is likely to be from biogenic DMS oxidation (Charlson et al., 1987). The high degree of correlation between CCN and ~~ss-SO~~ suggests the CCN were predominantly composed of ~~ss-SO~~₄, though the origin of these particles (i.e. coated primary or nucleated secondary particles) is again unclear. Accumulation mode OA had a similar mass spectrum to previous measurements of sea spray OA (Ovadnevaite et al., 2011), but a low correlation with CCN, meaning sea spray OA that was not internally-mixed with ~~ss-SO~~₄ was not a large source of CCN.

By comparing aerosol over the sea to that measured inland, we infer the influence terrestrial emissions had on the marine aerosol. The marine ~~ss-SO~~₄ was acidic over the sea, but was neutralised by terrestrial ammonia, which is likely to have been emitted by agricultural activities that are widespread throughout the peninsula. The increase in inorganics by this process was almost equally balanced out by a decrease in ~~ss-SO~~₄. Consequently, the differences between oceanic and inland CCN concentrations were less than the day-to-day variability in different marine cases. Additionally, large enhancements of nucleation mode particles were measured inland, which may also be related to agricultural emissions interacting with marine H₂SO₄, but these particles were too small to affect CCN concentrations.

The largest difference in aerosol mass loadings and CCN were between the marine flights and a case study where polluted urban emissions advected to the site from the east. The CCN concentrations were ~6 times higher in this air mass, and the inorganics were composed of equal amounts of ammonium nitrate and sulphate. The difference in CCN also corresponded to differences in the cloud base CDNC. In the SWW cases the peak CDNCs were ~450 cm⁻³, which is in agreement with the CCN measured at 0.9% SST, though most values of CDNC at cloud base were closer to 300 cm⁻³, which suggests the SST at cloud base was typically between 0.1% and 0.9%. In the polluted case, the peak CDNC was ~1300 cm⁻³, which was in agreement with CCN measured at 0.1% SST in this case.

Possible sources of INPs were assessed by comparing various different parameterisations, using our measured aerosol concentrations as input. We estimated ~~ININP~~ concentrations using measurements of total aerosol larger than 0.5 µm (DeMott et al., 2010; Tobo et al., 2013), mineral dust (DeMott et al., 2015; Niemand et al., 2012), fluorescent aerosol (Möhler et al., 2008; Tobo et al., 2013) and sea spray OA (Wilson et al., 2015). The parameterisations based on total aerosol produced ~~ININP~~ concentrations that agreed with measured first ice concentrations within an order of magnitude.

In contrast, the predicted ~~ININP~~ concentrations derived using all of the composition-specific parameterisations were several orders of magnitude lower. Either there was a source of INPs that was not characterised by our measurements, and/or one or more of the composition-specific parameterisations tested greatly underestimated ~~ININP~~ concentrations when used in the southwest peninsula of the UK. The lowest cloud-top temperatures during COPE were -15 °C. Better characterisation of INPs at these relatively warm temperatures would be required to gain any further insight into the sources of INPs during COPE. It is, however, very unlikely that the ice concentrations of up to several hundred per litre reported by Taylor et al. (2016) could be achieved without invoking the action of secondary ice processes.

The results presented in this paper serve as information for modellers simulating cloud-aerosol interactions for the COPE project, providing details on the aerosol size, composition, CCN and INPs in the different case studies. Our data also provide characterisation of Atlantic aerosol, and are likely to be particularly applicable to regions where marine air makes landfall in the absence of strong fossil fuel emissions.

Acknowledgements

The authors wish to thank all those involved in COPE. The BAe-146-301 Atmospheric Research Aircraft was flown by Directflight Ltd and managed by the Facility for Airborne Atmospheric Measurements (FAAM), which is a joint entity of the Natural Environment Research Council (NERC) and the Met Office. Processed data are available from the British Atmospheric Data Centre. Raw data are archived at the University of Manchester and available on request. COPE was supported by NERC under grant numbers NE/J022594/1 and NE/J023507/1, and the Met Office funded the operation of the BAe-146 FAAM aircraft. We also thank Jurgita Ovadnevaite and colleagues from the National University of Ireland, Galway for providing the sea spray mass spectrum. Finally, we thank the editor and two anonymous referees for helping improve the scientific quality of the manuscript.

References

Alfarra, M. R.: Insights Into Atmospheric Organic Aerosols Using An Aerosol Mass Spectrometer, University of Manchester Institute of Science and Technology, Manchester, UK., 2004.

Allan, J. D., Alfarra, M. R., Bower, K. N., Coe, H., Jayne, J. T., Worsnop, D. R., Aalto, P. P., Kulmala, M., Hyötyläinen, T., Cavalli, F. and Laaksonen, A.: Size and composition measurements of background aerosol and new particle growth in a Finnish forest during QUEST 2 using an Aerodyne Aerosol Mass Spectrometer, *Atmos. Chem. Phys.*, 6(2), 315–327, doi:10.5194/acp-6-315-2006, 2006.

Allan, J. D., Bower, K. N., Coe, H., Boudries, H., Jayne, J. T., Canagaratna, M. R., Millet, D. B., Goldstein, A. H., Quinn, P. K., Weber, R. J. and Worsnop, D. R.: Submicron aerosol composition at Trinidad Head, California, during ITCT 2K2: Its

- relationship with gas phase volatile organic carbon and assessment of instrument performance, *J. Geophys. Res. Atmos.*, 109(D23), n/a–n/a, doi:10.1029/2003JD004208, 2004.
- Allan, J. D., Jimenez, J. L., Williams, P. I., Alfarra, M. R., Bower, K. N., Jayne, J. T., Coe, H. and Worsnop, D. R.: Quantitative sampling using an Aerodyne aerosol mass spectrometer - 1. Techniques of data interpretation and error analysis, *J. Geophys. Res.*, 108, doi:4090 10.1029/2002jd002358, 2003.
- Allan, J. D., Morgan, W. T., Darbyshire, E., Flynn, M. J., Williams, P. I., Oram, D. E., Artaxo, P., Brito, J., Lee, J. D. and Coe, H.: Airborne observations of IEPOX-derived isoprene SOA in the Amazon during SAMBBA, *Atmos. Chem. Phys.*, 14(20), 11393–11407, doi:10.5194/acp-14-11393-2014, 2014.
- Almeida, J., Schobesberger, S., Kürten, A., Ortega, I. K., Kupiainen-Määttä, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppä, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhoffer, R., Seinfeld, J. H., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H. and Kirkby, J.: Molecular understanding of sulphuric acid-amine particle nucleation in the atmosphere., *Nature*, 502(7471), 359–63, doi:10.1038/nature12663, 2013.
- Asmi, A., Wiedensohler, A., Laj, P., Fjaeraa, A.-M., Sellegri, K., Birmili, W., Weingartner, E., Baltensperger, U., Zdimal, V., Zikova, N., Putaud, J.-P., Marinoni, A., Tunved, P., Hansson, H.-C., Fiebig, M., Kivekäs, N., Lihavainen, H., Asmi, E., Ulevicius, V., Aalto, P. P., Swietlicki, E., Kristensson, A., Mihalopoulos, N., Kalivitis, N., Kalapov, I., Kiss, G., de Leeuw, G., Henzing, B., Harrison, R. M., Beddows, D., O'Dowd, C., Jennings, S. G., Flentje, H., Weinhold, K., Meinhardt, F., Ries, L. and Kulmala, M.: Number size distributions and seasonality of submicron particles in Europe 2008–2009, *Atmos. Chem. Phys.*, 11(11), 5505–5538, doi:10.5194/acp-11-5505-2011, 2011.
- Augustin-Bauditz, S., Wex, H., Denjean, C., Hartmann, S., Schneider, J., Schmidt, S., Ebert, M. and Stratmann, F.: Laboratory-generated mixtures of mineral dust particles with biological substances: characterization of the particle mixing state and immersion freezing behavior, *Atmos. Chem. Phys.*, 16(9), 5531–5543, doi:10.5194/acp-16-5531-2016, 2016.
- Bahreini, R., Ervens, B., Middlebrook, A. M., Warneke, C., de Gouw, J. A., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., Neuman, J. A., Ryerson, T. B., Stark, H., Atlas, E., Brioude, J., Fried, A., Holloway, J. S., Peischl, J., Richter, D., Walega, J., Weibring, P., Wollny, A. G. and Fehsenfeld, F. C.: Organic aerosol formation in urban and industrial plumes near Houston and Dallas, Texas, *J. Geophys. Res.*, 114, D00F16, doi:10.1029/2008JD011493, 2009.
- Baumgardner, D., Popovicheva, O., Allan, J., Bernardoni, V., Cao, J., Cavalli, F., Cozic, J., Diapouli, E., Eleftheriadis, K., Genberg, P. J., Gonzalez, C., Gysel, M., John, A., Kirchstetter, T. W., Kuhlbusch, T. A. J., Laborde, M., Lack, D., Müller, T., Niessner, R., Petzold, A., Piazzalunga, A., Putaud, J. P., Schwarz, J., Sheridan, P., Subramanian, R., Swietlicki, E., Valli, G., Vecchi, R. and Viana, M.: Soot reference materials for instrument calibration and intercomparisons: a workshop

- p>summary with recommendations,
- Atmos. Meas. Tech.*
- , 5(8), 1869–1887, doi:10.5194/amt-5-1869-2012, 2012.
- Beswick, K. M., Gallagher, M. W., Webb, A. R., Norton, E. G. and Perry, F.: Application of the Aventech AIMMS20AQ airborne probe for turbulence measurements during the Convective Storm Initiation Project, *Atmos. Chem. Phys.*, 8(17), 5449–5463, doi:10.5194/acp-8-5449-2008, 2008.
- 5 Blyth, A. M., Lowenstein, J. H., Huang, Y., Cui, Z., Davies, S. and Carslaw, K. S.: The production of warm rain in shallow maritime cumulus clouds, *Q. J. R. Meteorol. Soc.*, 139(670), 20–31, doi:10.1002/qj.1972, 2013.
- Bohnenstengel, S. I., Belcher, S. E., Aiken, A., Allan, J. D., Allen, G., Bacak, A., Bannan, T. J., Barlow, J. F., Beddows, D. C. S., Bloss, W. J., Booth, A. M., Chemel, C., Coceal, O., Di Marco, C. F., Dubey, M. K., Faloony, K. H., Fleming, Z. L., Furger, M., Gietl, J. K., Graves, R. R., Green, D. C., Grimmond, C. S. B., Halios, C. H., Hamilton, J. F., Harrison, R. M.,
10 Heal, M. R., Heard, D. E., Helfter, C., Herndon, S. C., Holmes, R. E., Hopkins, J. R., Jones, A. M., Kelly, F. J., Kotthaus, S., Langford, B., Lee, J. D., Leigh, R. J., Lewis, A. C., Lidster, R. T., Lopez-Hilfiker, F. D., McQuaid, J. B., Mohr, C., Monks, P. S., Nemitz, E., Ng, N. L., Percival, C. J., Prévôt, A. S. H., Ricketts, H. M. A., Sokhi, R., Stone, D., Thornton, J. A., Tremper, A. H., Valach, A. C., Visser, S., Whalley, L. K., Williams, L. R., Xu, L., Young, D. E. and Zotter, P.: Meteorology, Air Quality, and Health in London: The ClearfLo Project, *Bull. Am. Meteorol. Soc.*, 96(5), 779–804, doi:10.1175/BAMS-D-
15 12-00245.1, 2015.
- Burrows, S. M., Hoose, C., Pöschl, U. and Lawrence, M. G.: Ice nuclei in marine air: biogenic particles or dust?, *Atmos. Chem. Phys.*, 13(1), 245–267, doi:10.5194/acp-13-245-2013, 2013.
- Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P. and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer., *Mass Spectrom. Rev.*, 26(2), 185–222, doi:10.1002/mas.20115, 2007.
- 20 Cavalli, F., Facchini, M. C., Decesari, S., Mircea, M., Emblico, L., Fuzzi, S., Ceburnis, D., Yoon, Y. J., O’Dowd, C. D., Putaud, J.-P. and Dell’Acqua, A.: Advances in characterization of size-resolved organic matter in marine aerosol over the North Atlantic, *J. Geophys. Res. Atmos.*, 109(D24), doi:10.1029/2004JD005137, 2004.
- 25 Chang, R. Y.-W., Leck, C., Graus, M., Müller, M., Paatero, J., Burkhardt, J. F., Stohl, A., Orr, L. H., Hayden, K., Li, S.-M., Hansel, A., Tjernström, M., Leaitch, W. R. and Abbatt, J. P. D.: Aerosol composition and sources in the central Arctic Ocean during ASCOS, *Atmos. Chem. Phys.*, 11(20), 10619–10636, doi:10.5194/acp-11-10619-2011, 2011.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O. and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326(6114), 655–661, doi:10.1038/326655a0, 1987.
- 30 Clark, P., Roberts, N., Lean, H., Ballard, S. P. and Charlton-Perez, C.: Convection-permitting models: a step-change in rainfall forecasting, *Meteorol. Appl.*, 23(2), 165–181, doi:10.1002/met.1538, 2016.

- Connolly, P. J., Möhler, O., Field, P. R., Saathoff, H., Burgess, R., Choularton, T. and Gallagher, M.: Studies of heterogeneous freezing by three different desert dust samples, *Atmos. Chem. Phys.*, 9(8), 2805–2824, doi:10.5194/acp-9-2805-2009, 2009.
- Crawford, I., Ruske, S., Topping, D. O. and Gallagher, M. W.: Evaluation of hierarchical agglomerative cluster analysis methods for discrimination of primary biological aerosol, *Atmos. Meas. Tech. Discuss.*, 8(7), 7303–7333, doi:10.5194/amtd-8-7303-2015, 2015.
- Cross, E. S., Slowik, J. G., Davidovits, P., Allan, J. D., Worsnop, D. R., Jayne, J. T., Lewis, D. K., Canagaratna, M. and Onasch, T. B.: Laboratory and ambient particle density determinations using light scattering in conjunction with aerosol mass spectrometry, *Aerosol Sci. Technol.*, 41, 343–359, doi:10.1080/02786820701199736, 2007.
- 10 Dall'Osto, M., Ceburnis, D., Martucci, G., Bialek, J., Dupuy, R., Jennings, S. G., Berresheim, H., Wenger, J., Healy, R., Facchini, M. C., Rinaldi, M., Giulianelli, L., Finessi, E., Worsnop, D., Ehn, M., Mikkilä, J., Kulmala, M. and O'Dowd, C. D.: Aerosol properties associated with air masses arriving into the North East Atlantic during the 2008 Mace Head EUCAARI intensive observing period: an overview, *Atmos. Chem. Phys.*, 10(17), 8413–8435, doi:10.5194/acp-10-8413-2010, 2010.
- 15 Dall'Osto, M., Monahan, C., Greaney, R., Beddows, D. C. S., Harrison, R. M., Ceburnis, D. and O'Dowd, C. D.: A statistical analysis of North East Atlantic (submicron) aerosol size distributions, *Atmos. Chem. Phys.*, 11(24), 12567–12578, doi:10.5194/acp-11-12567-2011, 2011.
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T. and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate., *Proc. Natl. Acad. Sci. U. S. A.*, 107(25), 11217–22, doi:10.1073/pnas.0910818107, 2010.
- 20 DeMott, P. J., Prenni, A. J., McMeeking, G. R., Sullivan, R. C., Petters, M. D., Tobo, Y., Niemand, M., Möhler, O., Snider, J. R., Wang, Z. and Kreidenweis, S. M.: Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles, *Atmos. Chem. Phys.*, 15(1), 393–409, doi:10.5194/acp-15-393-2015, 2015.
- Diehl, K., Simmel, M. and Wurzler, S.: Numerical sensitivity studies on the impact of aerosol properties and drop freezing modes on the glaciation, microphysics, and dynamics of clouds, *J. Geophys. Res.*, 111(D7), D07202, doi:10.1029/2005JD005884, 2006.
- 25 Drewnick, F., Diesch, J.-M., Faber, P. and Borrmann, S.: Aerosol mass spectrometry: particle–vaporizer interactions and their consequences for the measurements, *Atmos. Meas. Tech. Discuss.*, 8(4), 3525–3570, doi:10.5194/amtd-8-3525-2015, 2015.
- 30 Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S. and Worsnop, D. R.: A New Time-of-Flight Aerosol Mass Spectrometer (TOF-AMS)- Instrument Description and First Field Deployment, *Aerosol Sci. Technol.*, 39(7), 637–658, doi:10.1080/02786820500182040, 2005.

- Fan, J., Zhang, R., Li, G. and Tao, W.-K.: Effects of aerosols and relative humidity on cumulus clouds, *J. Geophys. Res.*, 112(D14), D14204, doi:10.1029/2006JD008136, 2007.
- Formenti, P., Rajot, J. L., Desboeufs, K., Caqueneau, S., Chevaillier, S., Nava, S., Gaudichet, A., Journet, E., Triquet, S., Alfaro, S., Chiari, M., Haywood, J., Coe, H. and Highwood, E.: Regional variability of the composition of mineral dust from western Africa: Results from the AMMA SOP0/DABEX and DODO field campaigns, *J. Geophys. Res.*, 113, D00C13, doi:10.1029/2008JD009903, 2008.
- Gerbig, C., Schmitgen, S., Kley, D., Volz-Thomas, A., Dewey, K. and Haaks, D.: An improved fast-response vacuum-UV resonance fluorescence CO instrument, *J. Geophys. Res.*, 104(D1), 1699, doi:10.1029/1998JD100031, 1999.
- Gershey, R. M.: Characterization of seawater organic matter carried by bubble-generated aerosols, *Limnol. Oceanogr.*, 28(2), 309–319, doi:10.4319/lo.1983.28.2.0309, 1983.
- Golding, B., Ed.: Boscastle and north Cornwall post flood event study–meteorological analysis of the conditions leading to flooding on 16 August 2004, Met Office., 2005.
- Golding, B., Clark, P. and May, B.: The Boscastle flood: Meteorological analysis of the conditions leading to flooding on 16 August 2004, *Weather*, 60(8), 230–235, doi:10.1256/wea.71.05, 2005.
- Hand, V. L., Capes, G., Vaughan, D. J., Formenti, P., Haywood, J. M. and Coe, H.: Evidence of internal mixing of African dust and biomass burning particles by individual particle analysis using electron beam techniques, *J. Geophys. Res.*, 115(D13), D13301, doi:10.1029/2009JD012938, 2010.
- Harrison, R. M. and Pio, C. A.: Size-differentiated composition of inorganic atmospheric aerosols of both marine and polluted continental origin, *Atmos. Environ.*, 17(9), 1733–1738, doi:10.1016/0004-6981(83)90180-4, 1983.
- Healy, D. A., Huffman, J. A., O'Connor, D. J., Pöhlker, C., Pöschl, U. and Sodeau, J. R.: Ambient measurements of biological aerosol particles near Killarney, Ireland: a comparison between real-time fluorescence and microscopy techniques, *Atmos. Chem. Phys.*, 14(15), 8055–8069, doi:10.5194/acp-14-8055-2014, 2014.
- Hering, S. V., Stolzenburg, M. R., Quant, F. R., Oberreit, D. R. and Keady, P. B.: A Laminar-Flow, Water-Based Condensation Particle Counter (WCPC), *Aerosol Sci. Technol.*, 39(7), 659–672, 2005.
- Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12(20), 9817–9854, doi:10.5194/acp-12-9817-2012, 2012.
- Hoppel, W. A., Frick, G. M. and Larson, R. E.: Effect of nonprecipitating clouds on the aerosol size distribution in the

- marine boundary layer, *Geophys. Res. Lett.*, 13(2), 125–128, doi:10.1029/GL013i002p00125, 1986.
- Huang, Y., Blyth, A. M., Brown, P. R. A., Choularton, T. W., Connolly, P., Gadian, A. M., Jones, H., Latham, J., Cui, Z. and Carslaw, K.: The development of ice in a cumulus cloud over southwest England, *New J. Phys.*, 10(10), 105021, doi:10.1088/1367-2630/10/10/105021, 2008.
- 5 Huebert, B. J., Zhuang, L., Howell, S., Noone, K. and Noone, B.: Sulfate, nitrate, methanesulfonate, chloride, ammonium, and sodium measurements from ship, island, and aircraft during the Atlantic Stratocumulus Transition Experiment/Marine Aerosol Gas Exchange, *J. Geophys. Res.*, 101(D2), 4413, doi:10.1029/95JD02044, 1996.
- Huffman, J. A., Treutlein, B. and Pöschl, U.: Fluorescent biological aerosol particle concentrations and size distributions measured with an Ultraviolet Aerodynamic Particle Sizer (UV-APS) in Central Europe, *Atmos. Chem. Phys.*, 10(7), 3215–3233, doi:10.5194/acp-10-3215-2010, 2010.
- 10 3233, doi:10.5194/acp-10-3215-2010, 2010.
- Jensen, T. L., Kreidenweis, S. M., Kim, Y., Sievering, H. and Pszenny, A.: Aerosol distributions in the North Atlantic marine boundary layer during Atlantic Stratocumulus Transition Experiment/Marine Aerosol and Gas Exchange, *J. Geophys. Res.*, 101(D2), 4455, doi:10.1029/95JD00506, 1996.
- Johnson, D. B.: The Role of Giant and Ultragiant Aerosol Particles in Warm Rain Initiation, *J. Atmos. Sci.*, 39(2), 448–460, doi:10.1175/1520-0469(1982)039<0448:TROGAU>2.0.CO;2, 1982.
- 15 doi:10.1175/1520-0469(1982)039<0448:TROGAU>2.0.CO;2, 1982.
- Klett, J. D. and Davis, M. H.: Theoretical Collision Efficiencies of Cloud Droplets at Small Reynolds Numbers, *J. Atmos. Sci.*, 30(1), 107–117, doi:10.1175/1520-0469(1973)030<0107:TCEOCD>2.0.CO;2, 1973.
- Koren, I., Kaufman, Y. J., Rosenfeld, D., Remer, L. A. and Rudich, Y.: Aerosol invigoration and restructuring of Atlantic convective clouds, *Geophys. Res. Lett.*, 32(14), n/a–n/a, doi:10.1029/2005GL023187, 2005.
- Laborde, M., Mertes, P., Zieger, P., Dommen, J., Baltensperger, U. and Gysel, M.: Sensitivity of the Single Particle Soot Photometer to different black carbon types, *Atmos. Meas. Tech.*, 5(5), 1031–1043, doi:10.5194/amt-5-1031-2012, 2012a.
- 20 Laborde, M., Schnaiter, M., Linke, C., Saathoff, H., Naumann, K.-H., Möhler, O., Berlenz, S., Wagner, U., Taylor, J. W., Liu, D., Flynn, M., Allan, J. D., Coe, H., Heimerl, K., Dahlkötter, F., Weinzierl, B., Wollny, A. G., Zanatta, M., Cozic, J., Laj, P., Hitenberger, R., Schwarz, J. P. and Gysel, M.: Single Particle Soot Photometer intercomparison at the AIDA chamber, *Atmos. Meas. Tech.*, 5(12), 3077–3097, doi:10.5194/amt-5-3077-2012, 2012b.
- 25 Laborde, M., Leaitch, W. R., Lohmann, U., Shantz, N. C. and Worsnop, D. R.: Contributions from DMS and ship emissions to CCN observed over the summertime North Pacific, *Atmos. Chem. Phys.*, 10(3), 1287–1314, doi:10.5194/acp-10-1287-2010, 2010.

- Leon, D. C., French, J. R., Lasher-Trapp, S., Blyth, A. M., Abel, S. J., Ballard, S., Barrett, A., Bennett, L. J., Bower, K., Brooks, B., Brown, P., Charlton-Perez, C., Choularton, T., Clark, P., Collier, C., Crosier, J., Cui, Z., Dey, S., Dufton, D., Eagle, C., Flynn, M. J., Gallagher, M., Halliwell, C., Hanley, K., Hawkness-Smith, L., Huang, Y., Kelly, G., Kitchen, M., Korolev, A., Lean, H., Liu, Z., Marsham, J., Moser, D., Nicol, J., Norton, E. G., Plummer, D., Price, J., Ricketts, H., Roberts, N., Rosenberg, P. D., Simonin, D., Taylor, J. W., Warren, R., Williams, P. I. and Young, G.: The COncvective Precipitation Experiment (COPE): Investigating the origins of heavy precipitation in the southwestern UK, *Bull. Am. Meteorol. Soc.*, doi:10.1175/BAMS-D-14-00157.1, 2015.
- Liu, P. S. K., Leaitch, W. R., Banic, C. M., Li, S.-M., Ngo, D. and Megaw, W. J.: Aerosol observations at Chebogue Point during the 1993 North Atlantic Regional Experiment: Relationships among cloud condensation nuclei, size distribution, and chemistry, *J. Geophys. Res.*, 101(D22), 28971, doi:10.1029/96JD00445, 1996.
- Lorv, J. S. H., Rose, D. R. and Glick, B. R.: Bacterial ice crystal controlling proteins., *Scientifica (Cairo)*., 2014, 976895, doi:10.1155/2014/976895, 2014.
- Mason, R. H., Si, M., Li, J., Chou, C., Dickie, R., Toom-Sauntry, D., Pöhlker, C., Yakobi-Hancock, J. D., Ladino, L. A., Jones, K., Leaitch, W. R., Schiller, C. L., Abbatt, J. P. D., Huffman, J. A. and Bertram, A. K.: Ice nucleating particles at a coastal marine boundary layer site: correlations with aerosol type and meteorological conditions, *Atmos. Chem. Phys.*, 15(21), 12547–12566, doi:10.5194/acp-15-12547-2015, 2015.
- Middlebrook, A. M., Bahreini, R., Jimenez, J. L. and Canagaratna, M. R.: Evaluation of Composition-Dependent Collection Efficiencies for the Aerodyne Aerosol Mass Spectrometer using Field Data, *Aerosol Sci. Technol.*, 46(3), 258–271, doi:10.1080/02786826.2011.620041, 2012.
- Möhler, O., Georgakopoulos, D. G., Morris, C. E., Benz, S., Ebert, V., Hunsmann, S., Saathoff, H., Schnaiter, M. and Wagner, R.: Heterogeneous ice nucleation activity of bacteria: new laboratory experiments at simulated cloud conditions, *Biogeosciences*, 5(5), 1425–1435, doi:10.5194/bg-5-1425-2008, 2008.
- Morgan, W. T., Allan, J. D., Bower, K. N., Highwood, E. J., Liu, D., McMeeking, G. R., Northway, M. J., Williams, P. I., Krejci, R. and Coe, H.: Airborne measurements of the spatial distribution of aerosol chemical composition across Europe and evolution of the organic fraction, *Atmos. Chem. Phys.*, 10(8), 4065–4083, doi:10.5194/acp-10-4065-2010, 2010.
- Morgan, W. T., Ouyang, B., Allan, J. D., Aruffo, E., Di Carlo, P., Kennedy, O. J., Lowe, D., Flynn, M. J., Rosenberg, P. D., Williams, P. I., Jones, R., McFiggans, G. B. and Coe, H.: Influence of aerosol chemical composition on N₂O₅ uptake: airborne regional measurements in North-Western Europe, *Atmos. Chem. Phys. Discuss.*, 14(13), 19673–19718, doi:10.5194/acpd-14-19673-2014, 2014.
- Morton, D., Rowland, C. S., Wood, C. M., Meek, L., Marston, C., Smith, G., Wadsworth, R. A. and Simpson, I. C.: Final Report for LCM2007 - the new UK land cover map. Countryside Survey Technical Report No 11/07, 2011.
- Murray, B. J., O’Sullivan, D., Atkinson, J. D. and Webb, M. E.: Ice nucleation by particles immersed in supercooled cloud

- p>droplets.,
- Chem. Soc. Rev.*
- , 41(19), 6519–54, doi:10.1039/c2cs35200a, 2012.
- Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prevot, A. S. H., Dinar, E., Rudich, Y. and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, doi:10.5194/acp-10-4625-2010, 2010.
- Niemand, M., Möhler, O., Vogel, B., Vogel, H., Hoose, C., Connolly, P., Klein, H., Bingemer, H., DeMott, P., Skrotzki, J. and Leisner, T.: A Particle-Surface-Area-Based Parameterization of Immersion Freezing on Desert Dust Particles, *J. Atmos. Sci.*, 69(10), 3077–3092, doi:10.1175/JAS-D-11-0249.1, 2012.
- Nuaaman, I., Li, S.-M., Hayden, K. L., Onasch, T. B., Massoli, P., Sueper, D., Worsnop, D. R., Bates, T. S., Quinn, P. K. and McLaren, R.: Separating refractory and non-refractory particulate chloride and estimating chloride depletion by aerosol mass spectrometry in a marine environment, *Atmos. Chem. Phys. Discuss.*, 15(2), 2085–2118, doi:10.5194/acpd-15-2085-2015, 2015.
- O’Dowd, C., Ceburnis, D., Ovadnevaite, J., Vaishya, A., Rinaldi, M. and Facchini, M. C.: Do anthropogenic, continental or coastal aerosol sources impact on a marine aerosol signature at Mace Head?, *Atmos. Chem. Phys.*, 14(19), 10687–10704, doi:10.5194/acp-14-10687-2014, 2014.
- O’Dowd, C. D. and de Leeuw, G.: Marine aerosol production: a review of the current knowledge., *Philos. Trans. A. Math. Phys. Eng. Sci.*, 365(1856), 1753–74, doi:10.1098/rsta.2007.2043, 2007.
- O’Dowd, C. D., Smith, M. H., Consterdine, I. E. and Lowe, J. A.: Marine aerosol, sea-salt, and the marine sulphur cycle: a short review, *Atmos. Environ.*, 31(1), 73–80, doi:10.1016/S1352-2310(96)00106-9, 1997.
- O’Dowd, C., Monahan, C. and Dall’Osto, M.: On the occurrence of open ocean particle production and growth events, *Geophys. Res. Lett.*, 37(19), n/a–n/a, doi:10.1029/2010GL044679, 2010.
- Ovadnevaite, J., Ceburnis, D., Canagaratna, M., Berresheim, H., Bialek, J., Martucci, G., Worsnop, D. R. and O’Dowd, C.: On the effect of wind speed on submicron sea salt mass concentrations and source fluxes, *J. Geophys. Res.*, 117(D16), D16201, doi:10.1029/2011JD017379, 2012.
- Ovadnevaite, J., Ceburnis, D., Leinert, S., Dall’Osto, M., Canagaratna, M., O’Doherty, S., Berresheim, H. and O’Dowd, C.: Submicron NE Atlantic marine aerosol chemical composition and abundance: Seasonal trends and air mass categorization, *J. Geophys. Res. Atmos.*, 119(20), 11,850–11,863, doi:10.1002/2013JD021330, 2014.
- Ovadnevaite, J., O’Dowd, C., Dall’Osto, M., Ceburnis, D., Worsnop, D. R. and Berresheim, H.: Detecting high contributions of primary organic matter to marine aerosol: A case study, *Geophys. Res. Lett.*, 38(2), L02807, doi:10.1029/2010GL046083, 2011.

- Phillips, V. T. J., Blyth, A. M., Brown, P. R. A., Choularton, T. W. and Latham, J.: The glaciation of a cumulus cloud over New Mexico, *Q. J. R. Meteorol. Soc.*, 127(575), 1513–1534, doi:10.1002/qj.49712757503, 2001.
- Phinney, L., Richard Leaitch, W., Lohmann, U., Boudries, H., Worsnop, D. R., Jayne, J. T., Toom-Sauntry, D., Wadleigh, M., Sharma, S. and Shantz, N.: Characterization of the aerosol over the sub-arctic north east Pacific Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 53(20-22), 2410–2433, doi:10.1016/j.dsr2.2006.05.044, 2006.
- 5 Pierce, J. R., Chen, K. and Adams, P. J.: Contribution of primary carbonaceous aerosol to cloud condensation nuclei: processes and uncertainties evaluated with a global aerosol microphysics model, *Atmos. Chem. Phys.*, 7(20), 5447–5466, doi:10.5194/acp-7-5447-2007, 2007.
- Pirjola, L., O'Dowd, C. D., Brooks, I. M. and Kulmala, M.: Can new particle formation occur in the clean marine boundary layer?, *J. Geophys. Res.*, 105(D21), 26531, doi:10.1029/2000JD900310, 2000.
- 10 Politovich, M. K. and Cooper, W. A.: Variability of the Supersaturation in Cumulus Clouds, *J. Atmos. Sci.*, 45(11), 1651–1664, doi:10.1175/1520-0469(1988)045<1651:VOTSIC>2.0.CO;2, 1988.
- Pouleur, S., Richard, C., Martin, J.-G. and Antoun, H.: Ice Nucleation Activity in *Fusarium acuminatum* and *Fusarium avenaceum*, *Appl. Envir. Microbiol.*, 58(9), 2960–2964, 1992.
- 15 Quinn, P. K., Barrett, K. J., Dentener, F. J., Lipschultz, F. and Six, K. D.: Estimation of the air/sea exchange of ammonia for the North Atlantic Basin, *Biogeochemistry*, 35(1), 275–304, doi:10.1007/BF02179831, 1996.
- Roberts, G. C., Day, D. A., Russell, L. M., Dunlea, E. J., Jimenez, J. L., Tomlinson, J. M., Collins, D. R., Shinozuka, Y. and Clarke, A. D.: Characterization of particle cloud droplet activity and composition in the free troposphere and the boundary layer during INTEX-B, *Atmos. Chem. Phys.*, 10(14), 6627–6644, doi:10.5194/acp-10-6627-2010, 2010.
- 20 Roberts, G. C. and Nenes, A.: A Continuous-Flow Streamwise Thermal-Gradient CCN Chamber for Atmospheric Measurements, *Aerosol Sci. Technol.*, 39, 206–221, doi:10.1080/027868290913988, 2005.
- Rollins, A. W., Fry, J. L., Hunter, J. F., Kroll, J. H., Worsnop, D. R., Singaram, S. W. and Cohen, R. C.: Elemental analysis of aerosol organic nitrates with electron ionization high-resolution mass spectrometry, *Atmos. Meas. Tech.*, 3(1), 301–310, doi:10.5194/amt-3-301-2010, 2010.
- 25 Rosenberg, P. D., Dean, A. R., Williams, P. I., Dorsey, J. R., Minikin, A., Pickering, M. A. and Petzold, A.: Particle sizing calibration with refractive index correction for light scattering optical particle counters and impacts upon PCASP and CDP data collected during the Fennec campaign, *Atmos. Meas. Tech.*, 5(5), 1147–1163, doi:10.5194/amt-5-1147-2012, 2012.

- Schmale, J., Schneider, J., Nemitz, E., Tang, Y. S., Dragosits, U., Blackall, T. D., Trathan, P. N., Phillips, G. J., Sutton, M. and Braban, C. F.: Sub-Antarctic marine aerosol: dominant contributions from biogenic sources, *Atmos. Chem. Phys.*, 13(17), 8669–8694, doi:10.5194/acp-13-8669-2013, 2013.
- Schwarz, J. P., Spackman, J. R., Gao, R. S., Perring, A. E., Cross, E., Onasch, T. B., Ahern, A., Wrobel, W., Davidovits, P., Olfert, J., Dubey, M. K., Mazzoleni, C. and Fahey, D. W.: The Detection Efficiency of the Single Particle Soot Photometer, *Aerosol Sci. Technol.*, 44(8), 612–628, doi:10.1080/02786826.2010.481298, 2010.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics - From Air Pollution to Climate Change* (2nd Edition), 1998.
- Spracklen, D. V. and Heald, C. L.: The contribution of fungal spores and bacteria to regional and global aerosol number and ice nucleation immersion freezing rates, *Atmos. Chem. Phys.*, 14(17), 9051–9059, doi:10.5194/acp-14-9051-2014, 2014.
- Sutton, M. A., Tang, Y. S., Dragosits, U., Fournier, N., Dore, A. J., Smith, R. I., Weston, K. J. and Fowler, D.: A spatial analysis of atmospheric ammonia and ammonium in the U.K., *ScientificWorldJournal.*, 1 Suppl 2, 275–86, doi:10.1100/tsw.2001.313, 2001.
- Taylor, J. W., Allan, J. D., Allen, G., Coe, H., Williams, P. I., Flynn, M. J., Le Breton, M., Muller, J. B. A., Percival, C. J., Oram, D., Forster, G., Lee, J. D., Rickard, A. R., Parrington, M. and Palmer, P. I.: Size-dependent wet removal of black carbon in Canadian biomass burning plumes, *Atmos. Chem. Phys.*, 14(24), 13755–13771, doi:10.5194/acp-14-13755-2014, 2014.
- Taylor, J. W., Choularton, T. W., Blyth, A. M., Liu, Z., Bower, K. N., Crosier, J., Gallagher, M. W., Williams, P. I., Dorsey, J. R., Flynn, M. J., Bennett, L. J., Huang, Y., French, J., Korolev, A. and Brown, P. R. A.: Observations of cloud microphysics and ice formation during COPE, *Atmos. Chem. Phys.*, 16(2), 799–826, doi:10.5194/acp-16-799-2016, 2016.
- Tobo, Y., Prenni, A. J., DeMott, P. J., Huffman, J. A., McCluskey, C. S., Tian, G., Pöhlker, C., Pöschl, U. and Kreidenweis, S. M.: Biological aerosol particles as a key determinant of ice nuclei populations in a forest ecosystem, *J. Geophys. Res. Atmos.*, 118(17), 10,100–10,110, doi:10.1002/jgrd.50801, 2013.
- Topping, D., Connolly, P. and McFiggans, G.: Cloud droplet number enhanced by co-condensation of organic vapours, *Nat. Geosci.*, 6(6), 443–446, doi:10.1038/ngeo1809, 2013.
- Trembath, J.: *Airborne CCN measurements*, University of Manchester., 2013.
- Trembath, J., Bart, M. and Brooke, J.: Efficiencies of Modified Rosemount Housings for sampling Aerosol on a Fast Atmospheric Research Aircraft., FAAM Tech. Note [online] Available from: http://www.faam.ac.uk/index.php/component/docman/doc_download/1673-inlet-efficiency (Accessed 24 July 2014), 2012.

- Wang, S. C. and Flagan, R. C.: Scanning Electrical Mobility Spectrometer, *Aerosol Sci. Technol.*, 13(2), 230–240, 1990.
- Warren, R. A., Kirshbaum, D. J., Plant, R. S. and Lean, H. W.: A “Boscastle-type” quasi-stationary convective system over the UK Southwest Peninsula, *Q. J. R. Meteorol. Soc.*, 140(678), 240–257, doi:10.1002/qj.2124, 2014.
- 5 Wilson, T. W., Ladino, L. A., Alpert, P. A., Breckels, M. N., Brooks, I. M., Browse, J., Burrows, S. M., Carslaw, K. S., Huffman, J. A., Judd, C., Kilhau, W. P., Mason, R. H., McFiggans, G., Miller, L. A., Nájera, J. J., Polishchuk, E., Rae, S., Schiller, C. L., Si, M., Temprado, J. V., Whale, T. F., Wong, J. P. S., Wurl, O., Yakobi-Hancock, J. D., Abbatt, J. P. D., Aller, J. Y., Bertram, A. K., Knopf, D. A. and Murray, B. J.: A marine biogenic source of atmospheric ice-nucleating particles, *Nature*, 525(7568), 234–238, doi:10.1038/nature14986, 2015.
- 10 Young, G., Jones, H. M., Darbyshire, E., Baustian, K. J., McQuaid, J. B., Bower, K. N., Connolly, P. J., Gallagher, M. W. and Choularton, T. W.: Size-segregated compositional analysis of aerosol particles collected in the European Arctic during the ACCACIA campaign, *Atmos. Chem. Phys.*, 16(6), 4063–4079, doi:10.5194/acp-16-4063-2016, 2016.
- Zorn, S. R.: Chemical composition measurements of pristine aerosols in the southern atlantic and amazonian regions by means of on-line time-of-flight aerosol mass spectrometry, Johannes Gutenberg-University Mainz., 2009.

Table 1- Mean aerosol properties measured in NW Coast and inland on each of the case study flights. Where quoted, uncertainty values are standard deviations. All measurements are corrected to standard temperature (273.15 K) and pressure (1013.25 hPa). On 18 July, the CCN column 2 was set to 0.6% SST and 0.9% for all other flying days. Standard deviations are listed for the CCN and CN data, as they are representative of real variations that are much larger than the instrument uncertainty.

Date	Flight number	Run altitude (m)	Run length (min)	ss-SO ₄ (µg m ⁻³)	Org (µg m ⁻³)	MSA (µg m ⁻³)	NO ₃ (µg m ⁻³)	NH ₄ (µg m ⁻³)	BC (ng m ⁻³)	CCN SST (cm ⁻³)	0.1% CCN Column 2 (cm ⁻³)	CN 2 (cm ⁻³)	N _{GA} (cm ⁻¹)	N _{UGA}
NW Coast														
05 July	2013 B786	330±140	12	2.19	1.27	0.10	0.11	0.11	12	-	-	-	1.3	3
18 July	2013 B788	612±14	9	5.0	11.88	0.32	5.15	3.55	530	1540±150	2860±110	5030±220	0.71	3
25 July	2013 B789	388±16	9	2.34	0.55	0.13	0.06	0.16	10	230±21	500±100	990±150	0.83	3
28 July	2013 B790	382±13	8	1.31	0.96	0.04	0.17	0.48	26	-	-	-	1.4	4
29 July	2013 B791	396±22	8	1.48	0.52	0.05	0.07	0.28	18	168±25	440±140	1400±400	1.6	12
03 Aug	2013 B792	405±23	8	1.73	0.32	0.09	0.07	0.18	7	146±9	450±120	-	1.2	6
Inland														
05 July	2013 B786	540±80	64	1.56	2.62	0.18	0.71	0.8	97	-	-	-	1.60	4
18 July	2013 B788	615±11	20	4.0	10.13	0.28	4.55	3.28	470	1230±120	2740±160	7200±800	0.75	3
25 July	2013 B789	570±70	39	1.1	0.67	0.10	0.17	0.67	32	278±27	500±50	8000±6000	0.7	3
28 July	2013 B790	520±30	18	1.1	0.97	0.04	0.23	0.5	30	-	-	-	0.9	4
29 July	2013 B791	530±40	20	1.4	0.54	0.05	0.1	0.39	31	164±49	410±120	9000±5000	1.1	4
03 Aug	2013 B792	550±40	20	1.2	0.42	0.08	0.09	0.49	30	212±33	440±70	7000±2000	0.7	3

Formatted Table

Table 2 - Uncentred R (also known as normalised dot product) between marine OA measured on the SWW case studies and several literature spectra. m/z 28 was not used for the correlations due to gas-phase interference.

Uncentred R						
Date	Sea spray (Ovadnevaite et al., 2011)	Marine biogenic SOA (Chang et al., 2011)	Organic factor (Chang et al., 2011)	HOA (Morgan et al., 2010)	OOA-1 (Morgan et al., 2010)	OOA-2 (Morgan et al., 2010)
05 July	0.99	0.84	0.99	0.47	0.81	0.83
25 July	0.97	0.89	0.97	0.51	0.78	0.83
29 July	0.99	0.79	0.98	0.38	0.83	0.79
03 August	0.95	0.85	0.96	0.46	0.79	0.79

5


Table 3 – Comparison of the m/z 30/46 ratios measured in the AMS aboard the BAe-146.

Date	(m/z 30) / (m/z 46) ratio		
	(NH ₄)NO ₃ calibration	NW Coast	Inland
05 July	1.77	6.33	2.78
18 July	1.65	1.67	1.66
25 July	1.48	6.61	3.05
28 July	1.51	4.40	3.14
29 July	1.37	7.84	6.15
03 August	1.33	8.68	4.01

Table 4 - Mean diameter of distinct aerosol modes measured over the NW coast and inland on the different cast study flights.

<u>Mean diameter (µm)</u>			
<u>Date</u>	<u>Aitken mode</u>	<u>Accumulation mode</u>	<u>Coarse mode</u>
<u>NW Coast</u>			
<u>05 July</u>	<u>0.048</u>	<u>0.188</u>	<u>3.92</u>
<u>18 July</u>	<u>N/A^a</u>	<u>0.125</u>	<u>3.25</u>
<u>25 July</u>	<u>0.060</u>	<u>0.184</u>	<u>4.18</u>
<u>28 July</u>	<u>0.049</u>	<u>0.153</u>	<u>3.04</u>
<u>29 July</u>	<u>0.041</u>	<u>0.155</u>	<u>4.20</u>
<u>03 August</u>	<u>0.043</u>	<u>0.158</u>	<u>4.00</u>
<u>Inland</u>			
<u>05 July</u>	<u>N/A^a</u>	<u>N/A^a</u>	<u>3.71</u>
<u>18 July</u>	<u>N/A^a</u>	<u>0.114</u>	<u>3.13</u>
<u>25 July</u>	<u>0.055</u>	<u>0.169</u>	<u>3.76</u>
<u>28 July</u>	<u>0.051</u>	<u>0.145</u>	<u>3.56</u>
<u>29 July</u>	<u>N/A^a</u>	<u>0.151</u>	<u>3.61</u>
<u>03 August</u>	<u>0.057</u>	<u>0.155</u>	<u>3.87</u>

^aMode not distinct from surrounding distribution



5

Figure 1 - 5-day HYSPLIT back trajectories initiated during the boundary layer aerosol runs. Panels (a) – (f) show the back trajectories on different flying days. Individual trajectories are coloured by altitude. The black lines show the mean trajectories, with markers every 24 hours before sampling. The two gold lines in panel (f) (which lie approximately on top of each other on the plot) are the two trajectories released from the ground site at 1100UTC and 1200UTC. The altitude of these two trajectories remained below 200 m in the first 48 hours, and below 1000 m in the 120 hours of the runs.




Figure 2 - Spatial location of the below-cloud aerosol runs on each case study flight. Panels (a) – (f) show the six case study flights. The arrow markers show the measured wind speed and direction. The black square shows the location of the Davidstow ground site.




Figure 3 - Aerosol composition measured over the NW coast during the six case study flights. ~~In this plot only, SO_4 and NH_4 are simply referred to as 'SO₄' and 'NH₄'.~~ The definitions of each chemical species are given in the text. The lower and middle dashed lines show the concentration of NH₄ required to neutralise the SO_4 to ammonium bisulphate and sulphate respectively. The top dashed line shows the concentration of NH₄ required to fully neutralise all measured inorganic components. The reader should consult the text for discussion of the NO₃ measurements.




Figure 4 - As in Fig. 3 but for measurements taken inland over the southwest peninsula of the UK.




Figure 5 – Normalised mass spectrum of marine OA measured over the NW coast on 05 July.




Figure 6 - Relationship between the fraction of OA signal at (m/z) = 44 and 43, in coastal and inland areas. The diagonal dotted lines define the region where ambient OOA measurements tend to fall (Ng et al., 2010). The data points labelled “All marine” are the NW coast data from 05, 25 and 29 July and 03 August, whereas the points labelled “All inland” are from all six flights. The literature values are the same as those in Table 2.




Figure 7 – Submicron aerosol size distributions measured by the SMPS (dashed lines) and PCASP (solid lines) probes aboard the BAe-146. Panels (a) – (f) show the inland and NW coast distributions for each of the case study flights, while panels (g) and (h) show comparisons of all the NW coast and inland distributions respectively. The error bars in panels (a) – (f) show the standard deviations over each measurement period.




Figure 8 – Comparison of particle concentrations measured in marine air (triangles) and inland (squares). Panel (a) shows the CN concentrations, while panel (b) compares the CCN concentrations to the CDNC measured in updrafts $>2 m s^{-1}$ within 500 m of cloud base. The boxes and whiskers show the 25/75th and the maximum/minimum. Panel (c) shows the distribution of vertical wind in cloud penetrations (liquid water content $>0.01 g m^{-3}$) within 500 m of cloud base.




Figure 9 - Scatter plots of measured CCN concentration at 0.1% SST with concentrations of aerosol species. The data shown are from 18, 25 and 29 July, and 03 August, as CCN data were only available on these days. The blue data points were

measured over the NW coast, and the red points were measured inland, as defined in




Figure 2. Data from the SWW case and anthropogenic pollution (18 July) cases are separated due to the different meteorological conditions and greatly differing CCN concentrations.




Figure 10 – As in Fig. 9 but at 0.9% SST. The data shown are from 25 and 29 July, and 03 August only. Data for 18 July are not included as the second CCN column was set to a different SST for this flight.




Figure 11 - Comparison of ~~IN~~INP calculations using different parameterisations. The solid lines show the temperature ranges in which the ~~IN~~INP parameterisations are deemed valid, while the dashed lines are extrapolations. The inland aerosol measurements were used as input for the parameterisations, other for Wilson et al. (2015) where the marine OA concentration was used. The blue and red markers show measured first ice concentrations from Taylor et al. (2015) at different cloud top temperatures.