Reply by the authors to co-editor's comments

Manuscript # acp-2016-844: "An improved parameterisation of ozone dry deposition to the ocean and its impact in a global climate-chemistry model" by Ashok K. Luhar et al.

5

10

20

25

We thank the co-editor, Dr Laurens Ganzeveld, for his comments. They have been particularly useful in addressing some loose ends in the paper. In the following, we provide a response to his comments. The reference to the changes made in the manuscript points to those in the 'Track Changes' version of the paper.

Comment: "I already wanted to provide this editors comment earlier so that you could have potentially included some of these in your response to the reviewer (editors) comments and for the revision of your paper. Unfortunately due to the workload this further feedback just comes after you have provided your

15 response and revision.

Some of these comments from my side also result from some interesting discussions I have had last week visiting some colleagues who also work on ocean-atmosphere exchange processes, in particular O3 and still want to introduce some of the points that come out of those discussions but also want to provide feedback on a couple of issues that already came across reading your ms after its initial submission and add up to the comments provided by the two reviewers."

Response: No problem. Thanks.

Changes in manuscript: None.

Comment: "Pp2; line 30: it is Wesely"

Changes in manuscript: Corrected.

Comment: "Pp3; line 11: I would leave out here the reference to Ganzeveld et al. 2009 since there we actually introduced the more mechanistic model. The one study that was discussed in that paper was the Kuhlmann et al., 2003 reference."

Changes in manuscript: The reference to Ganzeveld et al. (2009) has been removed.

Comment: "Pp4, line 25 and beyond: "but a detailed testing against deposition data was not reported, presumably because of the lack of suitable data at the time"

I don't agree with that statement. In the 2009 study we actually show a comparison of the simulated VdO3 and O3 dry deposition fluxes for the measurements that were available that time (Table 2 in Ganzeveld et al., 2009) and where we also evaluated how the comparison improved including the

potential role of DOM-O3 chemistry. You are correct that we couldn't include yet all the results reported in the Helmig et al. 2012 paper that you have applied in your paper but so might like to revise this statement."

Changes in manuscript: We agree with the co-editor. The last paragraph on Page 4 has been modified to incorporate his suggestion.

5 Comment: "Pp6: lne 21: "The deposition velocity is uniformly distributed to all model levels contained within the atmospheric boundary layer". Reading over this statement it interprets that the removal rate in the surface layer is smaller than the overall deposition rate or is the same deposition velocity used in all layers representing the boundary layer? and is this approach applied to potentially correct for issues on the representation of vertical transport?"

Response: Our statement could have been written a bit better. The way the present version of UKCA is setup, a first order loss rate is calculated as the deposition velocity divided by the height of the atmospheric boundary layer (ABL), and this loss rate is applied at each model level contained within the ABL. A grid-box mean loss rate at each model level within the ABL is then obtained by weighting the individual loss rates with the fractions of the surface types present in the grid box. A more appropriate method would be to divide the deposition velocity by the height of the lowest model level (or by the height of the surface layer) and apply the loss rate only at that height. According to the information provided by the U.K. Met Office, who are a UKCA developer, there are legacy reasons as to why the first method of loss calculation is used, and these are in no way related to any potential issues to do with the representation of vertical transport. The U.K. Met Office has tested a version of UKCA that has dry deposition loss done from the lowermost model level only and their results show that compared to the default case where deposition is applied within the full ABL, deposition fluxes change only marginally, if at all, whereas near-surface concentrations of ozone improve somewhat in comparison to measurements. We plan to explore this point further in the future.

Changes in manuscript: The last paragraph on Page 6 has been modified to better explain how the deposition velocity is used to determine the corresponding loss rate in the model (lines 25–28 on Page 6).

10

Comment: "Pp 12, line 26: "This is not as..."

Changes in manuscript: Corrected.

Comment: "Pp 16: lines 25 and beyond; in this short discussion on potential issues involved in the role of the reactivity at this point, but actually also further throughout the paper, you only elude on the role
15 of the Iodide whereas the total reactivity might also be strongly dependent on for example some of the DOM-O3 chemistry as discussed in more detail in the Ganzeveld et al., 2009 paper. In the revised paper you now mention explicitly that you have excluded in your study the role of organic compounds in your study. Still I think it would be worthwhile to shortly discuss how inclusion of this feature could further

affect the presented modelling results also in comparison to the 1-layer scheme. This chemistry could be especially relevant in the coastal zones but also in regions of strong upwelling with ocean productivity, in some of the colder waters of the Arctic but also in the southern oceans with cold conditions and where your analysis mainly makes a point about seeing there the largest differences between the 1- and

2 layer approach." 5

> **Response:** We provide a short discussion on this in the revised paper, incorporating a comment on relevant findings by Chang et al. (2004) and Ganzeveld et al. (2009).

> **Changes in manuscript:** The 2nd paragraph on Page 17 has been modified and new material added (lines 3–19 on Page 17).

Comment: "Pp16: I noticed that you removed the statement indicating that in the 2009 study we applied the nitrate as a proxy to indeed infer the Iodide concentrations (also since by that time the SST dependence was not yet presented). It seems that you have done so also because of the demonstrated

- stronger correlation between SST and Iodide. However, I think that it is actually important to explicitly 10 mention this essential difference in the inversion of the Iodide fields comparing the deposition schemes and modelling results. I also agree with one of the reviewers suggestion that it would have been interesting to potentially compare the impact of the differences in inferring Iodide (also including a run with the 2-layer model with Iodide based on nitrate) also because, despite the apparent stronger
- correlation between SST and Iodide, this feature of getting realistic oceanic biogeochemical boundary 15 conditions for these deposition calculations seems to be a remaining essential challenge also due to limited observations."

Response: The statement indicating that Ganzeveld et al. (2009) used oceanic surface nitrate concentration as a proxy for iodide concentration has been reinstated. We need more observations to relate iodide with other parameters of the ocean biogeochemistry (such as nitrate). Chance et al. (2014) provide a good account and analysis of available data to examine iodide dependencies, and perhaps these data could be further analysed for better parameterisations. Of course, we also need more comprehensive deposition velocity measurements so that the impact of changes in iodide parameterisations as well as those of any additional chemical reactions could be discerned and quantified.

Changes in manuscript: The sentence "Ganzeveld et al. (2009) used oceanic surface nitrate concentration as a proxy for iodide concentration" is added subsequent to Eq. (11). In addition, a sentence in the last paragraph of Conclusions added (lines 18–20, Page 22).

Comment: "The following comment also relates to one of the comments made by reviewer #2. Referring to your discussion on the global and oceanic deposition budget. I noticed that with your 20 global model simulations with an oceanic ozone deposition sink which is 12% of the global sink, you arrive at a flux of about 80 TgO3 yr-1. You discuss that this for the 1-layer model it would be about 19% and for the fixed rc approach about ~21% or so. This would imply a global oceanic O3 deposition

budget of 180 TgO3 yr-1 which is substantially smaller than the 300 TgO3 yr-1 global oceanic O3 deposition term presented by Ganzeveld et al. (2009). Is raises a question how to appreciate these differences between the 1- and 2- layer oceanic O3 deposition model recognizing already the fact that there seems to be like a ~factor 2 difference among such global chemistry-climate models inferring this

5 ozone budget term. It also comes back to my remark about the deposition approach in your model using the vd uniformely distributed over all boundary layers. Rather than evaluating the O3 dry deposition velocities, it would be useful to also evaluate the O3 dry deposition fluxes by comparison with the observations. Have you conducted such an evaluation??"

Response: The figures mentioned by the co-editor are correct. There is quite a range of modelled total ozone dry deposition reported in the literature, and some of these ranges are mentioned in the 1st paragraph of Section 7.2. The range cited by Ganzeveld et al. (2009) is 600–1000 Tg yr-1. As mentioned in our paper, our deposition budget values are on the lower end of the values reported based on other models. We think the main reason for that is that UKCA underestimates tropospheric ozone concentrations, which results in an underestimation of the ozone dry deposition flux at the surface. However, note that the calculated deposition velocity is not influenced by the predicted ozone concentration, and hence the relative (not absolute) dry deposition budget contributions given in our paper in terms of percentages for the various deposition schemes should remain more or less the same even if the model was predicting the ozone concentrations fine. The two-layer scheme agrees the best with the Helmig et al. deposition velocity data, and we emphasize the relative change in oceanic dry deposition when this scheme is used. A model that predicts tropospheric ozone correctly and includes a correct description of deposition velocity should yield a realistic value of the total deposition budget. We agree that it would also be useful to evaluate the ozone dry deposition flux, but for that to be meaningful we first need to fix the issue of the underestimation of tropospheric ozone concentration in the model because such a flux is proportional to the concentration given that the deposition velocity would change little from the present values. In this regard, we think that our approach of using deposition velocity uniformly distributed within the boundary layer is not a problem (see our response to the related comment above).

Changes in manuscript: At the end of 1st paragraph of Section 7.2 we add "An underestimation of ozone would result in an underestimation of the ozone dry deposition flux near the surface. However, the calculated deposition velocity is not influenced by the predicted ozone concentration, so the relative dry deposition budget contributions given above in terms of percentages should remain representative." In addition, a sentence in the 1th paragraph (lines 7–9) of Page 22 added.

An improved parameterisation of ozone dry deposition to the ocean and its impact in a global climate-chemistry model

Ashok K. Luhar¹, Ian E. Galbally¹, Matthew T. Woodhouse¹, Marcus Thatcher¹

¹CSIRO Oceans and Atmosphere, Aspendale, 3195, Australia

5 Correspondence to: Ashok K. Luhar (ashok.luhar@csiro.au)

Abstract. Schemes used to parameterise ozone dry deposition velocity at the oceanic surface mainly differ in terms of how the dominant term of surface resistance is parameterised. We examine three such schemes and test them in a global climate-chemistry model that incorporates meteorological nudging and monthly-varying reactive-gas emissions. The default scheme invokes the commonly used assumption that the water surface resistance is constant. The other two schemes, named the one-

- 10 layer and two-layer reactivity schemes, include the simultaneous influence on the water surface resistance of ozone solubility in water, waterside molecular diffusion and turbulent transfer, and a first-order chemical reaction of ozone with dissolved iodide. Unlike the one-layer scheme, the two-layer scheme can indirectly control the degree of interaction between chemical reaction and turbulent transfer through the specification of a surface reactive layer thickness. A comparison is made of the modelled deposition velocity dependencies on sea surface temperature (SST) and wind speed with recently reported cruise
- 15 based observations. The default scheme overestimates the observed deposition velocities by a factor of 2 to 4 when the chemical reaction is slow (e.g. under colder SSTs in the Southern Ocean). The default scheme has almost no temperature, wind-speed and latitudinal variations in contrast with the observations. The one-layer scheme provides noticeably better variations, but it overestimates deposition velocity by a factor of 2 to 3 due to an enhancement of the interaction between chemical reaction and turbulent transfer. The two-layer scheme with a surface reactive layer thickness specification of 2.5
- 20 microns, which is approximately equal to the reacto-diffusive length scale of the ozone-iodide reaction, is able to simulate the field measurements most closely, with respect to absolute values as well as SST and wind-speed dependence. The <u>annual</u> global oceanic deposition of ozone determined using this scheme is approximately 80 Tg yr⁻¹. This is almost half of the original oceanic deposition obtained using the default scheme, and corresponds to a 10% decrease in the original estimate of the total global ozone deposition. The previously reported modelled estimate of oceanic deposition has been roughly one
- 25 third of total deposition and with this new parameterization is reduced to 12% of the modelled total global ozone deposition. Deposition parameterisation influences the predicted atmospheric ozone mixing ratios, especially in the Southern Hemisphere. For the latitudes 45–70°S, the two-layer scheme improves the prediction of ozone observed at an altitude of 1 km by 7% and that within the altitude range 1–6 km by 5% compared to the <u>defaultoriginal</u> scheme.

1 Introduction

Ozone (O_3) is not emitted directly into the atmosphere but is formed in both the stratosphere and troposphere by photochemical reactions involving natural and anthropogenic precursor species. Ozone is an oxidant as well as a precursor to the formation of hydroxyl and hydroperoxyl radicals that play a critical role in the chemical cycles of many trace gases in the

- 5 troposphere. The lifetime of ozone in the troposphere is relatively short, being about 22 days, compared to long-lived and globally well-mixed greenhouse gases such as carbon dioxide (CO₂). Ozone acts as a greenhouse gas, and adversely impacts human health and plant productivity (Monks et al., 2015). As a greenhouse gas, ozone has the third largest global warming effect after CO₂ and methane (CH₄), and thus plays an important role in the Earth's climate system. The tropospheric ozone burden is estimated to have increased by about 40% since preindustrial times as a result of increases in the emissions of
- 10 ozone precursors (Young et al., 2013). According to the Fifth Assessment Report (AR5) of the United Nations Intergovernmental Panel on Climate Change (IPCC), the anthropogenic component of the radiative forcing due to ozone is estimated to be 0.35 ± 0.2 W m⁻², of which 0.40 ± 0.2 W m⁻² is the tropospheric ozone contribution and -0.05 ± 0.1 W m⁻² is the stratospheric contribution. Ozone represents about 15% of the total anthropogenic radiative forcing (including aerosols) estimated for the year 2011 (Climate Change, The Physical Science Basis, 2013, IPCC, AR5, Chapter 8, Anthropogenic and
- 15 Natural Radiative Forcing).

20

Aspects of ozone have been investigated in numerous air quality and climate related studies, including its sources and sinks (e.g., Vingarzan, 2004; Wild, 2007; Stevenson et al., 2006, 2013; Young et al., 2013; Monks et al., 2015). In the troposphere, the ozone budget is determined by transport from the stratosphere, deposition at the Earth's surface, and chemical production and loss. Dry deposition is an important sink of ozone (Galbally and Roy, 1980), which influences near-surface concentration of ozone as well as its lifetime and long range transport. The 'present-day' (about year 2000) total global dry

- deposition of ozone estimated by the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) is of the order of 1094 ± 264 Tg yr⁻¹ which is almost double the flux of ozone from the stratosphere to the troposphere (477 ± 96 Tg yr⁻¹) (Climate Change, The Physical Science Basis, 2013, IPCC, AR5, Chapter 8, Anthropogenic and Natural Radiative Forcing, Table 8.1, page 672; Young et al., 2013). Although the average deposition velocity to the ocean is smaller than that
- 25 to terrestrial surfaces, the larger coverage of the Earth's surface by the oceans leads to considerable deposition to water previous modelling studies estimate that about one third of total ozone deposition is to water (Ganzeveld et al., 2009; Hardacre et al., 2015). Based on results from a multi-model analysis, Hardacre et al. (2015) observed that ozone deposition to the water surface has the largest uncertainty compared to other surface types. The deposition process needs to be properly accounted for when modelling ozone chemistry in the atmosphere, irrespective of whether the application is related to air
- 30 quality or climate chemistry.

Similar to the land surface, the commonly used parameterisation for dry deposition of gases to the ocean surface is to express deposition velocity (v_d), which is the flux of the gas to the surface divided by its concentration in air near the surface, as the linear sum of three resistances (Wes<u>eley</u>, 1989):

$$v_d = \frac{1}{r_a + r_b + r_c},\tag{1}$$

where r_a , r_b and r_c are the resistances exerted to the transfer process in three successive top-to-bottom layers: the aerodynamic resistance r_a is the resistance to mixing by turbulent transport in the atmospheric surface layer, the atmospheric viscous (or quasi laminar) sublayer resistance r_b is the resistance to movement across the thin layer (0.1 – 1 mm) of air that is in direct contact with the surface and not moving with the mean flow of the wind, and the surface

5 mm) of air that is in direct contact with the surface and not moving with the mean flow of the wind, and the surface resistance r_c is the resistance to uptake by the underlying surface that can be controlled by physical, chemical and/or biological processes depending on the surface type and species of interest.

For ozone deposition to oceanic surfaces, r_c is the dominant resistance and several approaches have been proposed to calculate it. A common approach is to use a constant value, normally $r_c = 2000$ s m⁻¹, based on Wesely's (1989) deposition

10 parameterisation. Several global chemical transport models use this approach by default, for example, MATCH-MPIC (von Kuhlmann et al., 2003; Ganzeveld et al., 2009), MESSy (Kerkweg et al., 2006), MOZART-4 (Emmons et al., 2010), CAMchem (Lamarque et al., 2012), GEOS-Chem (Mao et al., 2013), and UKCA (Abraham et al., 2012).

While a constant r_c may be a good first-order estimate of global and long-term average oceanic dry deposition, it does not include any spatial and temporal dependencies of the surface uptake process on any oceanic physical, chemical or biological

15 properties or processes.

The surface ocean acts as a gateway for molecules to enter the atmosphere or ocean medium, and contains complex chemical reactions of inorganic components and dissolved organic matter. Some of these reactions are important sources or sinks of climatically active trace gases. For example, ozone is known to react with a number of dissolved chemical compounds present in the seawater, with the reaction with dissolved iodide being by far the fastest one (Garland et al., 1980; Chang et

20 al., 2004) which needs to be considered. Apart from providing more accurate ozone predictions, a proper treatment of deposition is also important from the point of view of better accounting for feedback cycles; for example the ozone reaction with iodide in the seawater produces volatile iodine compounds, which may then pass into the atmosphere and participate in catalytic ozone destruction cycles in the marine atmosphere (Carpenter et al., 2013).

For the purpose of clarity we describe the layers of the ocean from the surface down as follows. The top 1 µm to 1 mm of the sea surface is termed the sea surface microlayer (SSM) (Carpenter et al., 2015). The physical and chemical properties of the SSM can be different to those of the bulk water. The SSM may consist of various sublayers or scales depending on the physical or chemical properties being considered. The upper part of the SSM is a region where turbulent mixing is drastically reduced and molecular diffusion dominates, resulting in strong gradients in gas concentrations and other properties. A reacto-diffusive length scale can be defined as $(D/a)^{1/2}$ (Carpenter et al., 2013), where *D* is the molecular diffusivity $(m^2 s^{-1})$ and *a* is the reaction rate constant (s^{-1}) , and is typically 3 µm (at ~ 25°C) for the ozone-iodide reaction in seawater. A molecular (or diffusive) sublayer can be defined as the region where ozone molecular diffusivity is greater than turbulent transfer, and is typically 50 µm thick. A viscous (or quasi laminar) sublayer is where viscous processes effectively dissipate the turbulent energy, and is of the order of 1 mm (Fairall et al., 2000). In the surface turbulent layer (~ 10–50 m) below the

5

10

2 Mechanistic approaches to determining rc

viscous sublayer, turbulent processes dominate.

Most early studies on ozone deposition to seawater either explored its dependence on dynamical factors related to turbulent transfer without considering any chemical reaction (e.g. Galbally and Roy, 1980). Other studies have considered molecular diffusion coupled with chemical reactions (e.g. with iodide) in the seawater without including any effects of turbulent transfer (e.g. Garland et al., 1980). Using the assumption of Wanninkhof (1992) that the enhancement of deposition due to chemical processes was additive to the turbulent transfer in water, Chang et al. (2004) formulated $1/r_c$ as a linear sum of two independent terms with the correct asymptotic behaviours: the first term representing the influence of molecular

diffusion coupled with chemical reaction (Garland et al., 1980) and the second term representing the influence of wind 15 induced turbulent transfer coupled with a chemical enhancement factor (Liss and Merlivat, 1986). More recently, by solving a simplified form of the budget equation for mass conservation that include turbulent and molecular transport and a chemical reaction term, Fairall et al. (2007) derived two formulations for r_c that are able to account for the simultaneous effects of oceanic physical and chemical processes (i.e. ozone solubility, molecular diffusion,

turbulent transfer and chemical reaction). Their approach signifies a development based on fundamental conservations laws.

- 20 Some of the processes affecting deposition act in opposite directions and some are interlinked (e.g. ozone solubility in water and reaction rate coefficient are a function of sea surface temperature (SST) but in opposite way). Thus, a proper way to evaluate Consequently, it has been difficult to properly test a mechanistic deposition scheme against field measurements is without to coupleing it to an atmospheric composition model and then analysetesting the response of the full coupled system that represents the conditions of the measurements. To our knowledge, there has not previously been such a detailed testing
- 25 in light of newer deposition observations and better modelling capabilities that could lead to potential improvements in deposition parameterisation. Ganzeveld et al. (2009) explored the response of a free-running global climate-chemistry model to the choice of a constant r_c and the one-layer reactivity parameterisation derived by Fairall et al. (2007) (described later in Section 5) and compared the modelled ozone deposition velocity with observations reported in the literature. but a detailed testing against deposition data was not reported, presumably because of the lack of suitable data at the time. Since then there
- 30 have been new deposition velocity data reported by Helmig et al. (2012), which together with the inclusion of atmospheric

model features such as meteorological nudging, could lead to potential improvements in parameterisation of deposition velocity.

Recently-Helmig et al. (2012) presented the first ship-borne open-ocean ozone flux measurements and at the same time the most extensive set of these measurements (1700 hours of observations) probably surpassing all previous data. The measurements covered the Gulf of Mexico, the southern and northern Atlantic, the Southern Ocean, and the eastern Pacific Ocean. These experiments gave medians of the oceanic ozone deposition velocity (v_d) from the five cruises of 0.009 to 0.034 cm s⁻¹. The measurements cover the range of 45°N to 50°S and show little wind-speed dependence but a marked sea surface temperature dependence. This dataset provides a unique opportunity to both test oceanic ozone deposition schemes and re-evaluate the ozone deposition rate to the world's oceans.

- 10 The aims of this paper are: (a) to examine schemes of ozone dry deposition to the ocean involving: the default constant r_c assumption, the one-layer reactivity model suggested by Fairall et al. (2007) and a development of their two-layer reactivity model, within a global climate-chemistry model, ACCESS-UKCA, incorporating meteorological nudging and monthly-varying reactive-gas emissions, and to compare the results with the ozone deposition data of Helmig et al. (2012). This comparison enables selection of an improved deposition formulation. Then (b) using the best fitting scheme, re-evaluate the
- 15 rate of ozone deposition over the ocean and for the combined ocean-land system, and finally (c) with the best fitting scheme, examine its influence on the comparison of modelled ozone compared with global ozone profile observations covering the troposphere.

3 ACCESS-UKCA chemistry-climate model

- The Australian Community Climate and Earth-System Simulator (ACCESS, see Bi et al., 2013) has been developed for both climate and numerical weather prediction purposes. The physical atmosphere component of ACCESS is the UK Met Office's Unified Model (MetUM). The UK Chemistry and Aerosol (UKCA, <u>http://www.ukca.ac.uk</u>, described below) atmospheric composition module (at UM vn8.4; see Abraham et al., 2012) is part of ACCESS, and the resulting model is referred to here as ACCESS-UKCA.
- Several different chemical schemes are available within UKCA. The configuration used here (at UM vn8.4) combines the tropospheric chemistry scheme described by O'Connor et al. (2014) and the stratospheric chemistry as described by Morgenstern et al. (2009). The resulting chemistry configuration is known as Chemistry of the Stratosphere and Troposphere (CheST). The tropospheric chemistry scheme accounts for O_x, HO_x, NO_x, CH₄ and other volatile organic carbon species (e.g. isoprene). The stratospheric chemistry scheme includes chlorine and bromine chemistry (for ozone depleting substances, ODS), and heterogeneous polar stratospheric cloud chemistry suitable for simulating stratospheric ozone. The Fast-JX
- 30 photolysis scheme (Neu et al., 2007) is used in the CheST configuration, and is applicable both in the troposphere and



stratosphere. For aerosols, the size-resolved aerosol microphysics scheme GLOMAP-mode, which includes sulfur chemistry, is used (Mann et al., 2010). The total number of reactions is 306 across 86 species. Interactive gas-phase dry and wet deposition processes are included. The chemical reactions together with relevant deposition rates are solved within the ASAD (A Self-contained Atmospheric chemistry coDe) framework.

- 5 UKCA is coupled to a radiation scheme (via O₃, CH₄, N₂O, and aerosol direct and indirect effects). Changes in atmospheric composition are thus allowed to impact on model physics, allowing the representation of feedbacks from chemical composition to atmospheric dynamics when operating in a free-running (un-nudged) climate configuration. ACCESS also has a coupled ocean model, but the present configuration used an atmosphere-only set up forced by observed sea surface temperature and sea ice fields.
- 10 The atmospheric model domain is global with 85 levels extending from the surface to approximately 85 km, and the horizontal resolution was 1.875° in longitude and 1.25° in latitude (the so called N96L85 configuration).

In order for a free-running global climate model to realistically reproduce the state of the atmosphere, especially for process studies at short timescales and subsequent comparison with measured data, meteorological nudging is often used. Nudging is a simple data assimilation technique that uses meteorological reanalyses data to relax dynamical variables of a model

15 towards the observed state of the atmosphere at a given time, thus minimizing meteorology as a source of uncertainty in the modelled fields. The variables nudged are horizontal wind components and potential temperature in the free troposphere by using the ERA-Interim reanalyses on pressure levels (Telford et al., 2008; Uhe and Thatcher, 2015).

A global monthly-varying emissions database for reactive gases and aerosols that includes both anthropogenic and natural components was compiled for ACCESS-UKCA and is described in detail by Woodhouse et al. (2015), with the exception

20 that here we used GFED4s (including small fires) biomass burning emissions data instead of the original ACCMIP database, the former includes inter-annual variability. For methane, nitrous oxide and ozone depleting substances, concentrations are prescribed instead of emissions.

The model has nine surface types (namely broad-leaf trees, needle-leaf trees, C3 and C4 grass, shrub, urban, water, bare soil, and land ice) and for a particular grid box the three resistances are calculated for each surface type and a corresponding

- 25 deposition velocity is then computed. <u>A first order loss rate is calculated as the deposition velocity divided by the height of</u> <u>the atmospheric boundary layer (ABL), and this loss rate is applied at The deposition velocity is uniformly distributed to</u> <u>eachall</u> model levels contained within the <u>ABLatmospheric boundary layer</u>. <u>A first order loss rate is calculated</u> <u>corresponding to each deposition velocity, and aA</u> grid-box mean loss rate <u>at each model level within the ABL</u> is <u>then</u> obtained by weighting the individual loss rates with the fractions of the surface types present in the grid box. The mean loss
- 30 rate is used in the species mass conservation equation. Currently there is only one water surface type in the model, so the same deposition scheme is used for both seawater and freshwater. In our analysis for the oceanic deposition, we do not consider any coastal grid cells and only consider those grid cells that are 100% covered by the water surface type.

3.1 Current scheme for ozone deposition to water

In ACCESS-UKCA, the aerodynamic resistance r_a is a function of sea surface roughness, wind speed and atmospheric stability, and is determined as

$$r_a = \frac{\ln(z_r / z_0) - \psi}{\kappa u_*}, \qquad (2)$$

where u_* is the friction velocity, z_r is a reference height (= 50 m), κ is the von Karman constant (= 0.4) and ψ is a stability 5 correction function. The aerodynamic roughness length z_0 for the sea surface is parameterised as the sum of viscous and gravity wave parts (Smith, 1988):

$$z_0 = \frac{0.11 \, v_a}{u_*} + \frac{a_c \, u_*^2}{g},\tag{3}$$

where the kinematic viscosity of air v_a a function of surface temperature and air density, a_c is a Charnock coefficient (= 0.016), and *g* is the acceleration due to gravity.

The resistance r_b is related to molecular properties of the gas as well as atmospheric turbulence through the following 10 relationship (Hicks et al., 1987):

$$r_b = \left(\frac{S_c}{P_r}\right)^{2/3} \frac{1}{\kappa \, u_*},\tag{4}$$

where S_c is the Schmidt number defined as the ratio of the kinematic viscosity of air to molecular diffusivity of the gas species in air and P_r is the Prandtl number of air (i.e. the ratio of momentum diffusivity to thermal diffusivity ≈ 0.72). The molecular diffusivity of ozone in air is taken as 1.4×10^{-5} m² s⁻¹.

The present version of ACCESS-UKCA uses $r_c = 2200$ s m⁻¹, which yields a ceiling value for v_d of $1/r_c = 0.0455$ 15 cm s⁻¹. This will be the first (or default) model specification for ozone deposition in our study.

4 Ozone loss in seawater

Garland et al. (1980) introduced the idea that ozone loss in the ocean was dominated by its reaction with dissolved iodide ions ($O_3 + I^- \rightarrow$ products) within the seawater. This bimolecular reaction is considered as a pseudo first order in several models of ozone loss in seawater because only a very small proportion of iodide is used and its concentration remains almost constant. The first-order rate coefficient a (s⁻¹) is equal to the pertinent second-order rate coefficient (k) times the iodide concentration $[I^-]$, i.e.

$$a = k.[I^-], \tag{5}$$

where $[I^-]$ is in mole per litre (or molar, M) and k is in M⁻¹ s⁻¹. If there are other first order reactions, then the total reaction rate can be determined as $a = \sum_{i} k_i C_i$, where k_i and C_i are the first order rate coefficient and concentration of the *i*th

5 species.

Eq. (5) represents the integrated reactivity given for the bulk reaction ($O_3 + I^- \rightarrow$ products) and is used in this model to calculate the ozone removal by reaction with iodide. However, it is noted that the chemistry of ozone reaction with iodide in aqueous solution is complex (e.g. Sakamoto et al., 2009). There can also be complicating factors including reactions of ozone with other halogens in seawater (Sarwar et al., 2015), the modification of bulk properties of seawater in the surface

10 microlayer, and the presence of organic compounds in seawater including those preferentially concentrated in the sea surface microlayer (Carpenter et al., 2015). These are not addressed here.

The second-order rate coefficient (k) in Eq. (5) is derived from the data of Magi et al. (1997) (see Section 5.1).

5 An alternative scheme for r_c in ACCESS-UKCA

Atmospheric ozone in passing into the ocean is transferred by molecular diffusion through the sea surface microlayer

15 (including viscous sublayer) and then by turbulent processes in the surface turbulent layer. At the same time ozone can be lost by chemical reaction.

Considering current knowledge, assuming that iodide does provide the major sink for ozone in the ocean, the resistance to ozone deposition by the ocean surface will be an unknown function of a number of known variables:

$$r_{c} = f([I^{-}], k, T_{s}, \alpha, D, u_{*_{w}}),$$
(6)

20

where α is the dimensionless solubility of ozone in water (which is the ratio of the aqueous-phase ozone concentration to its gas-phase concentration and is related to Henry's law coefficient), T_s is the water temperature, D is the molecular diffusivity of ozone in seawater, and u_{*w} is the waterside friction velocity. r_c is expressed as (Liss and Merlivat, 1986):

$$r_c = \frac{1}{\alpha v_{dw}},\tag{7}$$

where v_{dw} is the waterside deposition velocity of ozone. u_{*w} is calculated as $u_{*w} = (\rho_a / \rho_w)^{1/2} u_*$ by assuming that the atmospheric surface stress is equal to the waterside surface stress, where ρ_a is the air density and ρ_w is the water density.

Applying horizontal homogeneity and stationarity to the fundamental equation for the conservation of mass of a reacting and diffusing substance in water yields the following equation in the vertical dimension (z, i.e. depth from the water surface) (Fairall et al., 2007)

$$\frac{\partial}{\partial z} \left[(D + K_i) \frac{\partial C}{\partial z} \right] - a C = 0, \qquad (8)$$

Where *C* is concentration and K_t is the turbulent diffusivity. By assuming $K_t = \kappa u_{*_w} z$, Fairall et al. (2007) solved Eq. (8) to obtain

$$v_{dw} = (a D)^{1/2} \left[\frac{K_1(\xi_0)}{K_0(\xi_0)} \right],$$
(9)

where $K_n(\xi)$ is modified Bessel functions of order *n*, $\xi_0 = b(aD)^{1/2}$ and $b = 2/(\kappa u_{*_w})$. Eq. (9) explicitly includes 10 the role of waterside molecular diffusion, turbulent transfer and first-order chemical reaction. In the limit of fast reaction or negligible turbulence, it reduces to

$$v_{dw} = (a D)^{1/2}, (10)$$

which is the form proposed by Garland et al. (1980). The above scheme forms part of the COAREG group of gas transfer algorithms (Fairall et al., 2011).

There are four controlling parameters involved in the above scheme for r_c (Eqs. (7) and (9)): ozone solubility, molecular diffusion, turbulent transfer and chemical reaction. Deposition velocity increases as these parameters attain larger values. This scheme is termed the one-layer reactivity scheme due to the assumption of uniform reactivity (*a*) through the ocean depth. We incorporated the above scheme for r_c in ACCESS-UKCA. This will be the second model specification for ozone deposition.

5.1 Parameterisation of input quantities

5

20 The second-order rate coefficient k (M⁻¹ s⁻¹) is derived by fitting an Arrhenius-type expression for k as a function of water temperature T_s (K) to the data of Magi et al. (1997):

$$k = \exp\left(\frac{-8772.2}{T_s} + 51.5\right).$$
 (11)

Ganzeveld et al. (2009) used oceanic surface nitrate concentration as a proxy for iodide concentration. Chance et al. (2014) compiled available measurements of sea surface iodide which show highest iodide concentrations in tropical waters. They examined statistical relationships between iodide and parameters such as SST, nitrate, salinity, chlorophyll-a and mixed layer depth, and found that SST (T_s) was the strongest predictor of iodide followed by latitude. MacDonald at al. (2014) used data from several cruises in the Atlantic and Pacific oceans covering the latitudes 50°S to 50°N to derive the following parameterisation for iodide concentration:

$$[I^{-}](nM) = 1.46 \times 10^{15} \exp\left(\frac{-9134}{T_s}\right).$$
(12)

5

10

In their study on iodine's impact on tropospheric oxidants, Sherwen et al. (2016) found that the MacDonald at al. (2014) parameterisation provided better results than the iodide-SST regression parameterisation derived by Chance at al. (2014). We use Eq. (12) in our ACCESS-UKCA modelling, but a sensitivity to the iodide parameterisation derived by Chance at al. (2014) will also be explored later. Based on Eq. (12), the concentrations of iodide lie in the range 6 to 160 nM, and the reaction time-scale a lies in the range 2 to 1350 s⁻¹ for the experimental conditions modelled. Iodide concentrations are highest in warm tropical waters and lowest in cool waters at higher latitudes.

The dimensionless solubility of ozone in water α decreases with water temperature, and is parameterised as (Morris, 1988):

$$\log_{10}(\alpha) = -0.25 - 0.013(T_s - 273.16). \tag{13}$$

The molecular diffusivity D (m² s⁻¹) of ozone in water is expressed as (Johnson and Davis, 1996):

$$D = 1.1 \times 10^{-6} \exp\left(\frac{-1896}{T_s}\right).$$
 (14)

15 We ran ACCESS-UKCA with both the default scheme and the above one-layer reactivity scheme. The modified Bessel functions were determined using the numerical algorithms given by Press et al. (1992). The values of $(a D)^{1/2}$ (see Eq. (10)) increase by a factor of 30 from high latitude cold waters to tropical warm waters in the model calculations.

6 Comparison of observed and modelled ozone deposition velocity to the ocean surface

Helmig et al. (2012) reported the magnitude and variability of ozone deposition velocity measured over the open ocean from 20 a ship-based eddy-covariance ozone flux system during 2006–2008. The experiments were conducted on five cruises, namely: (1) TexAQS06 (July 7 to September 12, 2006), (2) STRATUS06 (October 9–27, 2006), (3) GOMECC07 (July 11 to August 4, 2007), (4) GasEx08 (February 29 to April 11, 2008), and (5) AMMA08 cruises (April 27 to May 18, 2008). The respective areas covered were: (1) North-western Gulf of Mexico, (2) the persistent stratus cloud region off Chile in the eastern Pacific Ocean, (3) the Gulf of Mexico and the US east coast, (4) the Southern Ocean, and (5) the southern and

5 northern Atlantic Ocean. Helmig et al. (2012) present a map of cruise tracks together with bin-averaged deposition velocity data as a function of SST and wind speed for each of the five cruises. Information on the actual timings and locations of the observations along the cruise tracks is not available.

Helmig et al. (2012) plotted the observed v_d values against SST, but in order to compare the data with the v_d values obtained from the offline one-layer reactivity scheme of Fairall et al. (2007) they removed the wind-speed dependence from their data.

- 10 This resulted in an idealized testing of the model scheme, omitting the joint influences of SST and wind speed that occur at the specific field locations. In the present case, this scheme has been coupled to an atmospheric chemistry model in which all input parameters are supplied and act simultaneously in determining v_d for the specific field program locations and durations, so the modelled value and data can be compared directly. We obtained the same v_d -SST data but without the wind speed dependence removed (Ludovic Bariteau, personal communication, 2016) for a consistent comparison with our model results
- 15 (although there is not a large difference between the data with or without the wind-speed dependence).

The ACCESS-UKCA model was run from January 2005 until the end of 2008. The model output was used for 2006 onwards and consists of monthly-averaged values of deposition parameters at each grid point. The simulation months and years were matched with the experimental periods. The model output was extracted at a series of grid-point locations with almost uniform spacings along the tracks of the above experimental cruises, and the modelled values at these points were used for

- 20 comparison with the measurements. It is noted that the deposition velocity data and modelled values cannot be matched exactly in terms of time and location. The cruise data are averaged with respect to SST or wind-speed bins over a given experimental period. They may vary in both space and time (but information on any such variation is not available).. On the other hand, the model values are monthly averages extracted along the cruise tracks, with the model months matched to the months corresponding to the experimental periods, which means the binning is dominated by spatial changes. This matching
- 25 difference is not expected to pose significant problem for comparison purposes because the deposition velocity variations are mainly examined in terms of SST which is not expected to vary greatly with time at a given location for a typical experimental period under consideration, and the ozone concentration in marine air has only a minor diurnal variation (Galbally et al., 2000). (Note, however, that in contrast to oceanic deposition, land-based deposition over vegetated surfaces typically has a strong diurnal dependence as a result of the strong diurnal variations in ozone concentrations and in the

30 stomatal uptake.)

The v_d -SST plot in Figure 1 shows that although there are large fluctuations in the data from the various field experiments, a trend of v_d increasing with SST is apparent. The lowest deposition velocities are observed in the Southern Ocean (GasEx08) and the largest from the TexAQS06 experiment in the North-western Gulf of Mexico. An interpretation of this observed

variation is that despite the stronger wind and wave conditions normally present at Southern Ocean latitudes and the higher ozone solubility due to cooler temperatures there, the deposition velocities are lower because of the lower iodide concentrations at such temperatures which results in a slower ozone loss rate. On the other hand, the observed deposition velocities in the North-western Gulf of Mexico are larger due to warmer waters and higher iodide concentrations leading to

5 faster ozone loss rate despite relatively weak winds and lower ozone solubility there.

Figure 1 shows the ACCESS-UKCA predictions obtained using the default (i.e. constant r_c) deposition scheme for the five experiments (diamonds – same colour as the data). The range of SSTs from the model for different experiments is in agreement with the measurements, with the lowest values predicted for GasEx08 and the highest for TexAQS06/GOMECC07. However, the modelled v_d is virtually constant at 0.038 cm s⁻¹, and the observed variation is

- 10 progressively overestimated by a factor of 2 to 4 as water temperatures get cooler. This model behaviour with SST is a consequence of the model using a constant value for the dominant resistance r_c , resulting in a near-constant v_d with a ceiling value of $1/r_c = 100/2200 = 0.0455$ cm s⁻¹. The difference between the ceiling value and a modelled level corresponds to the contribution made by the airside resistances (r_a and r_b in Eq. (1)) to lower the deposition velocity. The comparison in Figure 1 demonstrates that the constant r_c assumption is not satisfactory both with respect to its absolute value and its lack of
- 15 variation with SST.

Figure 2a shows the same v_d -SST plot as in Figure 1 but with the modelled values obtained using the one-layer reactivity scheme. Even though the molecular diffusion controlled deposition component, $(a D)^{1/2}$, increases by a factor of 30 from high latitude cold waters to tropical warm waters, a variation to that extent is not evident in Figure 2a because turbulent transfer starts to dominate deposition as surface temperatures decrease (this is illustrated in the sensitivity analysis presented

- 20 later in Section 6.1.1). Compared to Figure 1, the one-layer deposition scheme is able to reproduce the observed trend of v_d increasing with SST noticeably better, accompanied by larger fluctuations in v_d , but the model overestimates v_d by a factor of 2 to 3 with the model-data differences getting progressively larger as the water temperatures get cooler. In Figure 2b we plot the behaviour of the one-layer reactivity scheme with the contribution of waterside turbulence removed. Compared to Figure 2a, the modelled deposition velocities are lower in magnitude and compare better with the data. However, it is clear that for
- 25 SSTs lower than about 15°C, the model without the waterside turbulence underestimate the deposition velocities compared with the measurements because the influence of waterside turbulence becomes important compared to weaker chemical reaction at lower temperatures. Overall, what this comparison suggests is that the influence of waterside turbulence in the one-layer reactivity scheme is overestimated, and this point will be discussed later on in the paper.

In Figure 3, the modelled variation of v_d with wind speed (at 10 m height) obtained using the default deposition scheme does not agree with the measurements, with the modelled v_d values almost constant as in Figure 1, whereas the data show an increasing v_d with wind speed for TexAQS06 and to a lesser extent for GOMECC07. For the other three experiments the variation is almost constant, but is much lower in magnitude than that estimated by the model. It is apparent that the higherend range of the modelled wind speed is smaller than that of the data, largely because the wind extremes are averaged out in the modelled monthly averages. This <u>is</u> not as much an issue in the earlier comparison involving SST because this parameter does not change as much as wind speed temporally. Hence, in the present case, the v_d -wind speed comparison is perhaps not as robust as the v_d -SST comparison.

5 Figure 4 compares the measured v_d variations with wind speed with those computed using the one-layer deposition scheme. The model is able to perform somewhat better than that in Figure 3 for a constant r_c , with a hint of the model curves for the five experiments separating out in a manner that is qualitatively consistent with the data.

It is apparent from the above comparison that the one-layer mechanistic scheme given by Eq. (9) is able to yield trends in deposition velocity in the right direction, but the modelled deposition velocities are still overestimated compared to the

10 observed values overall.

The likely reason for the overestimation of deposition velocity by the one-layer reactivity scheme is that it overestimates the influence of waterside turbulence and, consequently, its interaction with chemical reaction as a results of the use of a turbulent diffusivity parameterisation $K_t = \kappa u_{*w} z_w$ that is linear in z_w , the distance (or depth) from the water surface. This parameterisation is valid in the surface layer with fully developed turbulence, but in the small viscous sublayer ($z_w < \delta_u$)

- 15 whose thickness $\delta_u = 11 \nu / u_{*w}$ where ν is the kinematic viscosity of seawater, is of the order of 1 mm under typical conditions, the above K_t parameterisation would overestimate turbulent transfer because it ignores the dissipation of smallscale turbulence by viscosity. This is important because the length scale of the ozone-iodide reaction falls within the viscous sublayer. It is suggested that K_t varies as z_w^{m+1} in the viscous sublayer, where m = 2 for a smooth surface (Fairall et al., 2000). A form for K_t that is valid for both $z_w < \delta_u$ and $z_w > \delta_u$, such as $K_t = \kappa u_{*w} z_w / [1 + (\delta_u / z_w)^m]$, may be
- 20 considered in Eq. (8) but an analytical solution is currently elusive. Based on this more general form for K_t , we can determine that for $z_w \ll \delta_u$ the form $K_t = \kappa u_{*w} z_w$ would overestimate K_t by a factor of $(\delta_u / z_w)^m$. A consequence of this would be to artificially elevate ozone deposition rates to the ocean, especially in regions where the ozone removal chemistry in the surface layer is slow (i.e. areas with cooler SSTs).

6.1 Two-layer reactivity scheme – improved formulation

Fairall et al. (2007) extended the one-layer scheme by considering a two-layer approach in which a thin layer of depth δ_m below the water surface has a high reactivity (*a*) and the region below δ_m has a very small background reactivity (*a*₀) that is present everywhere including in the first layer (so the total reactivity of the first layer is $a + a_0$). The assumption is that virtually all the chemical reaction takes place within this thin layer which is roughly identified with the surface microlayer.

At this stage we acknowledge that there is iodide below the surface microlayer and that ozone may transfer into that layer and be rapidly destroyed. Thus the two-layer scheme has an arbitrary constraint that differs somewhat from nature. We describe this model as a semi-empirical model where the appropriate physical and chemical variables have been included in the model, where the turbulence scheme has limitations already discussed and whereby an arbitrary condition, the depth of

5

the reactive layer is imposed. The usefulness of the model depends on two issues: 1) is the model able to accurately represent the available ozone seawater deposition data, and 2) is the model formulation suitable for efficient use within global climate composition models? These will be both addressed.

The expression for the waterside transfer velocity from the two-layer water reactivity model is:

$$v_{dw} = \left[(a+a_0) D \right]^{1/2} \left[\frac{-A_1 I_1(\xi_0) + B_1 K_1(\xi_0)}{A_1 I_0(\xi_0) + B_1 K_0(\xi_0)} \right],$$
(15)

where $I_n(\xi)$ is modified Bessel function of order *n*, $\xi_0 = b[(a + a_0)D]^{1/2}$ and $b = 2/(\kappa u_*)$. The coefficients A_1 and 10 B_1 are determined by imposing three boundary conditions: the flux at the water surface obtained using the solution of the mass conservation Eq. (8) should be equal to a constant specified value F_0 ; the concentration at the interface of the two layers should be continuous; and the flux at the interface of the two layers should be continuous. These boundary conditions lead to the following equations, respectively:

$$-A_{1}\xi_{0}I_{1}(\xi_{0}) + B_{1}\xi_{0}K_{1}(\xi_{0}) = \frac{2F_{0}}{\kappa u_{*}},$$
(16)

$$A_{1}I_{0}(\xi_{\delta 1}) + B_{1}K_{0}(\xi_{\delta 1}) - B_{2}K_{0}(\xi_{\delta 2}) = 0, \qquad (17)$$

$$-A_{1}\xi_{\delta 1}I_{1}(\xi_{\delta 1}) + B_{1}\xi_{\delta 1}K_{1}(\xi_{\delta 1}) - B_{2}\xi_{\delta 2}K_{1}(\xi_{\delta 2}) = 0, \qquad (18)$$

where $\xi_{\delta 1} = [2b(a + a_0)(\delta_m + (bD/2))]^{1/2}$ and $\xi_{\delta 2} = [2ba_0(\delta_m + (bD/2))]^{1/2}$. Although Fairall et al. (2007) allude to the condition corresponding to Eq. (18) they used a different equation in lieu of Eq. (18) by imposing the condition that the total depletion of concentration caused by diffusion and reaction through the depth of the medium must equal the surface flux F_0 . However, their equation involves integration terms that need to be determined numerically, thus increasing the computational time considerably when used in a large-scale atmospheric model such as ours. We found that the results obtained using both the equations are the same. In Eq. (16), F_0 can be taken as unity because it cancels out in the formula for deposition velocity expressed as flux divided by concentration at the surface.

To solve Eqs. (16)–(18) for the three unknowns A_1 , B_1 and B_2 , the expression for A_1 obtained from Eq. (16) is substituted in Eqs. (17) and (18), the last two are then solved for B_1 and B_2 analytically by inverting the 2×2 matrix of the coefficients B_1

and B_2 . A_1 is obtained by putting the solution for B_1 back in Eq. (16). The two-layer scheme has not been previously tested with any measurements.

6.1.1 Sensitivity analysis

Deposition velocity is sensitive to the value of δ_m in the two-layer scheme. To explore this, we tested this scheme in 5 stand-alone mode (i.e. outside ACCESS-UKCA). The oceanic component of dry deposition velocity, i.e. αv_{dw} (=1/ r_c) was calculated using Eq. (15) as a function of sea surface temperature and reactivity for a range of δ_m values for a typical value of the waterside friction velocity (u_{*w}) of 0.01 m s⁻¹ (which is equivalent to the airside $u_* \approx 0.3$ m s⁻¹). The parameterisations of [I^-], k, D and α specified earlier and $a_0 = 10^{-4}$ s⁻¹ (Fairall et al., 2007) were used. The results are presented in Figure 5a and Figure 5b, along with the curves obtained using the one-layer scheme with and without waterside

- 10 turbulence (i.e. $K_t = 0$). These results show that: 1) the variation of αv_{dw} with SST is very similar to that with reactivity (the latter on logarithmic scale), 2) the thicker the δ_m the higher the deposition velocity and that for a sufficiently large δ_m of about 1 mm the two-layer solution approaches the one-layer solution, 3) turning off waterside turbulence in the one-layer scheme, which leads to the limiting behaviour $\alpha v_{dw} = \alpha (a D)^{1/2}$, makes the deposition velocity for the one-layer scheme without turbulence diverge further and further to become a very small fraction of the dry deposition velocity for the one-
- 15 layer scheme with turbulence as SST or reactivity gets smaller, 4) except for cooler SSTs or lower reactivity, the two-layer scheme results in deposition velocity values that are even smaller than those from the one-layer scheme without turbulence for smaller values of δ_n (such δ_n values are obviously not realistic because the no-turbulence behaviour is the lower limit), and 5) at the lower end of SST or reactivity values, there is a slight increase in αv_{dw} with decreasing SST or reactivity, which is mostly due to the larger influence of solubility which increases with decreasing temperature.
- 20 Sensitivity of v_{dw} to oceanic iodide concentration can also be explored. Based on their compilation of available measurements of sea surface iodide covering latitudes 69°S to 66°N, Chance at al. (2014) suggest the following parameterisation for iodide concentration:

$$[I^{-}](nM) = 0.225(T_{s} - 273.16)^{2} + 19.$$
⁽¹⁹⁾

The iodide concentrations and also the reactivity values (*a*) determined using Eq. (19) are higher by 207, 190, 154, 84 and 25 51% compared to Eq. (12) derived by MacDonald at al. (2014) for the SSTs of 5, 10, 20, 30 and 35°C, respectively. Comparison of the curves in Figure 5c obtained using Eq. (19) with those in Figure 5a shows, as expected, that the values of αv_{dw} are all higher in the former due to higher iodide levels: the one-layer curve is higher by 18, 21, 27, 24 and 17%, the

no-turbulence curve is higher by 75, 70, 59, 36 and 23%, and, as an example, the two-layer curve for $\delta_m = 2 \ \mu m$ is higher by 6, 15, 60, 48 and 25% for the above temperatures, respectively. The two-layer curves from the two iodide concentration formulae require different δ_m values to be similar: for example, the curve for $\delta_m = 2 \ \mu m$ in Figure 5c almost coincides with that for $\delta_m = 5 \ \mu m$ in Figure 5a. We use the parameterisation Eq. (12) for iodide concentration in all subsequent

5 calculations. Nevertheless, it is clear that the use of Eq. (19) in the one-layer scheme would not describe the data in Figure 2a properly.

Other sensitivity results show that as u_{*_w} gets larger it requires larger values of δ_m to approach the one-layer behaviour, and that the solution obtained by the two-layer model approaches the one-layer model when $a_0 \rightarrow a$.

6.1.2 Comparison with data

- 10 The two-layer reactivity scheme was incorporated into ACCESS-UKCA. The value of δ_n is selected based on the best agreement with data, given the constraint that it cannot be smaller than the reacto-diffusive length scale over which chemical reaction controls deposition. In other words, the variation of deposition yielded by a selected δ_m cannot be lower than the no turbulence case (i.e., $K_t = 0$) for the same input conditions (see, for example, Figure 5). Based on the above considerations, a value of $\delta_m = 2 - 3 \ \mu m$ was found to be appropriate (which is of the order of the reacto-diffusive length
- 15 scale) and we selected $\delta_m = 2.5 \ \mu\text{m}$. The model variation of v_d with SST determined using the two-layer reactivity scheme is presented in Figure 6 along with the observed data. There is now a substantial improvement in the prediction of the observed v_d compared to the results from the default and one-layer schemes presented earlier. The modelled variation passes almost through the middle of the data. The variation of v_d with wind speed in Figure 7 also shows considerable improvement over the previous model results, but as mentioned earlier the higher-end range of the modelled wind speed is smaller due to
- 20 the fact that the wind extremes are averaged out in the modelled monthly averages. There is a relatively large scatter in the GOMECC07 predictions, but the modelled values for the rest of the experiments are similar to the measurements. These results imply that the two-layer scheme has got the combined influences of solubility, molecular diffusion, chemical reaction and turbulent transfer into better balance than the other two schemes.

The reason the two-layer scheme works is as follows. As mentioned earlier, there is an overestimation of the interaction 25 between the chemical reaction term (*a*) and the total diffusivity term $(D + K_t)$ in Eq. (8) within the viscous sublayer because of the overestimation of K_t as a result of the assumption of its linearly increasing variation with depth. This leads to larger values of deposition velocity than observed. In the two-layer scheme, we effectively compensate for this overestimation of turbulent transfer by constraining the extent of chemical reaction by limiting the iodide reactivity to a

thickness of δ_m equivalent to the reacto-diffusive length scale even though in reality iodide is present through the depth of the oceanic surface layer.

There could be both We also speculate that there may be some overestimate and underestimate to some degree in the current reaction pathway for ozone within the ocean. Most of the chemistry underlying the kinetics has been studied at iodide concentrations varying from mM to M, whereas in the ocean the iodide concentration is approximately a million times more dilute, being in the nM range. The fundamental step by step chemistry of ozone reaction with iodide has not been fully characterized. For example if the bonding of the iodate complex in the reaction step involving ozone and iodide yielding iodate (Sakamoto et al., 2009) is weak, there could be a reverse reaction freeing the bound ozone. This is a speculative example of what could lead to a reduction in deposition rates. On the other hand, reaction of ozone with compounds other

- 10 than iodide in seawater could lead to a significant increase in deposition velocity. The analysis by Chang et al. (2004) considered reaction of ozone with iodide, dimethyl sulfide (DMS), ethene and propene and showed that the reaction with iodide was by far the fastest of all (the reaction with DMS was comparably fast only when it was present at extremely high concentrations). In one of their model simulations, in addition to the reactions with iodide, DMS and alkenes, Ganzeveld et al. (2009) also included ozone reaction with chlorophyll-a as a first order approximation to examine the possible role of
- 15 dissolved organic matter (DOM) (e.g. in coastal waters). They found that the inclusion of the O₃- chlorophyll-a chemistry in the model significantly increased deposition velocity at coastal sites but gave mixed results compared to observations. For open ocean sites, there were only small changes to deposition velocity. Clearly more observations of deposition velocity are needed for different oceanic regions to establish the relative role of additional ozone reactions in explaining the observed trends as well as related scatter.
- 20 <u>Based on the above comparison with data from Helmig et al. (2012), Overall</u> the two-layer mechanism provides a good <u>overall</u> representation of the temperature and wind speed dependence and absolute values of ozone deposition to the ocean. It is also sufficiently simple mathematically to require relatively little extra time to compute within the framework of the ACCESS chemistry-climate model.

7 Global fields of ozone deposition velocity, loss and concentration

25 7.1 Deposition velocity

30

The modelled global distribution of ozone deposition velocity (cm s⁻¹) averaged over the year 2006 in Figure 8 shows that the two-layer reactivity scheme (top panel) gives the largest deposition velocities, as high as 0.03 cm s⁻¹, in the tropics. v_d decreases with increasing latitude, but at around 60°S it increases again slightly mainly due to the increase in ozone solubility with decreasing water temperature starts to exert more influence coupled with higher wind speeds in this latitudinal zone. From the tropics to mid latitudes the global distribution of v_d is qualitatively similar to the global SST distribution (plot not shown) because of the dominance of the ozone-iodide reaction in determining v_d in which iodide concentration has a proportional dependence on SST. The distribution from the one-layer reactivity scheme (middle panel) is qualitatively similar to the two-layer result but with v_d values almost 2 to 3 times as large. The default ACCESS-UKCA scheme with $r_c = 2200$ s m⁻¹ (bottom panel) provides a much flatter variation of v_d across the globe. The magnitude of deposition velocity predicted by this scheme within the tropics is similar to that by the one-layer scheme, but beyond the tropics the values predicted by the constant surface resistance model are larger by approximately a factor of two.

5

The relative difference (%) between v_d predicted by the one-layer (v_{d1}) and two-layer (v_{d2}) reactivity schemes (defined as $(v_{d1} - v_{d2}) \times 100 / v_{d2}$) is presented in Figure 9 (top panel), where differences as high as 150% can be seen in the mid latitudes, and as low as 25–50% in the tropics. The differences between the default scheme and the two-layer reactivity scheme (bottom panel) in the mid latitudes are even greater (up to 300%).

In Table 1, the average deposition velocity for seawater (which excludes sea ice and coastal grid cells) obtained using the two-layer reactivity scheme is almost half that of the default scheme and two-third that of the one-layer scheme, which leads to a small lowering of the globally averaged value for the two-layer scheme. The averaged Southern Hemisphere deposition velocity values are generally lower than the Northern Hemisphere values. It is important to note that the two-layer model is the best fit to the available observational data. The observations as well as the model are indicating the need for a decrease in the oceanic deposition velocities used for ozone in atmospheric chemistry models.

To explore seasonal differences, we plot Figure 10 which is the same as Figure 8 except that the left panels are for January 2006 and the right panels are for July 2006. The distributions are similar to the corresponding ones in Figure 8 but there is more spatial variability in Figure 10 because of shorter averaging (i.e. one month). For the two-layer scheme (top panels), in January the deposition velocities are generally larger in the mid to high latitudes in the Northern Hemisphere than those in

- 20 the Southern Hemisphere, and as expected this behaviour is reversed in July mostly as a result of seasonal SST changes. For such latitudes in the Northern Hemisphere, the January deposition velocities are higher than those in July, and the reverse occurs in the Southern Hemisphere. In the tropics and subtropics, there is an upward latitudinal shift of high deposition velocity regions in July compared to January. In qualitative terms, the above behaviour is also evident in the results from the one-layer scheme (middle panels). For the default scheme (bottom panels), the seasonal variations in v_d are smaller and less
- 25 apparent, and would be mainly dominated by seasonal wind-speed variations. In all plots, the white region around the poles is the extent of sea ice which is larger in July around the South Pole and in January around the North Pole.

Ganzeveld et al. (2009) presented January and July deposition velocities simulated using the one-layer reactivity scheme in their free running ECHAM5/MESSy climate-chemistry model (see their Figures 3a and 3b, respectively). Their plots are significantly different to the corresponding middle panels in Figure 10. Our results show maximum values of deposition

30 velocity in the tropics and subtropics (largely as a result of enhancement due to chemical reaction because of higher iodide associated with higher SSTs) whereas the maxima in their plots are mostly located in mid-latitudes which they attribute to

high wind speeds in the storm track regions located at such latitudes. Apart from potentially significant differences in the formulation of the two climate-chemistry models, there are other differences as well such as resolution, use of nudging in our model, emissions, and the parameterisation of iodide concentration.

7.2 Ozone loss

- 5 The modelled global distribution of dry deposition velocity to the ocean can be combined with the modelled ozone concentration fields to enable the loss of ozone to the ocean surface to be estimated. The deposition budgets obtained using ACCESS-UKCA are presented in Table 2. Compared to the default scheme the global ozone deposition to seawater, excluding sea ice and coastal grid cells, is 85% for the one-layer reactivity scheme and it is almost halved for the two-layer reactivity scheme. Deposition to the sea is 12% of the total deposition for the two-layer scheme, 19% for the one-layer
- 10 scheme and 21% for the default scheme. The reduction in oceanic deposition using the two-layer scheme corresponds to a 10% decrease in the total global estimate of ozone deposition relative to the original (default) scheme<u>and this is consistent</u> with the current observations of deposition velocity to the ocean surface. The seawater deposition in the Northern Hemisphere is slightly larger than that in the Southern Hemisphere due to the higher average seawater deposition velocity in the former (Table 1). Of the total global ozone deposition, about 70% is in the Northern Hemisphere and 30% in the
- 15 Southern Hemisphere, the former being larger due to the much larger terrestrial contribution. The total ozone deposition obtained from ACCESS-UKCA is on the lower end of the values reported from other models: 1094 ± 264 Tg yr⁻¹ (Young et al., 2013), 1003 ± 200 Tg yr⁻¹ (Stevenson et al., 2006) and 949 ± 222 Tg yr⁻¹ (Wild, 2007) (see also Climate Change, The Physical Science Basis, 2013, IPCC, AR5, Chapter 8, Anthropogenic and Natural Radiative Forcing, Table 8.1, page 672; Young et al., 2013). This is likely due to the general underestimation of tropospheric ozone by ACCESS-UKCA
- 20 (Woodhouse et al., 2015), the exact reason(s) for which we have not been able to pinpoint but it is an ongoing area of study. An underestimation of ozone would result in an underestimation of the ozone dry deposition flux near the surface. However, the calculated deposition velocity is not influenced by the predicted ozone concentration, so the relative dry deposition budget contributions given above in terms of percentages should remain representative.

These <u>relative</u> decreases in ozone oceanic deposition, calculated here with the two-layer model, are not model artefacts, but through the model validation process, are based on the best available representation of the observations of oceanic ozone deposition rates available today.

7.3 Ozone concentration

While we have previously noted that ACCESS-UKCA underestimates tropospheric ozone, it is worthwhile to examine how an oceanic dry deposition parameterisation affects tropospheric ozone distributions. The top panel in Figure 11 shows the modelled annual, near-surface (at 20 m) ozone concentration (ppbv) based on the two-layer reactivity scheme. Relatively high concentrations in the Northern Hemisphere, especially within 0–50°N, are evident, which can be related to the larger

30

precursor emissions in these regions. The middle panel presents the relative difference (%) between the ozone concentration predicted by the one-layer (c_1) and two-layer (c_2) reactivity schemes (defined as $(c_1 - c_2) \times 100/c_2$). It can be seen that the largest underestimation of the near surface ozone concentration by the one-layer scheme is 11% for 45–70°S. The bottom panel in Figure 11 is the relative difference between the concentration predicted by the default scheme and the two-layer

5 reactivity scheme, which shows the largest underestimation by the default scheme of 17% within the same latitudinal band. The bottom panel can be approximately compared with the results in Figure 10 of the paper by Ganzeveld et al. (2009) in which they present the relative difference between a constant r_c scheme (with $r_c = 2000$ s m⁻¹) and the one-layer scheme. The largest underestimation by their constant r_c scheme is about 4% in the latitudes 45–70°S; this latitudinal band is very similar to what is suggested by the present results but with an underestimation of 17%. Their results show that there is also a 10 significant overestimation by their constant r_c scheme, the largest of which is about 4% over the tropical and subtropical

We also examine the influence of the various schemes on the modelled ozone distribution throughout the troposphere. The top left plot in Figure 12 is the observed zonal distribution of ozone concentration as a function of altitude for the year 2006 based on the monthly mean ozone profile database available from and described at

(30°S–30°N) waters and in high latitude regions in the Northern Hemisphere. This, however, is not apparent in our results.

- 15 <u>http://www.bodekerscientific.com/data/monthly-mean-global-vertically-resolved-ozone</u>. These gap filled data (Tier 1.4) are based on the raw individual ozone data sourced from the so-called Binary Data Base of Profiles (BDBP) database (Hassler et al., 2009) and cover the whole globe. The top right plot is the corresponding distribution obtained from ACCESS-UKCA using the two-layer reactivity scheme. There are similarities between the observed and modelled fields; for example, the tropospheric ozone levels are lower in the Southern Hemisphere in both plots. The high ozone concentration areas at the top
- 20 corners correspond to the lower edges of the stratosphere which starts at an altitude of about 10 km at the poles and increase to about 15 km at the Equator. In the troposphere, the modelled concentrations are generally lower compared to the observations, for example in the high latitudes, which is a known feature of the model (Woodhouse et al., 2015). This is more clearly evident in the bottom-left plot which presents the relative difference (= $[(\overline{M} - \overline{O})/\overline{O}] \times 100\%$) between the time-averaged modelled concentration (\overline{M}) determined using the two-layer reactivity scheme and the time-averaged
- 25 observed concentration (\overline{O}). The bottom-right figure is the same plot except for the default scheme. There are improvements in the ozone prediction with the two-layer scheme south of 40°S, both at the lowest altitudes and at upper levels as high as 6 km. There is also some improvement in the tropical south at a height of about 8 km. There is a slight model improvement in the high latitudes 70–80°N between the altitudes 2–6 km. Overall, for the latitudes 45–70°S, the twolayer scheme improves the prediction of the observed zonal ozone mixing ratios at an altitude of 1 km by 7% compared to
- 30 the original default scheme. For the altitude range 1-6 km, this improvement is 5%.

8 Conclusions

Ozone dry deposition parameterisation schemes are best evaluated within an integrated modelling framework that includes multi-parameter interdependencies of deposition as exists in the field. Using recent measurements, we assessed the performance of three ozone dry deposition schemes for seawater within the global climate-chemistry model ACCESS-

- 5 UKCA incorporating meteorological nudging and monthly-varying emissions. The default scheme assumes a constant water surface resistance of 2200 s m⁻¹, which is commonly used in most atmospheric chemistry/composition models. The second scheme is a mechanistic, one-layer reactivity scheme proposed by Fairall et al. (2007) in which the surface resistance formulation includes the simultaneous influence of ozone solubility in water, waterside molecular diffusion and turbulent transfer, and a first-order chemical reaction of ozone with dissolved iodide. The third scheme is a development of Fairall et
- 10 al.'s (2007) two-layer reactivity scheme in which, unlike the uniform reactivity assumption in the one-layer scheme, the water surface has a high reactivity and the region below has a very small background reactivity.

A comparison of the observed deposition velocity dependencies on sea surface temperature and wind speed with those obtained from ACCESS-UKCA using the three schemes showed that the two-layer scheme is able to describe the absolute magnitude and the sea surface temperature and wind speed dependence of the field measurements most closely. The two-

- 15 layer scheme results are sensitive to the value of the thickness of the surface reactive layer a value between 2 to 3 microns, which is about the same as the typical reacto-diffusive length scale over which the ozone-iodide chemical reaction dominates deposition, works well and we chose 2.5 microns. The assumption in the first scheme of a constant water surface resistance overestimates the observed deposition velocity by a factor of 2 to 4 and does not describe its variability with SST and wind speed. The one-layer scheme performs somewhat better than the default scheme. Although the one-layer scheme includes all
- 20 the important processes as the two-layer scheme, it still overestimates the observed deposition velocity, by a factor of 2 to 3 when the reaction is slow under colder SSTs. The likely reason for this problem is an overestimation of the influence of waterside turbulent transfer on chemical reaction via the assumption that turbulent diffusivity is a linear function of depth. This assumption, valid for a fully turbulent flow, overestimates turbulent transfer in the viscous sublayer (within which the bulk of the ozone-iodide reaction takes place) because it does not account for the viscous dissipation of turbulence in this
- 25 layer. The two-layer reactivity scheme limits the dissolved iodide concentration to a specified depth from the water surface, thereby restricting the ozone-iodide reaction and its interaction with turbulence to that depth. Although this restriction is artificial, it works as surrogate to compensate for the overestimation of turbulent transfer in the viscous sublayer. A better parameterisation of turbulent mixing near the interface in the concentration conservation equation could overcome the issue.

The modelled global distributions show that the two-layer reactivity scheme yields maximum ozone deposition velocities in

30 the tropics, they decrease with increasing latitude and then increase again slightly in the Southern Hemisphere. Significant local variations are also evident. This latitudinal behaviour is the result of the combined effects of chemical reaction, solubility, molecular diffusion and turbulent transfer, and because all these processes except turbulent transfer are direct functions of SST, which in turn is a function of latitude, there is a strong latitudinal dependence.



Previous model estimates in the scientific literature suggest approximately one third of the total ozone dry deposition is to the ocean. The deposition velocities obtained using the two-layer scheme and <u>also</u> the recent oceanic ozone deposition observations point to a substantially lower ozone deposition to the ocean. A new modelled estimate of ozone deposition to the ocean of about 80 Tg yr⁺ was obtained (which excludes sea ice and coastal grid cells), which is 12% of the modelled

- 5 total global ozone deposition, is almost half of the original oceanic deposition obtained using the default scheme, and corresponds to a 10% decrease in the total global estimate of ozone deposition relative to the original (default) scheme, making a significant relative change to the overall tropospheric ozone budget. Although it does not influence the modelled deposition velocity, the underestimation of tropospheric ozone by ACCESS-UKCA needs to be investigated as it leads to lower absolute value of annual global deposition.
- 10 The effects of the choice of a deposition scheme in the global model were noticeable in the predicted atmospheric ozone mixing ratios, especially in the Southern Hemisphere. For the latitudes 45–70°S, the two-layer scheme improved the ozone predictions near the surface by 7% and those within the altitude range 1–6 km by 5% compared to the original default scheme.

Based on all the evaluation results presented above, the two-layer reactivity scheme as formulated in this paper is the best

15 performing scheme, compared to the available observations, for describing the ozone deposition to seawater in a global modelling framework.

Further observations on deposition velocity with greater temporal and spatial coverage would help constrain <u>oceanic</u> dry deposition schemes better. Such work would also look at potential improvement to the parameterisation of iodide in terms of physical and chemical properties of seawater and an assessment of the relative impact of any additional chemical reactions to

20 <u>ozone uptake at sea surface</u>. Studies involving observations of the sea surface microlayer, the role of wave breaking and bubbles on deposition processes, and the degree of relevant feedbacks in climate-chemistry models (e.g., the ozone reaction with oceanic iodide which produces iodine compounds that participate in atmospheric ozone destruction cycles in the troposphere) need to be explored.

Acknowledgements

- 25 The work was supported by funding from the Australian Climate Change Science Program (ACCSP). The model runs were performed on the National Computational Infrastructure (NCI) Facility in Canberra, Australia, which is supported by the Australian Commonwealth Government. Communication with Chris Fairall, Detlev Helmig, Laurens Ganzeveld and Rosie Chance on their works was very useful. We are indebted to Ludovic Bariteau for kindly supplying data related to their published work. Fiona O'Connor, Mohit Dalvi, Steve Rumbold and Colin Johnson of the U.K. Met Office are thanked for their assistance with the UKCA emission methodology and answering questions about UM-UKCA. Luke Abraham of
- University of Cambridge is thanked for his help with several technical questions on UKCA. Martin Dix and Scott Wales are

acknowledged for their help with model configuration issues, and Lauren Stevens for help with emissions data. ERA-Interim data from the European Centre for Medium-Range Weather Forecasts was used in this research. We would like to thank Greg Bodeker (Bodeker Scientific) and Birgit Hassler (NOAA) for providing the combined vertical ozone profile database. We thank the two reviewers<u>and co-editor</u> for their helpful comments.

5 References

10

15

Abraham, N. L., Archibald, A. T., Bellouin, N., Boucher, O., Braesicke, P., Bushell, A., Carslaw, K. S., Collins, W., Dalvi,
M., Emmerson, K. M., Folberth, G., Haywood, J., Johnson, C., Kipling, Z., Macintyre, H., Mann, G. W., Telford, P. J.,
Merikanto, J., Morgenstern, O., O'Connor, F., Ordonez, C., Osprey, S., Pringle, K. J., Pyle, J. A., Rae, J. G. L., Reddington,
C. L., Savage, D., Spracklen, D., Stier, P., and West, R.: Unified Model Documentation Paper No. 84: United Kingdom
Chemistry and Aerosol (UKCA) Technical Description MetUM Version 8.4. UK Met Office, Exeter (UK), pp. 74, 2012.

Bi, D. H., Dix, M., Marsland, S. J., O'Farrell, S., Rashid, H. A., Uotila, P., Hirst, A. C., Kowalczyk, E., Golebiewski, M.,
Sullivan, A., Yan, H. L., Hannah, N., Franklin, C., Sun, Z. A., Vohralik, P., Watterson, I., Zhou, X. B., Fiedler, R., Collier,
M., Ma, Y. M., Noonan, J., Stevens, L., Uhe, P., Zhu, H. Y., Griffies, S. M., Hill, R., Harris, C., and Puri, K.: The ACCESS coupled model: description, control climate and evaluation, Australian Meteorological and Oceanographic Journal, 63, 41–64, 2013.

Carpenter, L. J., MacDonald, S. M., Shaw, M. D., Kumar, R., Saunders, R. W., Parthipan, R., Wilson, J., and Plane, J. M. C.: Atmospheric iodine levels influenced by sea surface emissions of inorganic iodine, Nature Geoscience, 6, 108–111, DOI: 10.1038/NGEO1687, 2013.

Carpenter, L. J. and Nightingale, P. D.: Chemistry and release of gases from the surface ocean, Chemical Reviews, 115, 4015–4034, DOI: 10.1021/cr5007123, 2015.

Chance, R., Baker, A. R., Carpenter, L., and Jickells, T. D.: The distribution of iodide at the sea surface, Environmental Science: Processes & Impacts, 16, 1841–1859, DOI: 10.1039/c4em00139g, 2014.

Chang, W. N., Heikes, B. G., and Lee, M.: Ozone deposition to the sea surface: Chemical enhancement and wind speed dependence, Atmospheric Environment, 38, 1053–1059, doi:10.1016/j.atmosenv.2003.10.050, 2004.

25 Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), Geosci. Model Dev., 3, 43–67, doi:10.5194/gmd-3-43-2010, 2010.

Fairall, C. W., Hare, J. E., Edson, J. B., and McGillis, W.: Parameterization and micrometeorological measurements of air-30 sea gas transfer, Bound.-Layer Meteorol., 96, 63–105, DOI: 10.1023/A:1002662826020, 2000. Fairall, C. W., Hare, J. E., Helmig, D., and Ganzveld, L.: Water-side turbulence enhancement of ozone deposition to the ocean, Atmos. Chem. Phys., 7, 443-451, doi:10.5194/acp-7-443-2007, 2007.

Fairall, C. W., Yang, M., Bariteau, L., Edson, J., Helmig, D., McGillis, W., Pezoa, S., Hare, J. E., Huebert, B., and Blomquist, B.: Implementation of the coupled ocean-atmosphere response experiment flux algorithm with CO₂, dimethyl sulfide, and O₃, Journal of Geophysical Research 116, C00F09, doi:10.1029/2010JC006884, 2011.

Galbally, I. E. and Roy C. R.: Destruction of ozone at the Earth's surface, Q. J. R. Meteorol. Soc., 106, 599-620, doi:10.1002/gj.49710644915, 1980.

Galbally, I. E., Bentley, S. T., and Meyer, C. P.: Mid-latitude marine boundary-layer ozone destruction at visible sunrise observed at Cape Grim, Tasmania, 41°S, Geophysical Research Letters, 27, 3841-3844, DOI: 10.1029/1999GL010943 2000.

Ganzeveld, L., Helmig, D., Fairall, C., Hare, J. E., and Pozzer, A.: Atmosphere-ocean ozone exchange: A global modeling study of biogeochemical, atmospheric and water-side turbulence dependencies, Global Biogeochem. Cycles, 23, GB4021, doi:10.1029/2008GB003301, 2009.

Garland, J. A., Elzerman, A. W., and Penkett, S. A.: The mechanism for dry deposition of ozone to seawater surfaces, J. 15 Geophys. Res., 85, 7488-7492, doi:10.1029/JC085iC12p07488, 1980.

Hardacre, C., Wild, O., and Emberson, L.: An evaluation of ozone dry deposition in global scale chemistry climate models, Atmospheric Chemistry and Physics, 15, 6419–6436, doi:10.5194/acp-15-6419-2015, 2015.

Hassler, B., Bodeker, G. E., Cionni, I., and Dameris, M.: A vertically resolved, monthly mean, ozone database from 1979 to 2100 for constraining global climate model simulations, International Journal of Remote Sensing, 30 (15-16), 4009-4018,

DOI: 10.1080/01431160902821874, 2009. 20

5

10

Helmig, D., Lang, E. K., Bariteau, L., Boylan, P., Fairall, C. W., Ganzeveld, L., Hare, J. E., Hueber, J., and Pallandt, M.: Atmosphere-ocean ozone fluxes during the TexAQS 2006, STRATUS 2006, GOMECC 2007, GasEx 2008, and AMMA 2008 cruises, J. Geophys. Res., 117, D04305, doi:10.1029/2011JD015955, 2012.

Hicks, B. B., Baldocchi, D. D., Meyers, T. P., Hosker Jr., R. P., and Matt, D. R.: A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities, Water, Air, and Soil Pollution, 36, 311-330, DOI: 25 10.1007/BF00229675, 1987.

Johnson, P. N. and Davis, R. A.: Diffusivity of ozone in water, Journal of Chemical & Engineering Data, 41, 1485–1487, DOI: 10.1021/je9602125, 1996.

Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jockel, P., Technical Note: An implementation of the dry removal processes DRY DEPosition and SEDImentation in the Modular Earth Submodel System (MESSy), Atmos. Chem. Phys., 6, 4617–4632, doi:10.5194/acp-6-4617-2006, 2006.

Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P.

5 H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, Geosci. Model Dev., 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.

Liss, P. and Merlivat, L.: Air–gas exchange rate: introduction and synthesis. In: Buat-Menard, P. (Ed.), The Role of Air–Sea Exchange in Geochemical Cycling. D. Reidel, Hingham, MA, pp. 113–129, 1986.

10 Magi, L., Schweitzer, F., Pallares, C., Cherif, S., Mirabel, P., and George, C.: Investigation of the uptake rate of ozone and methyl hydroperoxide by water surfaces, J. Phys. Chem. A, 101, 4943–4949, DOI: 10.1021/jp970646m, 1997.

Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, Geoscientific Model Development, 3, 519–551, doi:10.5194/gmd-3-519-2010, 2010.

15 MacDonald, S. M., Gómez Martín, J. C., Chance, R., Warriner, S., Saiz-Lopez, A., Carpenter, L. J., and Plane, J. M. C.: A laboratory characterisation of inorganic iodine emissions from the sea surface: dependence on oceanic variables and parameterisation for global modelling, Atmos. Chem. Phys., 14, 5841–5852, doi:10.5194/acp-14-5841-2014, 2014.

- Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., Fowler, D., Granier, C., Law, K. S., Mills, G. E., Stevenson, D. S., Tarasova, O., Thouret, V., von Schneidemesser, E., Sommariva, R., Wild, O., and Williams, M. L.: Tropospheric ozone and its precursors from the urban to the global scale from air quality to short-lived climate forcer, Atmos. Chem. Phys., 15, 8889–8973, doi:10.5194/acp-15-8889-2015, 2015.
- 25 Morgenstern, O., Braesicke, P., O'Connor, F. M., Bushell, A. C., Johnson, C. E., Osprey, S. M., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model - Part 1: The stratosphere, Geoscientific Model Development, 2, 43–57, doi:10.5194/gmd-2-43-2009, 2009.

Morris, J.C.: The aqueous solubility of ozone—A review, Ozone news, 1, 14-16, 1988.

20

Neu, J. L., Prather, M. J., and Penner, J. E.: Global atmospheric chemistry: Integrating over fractional cloud cover, J.
30 Geophys. Res., 112, D11306, doi:10.1029/2006JD008007, 2007.

Mao, J., Paulot, F., Jacob, D. J., Cohen, R. C., Crounse, J. D., Wennberg, P. O., Keller, C. A., Hudman, R. C., Barkley, M. P., and Horowitz, L. W.: Ozone and organic nitrates over the eastern United States: Sensitivity to isoprene chemistry, Journal of Geophysical Research: Atmospheres, 118, 11,256-211,268, doi:10.1002/jgrd.50817, 2013.

O'Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., Dalvi, M., Folberth, G. A., Sanderson, M. G., Telford, P. J., Voulgarakis, A., Young, P. J., Zeng, G., Collins, W. J., and Pyle, J. A.: Evaluation of the new UKCA climate-composition model - Part 2: The Troposphere, Geoscientific Model Development, 7, 41–91, doi:10.5194/gmd-7-41-2014, 2014.

5 Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P.: Numerical Recipes in Fortran 77: the Art of Scientific Computing. 2nd Edition, Cambridge University Press, Cambridge (UK), 973 pp., 1992.

Sakamoto, Y., Yabushita, A., Kawasaki, M., and Enami, S.: Direct emission of I₂ molecule and IO radical from the heterogeneous reactions of gaseous ozone with aqueous potassium iodide solution, J. Phys. Chem. A, 113, 7707–7713, DOI: 0.1021/jp903486u, 2009

10 Sarwar, G., Gantt, B., Schwede, D., Foley, K., Mathur, R., and Saiz-Lopez, A.: Impact of enhanced ozone deposition and halogen chemistry on tropospheric ozone over the Northern Hemisphere, Environ. Sci. Technol., 49, 9203–9211, doi: 10.1021/acs.est.5b01657, 2015.

Sherwen, T., Evans, M. J., Carpenter, L. J., Andrews, S. J., Lidster, R. T., Dix, B., Koenig, T. K., Sinreich, R., Ortega, I., Volkamer, R., Saiz-Lopez, A., Prados-Roman, C., Mahajan, A. S., and Ordóñez, C.: Iodine's impact on tropospheric oxidants: a global model study in GEOS-Chem, Atmos. Chem. Phys., 16, 1161–1186, doi:10.5194/acp-16-1161-2016, 2016.

15

Smith, S. D.: Coefficients for sea surface wind stress, heat flux, and wind profiles as a function of wind speed and temperature, J. Geophys. Res., 93, 15467–15472, DOI:10.1029/JC093iC12p15467, 1988.

Stevenson, D. S., Dentener, F. J., Schultz, M. G., Ellingsen, K., van Noije, T. P. C., Wild, O., Zeng, G., Amann, M., Atherton, C. S., Bell, N., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M.,

- 20 Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Krol, M. C., Lamarque, J.-F., Lawrence, M. G., Montanaro, V., Müller, J.-F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Shindell, D. T., Strahan, S. E., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, Journal of Geophysical Research, 111, D08301, doi:10.1029/2005JD006338, 2006.
- 25 Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B. et al.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), Atmos. Chem. Phys., 13, 3063–3085, doi:10.5194/acp-13-3063-2013, 2013.

Telford, P. J., Braesicke, P., Morgenstern, O., and Pyle, J. A.: Technical Note: Description and assessment of a nudged version of the new dynamics Unified Model, Atmospheric Chemistry and Physics, 8, 1701–1712, doi:10.5194/acp-8-1701-30 2008, 2008.

Uhe, P. and Thatcher, M.: A spectral nudging method for the ACCESS1.3 atmospheric model, Geoscientific Model Development, 8, 1645–1658, doi:10.5194/gmd-8-1645-2015, 2015.

Vingarzan, R.: A review of surface ozone background levels and trends, Atmospheric Environment, 38, 3431–3442, doi:10.1016/j.atmosenv.2004.03.030, 2004.

5 von Kuhlmann, R., Lawrence, M. G., Crutzen, P. J., and Rasch, P. J.: A model for studies of tropospheric ozone and nonmethane hydrocarbons: Model description and ozone results, J. Geophys. Res., 108(D9), 4294, doi:10.1029/2002JD002893, 2003.

Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, Journal of Geophysical Research, 97 (C5), 7373–7382, 10.1029/92JC00188, 1992.

10 Wesely, M.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical-models, Atmospheric Environment, 23, 1293–1304, doi:10.1016/0004-6981(89)90153-4, 1989.

Wild, O.: Modelling the global tropospheric ozone budget: Exploring the variability in current models, Atmos. Chem. Phys., 7, 2643–2660, doi:10.5194/acp-7-2643-2007, 2007.

Woodhouse, M. T., Luhar, A. K., Stevens, L., Galbally, I., Thatcher, M., Uhe, P., Wolff, H., Noonan, J., and Molloy, S.:

15 Australian reactive-gas emissions in a global chemistry-climate model and initial results, Air Quality and Climate Change, 49, 31-38, 2015.

Young, P. J., Archibald, A. T., Bowman, K. W., Lamarque, J.-F., Naik, V., Stevenson, D. S., Tilmes, S., Voulgarakis, A., Wild, O., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Horowitz, L. W., Josse, B., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M.,

20 Rumbold, S. T., Skeie, R. B., Shindell, D. T., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Pre-industrial to end 21st century projections of tropospheric ozone from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), Atmos. Chem. Phys., 13, 2063–2090, doi:10.5194/acp-13-2063-2013, 2013.

Table 1: Modelled average ozone dry deposition velocities for the year 2006 (cm s $^{\text{-}1}$)

Deposition scheme	Ocean			Ocean, land and iceGlobal			
	Northern	Southern	Average	Northern	Southern	Aueraga	
	Hemisphere	Hemisphere		Hemisphere	Hemisphere	Avelage	
Two layer	0.020	0.017	0.018	0.097	0.051	0.074	
One layer	0.033	0.032	0.032	0.104	0.062	0.083	
Default (constant r_c)	0.039	0.039	0.039	0.108	0.067	0.087	

5

Table 2: Modelled ozone dry deposition for the year 2006 $(Tg\ yr^{\text{-}1})$

Deposition scheme	Ocean			GlobalOcean, land and ice			
	Northern	Southern	<u>Global</u> Total	Northern	Southern	GlobalTotal	
	Hemisphere	Hemisphere		Hemisphere	Hemisphere	<u>0100ai</u> 10tai	
Two layer	44.8	37.7	82.5	500.5	209.2	709.7	
One layer	76.5	67.5	144.0	530.4	237.7	768.1	
Default (constant r_c)	89.2	78.5	167.7	541.5	248.0	789.5	

28



Figure 1: Ozone dry deposition velocity (v_d) as a function of sea surface temperature (SST) from five field experiments (Helmig et al., 2012; Ludovic Bariteau, personal communication, 2016) and the corresponding values obtained from the ACCESS-UKCA model using the default parameterisation for ozone deposition to the ocean involving a constant surface resistance of $r_c = 2200$ s m⁻¹. The dashed line is the ceiling value corresponding to $1/r_c = 100/2200 = 0.0455$ cm s⁻¹.



Figure 2: Ozone dry deposition velocity (v_d) as a function of sea surface temperature (SST) from five field experiments (Helmig et al., 2012; Ludovic Bariteau, personal communication, 2016) and the corresponding values obtained from the ACCESS-UKCA model using: (a) the one-layer reactivity scheme for ozone deposition to the ocean, and (b) the one-layer reactivity scheme without waterside turbulence.



Figure 3: Ozone dry deposition velocity (v_d) as a function of wind speed at 10 m height from five field experiments (Helmig et al., 2012) and the corresponding values obtained from the ACCESS-UKCA model with the default parameterisation for ozone deposition to the ocean.





Figure 4: Ozone dry deposition velocity (v_d) as a function of wind speed at 10 m height from five field experiments (Helmig et al., 2012) and the corresponding values obtained from the ACCESS-UKCA model using the one-layer reactivity scheme for ozone deposition to the ocean.



Figure 5: Variation of the oceanic component of ozone dry deposition velocity αv_{dw} (=1/ r_c) as a function of (a) sea surface temperature (SST, °C), and (b) reactivity *a* (s⁻¹), determined using the one-layer reactivity deposition scheme with and without waterside turbulence, and the two-layer reactivity deposition scheme for a range of δ_n values. Eq. (12) was used for the iodide concentration and the waterside friction velocity (u_{*w}) used was 0.01 m s⁻¹. Plot (c) is the same as (a) except that Eq. (19) was used for the iodide concentration.



5 Figure 6: Ozone dry deposition velocity (v_d) as a function of sea surface temperature (SST) from five field experiments (Helmig et al., 2012; Ludovic Bariteau, personal communication, 2016) and the corresponding values obtained from the ACCESS-UKCA model using the two-layer reactivity scheme for ozone deposition to the ocean.





Figure 7: Ozone dry deposition velocity (v_d) as a function of wind speed at 10 m height from five field experiments (Helmig et al., 2012) and the corresponding values obtained from the ACCESS-UKCA model using the two-layer reactivity scheme for ozone deposition to the ocean.



Figure 8: Modelled ozone dry deposition velocity (v_d , cm s⁻¹) averaged over the year 2006 obtained using the two-layer reactivity 5 scheme (top panel), the one-layer reactivity scheme (middle panel), and the default scheme with surface resistance $r_c = 2200$ s m⁻¹ (bottom panel). Note differences in scale.



5 Figure 9: Average relative difference (%) for the year 2006 between the ozone dry deposition velocity (v_d , cm s⁻¹) modelled using the one-layer reactivity scheme and the two-layer reactivity scheme (top panel), and that between the default scheme (with surface resistance $r_c = 2200$ s m⁻¹) and the two-layer reactivity scheme (bottom panel).



5 Figure 10: Modelled ozone dry deposition velocity (v_d , cm s⁻¹). The left panels are for January 2006 and right panels are for July 2006. Top panels: the two-layer reactivity scheme, middle panels: the one-layer reactivity scheme, and bottom panels: the default scheme with surface resistance $r_c = 2200$ s m⁻¹. Note differences in scale.



5 Figure 11: Top panel: Modelled annual ozone concentration (ppbv) for 2006 obtained using the two-layer reactivity scheme; middle panel: average relative difference (%) between the ozone concentration predicted using the one-layer reactivity scheme and the two-layer reactivity scheme; and bottom panel: average relative difference (%) between the ozone concentration predicted using the default scheme with surface resistance $r_c = 2200$ s m⁻¹ and the two-layer reactivity scheme.



Figure 12: Zonal distribution of ozone concentration (ppbv) for the year 2006. Top left: observed distribution based on the global 5 monthly mean vertical ozone profile database available from <u>http://www.bodekerscientific.com</u>; top right: the corresponding ACCESS-UKCA model distribution using the two-layer reactivity scheme; bottom left: the relative difference (%) between the modelled concentration determined using the two-layer reactivity scheme and the observations; and bottom right: the relative difference (%) between the modelled concentration determined using the default scheme and the observations.

