

## Referee #1 reply

The authors thank the anonymous referee #1 for the constructive comments and corrections. They helped us to improve our paper.

1. In the manuscript the sources of benzene and toluene along the Trans-Siberian railway were investigated mainly using the T/B ratios, and propene-equivalents and OFPs were also discussed. As emission sources of benzene and toluene include not only motor vehicle exhaust and industry emission, but also biomass burning, coal burning and gasoline evaporation, the source attribution need consider contributions from sources other than vehicle exhaust and industry emission.

**Reply:** The referee is certainly right. Biomass burning, coal burning and gasoline evaporation are very important sources of benzene and toluene emissions in the atmosphere. However, taking into account the measurements on a mobile platform, it is difficult to separate the data corresponding to the emissions from different sources (we will be very grateful for more exact recommendations). Therefore, taking into account the fact that vehicle exhaust is one of the main sources of benzene and toluene in anthropogenically polluted atmosphere, we use the well known criteria and the simultaneous measurements of CO, NO<sub>x</sub>, SO<sub>2</sub> and NMHC in the campaign to distinguish the impact of vehicle emissions as well as other local and regional-scale sources (including industrial emissions, coal burning and evaporative emissions) on benzene and toluene levels. This information is also very important for the immense Russian territory. It should be noted, however, that in TROICA-12 experiment no significant biomass burning along the Tran-Siberian railway was observed, so this emission source is not expected to impact significantly on benzene and toluene levels (see corrections at page 9, lines 14-16, and page 10, lines 30-32).

2. For the comparisons of the ozone formation rate from aromatics relative to that due to isoprene in the rural and urban areas, it should be noted that aromatic hydrocarbons are quite different from isoprene in reactivity and atmospheric lifetimes, and in their source regions and emission patterns. Therefore, it is important to figure out new understandings and new findings other than somewhat common senses like larger contributions of benzene and toluene to ozone formations in the urban areas.

**Reply:** It is well known that aromatic VOCs differ from isoprene by reactivity and sources. However, isoprene is one of the main biogenic VOCs forming ozone in the polluted surface air. So, it was important to reveal a significance of the benzene and toluene impact in ozone formation along the Trans-Siberian railway in contrast to isoprene. These estimations are very important because of a lack of such information for Russian regions. Although some new understandings are certainly necessary.

### Details

1. Page 3, the authors should give more details about where the mobile laboratory was located, in the front of the train or in the end or elsewhere? How to eliminate the interferences from emission inside the train or human activities in the train?

**Reply:** The location of the laboratory is described in the section 2.1 TROICA experiments, page 3, lines 4-6. To reduce a possible influence of emission from human activities in the train, all conveniences were placed at the end part of the train. Thus, their impact is expected to be generally non-significant (Panin et al., 2001). This information was added to the text (page 3, lines 7-10).

2. Page 4, the calibration method and frequencies for measuring VOCs by PTR-MS and APHA-360 should be stated.

**Reply:** Agreed. We provide more information about the instruments in the final version of the paper (see the sections 2.2 VOC measurements and 2.3 Other components and meteorology).

3. Page 8, lines 17 -20 were repeating lines 13-16

**Reply:** Agreed. Lines 13-16 are removed from the text.

4. Page 8, reference Karl et al., 2009 was not listed in the reference.

**Reply:** Agreed. The reference is added to the paper, page 15, line 14.

5. Page 8, section 3.2, as the benzene and toluene data were measured along the Trans-Siberian railway instead of at a fixed station, the discussions about diurnal variations of benzene and toluene should be careful. It is difficult to say whether emission sources, photochemistry or meteorological conditions had led to the variations. I'd like to suggest deleting this section.

**Reply:** The referee is certainly right about the influence of wide range of factors on the diurnal variations of benzene and toluene obtained from the measurements on a mobile laboratory. However, the statistics calculated for hourly mean values and presented in Fig. 4 allows some differentiating among these factors. Furthermore, the study of diurnal variations of benzene and toluene in the campaign is necessary to determine their contribution to spatial variability. The absence of significant diurnal variations of benzene and toluene allows us to neglect them and summarize the data in different spatial scales (for different regions and cities of Russia), as pointed out in the section 3.2.

6. **Page 9**, The correlations coefficients (R) between benzene, toluene, NMHC, CO, NO<sub>x</sub>, and SO<sub>2</sub> were all less than 0.6, that means their R<sup>2</sup> were all lower than 0.36. I don't think these can suggest the high or significant correlations between them. Section 3.3 should be rearranged.

**Reply:** As we know, the significance of the correlation coefficients is determined by t-test. According to our statistical calculations, all coefficients presented in Table 6 and shown by asterisks are statistically significant including the relationships between benzene, toluene, NMHC, CO, NO<sub>x</sub>, and SO<sub>2</sub>. Some of the coefficients are a little bit less than 0.5. It means that the strength of correlation is moderate (R= 0.3 to 0.5). The correlation coefficients for the relationships between benzene and toluene and CO and benzene in urban areas are > 0.5. The latter may evidence for (potentially) strong relationship (R= 0.5 to 1.0) between these compounds. Thus, the correlation analysis presented in section 3.3 allows us to see a relation between some compounds in urban atmosphere determining by their common pollution sources even in the large spatial scale. Furthermore, we noted that in rural areas correlation between all the species studied is very poor except for that between benzene and toluene. Some terminological corrections are included in the section 3.3 (page 10, lines 10-15).

7. Table 5 and Table 6, I think the authors wanted to list C<sub>7</sub>H<sub>8</sub> (Toluene) instead of C<sub>5</sub>H<sub>8</sub> (isoprene) as they discussed in section 3.3.

**Reply:** The referee is right. It is corrected.

8. Figures 3a, 3b and 8 in the manuscripts are really hard to read. I suggest plotting them in a different way.

**Reply:** The experience of previous presentations of a big TROICA dataset showed that it is rather difficult to present the data in a lot better way, however, we tried to improve the quality of the figures a little bit.

## Referee #2 reply

The authors thank the anonymous referee #2 for the constructive comments and corrections.

1. Have the authors considered plotting the concentrations in Figure 3 in a logarithmic rather than in a linear scale? In its current form, only the major concentration pikes can be identified from the figure, while potential differences in "background" concentration C1 ACPD Interactive comment Printer-friendly version Discussion paper between the different regions are very difficult to see.

**Reply:** Thank you for this recommendation. We plotted the concentrations in a logarithmic scale.

2. The authors could explain a bit more why and how the T/B ratio can be used as an indicator of chemical ageing. Does the chemical activity of toluene mentioned in the text refer to its OH-reactivity? Is OH the only important oxidant for benzene and toluene, and if not, what does this mean in terms of chemical ageing?

**Reply:** The chemical activity of toluene mentioned in the text certainly refers to its OH-reactivity. OH-radicals are the main oxidants of all VOCs in the atmosphere, especially at the daytime. Toluene has a shorter atmospheric lifetime than benzene due to its faster photochemical removal by OH (rate constants of  $1.23 \times 10^{-12}$  and  $5.96 \times 10^{-12}$   $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$  for benzene and toluene, respectively, at 298K by Atkinson and Arey, 2003). Thus, as toluene is more rapidly removed by oxidation, the T/B ratio decreases as air is transported over longer distances away from the pollution source. The chemical oxidation of benzene and toluene by  $\text{NO}_3$  and  $\text{O}_3$  is very weak ( $k \times 10^{-17}$   $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$  and  $k \times 10^{-18}$   $\text{cm}^3 \text{molecule}^{-1} \text{s}^{-1}$ , respectively for  $\text{NO}_3$  and  $\text{O}_3$ ) with faster toluene oxidation comparing with benzene as well.

3. I do not understand the meaning of statement "with similar relation for benzene" following Eq. 1.

**Reply:** Agreed. It is corrected.

4. The same sentence is repeated starting from lines 13 and 17 on page 8. Concerning the follow up of the latter sentence, one cannot infer from this information that benzene emission from motor vehicle exhaust is 25% lower than toluene emission. The logic here is incorrect!

**Reply:** Agreed. It is corrected, page 9, lines 4-5.

5. In the first sentence of page 9, do you mean "Observations in several locations have reported. . ."? If yes, then also the beginning of the next sentence need to be modified: "Contrary to these observations, no . . .".

A high pollution level itself cannot be a reason for the lack of observed diurnal cycle because, in principle, also high concentrations could be relatively evenly distributed. I would rather thing that a lack of strong local pollution sources (or lack of very high concentrations above the mixed layer) would be the reason.

The times are not usually given in a.m. or p.m (rather 04:00 and 23:00)

**Reply:** Agreed. It is corrected. Page 9, lines 22-28.

6. I do not think "meaningful correlation" is proper statistical language. Furthermore, I am not confident that  $R=0.6$  can be considered as a high correlation.

**Reply:**  $R > 0.5$  may be an evidence for (potentially) strong relationship ( $R= 0.5$  to  $1.0$ ) between the compounds observed. Some terminological corrections are included in the text (Section 3.3).

7. What is meant by "one-time maximum permissible concentration"? Also mentioned in section 3.1 and in abstract.

**Reply:** We meant a maximum concentration of the pollutant, short-term exposure of which (about 20 minutes) does not cause negative human effects. "Short-term exposure limit" would possibly be better. The corrections are added in the text.

8. Can the statement made on lines 20-23 considered general, as indicated here? These results are based on few measurement data points, so one would expect somewhat different numbers at some other time when travelling the same measurement route.

**Reply:** Agreed. It is corrected. Page 12, lines 21-24.

9. There are a few sentences that need to be re-written to make the text more understandable for the reader. A list of these sentences is given below.

**Reply:** Agreed. We tried to make them more understandable.

10. Finally, there are a number of minor grammatical issues that need to be corrected. Below is a list of suggestions for such corrections:

**Reply:** Agreed. We thank the referee for the corrections. All of them were included in the text.

# Benzene and Toluene in the surface air of North Eurasia from TROICA-12 campaign along the Trans-Siberian railway

5 Andrey I. Skorokhod<sup>1</sup>, Elena V. Berezina<sup>1</sup>, Konstantin B. Moiseenko<sup>1</sup>, Nikolay F. Elansky<sup>1</sup>, Igor B. Belikov<sup>1</sup>

<sup>1</sup>A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences, Moscow, 119017, Russia

*Correspondence to:* Andrey I. Skorokhod (askorokhod@mail.ru) and Elena V. Berezina (e\_berezina\_83@mail.ru)

**Abstract.** Volatile organic compounds (VOCs) were measured by proton transfer reaction – mass spectrometry (PTR-MS) on a mobile laboratory in a transcontinental TROICA-12 (21.07.2008 – 04.08.2008) campaign along the Trans-Siberian railway from Moscow to Vladivostok. Surface concentrations of benzene (C<sub>6</sub>H<sub>6</sub>) and toluene (C<sub>7</sub>H<sub>8</sub>) along with non-methane hydrocarbons (NMHCs), CO, O<sub>3</sub>, SO<sub>2</sub>, NO, NO<sub>2</sub> and meteorology are analyzed in this study to identify the main sources of benzene and toluene along the Trans-Siberian railway. The most measurements in the TROICA-12 campaign were conducted under low-wind/stagnant conditions in moderately (~78% of measurements) to weakly polluted (~20% of measurements) air directly affected by regional anthropogenic sources adjacent to the railroad. Only 2% of measurements were identified as characteristic of highly polluted urban atmosphere. Maximum values of benzene and toluene during the campaign reached 36.5 ppb and 45.6 ppb, correspondingly, which is significantly less than their short-term exposure limits (94 and 159 ppb for benzene and toluene, correspondingly). About 90% of benzene and 65% of toluene content is attributed to motor vehicle transport and 10% and 20%, correspondingly, provided by the other local and regional-scale sources. The highest average concentrations of benzene and toluene are measured in the industrial regions of the European Russia (up to 0.3 and 0.4 ppb for benzene and toluene, correspondingly) and south Siberia (up to 0.2 and 0.4 ppb for benzene and toluene, correspondingly). Total contribution of benzene and toluene to photochemical ozone production along the Trans-Siberian railway is about 16% compared to the most abundant organic VOC – isoprene. This contribution, however, is found to be substantially higher (up to 60–70%) in urbanized areas along the railroad suggesting important role of anthropogenic pollutant sources in regional ozone photochemistry and air quality.

## 1 Introduction

Aromatic compounds is an important class of hydrocarbons taking a significant part in the formation of tropospheric ozone and the secondary pollutants, including organic aerosols, that can lead to photochemical smog (Wayne, 2000; Baltaretu et al., 2009). Being toxic and carcinogenic, these gases, even without chemical transformation, have multiple impacts on the environment and human health (Derwent et al., 2000; Calvert et al., 2002; Lippmann, 2009). Thus, it is very important to study the role of aromatic compounds in air pollution for improving air quality.

A dominant source of aromatic compounds, particularly in urban environments, is automobile emissions with significant emissions being also from their use as solvents in various manufacturing processes (Wayne, 2000). Aromatic compounds can make up to 30-40% of the total nonmethane hydrocarbon (NMHC) content in the atmosphere, and be responsible for about 40% of the ozone producing potential of all NMHCs (Derwent et al., 2000; Calvert et al., 2002; Mugica et al., 2003; North et al., 2010).

In anthropogenically loaded regions, the most abundant aromatic compounds usually include benzene and toluene, which we focused our effort on in this study. They are often associated with air emissions from petroleum production industries (motor vehicle exhaust, incomplete combustion of fossil fuels, oil and gas service stations and other industrial and human activities). Filella et al., 2006, White et al., 2009 and Parra et al., 2006 also point out on the biogenic sources of benzene and toluene in summer rural conditions.

Many studies on benzene and toluene in urban and rural atmosphere have been carried out around the world (e.g. Keymeulen et al., 2001; Barletta et al., 2005; Parra et al., 2006; Velasco et al., 2007; Tivary et al., 2010; Miller et al., 2011; Civan et al., 2012; Chaudhary et al., 2012; Shaw et al., 2015). However, there is still a substantial lack of information on benzene and toluene abundance and spatial localization in remote areas of the North Eurasia. This gap is partially filled from the TROICA-12 campaign on a mobile carriage-laboratory in summer 2008 during which a complex measurements of surface air chemical composition were conducted along the Trans-Siberian railway from Moscow to Vladivostok (~ 9300 km long). This study aims to investigate variation of benzene and toluene in the surface air over Russia from PTR-MS measurements to determine their characteristic levels in urban, rural, and remote atmosphere in different geographic regions with respect to their major regional sources including large towns and industrial areas along the railroad. The relative importance of benzene and toluene emissions in photochemical near-surface ozone production is then assessed with respect to a major biogenic ozone precursor, isoprene. (Chameides et al., 1992; Geng et al., 2011).

## 2 Methods

### 2.1 TROICA experiments

TROICA experiments over Russia on a mobile laboratory have been carried out regularly since 1995 (Elansky et al., 2009). About 10 inorganic compounds (ozone, nitric oxides, carbon oxides, methane, sulphur dioxide etc.) as well as aerosols and

meteorological parameters (air temperature, pressure and humidity, solar radiation, vertical temperature profiles, wind speed and direction (at stops)) were measured continuously and simultaneously by a specially constructed automated system. The system was built on a railway carriage with air inlets at the height of about 4 m above the ground. VOC concentrations have been measured routinely since 2008 (TROICA-12, -13 and -14 campaigns). The TROICA carriage–laboratory is equipped in accordance with the measurement requirements of the Global Atmospheric Watch (WMO), and is located just after the electric locomotive to minimize various effects of near-surface air perturbations due to moving train. The possible impact of oncoming trains, e.g. those transporting oil products as well as human activities in the train (all conveniences were placed at the end part of the train) on the measurements has been removed through the respective time filtering of the original data. We also note that this impact is expected to be generally non-significant as demonstrated previously in Crutzen et al., 1996, Elansky et al., 2000 and Panin et al., 2001.

In present study, the data from the summer experiment TROICA-12 (21.07.2008 – 04.08.2008) along the Trans-Siberian railway (Fig.1) are analyzed. The train covers the total length of the route from Moscow to Vladivostok (~ 9288 km) for approximately 6 days, so the total duration of a single campaign is about two weeks. (Henceforth, we denote forward path from Moscow to Vladivostok, and return path from Vladivostok to Moscow, as east and west segments of the whole route, correspondingly).

Undoubtedly, the results of the observations at each particular location performed from the moving carriage are strongly influenced by specific weather conditions (synoptic patterns), as well as by complex interplay of local pollution sources and atmospheric transport by turbulent eddies on a variety of scales, the latter being considered as a source of random noise in the measurement data. Generally, straightforward quantification of the effects of atmospheric dilution and absolute strength of the associated nearby emission sources is inhibited in data analyses. Yet, the passage of each location twice (in the forward and return paths of the TROICA campaign) allows for some qualitative assessment of the impact of weather conditions, since the time between the two subsequent measurements is of the order of one week, which is comparable to the characteristic time of the boundary layer ventilation in mid-latitudes.

Various types of meteorological conditions along the railroad can be generalized into three distinctive weather patterns when traversing mountain area adjacent to Baikal Lake (~110° E) in east and west segments. There was clear and warm (>20° C at noon) weather on the route from Moscow to Baikal, hot weather (>24° C at noon) between Baikal Lake area and Vladivostok (east and west segments), and cool (daytime temperatures of 15 – 20° C) and rainy weather between Baikal Lake and Moscow on the return west segment of the route. Nighttime surface temperature inversions and stagnant air conditions were common for the east segment, although light winds were typical for the both east and west segments of TROICA. This feature is clearly seen in Fig.1 where 2-day back trajectories along the TROICA route calculated with the use of NOAA HIGHSPLIT model [Stein et al., 2015; Rolph, 2016] based on 3D wind fields are shown as averages of corresponding ensembles of trajectories originated within a height range from 0 – 400 m a.g.l. According to Fig.1, the characteristic distances of transport within a planetary boundary layer does not exceed 500 – 600 km in the two days preceding measurement time, so the measured chemical composition of the respective air masses can be considered as



representative of the cumulative impact of pollutant sources at local to regional scales. Relatively low wind speeds and short chemical lifetimes of the analyzed species (~12 days for benzene and ~2-4 days for toluene) support a regional approach to data analyzes implemented in present study (see Section 3) to quantify impact of various anthropogenic sources (large towns and industrial facilities) localized primarily along the railroad. The exception is the Far East Region where regional advection by southerly winds may contribute to measured pollutant concentrations from highly urbanized areas of the north-east China.

## 2.2 VOC measurements

VOC concentrations were measured by a Compact Proton Transfer Reaction Mass Spectrometer (Compact PTR-MS) from Ionicon Corp., Austria. One of the advantages of the Compact PTR-MS is a portability, which allows its using for continuous measurements on a mobile moving platform. Another big advantage of the Compact PTR-MS is that the volatile compound samples do not need to be specially prepared before the measurement, e.g. involving preconcentration procedures, thus headspace samples can be introduced directly into the reaction chamber consisting of the drift tube. The fact that, due to their low proton affinities,  $H_3O^+$  ions do not react with any of the major components present in clean air is an additional advantage as it allows the analysed air to be used directly as the buffer gas. To set a correct transmission in the software, a gas standard (Ionicon Analytic GmbH, Innsbruck) containing 17 VOCs including benzene (m79), toluene (m93) and isoprene (m69) was used. Error in calibration by the gas standard did not exceed 15%. The instrument allows the simultaneous on-line monitoring of 20 VOCs, including benzene, toluene, xylene, propilbenzene, isoprene, monoterpenes and others.

The Compact PTR-MS consists of three main components: an ion source, a drift tube, and a mass analyser (quadrupole mass spectrometer) in conjunction with an ion detector/amplifier.  $H_3O^+$  ions are produced in high concentrations from pure water vapour within a hollow cathode discharge ion source. These primary ions pass into the drift tube, where they undergo (mostly) non-dissociative proton transfer to the VOCs that are to be detected. The air sample to be analysed is introduced into the drift tube (which is maintained at about 2 mbar and 40 °C with a drift voltage of 600 V) close to its entrance at a typical flow rate of about 11 ml/min.

The Compact PTR-MS measurement range depends on the substances measured, integration time and system set-up. Its detection limits for the investigated VOCs are in order of pptv. The ions associated with benzene and toluene are free from significant interferences of other VOC species (Warneke et al., 2003). In the case of isoprene, other aldehydes and ketones, which are known to be detectable at this mass (Warneke et al., 2003; Fortner et al., 2009). However, isoprene has been found to be the dominant species at mass 69 within various kinds of air masses (de Gouw and Warneke, 2007). Nevertheless, the isoprene signal should be interpreted carefully, particularly from coniferous forests where emissions of 2-Methyl-3-buten-2-ol, which also determined at mass 69, take place.

## 2.3 Other components and meteorology

Concentrations of nonmethane hydrocarbons (NMHCs) were measured with an APHA-360 instrument (Horiba Company, Japan). This gas analyzer separates CH<sub>4</sub> and NMHC by using selective catalytic absorbers and measures the gas concentrations with a flame-ionization detector. The total error in the CH<sub>4</sub> and NMHC concentration measurements does not exceed ±5 ppb. To supply the flame ionization detector with hydrogen, which is necessary for its operation, the instrumentation set has hydrogen generators of different types. To make proper measurement accuracy, the zero calibration for every 20 minutes of the measurements and daily span calibrations were carried out in the campaign. The calibrations were performed using the gas standards provided by the D.I. Mendeleev All-Russian Institute for Metrology.

CO concentration was measured with a TE48S instrument (Thermo Electron Corp). This instrument uses an advanced method based on the measurements with the correlation of gas filters. It allows for measuring background CO concentrations at a level of less than 100 ppb with the total error of ±10 ppb.

SO<sub>2</sub> concentration was measured with an APSA-360 instrument (Horiba Company, Japan). This instrument measures SO<sub>2</sub> chemiluminescence induced by UV-radiation. For scheduled calibrations zero-air generator is used.

NO and NO<sub>2</sub> concentrations were measured at different times with a TE42C-TL instrument (Thermo Electron Corp., USA) and with a M200AU instrument produced by Teledyne Corp. (USA). These instruments apply the chemiluminescent method. The minimum NO and NO<sub>2</sub> concentrations detectable with these instruments are equal to 0.05 ppb, which makes it possible to measure the so-called background concentrations not influenced by the pollution sources. NO<sub>x</sub> concentration is the sum of NO and NO<sub>2</sub> concentrations.

Ozone concentration was measured with Dasibi 1008RS and 1008AH gas analyzers. These instruments are based on the photometric method. They allow measuring the ozone concentration in the range from 1 to 1000 ppb with a total error of ±1 ppbv. These instruments undergo scheduled calibrations against the secondary standard, the O3-41M No. 1294 instrument, which undergoes in its turn annual calibrations against the SRP No. 38 standard owned by the Mendeleev Research Metrology Institute (Russia).

For monitoring of meteorological parameters, the instrumentation set was constructed including the following instruments: acoustic anemometer ACAT-3M (Russia), Scanex (Russia), Driesen&Kem (Germany), and Vaisala (Finland). The temperature profile up to a height of 600 m was measured with MTP-5 meteorological temperature profiler (ATTEX Company, Russia).

## 2.4 Data processing

The major problem of the TROICA data analyses is a correct elimination of screening effects (relative to regional scale pollution sources) produced by local pollution sources along the railroad. Except for small areas of biomass burning and smoldering in the vicinity of railway, such sources are mainly of anthropogenic origin and characterized by highly limited spatial extents (and, hence, an impact upon a chemical composition), so they can be effectively filtered out by applying some objective criteria to the original 10-second dataset. Commonly, the chemical impact is characterized by strong short-time variations in CO, O<sub>3</sub>, and NO<sub>x</sub> concentrations and by an increase in NO/NO<sub>2</sub> ratio well above its regional characteristic value

of ~0.1. Upon a set of trials, we found that eliminating data points satisfying the criteria  $NO/NO_2 > 0.2$  was sufficient to obtain a robust statistics on measurement data concerning the regional effects of anthropogenic pollution. According to Fig. 2, such an approach results in eliminating ~25% data from the subsequent analyses with the remaining data (we call it “filtered dataset” henceforth), being safely attributed to the local to regional-scale effects produced by sustained anthropogenic sources.

Statistical and graphical data analysis was performed with a free software environment for statistical computing and graphics, R (<http://www.r-project.org/>), and with the data analysis and graphing software, OriginPro (OriginLab Corporation).

A statistical summary of the filtered dataset is shown in Table 1. In the east segment, toluene and benzene concentrations and their variations are somewhat higher compared to those for the west segment. As discussed above, this feature is most probably attributed to the observed differences in meteorological conditions, as the regional anthropogenic emission sources are not expected to vary at any appreciable rate on sub-seasonal scale. Although atmospheric dilution rates and advection paths (see Figure 1) vary significantly between the east and west segments, spatial localization of high levels of toluene and benzene near their emission sources (mostly large towns and their suburbs) strongly restricts the impact of meteorology on their near-surface abundance, at least at regional scale, due to their relatively short chemical lifetime. Consequently, the observed systematic differences in toluene and benzene statistics between east and west segments were relatively small for both mean and percentile values.

### 3 Results and discussion

#### 3.1 Spatial distribution of benzene and toluene along the Trans-Siberian railway

The areas adjacent to the Trans-Siberian railway are markedly different in amount of urbanization and anthropogenic load. Eighty-seven towns are located immediately on the railway, sixty-eight towns are in the Ural mountain region and West Siberia with the remaining ones located in the East Siberia and the Far East. Yet, it is in the area of first tens to hundreds of kilometers from the Trans-Siberian railway where the most significant regional anthropogenic sources are commonly located in all the regions considered. Fig. 3 shows 10-min averages of the surface concentrations of benzene and toluene derived from the filtered 10-second dataset for the east and west segments of the TROICA-12 route. Simultaneous measurements of surface NMHCs, CO, and SO<sub>2</sub> concentrations are also shown in the figure.

The observed simultaneous peaks in the concentrations of VOCs and inorganic compounds shown in the figure are spatially connected to the most significant regional anthropogenic sources along the railway (mostly in urban environment), which are mainly expected to be motor vehicle transport, industry, central heating, and power plants.

Although exact quantitative input of various types of toluene and benzene sources into the observed concentrations remains unknown, some qualitative assessment of their relative importance can be done from a ratio of toluene to benzene (T/B). The

T/B ratio is frequently used as an indicator of motor transport emissions, since toluene content in the gasoline and automobile exhausts is in 3–4 times higher than the benzene content (D. Brocco et al., 1997). Therefore,  $T/B \approx 1-3$  is widely accepted to indicate motor vehicle transport, whereas  $T/B > 3$  points out to industrial exhausts (Tiwari et al., 2010; Shaw et al., 2014; Carballo-Pat, 2014). Furthermore, the chemical activity of toluene in the atmosphere is approximately 5 times higher than that of benzene. Hence, the T/B ratio can serve as indicator of chemical aging and, consequently, proximity of the sampled air to the associated pollution source (Mugica et al., 2003; Tiwari et al., 2010; Carballo-Pat, 2014; Shaw et al., 2014).

Following our previous study on greenhouse gases emissions (Berezina et al., 2014), in present analysis, we divide the whole path from Moscow to Vladivostok into 6 lengthy segments according to climatological conditions and anthropogenic load intensity: European Russia (ER), Ural mountain region (UR), southern parts of West (SWS), Central (SCS), and East (SES) Siberia, and Far East (FE) (see Fig.1). Statistics for benzene, toluene, and T/B from 10-second filtered data for different regions are shown in Table 2. The highest concentrations of benzene and toluene were observed in ER, UR, and SCS, which are the regions of most significant anthropogenic emission sources and proximity of the TROICA route to the strong pollution sources. Since low/high T/B ratio measured at a given location can be equally attributed (in the absence of a prior information) either to a photochemically aged/young air mass or specific chemical composition of the primary pollutants affected the measured air mass, we must distinguish between the both factors to use T/B ratio as characteristic of the associated emission source. The problem is further complicated by that the measured air represents commonly a mixture of air parcels with different photochemical age and/or anthropogenic loading. One partial solution consists in separating air masses according to their chemical aging (transport times from the regional pollution sources) based on some additional data on their chemical composition and/or transport times.

In present work we utilized the former approach by invoking contemporary measurements of NO and NO<sub>2</sub> to distinguish between clean remote air ( $NO_x < 2$  ppb), moderately polluted air ( $2 \leq NO_x < 20$  ppb) and highly polluted air ( $NO_x \geq 20$  ppb), the latter being representative for urban and suburban environment. Here, the exact threshold values of NO<sub>x</sub> were chosen based on our experience in processing multiple data sets from TROICA campaigns as well as continuous measurements of NO<sub>x</sub> at ZOTTO Tall Tower – remote site in Central Siberia, which is occasionally affected by transport of polluted air from major regional anthropogenic sources in the south Siberia (Vasileva et al., 2011). The statistical analysis of the filtered dataset showed that about 80% of the data accounts for moderately polluted air, 18% - for clean remote air and only about 2% - for highly polluted air. Thus, the measurements in TROICA-12 campaign were performed mostly in moderately polluted urban atmosphere ( $2 \leq NO_x < 20$  ppb), with maximum concentrations of toluene and benzene reaching the values of 45.6 ppb and 36.5 ppb, correspondingly, which is significantly less than the short-term exposure limits (94 and 159 ppb for benzene and toluene, correspondingly).

Since high ( $\geq 2$  ppb) NO<sub>x</sub> are found to be a robust characteristic of photochemically young air (Vasileva et al., 2011), we use associated T/B ( $NO_x > 2$  ppb) values to infer some qualitative information on the pollution sources characteristic for the given region. According to Table 3, average and median values of T/B ( $NO_x > 2$  ppb) were in the ranges of 1.8–2.6 and 1.1–1.6,

respectively, which is well below commonly accepted threshold value for the motor vehicle exhausts ( $T/B < 3$ ). Yet, significantly higher T/B ratios (the column  $P_{90}$  of the table) were found in all the regions, suggesting other important regional sources of air contamination. These high values were measured commonly as short-lasting events where the mobile laboratory crossed the plumes of strongly contaminated air originated from large upwind emission sources, as evidenced from air trajectory analyses and a prior data on industrial facilities. In such cases, high toluene and benzene concentrations are accompanied with enhanced levels of NMHC, CO, and SO<sub>2</sub> as well, indicating the petrochemical and refining exhausts. These include heat and power plants in the vicinities of large towns: Perm, Tyumen, Omsk, Ulan-Ude, Chita, Khabarovsk (Fig.3a), and Ekaterinburg, Tyumen, Krasnoyarsk, Kansk, Irkutsk, Mogocho, Birobidzhan, Khabarovsk (Fig.3b). Assuming that high ( $>2$  ppb) NO<sub>x</sub> is a signature of freshly contaminated air, we estimated the bulk contribution of motor vehicle exhausts,  $\kappa(T)$ , to the near-surface abundance of toluene and benzene along the Trans-Siberian railway from a simple relation:

$$\kappa(T) = \frac{T_{low} \cdot t_{low}}{T_{low} \cdot t_{low} + T_{htgh} \cdot t_{htgh}} \cdot 100\% \quad (1)$$

where  $t_{low}$ ,  $t_{high}$  are the fractions of measurement time within air masses having low ( $< 3$  ppb) and high ( $\geq 3$  ppb) T/B ratios and  $NO_x > 2$  ppb ( $t_{low} + t_{high} = 1$ ), and  $T_{low}$ ,  $T_{high}$  are average toluene concentrations in the low- and high-T/B air masses, respectively (see Table 4).

5 One can see from Table 4, that motor vehicle exhausts were responsible for  $\sim 90\%$  of benzene levels and  $\sim 65\%$  of toluene levels during the campaign, that is, toluene emission from motor vehicle exhausts is 25% lower than the benzene one. It is comparable with the relative source contributions for benzene, and toluene presented in (Karl et al., 2009). Thus, the motor vehicle exhausts was the most significant anthropogenic source of air pollution by toluene, and benzene in all the areas adjacent to the TROICA-12 route (densely populated areas along the Trans-Siberian Railroad).

10 Benzene levels from the TROICA-12 campaign are broadly comparable in magnitude with other published data on their abundance in the summer urban and rural continental surface air (Elansky et al., 2000; Barletta et al., 2005; Na et al., 2005; Parra et al., 2006; Hoque et al., 2008; Tiwary et al., 2010; Seco et al., 2013; Wagner et al., 2014). However, toluene levels tended to be lower than those reported in earlier publications. It is possibly due to the more significant contribution of evaporative and industrial emissions in toluene levels (Karl et al., 2009) than this of mobile transport exhausts (which are about 65% for toluene from TROICA-12 measurements). It should be noted that in the campaign no significant biomass  
15 burning along the Tran-Siberian railway was observed, so this emission source is not expected to impact significantly on benzene and toluene levels.

### 3.2 Diurnal variations of benzene and toluene

To determine the contribution of diurnal variations of benzene and toluene to their surface levels and spatial variability, we analyzed their hourly mean concentrations measured in the campaign.

20 Observations in several locations (Filella et al., 2006, Zalel et al., 2008, Tiwary et al., 2010; Wagner et al., 2014) reported the highest levels of benzene and toluene being at the morning and evening hours due to increase in the motor vehicle transport exhausts at this time. Contrary to these studies, no clear diurnal variation of benzene and toluene was observed in the TROICA-12 campaign (Fig.4). We suppose that it is due to spatially smoothing of their diurnal variations in conditions of a lack of strong local pollution sources, which is clearly seen from median values. Somewhat higher levels in the morning  
25 (at 04-05 LST (local standard time)) and in the evening (at 20 and 23 LST) are most probably due to accumulation of benzene and toluene in the stable atmospheric conditions in the vicinities of their regional sources transected on the TROICA route. Episodic crossing of anthropogenic pollution plumes during the campaign caused the midday peak (which falls occasionally at  $\sim 12$  LST), seen on the Figure 4 from  $P_{90}$  profile.

30 Thus, diurnal variations of planetary boundary layer mixing regime did not contribute significantly to the spatial variability of benzene and toluene along the Trans-Siberian railway in the TROICA-12 campaign. It allows us to neglect them further summarizing the data in different spatial scales (for example, by different Russian regions and cities).

### 3.3 Benzene and toluene levels in urban and rural surface air

We found that the surface concentrations of VOCs, NO<sub>x</sub>, CO, and SO<sub>2</sub> were, in general, notably higher in urban areas (as it would be expected), as most of regional pollution sources are located in cities and their suburbs (Table 5). It was found, however, that the highest (>P95) concentrations of all the pollutants including benzene and toluene were measured outside the cities so they can not be attributed to direct impact of urban pollution sources. A closer examination showed that these events were most probably connected to specific transport conditions favorable for maintaining anthropogenic plumes from large upwind sources, i.e. towns and industrial manufactures away from the railroad, as highly coherent structures at time scales of the order of few to ten hours. The most prominent events of crossing industrial plumes took place in ETR (up to 37 ppb for benzene), SCS (up to 39 ppb for benzene), SES (up to 46 ppb for toluene) and FE (up to 41 ppb for benzene).

10 One can see from Table 6 that there is a statistically significant moderate correlation ( $R \approx 0.5$ ) between benzene and toluene both in urban and rural areas. In urban environments, there is a stronger correlation between benzene and toluene and CO ( $R \approx 0.6$ ), as well as between NO<sub>x</sub> and CO, which can point out to the motor vehicle transport as their main common emission source. In urban areas, benzene is also in a significant moderate correlation with NMHC, which can indicate partial input from the industrial exhausts: hydrocarbon processing, refining industry, fuel transportation and storage, tanks and pipeline  
15 leaks, etc.

In rural areas (identified as those outside the towns with additional constraint NO<sub>x</sub> < 0.2 ppb, see p.3.1), correlation between all the species studied was very poor, except for that between benzene and toluene. Evidently, the diversity of transport pathways, photochemical aging, as well as irreversible mixing of air masses subjected to different rates of anthropogenic contamination precludes direct quantification of primary pollutant sources for the substantial part of the TROICA route away  
20 from areas of their immediate impact.

To study atmospheric pollution in Russian cities along the Trans-Siberian railway, 29 cities were selected, for which the total amount of measurement time was at least 25% of the whole residence time in the city. The highest concentrations of benzene and toluene (up to 5 ppb) were observed in the industrial towns: Perm, Kirov (European Russia); Kungur and Yekaterinburg (south Ural mountain region), Tyumen (West Siberia), Angarsk, Irkutsk, and Ulan-Ude (East Siberia), Birobidzhan,  
25 Khabarovsk (Far East region) (Fig. 5). In these cities the highest levels of NMHC, NO<sub>x</sub>, and CO were also measured (Fig.6). Evidently, high CO abundances found in some of these towns point out to the significant contribution to the overall pollution rates from refineries and central heating and power plans, which was also confirmed by the high T/B ratios (> 3 – 4 basing on P90 regional values). The specific T/B ratios also indicate that Khabarovsk, Birobidzhan, Skovorodino, Tulun, Tajshet, and Tyumen are mainly polluted by industrial emissions, whereas Vladimir, Kungur, Yurga, and Krasnoyarsk – by transport  
30 exhausts. In other cities, motor vehicle transport was found to be a main pollution source, even though the contribution from other sources (including industrial emissions, coal burning and gasoline evaporation) was also important, as seen from the significantly higher P90 values comparing to the average one calculated for rural regions (see Fig.5). Benzene and toluene

surface levels in the medium-sized towns of Siberia (Achinsk, Taishet, Nizhneudinsk, et al.) were close to an average rural concentration calculated for the campaign (less than 0.3 ppb).

Unfortunately, exact quantification of inputs from various types of sources into anthropogenic contamination of urban air was inhibited when using the TROICA data due to the very limited amount of observations collected within a particular town. Considering that transport emissions occur when the T/B is in the range from 1 – 3 (Mugica et al., 2003; Tiwari et al., 2010; Carballo-Pat, 2014; Shaw et al., 2014), and supposing well mixed conditions such that each measured air parcel represent a uniform mixture of pollutants from various sources within a town, we found that motor vehicle transport accounts for approximately 75% of anthropogenic emissions in the Russian cities along the Trans-Siberian railway with the remaining 25% are attributed to other sources (Fig.7). These estimates correspond well to those derived in section 3.1 based on the whole TROICA dataset. As seen from Figure 8, T/B ratio for vehicle urban exhausts in the Russian cities along the Trans-Siberian railway is usually in the range of 2.3–2.8.

### 3.4 Contribution of VOCs to ozone formation potential over Russia

Along the whole route of the TROICA campaign, the lower troposphere chemical regime was found to be essentially a NO<sub>x</sub> sensitive, both in rural and urban environments, with typical morning NMHC/NO<sub>x</sub> ratios being well above 20. Hence, ozone production rates are expected to be controlled by regional NO<sub>x</sub> emissions (Silman, 1999) and their complex interplay with both natural and anthropogenic sources of VOCs. As mentioned above, the meteorological conditions during the most of the TROICA campaign were favorable for both studying chemical composition of fresh air masses contaminated by regional sources as well as for ozone production from the emitted precursors due to high daytime surface air temperatures and solar radiation. To estimate the impact of the measured VOCs on regional ozone production, we employed the widely used quantities: propylene-equivalent concentration (PE) and ozone-forming potential (OFP) (Carter, 1994; So and Wang, 2004), which utilize the measured concentrations of VOCs along with their reactivity with hydroxyl radical. These coefficients are defined as:

$$PE_{\text{VOC}} [\text{ppbC}] = C_{\text{VOC}} \times k_{\text{OH,VOC}} / k_{\text{OH,propylene}}, \quad (2)$$

$$OFP_{\text{VOC}} [\mu\text{g}/\text{m}^3] = C_{\text{VOC}} \times MIR_{\text{VOC}}, \quad (3)$$

where  $C_{\text{VOC}}$  is a VOC concentration having the dimension of ppbC and  $\mu\text{g}/\text{m}^3$  in (2) and (3), respectively,  $k_{\text{OH,VOC}}$  is the rate constant for the reaction of VOC with OH-radical,  $k_{\text{OH,propylene}}$  is the rate constant for the reaction between OH and propylene, and  $MIR_{\text{VOC}}$  is a maximum incremental reactivity. The latter is a dimensionless quantity defined as gram of O<sub>3</sub> produced per gram of the VOC, which equals to the maximum ozone concentration formed from chemical destruction of the given VOC.

We calculated PE and OFP values for benzene and toluene based on the daytime observations from 12 a.m. – 5 p.m., the time for which the highest correlations between ozone and its precursor species were observed in TROICA as well. The calculated PE and OFP values were compared against those for isoprene, the latter being known as one the most important



biogenic ozone precursors in rural as well as urban settings (Chameides et al., 1988; Fuentes et al., 2000; Wagner et al., 2014).

As seen from Table 7, the average value of OFP of isoprene along the TROICA route is much higher compared to those for benzene and toluene, owing to relatively high near-surface abundances of isoprene (approx. 3 times as much as that for the sum of benzene and toluene) and its exceptionally high reactivity with hydroxyl radical. According to Fig.8, the process of oxidizing of isoprene proves to be the most important chemical source of ozone in all the regions along the TROICA route, as it could be expected if one takes into account that the most part of the railway crosses the areas with very weak to moderate anthropogenic load. The highest OFPs of isoprene seen on the figure in the Far East are due to its high biogenic emissions from broad-leaved forests as well as high surface air temperatures measured in this region in the both east and west segments of the route.

As seen from figure 8, the OFPs of benzene and toluene do not have significant large-scale spatial variations along the railway, contrary to that for isoprene. The highest OFPs shown as peaks on figure 8 are spatially connected to large towns and their vicinities along the railway where the relative input of benzene and toluene into ozone production reaches as high as 60–70% compared to that of isoprene. This supports our general notion of the reduced impact of regional anthropogenic sources on the regional ozone budget compared to long-range advection and regional biogenic VOC emissions (Shtabkin et al., 2016). Taking into account the important role of biogenic emissions of isoprene in the regional ozone photochemistry, the detailed analyses of isoprene observations from the TROICA campaigns should be done in a separate publication.

#### 4 Summary

Surface concentrations of important anthropogenic VOCs, benzene and toluene, as well as inorganic compounds were measured simultaneously along the Trans-Siberian railway on a mobile railway laboratory in the TROICA-12 campaign in summer 2008. It is demonstrated that the TROICA-12 measurements were carried out mostly in moderately polluted ( $2 \leq \text{NO}_x < 20$  ppb) environment (~78% of measurements) with the remaining part of measurement time divided between weakly polluted ( $\text{NO}_x \leq 2$  ppb) and highly polluted ( $\text{NO}_x > 20$  ppb) urban environment (20 and 2% of measurements, correspondingly). Maximum values of benzene and toluene during the campaign reached 36.5 ppb and 45.6 ppb, correspondingly, which is significantly less than their short-term exposure limits (94 and 159 ppb for benzene and toluene, correspondingly). Although, the weather conditions during the major part of the TROICA campaign were favorable for accumulating anthropogenic pollutants in the lower atmosphere, the absence of clear diurnal variations of benzene and toluene along with their low abundances apart from the immediate vicinity of large towns and industrial manufactures points out to mostly unpolluted air conditions along the Trans-Siberian railway during the campaign.

We estimated that motor vehicle exhausts accounts for ~ 90% of benzene levels and ~ 65% of toluene levels during the campaign, with the remaining 10% and 25%, correspondingly, provided by other important regional anthropogenic sources: industrial enterprises, coal burning and evaporative emissions of VOCs etc.

5 The highest near-surface abundances of benzene and toluene, both in urban environment and on the regional scale, were observed in areas with the highest anthropogenic burden. They are industrial regions of the Southern Urals, European Russia, and the southern part of Central Siberia, where spatially averaged benzene and toluene levels, representative for rural conditions, equal to ~0.3 and ~0.4 ppb, respectively. Vehicle emissions constitute the major part of total anthropogenic pollution in these regions, with typical ration of T/B be 2.2 – 2.3. Similarly, the highest concentrations of benzene (up to 5 ppb) and toluene (up to 7 ppb) along with high levels of NMHC, CO and NO<sub>x</sub> were observed in the following industrial towns: Perm, Kirov (European Russia); Kungur and Yekaterinburg (south Ural mountain region), Tyumen (West Siberia), Angarsk, Irkutsk, and Ulan-Ude (East Siberia), Birobidzhan, Khabarovsk (Far East region).

10 Considering that transport emissions occur when the T/B is in the range from 1 – 3 and supposing well mixed conditions such that each measured air parcel represent a uniform mixture of pollutants from various sources within a town, we found that motor vehicle transport accounts for approximately 75% of anthropogenic emissions in the Russian cities along the Trans-Siberian railway with the remaining 25% are attributed to industrial sources. T/B ratio for vehicle urban exhausts in the Russian cities along the Trans-Siberian railway is usually in the range of 2.3–2.8.

15 The contribution of benzene and toluene to the local photochemical ozone production along the Trans-Siberian railway is generally not significant compared to biogenic VOCs in rural environment and reaches as much as 16% of that of isoprene. However, in large towns the contribution of benzene and toluene to ozone formation reaches 60–75% compared to isoprene, supporting important role of anthropogenic sources in local pollution.

*Acknowledgements.* The authors thank Shumsky R.A. for an active participation in designing of the measurement set of the mobile laboratory and controlling of its correct work and Lavrova O.V. for an active participation in the campaign and careful diary observations. This study was supported by the Russian Science Foundation (grant no. 14-47-00049), by the Russian Foundation for Basic Research (grant no. 16-35-00158) and contributes to the Pan-Eurasian Experiment (PEEX) Program research agenda.

## References

- 30 Baltaretu, C. O., Lichtman, E. I., Hadler, A. B., and Elrod, M. J.: Primary atmospheric oxidation mechanism for toluene., J. Phys. Chem. A, 113, 221–30, doi:10.1021/jp806841t, 2009.
- Barletta, B., Meinardi, S., Rowland, F.S., Chan, C.Y., Wang, X.M., Zou, S.C., Chan, L.Y., Blake, D.R.: Volatile organic compounds in 43 Chinese cities. Atmospheric Environment 39 (32): 5979-5990, 2005.

- Berezina, E.V., Elansky, N.F., Moiseenko, K.B., Safronov, A.S., Skorokhod, A.I., Lavrova, O.V., Belikov, R.A., Shumsky, I. B. Estimation of biogenic CH<sub>4</sub> and CO<sub>2</sub> emissions and dry deposition of O<sub>3</sub> using <sup>222</sup>Rn measurements in TROICA expeditions // *Izvestiya. Atmospheric and Oceanic Physics*, vol. 50, no. 6, pp. 583-594, 2014.
- Calvert, J. G., Atkinson, R., Becker, K. H., Kamens, R. M., Seinfeld, J. H., Wallington, T. J., Yarwood, G., Eds. The mechanisms of atmospheric oxidation of aromatic hydrocarbons; Oxford University Press: Oxford, U.K., 2002.
- Carballo-Pat., C.G., Cerón-Bretón, J.G., Cerón-Bretón, R.M., Ramírez-Lara, E., Aguilar-Ucán, C.A., Montalvo-Romero, C., Guevara-Carrió, E., Córdova-Quiroz, A.V., Gamboa-Fernández, J.M., Uc-Chi, M.P.: Latest trends in Energy. Environment and Development. Proceedings of the 7-th International Conference on Environmental and Geological Sciences and Engineering (EG'14). Salerno. Italy. June 3-5, 132-140, 2014.
- Chameides, W.L., Fehsenfeld, M.O. Rodgers, C. Cardelino, J. Martinez, D. Parrish, W. Lonneman, D.R. Lawson, R.A. Rasmussen, P. Zimmerman, J. Greenberg, P. Middleton, and T. Wang: Ozone precursor relationship in the ambient atmosphere, *Journal of Geophysical Research-Atmospheres*, 97 (D5), 6037- 6055, 1992.
- Chaudhary, S. and Kumar, A.: Monitoring of benzene, toluene, ethylbenzene and xylene (BTEX) concentrations in ambient air in Firozabad, India. *Int Arch ApplSciTechnol* 3(2): 92–96, 2012.
- Civan, M.Y., Kuntasal, O.O., Tuncel, G.: Source apportionment of ambient volatile organic compounds in Bursa, a heavily industrialized city in turkey. *Environ. Forensics* 12 (4), 357-370, 2011.
- Crutzen, P.J., N.F. Elansky, M. Hahn, G.S. Golitsyn, C.A.M. Brenninkmeijer, D. Scharffe, I.B. Belikov, M. Maiss, P. Bergamaschi, T. Rockmann, A.M. Grisenko and V.V. Sevastyano. Trace gas measurements between Moscow and Vladivostok using the Trans-Siberian Railroad // *J. Atm. Chemistry*, No 29, 179–194, 1998.
- de Gouw, J. and Warneke, C.: Measurements of volatile organic compounds in the Earth's atmosphere using proton-transferreaction mass spectrometry, *Mass Spectrom. Rev.*, 26, 223–257, doi:10.1002/mas.20119, 2007.
- Derwent, R.G., Davies, T.J., Delaney, M., Dollard, G.J., Field, R.A., Dumitrean, P., Nason, P.D., Jones, B.M.R., Pepler, S.A.: Analysis and interpretation of the continuous hourly monitoring data for 26 C<sub>2</sub>-C<sub>6</sub> hydrocarbons at 12 United Kingdom sites during 1996. *Atmospheric Environment* 34, 297-312, 2000.
- Elansky, N.F., Belikov, I.B., Berezina, E.V.: Atmospheric composition observations over Northern Eurasia using the mobile laboratory: TROICA experiments, edited by N.F. Elansky, Moscow, 2009, 73 p.
- Elansky, N. F.: Observations of the atmospheric composition over Russia using a mobile laboratory: the TROICA experiments // *International Global Atmospheric Chemistry. Newsletter*, No 37, 31–36, 2007.
- Elansky, N. F., G. S. Golitsyn, T. S. Vlasenko, and A.A. Volokh: Concentrations of Volatile Organic Compounds in Surface Air along the Trans-Siberian Railroad//*Izvestiya, Atmospheric and Oceanic Physics*, V. 37, Suppl. 1, S10–S23, 2001a.
- Elansky, N.F., G.S. Golitsyn, T.S. Vlasenko, and A.A. Volokh: Volatile organic compounds observed in the atmospheric surface layer along the Trans-Siberian Railroad//*Dokl. Akad. Nauk*, V. 373, No 6, 816–821, 2000.
- Filella, I., Penñuelas, J.: Daily, weekly, and seasonal time courses of VOC concentrations in a semi-urban area near Barcelona. *Atmospheric Environment* 40, 7752–7769, 2006.

- Fortner, E. C., Zheng, J., Zhang, R., Berk Knighton, W., Volkamer, R. M., Sheehy, P., Molina, L., and André, M.: Measurements of Volatile Organic Compounds Using Proton Transfer Reaction – Mass Spectrometry during the MILAGRO 2006 Campaign, *Atmos. Chem. Phys.*, 9, 467-481, doi:10.5194/acp-9-467-2009, 2009.
- Fuentes, J.D., M. Lerda, R. Atkinson, D. Baldocchi, J.W. Bottenheim, P. Ciccioli, B. Lamb, C. Geron, L. Gu, A. Guenther, T.D. Sharkey, W. Stockwell: Biogenic hydrocarbons in the atmospheric boundary layer: A review *Bulletin of the American Meteorological Society*, pp. 1537–1575, 2000
- Geng, F., Tie, X., Guenther, A., Li, G., Cao, J., and Harley, P.: Effect of isoprene emissions from major forests on ozone formation in the city of Shanghai, China, *Atmos. Chem. Phys.*, 11, 10449– 10459, doi:10.5194/acp-11-10449-2011, 2011.
- Guerra, G., Iemma, A., Lerda, D., Martines, C., Salvi, G., and Tamponi, M.: Benzene Emissions from Motor-Vehicle Traffic in the Urban Area of Milan – Hypothesis of Health Impact Assessment, *Atmos. Environ.*, 29, 3559–3569, doi:10.1016/13522310(95)00205-D, 1995.
- Hoque, R. R., P. S. Khillare, et al. "Spatial and temporal variation of BTEX in the urban atmosphere of Delhi, India." *Science of The Total Environment* 392(1): 30-40, 2008
- Karl, T., Apel, E., Hodzic, A., Riemer, D. D., Blake, D. R., Wiedinmyer, C.: Emissions of volatile organic compounds inferred from airborne flux measurements over a megacity, *Atmos. Chem. Phys.*, 9, 271–285, doi:10.5194/acp-9-271-2009, 2009.
- Kalogridis C, Gros V, Sarda-Estevé R, Langford B, Loubet B, Bonsang B, Bonnaire N, Nemitz E, Genard AC, Boissard C: Concentrations and fluxes of isoprene and oxygenated VOCs at a French Mediterranean oak forest. *AtmosChem Phys* 14: 10085–10102, 2014
- Keymeulen, R., Gögényi, M., Héberger, K, Priksane, A, and H.V. Lagenhove: Benzene, toluene, ethylbenzene and xylenes in ambient air and Pinussylvestris L. needles: a comparative study between Belgium. Hungary and Latvia. *Atmos. Environ.* Vol. 35, pp. 6327-6335, 2001.
- Lippmann M., *Environmental Toxicants: Human Exposures and Their Health Effects*: Wiley, 2009.
- Martín-Reviejo, M. and Wirtz, K.: “Is benzene a precursor for secondary organic aerosol?” *Environ. Sci. Technol.* 39, 1045–1054, 2005.
- Miller L., Xu, X.H., Wheeler, A., Atari, D.O., GrgicakMannion, A., Luginaah, I.: Spatial Variability and Application of Ratios between BTEX in Two Canadian Cities, *The Scientific World Journal*, 11, 2536-2549, 2011.
- Mugica, V., Ruiz, M.E., Watson, J., and J. Chow: Volatile Aromatic Compounds in Mexico City atmosphere: Levels and Source apportionment. *Atmosfera*. Vol. 16, 15-27, 2003.
- Na, K., Moon, K., Kim, Y.P.: Source contribution to aromatic concentration and ozone formation potential in the atmosphere of Seoul. *Atmospheric Environment* 39, 5517-5524, 2005.
- Ng N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and J. H. Seinfeld: “Secondary organic aerosol formation from m-xylene, toluene, and benzene,” *Atmos. Chem. Phys.* 7, 3909–3922, 2007.
- North G.R., Pyle J. and Zhang F. Ed.: *Encyclopedia of Atmospheric Sciences*, Second Edition, 90–109, 2015.

- Panin L. V., Elansky N. F., Belikov I. B., Granberg I. G., Andronova A. V., Obvintsev Yu. I., Bogdanov V. M., Grisenko A. M., and Mozgrin V. S: Estimation of Reliability of the Data on Pollutant Content Measured in the Atmospheric Surface Layer in the TROICA Experiments//Izvestiya, Atmospheric and Oceanic Physics, V. 37, Suppl. 1, 81-91, 2001.
- Parra, MA, González, L, Elustondo, D, Garrigó, J, Bermejo, R, Santamaría, J.M.: Spatial and temporal trends of volatile organic compounds (VOC) in a rural area of Northern Spain. *Sci Total Environ*, 370:157–67, 2006.
- Peñuelas, J. and Staudt, M.: BVOCs and global change, *Trends Plant Sci.*, 15, 133–144, doi:10.1016/j.tplants.2009.12.005, 2010.
- Rolph, G.D.: Real-time Environmental Applications and Display System (READY) Website (<http://www.ready.noaa.gov>). NOAA Air Resources Laboratory, College Park, MD, 2016.
- Seco, R., Peñuelas, J., Filella, I., Llusia, J., Schallhart, S., Metzger, A., Müller, M., and Hansel, A.: Volatile organic compounds in the western Mediterranean basin: urban and rural winter measurements during the DAURE campaign, *Atmos. Chem. Phys.*, 13, 4291–4306, doi:10.5194/acp-13-4291-2013, 2013.
- Shaw, M. D., Lee, J. D., Davison, B., Vaughan, A., Purvis, R. M., Harvey, A., Lewis, A. C., and Hewitt, C. N.: Airborne determination of the temporo-spatial distribution of benzene, toluene, nitrogen oxides and ozone in the boundary layer across Greater London, UK, *Atmos. Chem. Phys.*, 15, 5083-5097, doi:10.5194/acp-15-5083-2015, 2015.
- Sillman, S.: The relation between ozone, NO<sub>x</sub> and hydrocarbons in urban and polluted rural environments, *Atmos. Environ.*, 33, 1821–1845, doi:10.1016/S1352-2310(98)00345-8, 1999.
- Timkovsky, I.I., Elanskii, N.F., Skorkhod, A.I. and Shumskii, R.A.: Studying of biogenic volatile organic compounds in the atmosphere over Russia. *Izvestiya Atmospheric and Oceanic Physics* 46: 319-327, 2010.
- Tiwari, V., Y. Hanai, et al.: "Ambient levels of volatile organic compounds in the vicinity of petrochemical industrial area of Yokohama, Japan." *Air Qual Atmos Health* 3(2): 65-75, (2010).
- Vasileva, A.V., Moiseenko, K.B., Mayer, J.-C., Jurgens, N., Panov, A., Heimann, M., Andreae, M.O: Assessment of the regional atmospheric impact of wildfire emissions based on CO observations at the ZOTTO tall tower station in central Siberia // *J.Geophys. Res.*, 116, D07301, 2011, doi:10.1029/2010JD014571.
- Velasco, E., Lamb, B., Westeberg, H., Allwine, E., Sosa, G., Arriaga-Colina, J. L., Jonson, B. T., Alexander, M. L., Prazeller, P., Knighton, W. B., Rogers, T. M., Grutter, M., Herndon, S. C., Kolb, C. E., Zavala, M., de Foy, B., Volkamer, R., Molina, L. T., and Molina, M. J.: Distribution, magnitudes, reactivities, ratios and diurnal patterns of volatile organic compounds in the Valley of Mexico during the MCMA 2002 & 2003 field campaigns, *Atmos. Chem. Phys.*, 7, 329–353, 2007.
- Wagner, P., and W. Kuttler: Biogenic and anthropogenic isoprene in the near-surface urban atmosphere—A case study in Essen, Germany, *Sci. Total Environ.*, 475, 104–115, 2014, doi:10.1016/j.scitotenv.2013.12.026.
- Warneke, C., van der Veen, C., Luxembourg, S., de Gouw, J.A., Kok, A.: Measurements of benzene and toluene in ambient air using proton-transfer-reaction mass spectrometry: Calibration humidity dependence and field intercomparison.// *Int J Mass Spectrom.* V.207, P.167–182, 2001.

- Warneke, C., de Gouw, J.A., Kuster, W.C., Goldan, P.D., Fall, R.: Validation of atmospheric VOC measurements by proton-transfer-reaction mass spectrometry using a gaschromatographicpreparation method. *Environmental Science & Technology* 37 (11), 2494–2501, 2003.
- Wayne, R. P. Ed.; *Chemistry of atmospheres*; Oxford University Press: Oxford, U.K., 2000.
- 5 White, M. L., Russo, R. S., Zhou, Y., Ambrose, J. L., Haase, K., Frinak, E. K., Varner, R. K., Wingenter, O. W., Mao, H., Talbot, R., and Sive, B. C.: Are biogenic emissions a significant source of summertime atmospheric toluene in the rural Northeastern United States?, *Atmos. Chem. Phys.*, 9, 81–92, doi:10.5194/acp9-81-2009, 2009. Parra et al., 2006.
- Wilson, R. C., Fleming, Z. L., Monks, P. S., Clain, G., Henne, S., Konovalov, I. B., Szopa, S., and Menut, L.: Have primary emission reduction measures reduced ozone across Europe? An analysis of European rural background ozone trends 1996–
- 10 2005, *Atmos. Chem. Phys.*, 12, 437–454, doi:10.5194/acp-12-437-2012, 2012.
- Xue, L. K., Wang, T., Gao, J., Ding, A. J., Zhou, X. H., Blake, D. R., Wang, X. F., Saunders, S. M., Fan, S. J., Zuo, H. C., Zhang, Q. Z., and Wang, W. X.: Ground-level ozone in four Chinese cities: precursors, regional transport and heterogeneous processes, *Atmos. Chem. Phys.*, 14, 13175–13188, doi:10.5194/acp14-13175-2014, 2014.
- Zalel, A, Yuval Yuval, Broday, D.M.: "Revealing source signatures in ambient BTEX concentrations." *Environmental*
- 15 *Pollution* 156(2): 553-562, 2008.
- Zhang, Y. H., Su, H., Zhong, L. J., Cheng, Y. F., Zeng, L. M., Wang, X. S., Xiang, Y. R., Wang, J. L., Gao, D. F., and Shao, M.: Regional ozone pollution and observation-based approach for analyzing ozone–precursor relationship during the PRIDE-PRD2004 campaign, *Atmos. Environ.*, 42, 6203–6218, doi:10.1016/j.atmosenv.2008.05.002, 2008
- Zou, Y., Deng, X. J., Zhu, D., Gong, D. C., Wang, H., Li, F., Tan, H. B., Deng, T., Mai, B. R., Liu, X. T., and Wang, B. G.:
- 20 *Characteristics of 1 year of observational data of VOCs, NO<sub>x</sub> and O<sub>3</sub> at a suburban site in Guangzhou, China*, *Atmos. Chem. Phys.*, 15, 6625-6636, doi:10.5194/acp-15-6625-2015, 2015.

**Table 1. Statistics for the filtered original 10-sec data of benzene and toluene (in ppb) from the TROICA-12 campaign (E – East segment; W – West segment). N – total number of the original 10-sec filtered samples;  $\sigma$  – standard deviation, P – percentiles.**

5

TROICA segment	N	Mean	$\sigma$	Min	Max	P10	P50	P90
<b>Benzene</b>								
<b>E</b>	39873	0.23	0.60	0.01	36.51	0.05	0.15	0.40
<b>W</b>	36322	0.18	0.27	0.01	14.32	0.05	0.13	0.35
<b>Toluene</b>								
<b>E</b>	39873	0.34	0.79	0.01	45.58	0.07	0.21	0.62
<b>W</b>	36322	0.27	0.48	0.01	25.89	0.06	0.17	0.50

**Table 2. The regional statistics of benzene, toluene (in ppb) and T/B along the Trans-Siberian railway: ER - European Russia, UR - Ural mountain region, SWS - southern parts of West Siberia, SCS - southern parts of Central Siberia, SES - southern parts of East Siberia, FE - Far East.**

Region	N	Mean	$\sigma$	Min	Max	P <sub>10</sub>	P <sub>50</sub>	P <sub>90</sub>
Benzene								
ER	7456	0.28	1.10	0.01	36.51	0.08	0.16	0.43
UR	5464	0.28	0.42	0.01	12.22	0.08	0.20	0.49
SWS	10714	0.22	0.44	0.01	22.28	0.07	0.14	0.37
SCS	13120	0.25	0.64	0.01	35.35	0.07	0.16	0.45
SES	26228	0.20	0.26	0.01	14.99	0.07	0.15	0.36
FE	13212	0.22	0.28	0.01	15.81	0.08	0.17	0.38
Toluene								
ER	7456	0.35	0.65	0.01	32.10	0.09	0.23	0.67
UR	5464	0.39	0.86	0.01	32.82	0.10	0.27	0.70
SWS	10714	0.28	0.56	0.01	17.26	0.07	0.17	0.48
SCS	13120	0.37	0.89	0.01	38.58	0.08	0.21	0.71
SES	26228	0.31	0.59	0.01	45.58	0.09	0.21	0.55
FE	13212	0.28	0.63	0.01	40.67	0.08	0.18	0.45



**Table 3. T/B ratio for moderately to strongly polluted air (NO<sub>x</sub> > 2 ppb).**

Region	N	Mean	$\sigma$	Min	Max	P <sub>10</sub>	P <sub>50</sub>	P <sub>90</sub>
ER	4247	2.20	2.63	0.04	41.71	0.49	1.53	4.26
UR	3801	2.26	2.90	0.04	49.00	0.52	1.51	4.40
SWS	6685	2.27	3.28	0.01	57.65	0.40	1.37	4.63
SCS	10732	2.28	3.08	0.02	72.13	0.48	1.50	4.44
SES	21824	2.58	3.73	0.03	187.77	0.49	1.61	5.30
FE	11645	1.84	3.01	0.01	87.86	0.33	1.10	3.62

**Table 4. The bulk contribution of motor vehicle exhausts,  $\chi$  (%), to the near-surface abundance of T and B along the Trans-Siberian Railroad (see Eq. 1).**

TROICA Segment	$\bar{T}_{low}$	$t_{low}$	$\bar{T}_{high}$	$t_{high}$	$\chi$ (%)
			Benzene		
East	0.274	0.77	0.123	0.22	89
West	0.274	0.80	0.119	0.20	90
			Toluene		
East	0.316	0.77	0.603	0.22	65
West	0.333	0.80	0.704	0.20	65

5

**Table 5. Surface levels of the studied impurities in urban and rural areas along the Trans-Siberian railway. All impurities in ppb except for NMHC and CO (ppm).**

Compound	N	Mean	$\sigma$	Min	Max	P10	P50	P90
<b>Urban</b>								
C <sub>6</sub> H <sub>6</sub>	10492	0.37	0.33	0.01	2.94	0.12	0.26	0.67
C <sub>7</sub> H <sub>8</sub>	10492	0.70	1.34	0.04	22.56	0.17	0.39	1.30
NMHC	7571	0.26	0.24	0.09	2.65	0.14	0.20	0.39
NO <sub>x</sub>	11052	11.95	16.14	0.75	205.64	2.78	7.57	23.82
CO	7239	0.27	0.08	0.14	0.68	0.20	0.25	0.36
SO <sub>2</sub>	7518	1.38	1.16	0.04	10.74	0.35	1.16	2.48
<b>Rural</b>								
C <sub>6</sub> H <sub>6</sub>	65703	0.21	0.43	0.01	36.51	0.07	0.15	0.37
C <sub>7</sub> H <sub>8</sub>	65703	0.28	0.47	0.01	45.58	0.08	0.20	0.51
NMHC	51497	0.16	0.09	0.03	3.42	0.11	0.14	0.21
NO <sub>x</sub>	79941	4.38	5.07	0.56	237.98	1.46	3.38	7.70
CO	50256	0.23	0.07	0.05	2.77	0.17	0.22	0.28
SO <sub>2</sub>	53502	1.22	0.82	0.10	9.80	0.30	1.10	2.20

Table 6. Pearson correlation matrix for urban and rural measurements along the Trans-Siberian railway.  $R \geq 0.5$  are shown in bold. Asterisks show statistically significant correlations for  $P = 0.05$ . All impurities in ppb except for NMHC and CO (ppm).

5

	$C_6H_6$	$C_5H_8$	$NO_x$	CO	$SO_2$	NMHC
<b>Urban</b>						
$C_6H_6$	1	<b>0.53*</b>	0.23*	<b>0.57*</b>	0.35*	<b>0.47*</b>
<b><math>C_7H_8</math></b>	<b>0.53*</b>	1	0.15*	<b>0.49*</b>	0.18*	0.21*
$NO_x$	0.23*	0.15*	1	<b>0.47*</b>	0.25*	0.21*
CO	<b>0.57*</b>	<b>0.49*</b>	<b>0.47*</b>	1	0.25*	0.21*
$SO_2$	0.35*	0.18*	0.25*	0.25*	1	0.01
NMHC	<b>0.47*</b>	0.21*	0.21*	0.21*	0.01	1
<b>Rural</b>						
$C_6H_6$	1	<b>0.52*</b>	0.10*	0.11*	0.08*	0.21*
<b><math>C_7H_8</math></b>	<b>0.52*</b>	1	0.17*	0.14*	0.07*	0.17*
$NO_x$	0.10*	0.17*	1	0.22*	0.16*	0.16*
CO	0.11*	0.14*	0.22*	1	0.13*	0.18*
$SO_2$	0.08*	0.07*	0.16*	0.13*	1	0.13*
NMHC	0.21*	0.17*	0.16*	0.18*	0.13*	1


**Table 7. Averaged concentrations and photochemical properties of benzene, toluene and isoprene ( $\pm$  standard deviation) from the TROICA-12 campaign.**

<b>VOC</b>	<b><math>10^{12} * K_{OH}^a</math></b>	<b>MIR<sup>b</sup></b>	<b>Concentration, ppb</b>	<b>OFP<sup>b</sup> (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>PE<sup>r</sup> (<math>\mu\text{g}/\text{m}^3</math>)</b>
Benzene	1.23	0.42	0.20 $\pm$ 0.33	0.29 $\pm$ 0.48	0.06 $\pm$ 0.09
Toluene	5.96	2.70	0.28 $\pm$ 0.51	3.10 $\pm$ 5.69	0.44 $\pm$ 0.81
Isoprene	101.00	9.10	0.60 $\pm$ 0.55	16.65 $\pm$ 15.19	11.52 $\pm$ 10.51

5

<sup>a</sup> Rate constants of VOCs with OH at 298 K ( $\text{sm}^3 \text{molecule}^{-1} \text{s}^{-1}$ ) (Atkinson, 1989; Atkinson and Arey, 2003).

<sup>b</sup> Maximum incremental reactivity (g O<sub>3</sub>/g VOC) (Carter, 1994, 1997).



5 Figure 1: Schematic representation of the TROICA-12 route from Moscow (MSW) to Vladivostok (VLK). Thin solid lines across the route represent approximate boundaries of various geographic regions: European Russia (ER), Ural mountain region (UR), southern parts of West (SWS), Central (SCS), and East (SES) Siberia, and Far East (FE). Back 2-day trajectories with endpoints at Trans-Siberian Railroad at local noon of each successive day of carriage movement are shown for East (black solid) and West (gray solid) routes of the campaign. Open circles mark air particles positions at 0, 24, and 48 hours along the each trajectory.






Figure 2: Relative frequency of NO/NO2 values in the TROICA-12 campaign.



5 Figure 3a: 10-minute concentrations of benzene, toluene, T/B ratio as well as NMHC, CO and SO<sub>2</sub> along the Trans-Siberian railway in the East segment of the TROICA-12 campaign. Some parameters are plotted in a logarithmic scale for a better data presentation. White circles on the top of the figure – the times of local noon. The cities with the population density from 250000 to 1 million and more are shown by large font. A dashed line shows T/B boundary between transport and industrial emissions (Tiwary et al., 2010; Carballo-Pat, 2014).



b)



- 5 Figure 3b: 10-minute concentrations of benzene, toluene, T/B ratio as well as NMHC, CO and SO<sub>2</sub> along the Trans-Siberian railway in the West segment of the TROICA-12 campaign. Some parameters are plotted in a logarithmic scale for a better data presentation. White circles on the top of the figure – the times of local noon. The cities with the population density from 250000 to 1 million and more are shown by large font. A dashed line shows T/B boundary between transport and industrial emissions (Tiwary et al., 2010; Carballo-Pat, 2014).






Figure 4: Diurnal profiles of the surface levels of benzene and toluene from TROICA-12 (hourly mean values): average (solid) and percentiles (dashed).



5 Figure 5: Benzene, toluene and T/B in Russian cities along the Trans-Siberian railway. The cities with the population density from 250 000 to 1 million and more are shown by large font, the cities with the population density from 50 000 to 250 000 – by small font. Asterisks show the cities with the population density less than 50 000. The cities are shown in accordance with their location along the railway. The shaded area – mean, 10-th and 90-th percentiles calculated from the data measured in rural regions. Dotted line – is a border between vehicle and industrial exhausts (Mugica et al., 2003; Tiwari et al., 2010; Carballo-Pat, 2014; Shaw et al., 2014).



5 Figure 6: Inorganic impurities in Russian cities along the Trans-Siberian railway. The cities with the population density from 250 000 to 1 million and more are shown by large font, the cities with the population density from 50 000 to 250 000 – by small font. Asterisks show the cities with the population density less than 50 000. The cities are shown in accordance with their location along the railway. The shaded area – mean, 10-th and 90-th percentiles calculated from the data measured in rural regions.




Figure 7: Frequency distribution of T/B ratio for the Russian cities along the Trans-Siberian railway.




Figure 8: Ozone formation potential (OFP) along the Trans-Siberian railway from daytime measurements (12 – 17 LST).

5