Long-term analysis of clear-sky new particle formation events and non-events in Hyytiälä

4 Lubna Dada¹, Pauli Paasonen¹, Tuomo Nieminen^{1,2}, Stephany Buenrostro Mazon¹, Jenni

5 Kontkanen¹, Otso Peräkylä¹, Katrianne Lehtipalo^{1,4}, Tareq Hussein^{1,5}, Tuukka Petäjä¹, Veli-Matti

6 Kerminen¹, Jaana Bäck³, and Markku Kulmala¹

7

14

3

8 ¹Department of Physics, University of Helsinki, P.O. Box 64, FIN-00014 Helsinki, Finland

⁹ ²Department of Applied Physics, University of Eastern Finland, P.O. Box 1627, FI-70211 Kuopio, Finland

³Department of Forest Sciences, University of Helsinki, P.O. Box 27, FIN-00014 Helsinki, Finland

⁴ Laboratory of Atmospheric Chemistry, Paul Scherrer Institute (PSI), 5232 Villigen PSI, Switzerland

⁵Department of Physics, the University of Jordan, Amman 11942, Jordan

13 Correspondence to: Lubna Dada (<u>lubna.dada@helsinki.fi</u>)

15 Abstract. New particle formation (NPF) events have been observed all around the world and are known to be a major 16 source of atmospheric aerosol particles. Here we combine 20 years of observations in a boreal forest at the SMEAR II station (Station for Measuring Ecosystem-Atmosphere Relations) in Hyytiälä, Finland, by building on previously 17 18 accumulated knowledge, and by focusing on clear-sky (non-cloudy) conditions. We first investigated the effect of 19 cloudiness on NPF and then compared the NPF event and non-event days during clear-sky conditions. In this comparison 20 we considered, for example, the effects of calculated particle formation rates, condensation sink, trace gas concentrations 21 and various meteorological quantities in discriminating NPF events from non-events. The formation rate of 1.5 nm 22 particles was calculated by using proxies for gaseous sulfuric acid and oxidized products of low volatile organic 23 compounds, together with an empirical nucleation rate coefficient. As expected, our results indicate an increase in the 24 frequency of NPF events under clear-sky conditions in comparison to cloudy ones. Also, focusing on clear-sky conditions 25 enabled us to find a clear separation of many variables related to NPF. For instance, oxidized organic vapors showed 26 higher concentration during the clear-sky NPF event days, whereas the condensation sink (CS) and some trace gases had 27 higher concentrations during the non-event days. The calculated formation rate of 3 nm particles showed a notable 28 difference between the NPF event and non-event days during clear-sky conditions, especially in winter and spring. For 29 spring time, we are able to find a threshold equation for the combined values of ambient temperature and CS, (CS (s-1) > 30 $-3.091 \times 10^{-5} \times T$ (in Kelvin) + 0.0120), above which practically no clear-sky NPF event could be observed. Finally, we 31 present a probability distribution for the frequency of NPF events at a specific CS and temperature.

32

33 Keywords: Boreal forest, formation rate, atmospheric aerosols, aerosol dynamics, condensation sink, cloudiness

34 parameter

35

- 37 1 Introduction
- 38

39 The effects of atmospheric aerosol particles on the climate system, human health and environmental interactions have 40 raised the interest in various phenomena associated with the formation, growth and loss of these particles (Pöschl, 2005; 41 Seinfeld and Pandis, 2012; Apte et al., 2015). While primary emissions are a very important source of atmospheric aerosol 42 particles, especially in terms of the aerosol mass loading, the particle number concentration is greatly affected by 43 atmospheric new particle formation (NPF). During the last couple of decades, NPF has been observed to take place almost 44 all over the world (Kulmala et al., 2004; Zhang et al., 2011; Bianchi et al., 2016; Kontkanen et al., 2016a; Kontkanen et 45 al., 2017). Atmospheric NPF is thought to be the dominant source of the total particle number concentration (Kulmala et 46 al., 2016), and a major source of cloud condensation nuclei, in the global troposphere (Merikanto et al., 2009; Yu et al., 47 2010; Kerminen et al., 2012; Salma et al., 2016).

48

49 Understanding the NPF phenomenon requires understanding its precursors and pathways involved under different 50 atmospheric conditions. For instance, high concentrations of low-volatility vapors result in a higher probability for NPF 51 (Nieminen et al., 2015), whereas a high relative humidity and condensation sink tend to suppress NPF (Hyvönen et al., 52 2005; Nieminen et al., 2014). Recent laboratory experiments have shown the importance of sulfuric acid and low-volatile 53 oxidized organic vapors to NPF (Metzger et al., 2010; Kirkby et al., 2011; Petäjä et al., 2011; Kulmala et al., 2013; Ehn 54 et al., 2014; Riccobono et al., 2014). Additionally, atmospheric observations confirm the importance of these precursor 55 vapors in the initial steps of NPF and in the further growth of newly-formed particles (Kulmala et al., 1998; Smith et al., 56 2005; Kerminen et al., 2010; Paasonen et al., 2010; Ahlm et al., 2012; Bzdek et al., 2014; Nieminen et al., 2014; Vakkari 57 et al., 2015). The Station for Measuring Forest Ecosystem-Atmosphere Relations (SMEAR II), located in Hyytiälä, 58 southern Finland, compiles almost 21 years of particle number size distribution and extensive complementary data, 59 providing the longest size distribution time series in the world, and hence allows for robust NPF analysis which is not 60 readily possible at other sites. The station is located in a homogenous Scots pine forest far from major pollution sources. 61 Hyytiälä is therefore classified as a background site representative of the semi-clean northern hemisphere boreal forests.

62

63 Many studies have investigated the role of different variables in causing, enhancing or preventing new particle formation 64 (Hyvönen et al., 2005; Baranizadeh et al., 2014; Nieminen et al., 2014). In particular, Baranizadeh et. al (2014) studied 65 the effect of cloudiness on NPF events observed at SMEAR II in Hyytiälä. They concluded, in agreement with some other 66 studies, that clouds tend to attenuate or interrupt NPF events (Sogacheva et al., 2008; Boulon et al., 2010; Baranizadeh et 67 al., 2014; Nieminen et al., 2015). In this study, we eliminated one variable that limits NPF (cloudiness), in order to 68 provide a better insight into the other quantities related to atmospheric NPF. Based on 20 years of observations and data 69 analysis for the SMEAR II station in Hyytiälä, we aim to i) quantify the effect of cloudiness on new particle formation 70 frequency, ii) characterize the differences between NPF event and non-event days during clear-sky conditions iii) explore 71 the connections between new particle formation rates calculated from precursor vapor proxies and the occurrence of NPF 72 events, iv) formulate an equation that predicts whether a clear-sky day with specific temperature and CS is classified as 73 an event; v) use the clear-sky data set to calculate the NPF probability distribution based on temperature and CS.

75 2 Materials and methods

76 2.1 Measurements

77

78 The data used for the analysis in this study is from the University of Helsinki SMEAR (Station for Measurement of 79 Ecosystem – Atmosphere Relations) II station (Hari and Kulmala, 2005). The station provides long-term continuous 80 comprehensive measurements of quantities describing atmospheric-forest-ecosystem interactions. The SMEAR II station 81 is located in the boreal forest in Hyytiälä, southern Finland (61°51N', 24°17E', 181 m a.s.l.), 220 km NW of Helsinki. 82 Tampere (200,000 inhabitants) is the largest city nearest to the station and is located 60 km SW of the site. Being far from 83 major human activities and surrounded by a homogenous scots pine belt, Hyytiälä is considered as a rural background 84 site due to the low levels of air pollutants (Asmi et al., 2011). A more detailed overview of the measurements at the station 85 can be found in Hari and Kulmala (2005) and Nieminen et al. (2014).

86

87 In this study, the data analysis is based on four types of measurements: (i) aerosol particle number size distributions, (ii) 88 concentration of the trace gases (CO, NO, NO₂, NO_x, SO₂ and O₃), (iii) meteorological parameters (solar radiation, 89 temperature and relative humidity), and (iv) precursor vapor concentrations from previously-developed proxies. The 90 collection of data started in January 1996. Trace gas concentrations are measured at 6 different heights on a 74-m-high 91 mast (extended to 126 m in summer 2010). Gas concentrations used in this study are collected from the middle level on 92 the mast above the forest (at 16.8 m).

93

94 The aerosol number concentration size distributions were measured with a twin-DMPS (Differential Mobility Particle 95 Sizer) system (Aalto et al., 2001) for the size ranges 3-500 nm until year 2004 and 3-1000 nm from 2005 onwards. These 96 data were used to classify days as NPF events and non-events following the method proposed by Dal Maso et al. (2005). 97 The size distributions obtained from the DMPS measurements were used to calculate the condensation sink, CS, which 98 is equal to the rate at which non-volatile vapors condense onto a pre-existing aerosol particle population (Kulmala et al., 99 2012).

100

101 The CO concentration is measured with one infrared light absorption analyzer (API 300EU, Teledyne Monitor Labs, 102 Englewood, CO, USA). The NO and NOx concentrations are monitored with a chemiluminescence analyzer (TEI 42C 103 TL, Thermo Fisher Scientific, Waltham, MA, USA). The NO₂ concentration is calculated from the difference NOx–NO. 104 The detection limit is about 0.05 ppb. SO₂ measurements are made through a UV-fluorescence analyzer (TEI 43 CTL, 105 Thermo Fisher Scientific, Waltham, MA, USA) that has a detection limit of 0.1 ppb. The O₃ concentration is measured 106 with an UV light absorption analyzer (TEI 49C, Thermo Fisher Scientific, Waltham, MA, USA) that has a detection limit 107 of about 1 ppb. The data for trace gases are available as 30-minute arithmetic means.

- 109 Solar radiation in the wavelengths of UV-B (280 320nm) and global radiation ($0.30 4.8 \mu$ m) are monitored using
- 110 pyranometers (SL 501A UVB, Solar Light, Philadelphia, PA, USA; Reeman TP 3, Astrodata, Tõravere, Tartumaa, Estonia
- until June 2008, and Middleton Solar SK08, Middleton Solar, Yarraville, Australia since June 2008) above the forest at
- 112 18 m. The air temperature is measured with 4-wired PT-100 sensors, and the relative humidity (in percent) is measured

with relative humidity sensors (Rotronic Hygromet MP102H with Hygroclip HC2-S3, Rotronic AG, Bassersdorf,Switzerland). These data are provided as 30-minute averages.

115

116 2.2 Data analysis

117 2.2.1 New particle formation events classification

118

119 Formation of new aerosol particles in Hyytiälä is typically observed in the time window of several hours around noon, 120 while this phenomenon seems to be rare during nighttime (Junninen et al., 2008; Buenrostro Mazon et al., 2016). 121 Accordingly, aerosol number size distributions data from the DMPS measurements at around this time window are used 122 for classifying individual days as new particle formation event or non-event days. The classification follows the guidelines 123 presented by Kulmala et al. (2012), and the procedure presented in Dal Maso et al. (2005). The latter uses a decision 124 criterion based on the presence of particles < 25 nm in diameter and their consequent growth to Aitken mode. Event days 125 are days on which sub 25 nm particle formation and growth are observed. Non-event days are days on which neither 126 modes are present. Undefined days are the days which do not fit either criterion.

127

128 2.2.2 Selecting non-cloudy days

129

Cloudiness parameter (*P*) is the ratio of measured global radiation (Rd) divided by the theoretical global irradiance (Rg):

132 $P = \frac{\mathrm{Rd}}{\mathrm{Rg}}(1)$

The theoretical maximum of global radiation (Rg) is calculated by taking into consideration the latitude of the 133 134 measurement station and the seasonal solar cycle. While a complete cloud coverage is classified as P < 0.3, a clear-sky is 135 classified as P > 0.7 (Perez et al., 1990; Sogacheva et al., 2008; Sánchez et al., 2012). In Hyytiälä, the great majority of 136 NPF events are initiated during the morning hours after the sunrise, yet before the noon (Dada et al., 2017, in preparation). 137 Since the time of the sunrise varies widely in Hyytiälä between the different seasons, the time window 9:00-12:00 seems 138 a reasonable compromise for considering whether NPF did occur or not. We found that NPF events occurring outside our 139 selected time window were very few. Accordingly, in this work the days were classified as cloudy or clear-sky days based on the median value of P during 9:00-12:00 each day, corresponding to the time window for new particle formation. 140 141 Clear-sky days were those with a median of P > 0.7 between 9:00 and 12:00 and are the focus of this study. The median 142 value ensures that at least half of our selected time window is clear-sky while the rest can vary between clear-sky and 143 minor scattered clouds. The median is useful also because NPF is a regional-scale phenomenon, so for instance scattered 144 clouds on an otherwise sunny day affecting the local radiation measurements (and leading to a momentarily drop in P) do 145 not usually interrupt the regional NPF process. Clear-sky days were those with a median of P > 0.7 between 9:00 and 146 12:00 and are the focus of this study. For consistency, the variables compared in our study are taken from the same time 147 window, 9:00-12:00.

149 2.2.3 Sulfuric acid and oxidized organics proxies

150

The gaseous sulfuric acid concentration is estimated from a pseudo-steady-state-approximation proxy developed by Petäjä
et al. (2009). This proxy takes into consideration the sulfuric acid source and sink terms as

153

$$[H_2SO_4]_{\text{proxy}} = k \cdot \frac{[SO_2] \cdot UVB}{CS} (2)$$

154 155

Here, UVB (W m⁻²) is the fraction of the UV radiation reaching earth after being screened by ozone (280 - 320 nm) and the coefficient *k* (m² W⁻¹ s ⁻¹) is obtained from the comparison of the proxy concentration to the available measured H₂SO₄ data, and has a median value of 9.9×10^{-7} m² W⁻¹ s ⁻¹.

159

160 The concentration of monoterpene oxidation products, called oxidized organic compounds (OxOrg) here, is estimated 161 using a proxy developed by Kontkanen et al. (2016b). This proxy is calculated by using the concentrations of different 162 oxidants (the measured ozone concentration $[O_3]$ and parameterizations for the hydroxyl and nitrate radical concentration, 163 [OH] and $[NO_3]$, respectively) and their reaction rates, k_i , with the monoterpenes. The MT _{proxy} (in this case MT_{proxy1,doy}) 164 is calculated by taking into account the effect of temperature-driven emissions, mixing of the boundary layer and the

165 oxidation of monoterpenes, (Kontkanen et al., 2016b).

166

$$[0x0rg]_{proxy} = \frac{(k_{OH+MT}[OH]+k_{O3+MT}[O_3]+k_{NO3+MT}[NO_3]) \cdot MT_{proxy}}{CS} (3).$$

168

169 2.2.4 Particle formation rates

170

171 The formation rate of nucleation mode particles ($J_{3,C}$, particle diameter > 3 nm) was calculated based on the method 172 suggested by Kerminen and Kulmala's equation (Kerminen and Kulmala, 2002). This quantity is a function of the 173 calculated formation rate of 1.5 nm sized particles ($J_{1.5,C}$), their growth rate (GR) and the condensation sink (CS):

174
$$J_{3,C} = J_{1.5,C} \exp\left(-\gamma \frac{CS'}{GR_{1.5-3}} \left(\frac{1}{1.5} - \frac{1}{3}\right)\right), (4)$$

where γ is a coefficient with an approximate value of 0.23 m³ nm² s⁻¹. The value of $J_{1.5,C}$ was calculated by assuming heteromolecular nucleation between SA and OxOrg as follows:

177 178

$$J_{1.5,C} = K_{het} [H_2 SO_4]_{proxy} [OxOrg]_{proxy}, (5)$$

179

The heterogeneous nucleation coefficient used in Equation 5 is the median estimated coefficient for Hyytiälä scaled from Paasonen et al. (2010): $K_{het} = 9.2 \times 10^{-14}$ cm³ s⁻¹. The scaling was made in order to fit the current data. The median value of [OxOrg] during the event days in April and May was found to be 1.6×10^7 cm⁻³(Paasonen et al., 2010), whereas the revised median value of [OxOrg] by Kontkanen et al. (2016b) is 1.3×10^8 cm⁻³. The scaling factor is the ratio between new and original [OxOrg] (0.1194). Accordingly, while the value of K_{het} from Paasonen et al. (2010) is 1.1×10^{-14} cm³ s⁻¹, after the scaling by 0.1194 we obtain the revised $K_{het} = 9.2 \times 10^{-14}$ cm³ s⁻¹.

- 186 The particle growth rate over the particle diameter range of 1.5-3 nm was calculated by taking into account the size of
- 187 the condensing vapor molecule size and the thermal speed of the particle (Nieminen et al., 2010). The growth rates (1.5 188 -3 nm) were calculated as 30-minute averages and as the sum of the growth rates due to the sulfuric acid (SA) vapor and
- 189 OxOrg vapor condensation. The density of the particle was assumed to be constant (1440 kg/m³). For SA, we first
- 190 determined the SA concentration needed to make the particles grow at the rate of 1 nm/h by taking into account the mass
- 191 of hydrated SA at the present RH and its density (Kurtén et al., 2007). Then, we calculated the GR of the particles due to
- 192 SA condensation by using the SA proxy concentration. The same method was used for GR due to OxOrg condensation,
- 193 where the vapor density was assumed to be 1200 kg/m³ (Hallquist et al., 2009; Kannosto et al., 2008). Similarly, the GR
- 194 due to OxOrg was calculated by using OxOrg proxy concentrations divided by the concentration needed for 1 nm/h GR.
- 195

2.2.5 Calculation of backward air-mass trajectories

196

197 Air mass trajectories were calculated using Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT_4) Model 198 at 96-hour backward trajectories at 100, 250 and 500 m arrival heights once per hour. Free access to transport model is developed and provided by NOAA (http://www.ready.noaa.gov/HYSPLIT.php). Input meteorological data required for 199 200 the model were collected from GDAS (Global Data Assimilation System) archives.

- 201
- 202 3 **Results and discussion**
- 203

204 3.1 Effect of cloudiness on NPF

205

206 We studied NPF events as a function of cloudiness. Figure 1a shows the fraction of event, non-event and undefined days 207 as a function of cloudiness parameter. We can see that clear-sky conditions favor the occurrence of NPF: the less clouds 208 there were, the higher was the fraction of NPF event days. For instance, for days with the cloudiness parameter of 0.3 or 209 less, the fraction of event days was less than 0.1 of the total classified days. However, the fraction of NPF event days 210 reached a maximum of around 0.55 during complete clear-sky conditions (P > 0.7), with 877 days classified as NPF events, 560 undefined days and only 229 as non-events. On the NPF event days, the median cloudiness parameter P 211 212 during the time window 9:00-12:00 was found to be 0.75 (Fig. 1b), while the non-event days were characterized by lower 213 values of P (a median of around 0.25). Also, 75% of the NPF event days were found to have a cloudiness parameter larger 214 than 0.5. The pattern found in Figure 1a follows from the fact that radiation seems essential for NPF at this site, as the 215 events occur almost solely during daylight hours (Kulmala et al., 2004b). NPF is favored under abundant radiation 216 conditions since the main components of freshly formed particles, are mainly formed photochemically (Petäjä et al., 2009; 217 Ehn et al., 2014). The fraction of undefined days, however, remained constant regardless of cloudiness conditions.

218

219 Our results emphasize the fact that radiation favors NPF to occur, while clouds tend to decrease the probability of NPF.

220 Undefined days were observed under cloudiness conditions that fell between those for NPF events and non-events. In

- 221 general, undefined days can be interrupted NPF events, or unclassified plumes of small particles due to pollution
- 222 (Buenrostro Mazon et al., 2009). The interruption of a NPF event can be due to a change in the measured air mass, or due
- 223 to attenuation of solar radiation caused by the appearance of a cloud during the event. We will not consider undefined
- 224 days further in our analyses.

- The monthly variation of daily median cloudiness parameter within the time window of 9:00-12:00 during the classified
- days is shown in Figure 2. Spring showed the best separation between the events and non-events in terms of the cloudiness
- 227 parameter, while the separation became weaker during the summer and specifically for June and July. Taken together,
- Figures 1 and 2 emphasize the observation that the presence of clouds decreases the probability of NPF events.
- 229
- 230 3.2 General character of NPF on clear-sky days
- 231

232 Upon visualizing the cloudiness conditions during events and non-events, we chose a fixed constraint for clear-sky 233 conditions (P > 0.7) during the time window of NPF (9:00-12:00) and will next focus on other parameters that distinguish 234 NPF events from non-events.

235

The monthly distribution of the event fraction on clear-sky days appeared as double peaks in spring and autumn, with spring having a higher fraction of events (Figure 3a). The minimum fraction of NPF events was recorded in December. The fraction of non-event days peaked during winter with another peak in summer. The total number of NPF events varied from year to year between 1996 and 2015. However, this variation did not show any specific trend of frequency (Figure 3b), which is in agreement with previous statistics reported from studies that did not consider clear-sky classification (Nieminen et al., 2014).

242 3.2.1 Backward air mass trajectories during clear-sky NPF events and non-events

243

244 Since NPF is most frequent in spring, we dedicated our focus into this season (Figure 3a). The springtime medians and 245 percentiles of air-mass trajectories arriving at Hyytiälä during clear-sky NPF events and non-events were calculated 96 246 hours backward in time at the 100-m, 250-m and 500-m arrival heights for the years 1996-2015. The medians and similarly 247 the percentiles were calculated by taking the median compass direction at every point on the trajectory (1 hour between 248 every two points), arriving every half an hour at Hyytiälä. The trajectories arriving at Hyytiälä at these three heights were 249 quite similar, and those arriving at the 500-m height are shown in Figure 4. Medians and percentiles of the routes were 250 calculated by taking the median of the trajectories at every half hour for spring time NPF event days and non-event days 251 separately. During the NPF event days, the measured air masses were found to originate mainly from the north and to 252 pass over Scandinavia before arriving at Hyytiälä. Similar to previously reported results, air masses arriving from the 253 north and north-west directions result in clean air with low pollutant (particulate matter and trace gas) concentrations 254 (Nieminen et al., 2015). During NPF the non-event days, air masses originated from more polluted areas in Europe and 255 Russia, resulting in elevated levels of condensation sink and other air pollutants in Hyytiälä, as also seen in previous 256 studies (Sogacheva et al., 2005).

257 3.2.2 Influences of CS, meteorological parameters and trace gases

258

259 In Figure 5a we present the monthly variation of condensation sink during NPF events and non-events under daytime 260 clear-sky conditions. NPF events tended to be favored by low values of CS throughout the year. In all the months except

- 261 during summer, the 75th percentile of the event day values of CS was lower than the 25th percentile of the non-event day
- values of CS. On the NPF event days, CS had its maximum in summer, which might be one of the main reasons for the

local minimum in the NPF event frequency during the summer months (Figure 3a). However, the monthly cycle of CS during non-event days had two maxima, one in spring and another one in autumn, which might suggest that during these seasons, high values of CS prevented NPF to occur on particular days. The difference in the value of CS between the NPF event and non-event days was the highest in March and the lowest during the summer months.

267

268 Figure 5b shows the monthly temperature conditions (T) during the daytime NPF events and non-events. While higher 269 temperatures favored NPF during the months when the average temperature was below 273.15 K (0° C) (months 1, 2, 3, 270 11 and 12), the opposite was true at average temperatures above 273.15 K (0° C). The highest recorded temperature at 271 which an event occurred during P > 0.7 sky was 300 K (25 °C) and the minimum temperature was 252 K (-21 °C). 272 Accordingly, both very high and very low temperatures were not favorable conditions for NPF to occur. Although an 273 increase in the ambient temperature results in higher concentrations of monoterpenes due to increased emissions, thereby 274 favoring new particle formation and growth (Kulmala et al., 2004), Figure 5b shows that very high temperatures tend to 275 suppress NPF. This latter feature is at least partly related to the positive relation between the ambient temperature and 276 pre-existing aerosol loading (and hence CS) in Hyytiälä (Liao et al., 2014), even though it might also be attributed to the 277 increase in vapor evaporation coefficients, which results in less stable clusters at high temperatures (Paasonen et al., 278 2012).

279

280 Consistent with an earlier study (Hamed et al., 2011), our results indicate that NPF is favored by low values of ambient 281 relative humidity in Hyytiälä (Fig. 5c). This observation does not conflict with chamber experiments (e.g.Duplissy and 282 Flagan, 2016) or theory (Merikanto et al., 2016; Vehkamäki et al., 2002), which suggest higher nucleation rates at higher 283 values of RH, because binary H₂SO₄-water nucleation is not expected take place in Hyytiälä. Other studies have proposed 284 that increased RH limits some VOC (Volatile Organic Compounds) ozonolysis reactions, preventing the formation of 285 come condensable vapors necessary for nucleation (Boy and Kulmala, 2002). This might partially explain the observed 286 anti-correlation between RH and particle formation rates. Therefore, it seems plausible that RH affects NPF via 287 atmospheric chemistry rather than via changing the sink term for condensing vapors and small clusters. Additionally, we 288 found clear differences in how trace gas concentrations were associated with RH between the NPF event and non-event 289 days (Table 1). For instance, O₃ showed a strong negative correlation with RH during events and non-events. However, 290 during non-event days, a positive correlation appears between RH and each of CO, SO₂ and NOx while the correlation 291 between those seems to be absent during event days. Our results show that air masses coming from central Europe and 292 passing over the Baltic Sea tend to have higher values of RH.

293

After looking at the characteristics of clear-sky NPF event and non-event days in terms of meteorological parameters and CS, we looked at the variation of trace gas (CO, SO₂, NOx and O₃) concentrations during these conditions (Figure 6). Out of these gases, at least SO₂ and O₃ are expected to enhance NPF, SO₂ as a precursor for sulfuric acid and O₃ as an oxidant forming ELVOCs (Extremely Low Volatile Organic Compounds) (Donahue et al., 2012; Ehn et al., 2014). However, none of these vapors seemed to have higher concentrations during NPF event days. This suggests that, as tracers of pollution, these gases are strongly linked with high anthropogenic CS, so air masses having high trace gas concentrations often do not result in NPF in Hyytiälä.

302 3.3 Connection of nucleating precursor vapors with new particle formation rate

303 3.3.1 Precursor vapor proxies

304

In this study, we determined $J_{1.5,C}$ using the proxies for both SA and OxOrg. The monthly variations of these precursors (in the time window 9:00-12:00) are shown in the Figure 7. During clear-sky conditions, the SA proxy tended to have the highest median daytime values during the winter months with a maximum in February (Figure 7a). Contrary to this, the seasonal distribution of the SA proxy reported in Hyytiälä appears as double peaks with an absolute maximum in spring and a smaller one in autumn when presenting the data without excluding cloudy days (Nieminen et al., 2014). During winter, both condensation sink and boundary layer height are lower than in the summer (Paasonen et al., 2013), which might explain the higher concentrations of SA during the winter months.

312

Being a function of temperature, the OxOrg proxy concentration was generally found to follow the monthly cycle of the ambient temperature. The median value of [OxOrg] was higher on NPF events days during every month compared with non-event days (Figure 7b). The largest difference in [OxOrg] between the NPF events and non-events, in terms of its median value, was recorded for January and the least difference for May. It is to be noted that the proxy values represent the measured values less accurately during the winter than during the other periods (Kontkanen et al., 2016b).

318

319 3.3.2 Particle formation rates

320

321 The calculated new particle formation rate, $J_{1.5,C}$, approximated with Eq. (5) shows a similar behavior as the [OxOrg] (see 322 Figures 7 and 8), being higher for the clear-sky NPF event days in comparison with non-event days. Also, the difference 323 in the value of $J_{1.5,C}$ between the NPF events and non-events was the highest in the winter, and the lowest in summer. The 324 monthly cycle of $J_{1.5,C}$ followed closely that of [OxOrg], as the latter had a higher seasonal variability than the sulfuric 325 acid proxy concentration, being thereby capable of affecting the seasonal pattern of $J_{1.5,C}$ (Figure 8a). The diurnal cycle 326 of $J_{1.5,C}$ during the NPF event days showed an increase along with sunrise, a peak at midday and decrease along with 327 sunset. However, for non-event days the $J_{1.5,C}$ value was relatively constant throughout the day and had clearly lower 328 values than during the NPF event days (Figure 8b).

329

330 Since previous studies have shown that there is a clear difference in observed J_3 between the event and non-event days, 331 and much less difference in observed $J_{1,5}$ (Kulmala et al. 2013), we decided to focus on J_3 in our event to non-event 332 discrimination. Previous studies which did not consider clear-sky conditions have reported values of observed spring time J_3 between 0.01 and 5 cm⁻³ s⁻¹ (median = 0.94 cm⁻³ s⁻¹) during the period of active NPF (Kulmala et al. 2013). Our values 333 334 of $J_{3,C}$ fit between the extremes of these values for the spring time and time window 9:00 to 12:00, with a slightly higher 335 median value of $1.9 \text{ cm}^{-3} \text{ s}^{-1}$ (Figures 9 a,b). The formation rate of 3 nm particles is affected not only by the new particle 336 formation rate $(J_{1.5})$ but also by the scavenging of newly-formed particles by coagulation into pre-existing particles. We 337 found that, in general, the values of $J_{3,C}$ calculated using equations (4) and (5) were higher on NPF event days compared 338 with non-event days in all months (Figure 9a). The difference between the event and non-event days was the largest in 339 winter and decreased towards summer. However, the diurnal cycles of percentiles and medians of $J_{3,C}$ during each month 340 peaked around noon for both NPF events and non-events. One example is presented in Figure 9b, showing that $J_{3,C}$ tended

- to increase after the sunrise, to peak at about midday and to diminish after sunset. This kind of diurnal cycle was similar for all the months. Hourly values of $J_{3,C}$ calculated during the NPF event days were higher than those during the nonevent days. During the spring months, the difference in the median $J_{3,C}$ between the NPF events and non-events, calculated for every half an hour, appeared to increase at about 10:00 and then it started to decrease again at about 13:00 (Figure 9b). On NPF event days, in comparison to springtime $J_{1.5,C}$ which peaked at around 10:45 (Figure 8b), $J_{3,C}$ peaked typically about half-an hour later. This time delay indicates how long it takes for the particles grow from 1.5 nm to 3 nm. This growth is a critical step of NPF (Kulmala et al. 2013), and depends on concentrations of available vapor precursors.
- 348

349 In Figure 10 we present the median diurnal cycles of $J_{3,C}$ plotted against the median diurnal cycles of CS during classified 350 clear-sky NPF events and non-events. The diurnal cycle was calculated by taking the median CS at every half hour 351 throughout the season. On the NPF event days, the CS had higher values during the nighttime and lower values during 352 daytime with a minimum at noon. It is important to remember that J was calculated only for daytime when the SA proxy 353 was available (UV-B radiation is needed for the proxy). On non-event days, the values of CS showed no clear diurnal 354 pattern, had practically no difference between the daytime and nighttime hours, and were roughly twice those recorded 355 during the clear-sky NPF event days. The difference in CS between NPF events and non-events follows from the distinctly 356 different air masses arriving at Hyytiälä. For instance, it has been shown that air-masses originating from the north and 357 passing over Scandinavia have, on average, lower values of CS than the air masses passing over Russia and central Europe 358 (Sogacheva et al., 2005; Nieminen et al., 2015).

359

360 On NPF event days, the median approximated formation rate of 3 nm particles had its maximum value at about midday 361 and was significantly higher than on non-events days (Figures 9b and 10). A clear negative relation could be seen between 362 the median seasonal diurnal cycles of CS and $J_{3,C}$ on NPF event days (specifically during spring daytime) (Figure 10). 363 This kind of relation was not observed during non-event days when these two quantities seemed to be independent of 364 each other (Figure 10). In summer, the median value of $J_{3,C}$ was roughly similar between NPF events and non-events, 365 whereas the median value of CS was almost ten times higher during the non-event days compared with event days. The 366 high values of $J_{3,C}$ for the non-event days in summer, despite the high CS values, seem to suggest that some other factor 367 limits the actual NPF rate. One possibility is that freshly-formed clusters are rapidly evaporated due to higher ambient 368 temperatures (see Fig. 5b). This will be discussed in a more detail in the following section. Higher values of CS on non-369 event days are expected, bearing in mind that these particles act as surfaces for scavenging precursor gases and freshly 370 formed particles (Hussein et al., 2008). The association of a high CS with the lower NPF probability has been observed 371 in many studies conducted in Hyytiälä (Boy and Kulmala, 2002; Hyvönen et al., 2005; Baranizadeh et al., 2014), as well 372 as in other rural and urban areas, including Egbert and Toronto in Canada (Jun et al., 2014), Preila in Lithuania (Mordas 373 et al., 2016), Po Valley in Italy (Hamed et al., 2007) and Budapest and K-puszta in Hungary (Salma et al., 2016).

374

375 3.3.3 Threshold separating the NPF events and non-events

376

Since quite a visible separation could be observed in the calculated values of $J_{3,C}$ between the spring-time clear-sky NPF events and non-events, and since $J_{3,C}$ had its maximum at around midday, the plot of CS versus temperature at midday (11:00-12:00) in spring provides an equation that effectively separates the NPF events from non-events during this season (Figure 11). This equation was determined using a linear discriminant analysis (LDA) similar to Hyvönen et al. (2005). The equation provides a line that separates NPF events from non-events at 95% confidence towards non-events. Basedon their midday CS and Temperature, the data point follows either classes. More specifically, the days with

383

384 CS (s⁻¹) > -3.091×10⁻⁵ × T (in Kelvin) + 0.0120, (6)

385

lie above the threshold line. Almost no non-event days fall below this line (< 5%). The points above the line were also
characterized with higher trace gases concentrations and lower calculated formation rates of 3 nm particles than the rest
of the points.

389

The separation between the clear-sky NPF events and non-events in the CS versus *T* plot was less evident in autumn and disappeared completely in the summer and winter (Figure 12). Interestingly, a large number of NPF event days during these seasons still fell below the threshold line given by Equation 6. Furthermore, we analyzed the effect of RH in separating the events from nonevents, similar to the study done on RH by Hyvönen et al. (2005). We found that compared with CS vs temperature data, depicting CS vs RH (data not presented) did not work better in separating NPF events from non-events during clear-sky conditions.

- 396 3.3.4 Probability of NPF events and non-events
- 397

398 Since the biggest difference in the calculated 3 nm particle formation rates between the NPF events and non-events was 399 observed around noon (Figure 9b), and since CS and temperature showed promising threshold values for predicting the 400 occurrence of NPF non-events during spring (up to 95%) (Figure 11), Figure 13 presents the probability of having a NPF 401 event in Hyytiälä at a specific CS and temperature within the time window 11:00-12:00. The probability was calculated 402 taking the fraction of events to the total events and non-events in every cell which is 2.5 K on the x-axis and a ratio of 403 1.14 on the y-axis between every two consecutive CS values. The highest probability of having a NPF event corresponded 404 to conditions having moderate temperatures and low values of CS. At high values of CS, there was a zero probability for 405 NPF regardless of the temperature. However, at moderate and low values of CS, the probability of having a NPF event 406 decreases as we go to lower temperature. This could be explained by lower emissions of VOCs, and thus lower OxOrg 407 concentrations, at lower temperatures. Similarly, the probability of NPF decreases as we go to higher temperatures at 408 constant values of CS. This latter feature might be attributed to conditions unfavorable for clustering due to high 409 temperatures. Although previous studies have developed criteria for NPF probability which could work in diverse 410 environments(Kuang et al., 2010), they did not explore the dependency of their parameter on atmospheric conditions.

411

412 4 Conclusion

413

In this study we combined 20 years of data collected at the SMEAR II station in order to characterize the conditions affecting the frequency of NPF events in that location. By focusing only on clear-sky conditions, we were able to get new insight into differences between the NPF events and non-events. In clear-sky conditions, the meteorological conditions, trace gas concentrations and other studied variables on NPF event days appeared to be similar to those presented in the previous studies which did not consider clear-sky classification. Furthermore, the monthly data refined the analysis so that the differences caused by different quantities became more visible compared the previous studies conducted for this site. Our work confirms, with a complementary dataset, the conclusions of Baranizadeh et al. (2014) that NPF events and
 non-events are typically associated with clear-sky and cloudy conditions, respectively.

422

423 Our results showed that using SA and OxOrg proxies to calculate the apparent formation rates of 1.5 and 3 nm particles 424 works well in differentiating the clear-sky NPF events from non-events. Moreover, during clear-sky conditions the effect 425 of CS on attenuating or even preventing NPF was quite visible: CS was, on average, two times higher on the non-event 426 days compared with the NPF event days. Similarly, many other meteorological variables affected NPF. By using CS and 427 ambient temperature, we were able to find a threshold above which no clear-sky NPF events occurred. This threshold is 428 described with an equation that is able to separate 97.4% of the NPF events from non-events during spring time. In clear 429 sky conditions, when there is plenty of radiation available, NPF events take place as long as the CS is low enough and 430 temperature is moderate. Although a weaker separation was observed in the other seasons, considering only clear-sky 431 conditions enabled us to form a map of the probability of having a NPF event within specific CS and temperature 432 conditions. Using clear-sky conditions appears to bring us one step forward towards understanding NPF and predicting 433 their occurrences in Hyytiälä. Our study serves as a basis to future detailed comparisons with observations to formulate 434 even more robust conclusions.

435

436 5 Acknowledgements

437

This work was supported by the Academy of Finland Centre of Excellence program (grant no. 272041) and Nordic Toplevel Research Initiative (TRI) Cryosphere-Atmosphere Interactions in a Changing Arctic Climate (CRAICC). The authors thank the division of atmospheric sciences at the University of Helsinki. We also thank Mrs. Ksenia Tabakova for providing air-mass trajectory data.

448

464

470

483

487

443 6 References

- 445 Aalto, P., Hämeri, K., Becker, E., Weber, R., Salm, J., Mäkelä, J. M., Hoell, C., O'dowd, C. D., Hansson, H.-C., Väkevä, 446 M., Koponen, I. K., Buzorius, G., and Kulmala, M.: Physical characterization of aerosol particles during 447 nucleation events, Tellus B, 53, 344-358 10.1034/j.1600-0889.2001.530403.x, 2001.
- 449 Ahlm, L., Liu, S., Day, D. A., Russell, L. M., Weber, R., Gentner, D. R., Goldstein, A. H., DiGangi, J. P., Henry, S. B., 450 Keutsch, F. N., VandenBoer, T. C., Markovic, M. Z., Murphy, J. G., Ren, X., and Scott, S.: Formation and 451 growth of ultrafine particles from secondary sources in Bakersfield, California, J. Geophys. Res. Atmos, 117, 452 10.1029/2011JD017144, 2012. 453
- 454 Apte, J. S., Marshall, J. D., Cohen, A. J., and Brauer, M.: Addressing global mortality from ambient PM2. 5, 455 Environmental science & technology, 49, 8057-8066, 10.1021/acs.est.5b01236, 2015. 456
- 457 Asmi, A., Wiedensohler, A., Laj, P., Fjaeraa, A.-M., Sellegri, K., Birmili, W., Weingartner, E., Baltensperger, U., Zdimal, 458 V., and Zikova, N.: Number size distributions and seasonality of submicron particles in Europe 2008–2009, 459 Atmos. Chem. Phys., 11, 5505-5538, 10.5194/acp-11-5505-2011, 2011. 460
- 461 Baranizadeh, E., Arola, A., Hamed, A., Nieminen, T., Mikkonen, S., Virtanen, A., Kulmala, M., Lehtinen, K., and 462 Laaksonen, A.: The effect of cloudiness on new-particle formation: investigation of radiation levels, Boreal Env. 463 Res., 19, 343-354, 2014.
- 465 Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki, 466 N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H. 467 E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., 468 Kulmala, M., Dommen, J., and Baltensperger, U.: New particle formation in the free troposphere: A question of 469 chemistry and timing, Science, 352, 1109-1112, 10.1126/science.aad5456, 2016.
- 471 Boulon, J., Sellegri, K., Venzac, H., Picard, D., Weingartner, E., Wehrle, G., Collaud Coen, M., Bütikofer, R., Flückiger, 472 E., and Baltensperger, U.: New particle formation and ultrafine charged aerosol climatology at a high altitude 473 site in the Alps (Jungfraujoch, 3580 m asl, Switzerland), Atmos. Chem. Phys., 10, 9333-9349, 10.5194/acp-10-474 9333-2010, 2010. 475
- 476 Boy, M., and Kulmala, M.: Nucleation events in the continental boundary layer: Influence of physical and meteorological 477 parameters, Atmos. Chem. Phys., 2, 1-16, 10.5194/acp-2-1-2002, 2002. 478
- 479 Buenrostro Mazon, S., Riipinen, I., Schultz, D., Valtanen, M., Maso, M. D., Sogacheva, L., Junninen, H., Nieminen, T., 480 Kerminen, V.-M., and Kulmala, M.: Classifying previously undefined days from eleven years of aerosol-particle-481 size distribution data from the SMEAR II station, Hyytiälä, Finland, Atmos. Chem. Phys., 9, 667-676, 482 10.5194/acp-9-667-2009, 2009.
- 484 Buenrostro Mazon, S., Kontkanen, J., Manninen, H. E., Nieminen, T., Kerminen, V.-M., and Kulmala, M.: A long-term 485 comparison of nighttime cluster events and daytime ion formation in a boreal forest, Boreal Env. Res., 21, 242-486 261, 2016.
- 488 Bzdek, B. R., Lawler, M. J., Horan, A. J., Pennington, M. R., DePalma, J. W., Zhao, J., Smith, J. N., and Johnston, M. 489 V.: Molecular constraints on particle growth during new particle formation, Geophys. Res. Lett., 41, 6045-6054, 490 10.1021/ac100856j, 2014. 491
- 492 Dada, L., Chellapermal, R., Buenrostro Mazon, S., Junninen, H., Kerminen, V. M., Paasonen, P., and Kulmala, M.: 493 Method for identifying NPF event start and end times as well as NPF types (ion-initiated, particle initiated, 494 transported..) using characteristic nucleation-mode particles and air ions, 2017, in preparation. 495
- 496 Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P., and Lehtinen, K. E.: Formation and growth 497 of fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II, Hyytiala, Finland, 498 Boreal Env. Res., 10, 323, 2005. 499
- 500 Donahue, N. M., Kroll, J., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set-Part 2: Diagnostics 501 of organic-aerosol evolution, Atmos. Chem. Phys., 12, 615-634, 10.5194/acp-12-615-2012, 2012.

Duplissy, J., and Flagan, R.: Effect of ions on sulfuric acid-water binary particle formation: 2. Experimental data and comparison, J. Geophys. Res. Atmos, 121, 1752-1775, 10.1002/2015JD023539, 2016.

502

512

516

520

- Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee,
 B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J.,
 Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M.,
 Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and
 Mentel, T. F.: A large source of low-volatility secondary organic aerosol, Nature, 506, 476-479,
 10.1038/nature13032, 2014.
- Hallquist, M., Wenger, J., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N., George,
 C., and Goldstein, A.: The formation, properties and impact of secondary organic aerosol: current and emerging
 issues, Atmos. Chem. Phys., 9, 5155-5236, 10.5194/acp-9-5155-2009, 2009.
- Hamed, A., Joutsensaari, J., Mikkonen, S., Sogacheva, L., Maso, M. D., Kulmala, M., Cavalli, F., Fuzzi, S., Facchini, M.,
 and Decesari, S.: Nucleation and growth of new particles in Po Valley, Italy, Atmos. Chem. Phys., 7, 355-376,
 10.5194/acp-7-355-2007, 2007.
- 521 Hari, P., and Kulmala, M.: Station for measuring ecosystem-atmosphere relations, Boreal Env. Res., 10, 315-322, 2005.
- Hussein, T., Martikainen, J., Junninen, H., Sogacheva, L., Wagner, R., Dal Maso, M., Riipinen, I., Aalto, P. P., and Kulmala, M.: Observation of regional new particle formation in the urban atmosphere, Tellus B, 60, 509-521, 10.1111/j.1600-0889.2008.00365.x, 2008.
- Hyvönen, S., Junninen, H., Laakso, L., Maso, M. D., Grönholm, T., Bonn, B., Keronen, P., Aalto, P., Hiltunen, V., Pohja,
 T., Launiainen, S., Hari, P., Mannila, H., and Kulmala, M.: A look at aerosol formation using data mining
 techniques, Atmos. Chem. Phys., 5, 3345-3356, 10.5194/acp-5-3345-2005, 2005.
- Jun, Y.-S., Jeong, C.-H., Sabaliauskas, K., Leaitch, W. R., and Evans, G. J.: A year-long comparison of particle formation
 events at paired urban and rural locations, Atmospheric Pollution Research, 5, 447-454, 10.5094/APR.2014.052,
 2014.
- Junninen, H., Hulkkonen, M., Riipinen, I., Nieminen, T., Hirsikko, A., Suni, T., Boy, M., LEE, S. H., Vana, M., Tammet,
 H., KERMINEN, V.-M., and KULMALA, M.: Observations on nocturnal growth of atmospheric clusters, Tellus
 B, 60, 365-371, 10.1111/j.1600-0889.2008.00356.x, 2008.
- Kannosto, J., Virtanen, A., Lemmetty, M., Mäkelä, J. M., Keskinen, J., Junninen, H., Hussein, T., Aalto, P., and Kulmala,
 M.: Mode resolved density of atmospheric aerosol particles, Atmos. Chem. Phys., 8, 5327-5337, 10.5194/acp8-5327-2008, 2008.
- Kerminen, V.-M., and Kulmala, M.: Analytical formulae connecting the "real" and the "apparent" nucleation rate and the nuclei number concentration for atmospheric nucleation events, Journal of Aerosol Science, 33, 609-622, 10.1016/S0021-8502(01)00194-X, 2002.
- 547 Kerminen, V.-M., Petäjä, T., Manninen, H., Paasonen, P., Nieminen, T., Sipilä, M., Junninen, H., Ehn, M., Gagné, S.,
 548 Laakso, L., Riipinen, I., Vehkamäki, H., Kurten, T., Ortega, I. K., Maso, M. D., Brus, D., Hyvärinen, A.,
 549 Lihavainen, H., Leppä, J., Lehtinen, K. E. J., Mirme, A., Mirme, S., Hõrrak, U., Berndt, T., Stratmann, F.,
 550 Birmili, W., Wiedensohler, A., Metzger, A., Dommen, J., Baltensperger, U., Kiendler-Scharr, A., Mentel, T. F.,
 551 Wildt, J., Winkler, P. M., Wagner, P. E., Petzold, A., Minikin, A., Plass-Dülmer, C., Pöschl, U., Laaksonen, A.,
 553 Chem. Phys., 10, 10829-10848, 10.5194/acp-10-10829-2010, 2010.
- Kerminen, V.-M., Paramonov, M., Anttila, T., Riipinen, I., Fountoukis, C., Korhonen, H., Asmi, E., Laakso, L.,
 Lihavainen, H., Swietlicki, E., Svenningsson, B., Asmi, A., Pandis, S. N., Kulmala, M., and Petäjä, T.: Cloud
 condensation nuclei production associated with atmospheric nucleation: a synthesis based on existing literature
 and new results, Atmos. Chem. Phys., 12, 12037-12059, 10.5194/acp-12-12037-2012, 2012.
- Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A.,
 Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Georgios Tsagkogeorgas, Daniela

562 563 564 565 566 567 568 569 570	Wimmer, Antonio Amorim, Bianchi, F., Martin Breitenlechner, André David, Josef Dommen, Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Sandra Mogo, Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, Nature, 476, 429- 433, 10.1038/nature10343, 2011.
571 572 573 574 575	Kontkanen, J., Järvinen, E., Manninen, H. E., Lehtipalo, K., Kangasluoma, J., Decesari, S., Gobbi, G. P., Laaksonen, A., Petäjä, T., and Kulmala, M.: High concentrations of sub-3nm clusters and frequent new particle formation observed in the Po Valley, Italy, during the PEGASOS 2012 campaign, Atmos. Chem. Phys., 16, 1919-1935, 10.5194/acp-16-1919-2016, 2016a.
576 577 578 579	Kontkanen, J., Paasonen, P., Aalto, J., Bäck, J., Rantala, P., Petäjä, T., and Kulmala, M.: Simple proxies for estimating the concentrations of monoterpenes and their oxidation products at a boreal forest site, Atmos. Chem. Phys., 16, 13291-13307, 10.5194/acp-16-13291-2016, 2016b.
580 581 582 583 584	 Kontkanen, J., Lehtipalo, K., Ahonen, L., Kangasluoma, J., Manninen, H. E., Hakala, J., Rose, C., Sellegri, K., Xiao, S., Wang, L., Qi, X., Nie, W., Ding, A., Yu, H., Lee, S., Kerminen, V. M., Petäjä, T., and Kulmala, M.: Measurements of sub-3 nm particles using a particle size magnifier in different environments: from clean mountain top to polluted megacities, Atmos. Chem. Phys., 17, 2163-2187, 10.5194/acp-17-2163-2017, 2017.
585 586 587 588	Kuang, C., Riipinen, I., Sihto, SL., Kulmala, M., McCormick, A., and McMurry, P.: An improved criterion for new particle formation in diverse atmospheric environments, Atmos. Chem. Phys., 10, 8469-8480, 10.5194/acp-10- 8469-2010, 2010.
589 590 591	Kulmala, M., Toivonen, A., Mäkelä, J. M., and Laaksonen, A.: Analysis of the growth of nucleation mode particles observed in Boreal forest, Tellus B, 50, 449-462, 10.3402/tellusb.v50i5.16229, 1998.
592 593 594 595	Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, VM., Birmili, W., and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, Journal of Aerosol Science, 35, 143-176, 10.1016/j.jaerosci.2003.10.003, 2004.
596 597 598 599 600	 Kulmala, M., Suni, T., Lehtinen, K. E. J., Dal Maso, M., Boy, M., Reissell, A., Rannik, Ü., Aalto, P., Keronen, P., Hakola, H., Bäck, J., Hoffmann, T., Vesala, T., and Hari, P.: A new feedback mechanism linking forests, aerosols, and climate, Atmos. Chem. Phys. , 4, 557-562, 10.5194/acp-4-557-2004, 2004b.
601 602 603 604	Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H., and Paasonen, P.: Measurement of the nucleation of atmospheric aerosol particles, Nature protocols, 7, 1651- 1667, 10.1038/nprot.2012.091, 2012.
605 606 607 608 609 610 611	 Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. r., Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M., Smith, J., Ehn, M., Mentel, T., Lehtinen, K., Laaksonen, A., Kerminen, V., and Worsnop, D.: Direct observations of atmospheric aerosol nucleation, Science, 339, 943-946, 10.1126/science.1227385, 2013.
612 613 614	Kulmala, M., Luoma, K., Virkkula, A., Petäjä, T., Paasonen, P., Kerminen, VM., Nie, W., Qi, X., Shen, Y., and Chi, X.: On the mode-segregated aerosol particle number concentration load, Boreal Env. Res., 2016.
615 616 617	Kurtén, T., Torpo, L., Ding, C. G., Vehkamäki, H., Sundberg, M. R., Laasonen, K., and Kulmala, M.: A density functional study on water-sulfuric acid-ammonia clusters and implications for atmospheric cluster formation, J. Geophys. Res. Atmos, 112, 10.1029/2006JD007391, 2007.
619 620 621	Liao, L., Kerminen, VM., Boy, M., Kulmala, M., and Dal Maso, M.: Temperature influence on the natural aerosol budget over boreal forests, Atmos. Chem. Phys., 14, 8295-8308, 10.5194/acp-14-8295-2014, 2014.

Merikanto, J., Spracklen, D., Mann, G., Pickering, S., and Carslaw, K.: Impact of nucleation on global CCN, Atmos.
 Chem. Phys., 9, 8601-8616, 10.5194/acp-9-8601-2009, 2009.

624

629

638

642

661

- Merikanto, J., Duplissy, J., Määttänen, A., Henschel, H., Donahue, N. M., Brus, D., Schobesberger, S., Kulmala, M., and
 Vehkamäki, H.: Effect of ions on sulfuric acid-water binary particle formation: 1. Theory for kinetic-and
 nucleation-type particle formation and atmospheric implications, J. Geophys. Res. Atmos, 121, 1736-1751,
 10.1002/2015JD023538, 2016.
- Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S., Weingartner, E., Riipinen, I., Kulmala, M.,
 Spracklen, D. V., and Carslaw, K. S.: Evidence for the role of organics in aerosol particle formation under
 atmospheric conditions, Proceedings of the National Academy of Sciences, 107, 6646-6651,
 10.1073/pnas.0911330107, 2010.
- Mordas, G., Plauškaitė, K., Prokopciuk, N., Dudoitis, V., Bozzetti, C., and Ulevicius, V.: Observation of new particle
 formation on Curonian Spit located between continental Europe and Scandinavia, Journal of Aerosol Science,
 10.1016/j.jaerosci.2016.03.002, 2016.
- Nieminen, T., Lehtinen, K., and Kulmala, M.: Sub-10 nm particle growth by vapor condensation–effects of vapor molecule size and particle thermal speed, Atmos. Chem. Phys., 10, 9773-9779, 10.5194/acp-10-9773-2010, 2010, 2010.
- Nieminen, T., Asmi, A., Dal Maso, M., Aalto, P. P., Keronen, P., Petäjä, T., Kulmala, M., and Kerminen, V.-M.: Trends
 in atmospheric new-particle formation: 16 years of observations in a boreal-forest environment, Boreal Env.
 Res., 19, 2014.
- Nieminen, T., Yli-Juuti, T., Manninen, H., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Technical note: New particle
 formation event forecasts during PEGASOS–Zeppelin Northern mission 2013 in Hyytiälä, Finland, Atmos.
 Chem. Phys., 15, 12385-12396, 10.5194/acp-15-12385-2015, 2015.
- Paasonen, P., Nieminen, T., Asmi, E., Manninen, H., Petäjä, T., Plass-Dülmer, C., Flentje, H., Birmili, W., Wiedensohler,
 A., Horrak, U., Metzger, A., Hamed, A., Laaksonen, A., Facchini, M. C., Kerminen, V.-M., and Kulmala, M.:
 On the roles of sulphuric acid and low-volatility organic vapours in the initial steps of atmospheric new particle
 formation, Atmos. Chem. Phys., 10, 11223-11242, 10.5194/acp-10-11223-2010, 2010.
- Paasonen, P., Olenius, T., Kupiainen, O., Kurtén, T., Petäjä, T., Birmili, W., Hamed, A., Hu, M., Huey, L., Plass-Duelmer,
 C., Smith, J. N., Wiedensohler, A., Loukonen, V., McGrath, M. J., Ortega, I. K., Laaksonen, A., Vehkamäki, H.,
 Kerminen, V.-M., and Kulmala, M.: On the formation of sulphuric acid–amine clusters in varying atmospheric
 conditions and its influence on atmospheric new particle formation, Atmos. Chem. Phys., 12, 9113-9133,
 10.5194/acp-12-9113-2012, 2012.
- Paasonen, P., Asmi, A., Petäjä, T., Kajos, M. K., Äijälä, M., Junninen, H., Holst, T., Abbatt, J. P., Arneth, A., Birmili,
 W., Gon, H. D. v. d., Hamed, A., Hoffer, A., Laakso, L., Laaksonen, A., Leaitch, W. R., Plass-Dülmer, C., Pryor,
 S. C., Räisänen, P., Swietlicki, E., Wiedensohler, A., Worsnop, D. R., Kerminen, V.-M., and Kulmala, M.:
 Warming-induced increase in aerosol number concentration likely to moderate climate change, Nature
 Geoscience, 6, 438-442, 10.1038/ngeo1800, 2013.
- Perez, R., Ineichen, P., Seals, R., and Zelenka, A.: Making full use of the clearness index for parameterizing hourly
 insolation conditions, Solar Energy, 45, 111-114, 10.1016/0038-092X(90)90036-C, 1990.
- Petäjä, T., Mauldin Iii, R., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, Atmos. Chem. Phys., 9, 7435-7448, 10.5194/acp-9-7435-2009, 2009.
- Petäjä, T., Sipilä, M., Paasonen, P., Nieminen, T., Kurtén, T., Ortega, I. K., Stratmann, F., Vehkamäki, H., Berndt, T.,
 and Kulmala, M.: Experimental observation of strongly bound dimers of sulfuric acid: Application to nucleation
 in the atmosphere, Physical review letters, 106, 228302, 10.1103/PhysRevLett.106.228302, 2011.
- Pöschl, U.: Atmospheric aerosols: composition, transformation, climate and health effects, Angewandte Chemie
 International Edition, 44, 7520-7540, 10.1002/anie.200501122, 2005.
 - 16

- 682 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, 683 F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, 684 A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kürten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petäjä, T., Praplan, A. P., Santos, F. D., 685 Schallhart, S., Seinfeld, J. H., Sipilä, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tomé, A., 686 Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, 687 688 D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R., and Baltensperger, 689 U.: Oxidation products of biogenic emissions contribute to nucleation of atmospheric particles, Science, 344, 690 717-721, 10.1126/science.1243527, 2014.
- Salma, I., Németh, Z., Kerminen, V.-M., Aalto, P., Nieminen, T., Weidinger, T., Molnár, Á., Imre, K., and Kulmala, M.:
 Regional effect on urban atmospheric nucleation, Atmos. Chem. Phys., 16, 8715-8728, 10.5194/acp-16-87152016, 2016.
- Sánchez, G., Serrano, A., and Cancillo, M.: Effect of cloudiness on solar global, solar diffuse and terrestrial downward
 radiation at Badajoz (Southwestern Spain), Optica pura y aplicada, 45, 33-38, 2012.
- Seinfeld, J. H., and Pandis, S. N.: Atmospheric chemistry and physics: from air pollution to climate change, John Wiley & Sons, 2012.
 701
- Smith, J. N., Moore, K. F., Eisele, F. L., Voisin, D., Ghimire, A. K., Sakurai, H., and McMurry, P. H.: Chemical composition of atmospheric nanoparticles during nucleation events in Atlanta, J. Geophys. Res. Atmos, 110, 10.1029/2005JD005912, 2005.
- Sogacheva, L., Dal Maso, M., Kerminen, V.-M., and Kulmala, M.: Probability of nucleation events and aerosol particle
 concentration in different air mass types arriving at Hyytiälä, southern Finland, based on back trajectories
 analysis, Boreal Env. Res., 10, 2005.
- Sogacheva, L., Saukkonen, L., Nilsson, E., Dal Maso, M., Schultz, D. M., De Leeuw, G., and Kulmala, M.: New aerosol particle formation in different synoptic situations at Hyytiälä, southern Finland, Tellus B, 60, 485-494, 10.1111/j.1600-0889.2008.00364.x, 2008.
- Vakkari, V., Tiitta, P., Jaars, K., Croteau, P., Beukes, J. P., Josipovic, M., Kerminen, V. M., Kulmala, M., Venter, A. D.,
 and Zyl, P. G.: Reevaluating the contribution of sulfuric acid and the origin of organic compounds in atmospheric
 nanoparticle growth, Geophys. Res. Lett., 42, 10.1002/2015GL066459, 2015.
- Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E., Timmreck, C., Noppel, M., and Laaksonen, A.: An improved parameterization for sulfuric acid–water nucleation rates for tropospheric and stratospheric conditions, J. Geophys. Res. Atmos, 107, 10.1029/2002JD002184, 2002.
 721
- Yu, F., Luo, G., Bates, T. S., Anderson, B., Clarke, A., Kapustin, V., Yantosca, R. M., Wang, Y., and Wu, S.: Spatial distributions of particle number concentrations in the global troposphere: Simulations, observations, and implications for nucleation mechanisms, J. Geophys. Res. Atmos, 115, 10.1029/2009JD013473, 2010.
- Zhang, R., Khalizov, A., Wang, L., Hu, M., and Xu, W.: Nucleation and growth of nanoparticles in the atmosphere, Chem.
 Rev., 112, 1957-2011, 10.1021/cr2001756, 2011.
- 728 729

698

705

- Table 1 Correlation coefficients between different meteorological parameters, gas concentrations and condensation sink (CS) during clear-sky events and non-events during spring (Mar-May, 1996-2015) and time window 9:00 12:00. The light blue refers to medium positive correlation (>0.45) and the dark blue to high positive correlation (>0.7), light orange refers to medium
- 731 732 733

negative correlation (<-0.45) and dark red to high negative ones (<-0.7).

	CS	Τ	RH	CO	NOx	SO ₂	O 3	
	Events							
CS	1							
Т	0.28	1						
RH	-0.06	-0.64	1					
СО	0.33	-0.37	0.26	1				
NOx	0.53	-0.19	0.21	0.47	1			
SO ₂	0.4	-0.29	0.14	0.36	0.58	1		
O ₃	0.23	0.52	-0.51	-0.06	-0.08	-0.08	1	
	Non-Events							
CS	1							
Т	0.15	1						
RH	-0.12	-0.81	1					
СО	0.53	-0.68	0.5	1				
NOx	0.34	-0.51	0.45	0.7	1			
SO ₂	0.23	-0.55	0.42	0.56	0.41	1		
O 3	0.43	0.62	-0.64	-4E-04	-0.07	-0.13	1	



Figure 1: (a) Figure showing the fraction of days which are classified as NPF events, non-events, and undefined days during different sky cloudiness conditions. (b) Cloudiness daily (9:00 - 12:00) medians and percentiles recorded during NPF event, undefined and non-event days. The red line represents the median of the data and the lower and upper edges of the box represent 25th and 75th percentiles of the data respectively. The length of the whiskers represent 1.5 x interquartile range which includes 99.3% of the data. Data outside the whiskers are considered outliers and are marked with red crosses.



Figure 2: Monthly variation of cloudiness daily (9:00 – 12:00) medians and percentiles recorded during NPF events (E; white)
 and non-events (N; shaded). Numbers below the plot correspond to the number of data points included in each boxplot. Number
 of clear-sky events (E (P>0.7)) and clear-sky non-events (N (P>0.7)) accompany the plot. See Figure 1 for explanation of
 symbols.



Figure 3: (a) Monthly and (b) yearly fraction of clear-sky days classified as NPF Events, undefined and non-events. In year 1998, global radiation data is limited to 5.4%, making the classification bias.





Figure 4: Median and percentiles of 96 hours backward air-mass trajectories arriving to Hyytiälä during spring time (9:00-12:00).





Figure 5: Median and percentiles of monthly variation (9:00 – 12:00) at P>0.7 of (a) CS (b) Temperature and (c) RH during NPF events (E, white) and non-events (N, shaded). See Figure 1 for explanation of symbols.



Figure 6: Spring time (months 3,4,5) medians and percentiles of trace gases during clear-sky events (E, white) and non-events (n, shaded) during daytime (9:00 – 12:00). See Figure 1 for explanation of symbols.





Figure 7: Monthly variation of medians and percentiles of (a) SA proxy and (b) OxOrg proxy at P>0.7 during the time window 9:00 – 12:00 of NPF events (E, white) and non-events (N, shaded). See Figure 1 for explanation of symbols.



Figure 8: (a) Monthly variation of medians and percentiles of $J_{1.5,C}$ during the time window 9:00 – 12:00 of NPF events (E, white) and non-events (N, shaded). See Figure 1 for explanation of symbols. (b) The diurnal cycle of $J_{1.5,C}$ during Spring. The nighttime is missing in this plot due to unavailable SA proxy which uses UVB to be calculated.



b

Figure 9: (a) $J_{3,C}$ medians and percentiles during different months separated classified NPF events (E, white) and non-events (N, shaded) (9:00 -12:00). See Figure 1 for explanation of symbols. (b) The diurnal cycle of $J_{3,C}$ during spring. The nighttime is missing in this plot due to unavailable SA proxy which uses UVB to be calculated.



Figure 10: Diurnal cycle of median values of calculated formation rate of 3 nm particles (*J*_{3,C}) and condensation sink (CS)
 during different seasons on clear-sky events and non-events.



Figure 11: Relationship between temperature and CS during spring time (11:00 – 12:00) NPF clear-sky (P > 0.7) event days and non-event days color-coded with $J_{3,C}$. Horizontal line is calculated from LDA at 95% confidence relative to nonevents and is demonstrated by equation (6).



Figure 12: Relationship between CS and Temperature (time window: 11:00 – 12:00) NPF clear-sky event days and non-event days. Horizontal line is calculated from spring LDA at 95% confidence relative to non-events and is demonstrated by equation (6).



Figure 13: NPF probability distribution based on the CS and temperature conditions during clear-sky days (11:00 -12:00).
 Marker size indicates number of days included in the probability calculation within every cell.