

1 **Chemical oxidative potential of secondary organic aerosol (SOA) generated from the**
2 **photooxidation of biogenic and anthropogenic volatile organic compounds**

3 *Wing Y. Tuet¹, Yunle Chen², Lu Xu¹, Shierly Fok¹, Dong Gao³, Rodney J. Weber⁴, Nga L. Ng^{1,4*}*

4 ¹School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA

5 ²School of Materials Science and Engineering, Georgia Institute of Technology, Atlanta, GA

6 ³School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA

7 ⁴School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA

8 **Corresponding Author**

9 *email: ng@chbe.gatech.edu

10 Keywords: oxidative potential, particulate matter, secondary organic aerosol, chemical composition

11 Abstract

12 Particulate matter (PM), of which a significant fraction is comprised of secondary organic
13 aerosols (SOA), has received considerable attention due to their health implications. In this study,
14 the water-soluble oxidative potential (OP^{WS}) of SOA generated from the photooxidation of
15 biogenic and anthropogenic hydrocarbon precursors (isoprene, α -pinene, β -caryophyllene,
16 pentadecane, *m*-xylene, and naphthalene) under different reaction conditions (“RO₂ + HO₂”/“RO₂
17 + NO” dominant, dry/humid) was characterized using dithiothreitol (DTT) consumption. The
18 measured intrinsic OP^{WS-DTT} ranged from 9–205 pmol min⁻¹ μ g⁻¹ and were highly dependent on
19 the specific hydrocarbon precursor, with naphthalene and isoprene SOA generating the highest
20 and lowest OP^{WS-DTT}, respectively. Humidity and RO₂ fate affected OP^{WS-DTT} in a hydrocarbon-
21 specific manner, with naphthalene SOA exhibiting the most pronounced effects, likely due to the
22 formation of nitroaromatics. Together, these results suggest that precursor identity may be more
23 influential than reaction condition in determining SOA oxidative potential, demonstrating the
24 importance of sources, such as incomplete combustion, to aerosol toxicity. In the context of other
25 PM sources, all SOA systems with the exception of naphthalene SOA were less DTT active than
26 ambient sources related to incomplete combustion, including diesel and gasoline combustion as
27 well as biomass burning. Finally, naphthalene SOA was as DTT active as biomass burning aerosol,
28 which was found to be the most DTT active OA source in a previous ambient study. These results
29 highlight a need to consider SOA contributions (particularly from anthropogenic hydrocarbons) to
30 health effects in the context of hydrocarbon emissions, SOA yields, and other PM sources.

31 Introduction

32 Numerous epidemiological studies have found associations between elevated particulate
33 matter (PM) concentrations and increased incidences of cardiopulmonary disease, including
34 increases in lung cancer, asthma, chronic obstructive pulmonary disease, arrhythmia, and ischemic
35 heart disease (Li et al., 2008; Pope III and Dockery, 2006; Brunekreef and Holgate, 2002; Dockery
36 et al., 1993; Hoek et al., 2013; Anderson et al., 2011; Pope et al., 2002). Furthermore, ambient PM
37 pollution ranked among the top 10 global risk factors in the 2010 Global Burden of Disease Study,
38 with significant contributions from cardiopulmonary diseases and lower respiratory infections
39 (Lim et al., 2012). Recent epidemiological studies have also found an association between particle
40 oxidative potential and various cardiopulmonary health endpoints (Bates et al., 2015; Fang et al.,
41 2016; Yang et al., 2016; Weichenthal et al., 2016), and results from toxicology studies suggest that
42 PM-induced oxidant production, including reactive oxygen and nitrogen species (ROS/RNS), is a
43 possible mechanism by which PM exposure results in adverse health effects (Li et al., 2003a; Tao
44 et al., 2003; Castro and Freeman, 2001; Gurgueira et al., 2002). These species can initiate
45 inflammatory cascades, which may ultimately lead to oxidative stress and cellular damage
46 (Wiseman and Halliwell, 1996; Hensley et al., 2000). Prolonged stimulation of inflammatory
47 cascades may also lead to chronic inflammation, for which there is a well-established link between
48 chronic inflammation and cancer (Philip et al., 2004). Collectively, these findings suggest a
49 possible link between PM exposure and epidemiologically associated health endpoints as PM can
50 contain ROS/RNS and generate ROS/RNS via redox reactions and by inducing cellular pathways
51 that produce ROS/RNS.

52 Chemical assays in which an anti-oxidant is used to simulate redox reactions that would
53 occur in biological systems have been developed to study the oxidative potential of PM samples

54 (Kumagai et al., 2002; Cho et al., 2005). In these assays, redox-active species in PM samples
55 catalyze electron transfer from the anti-oxidant (e.g., dithiothreitol, DTT; ascorbic acid, AA; etc.)
56 to oxygen, and anti-oxidant decay provides a measure of the concentration of redox-active species
57 in the sample (Fang et al., 2015b). These assays have been utilized extensively to characterize
58 ambient PM samples and source apportionment regressions have been applied to DTT activity
59 results to identify PM sources that may be detrimental to health (Bates et al., 2015; Fang et al.,
60 2015a; Verma et al., 2015a; Verma et al., 2014). Results from these regressions, as well as
61 inhalation and exposure studies, suggest that organic carbon constituents may play a significant
62 role in PM-induced health effects (Li et al., 2003b; Kleinman et al., 2005; Hamad et al., 2015;
63 Verma et al., 2015b). In particular, humic-like substances (HULIS) and oxygenated polyaromatic
64 hydrocarbons (PAH) have been shown to contribute significantly to the redox activity of water-
65 soluble PM samples (Verma et al., 2012; Verma et al., 2015a; Dou et al., 2015; Verma et al.,
66 2015b; Lin and Yu, 2011). Recently, Tuet et al. (2016) also showed that there is a significant
67 correlation between intracellular ROS/RNS production and organic species (water-soluble organic
68 carbon and brown carbon) for summer ambient samples, which suggests that photochemically-
69 driven secondary organic aerosols (SOA) may be important in PM-induced oxidative stress.

70 Many prior studies have focused on the health effects of primary emissions, such as PM
71 directly emitted from diesel and gasoline engines (Bai et al., 2001; Kumagai et al., 2002;
72 McWhinney et al., 2013a; Turner et al., 2015). Conversely, few studies have explored the potential
73 health implications of SOA, which are formed from the oxidation of volatile organic compounds
74 (VOCs) (McWhinney et al., 2013b; Rattanavaraha et al., 2011; Kramer et al., 2016; Lund et al.,
75 2013; McDonald et al., 2010; McDonald et al., 2012; Baltensperger et al., 2008; Arashiro et al.,
76 2016; Platt et al., 2014), even though field studies have shown that SOA often dominate over

77 primary aerosols even in urban environments (Zhang et al., 2007; Jimenez et al., 2009; Ng et al.,
78 2010). The few studies that exist focus on SOA generated from a single class of hydrocarbon
79 precursor or on SOA formed in a simulated urban background (Kramer et al., 2016; McWhinney
80 et al., 2013b; Rattanavaraha et al., 2011; Arashiro et al., 2016; McDonald et al., 2012). While
81 studies on oxidative potential have shown that SOA is indeed redox active, the combined range of
82 oxidative potentials observed for individual SOA systems is quite large and remains unexplored
83 (McWhinney et al., 2013b; Kramer et al., 2016). Furthermore, results from cellular exposure
84 studies are inconclusive, with some studies finding significant response from SOA exposure and
85 others finding little to no response. The exposure dose also differed from study to study, which
86 may result in inconclusive results. This also highlights a need to consider dose-response
87 relationships as demonstrated recently in Tuet et al. (2016). Comparisons between the observed
88 cellular endpoints from exposure to SOA formed from individual precursors are also lacking
89 (Baltensperger et al., 2008; Lund et al., 2013; McDonald et al., 2010; McDonald et al., 2012;
90 Arashiro et al., 2016). As such, there is a lack of perspective in terms of different individual SOA
91 systems and their contributions to PM-induced health effects, making it unclear whether certain
92 responses are indeed toxic for a range of sources and subtypes of PM. However, as cellular assays
93 and animal inhalation experiments are more complex, a systematic study on the oxidative potential
94 of individual SOA systems may be warranted first.

95 In the present study, the water-soluble oxidative potential of SOA generated from various
96 precursors under different reaction conditions was measured using the DTT assay (henceforth
97 referred to as OP^{WS-DTT}). While numerous cell-free assays have been developed to measure
98 oxidative potential, the DTT assay is well-suited for the purposes of this study due to its proven
99 sensitivity to organic carbon constituents and correlation with organic carbon (Janssen et al., 2014;

100 Visentin et al., 2016). Furthermore, there are many previous studies reporting the DTT activities
101 of laboratory-generated SOA and ambient samples for comparison purposes (Kramer et al., 2016;
102 Bates et al., 2015; McWhinney et al., 2013a; McWhinney et al., 2013b; Verma et al., 2015a; Xu
103 et al., 2015a; Xu et al., 2015b; Fang et al., 2015b; Lu et al., 2014). VOCs were chosen to represent
104 the major classes of compounds known to produce SOA upon oxidation by atmospheric oxidants
105 and to include precursors of both anthropogenic and biogenic origins (Table S1). Biogenic
106 precursors include isoprene, α -pinene, and β -caryophyllene, while anthropogenic precursors
107 include pentadecane, *m*-xylene, and naphthalene. Isoprene was chosen as it is the most abundant
108 non-methane hydrocarbon, with estimated global emissions around 500 Tg yr⁻¹ (Guenther et al.,
109 2006). α -pinene and β -caryophyllene were chosen as representative, well-studied monoterpenes
110 and sesquiterpenes, respectively. Both classes of compounds contribute significantly to ambient
111 aerosol (Eddingsaas et al., 2012; Hoffmann et al., 1997; Tasoglou and Pandis, 2015; Goldstein and
112 Galbally, 2007). α -pinene emissions (~50 Tg yr⁻¹) are also on the same order of global
113 anthropogenic emissions (~110 Tg yr⁻¹) (Guenther et al., 1993; Piccot et al., 1992). Similarly,
114 anthropogenic precursors were chosen to include a long-chain alkane (pentadecane), a single-ring
115 aromatic (*m*-xylene), and a poly-aromatic (naphthalene). These classes of compounds are emitted
116 as products of incomplete combustion (Robinson et al., 2007; Jia and Batterman, 2010; Bruns et
117 al., 2016) and have been shown to have considerable SOA yields (e.g., Chan et al., 2009; Ng et
118 al., 2007b; Lambe et al., 2011). In addition to precursor identity, the effects of humidity (dry vs.
119 humid) and NO_x (differing peroxy radical (RO₂) fates, RO₂ + HO₂ vs. RO₂ + NO) on OP^{WS-DTT}
120 were investigated, as these conditions have been shown to affect the chemical composition and
121 mass loading of SOA formed (Chhabra et al., 2010; Chhabra et al., 2011; Eddingsaas et al., 2012;
122 Ng et al., 2007b; Loza et al., 2014; Ng et al., 2007a; Chan et al., 2009; Boyd et al., 2015). Finally,

123 intrinsic OP^{WS-DTT} was compared with bulk aerosol composition, specifically elemental ratios, to
124 investigate whether there is a link between OP^{WS-DTT} and aerosol composition.

125 Methods

126 **Chamber experiments.** SOA from the photooxidation of biogenic and anthropogenic
127 VOCs were generated in the Georgia Tech Environmental Chamber (GTEC) facility. Details of
128 the facility are described elsewhere (Boyd et al., 2015). Briefly, the facility consists of two 12 m³
129 Teflon chambers suspended inside a 21 ft x 12 ft temperature-controlled enclosure, surrounded by
130 black lights (Sylvania 24922) and natural sunlight fluorescent lamps (Sylvania 24477). Multiple
131 sampling ports from each chamber allow for gas- and aerosol-phase measurements, as well as
132 introduction of reagents. Gas-phase measurements include O₃, NO₂, and NO_x concentrations as
133 measured by an O₃ analyzer (Teledyne T400), a cavity attenuated phase shift (CAPS) NO₂ monitor
134 (Aerodyne), and a chemiluminescence NO_x monitor (Teledyne 200EU) respectively. Additionally,
135 a gas chromatography-flame ionization detector (GC-FID, Agilent 7890A) was used to monitor
136 hydrocarbon decay and estimate hydroxyl radical (OH) concentration. In terms of aerosol-phase
137 measurements, aerosol volume concentrations and distributions were measured using a Scanning
138 Mobility Particle Sizer (SMPS, TSI), while bulk aerosol composition was determined using a High
139 Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS, Aerodyne; henceforth
140 referred to as the AMS) (DeCarlo et al., 2006). HR-ToF-AMS data was analyzed using the data
141 analysis toolkit SQUIRREL (v. 1.57) and PIKA (v. 1.16G). Elemental ratios (O:C, H:C, and N:C)
142 were obtained using the method outlined by Canagaratna et al. (2015), and used to calculate the
143 average carbon oxidation state (\overline{OS}_c) (Kroll et al., 2011). Temperature and relative humidity (RH)
144 were monitored using a hydro-thermometer (Vaisala HMP110).

145 Experimental conditions, given in Table 1, were designed to probe the effects of humidity,
146 RO₂ fate, and precursor hydrocarbon on OP^{WS-DTT}. All experiments were performed at ~25 °C
147 under dry (RH < 5%) or humid (RH ~ 45%) conditions. Prior to each experiment, the chambers
148 were flushed with pure air for ~24 hrs. For humid experiments, the chambers were also humidified
149 by means of a bubbler filled with deionized (DI) water during this time. Seed aerosol was then
150 injected by atomizing a 15 mM (NH₄)₂SO₄ seed solution (Sigma Aldrich) until the seed
151 concentration was approximately 20 μg m⁻³. It is noted that for experiment 7 (isoprene SOA under
152 RO₂ + HO₂ dominant, “humid” conditions), experimental conditions deviated due to extremely
153 low SOA mass yields. For this experiment, an acidic seed solution (8 mM MgSO₄ and 16 mM
154 H₂SO₄) was used under dry conditions to promote SOA formation via the IEPOX uptake pathway,
155 which has a higher SOA mass yield compared to the IEPOX + OH pathway and contributes
156 significantly to ambient OA (Surratt et al., 2010; Lin et al., 2012).

157 Once the seed concentration stabilized, hydrocarbon was added by injecting a known
158 volume of hydrocarbon solution [isoprene, 99%; α-pinene, ≥ 99%; β-caryophyllene, > 98.5%;
159 pentadecane, ≥ 99%; *m*-xylene, ≥ 99%; naphthalene, 99% (Sigma Aldrich)] into a glass bulb and
160 passing zero air at 5 L min⁻¹ over the solution until fully evaporated (~10 min). For pentadecane
161 and β-caryophyllene, the glass bulb was gently heated to ensure full evaporation (Tasoglou and
162 Pandis, 2015). Naphthalene was injected by passing pure air over the solid, as outlined in previous
163 studies (Chan et al., 2009). After hydrocarbon injection, OH precursor was added to the chamber.
164 Experiments were conducted under various NO_x conditions where different RO₂ reaction pathways
165 prevailed. For RO₂ + HO₂ experiments, hydrogen peroxide (H₂O₂) was used as the OH precursor.
166 H₂O₂ (50% aqueous solution, Sigma Aldrich) was injected using the method described for
167 hydrocarbon injection to achieve an H₂O₂ concentration of 3 ppm, which yielded OH

168 concentrations on the order of 10^6 molec cm^{-3} . For $\text{RO}_2 + \text{NO}$ experiments, nitrous acid (HONO),
169 was used as the OH precursor. HONO was prepared by adding 10 mL of 1%wt aqueous NaNO_2
170 (VWR International) dropwise into 20 mL of 10%wt H_2SO_4 (VWR International) in a glass bulb.
171 Zero air was then passed over the solution to introduce HONO into the chamber (Chan et al., 2009;
172 Kroll et al., 2005). Photolysis of HONO yielded OH concentrations on the order of 10^7 molec cm^{-3} .
173 NO and NO_2 were also formed as byproducts of HONO synthesis. Once all the H_2O_2 evaporated
174 ($\text{RO}_2 + \text{HO}_2$ experiments) or NO_x concentrations stabilized ($\text{RO}_2 + \text{NO}$ experiments), the UV lights
175 were turned on to initiate photooxidation.

176 **Aerosol collection and extraction.** Aerosol samples were collected onto 47 mm Teflon™
177 filters (0.45 μm pore size, Pall Laboratory) for approximately 1.5 hrs at a flow rate of 28 L min^{-1} .
178 For each experiment, two filters (front filter and backing filter) were loaded in series to account
179 for possible sampling artifacts (Conny and Slater, 2002). Total mass collected was determined by
180 integrating the SMPS volume concentration as a function of time over the filter collection period
181 and using the total volume of air collected. Volume concentrations were integrated using time-
182 dependent data. Background filters containing seed and OH precursor (H_2O_2 or HONO) only at
183 experimental conditions were also collected to account for potential H_2O_2 or HONO uptake, which
184 may influence oxidative potential. Collected filter samples were placed in sterile petri dishes,
185 sealed with Parafilm M®, and stored at $-20 \text{ }^\circ\text{C}$ until extraction and analysis (Fang et al., 2015b).
186 Prior to determining $\text{OP}^{\text{WS-DTT}}$, collected particles were extracted in DI water by submerging the
187 filter and sonicating for 1 hr using an Ultrasonic Cleanser (VWR International) (Fang et al., 2015a).
188 Sonication steps were performed in 30 min intervals with water replacement after each interval to
189 reduce bath temperature. After sonication, extracts were filtered using 0.45 μm PTFE syringe

190 filters (Fisherbrand™) to remove insoluble material (Fang et al., 2015b). All filter samples were
191 extracted within 1-2 days of collection and analyzed immediately following extraction.

192 **Oxidative potential.** The decay of DTT, a chemical species that reacts with redox-active
193 species in a sample via electron transfer reactions, was used as a measure of oxidative potential
194 (Cho et al., 2005; Kumagai et al., 2002). The intrinsic OP^{WS-DTT} of aerosol samples, as well as
195 method blanks and positive controls (9,10-phenanthraquinone), was determined using a semi-
196 automated DTT system. Specifics of the high-throughput system are detailed in Fang et al. (2015b)
197 Briefly, the method consisted of three main steps: (1) oxidation of DTT by redox-active species in
198 the sample, (2) reaction of residual DTT with DTNB to form 2-nitro-5-mercaptobenzoic acid
199 (TNB), repeated at specific time intervals, and (3) measurement of TNB to determine DTT
200 consumption. After each time interval and between samples, the system was flushed with DI water.

201 Results and Discussion

202 **Laboratory-generated aerosol.** Over the course of each experiment, gas and aerosol
203 composition was continuously monitored. A typical time series for NO, NO₂, gas-phase
204 hydrocarbon concentration, and aerosol mass concentration is shown in Fig. 1 for naphthalene
205 photooxidation under RO₂ + NO dominant reaction conditions. Hydrocarbon decay was monitored
206 using GC-FID, while initial gas-phase hydrocarbon concentrations were determined using the
207 chamber volume and mass of hydrocarbon injected. Following irradiation, NO decreased due to
208 reaction with RO₂ from hydrocarbon oxidations. Nevertheless, ozone formation was suppressed
209 owing to the high NO concentration throughout the experiment. Aerosol growth is observed
210 shortly after initiation of photooxidation (i.e., turning on the lights) due to the efficient photolysis
211 of HONO, which produced a high OH concentration on the order of 10^7 molec cm⁻³. Once HONO

212 was completely consumed, no further decay in the parent hydrocarbon and growth in aerosol mass
213 were observed.

214 For each experiment, aerosol chemical composition was also monitored using the AMS.
215 The average AMS mass spectra (Fig. S1) for all VOC systems were consistent with those reported
216 in previous studies (Chhabra et al., 2010; Chhabra et al., 2011). For RO₂ + NO dominant
217 experiments, the NO⁺:NO₂⁺ ratio has been used extensively in previous studies to differentiate
218 between organic and inorganic nitrates (Farmer et al., 2010; Fry et al., 2009; Boyd et al., 2015; Xu
219 et al., 2015b). The observed NO⁺:NO₂⁺ ratio for all RO₂ + NO dominant experiments (4.2–6.1)
220 was higher than that observed for inorganic (ammonium) nitrates (~2.3), which indicates that these
221 peaks are likely from organic nitrates rather than inorganic nitrates. The observed range is also
222 consistent with values measured in previous organic nitrate studies for similar VOC systems and
223 ambient studies (Bruns et al., 2010; Sato et al., 2010; Xu et al., 2015b). Elemental ratios (O:C,
224 H:C, and N:C) were also obtained for each SOA system using the AMS. The aerosol systems
225 investigated span a wide range of O:C ratios, as observed in previous laboratory and field studies
226 (Chhabra et al., 2011; Lambe et al., 2011; Jimenez et al., 2009; Ng et al., 2010).

227 **Effect of hydrocarbon precursor and reaction condition on oxidative potential.** To
228 investigate whether different types of SOA differ in toxicity, the OP^{WS-DTT}, a measure of the
229 concentration of redox-active species present in a sample, was measured for SOA generated from
230 six VOCs under three conditions (see Table 1 for specifics). The blank-corrected OP^{WS-DTT},
231 represented on a per mass (μg) basis, are shown in Fig. 2. Uncertainties associated with OP^{WS-DTT}
232 determination were approximated using a 15% coefficient of variation, in accordance with
233 previous studies using the same semi-automated system (Fang et al., 2015b). The OP^{WS-DTT} of all
234 backing filters and background filters were also measured and found to be within the uncertainty

235 for blank Teflon filters, which indicates that there were no observable sampling artifacts, gaseous
236 absorption onto Teflon filters, or H₂O₂/HONO uptake onto seed particles.

237 Overall, it is clear that the hydrocarbon precursor identity influenced OP^{WS-DTT}, with
238 naphthalene having the highest intrinsic DTT activity (Fig. 2). All other hydrocarbon precursors
239 investigated produced SOA with relatively low intrinsic OP^{WS-DTT} (~9 – 45 pmol min⁻¹ μg⁻¹). For
240 isoprene, the SOA in this study was generated through different reaction pathways, including
241 isoprene photooxidation under different RO₂ fates and IEPOX reactive uptake to acidic seed
242 particles. Although these different conditions produced different products and SOA compositions
243 (Xu et al., 2014; Surratt et al., 2010; Chan et al., 2010), the OP^{WS-DTT} is very similar. It is important
244 to note that the intrinsic OP^{WS-DTT} for SOA generated under all conditions in this study are in
245 agreement with the isoprene-derived OA factor resolved from positive matrix factorization (PMF)
246 analysis of ambient AMS data (Fig. 4) (Xu et al., 2015a; Xu et al., 2015b; Verma et al., 2015a).
247 The isoprene-derived OA from ambient measurements is largely attributed to IEPOX uptake, but
248 possibly contains some contribution from other isoprene oxidation pathways (Xu et al., 2015a; Xu
249 et al., 2015b). The similarity between laboratory-generated and ambient isoprene SOA suggests
250 that isoprene SOA may have low OP^{WS-DTT} regardless of reaction conditions. A previous
251 laboratory chamber study by Kramer et al. (2016) also measured the DTT activity of isoprene SOA
252 produced via different pathways, including SOA formed from direct photooxidation of isoprene.
253 It was found that isoprene SOA formed under “high-NO_x” conditions was more DTT active than
254 that formed under “low-NO_x” conditions. These results are in contrast with those obtained in this
255 study, where the OP^{WS-DTT} of isoprene SOA was similar regardless of reaction condition. However,
256 we caution that 1) the SOA measured in Kramer et al. (2016) was formed under different
257 experimental conditions, and 2) they utilized a different method for measuring DTT consumption

258 (i.e., different extraction solvent, different initial DTT concentration, different method for
259 quantifying DTT activity), therefore the results from their study and ours may not be directly
260 comparable. For instance, for isoprene photooxidation experiments, the “low-NO_x” conditions in
261 Kramer et al. (2016) corresponded to “5 ppm isoprene and 200 ppb NO”, where the reaction regime
262 was largely defined by the VOC/NO_x ratio. It has been shown previously that SOA formed under
263 the same VOC/NO_x conditions can be drastically different and the use of this metric might not
264 necessarily reflect the actual peroxy radical fate (Ng et al., 2007b; Kroll and Seinfeld, 2008;
265 Wennberg, 2013). In our study, the “low-NO_x” experimental condition is defined by the fate of
266 peroxy radicals directly, i.e., no NO_x added, but with the presence of H₂O₂ to enhance the RO₂ +
267 HO₂ reaction pathway, which is dominant in ambient environments when NO_x levels are low.

268 α-pinene, β-caryophyllene, and pentadecane produced low OP^{WS-DTT} across all conditions
269 explored in this study (Fig. 2). Specifically, the SOA formed under different reaction conditions
270 do not appear to have significantly different OP^{WS-DTT}, even though different NO_x conditions have
271 been shown to affect SOA loading and composition due to competing RO₂ chemistry (Chan et al.,
272 2009; Eddingsaas et al., 2012; Loza et al., 2014; Ng et al., 2007a). For instance, under conditions
273 that favor RO₂ + NO, organic nitrates are formed, whereas under conditions that favor RO₂ + HO₂,
274 organic peroxides are the predominant products. In this study, the formation of organic nitrates is
275 evident in the RO₂ + NO experiments with the relatively higher NO⁺:NO₂⁺ ratio in the AMS mass
276 spectra. It is possible that the organic peroxides and organic nitrates formed from the oxidation of
277 these precursors are both not highly redox active, such that the overall OP^{WS-DTT} is similar even
278 though the products differ. Further studies are required to establish this.

279 Similarly, the OP^{WS-DTT} of SOA formed from *m*-xylene under conditions that favor
280 different RO₂ fates were not significantly different. Since OP^{WS-DTT} is intended as a measure of

281 redox activity, the reaction products' ability to participate in electron transfer may explain this lack
282 of difference (e.g., lack of conjugated systems and associated pi bonds with unbound electrons).
283 Under both $\text{RO}_2 + \text{HO}_2$ and $\text{RO}_2 + \text{NO}$ pathways, a large portion of *m*-xylene oxidation products
284 do not retain the aromatic ring (Vivanco and Santiago, 2010; Jenkin et al., 2003). Therefore, these
285 products may have similar $\text{OP}^{\text{WS-DTT}}$ as reaction products of α -pinene, β -caryophyllene, and
286 pentadecane, which also do not contain an aromatic ring. Under humid conditions, aerosol formed
287 from the oxidation of *m*-xylene were more DTT active than those formed under dry conditions.
288 The AMS mass spectra for aerosol formed under humid conditions also differs notably for several
289 characteristic fragments (Fig. S2), which may explain the difference observed in $\text{OP}^{\text{WS-DTT}}$. More
290 specifically, m/z 44, which serves as an indication of oxidation (O:C ratio) (Ng et al., 2010), is
291 very different for this experiment (dry signal: 0.098 vs. humid signal: 0.15). It is possible that the
292 degree of oxidation may be an important factor for SOA formed from the same hydrocarbon, and
293 systematic chamber studies investigating changes in O:C for SOA formed from a single
294 hydrocarbon precursor would be valuable. Previous studies involving the effect of humidity on
295 SOA composition also yield mixed results, with some finding significant changes in SOA
296 composition and yields (Nguyen et al., 2011; Wong et al., 2015; Healy et al., 2009; Stirnweis et
297 al., 2016) and others reporting little difference (Boyd et al., 2015; Edney et al., 2000; Cocker III
298 et al., 2001). Humidity effects are therefore highly hydrocarbon-dependent. Further study into the
299 specific oxidation mechanisms and products in the photooxidation of aromatic hydrocarbon under
300 dry and humid conditions may be warranted to understand the difference in DTT activity.

301 For naphthalene, the $\text{OP}^{\text{WS-DTT}}$ measured for SOA generated under dry, $\text{RO}_2 + \text{HO}_2$
302 dominant conditions is in agreement with that measured by McWhinney et al. (2013b) (Fig. 4),
303 which generated naphthalene SOA under similar chamber conditions using the same OH radical

304 precursor. These values should be directly comparable as the same standard method described by
305 Cho et al. (2005) was used to obtain the oxidative potentials in both McWhinney et al. (2013b)
306 and this study. The OP^{WS-DTT} of naphthalene aerosol also appears to be strongly influenced by
307 humidity and RO_2 fate (Fig. 2), with higher toxicities observed for aerosol formed under both
308 humid and $RO_2 + NO$ dominant conditions. The effect of RO_2 fate may be explained by the
309 different products known to form from $RO_2 + HO_2$ and $RO_2 + NO$ reaction pathways. Many of the
310 same products, including naphthoquinones and all of the ring-opening derivatives of 2-
311 formylcinnamaldehyde, are formed under both reaction conditions (Kautzman et al., 2010).
312 Naphthoquinones are also known to be DTT active and have been shown to account for
313 approximately 21% of the DTT activity observed for naphthalene SOA (Charrier and Anastasio,
314 2012; McWhinney et al., 2013b). In addition to these products, nitroaromatics including
315 nitronaphthols and nitronaphthalenes are formed under $RO_2 + NO$ conditions (Kautzman et al.,
316 2010). The nitrite group next to the aromatic ring in these products may further promote electron
317 transfer between nitroaromatics and DTT, resulting in more DTT consumption and a higher OP^{WS-}
318 DTT . This effect was not observed for *m*-xylene SOA due to the formation of predominantly ring-
319 opening products (Vivanco and Santiago, 2010; Jenkin et al., 2003). The presence of an aromatic
320 ring in SOA products may therefore be important for determining oxidative potentials and
321 polyaromatic precursors may yield products of substantial toxicity. This is further supported by
322 the observation that the AMS mass spectra for highly DTT active naphthalene SOA contains peaks
323 at m/z 77 and m/z 91, which are indicative of aromatic phenyl and benzyl ions (Chhabra et al.,
324 2010; McLafferty and Tureček, 1993). Additionally, peaks indicative of aromatic compounds
325 greater than m/z 120 were observed with similar mass spectral features as those reported for aerosol
326 generated from naphthalene oxidation by OH radicals in previous studies (Riva et al., 2015).

327 Aromatic species are also exclusive to HULIS (Sannigrahi et al., 2006), and ambient data have
328 shown that HULIS is a significant aerosol component contributing to OP^{WS-DTT} (Verma et al.,
329 2015b; Verma et al., 2012; Dou et al., 2015; Lin and Yu, 2011).

330 Bulk aerosol elemental ratios (O:C, H:C, and N:C) were also determined for each SOA
331 system as different types of aerosol are known to span a wide range of O:C (Chhabra et al., 2011;
332 Lambe et al., 2011). All elemental ratios were stable during the filter collection period and could
333 thus be represented by a single value. To visualize these differences in oxidation, the van Krevelen
334 diagram was utilized (Fig. 3), as changes in the slope of data points within the van Krevelen space
335 can provide information on SOA functionalization (Heald et al., 2010; Van Krevelen, 1950; Ng et
336 al., 2011). Starting from the precursor hydrocarbon, a slope of 0 indicates addition of alcohol
337 groups, a slope of -1 indicates addition of carbonyl and alcohol groups on separate carbons or
338 addition of carboxylic acids, and a slope of -2 indicates addition of ketones or aldehydes. Previous
339 studies show that both laboratory-generated and ambient OA occupy a narrow van Krevelen space
340 with a slope of $\sim -1 - -0.5$ (Heald et al., 2010; Ng et al., 2011). Ambient data included in Fig. 3 are
341 for different organic aerosol subtypes resolved from PMF analysis of AMS data collected in the
342 southeastern U.S. (Verma et al., 2015a; Xu et al., 2015a; Xu et al., 2015b).

343 The laboratory-generated aerosols span the range of H:C and O:C observed in the ambient.
344 As seen in Fig. 3 (data points sized by intrinsic OP^{WS-DTT}), while different reaction conditions
345 produced aerosol of differing composition (i.e., different O:C and H:C), the intrinsic OP^{WS-DTT}
346 does not appear to be affected by these differences. On the other hand, the hydrocarbon precursor
347 identity influences OP^{WS-DTT} substantially. It has been shown that ambient OA from different
348 sources can become increasingly oxidized (increasing O:C ratio) with atmospheric aging (Jimenez
349 et al., 2009; Ng et al., 2011). Based on the results shown in Fig. 3, it appears that a higher O:C

350 ratio did not correspond to a higher OP^{WS-DTT} . This is true for both the laboratory-generated SOA
351 in this study and the different OA subtypes resolved from ambient data (Verma et al., 2015a; Xu
352 et al., 2015a; Xu et al., 2015b). Nevertheless, the O:C ratios for individual systems (i.e. SOA
353 formed from the same hydrocarbon precursor) may affect the intrinsic OP^{WS-DTT} . Indeed, for
354 several SOA systems (β -caryophyllene, pentadecane, and *m*-xylene), SOA with higher O:C ratios
355 also had a higher intrinsic OP^{WS-DTT} (Fig. 2, 3). For SOA systems formed under $RO_2 + NO$
356 dominant conditions, N:C ratios were also determined to investigate if there is a link between N:C
357 and intrinsic DTT activity (Fig. S3). Again, with the exception of naphthalene SOA, the intrinsic
358 OP^{WS-DTT} does not appear to be affected by N:C ratio even though the systems explored span a
359 wide range of N:C. This is consistent with that observed in the van Krevelen diagram and further
360 emphasizes the importance of hydrocarbon identity in determining oxidative potentials.

361 **Comparison to other types of PM.** In order to evaluate how the oxidative potential of
362 individual SOA systems compares to other sources and subtypes of PM, the intrinsic OP^{WS-DTT}
363 from this study are compared to values reported in the literature (Fig. 4). Comparatively, SOA
364 formed from the photooxidation of isoprene, α -pinene, β -caryophyllene, pentadecane, and *m*-
365 xylene were not very DTT active and produced low intrinsic OP^{WS-DTT} . The OP^{WS-DTT} of these
366 aerosol systems were also within the range of various OA subtypes resolved from ambient data.
367 The method for determining intrinsic OP^{WS-DTT} for various OA subtypes is provided in the
368 Supplement. As noted earlier, the OP^{WS-DTT} for isoprene SOA generated in this study is similar to
369 the isoprene-derived OA factor from ambient data. The other ambient OA factors include a highly
370 oxidized MO-OOA (more-oxidized oxygenated OA) factor resolved from PMF analysis of
371 ambient OA data, as well as an oxidized organic aerosol factor containing contributions from
372 biogenic SOA (other OC) resolved using the chemical mass balance (CMB) method with

373 ensemble-averaged source impact profiles (Bates et al., 2015; Xu et al., 2015a; Xu et al., 2015b;
374 Verma et al., 2014). While sources of MO-OOA have not been identified, studies have shown that
375 the aerosol mass spectra for various sources of OA approach that of MO-OOA as it ages (Ng et
376 al., 2010) and it has been speculated that MO-OOA may contain aerosol from multiple aged
377 sources (Xu et al., 2015b). Furthermore, MO-OOA has been shown to have widespread
378 contributions across urban and rural sites, as well as different seasons (Xu et al., 2015a; Xu et al.,
379 2015b). On the other hand, naphthalene SOA was highly DTT active with an OP^{WS-DTT} on the
380 order of biomass burning OA [BBOA (Verma et al., 2015a), BURN (Bates et al., 2015)]. The
381 BBOA and BURN factors were resolved using different source apportionment methods and as
382 such, the range for comparison is large. Here, we focus on BBOA as Verma et al. (2015a)
383 previously showed that BBOA had the highest intrinsic DTT activity among all OA subtypes
384 resolved from PMF analysis of ambient AMS data collected in the southeastern U.S. (see Fig. 4
385 for comparison). Because naphthalene aerosol formed under $RO_2 + NO$ dominant conditions may
386 be even more redox active than BBOA and anthropogenic emissions are more abundant in urban
387 environments with higher NO_x , this system warrants further systematic studies. It should however
388 be noted that comparisons of intrinsic DTT activities between SOA from a pure VOC and an
389 ambient source is difficult. BBOA is a source that contains many compounds, some of which may
390 not be redox active. Thus, although it may contain highly DTT-active components with high
391 intrinsic activities, the overall intrinsic activity will be much lower. As a result, a direct comparison
392 with pure naphthalene SOA on a per mass basis is tenuous. However, naphthalene SOA formed
393 under urban conditions ($RO_2 + NO$) also produces nitroaromatics, which may induce DNA breaks
394 and induce other mutagenic effects (Baird et al., 2005; Helmig et al., 1992). As such, aerosols

395 formed from photooxidation of PAHs may be a particularly important OA source in terms of PM
396 health effects.

397 Other common sources of PM are those related to traffic. Previous studies have determined
398 that products of incomplete combustion include quinones capable of participating in redox
399 reactions, including the oxidation of DTT (Kumagai et al., 2002; McWhinney et al., 2013a). The
400 SOA systems investigated, including isoprene, α -pinene, β -caryophyllene, pentadecane, and *m*-
401 xylene produced SOA that were less DTT active than diesel exhaust particles (DEP) collected from
402 light-duty diesel vehicle (LDDV) engines operated under various conditions (McWhinney et al.,
403 2013a) and resolved for heavy-duty diesel vehicles (HDDV) from ambient data (Bates et al., 2015).
404 It should be noted that the DTT activity reported for DEP includes both water-soluble and water-
405 insoluble fractions (total DTT activity), whereas the DTT activity measured for SOA is water-
406 soluble. However, there should be very little contribution from water-insoluble species to SOA
407 (McWhinney et al., 2013a). Conversely, the intrinsic OP^{WS-DTT} of naphthalene SOA was on par
408 with that of light-duty gasoline vehicles (LDGV) and higher than that of HDDV and DEP (Verma
409 et al., 2014; Bates et al., 2015). Since naphthalene may also be emitted from gasoline and diesel
410 combustion (Jia and Batterman, 2010), traffic-related controls may be extremely important to
411 control these highly DTT active sources. Furthermore, since SOA often dominate over POA even
412 in urban centers (Zhang et al., 2007; Ng et al., 2011), even SOA that is only slightly DTT active
413 may contribute significantly to PM-induced health effects.


414 **Implications.** The water-soluble oxidative potential, as measured by DTT consumption,
415 was determined for SOA generated from six different hydrocarbon precursors under three
416 conditions of varying humidity and RO_2 fate. Results from this study demonstrate that hydrocarbon
417 precursor identity influenced intrinsic SOA oxidative potential substantially. The biogenic and

418 anthropogenic precursors investigated yielded SOA with OP^{WS-DTT} ranging from 9–205 pmol min^{-1}
419 μg^{-1} , with isoprene SOA and naphthalene SOA having the lowest and highest intrinsic OP^{WS-DTT}
420 respectively. In general, OP^{WS-DTT} for biogenic SOA were lower than those for anthropogenic
421 SOA. Therefore, to evaluate overall oxidative potentials of ambient SOA, hydrocarbon precursor
422 emissions and their corresponding SOA formation potential must be considered. Moreover, it may
423 be possible to roughly estimate regional oxidative potentials using individual intrinsic OP^{WS-DTT}
424 of different types of SOA in conjunction with VOC emissions and SOA loadings in models. For
425 instance, DTT activities of aerosols collected in Beijing, China (77–111 $\text{pmol min}^{-1} \mu\text{g}^{-1}$) (Lu et
426 al., 2014), where anthropogenic emissions dominate, more closely resemble the OP^{WS-DTT} of
427 naphthalene SOA, whereas ambient aerosols collected in the southeastern U.S. have DTT activities
428 (25–36 $\text{pmol min}^{-1} \mu\text{g}^{-1}$) (Fang et al., 2015b) that more closely resemble those of biogenic SOA.
429 It may therefore be informative to investigate whether concentration addition can be applied to
430 DTT consumption by exploring well-characterized PM mixtures.

431 Chamber reaction conditions, including relative humidity and specific RO_2 fate, influenced
432 SOA elemental composition substantially and affected OP^{WS-DTT} in a hydrocarbon-specific
433 manner, although hydrocarbon identity was by far the most influential in determining OP^{WS-DTT} .
434 For several VOCs (isoprene, α -pinene, β -caryophyllene, and pentadecane), the reaction conditions
435 had a negligible effect on OP^{WS-DTT} , which suggests that the organic peroxides and organic nitrates
436 formed from the oxidation of these precursors may have similarly low redox activity. An
437 investigation on the redox activity of individual known photooxidation products, including organic
438 peroxides and organic nitrates, may elucidate further information on the lack of reaction condition
439 effect. Similarly, nitroaromatics may explain the difference observed between naphthalene aerosol
440 formed under different RO_2 reaction pathways as the nitrite group may promote electron transfer


441 and result in a higher OP^{WS-DTT} . This effect was not observed for *m*-xylene SOA, due to the
442 formation of predominantly ring-opening products. The loss of the aromatic ring may also explain
443 the differences in intrinsic OP^{WS-DTT} . For instance, naphthalene SOA, which contains many
444 aromatic ring-retaining products, is as redox active as BBOA, one of the most DTT active aerosol
445 subtypes found in ambient studies. On the other hand, *m*-xylene SOA with predominantly aromatic
446 ring-breaking products is much less redox active and the measured OP^{WS-DTT} is lower than that of
447 traffic-related sources and several OA subtypes (BBOA and cooking OA, COA). This further
448 supports earlier findings (Verma et al., 2015b) that the poly-aromatic ring structure may be an
449 important consideration for understanding SOA redox activity, which may have implications for
450 cellular redox imbalance (Tuet et al., 2016). Furthermore, nitroaromatics and polyaromatics may
451 also have significant health effects beyond redox imbalance, including various mutagenic effects
452 (Baird et al., 2005; Helmig et al., 1992). As such, hydrocarbon precursors forming aromatic ring-
453 retaining products may be the most important to consider in PM-induced health effects, in terms
454 of oxidative potential. This is consistent with many studies using DTT to show oxidative potential
455 associated with sources related to incomplete combustion (Bates et al., 2015; Verma et al., 2014;
456 McWhinney et al., 2013b) and the identification of HULIS (Verma et al., 2015b; Dou et al., 2015;
457 Lin and Yu, 2011), and more specifically, quinones as key components contributing to oxidative
458 potential (Verma et al., 2014). Finally, redox-active metals are also emitted by traffic through
459 mechanical processes, such as brake and tire wear (Charrier and Anastasio, 2012; Fang et al.,
460 2015a). These species have not be considered in the chamber experiments explored in this study.
461 Inclusion of redox-active metals in future SOA experiments may be valuable to further understand
462 the roles of SOA and metal species in overall redox activity.

463




464

465 **Figure 1.** Typical reaction profile for a chamber experiment under RO₂ + NO dominant conditions.
 466 NO and NO₂ concentrations were monitored by CAPS NO₂ and chemiluminescence NO_x monitors,
 467 respectively. Hydrocarbon decay was monitored using GC-FID, while initial hydrocarbon
 468 (naphthalene) concentrations were determined using the chamber volume and mass of hydrocarbon
 469 injected. Aerosol mass concentrations were determined using volume concentrations obtained
 470 from SMPS and assuming an aerosol density of 1 g cm⁻³. Mass concentrations have been corrected
 471 for particle wall loss (Nah et al., 2016).




472

473 **Figure 2.** Intrinsic DTT activities for SOA generated from various hydrocarbon precursors (ISO:
 474 isoprene, AP: α -pinene, BCAR: β -caryophyllene, PD: pentadecane, MX: *m*-xylene, and NAPH:
 475 naphthalene) under various conditions (**red circles**: dry, RO₂ + HO₂; **blue squares**: humid, RO₂
 476 + HO₂; and **black triangles**: dry, RO₂ + NO). Dry, RO₂ + HO₂ experiments were repeated to
 477 ensure reproducibility in SOA generation and collection. Error bars represent a 15% coefficient
 478 of variation (Fang et al., 2015b).



479
 480 **Figure 3.** van Krevelen plot for various SOA systems. Data points are colored by SOA system
 481 (red: isoprene, yellow: α -pinene, green: β -caryophyllene, light blue: pentadecane, blue: *m*-xylene,
 482 and purple: naphthalene), shaped according to reaction conditions (circle: dry, $\text{RO}_2 + \text{HO}_2$; square:
 483 humid, $\text{RO}_2 + \text{HO}_2$; and triangle: dry, $\text{RO}_2 + \text{NO}$), and sized by intrinsic DTT activity. OA factors
 484 resolved from PMF analysis of ambient AMS data are shown as black markers, also sized by
 485 intrinsic DTT activity. Hydrocarbon precursors are shown as stars, colored by SOA system.
 486 Specifics on site locations and factor resolution methods are described elsewhere. COA: cooking
 487 OA, BBOA: biomass burning OA, Isoprene-OA: isoprene-derived OA, MO-OOA: more-oxidized
 488 oxygenated OA (Verma et al., 2015a; Xu et al., 2015a; Xu et al., 2015b).



489

490 **Figure 4.** Intrinsic DTT activities for chamber SOA, various PM subtypes resolved from
 491 ambient data, and diesel exhaust particles. It should be noted that the DTT activity for isoprene
 492 SOA in Kramer et al. (2016) was determined using a different DTT method and may not be
 493 directly comparable. All other studies shown used the method outlined in Cho et al. (2005). DTT
 494 activities obtained in this study are shaped by reaction condition (circle: dry, RO₂ + HO₂; square:
 495 humid, RO₂ + HO₂; triangle: dry, RO₂ + NO). Specifics on site locations and factor resolution
 496 methods are described elsewhere. DTT activities for Beijing and Atlanta are averages obtained
 497 across multiple seasons. Isoprene-OA: isoprene-derived OA, MO-OOA: more-oxidized
 498 oxygenated OA, BBOA: biomass burning OA, COA: cooking OA, LDGV: light-duty gasoline
 499 vehicles, HDDV: heavy-duty diesel vehicles, BURN: biomass burning, DEP: diesel exhaust

500 particles (Kramer et al., 2016; McWhinney et al., 2013b; Verma et al., 2015a; Bates et al., 2015;
501 McWhinney et al., 2013a; Xu et al., 2015a; Xu et al., 2015b; Lu et al., 2014; Fang et al., 2015b).

502 **Table 1.** Experimental conditions.

Experiment	Compound	OH precursor	Relative humidity (%)	[HC] ₀ (ppb)	[SOA] ^c (µg m ⁻³)
1 ^a	isoprene	H ₂ O ₂	<5%	97	5.73
2 ^a	α-pinene	H ₂ O ₂	<5%	191	119
3 ^a	β-caryophyllene	H ₂ O ₂	<5%	36	221
4 ^a	pentadecane	H ₂ O ₂	<5%	106	9.71
5 ^a	<i>m</i> -xylene	H ₂ O ₂	<5%	450	89.3
6 ^a	naphthalene	H ₂ O ₂	<5%	178	128
7	isoprene	H ₂ O ₂	<5% ^b	97	17.1
8	α-pinene	H ₂ O ₂	40%	334	154
9	β-caryophyllene	H ₂ O ₂	42%	63	230
10	pentadecane	H ₂ O ₂	45%	106	23.5
11	<i>m</i> -xylene	H ₂ O ₂	45%	450	13.9
12	naphthalene	H ₂ O ₂	44%	431	132
13	isoprene	HONO	<5%	970	148
14	α-pinene	HONO	<5%	174	166
15	β-caryophyllene	HONO	<5%	21	80.8
16	pentadecane	HONO	<5%	74	35.7
17	<i>m</i> -xylene	HONO	<5%	431	153
18	naphthalene	HONO	<5%	145	142

503 ^a These experiments were repeated to establish reproducibility; ^b Acidic seed (8 mM MgSO₄ and
504 16 mM H₂SO₄) was used instead of 8 mM (NH₄)₂SO₄; ^c Average SOA concentration in the
505 chamber during filter collection

506 ACKNOWLEDGMENT

507 This work was supported by the Health Effects Institute under research agreement No. 4943-
508 RFA13-2/14-4. Wing Y. Tuet acknowledges support by the National Science Foundation
509 Graduate Research Fellowship under Grant No. DGE-1148903.

510 ABBREVIATIONS

511 PM: particulate matter; SOA: secondary organic aerosol; ROS/RNS: reactive oxygen and
512 nitrogen species; DTT: dithiothreitol; OP^{WS}: oxidative potential of water-soluble species

513 REFERENCES

- 514 Anderson, J. O., Thundiyil, J. G., and Stolbach, A.: Clearing the Air: A Review of the Effects of
515 Particulate Matter Air Pollution on Human Health, *Journal of Medical Toxicology*, 8, 166-175,
516 10.1007/s13181-011-0203-1, 2011.
- 517 Arashiro, M., Lin, Y. H., Sexton, K. G., Zhang, Z., Jaspers, I., Fry, R. C., Vizuete, W. G., Gold,
518 A., and Surratt, J. D.: In Vitro Exposure to Isoprene-Derived Secondary Organic Aerosol by
519 Direct Deposition and its Effects on COX-2 and IL-8 Gene Expression, *Atmos. Chem. Phys.*
520 *Discuss.*, 2016, 1-29, 10.5194/acp-2016-371, 2016.
- 521 Bai, Y., Suzuki, A. K., and Sagai, M.: The cytotoxic effects of diesel exhaust particles on human
522 pulmonary artery endothelial cells in vitro: role of active oxygen species, *Free Radical Biology*
523 *and Medicine*, 30, 555-562, [http://dx.doi.org/10.1016/S0891-5849\(00\)00499-8](http://dx.doi.org/10.1016/S0891-5849(00)00499-8), 2001.
- 524 Baird, W. M., Hooven, L. A., and Mahadevan, B.: Carcinogenic polycyclic aromatic
525 hydrocarbon-DNA adducts and mechanism of action, *Environmental and Molecular*
526 *Mutagenesis*, 45, 106-114, 10.1002/em.20095, 2005.
- 527 Baltensperger, U., Dommen, J., Alfarra, R., Duplissy, J., Gaeggeler, K., Metzger, A., Facchini,
528 M. C., Decesari, S., Finessi, E., Reining, C., Schott, M., Warnke, J., Hoffmann, T., Klatzer, B.,
529 Puxbaum, H., Geiser, M., Savi, M., Lang, D., Kalberer, M., and Geiser, T.: Combined
530 determination of the chemical composition and of health effects of secondary organic aerosols:
531 The POLYSOA project, *J. Aerosol Med. Pulm. Drug Deliv.*, 21, 145-154,
532 10.1089/jamp.2007.0655, 2008.
- 533 Bates, J. T., Weber, R. J., Abrams, J., Verma, V., Fang, T., Klein, M., Strickland, M. J., Sarnat,
534 S. E., Chang, H. H., Mulholland, J. A., Tolbert, P. E., and Russell, A. G.: Reactive Oxygen
535 Species Generation Linked to Sources of Atmospheric Particulate Matter and Cardiorespiratory
536 Effects, *Environmental Science & Technology*, 49, 13605-13612, 10.1021/acs.est.5b02967,
537 2015.

538 Boyd, C. M., Sanchez, J., Xu, L., Eugene, A. J., Nah, T., Tuet, W. Y., Guzman, M. I., and Ng, N.
539 L.: Secondary organic aerosol formation from the β -pinene+NO₃ system: effect of
540 humidity and peroxy radical fate, *Atmos. Chem. Phys.*, 15, 7497-7522, 10.5194/acp-15-7497-
541 2015, 2015.

542 Brunekreef, B., and Holgate, S. T.: Air pollution and health, *Lancet*, 360, 1233-1242, 2002.

543 Bruns, E. A., Perraud, V., Zelenyuk, A., Ezell, M. J., Johnson, S. N., Yu, Y., Imre, D.,
544 Finlayson-Pitts, B. J., and Alexander, M. L.: Comparison of FTIR and Particle Mass
545 Spectrometry for the Measurement of Particulate Organic Nitrates, *Environmental Science &*
546 *Technology*, 44, 1056-1061, 10.1021/es9029864, 2010.

547 Bruns, E. A., El Haddad, I., Slowik, J. G., Kilic, D., Klein, F., Baltensperger, U., and Prévôt, A.
548 S. H.: Identification of significant precursor gases of secondary organic aerosols from residential
549 wood combustion, *Scientific Reports*, 6, 27881, 10.1038/srep27881
550 <http://www.nature.com/articles/srep27881#supplementary-information>, 2016.

551 Canagaratna, M. R., Jimenez, J. L., Kroll, J. H., Chen, Q., Kessler, S. H., Massoli, P.,
552 Hildebrandt Ruiz, L., Fortner, E., Williams, L. R., Wilson, K. R., Surratt, J. D., Donahue, N. M.,
553 Jayne, J. T., and Worsnop, D. R.: Elemental ratio measurements of organic compounds using
554 aerosol mass spectrometry: characterization, improved calibration, and implications, *Atmos.*
555 *Chem. Phys.*, 15, 253-272, 10.5194/acp-15-253-2015, 2015.

556 Castro, L., and Freeman, B. A.: Reactive oxygen species in human health and disease, *Nutrition*,
557 17, 161-165, 2001.

558 Chan, A. W. H., Kautzman, K. E., Chhabra, P. S., Surratt, J. D., Chan, M. N., Crouse, J. D.,
559 Kurten, A., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol
560 formation from photooxidation of naphthalene and alkylnaphthalenes: implications for oxidation
561 of intermediate volatility organic compounds (IVOCs), *Atmos. Chem. Phys.*, 9, 3049-3060,
562 2009.

563 Chan, A. W. H., Chan, M. N., Surratt, J. D., Chhabra, P. S., Loza, C. L., Crouse, J. D., Yee, L.
564 D., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Role of aldehyde chemistry and
565 NO_x concentrations in secondary organic aerosol formation, *Atmos. Chem. Phys.*,
566 10, 7169-7188, 10.5194/acp-10-7169-2010, 2010.

567 Charrier, J. G., and Anastasio, C.: On dithiothreitol (DTT) as a measure of oxidative potential for
568 ambient particles: evidence for the importance of soluble transition metals, *Atmos. Chem. Phys.*,
569 12, 9321-9333, 10.5194/acp-12-9321-2012, 2012.

570 Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.: Elemental analysis of chamber organic aerosol
571 using an aerodyne high-resolution aerosol mass spectrometer, *Atmos. Chem. Phys.*, 10, 4111-
572 4131, 10.5194/acp-10-4111-2010, 2010.

573 Chhabra, P. S., Ng, N. L., Canagaratna, M. R., Corrigan, A. L., Russell, L. M., Worsnop, D. R.,
574 Flagan, R. C., and Seinfeld, J. H.: Elemental composition and oxidation of chamber organic
575 aerosol, *Atmos. Chem. Phys.*, 11, 8827-8845, 10.5194/acp-11-8827-2011, 2011.

576 Cho, A. K., Sioutas, C., Miguel, A. H., Kumagai, Y., Schmitz, D. A., Singh, M., Eiguren-
577 Fernandez, A., and Froines, J. R.: Redox activity of airborne particulate matter at different sites

578 in the Los Angeles Basin, *Environmental Research*, 99, 40-47, 10.1016/j.envres.2005.01.003,
579 2005.

580 Cocker III, D. R., Mader, B. T., Kalberer, M., Flagan, R. C., and Seinfeld, J. H.: The effect of
581 water on gas-particle partitioning of secondary organic aerosol: II. m-xylene and 1,3,5-
582 trimethylbenzene photooxidation systems, *Atmos. Environ.*, 35, 6073-6085,
583 [http://dx.doi.org/10.1016/S1352-2310\(01\)00405-8](http://dx.doi.org/10.1016/S1352-2310(01)00405-8), 2001.

584 Conny, J. M., and Slater, J. F.: Black carbon and organic carbon in aerosol particles from crown
585 fires in the Canadian boreal forest, *Journal of Geophysical Research: Atmospheres*, 107, AAC 4-
586 1-AAC 4-12, 10.1029/2001JD001528, 2002.

587 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin,
588 M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-
589 Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, *Analytical Chemistry*,
590 78, 8281-8289, 10.1021/ac061249n, 2006.

591 Dockery, D. W., Pope, C. A., Xu, X., Spengler, J. D., Ware, J. H., Fay, M. E., Ferris, B. G., and
592 Speizer, F. E.: An Association between Air Pollution and Mortality in Six U.S. Cities, *New
593 England Journal of Medicine*, 329, 1753-1759, doi:10.1056/NEJM199312093292401, 1993.

594 Dou, J., Lin, P., Kuang, B.-Y., and Yu, J. Z.: Reactive Oxygen Species Production Mediated by
595 Humic-like Substances in Atmospheric Aerosols: Enhancement Effects by Pyridine, Imidazole,
596 and Their Derivatives, *Environmental Science & Technology*, 49, 6457-6465,
597 10.1021/es5059378, 2015.

598 Eddingsaas, N. C., Loza, C. L., Yee, L. D., Chan, M., Schilling, K. A., Chhabra, P. S., Seinfeld,
599 J. H., and Wennberg, P. O.: α -pinene photooxidation under controlled chemical conditions
600 – Part 2: SOA yield and composition in low- and high-NO_x environments, *Atmos. Chem.
601 Phys.*, 12, 7413-7427, 10.5194/acp-12-7413-2012, 2012.

602 Edney, E. O., Driscoll, D. J., Speer, R. E., Weathers, W. S., Kleindienst, T. E., Li, W., and
603 Smith, D. F.: Impact of aerosol liquid water on secondary organic aerosol yields of irradiated
604 toluene/propylene/NO_x/(NH₄)₂SO₄/air mixtures, *Atmos. Environ.*, 34, 3907-3919,
605 [http://dx.doi.org/10.1016/S1352-2310\(00\)00174-6](http://dx.doi.org/10.1016/S1352-2310(00)00174-6), 2000.

606 Fang, T., Guo, H., Verma, V., Peltier, R. E., and Weber, R. J.: PM_{2.5} water-soluble elements in
607 the southeastern United States: automated analytical method development, spatiotemporal
608 distributions, source apportionment, and implications for health studies, *Atmos. Chem. Phys.*, 15,
609 11667-11682, 10.5194/acp-15-11667-2015, 2015a.

610 Fang, T., Verma, V., Guo, H., King, L. E., Edgerton, E. S., and Weber, R. J.: A semi-automated
611 system for quantifying the oxidative potential of ambient particles in aqueous extracts using the
612 dithiothreitol (DTT) assay: results from the Southeastern Center for Air Pollution and
613 Epidemiology (SCAPE), *Atmos. Meas. Tech.*, 8, 471-482, 10.5194/amt-8-471-2015, 2015b.

614 Fang, T., Verma, V., Bates, J. T., Abrams, J., Klein, M., Strickland, M. J., Sarnat, S. E., Chang,
615 H. H., Mulholland, J. A., Tolbert, P. E., Russell, A. G., and Weber, R. J.: Oxidative potential of
616 ambient water-soluble PM_{2.5} in the southeastern United States: contrasts in sources and health
617 associations between ascorbic acid (AA) and dithiothreitol (DTT) assays, *Atmos. Chem. Phys.*,
618 16, 3865-3879, 10.5194/acp-16-3865-2016, 2016.

619 Farmer, D. K., Matsunaga, A., Docherty, K. S., Surratt, J. D., Seinfeld, J. H., Ziemann, P. J., and
620 Jimenez, J. L.: Response of an aerosol mass spectrometer to organonitrates and organosulfates
621 and implications for atmospheric chemistry, *Proceedings of the National Academy of Sciences*,
622 107, 6670-6675, 10.1073/pnas.0912340107, 2010.

623 Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H., Dubé,
624 W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H. P., Brauers, T., and Cohen, R. C.: Organic
625 nitrate and secondary organic aerosol yield from NO₃ oxidation of β-pinene
626 evaluated using a gas-phase kinetics/aerosol partitioning model, *Atmos. Chem. Phys.*, 9, 1431-
627 1449, 10.5194/acp-9-1431-2009, 2009.

628 Goldstein, A. H., and Galbally, I. E.: Known and Unexplored Organic Constituents in the Earth's
629 Atmosphere, *Environmental Science & Technology*, 41, 1514-1521, 10.1021/es072476p, 2007.

630 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of
631 global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols
632 from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.

633 Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and
634 monoterpene emission rate variability: Model evaluations and sensitivity analyses, *Journal of*
635 *Geophysical Research: Atmospheres*, 98, 12609-12617, 10.1029/93JD00527, 1993.

636 Gurgueira, S. A., Lawrence, J., Coull, B., Murthy, G. G. K., and Gonzalez-Flecha, B.: Rapid
637 increases in the steady-state concentration of reactive oxygen species in the lungs and heart after
638 particulate air pollution inhalation, *Environmental Health Perspectives*, 110, 749-755, 2002.

639 Hamad, S. H., Shafer, M. M., Kadhim, A. K. H., Al-Omran, S. M., and Schauer, J. J.: Seasonal
640 trends in the composition and ROS activity of fine particulate matter in Baghdad, Iraq, *Atmos.*
641 *Environ.*, 100, 102-110, <http://dx.doi.org/10.1016/j.atmosenv.2014.10.043>, 2015.

642 Heald, C. L., Kroll, J. H., Jimenez, J. L., Docherty, K. S., DeCarlo, P. F., Aiken, A. C., Chen, Q.,
643 Martin, S. T., Farmer, D. K., and Artaxo, P.: A simplified description of the evolution of organic
644 aerosol composition in the atmosphere, *Geophysical Research Letters*, 37, n/a-n/a,
645 10.1029/2010GL042737, 2010.

646 Healy, R. M., Temime, B., Kuprovskite, K., and Wenger, J. C.: Effect of Relative Humidity on
647 Gas/Particle Partitioning and Aerosol Mass Yield in the Photooxidation of p-Xylene,
648 *Environmental Science & Technology*, 43, 1884-1889, 10.1021/es802404z, 2009.

649 Helmig, D., Arey, J., Harger, W. P., Atkinson, R., and Lopez-Cancio, J.: Formation of mutagenic
650 nitrodibenzopyranones and their occurrence in ambient air, *Environmental Science &*
651 *Technology*, 26, 622-624, 10.1021/es00027a028, 1992.

652 Hensley, K., Robinson, K. A., Gabbita, S. P., Salsman, S., and Floyd, R. A.: Reactive oxygen
653 species, cell signaling, and cell injury, *Free Radical Biology and Medicine*, 28, 1456-1462,
654 [http://dx.doi.org/10.1016/S0891-5849\(00\)00252-5](http://dx.doi.org/10.1016/S0891-5849(00)00252-5), 2000.

655 Hoek, G., Krishnan, R. M., Beelen, R., Peters, A., Ostro, B., Brunekreef, B., and Kaufman, J. D.:
656 Long-term air pollution exposure and cardio-respiratory mortality: a review, *Environ Health*, 12,
657 43, 2013.

658 Hoffmann, T., Odum, J., Bowman, F., Collins, D., Klockow, D., Flagan, R., and Seinfeld, J.:
659 Formation of Organic Aerosols from the Oxidation of Biogenic Hydrocarbons, *Journal of*
660 *Atmospheric Chemistry*, 26, 189-222, 10.1023/A:1005734301837, 1997.

661 Janssen, N. A. H., Yang, A., Strak, M., Steenhof, M., Hellack, B., Gerlofs-Nijland, M. E.,
662 Kuhlbusch, T., Kelly, F., Harrison, R., Brunekreef, B., Hoek, G., and Cassee, F.: Oxidative
663 potential of particulate matter collected at sites with different source characteristics, *Sci. Total*
664 *Environ.*, 472, 572-581, <http://dx.doi.org/10.1016/j.scitotenv.2013.11.099>, 2014.

665 Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of
666 the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic
667 volatile organic compounds, *Atmos. Chem. Phys.*, 3, 181-193, 10.5194/acp-3-181-2003, 2003.

668 Jia, C., and Batterman, S.: A Critical Review of Naphthalene Sources and Exposures Relevant to
669 Indoor and Outdoor Air, *International Journal of Environmental Research and Public Health*, 7,
670 2903-2939, 10.3390/ijerph7072903, 2010.

671 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H.,
672 DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M.,
673 Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin,
674 C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M.,
675 Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, J., Huffman, J. A.,
676 Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F.,
677 Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A.,
678 Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J.
679 R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C.,
680 Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic
681 Aerosols in the Atmosphere, *Science*, 326, 1525-1529, 10.1126/science.1180353, 2009.

682 Kautzman, K. E., Surratt, J. D., Chan, M. N., Chan, A. W. H., Hersey, S. P., Chhabra, P. S.,
683 Dalleska, N. F., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of
684 Gas- and Aerosol-Phase Products from the Photooxidation of Naphthalene, *The Journal of*
685 *Physical Chemistry A*, 114, 913-934, 10.1021/jp908530s, 2010.

686 Kleinman, M. T., Hamade, A., Meacher, D., Oldham, M., Sioutas, C., Chakrabarti, B., Stram, D.,
687 Froines, J. R., and Cho, A. K.: Inhalation of concentrated ambient particulate matter near a
688 heavily trafficked road stimulates antigen-induced airway responses in mice, *Journal of the Air*
689 *& Waste Management Association*, 55, 1277-1288, 2005.

690 Kramer, A. J., Rattanavaraha, W., Zhang, Z., Gold, A., Surratt, J. D., and Lin, Y.-H.: Assessing
691 the oxidative potential of isoprene-derived epoxides and secondary organic aerosol, *Atmos.*
692 *Environ.*, <http://dx.doi.org/10.1016/j.atmosenv.2015.10.018>, 2016.

693 Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary organic
694 aerosol formation from isoprene photooxidation under high-NO_x conditions, *Geophysical*
695 *Research Letters*, 32, n/a-n/a, 10.1029/2005GL023637, 2005.

696 Kroll, J. H., and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and
697 evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593-3624,
698 10.1016/j.atmosenv.2008.01.003, 2008.

699 Kroll, J. H., Donahue, N. M., Jimenez, J. L., Kessler, S. H., Canagaratna, M. R., Wilson, K. R.,
700 Altieri, K. E., Mazzoleni, L. R., Wozniak, A. S., Bluhm, H., Mysak, E. R., Smith, J. D., Kolb, C.
701 E., and Worsnop, D. R.: Carbon oxidation state as a metric for describing the chemistry of
702 atmospheric organic aerosol, *Nat Chem*, 3, 133-139,
703 <http://www.nature.com/nchem/journal/v3/n2/abs/nchem.948.html#supplementary-information>,
704 2011.

705 Kumagai, Y., Koide, S., Taguchi, K., Endo, A., Nakai, Y., Yoshikawa, T., and Shimojo, N.:
706 Oxidation of proximal protein sulfhydryls by phenanthraquinone, a component of diesel exhaust
707 particles, *Chemical Research in Toxicology*, 15, 483-489, 10.1021/tx0100993, 2002.

708 Lambe, A. T., Onasch, T. B., Massoli, P., Croasdale, D. R., Wright, J. P., Ahern, A. T.,
709 Williams, L. R., Worsnop, D. R., Brune, W. H., and Davidovits, P.: Laboratory studies of the
710 chemical composition and cloud condensation nuclei (CCN) activity of secondary organic
711 aerosol (SOA) and oxidized primary organic aerosol (OPOA), *Atmos. Chem. Phys.*, 11, 8913-
712 8928, 10.5194/acp-11-8913-2011, 2011.

713 Li, N., Hao, M. Q., Phalen, R. F., Hinds, W. C., and Nel, A. E.: Particulate air pollutants and
714 asthma - A paradigm for the role of oxidative stress in PM-induced adverse health effects,
715 *Clinical Immunology*, 109, 250-265, 10.1016/j.clim.2003.08.006, 2003a.

716 Li, N., Sioutas, C., Cho, A., Schmitz, D., Misra, C., Sempf, J., Wang, M. Y., Oberley, T.,
717 Froines, J., and Nel, A.: Ultrafine particulate pollutants induce oxidative stress and mitochondrial
718 damage, *Environmental Health Perspectives*, 111, 455-460, 10.1289/ehp.6000, 2003b.

719 Li, N., Xia, T., and Nel, A. E.: The role of oxidative stress in ambient particulate matter-induced
720 lung diseases and its implications in the toxicity of engineered nanoparticles, *Free Radical*
721 *Biology and Medicine*, 44, 1689-1699, 10.1016/j.freeradbiomed.2008.01.028, 2008.

722 Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H., AlMazroa, M.
723 A., Amann, M., Anderson, H. R., Andrews, K. G., Aryee, M., Atkinson, C., Bacchus, L. J.,
724 Bahalim, A. N., Balakrishnan, K., Balmes, J., Barker-Collo, S., Baxter, A., Bell, M. L., Blore, J.
725 D., Blyth, F., Bonner, C., Borges, G., Bourne, R., Boussinesq, M., Brauer, M., Brooks, P., Bruce,
726 N. G., Brunekreef, B., Bryan-Hancock, C., Bucello, C., Buchbinder, R., Bull, F., Burnett, R. T.,
727 Byers, T. E., Calabria, B., Carapetis, J., Carnahan, E., Chafe, Z., Charlson, F., Chen, H., Chen, J.
728 S., Cheng, A. T.-A., Child, J. C., Cohen, A., Colson, K. E., Cowie, B. C., Darby, S., Darling, S.,
729 Davis, A., Degenhardt, L., Dentener, F., Des Jarlais, D. C., Devries, K., Dherani, M., Ding, E. L.,
730 Dorsey, E. R., Driscoll, T., Edmond, K., Ali, S. E., Engell, R. E., Erwin, P. J., Fahimi, S., Falder,
731 G., Farzadfar, F., Ferrari, A., Finucane, M. M., Flaxman, S., Fowkes, F. G. R., Freedman, G.,
732 Freeman, M. K., Gakidou, E., Ghosh, S., Giovannucci, E., Gmel, G., Graham, K., Grainger, R.,
733 Grant, B., Gunnell, D., Gutierrez, H. R., Hall, W., Hoek, H. W., Hogan, A., Hosgood, H. D., III,
734 Hoy, D., Hu, H., Hubbell, B. J., Hutchings, S. J., Ibeanusi, S. E., Jacklyn, G. L., Jasrasaria, R.,
735 Jonas, J. B., Kan, H., Kanis, J. A., Kassebaum, N., Kawakami, N., Khang, Y.-H., Khatibzadeh,
736 S., Khoo, J.-P., Kok, C., Laden, F., Lalloo, R., Lan, Q., Lathlean, T., Leasher, J. L., Leigh, J., Li,
737 Y., Lin, J. K., Lipshultz, S. E., London, S., Lozano, R., Lu, Y., Mak, J., Malekzadeh, R.,
738 Mallinger, L., Marcenes, W., March, L., Marks, R., Martin, R., McGale, P., McGrath, J., Mehta,
739 S., Memish, Z. A., Mensah, G. A., Merriman, T. R., Micha, R., Michaud, C., Mishra, V.,
740 Hanafiah, K. M., Mokdad, A. A., Morawska, L., Mozaffarian, D., Murphy, T., Naghavi, M.,
741 Neal, B., Nelson, P. K., Nolla, J. M., Norman, R., Olives, C., Omer, S. B., Orchard, J., Osborne,
742 R., Ostro, B., Page, A., Pandey, K. D., Parry, C. D. H., Passmore, E., Patra, J., Pearce, N.,

743 Pelizzari, P. M., Petzold, M., Phillips, M. R., Pope, D., Pope, C. A., III, Powles, J., Rao, M.,
744 Razavi, H., Rehfuess, E. A., Rehm, J. T., Ritz, B., Rivara, F. P., Roberts, T., Robinson, C.,
745 Rodriguez-Portales, J. A., Romieu, I., Room, R., Rosenfeld, L. C., Roy, A., Rushton, L.,
746 Salomon, J. A., Sampson, U., Sanchez-Riera, L., Sanman, E., Sapkota, A., Seedat, S., Shi, P.,
747 Shield, K., Shivakoti, R., Singh, G. M., Sleet, D. A., Smith, E., Smith, K. R., Stapelberg, N. J.
748 C., Steenland, K., Stöckl, H., Stovner, L. J., Straif, K., Straney, L., Thurston, G. D., Tran, J. H.,
749 Van Dingenen, R., van Donkelaar, A., Veerman, J. L., Vijayakumar, L., Weintraub, R.,
750 Weissman, M. M., White, R. A., Whiteford, H., Wiersma, S. T., Wilkinson, J. D., Williams, H.
751 C., Williams, W., Wilson, N., Woolf, A. D., Yip, P., Zielinski, J. M., Lopez, A. D., Murray, C. J.
752 L., and Ezzati, M.: A comparative risk assessment of burden of disease and injury attributable to
753 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the
754 Global Burden of Disease Study 2010, *The Lancet*, 380, 2224-2260, 10.1016/S0140-
755 6736(12)61766-8, 2012.

756 Lin, P., and Yu, J. Z.: Generation of Reactive Oxygen Species Mediated by Humic-like
757 Substances in Atmospheric Aerosols, *Environmental Science & Technology*, 45, 10362-10368,
758 10.1021/es2028229, 2011.

759 Lin, Y.-H., Zhang, Z., Docherty, K. S., Zhang, H., Budisulistiorini, S. H., Rubitschun, C. L.,
760 Shaw, S. L., Knipping, E. M., Edgerton, E. S., Kleindienst, T. E., Gold, A., and Surratt, J. D.:
761 Isoprene Epoxydiols as Precursors to Secondary Organic Aerosol Formation: Acid-Catalyzed
762 Reactive Uptake Studies with Authentic Compounds, *Environmental science & technology*, 46,
763 250-258, 10.1021/es202554c, 2012.

764 Loza, C. L., Craven, J. S., Yee, L. D., Coggon, M. M., Schwantes, R. H., Shiraiwa, M., Zhang,
765 X., Schilling, K. A., Ng, N. L., Canagaratna, M. R., Ziemann, P. J., Flagan, R. C., and Seinfeld,
766 J. H.: Secondary organic aerosol yields of 12-carbon alkanes, *Atmos. Chem. Phys.*, 14, 1423-
767 1439, 10.5194/acp-14-1423-2014, 2014.

768 Lu, Y., Su, S., Jin, W., Wang, B., Li, N., Shen, H., Li, W., Huang, Y., Chen, H., Zhang, Y.,
769 Chen, Y., Lin, N., Wang, X., and Tao, S.: Characteristics and cellular effects of ambient
770 particulate matter from Beijing, *Environmental Pollution*, 191, 63-69,
771 <http://dx.doi.org/10.1016/j.envpol.2014.04.008>, 2014.

772 Lund, A. K., Doyle-Eisele, M., Lin, Y. H., Arashiro, M., Surratt, J. D., Holmes, T., Schilling, K.
773 A., Seinfeld, J. H., Rohr, A. C., Knipping, E. M., and McDonald, J. D.: The effects of alpha-
774 pinene versus toluene-derived secondary organic aerosol exposure on the expression of markers
775 associated with vascular disease, *Inhal. Toxicol.*, 25, 309-324, 10.3109/08958378.2013.782080,
776 2013.

777 McDonald, J. D., Doyle-Eisele, M., Campen, M. J., Seagrave, J., Holmes, T., Lund, A., Surratt,
778 J. D., Seinfeld, J. H., Rohr, A. C., and Knipping, E. M.: Cardiopulmonary response to inhalation
779 of biogenic secondary organic aerosol, *Inhal. Toxicol.*, 22, 253-265,
780 10.3109/08958370903148114, 2010.

781 McDonald, J. D., Doyle-Eisele, M., Kracko, D., Lund, A., Surratt, J. D., Hersey, S. P., Seinfeld,
782 J. H., Rohr, A. C., and Knipping, E. M.: Cardiopulmonary response to inhalation of secondary
783 organic aerosol derived from gas-phase oxidation of toluene, *Inhal. Toxicol.*, 24, 689-697,
784 10.3109/08958378.2012.712164, 2012.

785 McLafferty, F. W., and Tureček, F.: Interpretation of mass spectra, University science books,
786 1993.

787 McWhinney, R. D., Badali, K., Liggio, J., Li, S.-M., and Abbatt, J. P. D.: Filterable Redox
788 Cycling Activity: A Comparison between Diesel Exhaust Particles and Secondary Organic
789 Aerosol Constituents, *Environmental Science & Technology*, 47, 3362-3369,
790 10.1021/es304676x, 2013a.

791 McWhinney, R. D., Zhou, S., and Abbatt, J. P. D.: Naphthalene SOA: redox activity and
792 naphthoquinone gas-particle partitioning, *Atmos. Chem. Phys.*, 13, 9731-9744, 10.5194/acp-13-
793 9731-2013, 2013b.

794 Nah, T., McVay, R. C., Pierce, J. R., Seinfeld, J. H., and Ng, N. L.: Constraining uncertainties in
795 particle wall-deposition correction during SOA formation in chamber experiments, *Atmos.*
796 *Chem. Phys. Discuss.*, 2016, 1-35, 10.5194/acp-2016-820, 2016.

797 Ng, N. L., Chhabra, P. S., Chan, A. W. H., Surratt, J. D., Kroll, J. H., Kwan, A. J., McCabe, D.
798 C., Wennberg, P. O., Sorooshian, A., Murphy, S. M., Dalleska, N. F., Flagan, R. C., and
799 Seinfeld, J. H.: Effect of NO_x level on secondary organic aerosol (SOA) formation from the
800 photooxidation of terpenes, *Atmos. Chem. Phys.*, 7, 5159-5174, 10.5194/acp-7-5159-2007,
801 2007a.

802 Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.:
803 Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem. Phys.*,
804 7, 3909-3922, 10.5194/acp-7-3909-2007, 2007b.

805 Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H.,
806 Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L.,
807 Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and
808 Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from
809 Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625-4641, 10.5194/acp-10-4625-2010,
810 2010.

811 Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Chhabra, P. S., Seinfeld, J. H., and Worsnop, D.
812 R.: Changes in organic aerosol composition with aging inferred from aerosol mass spectra,
813 *Atmos. Chem. Phys.*, 11, 6465-6474, 10.5194/acp-11-6465-2011, 2011.

814 Nguyen, T. B., Roach, P. J., Laskin, J., Laskin, A., and Nizkorodov, S. A.: Effect of humidity on
815 the composition of isoprene photooxidation secondary organic aerosol, *Atmos. Chem. Phys.*, 11,
816 6931-6944, 10.5194/acp-11-6931-2011, 2011.

817 Philip, M., Rowley, D. A., and Schreiber, H.: Inflammation as a tumor promoter in cancer
818 induction, *Seminars in Cancer Biology*, 14, 433-439,
819 <http://dx.doi.org/10.1016/j.semcancer.2004.06.006>, 2004.

820 Piccot, S. D., Watson, J. J., and Jones, J. W.: A global inventory of volatile organic compound
821 emissions from anthropogenic sources, *Journal of Geophysical Research: Atmospheres*, 97,
822 9897-9912, 10.1029/92JD00682, 1992.

823 Platt, S. M., Haddad, I. E., Pieber, S. M., Huang, R. J., Zardini, A. A., Clairotte, M., Suarez-
824 Bertoa, R., Barmet, P., Pfaffenberger, L., Wolf, R., Slowik, J. G., Fuller, S. J., Kalberer, M.,
825 Chirico, R., Dommen, J., Astorga, C., Zimmermann, R., Marchand, N., Hellebust, S., Temime-

826 Roussel, B., Baltensperger, U., and Prévôt, A. S. H.: Two-stroke scooters are a dominant source
827 of air pollution in many cities, *Nature Communications*, 5, 3749, 10.1038/ncomms4749
828 <http://www.nature.com/articles/ncomms4749#supplementary-information>, 2014.

829 Pope, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and Thurston, G. D.:
830 Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution,
831 *Jama-Journal of the American Medical Association*, 287, 1132-1141, 10.1001/jama.287.9.1132,
832 2002.

833 Pope III, C. A., and Dockery, D. W.: Health effects of fine particulate air pollution: Lines that
834 connect, *Journal of the Air and Waste Management Association*, 56, 709-742, 2006.

835 Rattanavaraha, W., Rosen, E., Zhang, H., Li, Q., Pantong, K., and Kamens, R. M.: The reactive
836 oxidant potential of different types of aged atmospheric particles: An outdoor chamber study,
837 *Atmos. Environ.*, 45, 3848-3855, <http://dx.doi.org/10.1016/j.atmosenv.2011.04.002>, 2011.

838 Riva, M., Robinson, E. S., Perraudin, E., Donahue, N. M., and Villenave, E.: Photochemical
839 Aging of Secondary Organic Aerosols Generated from the Photooxidation of Polycyclic
840 Aromatic Hydrocarbons in the Gas-Phase, *Environmental Science & Technology*, 49, 5407-
841 5416, 10.1021/acs.est.5b00442, 2015.

842 Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop,
843 A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking Organic Aerosols: Semivolatile
844 Emissions and Photochemical Aging, *Science*, 315, 1259-1262, 10.1126/science.1133061, 2007.

845 Sannigrahi, P., Sullivan, A. P., Weber, R. J., and Ingall, E. D.: Characterization of Water-Soluble
846 Organic Carbon in Urban Atmospheric Aerosols Using Solid-State ¹³C NMR Spectroscopy,
847 *Environmental Science & Technology*, 40, 666-672, 10.1021/es051150i, 2006.

848 Sato, K., Takami, A., Isezaki, T., Hikida, T., Shimono, A., and Imamura, T.: Mass spectrometric
849 study of secondary organic aerosol formed from the photo-oxidation of aromatic hydrocarbons,
850 *Atmos. Environ.*, 44, 1080-1087, <http://dx.doi.org/10.1016/j.atmosenv.2009.12.013>, 2010.

851 Stirnweis, L., Marcolli, C., Dommen, J., Barmet, P., Frege, C., Platt, S. M., Bruns, E. A., Krapf,
852 M., Slowik, J. G., Wolf, R., Prévôt, A. S. H., El-Haddad, I., and Baltensperger, U.: α -Pinene
853 secondary organic aerosol yields increase at higher relative humidity and low NO_x conditions,
854 *Atmos. Chem. Phys. Discuss.*, 2016, 1-41, 10.5194/acp-2016-717, 2016.

855 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S.
856 P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in
857 secondary organic aerosol formation from isoprene, *Proceedings of the National Academy of*
858 *Sciences*, 107, 6640-6645, 10.1073/pnas.091114107, 2010.

859 Tao, F., Gonzalez-Flecha, B., and Kobzik, L.: Reactive oxygen species in pulmonary
860 inflammation by ambient particulates, *Free Radical Biology and Medicine*, 35, 327-340,
861 [http://dx.doi.org/10.1016/S0891-5849\(03\)00280-6](http://dx.doi.org/10.1016/S0891-5849(03)00280-6), 2003.

862 Tasoglou, A., and Pandis, S. N.: Formation and chemical aging of secondary organic aerosol
863 during the β -caryophyllene oxidation, *Atmos. Chem. Phys.*, 15, 6035-6046, 10.5194/acp-15-
864 6035-2015, 2015.

865 Tuet, W. Y., Fok, S., Verma, V., Tagle Rodriguez, M. S., Grosberg, A., Champion, J. A., and
866 Ng, N. L.: Dose-dependent intracellular reactive oxygen and nitrogen species production from

867 particulate matter exposure: comparison to oxidative potential and chemical composition, *Atmos.*
868 *Environ.*, 10.1016/j.atmosenv.2016.09.005, 2016.

869 Turner, J., Hernandez, M., Snawder, J. E., Handorean, A., and McCabe, K. M.: A Toxicology
870 Suite Adapted for Comparing Parallel Toxicity Responses of Model Human Lung Cells to Diesel
871 Exhaust Particles and Their Extracts, *Aerosol Sci. Technol.*, 49, 599-610,
872 10.1080/02786826.2015.1053559, 2015.

873 Van Krevelen, D.: Graphical-statistical method for the study of structure and reaction processes
874 of coal, *Fuel*, 29, 269-284, 1950.

875 Verma, V., Rico-Martinez, R., Kotra, N., King, L., Liu, J. M., Snell, T. W., and Weber, R. J.:
876 Contribution of Water-Soluble and Insoluble Components and Their Hydrophobic/Hydrophilic
877 Subfractions to the Reactive Oxygen Species-Generating Potential of Fine Ambient Aerosols,
878 *Environmental Science & Technology*, 46, 11384-11392, 10.1021/es302484r, 2012.

879 Verma, V., Fang, T., Guo, H., King, L., Bates, J. T., Peltier, R. E., Edgerton, E., Russell, A. G.,
880 and Weber, R. J.: Reactive oxygen species associated with water-soluble PM 2.5 in the
881 southeastern United States: spatiotemporal trends and source apportionment, *Atmos. Chem.*
882 *Phys.*, 14, 12915-12930, 2014.

883 Verma, V., Fang, T., Xu, L., Peltier, R. E., Russell, A. G., Ng, N. L., and Weber, R. J.: Organic
884 Aerosols Associated with the Generation of Reactive Oxygen Species (ROS) by Water-Soluble
885 PM2.5, *Environmental Science & Technology*, 49, 4646-4656, 10.1021/es505577w, 2015a.

886 Verma, V., Wang, Y., El-Afifi, R., Fang, T., Rowland, J., Russell, A. G., and Weber, R. J.:
887 Fractionating ambient humic-like substances (HULIS) for their reactive oxygen species activity
888 – Assessing the importance of quinones and atmospheric aging, *Atmos. Environ.*, 120, 351-359,
889 <http://dx.doi.org/10.1016/j.atmosenv.2015.09.010>, 2015b.

890 Visentin, M., Pagnoni, A., Sarti, E., and Pietrogrande, M. C.: Urban PM2.5 oxidative potential:
891 Importance of chemical species and comparison of two spectrophotometric cell-free assays,
892 *Environmental Pollution*, 219, 72-79, <http://dx.doi.org/10.1016/j.envpol.2016.09.047>, 2016.

893 Vivanco, M. G., and Santiago, M.: Secondary Organic Aerosol Formation from the Oxidation of
894 a Mixture of Organic Gases in a Chamber, *Air Quality*, Ashok Kumar (Ed.), InTech, DOI:
895 10.5772/9761. Available from: [http://www.intechopen.com/books/air-quality/secondary-organic-](http://www.intechopen.com/books/air-quality/secondary-organic-aerosols-experiments-in-an-outdoor-chamber-)
896 [aerosols-experiments-in-an-outdoor-chamber-](http://www.intechopen.com/books/air-quality/secondary-organic-aerosols-experiments-in-an-outdoor-chamber-), 2010.

897 Weichenthal, S. A., Lavigne, E., Evans, G. J., Godri Pollitt, K. J., and Burnett, R. T.: Fine
898 Particulate Matter and Emergency Room Visits for Respiratory Illness. Effect Modification by
899 Oxidative Potential, *American Journal of Respiratory and Critical Care Medicine*, 194, 577-586,
900 10.1164/rccm.201512-2434OC, 2016.

901 Wennberg, P.: Let's abandon the "high NOx" and "low NOx" terminology, *IGAC news*, 50, 3-4,
902 2013.

903 Wiseman, H., and Halliwell, B.: Damage to DNA by reactive oxygen and nitrogen species: role
904 in inflammatory disease and progression to cancer, *Biochem. J.*, 313, 17-29, 1996.

905 Wong, J. P. S., Lee, A. K. Y., and Abbatt, J. P. D.: Impacts of Sulfate Seed Acidity and Water
906 Content on Isoprene Secondary Organic Aerosol Formation, *Environmental Science &*
907 *Technology*, 49, 13215-13221, 10.1021/acs.est.5b02686, 2015.

908 Xu, L., Kollman, M. S., Song, C., Shilling, J. E., and Ng, N. L.: Effects of NO_x on the Volatility
909 of Secondary Organic Aerosol from Isoprene Photooxidation, *Environmental Science &*
910 *Technology*, 48, 2253-2262, 10.1021/es404842g, 2014.

911 Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-
912 VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H., Hering, S. V.,
913 de Gouw, J., Baumann, K., Lee, S.-H., Nenes, A., Weber, R. J., and Ng, N. L.: Effects of
914 anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the
915 southeastern United States, *Proceedings of the National Academy of Sciences*, 112, 37-42,
916 10.1073/pnas.1417609112, 2015a.

917 Xu, L., Suresh, S., Guo, H., Weber, R. J., and Ng, N. L.: Aerosol characterization over the
918 southeastern United States using high-resolution aerosol mass spectrometry: spatial and seasonal
919 variation of aerosol composition and sources with a focus on organic nitrates, *Atmos. Chem.*
920 *Phys.*, 15, 7307-7336, 10.5194/acp-15-7307-2015, 2015b.

921 Yang, A., Janssen, N. A. H., Brunekreef, B., Cassee, F. R., Hoek, G., and Gehring, U.: Children's
922 respiratory health and oxidative potential of PM_{2.5}: the PIAMA birth cohort study, *Occupational*
923 *and Environmental Medicine*, 10.1136/oemed-2015-103175, 2016.

924 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R.,
925 Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P.
926 F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimonono, A., Hatakeyama, S., Takegawa,
927 N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams,
928 P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M.,
929 and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in
930 anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophysical Research Letters*,
931 34, n/a-n/a, 10.1029/2007GL029979, 2007.

932