

Responses to reviewers of the paper by

Klimont et al. (<http://www.atmos-chem-phys-discuss.net/acp-2016-880/>)

RESPONSES TO Reviewer 1:

We thank the reviewer for a very thorough and insightful review which we have used to improve the manuscript and provide additional results in further extended Supporting Information set. The responses to specific points raised by the reviewer are provided below.

REVIEWER:

The calculations often incorporate substantial technology and emission control details, however the results of this detail are only presented at a fairly aggregate level. Some key intermediate results here would be very useful to present. In particular, for some key sector/fuel combinations, I suggest that the emission factors over time for different regions (perhaps PM or BC in the main paper, and other species in the supplement). One key sector is onroad diesel, for example, where aggregate emission factors have changed over time in many regions. Other key sectors might be residential biomass, off-road diesel, etc. Where these emission factors are largely constant, this could be mentioned in the text without a figure, but where these have changed some figures and/or tables and some discussion would be useful

RESPONSE: Indeed, the model considers explicitly implementation of particular technologies to achieve required emission standards. We produced now a specific output for selected sectors (on-road diesel, off-road diesel, power –coal, industry-coal, residential biomass) and all key regions. On the basis this output we present a few examples where changes were relevant within the characterized period.

REVIEWER:

The main presentation focuses on emissions by region, and global emissions by sector. There is a lot of material here, and this is a reasonable choice for the main paper, however many readers will want to see emissions by sector for specific regions. It would be useful to present the equivalent of Figure 3 for the different regions in the supplement. (Hopefully the codes that generated that figure can readily be generalized.). As noted below the authors should also provide an electronic supplementary file with more detailed emissions. Some further suggestions for details that would be useful to supply are below.

RESPONSE: We added now two additional outputs:

- The set of tables presenting emissions for each of the 25 IMAGE regions by key sectors (power-coal, power-other fuels, industry, coke ovens, fossil fuel production, residential – biomass, res –coal, res – other fuels, road transport –diesel, road transport – other fuels, non-road – diesel, non-road – other fuels, other). This is an Excel file that will be part of the SI.

- A set of figures for key regions (as in Figure 3) with emissions for PM species over time as in Figure 4. We believe that the reviewer meant Figure 4 for key regions rather than Figure 3 since the latter is already regional. These additional figures are in the SI.

REVIEWER:

Overall, its not clear how equipment vintages are treated. There is a mention of old/new power plants, but not as much discussion for other sectors. For vehicles, for example, is the model based on aggregate emission factors by year, or are vintages of vehicles tracked over time? This should be clarified in the manuscript.

RESPONSE: The equipment vintages are dealt in GAINS in two ways. Explicit assumptions for power sector where activity (energy use) for existing and new (post 2000) power plants are included, and specifying pace of equipment replacement, using technical lifetime of technologies in GAINS database which is defined for each control technology, in the so called ‘control strategies’ where share of fuel use for a given technology is given. While for many sectors, the add-on control technologies can be applied at any time and vintage plays smaller role, for transport new standards most of the time are synonymous with a new vintage year of a particular vehicle. Vehicle lifetimes are region specific and affect the fleet turnover which in turn determines how quickly a new technology penetrates the market. As for vehicle emission factors, in GAINS they represent for each of the categories, e.g., EURO 1, EURO 2 or any other standard, the average lifetime emission rate, including typical deterioration factors. We add additional text in the manuscript to explain how vintages of installations and technologies are dealt with – see section 2.2.2 and 2.2.3.

REVIEWER:

Specific comments Section: 2.2.1 Residential combustion: cooking, heating, lighting: It is not clear if assumptions such as the use fraction of different technologies and also splits between end-use services (e.g. cooking, heating in particular) are constant over time within a given country or region?

RESPONSE: There is an explicit region-specific assumption in the model about the fuel use for cooking and heating. For Europe and North America solid fuels in residential sector are allocated to heating unless specific information exist, e.g., Switzerland provided their estimates for cooking. We are aware that there is some cooking (or cooking and heating using the same devices) in several countries but there is no data and the cooking share is most likely very small compared to demand for heating. For Asia, Africa, and Latin America cooking dominates although for some countries like China and few of the Latin American countries heating plays an important role; similarly in some provinces (states) in India, Pakistan, or Nepal. The data for that originates from GAINS-Asia related studies as well as the recent work Latin America with support of CCAC and UNEP (final report in preparation for publication; see summary for policy makers <http://www.ccacoalition.org/en/resources/integrated-assessment-short-lived-climate-pollutants-latin-america-and-caribbean-summary>). The share of cooking/heating is assumed constant in the 1990-2010 period as we have not found any data allowing to change that

assumption. As far as technology split is concerned; there is no firm statistical data but for several countries our exchange with national experts led to adjustment of assumptions how the technology (stoves/boilers/automatic boilers/pellet stoves/fireplaces) shares changed over time. Again, there is no model behind but an attempt to reflect on available information from local studies or expert contacts. For example sales statistics, for example for pellet stoves and boilers, resulted in adjustment of shares of biomass used in such installations in several European countries where strong growth has been observed towards the end of the period under investigation and continues into the future (another paper). Also for China, trends towards cleaner coal stoves and more household coal boilers (in specific provinces) were taken into account. Finally, residential use of kerosene was split into cooking and lighting and the shares change over time depending on the regional access to electricity, as described in the manuscript.

REVIEWER:

Specific comments Section: 2.2.1 Residential combustion: cooking, heating, lighting: There is quite a bit of good work here and some sort of summary later on in terms of the evolution of aggregate emission factors (either over time – if the above assumptions change over the historical period) would be useful. For example, do PM emissions per unit biomass fuel use for cooking change over time in a given region? (Some of this could be in the supplement, with main points summarized in the text). While there are too many details to do this comprehensively, it would be important to summarize where emission factors for important sectors change over time (either as a result of different technology fractions, or emission controls). This is clearly going to be the case for diesel fuel use in road transportation in many regions, but how much change was assumed for the various regions. How about off road, agriculture, or residential sectors? Did controls or technology mix (beyond shifts in type of fuel use) have a noticeable impact on emission factors in these regions?

RESPONSE: As already mentioned in one of the earlier points, we are developing implied emission factors for a number of sectors where changes in structure of the sector or increased penetration of control technology made an impact. Examples include heavy duty trucks where we estimate for BC implied emissions factor declined globally by nearly 20% (2010 to 1990) but in several regions like North America, Western Europe, Japan the reduction was over about 60-65%, Central Europe about 40-50%, but for Russia less than 10% and for most other regions no significant change was estimated (for several regions like China the impact of legislation is visible only in 2015 and later). Another example includes coal power plants where globally emission factors for PM2.5 dropped by about 45% with NA, Western Europe, Japan having over 80% decline and even for China we estimated over 70% reduction while in Russia and several FSU countries only 20-30% decline. For industrial coal use lower reductions were achieved with exception of Eastern Europe and some FSU where collapse of heavy industry in the period 1990-2000 resulted in decline of emission factors by over 90% compared to the 1990 period. Finally, for residential heating (fuelwood) the 'global average factor' declined by about 15% which is mostly due to moderate changes in North America, Japan but mostly Western Europe where nearly 40% decline was estimated owing to strong increase in biomass use but in new installations including hundred thousands

of pellet stoves and boilers. We will include brief discussion of these trends and illustrate it on a chart in the manuscript – see sections 221, 222,223,224 and figures 1,2,3,

REVIEWER:

Pg 14 - "the independent fuel estimate by Denier van der Gon et al. (2015)". van der Gon et al. state "A consistent set of PM10, PM2.5 and PM1 emission data for Europe was obtained from the GAINS model...", so these do not appear to be independent

RESPONSE: We need to point out that our statement refers to the 'independent fuel use' estimates by van der Gon et al. Indeed, they relied on GAINS emission factors but estimated fuelwood consumption independently rather than relying on statistical data. Therefore, we believe that this is an accurate statement.

REVIEWER:

pg 16 - "The resulting fuel use was compared and calibrated to the diesel consumption reported in the power and commercial sector." Not clear what this means.

RESPONSE: For DG sets, fuel use is estimated from the number and size of DG sets in some regions (i. e. Nepal, Nigeria). For some regions, share of diesel used in DG sets as a percentage of total diesel consumption in the country/region is available (i. e. India). Once the total diesel consumption by country/region is estimated then the fuel use for DG sets in GAINS model is allocated from commercial and power sector. We have revised the statement in the manuscript that reads now: *"The resulting fuel use was compared to the IEA statistics for the power and commercial sector and adjusted if necessary so that the overall energy use is consistent with the IEA."*

REVIEWER:

pg 17, last portion of diesel generator section. There is some discussion of emission reduction options, but no mention the extent to which these were assumed to be applied in the emission estimates.

RESPONSE: Currently we have not applied any of the post-combustion control measures for DG sets in the developing world where they matter most; at least regionally. We modify that section and add explicit statement that no post-combustion measures are implemented for developing countries in this period.

REVIEWER:

pg 17 - line 25. IEA energy statistics contain separate lines for agriculture and construction - while these are not available for all years, these data seem to be becoming more complete in

more recent years, even for non-OECD countries. It appears that this data is not used? Some further explanation would be useful.

RESPONSE: In fact we do make use of this information but this (IEA statistics) data is for total diesel consumption in agriculture or industry and not specifically in mobile machinery. For example, in agriculture diesel is used also for irrigation purposes (we attempt to allocate this to diesel generation sets in GAINS as the operating conditions are similar) but also in heating boilers used for example for drying or heating. The text in the manuscript does specifically refer to mobile machinery information and therefore we believe it is a valid statement.

REVIEWER:

pg 17 - line 31 "Also old and often poorly maintained vehicle fleet is reflected in measurements of emission factors" - not clear what is meant here.

RESPONSE: We agree, this statement is unclear and we replace it with: "*For all world regions we assume that a certain fraction of vehicles is badly maintained (e.g., Mancilla et al., 2012) or their emission controls tampered which is reflected as the share of so-called high-emitters (McClintock, 1999, 2007; Smit and Bluett, 2011; Yan et al., 2011, 2014); see further discussion in section 3.4.1.*"

REVIEWER:

pg 20 line 14. This "fuel consumption data for 2007 were extrapolated to 2010 using GDP". The result of this assumption should be compared to the fuel consumption estimates from IMO, who compared both bottom up and top down methodologies. ("Third IMO Greenhouse Gas Study 2014")

RESPONSE: We compared the result of our extrapolation for 2010 (1056 Mt CO₂ (13.83 EJ)) with the data in IMO 3rd GHG Study 2014 showing: [Table 1]

- For 2010: Total shipping: 915 mio t CO₂ / International shipping: 771 mio t CO₂
- Average 2007-2012: 1015 mio t CO₂ / 846 mio t CO₂

Our extrapolation is slightly higher than the average for the period 2007-2012 and about 10% higher than the reported fuel use in 2010 by IMO. We have added a remark and reference to IMO in the manuscript.

REVIEWER:

Page 23 top regarding coke ovens. It is not quite clear what the technology representation is here. "uncontrolled ovens" are mentioned, is the split between controlled and uncontrolled? Is this assumed to change over time? It appears there is little detail in terms of emission factors available in the literature (and emissions seem likely to depend on site specific characteristics in any event), so some comment might be useful on what are the most important data that would be needed to improve estimates.

RESPONSE: Indeed, coke industry PM emissions are poorly understood and this is of special importance for China in the last decade. We highlight the poor data availability with respect to measurements and add at the end a sentence about the assumption regarding the change in emission factor over time, i.e., *“Owing to lack of specific data for various world regions, we assume little change in emissions factors over time for the developing world, although the transition in China reported in Huo et al. (2012) was considered, and for OECD countries emission factor trend follows reported emissions, where available.”*

REVIEWER:

Page 26. Might be useful to also mention that Agricultural waste burning can also be seasonally concentrated, so that it might be particularly important in some months.

RESPONSE: We have rewritten this part which now reads: *“At the same time, for several regions this source might be even more important, e.g., for Brazil we estimate its contribution at up to 15% of PM_{2.5} and 10% of BC emissions. Finally, agricultural burning has a strong seasonal pattern (see also section 2.4.1.) and has been also linked with heavy smog and haze episodes (e.g., Mukai et al., 2015; Stohl et al., 2007).”*

REVIEWER:

Page 26 line 12. "This database has been further extended and updated" perhaps edit to clarify that this refers to the data presented in the current paper.

RESPONSE: We have made that explicit now by joining this sentence and the next one that reads now: *“Niemi (2007) compared various datasets for all open biomass sources and developed the first global activity set for the RAINS model drawing on EDGAR3.2FT2000 (Van Aardenne et al., 2005) which we have further extended and updated to accommodate other data sources allowing gaps to be filled for several countries.”*

REVIEWER:

Page 27 "3.9 Other sources" Is dust included? Its not mentioned until the discussion section

RESPONSE: We have added a sentence at the end of the first paragraph of this section: *“Note that windblown dust and emissions from unpaved roads are not included (see also introduction to section 3).”* As indicated here, there are few words of explanation about what is included and what not in the introductory section of the section 3 of the manuscript.

REVIEWER:

Page 28 line 13 - " for barbeques, a per capita emission factor is established, i.e.,...". Presumably this varies by region?

RESPONSE: Unfortunately, we found very little data allowing to differentiate between the countries and therefore for most countries the same factor is used, except for few countries in Europe where national experts contributed their input leading to adjustment. We added a comment in the text reflecting that.

REVIEWER:

Page 32 line 19-20. Probably useful note here that the emissions in Granier et al., past about 2000/2005 were based on projections, where as the estimates here up to 2010 are (to the extent available) based on reported data and practices.

RESPONSE: Indeed, the numbers reported for 2010 represent various projections in Granier et al; in fact including also RCP numbers; this justifies also the widening emission range shown towards the end of the period they have investigated. We add a comment about it at the end of the line the reviewer referred to, which now reads: "...stabilization shown in earlier studies; *note that values reported in Granier et al. (2011) for 2010 were results of projections.*"

REVIEWER:

Page 35 line 3, this presumably is a typo? "confidence intervals to be 160-500% for the developing countries"

RESPONSE: As a matter of fact it is not a typo; see table 8 and discussion on page 18 of Streets et al (2003) manuscript. However, we choose to delete this statement from the paper as in view of the newer work (Bond et al, 2004, 2013), quoted already, where also a similar emission methodology was used as here, this particular statement does not add any specific insight unless elaborated further discussing specifically reasons for very high uncertainties estimated in that study.

REVIEWER:

Page 35 while "error compensation" is, indeed important, it might be useful to note that this might only be partial compensation. (e.g., some errors, such as measurement or enforcement issues, could be correlated across sectors.)

RESPONSE: Indeed, the poor enforcement might be one of the factors that cuts across the sector while for measurements of emission factors it could be also the case, often measurement techniques and teams performing them will be different reducing such potential. Our statement in the paper highlights the fact that the error compensation 'works' when errors are not correlated and so it is either known or assumed based on well-founded knowledge. We have added a respective comment in the paper and now the whole statement reads: "*Additionally, the error compensation, which is especially relevant if calculated emissions are the sum of a large number of equally important source categories (and where the errors in input parameters are not correlated with each other), can lead to a further reduction of*

overall emission uncertainty (Schöpp et al., 2005). A careful assessment of the assumption about correlation between input parameters is essential as for example poor enforcement of legislation or measurements errors could affect several source sectors in a similar way."

REVIEWER:

Wasn't sure about the meaning of this sentence: "In fact, they could be even lower considering that they typically rely on a harmonized data set and include a simultaneous calculation of emissions of several species using the same principal activity and technology data." I would presume most country-level inventories are similar in this respect?

RESPONSE: Indeed, this statement is more relevant for the multipollutant inventory (whole GAINS framework) rather than PM alone. We delete this sentence in the final manuscript even though from our experience working with several regional and national inventories we see that often the methods applied to for example PM10 or PM2.5 are not the same as for BC or OC. The latter are often derived using simply shares of PM2.5 rather than absolute emissions factors representative for a given technology. Such shares are often derived from a large set of measurements representing a category of installations rather than a specific one for which PM2.5 emissions were calculated. In that way additional uncertainty is introduced.

REVIEWER:

Would be useful to mention in this section that there is also uncertainty in the speciation fractions (but this is constrained across species since these must sum to ≤ 1).

RESPONSE: Thank you for pointing it out, we add a sentence highlighting this point on page 35, line 9 [in the original submission] and it reads: "*Allocating total PM emissions into different size bins or chemical species (here BC and OC) is associated with uncertainties that for a specific source are determined by the measurement. Among others, Bond et al. (2013) discussed specific issues related to BC and OC aerosols, while for PM size distribution there exist specific analysis for particular measurement equipment (e.g., Armas et al., 2007; Coquelin et al., 2013) and most of the studies reporting measurements of size distribution estimate uncertainties for each size category. While the sum of all the PM species is constrained by the total mass, the single size distribution values rely on a large number of measurements reducing the overall uncertainty. Exceptions are source-sectors for which very few measurements exist, e.g., coke ovens, fireworks, handling of bulk materials."*

REVIEWER:

Page 36 In addition to this "Our new global estimate of BC emissions suggests higher numbers than previously published...", perhaps useful to also mention something about BC trends over time here (since there is substantial interest in BC, and it looks like BC trends can be different from PM trends).

RESPONSE: More detailed discussion of this is actually provided in section 4.1 on page 30 of the original manuscript. Here, in conclusions we have added a statement about the different PM2.5 and PM10 trends vs BC and slightly reformulated the concerned paragraph (originally p.36, from line 10) and it reads now: *“We estimate that global emissions of PM have not changed much between 1990 and 2010 but there are significantly different regional trends with North America, Pacific, and Europe reducing emissions by 30 to over 50% and Asia and Africa increasing by about 30%. While these regionally varying developments are clearly visible in PM2.5 and PM10 estimates, the BC regional changes were somewhat less dramatic, mostly because trends in power and industrial sector emissions of PM are much less relevant for total black carbon emissions. Globally, over 75% of anthropogenic PM10 and PM2.5 originates from residential combustion, power plants and industry while for BC residential combustion and transport represent more than 75% but the importance varies across regions with Europe and North America having transport as key and rest of the world residential combustion. Our new global estimate of BC emissions suggests higher numbers than previously published owing primarily to inclusion of new sources.”*

REVIEWER:

Supplement When GAINS values are listed in the tables, these are sometimes listed as ranges. I assume these are not uncertainty ranges (as in some of the other ranges in the table), but are GAINS central values and that the range represents the range used in different GAINS regions? It would be useful to clarify this.

RESPONSE: Indeed, these are ranges representing the spread of values across different regions rather than uncertainty ranges. While this is written for example on page 4 before the Table S2.1, we add a respective comment next to other tables in the SI.

REVIEWER:

I suggest providing a more detailed summary of the emissions data. It would make this data more readily useful to the community to have an electronic file (either csv or excel) that provides emissions of the various species by country/region and by sector and fuel (I appreciate that some aggregation with regard to sector/fuel might be necessary). I realize much of this (or perhaps all of this) would be available on-line, but providing this in a supplement will be more accessible and also provide for an archival record of these important results.

RESPONSE: As indicated in the responses to the initial comments of the reviewer, we have developed an additional set of tables with sectoral emissions (including split across key fuels) for the 25 global regions and all considered PM species over time. This is now included as the MS Excel file in the SI.

RESPONSES TO Reviewer 2:

We thank the reviewer for useful comments which have been helpful in improving the manuscript. The responses to specific points raised by the reviewer are provided below.

REVIEWER:

The title mentions that the paper focuses on anthropogenic emissions. However, the paper also discusses open fires. Since the paper is already very long, it might be better to only focus on anthropogenic emissions as stated in the title. The inclusion of emissions from fires (which come from other authors) is a bit confusing.

RESPONSE: Indeed, the paper documents the methodology for PM estimation in GAINS focusing on anthropogenic sources, including also open burning of agricultural waste, but at the same time documents also the complete dataset of ECLIPSE emissions. The latter includes explicit information about what has been used in modelling exercises (see for example Stohl et al., 2015; Eckhardt et al., 2016) and here the reference to the open fires, or more specifically forest and savannah fires is referred to. We allocate one section in the paper to agricultural burning (not all open fires) [see section 3.7] and make explicit references to work on forest and savannah fires on page 13 in the introduction to section 3. We feel that it is justified to provide full documentation of sources used in the entire ECLIPSE set, including forest savannah fires and allocate a page to discuss specific aspect of agricultural burning for which national and regional inputs were used beyond remote sensing data of GFED.

REVIEWER:

Abstract and line 24, page 2: the abstract claims this paper is " the first comprehensive assessment of historical (1990-2010) global anthropogenic particulate matter (PM): : :". However, the EDGAR4.3 inventory described in Crippa et al. (2016) provides emissions for 1970-2010 for PM2.5 and PM10. The statement about being "the first comprehensive assessment" is true for PM1, but not for the other species. Please rephrase.

RESPONSE: As a matter of fact, the word 'comprehensive' is referring here to the comprehensive assessment of several PM species (as well as forming the base for the development of the particulate number inventory referred to in the abstract) within one system assuring that consistent framework is used for the assessment of all of the considered species including PM1, PM2.5, PM10, PMTSP, BC, OC, and OM. But in order to avoid any possible misinterpretation or confusion we simply delete the word 'first' in the abstract as well as in the introduction.

REVIEWER:

Page 18, line 6: " exceptions are old vehicles running on leaded gasoline and preregulation 2-stroke mopeds : : : while latest gasoline direct injection engines have PM mass emissions comparable or even higher than latest diesel engines with particle

filter, however, the absolute level is about one order of magnitude lower than for older generations. This sentence is not clear. What does "absolute level" refer to?

RESPONSE: Thank you for pointing this out. We will rephrase this sentence making clear that the 'absolute level' refers here to the modern (Euro5 and Euro 6) diesel vehicles which have reduced their PM emissions significantly compared the pre or early control stages. Below an example for Italy to illustrate the point with COPERT data for emission factor for PM10 [in g/km]

	g/km	No-control	Advanced controls
Two-stroke		0.176	0.018 (Euro 3)
Light duty gasoline		0.0024	0.0010 (Euro 3 and younger)
Light duty diesel		0.216	0.0018 (Euro 5 and younger)

Interpretation:

=> old 2-strokes are as bad polluters as uncontrolled diesel cars.

Modern diesel cars have reduced their emission rate by a factor 100, such that they are today at the level of or even lower than modern gasoline cars.

REVIEWER:

Page 26, line 20: The authors use quite old data for emission factors for agriculture waste burning. Akagi et al. (Atmos. Chem. Phys., 11, 4039–4072, 2011) have published a more recent and detailed review of all data available on emission factors. The authors should indicate why they did not use this more recent review.

RESPONSE: The paper by Akagi et al (2011) is included in the references and in fact we have considered it while comparing and deriving emission factors for this work. Our emission factors derived from several studies listed compare well with the ones provided in the review by Akagi since they mostly refer to the same work already listed in our previous text. We add explicit reference to Akagi paper in the revised text.

REVIEWER:

Page 29, lines 1-4: these lines should be rephrased. Many recent global chemistrytransport and chemistry-climate models now include detailed aerosols schemes, and PMs distribution are calculated as the sum of the mass of all the components included in the models. Maybe a few older models use the "BC + 1.4 OC" formula to calculate the mass of PM, but the recent models are much more advanced and calculate the mass of PMs in a more accurate way.

RESPONSE: Thank you for pointing this out, in fact we would be interested to know which models are these so that we could an example reference. The mentioned paragraph was meant to highlight two elements, the issue of oversimplifying the total carbonaceous mass in PM where often 1.4 ratio was used to convert OC to POM and the fact that the BC+POM often represents total fine anthropogenic PM in global (not regional) models. Taking into account your comments we have revised one of the sentences in this paragraph that reads now: *"This total fine PM mass has been typically estimated as BC+1.4*OC and only recently a number of models included more detailed aerosol schemes accounting*

for varying BC/OC ratios while still largely neglecting the anthropogenic dust component (e.g., Philip et al., 2017)”

REVIEWER:

Page 31, lines 14-15: The sentence starting with "combined : : :" is unclear

RESPONSE: We have rewritten that sentence, specifically the second part starting with ‘combined’, and the whole sentence reads now: *“However, as further discussion shows, the largest discrepancy for PM₁₀ and PM_{2.5} is for China as well as Europe and Russia; the sum of the differences in these three regions represents about 90% and over 50% of all the difference for PM₁₀ and PM_{2.5}.”*

REVIEWER:

Page 25 of the supplement: the authors should add in their table the TNO-MACC and TNO-MACCII (Kuenen et al., ACP, 2014) inventories, which provide emissions of PM for Europe and neighboring countries. The TNO-MACC inventories are now becoming a reference for atmospheric modeling in Europe, and these emissions should be mentioned in the paper.

RESPONSE: Thank you for this suggestion. We have included MACCII reference and also added the respective emission estimates to the table

REVIEWER:

Page 25 of the supplement: The emissions provided by US EPA are given as the sum of anthropogenic and wildfires. The dataset provided by EPA (note that the last release of the emissions is 2016 and not 2011 as mentioned in the supplement) provides emissions with and without wildfires. It would be better to include the emissions without wildfires, in order to be consistent with the other data in the table.

RESPONSE: Thank you for suggesting the review of these numbers. We retrieved new numbers from the US EPA website (<https://www.epa.gov/air-emissions-inventories/national-emissions-inventory-nei>) for both 2011 and 2014. This allows now for a better comparison to our numbers constructing a similar sector set excluding wildfires. The reference to this EPA source is also included in the manuscript.

Global anthropogenic emissions of particulate matter including black carbon

Zbigniew Klimont¹, Kaarle Kupiainen^{1,2}, Chris Heyes¹, Pallav Purohit¹, Janusz Cofala¹, Peter Rafaj¹, Jens Borken-Kleefeld¹, Wolfgang Schöpp¹

¹International Institute for Applied Systems Analysis (IIASA), Laxenburg, 2361, Austria

²SYKE, Helsinki, Finland

Correspondence to: Zbigniew Klimont (klimont@iiasa.ac.at)

Abstract. This paper presents ~~at~~^{the first} comprehensive assessment of historical (1990-2010) global anthropogenic particulate matter (PM) emissions including consistent and harmonized calculation of mass-based size distribution (PM₁, PM_{2.5}, PM₁₀) as well as primary carbonaceous aerosols including black carbon (BC) and organic carbon (OC). The estimates were developed with the integrated assessment model GAINS, where source- and region-specific technology characteristics are explicitly included. This assessment includes a number of previously unaccounted or often misallocated emission sources, i.e., kerosene lamps, gas flaring, diesel generators, trash burning; some of them were reported in the past for selected regions or in the context of a particular pollutant or sector but not included as part of a total estimate. Spatially, emissions were calculated for 170 source regions (as well as international shipping), presented for 25 global regions, and allocated to 0.5° x 0.5° longitude-latitude grids. No independent estimates of emissions from forest fires and savannah burning are provided and neither windblown dust nor unpaved roads emissions are included.

We estimate that global emissions of PM have not changed significantly between 1990 and 2010, showing a strong decoupling from the global increase in energy consumption and consequently, CO₂ emissions but there are significantly different regional trends, with a particularly strong increase in East Asia and Africa and a strong decline in Europe, North America and Pacific. This in turn resulted in important changes in the spatial pattern of PM burden, e.g., European, North American, and Pacific contributions to global emissions dropped from nearly 30% in 1990 to well below 15% in 2010, while Asia's contribution grew from just over 50% to nearly 2/3 of the global total in 2010. For all considered PM species, Asian sources represented over 60% of the global anthropogenic total, and residential combustion was the most important sector contributing about 60% for BC and OC, 45% for PM_{2.5} and less than 40% for PM₁₀ where large combustion sources and industrial processes are equally important. Global anthropogenic emissions of BC were estimated at about 6.6 and 7.2 Tg in 2000 and 2010, respectively, and represent about 15% of PM_{2.5} but for some sources reach nearly 50%, i.e., transport sector. Our global BC numbers are higher than previously published owing primarily to inclusion of new sources.

This PM estimate fills the gap in emission data and emission source characterization required in air quality and climate modelling studies and health impact assessments at a regional and global level, as it includes both carbonaceous and non-carbonaceous constituents of primary particulate matter emissions. The developed emission data set has been used in several

regional and global atmospheric transport and climate model simulations within the ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants) project and beyond, serves better parameterization of the global integrated assessment models with respect to representation of black carbon and organic carbon emissions, and built a basis for recently published global particulate number estimates.

5 1 Introduction

Particulate matter (PM) or aerosols are solid and liquid particles small enough to remain airborne. PM can be directly emitted to the atmosphere (primary PM) or it can form from gaseous precursors (secondary PM). The size of PM stretches from clusters of molecules with a diameter of a few nanometers up to micrometer-sized abrasion products. This vast dimensional spectrum is reflected in the varying composition and characteristics of PM measured at source and receptor sites. PM species are important constituents of the atmosphere and they play a role in the earth's climate system. Some PM species, i.e. black carbon, absorb visible light and warm the atmosphere, whereas other species, i.e., sulphates and organics reflect sunlight back to space and cool the climate (Bond et al., 2013). PM also serve as condensation nuclei for water vapour to eventually form cloud droplets. There is well-documented evidence that exposure to PM results in adverse effects on human health (e.g., Anenberg et al., 2012; Lim et al., 2012; WHO, 2004).

15 Integrated assessment models, such as the GAINS (Greenhouse gas - Air pollution Interactions and Synergies) model (Amann et al., 2011), utilize data on economic development and corresponding emissions, estimate atmospheric concentrations and further assess the impacts on climate, human health and ecosystems. When this information is combined with potentials and costs for controlling the emissions it is possible to study the cost-efficiency of different policies to reduce the undesirable effects and meet environmental objectives on climate, human health and ecosystem impacts. Such an integrated modelling framework is particularly important for assessing the impacts of particulate matter owing to the multitude of sources, including primary and secondary, and effects on health and climate. All these aspects of PM call for consistent data to support the assessments of impacts and potential for formulating robust strategies to reduce emissions together with consequent concentrations and impacts.

25 This paper presents the first comprehensive assessment of historical (1990-2010) global anthropogenic particulate matter (PM) emissions including consistent and harmonized calculation of mass-based size distribution (PM_1 , $PM_{2.5}$, PM_{10}) as well as primary carbonaceous aerosols; black carbon (BC) and organic carbon (OC). The methodology draws on the earlier developed structure of the PM module in GAINS (Klimont et al., 2002b; Kupiainen and Klimont, 2004, 2007) but was extended to include new information as well as previously unaccounted sources, i.e., gas flaring, kerosene lamps, diesel generators.

30 A recent GAINS model development includes extension to include particulate number (PN) emissions (Paasonen et al., 2013). This builds on the emission methodology and estimates described in this paper making use of one of the datasets

Field Code Changed

(ECLIPSE V5) to calculate past and future PN emissions and their spatial distribution. The respective documentation and discussion paper is in review (Paasonen et al., 2016).

While the results presented in this paper focus on the outcomes included in the ECLIPSE V5a version of data, there were several datasets developed within the ECLIPSE project¹ (Stohl et al., 2015) and the key differences between the datasets are also briefly discussed. Table 1 gives an overview of the datasets that are accessible from the GAINS website²; the paper describing the projections is in review for this issue of ACP (Klimont et al., in preparation).

2 Method

The ECLIPSE emission data set was created with the GAINS (Greenhouse gas – Air pollution Interactions and Synergies; <http://gains.iiasa.ac.at>) model (Amann et al., 2011), which calculates emissions of air pollutants and Kyoto greenhouse gases (GHG; i.e., carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O) and the three F-gases) in a consistent framework. The GAINS model holds essential information about key sources of emissions, environmental policies, and further mitigation opportunities for 170 country-regions. The model relies on international and national statistics of activity data for energy use, industrial production, and agricultural activities (see section 3) for which it distinguishes all key emission sources and control measures. Several hundred technologies to control air pollutant and greenhouse gases emissions are represented allowing simulation of implemented air quality legislation (see section 2.3).

Since previous work (Cofala et al., 2007; Klimont et al., 2002b, 2009, Kupiainen and Klimont, 2004, 2007; Shindell et al., 2012) we have reviewed recent literature, including non-peer reviewed studies, to improve characterization of the source sectors and control technologies in the GAINS model, update the assumptions about penetration of control measures, and to include previously unaccounted or poorly allocated sources. Emission sources that have been recently added or for which the emission calculation has been refined include: flaring of associated petroleum gas in the oil and gas exploration sectors, kerosene lamps for lighting (further development of estimates originally presented by Lam et al. (2012)), diesel generator sets, high emitting vehicles, international shipping, trash burning as well as brick kilns (see section 3).

Further improvements in the emission model have been made especially for China (Klimont et al., 2013; Wang et al., 2014; Zhao et al., 2013), where large changes have occurred recently as well as new data becoming available, but also for Europe where results of the consultation with national experts during the review of the EU National Emission Ceilings Directive were considered in the last datasets (Amann et al., 2015). Finally, the regional resolution of the global GAINS model has been improved by distinguishing more countries in Latin America where five regions (Argentina, Brazil, Chile, Mexico, all remaining Latin America) were replaced with 13 regions in version V5a, including most countries of South America,

¹ European Commission FP7 project ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants); Project no. 282688; <http://eclipse.nilu.no>

² http://www.iiasa.ac.at/web/home/research/researchPrograms/air/Global_emissions.html

Mexico, Central America, and the Caribbean; a full list of country-regions in the global GAINS application is included in the supporting information (SI).

2.1 PM estimation method

The methodology to derive particulate matter (PM) emission factors and calculate emissions relies on the methods documented in (Klimont et al., 2002b; Kupiainen and Klimont, 2004, 2007). However, apart from updates to emission factors a number of modifications and extensions have been introduced since, especially for carbonaceous particles. We summarize the principles below allocating more space to discuss extensions.

The emissions of PM in the GAINS model are calculated for several size classes: submicron fraction (particles with diameter smaller than 1 μm ; $\leq\text{PM}_1$), fine fraction ($\leq\text{PM}_{2.5}$), coarse fraction ($\text{PM}_{2.5}>\text{PM}_{10}$), and large particles ($\geq\text{PM}_{10}$). PM_{10} is calculated as the sum of fine and coarse fractions. Total suspended particles (TSP) as the sum of fine, coarse, and $\geq\text{PM}_{10}$ fractions. Additionally, black carbon (BC) and organic carbon (OC) are calculated.

The methodology includes the following steps:

(i) region- (i), sector- (j) and fuel- (k) specific “raw gas = unabated” emission factors for total suspended particles (TSP) are derived. For solid fuels (excluding biomass and use of solid fuels in small residential installations) the mass balance approach is used where ash content (ac) and heat value (hv) of fuels, and ash retention in boilers (ar) for given combustion technologies are considered Eq. (1):

$$ef(TSP)_{i,j,k} = \frac{ac_{i,j,k}}{hv_{i,j,k}}(1 - ar_{j,k}) \quad (1)$$

For liquid fuels, biomass, solid fuels used in small residential installations, industrial processes, mining, storage and handling of bulk materials, waste incineration, agriculture³, and transport, TSP emission factors are taken from the literature;

(ii) considering fuel- and sector-specific size fraction profiles reported in the literature, “raw gas” emission factors for each of the size fractions and carbonaceous species are estimated;

(iii) the emission factors for organic carbon (OC), calculated in the previous step, are adjusted considering carbonaceous fraction in $\text{PM}_{2.5}$ and organic carbon (OM); see section 2.1.1 for discussion;

(iv) PM emissions are calculated for each size fraction and carbonaceous species applying the following equation Eq. (2), where also the application rates of control technologies (X) and size fraction specific emission removal efficiencies (eff) are taken into account:

$$E_{i,y} = \sum_{j,k,m} E_{i,j,k,m,y} = \sum_{j,k,m} A_{i,j,k} ef_{i,j,k,y} (1 - eff_{m,y}) X_{i,j,k,m}, \quad (2)$$

where i,j,k,m are region, sector, fuel, abatement technology; y size fraction, i.e., fine, coarse, $\text{PM}_{>10}$, or carbonaceous species (BC, OC); $E_{i,y}$ emissions in region i for size fraction y ; A activity in a given sector, e.g., coal consumption in power

³ For livestock, emission factors refer to housing period and therefore, information on the length of this period (one of the parameters in the GAINS model) is considered to derive annual animal- and country-specific values.

plants; ef “raw gas” emission factor; $eff_{m,y}$ reduction efficiency of the abatement option m for size fraction y , and X actual implementation rate of the considered abatement, e.g., percent of total coal used in power plants that are equipped with electrostatic precipitators. If no emission controls are applied, the abatement efficiency equals zero ($eff_{m,y} = 0$) and the application rate is one ($X = 1$). In that case, the emission calculation is reduced to simple multiplication of activity rate by the

5 “raw gas” emission factor.

There are a few source sectors where additional assumptions are made in order to develop emission factors used in the calculation. Specifically, for gas flaring additional information about the composition of associated gas is used (see section 3.6.3 for more details), and to estimate emissions from high emitting vehicles (or superemitters) assumptions about region-specific shares of high emitters as well as technology and pollutant specific increments, compared to the average fleet emissions factors (excluding high emitters), are made (see section 3.4.1).

10

2.1.1 Adjustments of carbonaceous particle emission factors

While we principally follow the definition of black carbon (BC) given by Bond et al. (2013), i.e., “...a distinct type of carbonaceous material that is formed primarily in flames, is directly emitted to the atmosphere, and has a unique combination of physical properties. It strongly absorbs visible light, is refractory with a vaporization temperature near

15 4000K, exists as an aggregate of small spheres, and is insoluble in water and common organic solvents”, the available measurement studies have not been consistent in this respect, and it has not been possible to systematically follow the definition in developing the input data for emission estimates; this has also been discussed in our previous papers (Kupiainen and Klimont, 2004, 2007).

Organic carbon (OC) refers to the carbon fraction in numerous organic compounds that contain hydrogen and, usually, oxygen, and are emitted to the air as particles (Bond et al., 2013). To attain the total mass associated with the organic compounds, organic matter (OM), OC needs to be multiplied with a fraction that depends on the suite of compounds emitted and varies between the emission sources. We introduce source specific OM to OC fractions for primary emissions found from literature, varying between 1.3 and 2.1 (Aiken et al., 2008; Tissari et al., 2007; Turpin and Lim, 2001). Due to the lack of a formal definition and available measurement studies we have not attempted so far to separate emissions of “brown carbon”, a group of absorbing compounds considered a subset of organic aerosol (Bond et al., 2013).

25

Emission factors of organic carbon (ef_{OC}) for each GAINS technology category are calculated using a mass balance equation Eq. (3). This equation has been introduced to ensure that the mass balance of the chemical species of particulate matter (black carbon and organic carbon) will still stay within physical limits of the PM mass metrics applied in GAINS. The calculation uses $PM_{2.5}$ as the limiting mass metric since the emissions of carbonaceous matter occur primarily in that size range. We introduce only few exceptions where larger carbonaceous particles are expected to be present, e.g., tyre wear.

30

$$ef_{OC} = (ef_{PM_{2.5}} \times f_{carb} - ef_{BC}) \div f_{OM}, \quad (3)$$

where f_{carb} is the mass fraction of the total carbonaceous matter, or black carbon and organic matter, in $PM_{2.5}$, f_{OM} the average organic molecular weight per carbon weight in particular matter, ef_{BC} the emission factor of BC, $ef_{PM_{2.5}}$ the emission factor of

PM_{2.5}. Emission factors of BC and PM_{2.5} as well as f_{carb} and f_{OM} are estimated based on emission measurement data. The final set of OC emission factors is checked for consistency with emission measurements.

Fraction of carbonaceous matter in PM_{2.5} (f_{carb}) varies significantly between source sectors. Highest fractions are usually found in residential combustion and transport sectors in technologies with poor combustion, where over 90 percent of the particulate matter is estimated to consist of carbonaceous matter. As the combustion process becomes more efficient and optimized, the fraction reduces drastically and, for example, in large modern power plants, which have optimized combustion processes and efficient air pollution abatement technologies, the fraction is typically negligible; see discussion in Kupiainen and Klimont (2007) and Sippula et al. (2009).

The average fraction of organic molecular weight per carbon weight (f_{OM}) varies also between different emission source sectors and fuels. For combustion of biomass, including wood, we use $f_{OM} = 1.8$, which represents approximately the middle of the range (1.6 to 2.1) of f_{OM} values available for combustion of different wood species in the literature (Aiken et al., 2008; Tissari et al., 2007; Turpin and Lim, 2001). For diesel and gasoline in transport sector, we use $f_{OM} = 1.3$, based on Aiken et al. (2008).

2.2 Model technology resolution

The GAINS model structure includes representation of key emission sources compatible with global and regional emission inventories but the calculation often distinguishes an additional level of detail where combustion technology (e.g., pulverized coal or grate firing boilers, fireplaces, various stoves, pellet boilers, etc.) as well as emission control technology (e.g., wet scrubbers, fabric filters, fan assisted stoves, diesel particulate filters, etc.) are explicitly distinguished (see also Eq. (2)). Such an approach has been an integral part of the GAINS model development for both particulate matter (e.g., Klimont et al., 2002b; Lükewille et al., 2001) and other pollutants (e.g., Amann et al., 2011; Cofala and Syri, 1998; Klimont et al., 2002a); the details for PM are documented in Klimont et al. (2002b) and the current structure can be reviewed in the on-line application of the GAINS model⁴. This approach has also been used in other emission assessment studies and often referred to as 'technology-based' (e.g., Bond et al., 2004; Lu et al., 2011; Zhao et al., 2013).

Implementation of such technology resolution requires additional assumptions about the shares of activity in a given sector falling into each subcategory and share of activity controlled with a specific mitigation measure. The following sections highlight and document briefly the assumptions for key sectors.

2.2.1 Residential combustion: cooking, heating, lighting

GAINS divides residential-commercial sector into several fuel-dependent categories (Table 2). The division is driven by varying emission characteristics and available control options (Table 3). While such a structure is fairly compatible with the available emission measurements (see section 3.1), it is challenging to distribute fuel consumption into these categories as

⁴ <http://gains.iiasa.ac.at>; select any of the accessible regional version to view the model structure

typically statistical data is available either as total residential sector or split into commercial/residential/other (e.g., IEA, 2015a, 2015b). We rely on a mix of sources and our own assessment to derive the respective shares. There have been several assessments at a global level where either total fuel demand for cooking and heating, allocation between various fuels, or stove types was attempted (Bonjour et al., 2013; Chafe et al., 2014; Fernandes et al., 2007). For Europe, such data are not readily available; however, within the work on revision of air quality legislation we were involved in several rounds of stakeholder consultations where national experts representing various sectors reviewed GAINS assumptions (Amann et al., 2015) and all data can be viewed in the on-line model. For the US and Canada, a similar discussion and exchange took place within the work of the Arctic Council where the GAINS model was used to develop unified emissions and scenarios (AMAP, 2015). For Australia and New Zealand a number of local studies were used (Driscoll et al., 2000; Scott, 2005; Todd, 2003).

The allocation of fuel between various categories varies between Europe, North America, and OECD Asia and Pacific where most solid fuel is used for heating (e.g., Chafe et al., 2015), and most of Asia, Africa and Latin America where cooking is the primary use. Consequently, nearly all solid fuels in South Asia, Africa, and Latin America are allocated to cooking stoves. For Asia, we draw on the past and ongoing collaboration on the development of the GAINS-Asia model (Amann et al., 2008; Klimont et al., 2009; Zhang et al., 2006; Zhao et al., 2013) where assumptions on the split between heating and cooking as well as fuel used in medium size boilers were made, as well as several peer-reviewed publications (e.g., Venkataraman et al., 2010). For Latin America, information about this sector structure originates from the discussions with the authors of various assessments of effectiveness of clean cooking programs (e.g., Pine et al., 2011; Ruiz-Mercado et al., 2011).

The GAINS model includes a number of mitigation measures in this sector (Table 3), although some of them might be seen more as different types of installations, e.g., various stove types already in place (for specific discussion of their assumed characteristics see Supplementary Information – section S2). While there has not been a lot of success in sustained replacement of traditional stoves with improved clean burning stoves (e.g., Foell et al., 2011; Pine et al., 2011; Ruiz-Mercado et al., 2011; Wickramasinghe, 2011), it is important to consider the varying level of implementation across the regions if such information is available. As with the allocation of fuel use (see discussion above), we rely on data and assessments collected within several bilateral projects (e.g., Amann et al., 2008, 2015), peer-reviewed papers (e.g., Klimont et al., 2009; Lewis and Pattanayak, 2012; Li et al., 2016; Pine et al., 2011; Ruiz-Mercado et al., 2011; Shrimali et al., 2011; Silk et al., 2012; Troncoso et al., 2011), and published reports (Adria and Bethge, 2013; Germain et al., 2008; Scott, 2005; Todd, 2003). Technology structure has impact on implied (average) emission factor for a given category distinguished in the model. While changes for biomass cooking stoves were rather limited at a larger scale, resulting in up to 10% decline in implied PM_{2.5} emission factor in Asia and up to 5% in Latin America, we estimate larger impact for residential biomass heating. We estimate that for PM_{2.5}, the 'global average emission factor' declined from 1990 to 2010 by about 15% which is mostly due to strong increase in sales of pellet stoves and boilers in Western Europe leading to nearly 40% reduction in implied emission factor (Fig. 1). Interestingly, the changes in emission factors for BC are less pronounced (Fig. 1) since the

Field Code Changed

[improved stoves are more efficient in reducing total level of particulate matter emissions rather than black carbon \(see further discussion in section 3.1 and SI \(S2\)\).](#)

One of the recent developments in the GAINS model was explicit distinction of use of kerosene for cooking and lighting (Table 2); earlier all kerosene was allocated to cooking. This modification was driven by the study highlighting the potentially high contribution of kerosene lamps to black carbon emissions (Lam et al., 2012). The emissions depend on what type of lamp is used, and for historical data we distinguish between wick and hurricane lamps, with the former representing the majority (Lam et al., 2012; Mills, 2005). As a default we assume 80% kerosene wick lamps in South Asia and 50% in other developing world regions. For discussion of how total activity data for kerosene lighting is calculated see section 3.2.

2.2.2 Transport

The GAINS model distinguishes several source categories within the road and non-road transport sectors. Road transport is disaggregated into six vehicle categories: 2- and 4-stroke two-wheelers, passenger cars and vans, light duty vehicles, heavy duty trucks, and buses. The non-road mobile sources are grouped into nine broad categories: agriculture and forestry, construction and mining, rail, inland navigation, coastal shipping, aviation (only landing and take-off), 2-stroke engines (e.g., in households, recreation, forestry, etc.), other land based engines. Each vehicle/machine category is associated with a fuel according to its propulsion type; several fuels are distinguished: diesel, gasoline, CNG, LPG, jet fuel or kerosene, heavy fuel oil as well as hydrogen and electricity. For each of the fuel-vehicle combinations, activity data (fuel consumption and km-driven for road vehicles) are sought and are usually available in national and international statistics for road transport categories while are often incomplete, allocated under other sectors, or even lacking for non-road sources. For a complete list of transport sources and fuels see Table S8.1.

[While we do not model specifically vehicle vintages, the new emission standards are typically synonymous with a new vintage year of a particular vehicle category.](#) In order to reflect existing legislation (section 2.3), each fuel-vehicle combination is further subdivided by its average emission level. The key proxy for the emission level is the exhaust emission legislation in force in the country (or region) at the time when the vehicle type is put into service or to which level it is retrofitted. The associated emission factors describe the emission rates for the pollutants averaged over the actual operating conditions, vehicle sizes, and machine types as well ages and model years within one emission standard. More details about the emission factors, control stages in GAINS, and discussion of high emitting vehicles are provided in section 3.4.

[Depending on the region, the implied \(average\) emission factors for key vehicle categories have been changing in the considered period. We estimate that the global average BC emission rate has declined by 2010 by nearly 20% for heavy and duty vehicles but in several regions like North America, Western Europe, developed Asia and Pacific the reduction was about 60-65%, Central Europe about 40-50%, while for most other regions small or no significant change was estimated \(Fig. 2\). Similar trends were found for light duty vehicles but the reductions are typically higher with a global average declining by nearly 35% \(Fig. 2\).](#)

2.2.3 Large scale industrial combustion

The available statistical data allows for allocation of fuel into key sectors, like power plants and industrial boilers, but owing to varying emission characteristics and often different legislation for different boiler types, the GAINS model distinguishes additionally a number of selected plant and boiler types (for more background discussion see Klimont et al. (2002b)).

5 Specifically, the power sector is divided into existing (constructed before 2005), new and modern plants for which additionally large and small plants (grate firing) are distinguished. Structural changes as well as increasing stringency of emission legislation resulted in declining emission factors. For example, we estimate that global average PM_{2.5} emission factor for coal power plants dropped by about 40% with Northern America, Europe, and Japan having 70-80% decline and even for China we estimate over 70% reduction while in Russia and several Former Soviet Union countries only 20-30% decline (Fig. 3). Industrial combustion is associated with several sectors for which also small boilers are included to capture the large numbers of often old and poorly controlled solid fuel grate firing boilers in the developing countries (e.g., Wang et al., 2014; Zhao et al., 2013), for example in China they accounted for about 85% of all industrial boilers (Wang et al., 2009). For industrial coal use lower reductions in average emission factors were achieved than for power plants with exception of Eastern Europe and some Former Soviet Union countries where collapse of heavy industry in the period 1990-2000 resulted in decline of emission factors by over 90% compared to the 1990. While the estimated changes in emission characteristics could be modelled more accurately if assumptions about equipment vintages were made, the GAINS model does not explicitly include it with exception of power sector (see above). Instead, GAINS defines technical lifetimes of the add-on control technologies (e.g., cyclones, electrostatic precipitators, fabric filters) and considers that these can be principally applied shortly after respective legislation is put in place.

10 Finally, the GAINS model structure has been extended to distinguish diesel generator sets; previous GAINS regional and global assessments of PM or carbonaceous particles (Cofala et al., 2007; Klimont et al., 2009; Kupiainen and Klimont, 2007) included their fuel consumption in power and residential combustion sectors. The new structure allows for better representation of emissions and mitigation opportunities, especially in regions with low reliability of electricity supply and poor emission standards, e.g., South Asia. The estimates of regional diesel generators fuel use is discussed in section 3.3.

25 2.2.4 Industrial processes

Most industrial processes are sources of particulate matter emissions. For the majority of them emissions are calculated using total production volumes without distinguishing specific stages of the processing chain. However, for a number of manufacturing processes we define default plant profile and distinguish between process and fugitive emissions, for details see Klimont et al. (2002). Additionally, for selected industries a more detailed structure was designed to reflect the significant differences between types of plants (kilns); this has been done for cement, coke, and brick manufacturing.

30 The key driver behind the extended structure for cement and coke manufacturing was developments in China where in the last decades strong growth resulted in often rapid transformation of the two sectors. For cement production rotary kilns with

Field Code Changed

precalciner and shaft kilns are distinguished for which the activity split has been developed in collaboration with Tsinghua University (Zhao et al., 2013). Such technological changes, often accelerated by political and economic transformation (e.g., Eastern Europe and Former Soviet Union), and legislation landscape resulted in rather significant change in average emission rates in cement production sector. We illustrate that in Fig. 3 where in several regions GAINS implied PM_{2.5} emission factors in 2010 are lower by up to 90% than in 1990. The coke production sector in China experiences rapid transformation from traditional ovens to mechanized integrated coke ovens which have different emission characteristics; the changes in the structure of the sector are discussed by Huo et al. (2012). Currently, the information about the comparable technology split is not available for other countries for which emissions are calculated without such distinction.

Brick manufacturing

There are strong regional differences in brick manufacturing sector structure that is especially relevant in the developing world where a large share of the market is occupied by traditional, heavy-polluting kilns. Our earlier work focused on characterizing the brick sector in Asia, by far the largest producer, and therefore the distinguished kiln types reflected practices in Asia (Klimont et al., 2009; UNEP/WMO, 2011). However, such design of the model did not allow to correctly address the structure of this sector in other regions like Africa or Latin America and the Caribbean. We have reviewed regional and national assessment studies to identify typical regional profiles (distribution of production by kiln types) of the brick manufacturing sector, including also typically used fuels; such profiles change over time and it has been considered where such information was found. Table 4 shows the kiln structure included in GAINS and highlights key representative technologies assumed for different world regions. The overview of studies used to develop the respective assumptions is provided in the SI (S5). The overall brick production data are discussed in section 3.6.2 and SI (Table S5.2).

2.3 Emission legislation

We have collected information about existing international and national requirements with respect to emission limit values for stationary and mobile sources and estimated control technology implementation rates required to achieve respective standards in all GAINS regions. The interpretation of the laws and translation into the set of GAINS technologies with the associated emission rates under average operating conditions has been discussed previously in a number of papers and assessments addressing regional (Amann et al., 2015; Klimont et al., 2009; Kupiainen and Klimont, 2007; Wang et al., 2014) and global (Amann et al., 2013; Cofala et al., 2007; Rao et al., 2013; Riahi et al., 2012; UNEP/WMO, 2011) emissions.

For a number of sources there exist global databases summarizing current laws and emission limit values, including power plants (IEA, 1997; IEA CCC, 2012), transport (Delphi Inc., 2013, 2015; ICCT & Dieselnet, 2014), and the cement industry (Edwards, 2014). Additionally, specific regional and national laws and policy implementation studies were reviewed, i.e., for the European Union a number of Directives was considered (Crippa et al., 2016; EC, 2001a, 2001b, 2010; Krasenbrink and Dobranskyte-Niskota, 2008), for Asia several peer-reviewed studies (Goel and Guttikunda, 2015; Guttikunda and Jawahar, 2014; Huo et al., 2011, 2012; Klimont et al., 2009; Liu et al., 2015; Lu et al., 2011; Wang et al., 2014; Zhang et al., 2006) as

well as other sources (CAI-Asia, 2011; CPCB, 2007; IIDFC, 2009); for Latin America and Caribbean additional information was obtained for the brick sector (e.g., Stratus Consulting, 2014) and for Argentina, Brazil, and Mexico also for the transport sector (e.g., Ministério do Meio Ambiente, 2011).

In the course of development of the several ECLIPSE datasets, the legislation information and mostly the rates of enforcement and implementation of actual measures have been revisited. The key updates in version *V4a* (see Table 1) include consideration of the initial round of consultations with European Union member states experts within the review of the National Emission Ceiling (NEC) directive (Amann et al., 2012), which included comparison of GAINS estimates with the emissions officially reported to the Centre on Emission Inventories and Projections (CEIP; www.ceip.at) under the Convention on Long-range Transboundary Air Pollution. A much more substantial update came with version *V5a* where for China the 12th Five Year Plan policies were introduced, resulting in revision of the implementation and enforcement rates of control measures for 2010 drawing also on analysis of progress in legislation implementation in China (e.g., Lin et al., 2010; Zhang et al., 2015). Furthermore, the legislation for cement industry was reviewed and updated (Edwards, 2014), emissions from international shipping were also calculated, the treatment of the non-road mobile machines were reviewed, and for Latin America and Caribbean (LAC) the GAINS model has been revised to include nearly all single countries⁵ and consequently required definition of control strategies reflecting current legislation for each country. Finally, also for the European Union an update was performed in *V5a* to include the latest status of discussion with the national experts (Amann et al., 2015), as well as new submissions of PM_{2.5} emissions (also for the past years) to CEIP, especially for 2010.

2.4 Spatial and temporal distribution

The GAINS model calculation is performed for 170 regions globally and for Europe and Asia the calculation and results are directly available by country or even subnational level from the online version of the model (<http://magcat.iiasa.ac.at>) for all ECLIPSE data sets. At a global level, the emissions and activity data are available online at the resolution of 25 global regions (see Supporting Information (SI), S7) and key sources (<http://magcat.iiasa.ac.at/gains/IAM/index.login>); the structure is compatible with most of the global integrated assessment models. Additionally, the total annual emissions were gridded and temporal (monthly) distributions were developed.

The GAINS particulate matter emissions were distributed into 0.5° x 0.5° longitude-latitude grids and stored in *netCDF* format files available from http://www.iiasa.ac.at/web/home/research/researchPrograms/air/Global_emissions.html as well as from the ECLIPSE project web: <http://eclipse.nilu.no>. The files contain several layers (Table 5), reflecting key sectors (consistent with Representative Concentration Pathways (RCP) used in the Intergovernmental Panel for Climate Change Fifth Assessment Report (IPCC AR5)), and a total emission layer. The spatial distribution was prepared from RCP-consistent proxies as used and further developed within the Global Energy Assessment project (GEA, 2012). These are in line with proxies applied within the RCP projections as described in Lamarque et al. (2010) and were modified to

⁵ Previous versions included five regions: Argentina, Brazil, Chile, Mexico, other LAC

accommodate more recent information where available, e.g., population distribution, open biomass burning, effectively making them year-specific (Klimont et al., 2013; Riahi et al., 2012).

In the process of preparing gridded emissions we have developed additional layers which were merged into the sector layers listed in Table 5. The primary example, relevant for particulate matter emissions, is the flaring layer which has been developed by IIASA using the information on flare location areas developed in the collaborative project of NOAA, NASA, and the World Bank (Elvidge et al., 2009, 2011). This layer contains emissions from flaring in oil/gas exploration and it is for the first time that a global PM emission assessment includes this source with explicit spatial allocation (Fig. 44); this dataset was used within the ECLIPSE project and highlighted the relevance of proper distribution of black carbon emissions from this source (Stohl et al., 2013). The flaring emissions are integrated in the *Energy* (Table 5) layer but a separate file with all emissions from flaring only is also available for download.

2.4.1 Temporal distribution

The GAINS model does not explicitly include any assumptions about temporal distribution and therefore all emissions are calculated as annual totals. However, within the MACEB⁶ and ECLIPSE projects we have developed monthly emission profiles for the gridded output, shares of emissions in different months in each grid, for a number of sources. The focus was on allocation of domestic heating and cooking emissions where the methodology combines the stove use assumptions from Streets et al. (2003) with the global gridded temperature fields from the CRU3.0 archive⁷ of monthly mean temperatures (Brohan et al., 2006). The shares were developed for six years (2000-2006) and an average was eventually used as a representative monthly fraction. Fig. S1 in SI compares this pattern with other existing estimates for selected countries. The importance of considering the temporal distribution of residential combustion emissions developed within ECLIPSE has been demonstrated in Stohl et al. (2013) for the Arctic.

For the energy sector, country-specific monthly patterns were created for selected regions based on available data; for Europe and Russia such data were originally developed in the GENEMIS project (Ebel et al., 1997) and are readily available in the EMEP database; for North America we used the US-EPA Clearinghouse for Emission Inventories (<http://www.epa.gov/ttnchie1/emch/temporal/>) and the US Energy Information Agency Monthly Energy Review (<http://www.eia.gov/totalenergy/data/monthly/>); for Thailand the information provided by (Vongmahadlek et al., 2008, 2009) was applied. For all other regions, the temporal distribution file includes constant emissions across the year.

The emissions from open burning of agricultural residues are seasonal since the activity is related to growing cycles and harvesting of different crop types. A global spatial and temporal representation was developed based on the timing and location of active fires on agricultural land in the Global Fire Database GFEDv3.1 (<http://www.globalfiredata.org/Data/index.html>) combined with annual emissions from GAINS. All active grid cells

⁶ MACEB - Mitigation of Arctic warming by Controlling European Black carbon emissions, European Union Life+ project no: LIFE09 ENV FI 572

⁷ <http://badc.nerc.ac.uk/data/cru/>

Field Code Changed

(0.5° x 0.5°) in the monthly data from 1997 to 2010 in GFED were summed up and normalized. Also for other agricultural activities several patterns were developed but they are more relevant for ammonia and methane emissions and therefore discussed in Klimont et al. (in preparation).

3 Emission sources – activity data and emission factors

5 Here we highlight the contribution of key sources to total emissions and document the sources of activity data and emission factors used in the GAINS model for all relevant sources of particulate matter (PM) emissions, including discussion of differences between several published ECLIPSE datasets. The technology splits and air pollution legislation is discussed in section 2.2 and 2.3.

The basic statistical data for energy consumption, industrial output, and agriculture originates from International Energy Agency (IEA, 2015a, 2015b), Eurostat (EUROSTAT, 2011), UN Food and Agriculture Organization (<http://faostat.fao.org>), and several national sources that have been used in the course of collaboration with several partners in Europe (e.g., Amann et al., 2012, 2015) and Asia (e.g., Amann et al., 2008; Purohit et al., 2010; Zhang et al., 2006; Zhao et al., 2013). For several sectors more specific regional data were used; see discussion in the following source-specific sections. There are also differences in data used for various versions of the ECLIPSE dataset; an overview is provided in Table 1. For activity data, the most significant changes are due to update of the historical data in version V5 and V5a where all IEA statistical data was imported at national level and processed for use in GAINS. Furthermore, for Europe the consultations with national experts during the National Emission Ceiling Directive (NEC) revision process led to a number of updates (including activity, emission factors, penetration of control technologies) for the EU-28, specifically in V4a (Amann et al., 2012) and then in the V5a (Amann et al., 2015) version. Both of these updates were most significant for the year 2010 as new information became available.

The GAINS model database has been developed for five-year periods starting in 1990 and extending to 2050 and as shown in Table 1, different ECLIPSE versions include estimates for either the whole time horizon or selected five-year periods. There is one exception; in the V3 dataset we estimated also global emissions for 2008 and 2009. In order to calculate emission fields for 2008 and 2009 we have used a number of additional sources of information to develop scaling factors for emissions of the year 2005. The exercise was performed at the finest possible sectoral resolution compatible with GAINS but for some regions only key aggregated sectors (see Table 5) were estimated. For most sectors, the country specific emission ratios were developed using officially reported emissions by US-EPA (<http://www.epa.gov>), Environment Canada (<http://www.ec.gc.ca/inrp-npri/>), within UNECE LRTAP Convention (<http://www.ceip.at>), and 2012 UNFCCC national inventory submissions (<http://unfccc.int/>). For countries where we found no submissions, emissions for key sectors (Table 5) were linearly interpolated between 2005 and 2010. Additionally, for flaring in the oil and gas industry the emissions for 2008 and 2009 were calculated using GAINS methodology and data on activities available from the NASA report (Elvidge et al.,

Field Code Changed

2011). Finally for open biomass burning we have used the data from the GFED v3.1 global database (<http://www.globalfiredata.org/>).

What is not included and where to find it

None of the ECLIPSE datasets includes estimated emissions from forest and savannah fires (note that emissions from open burning of agricultural residue are included; see section 3.7), which can be obtained from the GFED v3.1 global database (van der Werf et al., 2010) or a more recent version GFED v4 that was made available subsequently (Randerson et al., 2015). GFED contains emissions for BC, OC, PM_{2.5}, and total particle matter (TPM) for the period 1997-2014 in varying temporal and spatial distribution (including gridded dataset) depending on the version (<http://www.globalfiredata.org/>).

None of the ECLIPSE datasets includes emissions from international aviation but these can be acquired from the Representative Concentration Pathways (RCP) database available at e.g., <http://tntcat.iiasa.ac.at:8787/RcpDb/>. The data originates from a study by Lee et al. (2009) and were used in the development of the RCPs (Van Vuuren et al., 2011). However, only emissions of black carbon (BC) are included.

Versions *V3* and *V4a* do not include emissions from international shipping and at the time we recommended using datasets developed for the RCP process (Buhaug et al., 2009; Eyring et al., 2010). Version *V5* and *V5a* include international shipping estimates for all PM species (the RCP set contains only BC and OC), which we have developed drawing on the QUANTIFY⁸ project spatial distribution (Endresen et al., 2007) and activity data from Buhaug et al. (2009); for more details see section 3.4.2. The datasets for international shipping, aviation, and open burning have been extracted for use in the ECLIPSE project and can be downloaded (upon request) from the project website <http://eclipse.nilu.no>.

3.1 Residential sector

Several previous studies (e.g., Bond et al., 2004; Cofala et al., 2007; Kupiainen and Klimont, 2007; Lu et al., 2011; Venkataraman et al., 2005) showed that the residential sector is an important source of PM emissions at a regional and global level, especially of carbonaceous species. GAINS distinguishes a number of source categories for residential sector heating and cooking, i.e., fireplaces, stoves, single house boilers and medium sized boilers as well as a number of solid fuels, i.e., fuelwood, agricultural residues, dung, and coal as well as liquid and gaseous fuels, i.e. kerosene, fuel oil, LPG, and natural gas; see Table 2. The data about fuel consumption used in the GAINS model originates primarily from IEA statistics but is enriched with additional data from regional statistics and studies. This includes regional, rather than national, statistics of coal use in China (Zhao et al., 2013) but most of all additional assessments of biomass use for cooking and heating in several regions; for US, Canada, Finland, Sweden, and Norway drawing on the collaboration within the Arctic Council (AMAP, 2015); for Australia and New Zealand (Driscoll et al., 2000; Scott, 2005); Asia (Amann et al., 2008; Klimont et al., 2009; Purohit et al., 2010; Venkataraman et al., 2010); and finally for Europe where exchange with national experts led to

⁸ QUANTIFY - Quantifying the Climate Impact of Global and European Transport Systems; European Union Sixth Framework project (<https://www.pa.op.dlr.de/quantify/>)

Field Code Changed

consideration of several local datasets in the GAINS model (Amann et al., 2015). The data used in the last version of ECLIPSE (V5a) for Europe are comparable with the independent fuel estimate by Denier van der Gon et al. (2015). Beyond the total fuel use, the split by fuel and installation types is of high relevance (see discussion in section 2.2).

The global fuel use for cooking and heating used in GAINS ranges from about 2100±200 Tg in 1990 to 2600±200 Tg in 2010 and compares well with the total fuel demand estimated in other global studies; for example, Fernandes et al. (2007) estimated total biofuel use in 2000 at 2460 Tg, which compares with GAINS value of 2200-2500 Tg (the range given owing to uncertainties in assumptions about heat value of various biofuels).

The emission factors aim to reflect real world emissions (e.g., MacCarty et al., 2007; Roden et al., 2006, 2009), i.e., incorporate emission measurements of diluted samples, and have been recently compared and updated for Europe (Boman et al., 2011; Pettersson et al., 2011; Schmidl et al., 2011; Tissari et al., 2008, 2009), specifically for modern stoves and boilers, Asia (Cao et al., 2006; Chen et al., 2009; Habib et al., 2008; Li et al., 2009; Parashar et al., 2005; Venkataraman et al., 2005; Zhi et al., 2008, 2009), Latin America (Johnson et al., 2008).

Emission factors and shares of BC and OC in particulate mass emissions from selected measurement literature together with the range of values used in the GAINS model are presented in Tables S2.1 – S2.4 in the SI (S2), where also a brief characterization of stove and boiler categories used in GAINS is provided.

3.2 Kerosene lamps

Most of the previous emission studies did not highlight particulate matter emissions from kerosene used for lighting, primarily because the information about emission factors and fuel use was either not available or sparse. Only after Lam et al. (2012) reported very high black carbon emission factors, indicating that this is potentially an important ‘missing’ source, has more work been done to distinguish between kerosene used for cooking and lighting; the new estimates suggest this source might contribute 5-10% of global BC emissions.

Approximately 250 million households (about 1.3 to 1.5 billion people, mostly in developing Asia and Sub-Saharan Africa) lacked access to reliable electricity to meet basic lighting needs in 2010 (IEA, 2012b)(IEA, 2012). These households often rely on fuel-based lighting, with the majority burning kerosene in wick-type lamps (Lam et al., 2012; Mills, 2005); their consumption was estimated at up to 25 billion litres of kerosene per year (Lam et al., 2012). Growing evidence suggests that these light sources pose risks to health (Pokhrel et al., 2010) and environment (Lam et al., 2012), and improvements to lighting may provide numerous welfare benefits to households (Jacobson et al., 2013).

Annual kerosene consumption (K_i) for lighting in GAINS region i in year y was estimated by using the following expression

$$K_{i,y} = \left(\frac{POP_{i,y}}{HS_{i,y}} \right) (1 - ele_{i,y}) * 365 \sum_{j=1}^n (N_{i,j,y} h_{i,j,y} CV_k f_{i,j,y} SC_j), \quad (4)$$

where, POP represents population, HS household size, ele electrification rate, f share of device type j (either wick lamps or hurricane lanterns), N number of kerosene lamps, h daily operating hours, SC specific kerosene consumption of a device, and CV_k the calorific value of kerosene.

Field Code Changed

The population data originates from [\(IEA, 2012a\)](#)~~(IEA/ETP, 2012)~~, household size from (UN-Habitat, 2005), the electrification rates from OECD/IEA sources [\(IEA, 2007, 2011, 2012b\)](#)~~(IEA, 2007, 2011, 2012)~~ and national data/reports (ESMAP, 2005; GOI, 2011; NSSO, 2007). For India, information about the share of lighting devices (i.e., wick lamps, hurricane lanterns), operating hours, specific kerosene consumption are derived from regional studies (Desai et al., 2010; Mahapatra et al., 2009; Purohit and Michaelowa, 2008). Reported specific kerosene consumption in kerosene lamps varied from 0.005 to 0.042 litre per hour (e.g., Mills, 2003; Pode, 2010) and we assumed 0.006 and 0.02 litre per hour for wick lamps and hurricane lanterns, respectively. Further, we assumed that each household will use three lamps for 6 hours per day whereas the share of hurricane lanterns is 20 percent for South Asia and 50 percent for other regions.

In India, over 44 percent of rural and about seven percent of urban households reported kerosene as their primary source of lighting in 2004–2005 (NSSO, 2007) and in the lowest four socioeconomic deciles, 60 percent of households use kerosene for lighting (Parikh, 2010). In several of the most populated African countries, including Uganda, Ethiopia, and Kenya, more than 60 percent of the population relies on kerosene as the primary lighting fuel (Apple et al., 2010; IFC/WB, 2008; UBOS, 2010).

Less is known of the quantity of kerosene used for lighting, since it is often difficult to differentiate kerosene used for lighting from that used for other purposes, particularly cooking. The India Human Development Survey 2005 (Desai et al., 2010) results indicate that kerosene lighting accounts for approximately 65 percent (or 5–6 Tg year⁻¹) of residential kerosene consumption in India. Lam et al. (2014) observed that use of kerosene for lighting in electrified homes is substantial (due to intermittent and unreliable electricity supply), constituting an approximately equal share of demand as non-electrified households.

Particulate matter emission factors for kerosene lamps used in this work were derived from Lam et al. (2012). The PM_{2.5} emission factor for kerosene lighting (1.92 g GJ⁻¹) is approximately 13 times higher compared to that for kerosene used for cooking (0.15 g GJ⁻¹), whereas the OC emission factor for kerosene lighting is roughly one third of the kerosene stove. Furthermore, particulate emissions from kerosene lamps are mostly BC (~92%) (Lam et al., 2016).

3.3 Diesel generators

At a global scale, diesel generator (DG) sets are not a large source of pollution but locally, and especially in the developing world, they could be responsible for a significant share of air pollutant emissions, especially nitrogen oxides and black carbon. DG set are the prevailing option for backup power in facilities where continuous power is essential, based on their combination of reliability, durability, affordability, and overall efficiency (Shah et al., 2006). While increasing power deficit and instabilities in the electricity market resulted in rapid growth of the DG set market in several developing regions, DG have been in use all over the world as backup power facilities, primary electricity generation sources in small remote areas or at initial development stage of industrial plants, for irrigation purposes, etc. The DG sets range from small engines to large generators, are operated on very variable fuel quality, and the emission limit values have been typically lagging behind those for mobile engines.

Field Code Changed

Field Code Changed

There is no direct statistical data on fuel use in DG sets as their consumption is typically part of the energy use reported within power plants, commercial, and potentially agricultural sector. Therefore, fuel consumption was estimated from data on number and size of diesel generators as well as regional studies. The resulting fuel use was compared [to the IEA statistics for and calibrated to the diesel consumption reported in](#) the power and commercial sector [and adjusted if necessary so that the overall energy use is consistent with the IEA.](#)

According to a market review in India, annual DG sales in 2010 were about 150,000 units and they are likely to grow at a rate of about seven percent (Frost & Sullivan, 2010) driven by chronic power shortages and prolific growth in industries, infrastructure, telecommunication, information technology (IT), and IT enabled services. The DG market spans from small (15 – 75 kVA) to large (375.1 – 2000 kVA) sets with estimated diesel consumption of about 5 to 6 billion litres between 2008 (Anand, 2012) and 2010⁹. This represents about 8-9%¹⁰ of total diesel consumption (Anand, 2012; NIELSEN, 2013) and in peak periods up to 18% or even more in some regions (NIELSEN, 2013). In Nepal, electricity deficit has been estimated recently at almost 50% (NEA, 2012) massively increasing dependency on diesel generators. The share of diesel used for DG sets in Nepal is estimated at 15 percent for 2010 (World Bank, 2014a). In Nigeria, total electricity demand is estimated at between 8,000 and 10,000 MW while supply from the national grid is about 4,500 MW, which results in very heavy reliance on DG sets operating most times between 15 – 18 hours a day (Triple E., 2013; World Bank, 2014b). For South Asia (except Nepal), Cambodia, Indonesia and Myanmar we have used the Indian share of diesel consumption in DG sets whereas in other developing countries, the share of diesel use for DG sets is assumed to be one fourth of the Indian share due to high electrification rates and relatively low power deficit. For sub-Saharan Africa, due to very high power deficit (up to 50 percent), in some regions we have used the share of diesel use in DG sets from Nepal (World Bank, 2014a).

For South Korea, diesel consumption in DG sets was less than 0.2 percent of total diesel consumption (KEEI, 2011). In EU-28, the share of diesel consumption in DG sets is less than 0.4% of the total diesel consumption; however, the share of heavy fuel oil (HFO) use in DG sets is more than 3% of the total HFO used in EU. Similarly, in United States and Japan the share of diesel consumption is small while the share of HFO is approximately 0.5% and 2%, respectively.

Stationary DG sets are frequently operated in harsh conditions and until recently were rarely subject to emission regulation.

Information on DG set emissions factors is fairly limited and not necessarily representative for all regions. GAINS model emission factors were developed on the basis of data reported in a number of studies (Anayochukwu et al., 2013; Corbett and Koehler, 2003; Gilmore et al., 2006; Lee et al., 2011; Lin et al., 2008; Shah et al., 2004, 2006; Shi et al., 2006; Tsai et al., 2010; Uma et al., 2004; US EPA, 1996). While it is possible to achieve emissions reductions from diesel combustion through engine modifications [and](#) post-combustion [measures, we assume that in the period 1990-2010 DG sets operating in the developing world lack any such controls. In case new information will become available and for future implementation of respective policies, the GAINS model includes a number of post-combustion](#) control technologies such as diesel particulate filters (DPFs), diesel oxidation catalysts (DOC), and fuel-borne catalysts (FBC) [offering an array of options for mitigation or](#)

⁹ <http://ppac.org.in/>

¹⁰ <http://www.nipfp.org.in/newweb/sites/default/files/Diesel%20Price%20Reform.pdf>

~~elimination or reduction of~~ gaseous and particulate emissions, ~~and can be utilized for both on- and off-road applications~~ (Herzog, 2002; Yelverton et al., 2016). Shah et al. (2007) observed that DOC and DOC+FBC technologies were effective in reducing mainly organic carbon (OC) emissions (56-77%) while DPFs showed excellent performance in reducing both elemental carbon (EC) and OC emissions (>90%). The emission factors and shares of BC and OC in particulate mass emissions from measurement literature together with the range of values used in the GAINS model are presented in Table S3.1 in SI.

3.4 Transport

Globally, the transport sector, including international shipping, is estimated to contribute about 10% of total anthropogenic PM₁₀ and PM_{2.5} emissions and up to 25% of BC (Table 8). At a regional level, the role of transport in BC emissions varies strongly and, for example, in Europe and North America was estimated at over 60% in 1996 (Bond et al., 2004) and about 50% in 2005 (Kupiainen and Klimont, 2007) and 2010 in this study, while for East Asia its share grew from about 8 to 23% between 1990 and 2010 (this study). The key source of PM emissions in the transport sector is exhaust emissions from diesel engines with typically light- and heavy-duty trucks playing the largest role; Europe is an exception as policies favouring diesel fuels, in terms of both tax rates and emission limits, resulted in large a share of diesel cars (Cames and Helmers, 2013). Non-exhaust emissions (brake, tyre, and road wear) represent a relatively small share, especially for carbonaceous particles, but their importance grows over time owing to ever more stringent exhaust emission limits.

The overall energy consumption in the transport sector was taken from Eurostat (EUROSTAT, 2011) statistics for the 28 European Union (EU) member states and from the International Energy Agency (IEA, 2015a, 2015b) for all other countries. Fuel consumption of road vehicles is allocated to the different vehicle types through triangulation with data on the active fleet, their average annual mileage, and their average fuel efficiency. The IEA statistics provide fuel consumption figures separately for rail, aviation, and domestic shipping, however, not for mobile machinery used in agriculture, forestry, industry, and construction and mining sectors. Unless national information is available, as is the case for European countries, the US and Canada, we re-allocate 80% of diesel fuel consumption from the IEA categories “industry” and “agriculture” to construction and agricultural machinery, respectively. International shipping and aviation are not included in the GAINS model but were estimated for the ECLIPSE project separately; see section 3.4.2.

There is a vast literature on PM measurements of internal combustion engines used in road vehicles in both developing and developed countries, including also pre-regulation vehicles (e.g., Cadle et al., 2009; Cheung et al., 2009; Geller et al., 2006; Kirchstetter et al., 1999; Liu et al., 2009; Subramanian et al., 2009; Yanowitz et al., 2000). ~~For all world regions we assume that a certain fraction of vehicles is badly maintained (e.g., Mancilla et al., 2012) or their emission controls tampered which~~ Also old and often poorly maintained vehicle fleet is reflected in measurements of emission factors ~~(e.g., Mancilla et al., 2012) as well~~ as the share of so-called high-emitters (McClintock, 1999, 2007; Smit and Bluett, 2011; Yan et al., 2011, 2014); see further discussion in section 3.4.1. For Europe and the USA we draw the emission factors for road vehicles from

Formatted: English (U.S.)

Formatted: English (U.S.)

established emission factor models where experts already synthesized the information (HBEFA 3.1, 2010; Ntziachristos et al., 2009; US-EPA OTAQ, 2011). These emission factors are adjusted to conditions in other world regions.

Kupiainen and Klimont (2004, 2007), Bond et al. (2004), Maricq (2007) are examples of studies which summarized and compared emission factors for various vehicle categories. Most of exhaust PM is emitted in a submicron range, actually

5 within 100 nm, and diesel vehicles typically emit several times more (mass-based) PM than equivalent gasoline engines (e.g., Maricq, 2007); exceptions are old vehicles running on leaded gasoline and pre-regulation 2-stroke mopeds (Klimont et al., 2002b; Kupiainen and Klimont, 2004) while latest gasoline direct injection engines have PM mass emissions comparable or even higher than latest diesel engines with particle filter. It is important to highlight that properly functioning particulate

10 filters reduce PM emissions significantly however and consequently, the absolute level of the latest diesel vehicles is about two orders of magnitude lower than for older generations. The carbonaceous particles represent the largest share with the elemental carbon fraction higher for diesel (50–70%) than for gasoline vehicles (30–40%) (e.g., Kupiainen and Klimont, 2007; Maricq, 2007). Non-exhaust emissions, i.e., brake and tyre wear as well as road abrasion were updated based on (Denier van der Gon et al., 2013; EEA, 2013; Harrison et al., 2012). Recent roadside measurements showed that tyre wear produces essentially coarse particles, with only a small contribution (<0.5%) in the PM_{2.5} size fraction (Stein et al., 2012).

15 Road abrasion emissions significantly increase when studded tyres are used, e.g., a common practice in Scandinavian and some Baltic countries. Higher abrasion during winter and spring conditions, average usage period, and application shares are factored into the average abrasion emission factor for the Nordic countries (Kupiainen et al., 2005; Kupiainen and Pirjola, 2011).

20 PM emission factors for the diverse non-road mobile machinery are much less well established, and only seldom available for developing countries. Moreover, most measurements refer to the mandatory duty cycles rather than real-life operating conditions. For Europe and North America we use emission factors based on (EEA, 2013; OTAQ, 2004; Schäffeler and Keller, 2008) and transfer to other world regions assuming that technology performs similarly and the under comparable operating conditions.

25 The contribution from diesel engines used in agriculture, construction, mining, rail, shipping, and as back-up generators has been increasing, not least because the emission legislation lags behind that for road transport, but has been receiving more attention recently (e.g., Kholod et al., 2016). Diesel generators and shipping are discussed in separate sections (3.3 and 3.4.2), more recent emission factors for diesel locomotives (e.g., Johnson et al., 2013; Tang et al., 2015) are compared with GAINS in Table S4.3, and emission factors for other non-road machinery used in GAINS were summarized earlier (Klimont et al., 2002b; Kupiainen and Klimont, 2004, 2007) and are also included in the supplementary information (SI). Emission
30 factors for key diesel and gasoline engines in the transport sector from recent literature and the GAINS model are compared in Tables S4.1 to S4.5.

3.4.1 High-emitting vehicles

On-road remote sensing measurements of vehicles suggest that a relatively small fraction of the fleet is responsible for a relatively large fraction of emissions (e.g., Ban-Weiss et al., 2009; Cadle et al., 1997; Mazzoleni et al., 2004; Subramanian et al., 2009). In the literature, these vehicles have been referred to as: high emitters or high-emitting vehicles, heavy emitters, super emitters, gross emitters, excess emitters or smokers, but in principle highlighting the same problem (Shafizadeh et al., 2004). Reasons for their poor emission performance are variable and can be traced back to malfunctioning or totally inoperative emission controls, low combustion efficiency of the engine, engine oil that is entering the combustion chamber, and/or leakage in the exhaust system between the engine and the emissions control devices (Jimenez et al., 2000; Mazzoleni et al., 2004; Norris, 2001). The shares of high emitters and their contribution to total fleet emissions are variable across countries, with for instance only limited evidence in Europe for light duty vehicles (Borken-Kleefeld and Chen, 2015; Chen and Borken-Kleefeld, 2016) and more modern vehicles seem to have more durable emission controls (McClintock, 2007). Though there is no doubt in the existence of high emitting vehicles, quantifying their emissions is much more speculative.

According to Shafizadeh et al. (2004) two general definitions of high emitters can be identified from the literature: a group of vehicles that (i) account for a certain fraction, e.g., 50 percent, of air pollutant emissions, or (ii) have emissions above a certain emission threshold or cut-off. The GAINS estimation of high emitter emissions is based on the second general definition. The calculation requires two sets of information: (i) the amplification factor which is the ratio between the high and normal emitter emission factors, and (ii) the share of high emitters in the whole vehicle fleet.

The technology-specific amplification factors, i.e., for Euro 1 to 6, were developed based on existing studies mainly from the United States (Ban-Weiss et al., 2009; Durbin et al., 1999; Hsu and Mullen, 2007; Yanowitz et al., 2000) and Europe (Carslaw et al., 2011; Ekström et al., 2004) studying the 90-95th percentile as the cut-off between high and normal emitting behaviour. Similar datasets from Australia (Smit and Bluett, 2011), China (Guo et al., 2007), Thailand (Subramanian et al., 2009) and Mexico City (Jiang et al., 2005) were also studied in order to find which percentiles would represent the local fleets if the amplification factors identified, based on the 90-95th percentiles in the European and US studies, would be applied also there. The identified percentiles then determined what share of the vehicle fleet corresponded to the amplification factors specified for the high emitting vehicles. A global coverage of the parameterization was developed using the available studies and databases listed above as benchmarks representative for larger groups of countries and regions. We acknowledge that this definition of the high emitting vehicle class is based on a statistical analysis only and currently does not have a technical definition. However, the motivation of the exercise is to single out a portion of the vehicle fleet that might emit significantly more than the majority of the fleet and study the potential importance of such vehicles in total emissions. The amplification factors determined from the studies varied between pollutants, vehicle types and fuels. Table 6 demonstrates the derived amplification factors for light and heavy duty on-road vehicles that apply for all countries and all PM species, following the observations reported by Subramanian et al. (2009). We have noted the results by Lawson (2010)

who showed that the OC/BC ratio might be different for high emitters than for normal vehicles but have not introduced variable ratios for individual vehicle categories.

The default assumptions about the high-emitter shares are: about 5% for the EU-28, Japan, and Korea; 8% for Australia, Canada, and US; 5-10% for non-EU Europe, 12% for China (except some key cities with more modern fleet where 10% is assumed); 15% for India, and 20% for other developing Asia, Africa, Latin America. These assumptions are compatible with those used in other global studies (e.g., Bond et al., 2004, 2007, Yan et al., 2011, 2014). In addition we factor in that the durability of the emission controls have increased. Therefore we assume that failure rates decline for the more modern technologies, i.e., above the equivalent of Euro 4, which translates to halving the percent of high emitters for such vehicles. For example, for Europe or Japan for most recent years this results in a lower overall rate of about 2%, which is consistent with assessments for the US and Europe (Chen and Borken-Kleefeld, 2014; McClintock, 2007).

3.4.2 International shipping and aviation

Particulate matter emissions from international shipping contribute about 3-4% of the global total, and while, unlike for SO₂ and NO_x, this is a rather small share, it is also comparable to the contribution of road transport (e.g., Lack et al., 2009). Aviation contributes only a very small proportion of global PM emissions, e.g., for black carbon its share was estimated at about 0.1-0.2% (Lee et al., 2009; Stettler et al., 2013) of which about 14% were during landing and take-off (LTO) (Stettler et al., 2013).

The GAINS model does not include emissions from these sources and the gridded ECLIPSE datasets V3 and V4a refer to other sources, e.g., datasets developed for the RCP process (Buhaug et al., 2009; Eyring et al., 2010; Lee et al., 2009). However, the more recent ECLIPSE sets (V5 and V5a) include international shipping estimates developed using activity data from Buhaug et al. (2009); fuel consumption data for 2007 were extrapolated to 2010 using GDP. Our extrapolation for 2010 produced fuel consumption similar to the average estimated for the period 2007-2012 (Smith et al., 2015) but larger by about 10% than IMO estimate for 2010 (Smith et al., 2015). Emissions are estimated for all PM species (the RCP set contains only BC and OC) using emission factors shown in Fig. 52 and spatially distributed drawing on the QUANTIFY project¹¹, i.e., based on global ship traffic data (Endresen et al., 2007). The fuel consumption data includes assumptions about region-specific regulation with respect to fuel quality, i.e., sulphur content of fuels.

The shipping PM emission and their chemical, physical, and optical properties have been analysed for various types of fuels, engines, and vessels, as well as operating conditions, e.g., load factors (Agrawal et al., 2008, 2010, Lack et al., 2008, 2009; Moldanova et al., 2009; Murphy et al., 2009; Petzold et al., 2008, 2010). Further studies reviewed and compared emission factors (Buhaug et al., 2009; Dalsoren et al., 2009; Lack and Corbett, 2012). The particulate matter emission profile, including BC and OC, presented in Fig. 52, was developed on the basis of the studies listed above.

Field Code Changed

Field Code Changed

¹¹ <https://www.pa.op.dlr.de/quantify/>

3.5 Large scale combustion

Solid fuel combustion in large boilers used in power plants and industry has been a major source of primary particulate matter emissions and although efficient reduction technology exists and is typically required by law, about 15% of total global anthropogenic PM_{2.5} emissions in 2010 originated from this source. At the same time, since 1990's emissions declined by over 30% and its share dropped from over 20% to 15%. Primary PM from combustion can be divided into two major categories: (i) ash, formed from non-combustible mineral constituents in fuel which vary from few to over 30% depending on fuel quality, and (ii) carbonaceous particles, e.g., char, coke and soot, which are formed by pyrolysis of unburned fuel molecules (e.g., Seinfeld and Pandis, 2012). The largest particles remain in the boiler and are removed with bottom ash while smaller (typically <100 µm) are entrained in combustion gas forming fly ash. Emissions of elemental and organic carbon from such installations are small due to high combustion temperature, oxidizing conditions, and long residence times (e.g., Ohlström et al., 2000); only about 2% of global total black carbon was estimated to originate from this source (Bond et al., 2004, 2013; Cofala et al., 2007).

The principal statistical data for energy use in the power sector and industry used in GAINS originates from International Energy Agency (IEA, 2015a, 2015b), Eurostat (EUROSTAT, 2011), and national sources, especially for Europe (e.g., Amann et al., 2012, 2015) and Asia (e.g., Amann et al., 2008; Purohit et al., 2010; Zhang et al., 2006; Zhao et al., 2013). The national sources and consultations were especially useful to distribute fuel use among different types of plants; see discussion in section 2.2.3.

The PM emission factors in GAINS are calculated considering region-specific fuel properties (heat value, ash content), installation-specific parameters (ash retention in boiler, size distribution), size-specific efficiency of control equipment (cyclones, wet scrubbers, electrostatic precipitators, fabric filters); see Eq. 1, Eq. 2, and discussion in section 2.1. Detailed review of measurement studies, methodology and assumptions applied in GAINS has been documented in a number of earlier reports and papers (Klimont et al., 2002b, 2009, Kupiainen and Klimont, 2004, 2007; Zhang et al., 2006; Zhao et al., 2013). Key updates with respect to emission factors have been done for Europe within the work for the European Commission (Amann et al., 2015) and China, where latest information about efficiency and penetration of control measures was used (Zhao et al., 2013).

3.6 Industry

There are many industrial processes that emit particulate matter to the atmosphere and the origin of these emissions is often more complex than that of stationary combustion since there are several process stages, fugitive sources, and the process designs vary significantly across the world. The particular processes will also differ with respect to emission characteristics, i.e., PM size distribution and chemical speciation. The GAINS model distinguishes tens of industrial processes, including several within the iron and steel sector, non-ferrous metals, cement and lime, petroleum refining, coal mining, gas flaring, and production of bricks, coal briquettes, mineral fertilizers, glass, carbon black, and pulp. Extensive discussion of these

Field Code Changed

sources, including their particulate matter and carbonaceous aerosols emissions and mitigation measures in GAINS is available from previously published reports (Klimont et al., 2002b; Kupiainen and Klimont, 2004). The estimates presented in this paper rely for most sectors on the characteristics presented in those reports, however with updated emission factors for a number of regions and specifically a new structure of the three sectors mostly relevant for carbonaceous particles, i.e., coke ovens, brick making, and gas flaring.

While there are well established PM control technologies applicable to most of the sources (Klimont et al., 2002b; Kupiainen and Klimont, 2004; Maithel et al., 2012) and typically, even in the developing world, there exists legislation prescribing emission limit values, this sector remains among the most uncertain in terms of emission estimation of total PM as well as carbonaceous aerosols. We estimate that at a global scale industrial processes contributed between about 13 and 20% of PM_{2.5} emissions in 1990 and 2010 and total emissions grew in this period by over 60%. Regional shares might be much larger, e.g., for China were estimated at over 30% in 2010 and grew by nearly a factor of three compared to 2000, or significantly lower, e.g., for Africa less than 5%. For most regions, key PM_{2.5} sectors include cement and iron and steel production representing globally about 75% of industrial emissions of PM_{2.5}. For carbonaceous particles, this sector plays a slightly less important role from the global perspective; Bond et al. (2004) estimated its contribution at about 13% to BC emissions, primarily from coking and brick making. This is broadly consistent with our assessment, although we estimate a somewhat lower share of about 10% globally, of which about a third comes from gas flaring, and there is very strong regional variation from less than a percent to over 20%, especially in regions with high oil production, e.g., Middle East, Russia.

The principal statistical data used in GAINS originates from international sources (Elvidge et al., 2009; EUROSTAT, 2011; IEA, 2015a, 2015b), and national sources, especially for Europe (e.g., Amann et al., 2012, 2015) and Asia (e.g., Amann et al., 2008; Heierli and Maithel, 2008; Huo et al., 2012; Purohit et al., 2010; Zhang et al., 2006; Zhao et al., 2013).

The PM emission factors used in the GAINS model have been discussed in previously published reports (Klimont et al., 2002b; Kupiainen and Klimont, 2004) and key updates concern the region-specific primary technology allocation and implementation rates of control technologies – as discussed in section 2.2.4 and 2.3. For coke manufacturing (see 3.6.1), brick production (see 3.6.2), and gas flaring in oil and gas industry (see 3.6.3) more significant changes were introduced with new technology and region-specific emission factors.

3.6.1 Coke production

Total coke production grew by about a factor of two in the 1990-2010 period and most of the change took place after 2000 when China increased their production by about a factor four from just over a 100 Tg to about 400 Tg coke which represented over 60% of global production in 2010 (Huo et al., 2012) and see <http://www.statista.com>. China's coke sector undergoes a significant transformation moving from low efficiency and high emission indigenous ovens to highly mechanized recovery ovens, following the world trend (Huo et al., 2012; Polenske, 2006). Several of the other producing countries remained fairly constant or reduced their output in the last decade, e.g., US, Europe, former Soviet Union region,

Field Code Changed

and only few increased their production, e.g., India, but from the global perspective these changes were not very significant (<http://www.statista.com>).

There are only few measurements of PM emissions from coke plants and the established emission factors show a wide range.

This is partly driven by the varying technology but also owing to the sources of emissions from coke manufacturing since

5 they include several stack and fugitive sources. In the GAINS model, we have constructed a PM emission profile based on the US EPA Compilation of Air Pollutant Emission Factors (AP-42)¹² and SPECIATE¹³ (US EPA, 1995, 2002) as discussed in Klimont et al. (2002b) and Kupiainen and Klimont (2004), and updated it with more recent measurements discussed in Huo et al. (2012) and Weitkamp et al. (2005). For uncontrolled ovens, GAINS emission factors for PM_{2.5} range from about

10 2- to 4.8 kg t⁻¹ coke, the upper bound being representative for China and the range reflecting different oven types across the global regions. For BC and OC, the emission factor range is 0.28-1.3 kg t⁻¹ and 0.46-2.2 kg t⁻¹, respectively, with upper range values representing Chinese indigenous ovens. The PM emission factors for China are comparable to the ones used in recent Chinese studies (Huo et al., 2012; Lei et al., 2011) and the ratio of BC/OC of about 0.6 is also consistent with the

15 estimates by Weitkamp et al. (2005). Owing to lack of specific data for various world regions, we assume little change in emissions factors over time for the developing world, although the transition in China reported in Huo et al. (2012) was considered, and for OECD countries emission factor trend follows reported emissions, where available.

3.6.2 Brick kilns

The brick making industry is dominated by production in the developing countries where over 95% of global output, estimated at about 1.5 trillion bricks per year (e.g., Schmidt, 2013), is produced and most of it in fairly inefficient and polluting kilns. In India, over 70% of kilns, or about 100,000, are clamp kilns, the least efficient kiln that remains widespread

20 in the developing world. More than 1.2 trillion bricks per year are produced in Asia alone which is associated with use of over 100 million tons of coal as well as other fuels including agricultural residues, dung, and waste (Heierli and Maithel, 2008; Schilderman and Mason, 2009). The largest brick producing countries in Asia are China, India, Pakistan, Bangladesh and Vietnam (AIT, 2003; FAO, 1993; Heierli and Maithel, 2008; Maithel, 2014). Worldwide non-automated brick

25 production, including artisanal brick kilns, in developing countries is about 1.25 trillion bricks per annum and is distributed between three main regions (i) China – about 700 billion bricks or 56%, (ii) India – about 150 billion bricks or 12%, (iii) Asia, Africa, South America & Mexico – about 400 billion bricks or 32%. In contrast, worldwide machine made brick production using automated kilns, is approximately 125 billion bricks, with Australia's brick production accounting for only 2 billion, UK 4 billion, USA 8 billion, China 100 billion, and other developed countries approximately 11 billion bricks. A summary of studies used to compile the brick production data is provided in SI (S5) along with the activity data used in

30 ECLIPSE V5a for key global regions (Table S5.2).

¹² <https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emission-factors>

¹³ SPECIATE is the US EPA repository of volatile organic gas and particulate matter (PM) speciation profiles of air pollution sources: <https://www.epa.gov/air-emissions-modeling/speciate-version-45-through-32>

Even though from the global perspective, the brick manufacturing sector does not represent a major share of particulate matter emissions, about 1-2% for PM_{2.5} according to our estimates and less than 5% for BC (e.g., Bond et al., 2004, 2013), the regional impacts might be much more significant (Guttikunda et al., 2013; Le and Oanh, 2010; Skinder et al., 2014). And while many countries may have emissions standards, i.e., maximum permissible concentrations of several pollutants, including PM, the enforcement is difficult for several reasons including relatively few measurements available. Maithel et al. (2012), Weyant et al. (2014), and Rajarathnam et al. (2014) reported particulate matter measurements for key brick kiln production technologies in Asia (primarily India and Vietnam), and a few studies, focusing on toxics and black carbon, were performed in Mexico (Cardenas et al., 2012; Christian et al., 2010; Maíz, 2012); the latter covered several types of kilns including the Marquez kiln (MK) that is specific to Latin America. For the main brick producing technologies in South Asia, the PM emission factors derived from the above measurements are lower by over 30% for BC and 90% for PM_{2.5} than previously estimated values (Weyant et al., 2014), which were used in several regional (Klimont et al., 2009; Lu et al., 2011; Ohara et al., 2007; Zhang et al., 2009) and global inventories (Bond et al., 2004; Cofala et al., 2007; UNEP/WMO, 2011). Additionally, the BC/TC ratio appears higher than previously thought (Weyant et al., 2014).

The emission factor set used in GAINS to calculate ECLIPSE set is more in line with the currently available measurements although it was developed prior to the publication of measurements by Weyant et al. (2014); compare Table S5.1 in the SI, where current GAINS emission factors for PM_{2.5}, BC, and OC are compared with the previous GAINS dataset and recent measurements by Weyant et al. (2014). Also the EC/TC ratio in GAINS, from about 0.67 for zig-zag, about 0.75 for clamps, downdraft, moving chimney BTK, and 0.93 for fixed chimney BTK, resembles the measurements by Weyant et al. (2014)

3.6.3 Gas flaring

Understanding venting, flaring, and associated gas utilization practices in the oil industry has been of high relevance for the assessment of methane emissions while it was not considered as a potentially important source of air pollution. Consequently, non-CO₂ emissions from flaring of associated gas in oil industry were not part of previous inventories (e.g., Bond et al., 2013), including the datasets used in the IPCC assessments. We have developed the first global estimate of air pollutant emissions from this activity, including black carbon, which was used first in the studies focusing on the role of black carbon and other short-lived climate forcers in climate mitigation (Bond et al., 2013; Shindell et al., 2012; UNEP, 2011; UNEP/WMO, 2011; World Bank and ICCI, 2013). Within the ECLIPSE project, an update and future mitigation scenarios (Klimont et al., in preparation) were developed and used in several regional and global modelling exercises (Stohl et al., 2013, 2015).

Associated petroleum gas (APG) is gas that is associated with the oil in the reservoir and once oil is extracted, the dissolved gas follows and is commonly separated from the oil and either vented or flared. The volumes and composition of APG depend on several factors including the nature of the oil reservoir, degree of depletion, etc. (PFC Energy, 2007; Røland, 2010). While the APG could be utilized, the lack of developed markets, missing infrastructure, no legislation, etc. resulted in very low recovery rates before 1980; only in the last decades has the flaring trend been decoupled from oil production but

the level of gas utilization varies greatly among the producing regions. Globally, about 140-160 billion m³ (bcm) APG have been flared annually, which represents about 5% of 2009 global natural gas consumption or about 30% of European Union demand (Elvidge et al., 2009). Regions where the largest volumes of gas are flared include Middle East, Russia, Northern Africa, Nigeria, Venezuela representing about 70-80% of the global total (Elvidge et al., 2009, 2013). There are significant uncertainties in estimates of flared volumes as metering is rare and official estimates differ significantly from remote sensing data or even between different official versions, e.g., for Russia governmental sources reported for 2006 about 15-20 bcm of APG flared while Global Gas Flaring Reduction Initiative (GGFR) estimates were about 40-60 bcm (PFC Energy, 2007). The reported share of APG flared in Russia in 2006 varied from 27% (governmental sources) to 75% (NGOs) with 45% estimated by PFC Energy (2007) (Røland, 2010). For Nigeria, flaring volumes have been estimated or reported between 10 to 25 bcm indicating that up to 70% of APG is flared (Aghalino, 2009; Ite and Ibok, 2013). While for several countries APG utilization rates have been increasing (Elvidge et al., 2009; Haugland et al., 2013), Russia made relatively little progress [until 2010](#) in recent years in spite of new legislation requiring a 95% recovery rate (PFC Energy, 2007; Røland, 2010). For US, flaring volumes increased by about a factor three between 2006 and 2011 owing to the boom in unconventional gas and oil production (Elvidge et al., 2013). GAINS activity data relies on the time series of gas flaring volumes developed within the GGFR initiative (Elvidge et al., 2007, 2011)

There is a very limited number of measurements of flaring emissions allowing the establishment of a representative set of emission factors where local flare operating conditions and APG properties could be considered. Some of the earlier published PM emission factors (about 2.6 g m⁻³) referred to landfill (CAPP, 2007) or refinery flares (US EPA, 1995) and are generally considered inappropriate. A new technique for quantitatively measuring soot emission rates in flare plumes under field conditions has been reported by Carlton University group (Johnson et al., 2011) and while their average BC emission factor of 0.51 g m⁻³ (McEwen and Johnson, 2012) considers representative fuel mixtures, their measurements were performed on laboratory-scale flares, which might underestimate real-world emissions. The first ECLIPSE datasets include flaring emissions calculated with one BC emission factor of 1.6 g m⁻³ gas flared, assuming that real-life flares perform much worse than laboratory measurements. In the later ECLIPSE set V5a, region-specific PM emission factors were developed considering a more recent study measuring emissions from flares in the Bakken region (Schwarz et al., 2015) which confirmed the order of magnitude measured by McEwen and Johnson (2012) by establishing upper bound BC emission factor of 0.57±0.14 g m⁻³. We have assumed that such emission rates are representative for well operated flares, i.e., OCED countries. For other countries we retained the previously used value of 1.6 g m⁻³ but considered, where available, composition of flared gas that apart from methane includes several heavier hydrocarbons. The relationship between BC emission factors and heat value of flared gas has been proposed by McEwen and Johnson (2012) and was also applied in estimates for Norway (Aasestad, 2013) and Russia (Huang et al., 2015).

The range of current BC emission factors in GAINS is ~0.5-1.75 g m⁻³, the upper bound represents values for Russia, and the estimated heat value of APG varied from about 41 to 50 MJ m⁻³. Huang et al. (2015) suggested even higher BC emission factors for Russia (2.27 g m⁻³) assuming local APG composition with estimated heat value of about 75 MJ m⁻³ and

extrapolating linearly from the relationship from McEwen and Johnson (2012); well beyond the range presented there. Finally, the most recent measurements of BC from flaring, also in Bakken field, estimate much lower overall emission factors of $0.13 \pm 0.36 \text{ g m}^{-3}$ and characterize flares without visible smoke (Weyant et al., 2016) and therefore, likely not representative for regions with visible high-density smoke, e.g., Russia, Nigeria, Middle East, Northern Africa (e.g., Aghalino, 2009; Elvidge et al., 2013; Pederstad et al., 2015). We assume that all PM from flaring is $\text{PM}_{2.5}$ and BC and OC represent about 78% and 16%, respectively. These assumptions are broadly consistent with the results of McEwen and Johnson (2012) who reported BC/OC share of 80/20 and Fortner et al. (2012) measuring 4-20% of OC and over 95% of PM within $\text{PM}_{2.5}$.

3.7 Agricultural waste burning

Bond et al. (2004) estimated that globally about 7 and 15% of anthropogenic (excluding forest and savannah fires) BC and OC emissions originated from this source in 1996; our own estimates point to a slightly lower share in carbonaceous particles emissions but mostly because our total, not agricultural burning, estimates are higher. At the same time, for several regions this source might be even more important, e.g., for Brazil we estimate its contribution at up to 15% of $\text{PM}_{2.5}$ and 10% of BC emissions. Finally, agricultural burning has a strong seasonal pattern (see also section 2.4.1.) and ~~These emissions have~~ been also linked with heavy smog and haze episodes (e.g., Mukai et al., 2015; Stohl et al., 2007).

Typically assessment of global emissions from open field burning of agricultural residues is based either on a compilation of national reports/sources (e.g., Bond et al., 2004; EC-JRC/PBL, 2010) or on remote sensing data which characterize the magnitude and spatial distribution of open biomass burning including agricultural, savannah, and forest fires (e.g., van der Werf et al., 2010; Wiedinmyer et al., 2011), however, it has been shown the latter underestimates small open fires (e.g., Randerson et al., 2012). Niemi (2007) compared various datasets for all open biomass sources and developed the first global activity set for the RAINS model drawing on EDGAR3.2FT2000 (Van Aardenne et al., 2005) which we have ~~This database has been~~ further extended and updated to accommodate other data sources allowing gaps to be filled for several countries. Specifically, we have used estimates from the global studies (Bond et al., 2004), a number of regional estimates (Cao et al., 2008; Oanh et al., 2011; Pettus, 2009), reporting of emissions to EMEP (<http://www.ceip.at>), and bilateral discussions within the revision of the European air pollution policy (Amann et al., 2015). Our global estimate of open burning of agricultural residue has been fairly constant in the assessment period varying from about 485 to 515 Mt between 1990 and 2010; this estimate is comparable with 475 Mt for 1996 by Bond et al. (2004) and higher than the original EDGAR3.2FT2000 of 252 Mt of residue burned in 2000.

To derive particulate matter emission factors, we have relied on Akagi et al. (2011), Andreae and Merlet (2001), Turn et al. (1997), and Hegg et al. (1997); the latter for the OM/OC ratio which we assumed 1.7 as discussed in Kupiainen and Klimont (2004). The default emission factors used in GAINS (all values in g kg^{-1}) are 8.5 for TSP, 7.1 for PM_{10} , 6.3 for $\text{PM}_{2.5}$, 5.6 for PM_1 , 2.62 for OC and 0.83 g kg^{-1} for BC. Using data from Turn et al. (1997), these values were adjusted for specific

regions considering typical type of crops; for example, for regions with a high share of rice production (primarily Asia) the values of BC and OC factors were estimated at 0.6 and 2.2 g kg⁻¹.

3.8 Waste

Open burning of solid waste is a widespread method, especially in the developing world, to reduce the volume or odours of dumped or uncollected municipal solid wastes (EAWAG, 2008) and it has been identified as significant source of particulate matter and hazardous air pollutants to the atmosphere (Christian et al., 2010; Hodzic et al., 2012; Kumar et al., 2015; Wiedinmyer et al., 2014). The estimated magnitude of emissions and contribution to PM concentrations vary widely across the studies, ranging from a few percent to nearly 50% of the total contribution in particular regions. While large uncertainties remain owing to only few measurements and difficulties in finding reliable data on waste collection, recycling, and disposal rates, the open burning of residential waste is a potentially important source of PM, especially in the developing world.

To estimate the region-specific share of the municipal solid waste (MSW) that is burned, we used a mass balance approach described in the IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006b). As a starting point, we used the IPCC reported data on MSW generation and management and assumed that the category “*other MSW management, unspecified*” represents the upper limit for the open burning of residential waste. However, the IPCC values were not used directly in many cases, because the IPCC unspecified fractions are in some cases relatively high, up to 60 percent, and also because not all unspecified mass is necessarily burned. We have additionally used information on percentages of commonly used MSW disposal methods in other studies (CEPMEIP, 2002; EAWAG, 2008; Neurath, 2003); the final fraction of open burning from the total waste produced in the developed world was estimated to vary between 0.5 and 5% and for the developing world the region-specific fractions were estimated at 10-20%. The GAINS model estimate of the global MSW is about 1500 to 2150 Tg in the period 1990 to 2010 of which about 115 to 160 Tg were estimated as openly burned. While the total waste generation rate is consistent with other studies (e.g., Christian et al., 2010; Wiedinmyer et al., 2014), the open burned fraction differs significantly owing to different assumptions about the fraction burned and practices in urban and rural areas. For example Bond et al. (2004) and Wiedinmyer et al. (2014) estimated that 33 and 970 Tg of waste are burned; the latter is still about six times larger than GAINS. We were not able to consider the results of Wiedinmyer et al. (2014) in GAINS yet, but a comparison at the national level shows that GAINS has significantly lower estimates for most of the developing countries as well as Europe; for the latter, GAINS is consistent with national reporting and often a factor 5 to 10 lower than Wiedinmyer et al. (2014). For the US and Canada GAINS has a factor of 2-3 higher estimates (also consistent with the US EPA and Environment Canada).

The PM emission factors used in GAINS were derived from Akagi et al. (2011) and Christian et al. (2010) and are consistent with the ones used by Wiedinmyer et al. (2014). These are (all in g kg⁻¹) 9.5 for PM₁₀, 8.74 for PM_{2.5}, 6 for PM₁, 5.27 for OC, and 0.65 g kg⁻¹ for BC.

3.9 Other sources

The GAINS model also includes several other sources of PM which at a larger scale represent rather small contribution but could be of relevance locally. These are mostly non-combustion (fugitive) emission sources and include animal livestock, storage and handling of bulk industrial and agricultural products, arable land related agricultural activities, and construction works. Additionally, emissions from cigarette smoking, barbeques, and fireworks are considered. Note that windblown dust and emissions from unpaved roads are not included (see also introduction to section 3).

The predominant sources of PM from animal housing include feed and faecal material, bedding, skin, hair, mould, and pollen. Size-specific PM emission factors were developed in GAINS drawing on the results of measurements done in Europe (e.g., ICC & SRI, 2000; Louhelainen et al., 1987; Takai et al., 1998) which are discussed in more detail in Klimont et al. (2002b). The values presented in that report were adapted considering region-specific length of the housing period (time animals spend indoors) which is a regional parameter in the model, also relevant for estimation of ammonia emissions. For dairy cows the PM₁₀ factors range from 0.22-0.43 kg animal⁻¹ per year, for beef 0.11-0.43 kg animal⁻¹, for poultry about 0.05 kg animal⁻¹, and for pigs 0.4-0.45 kg animal⁻¹. The share of PM_{2.5} is about 22% with the exception of pigs where it was estimated at about 17%; no BC or OC emission factors were assumed. Emissions from arable farming include harvesting, ploughing, tilling, etc. The GAINS PM₁₀ emission factor varies from 0.8 to 2 kg ha⁻¹ and the PM_{2.5} is assumed to represent about 22% of PM₁₀. These revised numbers, compared to the earlier GAINS values discussed in Klimont et al. (2002b), draw on the more recent work in Germany and France discussed within the EU air quality consultation (Amann et al., 2015).

Emissions from storage and handling of bulk industrial (coal, iron ore, fertilizers, cement, other) and agricultural products as well as from construction activities are estimated using emission factors discussed in Klimont et al. (2002b). For the latter, some updates were made based on national consultations within work on the revision of the EU air quality policy (Amann et al., 2015) and the recent range for PM₁₀ is 0.07 – 0.22 Gg per million m² of constructed floor space, with a share of PM_{2.5} assumed at 12% and no primary carbonaceous particles.

For cigarette smoking we assume a PM_{2.5} emission factor of 0.01 – 0.0165 kg capita⁻¹ (equal to PM₁₀) and a share of BC and OC as 0.5% and 60%, respectively (Klimont et al., 2002b). Also for barbeques, a per capita emission factor is established, i.e., 0.02 – 0.075 kg capita⁻¹ with a share of BC and OC assumed as about 15% and 50%, respectively (Klimont et al., 2002b). Only very few regional estimates were available for these sources, specifically identified within the discussion in Europe (Amann et al., 2015), therefore for most countries the same emission rates are used.

4 Results and discussion

Global, regional and sectoral emissions of particulate matter (PM) distributed into several size bins (PM₁₀, PM_{2.5}, PM₁) as well as into black and organic carbon are shown in Table 7-8 for 2010 and Fig. 63-74 for the period 1990-2010; Table S6.2-S6.6 in the SI show global emissions of PM species for 25 global regions in the period 1990-2010. To our knowledge, these estimates represent the first global dataset of anthropogenic emissions where size-specific mass PM calculation, including

BC and OC, was performed using a uniform and consistent estimation framework. Emissions are also allocated into a $0.5^\circ \times 0.5^\circ$ (longitude-latitude) grid and available freely for a number of datasets¹⁴. Finally, the PM estimates are consistently linked with the emissions of other air pollutants and greenhouse gases for the same time period as well as their future projections developed with the GAINS model (Klimont et al., in preparation).

5 Total emissions of particulate matter (including open burning based on GFED3.1 database but excluding windblown dust) in 2010 are estimated at about 111 Tg for PM₁₀, 81 Tg for PM_{2.5}, 71 Tg for PM₁, 9.5 Tg for BC, and 33 Tg of OC. Anthropogenic contribution dominated all species except OC and OM, i.e., about 55% of PM₁, PM_{2.5}, and PM₁₀, 75% of BC, and 40% for OC and OM (Table 7). For all considered PM species, sources in Asia represented over 60% of the global anthropogenic total (Table 7) with residential combustion being the most important sector although its share declines with
10 increasing particle size: about 60% for BC and OC, 45% for PM_{2.5} and less than 40% for PM₁₀ for which large combustion sources and industrial processes are equally important (Table 8).

In contrast to several local and regional atmospheric modelling studies, the global modelling community has been relying so far on the assumption that anthropogenic PM_{2.5} emissions are sufficiently well represented by the sum of black carbon and primary organic PM, often referred to as POM. This total fine PM mass hasis been typically estimated as $BC+1.4*OC$ ¹⁵ and
15 only recently a number of models included more detailed aerosol schemes accounting for varying BC/OC ratios while still largely neglecting the anthropogenic dust component (e.g., Philip et al., 2017). Combining such estimates with windblown dust and open biomass fires to arrive at the total PM_{2.5} might be sufficient from the perspective of global climate impacts of primary PM aerosols; however, the health impacts could be severely underestimated in some regions where the non-carbonaceous share of anthropogenic fine particulate matter is significant (Fig. 63).

20 We argue that assessment of health impacts due to PM using results of the global emission projections developed in the first place for climate simulations, e.g., Representative Concentration Pathways (RCP), which included anthropogenic BC and OC, windblown dust, and open fires but not the non-carbonaceous component of primary PM_{2.5} and PM₁₀ emissions originating from combustion, industrial processes, and some fugitive sources, might lead to inconsistent results and underestimation of PM concentrations and regional impacts. This study provides the first global assessment of the role non-carbonaceous particle emissions play in total anthropogenic PM₁, PM_{2.5}, and PM₁₀ mass emissions and could prove more
25 appropriate to use in global modelling studies of health impacts as well as climate. Moreover, while at the global level, the ratio of anthropogenic emissions of PM₁ and PM_{2.5} to (BC+POM) is about 1.3 and over 1.6, there are important differences between the regions and the emission ratios have been changing over time (Fig. 63). For example, in 2010 we estimate for Asia an emission ratio of two for PM_{2.5}/(BC+POM) while for North America the same ratio is about 1.5 (Fig. 63, Table 7).
30 In Europe, including Russia, this ratio has changed from about three in the early 1990's, where primary PM emissions from poorly controlled coal power plants and heavy industry (not a large source of carbonaceous particles – compare Fig. 74) dominated the total, to below two in 2010 (Fig. 63). Even when the emissions from open biomass burning (forest and

¹⁴ http://www.iiasa.ac.at/web/home/research/researchPrograms/air/Global_emissions.html

¹⁵ The 1.4 has been the most commonly used OM/OC ratio (Aiken et al., 2008)

savannah fires) are taken into account, and most of these occur far from densely populated areas, the total PM_{2.5} mass emissions are over 20% larger than the BC+POM (Table 7).

We estimate that about 75% of global anthropogenic emissions of PM₁₀ are PM_{2.5} and while there was only little change in that ratio (slight increase) in the last decade at the global level, more significant variation has been observed across sectors

(Fig. 74). Combustion of liquid fuels, biomass, and waste produces typically over 90% of PM_{2.5} in PM₁₀ but for several industrial processes, power and industrial boilers burning coal, and coal production, distribution and storage, emissions of PM_{2.5} represent only 40-60%. Carbonaceous particles (BC+OM) emissions play a key role in PM_{2.5} representing over 60% with the largest contribution from residential combustion (about 80%) and transport and agriculture (each about 10%).

Nearly 90% of PM_{2.5} emissions from residential boilers and cooking and heating stoves is BC+OM of which over 20% is BC. Similarly high share of BC+OM is estimated for transport sector but it varies between about 95% for road transport and 80% for non-road vehicles, however, the share of BC is much larger than for residential combustion: 35 – 45% of PM_{2.5} emissions from transport (including non-exhaust) is BC. Few of the smaller sources, agricultural residue and trash burning, have also large share of BC+OM (over 80%) but rather small contribution of BC. Combustion of solid fossil fuels in power and industrial boilers as well as most industrial processes (except brick manufacturing in traditional kilns and possibly coke making) are characterized by very low share of carbonaceous particles (below 5%).

4.1 Regional distribution and temporal trends

Total anthropogenic emissions of PM_{2.5} and BC in 2010 have a similar spatial distribution (Fig. 85). Emission densities are generally the highest in Asia, however, there are some important differences in the contributions of various sectors to both species as well as across regions. Residential combustion plays a key role but appears far more important for BC where it represents nearly 60% of the global total (Table 8) and an even higher share for Asia and Africa; for PM_{2.5} this sector contributes globally about 45%. While for PM_{2.5} the energy and waste sector (incl. agricultural burning) and industry make most of the remaining emissions (25% and 17.5%, respectively), they represent just over 10% of BC emissions (Table 8 and

Fig. 85). Industrial emissions appear much more important in Asia (Fig. 85) and while there are several processes contributing to PM_{2.5} emissions, for BC brick and coke production make the most and represent up to 12% of Asian emissions, globally about 6%. Some sector contribution patterns are similar across continents, for example, for North America, Latin America, and Europe transport and the residential sector dominate BC emissions while for PM_{2.5} it is mostly energy and the waste sector, except Europe where also residential combustion appears important (Fig. 85). For Africa, residential combustion is the key source of all PM with the exception of a few areas like Republic of South Africa or oil producing countries where the energy sector is an important source. It is particularly striking to see the difference in the source contributions to BC emissions in Africa and Asia where the most important source is the residential sector, but while in Africa other sources are barely visible, for Asia there are important contributions from the transport and industry (Fig. 85).

The other feature worth highlighting is the difference in relative importance of the transport sector for PM_{2.5} and BC emissions (about 8% and 24% at a global level, respectively) which is clearly visible in the third row of maps in Fig. 85.

We estimate that global emissions of PM have changed little in the period 1990-2010 showing a strong decoupling from the global increase in energy consumption and consequently CO₂ emissions (Fig. 6.3). However, there are very different regional emission trends with a particularly strong increase in East Asia and Africa and a strong decline in Europe, North America and Pacific. The development of PM₁₀ and PM_{2.5} emissions is fairly similar with a slightly faster growth of PM_{2.5} (+8%) than PM₁₀ (+4%) at a global level. The difference is mostly due to reductions of industrial emissions in Europe and Russia following the political and economic transition in Eastern Europe that started already in the mid-80s. This economic restructuring resulted in closure or transformation of inefficient and polluting heavy industries which in turn brought in about 55 and 60% reduction of PM_{2.5} and PM₁₀ emissions between 1990 and 2010; most of which was achieved before 2000 (Fig. 6.3). Also North American and Pacific emissions declined in this period by about 30%. In contrast, PM₁₀ and PM_{2.5} emissions in East Asia and Africa increased by about 40-50% and those of Other Asia and Latin America by about 10%. The stark differences in regional trends resulted in important changes in the spatial pattern of PM burden. Europe, North American, and Pacific contribution to global emissions dropped from nearly 30% in 1990 to well below 15% in 2010 while Asia's contribution grew from just over 50% to nearly 2/3 of the global total in 2010 (Fig. 6.3, Table 7, Table S6.2-S6.3). For black carbon (BC), the regional changes were less dramatic but the global emissions are estimated to grow by about 15% by 2010 compared to 1990, mostly driven by increases in Asia (about 30%) and Africa (over 40%) (Fig. 6.3, Table 7, Table S6.5-S6.6). BC emissions in Europe, North America, and Pacific declined by about 30% but their share in the global total are estimated at below 15% in 2010 (from about 24% in 1990).

4.2 Comparison with other studies

This is the first assessment of the global anthropogenic emissions of PM₁₀, PM_{2.5}, and PM₁ using a consistent bottom up approach across all the sources and regions and therefore only limited comparison to other work at a global level can be made. In fact, the only global set where PM₁₀, PM_{2.5}, BC, and OC were published is the so called 'mosaic inventory', developed within the UNECE Task Force on Hemispheric Transboundary Air Pollution (HTAP) where a compilation of EDGAR and several regional inventories were put together (Janssens-Maenhout et al., 2015) for 2010. For most of the species the HTAP_v2 is lower than ECLIPSE V5a by about 20-30% except OC where the agreement is good (Table S8.1 in SI). It is difficult to easily conclude on the reasons for observed differences as the methods are not fully comparable and HTAP_v2 is a compilation where single products rely on different methods. However, as further discussion shows, the largest discrepancy for PM₁₀ and PM_{2.5} is for China as well as Europe and Russia; the sum of the differences in these three regions combined they represent about 90% and over 50% of all the difference for PM₁₀ and PM_{2.5}. There have been a number of global studies of BC and OC emissions as well as several regional assessments of PM₁₀, PM_{2.5}, BC, and OC which we discuss in a more detail below.

A seminal work by Bond et al. (2004) established a benchmark global inventory of BC and OC emissions for the year 1996 that was later updated to 2000 (Bond et al., 2013) and was also used as the basis for the development of BC and OC emission in the RCP scenarios (Lamarque et al., 2010; Van Vuuren et al., 2011). Bond et al. (2004) provided a thorough review of BC

and OC estimates to date and has been used as the primary reference since. We compare our results with Bond et al. (2004, 2013) in Table 9 and Fig. 96 for 1995 and 2000. At a global level, recent GAINS calculation (V5a) shows higher values, which is mostly due to inclusion and re-estimation of few sources: kerosene wick lamps, gas flaring, use of regional coal statistics for China; Fig. 96 shows the role of these sources in GAINS estimates for 2000, ECLIPSE version totals (see also Fig. S6.1 in SI), and compares them to the range presented in Bond et al. (2013). Even though the global totals fall within the same range, especially when considering the role of newly calculated emissions from kerosene lamps (version V4a did not include them), there are often larger differences at a source-sector level, particularly for residential combustion where largest uncertainties exist in fuel consumption, its allocation between uses and technologies, and emission factors (Table 9). Excluding kerosene lamps and gas flaring, which were not included in Bond et al. (2004, 2013), GAINS global estimates are larger by less than 5% and 15% for 1995 and 2000 than Bond et al. (2004, 2013). This difference is mostly due to residential sector where comparable source categories are larger in GAINS by 40-60% but the overall balance is partly offset by emissions from industrial coal use (including coke and brick production as well as industrial boilers) that are larger in Bond et al. (2004, 2013) (Table 9).

Emission characteristics for kerosene lamps, gas flaring, and diesel generators have been included in GAINS only recently (most of the previously published global work has not included these sources). For kerosene wick lamps we followed on the work of Lam et al. (2012) but developed an independent assessment of activity data and estimated global BC emissions from this source at 706 Gg in 2005. Our estimates are higher than the previous assessment of 270 Gg (Lam et al., 2012) and 580 Gg (Jacobson et al., 2013) because of larger kerosene consumption in our study, but compare well to Elisabeth (2013) who calculated 702 Gg BC from this activity. For gas flaring we estimated global BC emissions at about 270 Gg and 210 Gg in 2005 and 2010. A recent study of flaring emissions for Bakken field (Weyant et al., 2016) extrapolated their results to global estimates of 20 ± 6 Gg BC, assuming the same range of emission factors as measured by them at the Bakken field. This is over ten times less than our estimates but we argue that the Bakken flares are not necessarily representative for some of the other regions where strong variability and potentially high soot emissions have been shown by (Conrad and Johnson, 2017; Johnson et al., 2011)(Conrad and Johnson, in review; Johnson et al., 2011) and also speculated in Huang et al. (2015). We found no global estimates of PM emissions from diesel generators and our estimate of 113 Gg for $PM_{2.5}$ and 50 Gg for BC in 2010 confirms that it appears to be a rather small source from a global perspective, and although, important locally, it is expected that in the near future with reliable access to grid electricity use of DG sets will be limited particularly in residential, commercial and industrial sectors.

Granier et al. (2011) compared global and regional estimates of BC developed within global and regional modelling activities or inventories for the period 1980-2010. We compare the range presented in that study with the inventory used during development of RCP scenarios (Lamarque et al., 2010) and the GAINS model calculation for version V5a, highlighting the role of the newly included and re-estimated sources (Fig. 107). At a global level, the GAINS range overlaps with the span of estimates presented in other studies, although the GAINS total is actually higher than all previous estimates and the post 2000 trend is also different, implying a slight increase in emissions rather than a decline or stabilization shown

Field Code Changed

in earlier studies; note that values reported in Granier et al. (2011) for 2010 were results of projections. As shown in comparison to Bond et al. (2004, 2013) (Table 9), the GAINS values are higher primarily due to inclusion of kerosene lamps and gas flaring but also because of more recent statistical data for 2010 than used in the previously published work. Fig. 107 also includes results of selected global and regional studies which were not explicitly referred to in Granier et al. (2011); these are marked with 'black star' symbols and included in Table S8.1 in SI. The values for 1996 and 2000 refer to the Bond et al. (2004, 2013) and for 2010 to HTAP_v2 inventory (Janssens-Maenhout et al., 2015), none of which included emissions from kerosene wick lamps.

Fig. 107 shows also a similar comparison for selected countries: China, India, and US; note that the ranges presented in Granier et al. (2011) for regions/countries do not necessarily add up to the global total as the former included also selected regional studies which were not part of the comparison of the global totals. For China, a continuing growth in BC emissions has been reported in all investigated studies. GAINS is comparable with the RCP input (Lamarque et al., 2010) for 1990-1995, while for the last decade is consistently higher or at the top of the range, which in Granier et al. (2011) is representative of the upper estimates in the RCP scenarios rather than specific inventories. However, a number of recently published studies for China reported rather high BC, e.g., Zhang et al. (2009) estimated about 1.8 Tg for 2006, HTAP_v2 (based on the MEIC¹⁶ system developed by the Tsinghua University (Beijing, China)) 1.76 Tg for 2010, Lu et al. (2011) 1.84 Tg for 2010, and 1.92 Tg for 2008 using a top-down approach (Kondo et al., 2011); these results and other recent regional studies are marked with 'black star' symbols in Fig. 107 and included in Table S8.1. Several authors estimated also PM₁₀ and PM_{2.5} emissions for China and these compare reasonably well with GAINS, although are systematically lower by up to 15% with exception of the HTAP_v2 mosaic inventory (Janssens-Maenhout et al., 2015) which is lower by nearly 25% for 2010 (Table S8.1 in SI); the latter inventory relies on the data from the MEIC system where more optimistic assumptions about the penetration and achieved efficiency of wet scrubbers and electrostatic precipitators in industry are made. For India, all inventories suggest emissions have been increasing in the investigated period but there is a very large spread of estimates.

Current GAINS estimates are higher than Lamarque et al. (2010) and the range shown by Granier et al. (2011) (Fig. 107) – the overlap in the last decade is because the upper values are based on the earlier GAINS model estimates (e.g., Klimont et al., 2009) which are consistent with ECLIPSE set. Some recent papers have shown similar BC emissions as GAINS (e.g., Janssens-Maenhout et al., 2015; Lu et al., 2011; see also Table S8.1) but overall the range of published emission estimates for PM species for India varies greatly between studies, e.g., for BC from about 350 Gg to over 1000 Gg (Table S8.1). A lot of that variability links to different assumptions about biomass use for cooking (Venkataraman et al., 2005), efficiency of PM abatement in power and industry, large uncertainty in agricultural burning activity (Venkataraman et al., 2006). For the US, all studies indicate a declining trend in BC emissions (Fig. 107). However, in contrary to China and India, GAINS emissions are in the lower range of existing estimates (Fig. 107, Table S8.1) and difference in emissions from non-road machinery and agricultural (or prescribed) burning appears to be the key reason for observed discrepancies.

¹⁶ MEIC - Multi-resolution Emission Inventory for China; <http://www.meicmodel.org>

For Europe (including European part of Russia), the published studies of BC and OC (Bond et al., 2004; Kupiainen and Klimont, 2007; Schaap et al., 2004; see Table S8.1) compare well showing differences within $\pm 10\%$ or less with exception of EDGAR (Janssens-Maenhout et al., 2015) which shows much lower emissions but does not include any Russian territory. At the level of whole of Europe, GAINS calculates similar PM_{10} and $PM_{2.5}$ emissions as officially reported to UNECE LRTAP Convention (www.ceip.at) while EDGAR estimate is nearly 40% lower for both species, however, does not include Russia (Table S8.1). There have been only few published estimates of PM emissions in Russia (Table S8.1). For PM_{10} and $PM_{2.5}$ in 2010, GAINS calculates higher emissions than EDGAR (Janssens-Maenhout et al., 2015) or national inventory submitted to LRTAP Convention (www.ceip.at) which covers only European part of Russian Federation; remarkably the total EDGAR estimate is similar to the national submission for European part. The main reasons for discrepancy are significantly larger GAINS emissions from industrial processes, residential combustion (these are very low in the national submission – less than a quarter of EDGAR and GAINS estimates), agricultural burning, as well as inclusion of gas flaring. The uncertainties in volume of gas flared and actual emission factors are major reasons for the difference in estimated BC emissions in GAINS and Huang et al. (2015) who derived a much higher emission factor for this activity; for other sectors both studies report a fairly similar emissions of BC for 2010.

Yan et al. (2011) developed projections of PM_{10} emissions from road transport sector (exhaust only). Their PM_{10} estimates for 2000-2010 were about 1.65-1.75 Tg with a contribution from high emitters of about 0.3 Tg. The ECLISPE *V4a* results are comparable Yan et al. (2011), while in *V5* and *V5a*, updates to the emission factors (reflecting more recent measurements, poor fuel quality, and maintenance) and penetration rates of control measures for developing countries (often delayed or postponed implementation of legislation) led to higher estimates of about 2.4-2.6 Tg, including high emitters (0.4-0.5 Tg). Total GAINS model estimates for road transport also include non-exhaust emissions (brake, tyre, road abrasion) which add up to around 0.6 Tg PM_{10} .

Wiedinmyer et al. (2014) developed a new assessment of global emissions from burning of waste, including particulate matter. That study suggests that all current estimates largely underestimate emissions from this activity. Compared to GAINS, their emissions are nearly seven times larger and would make open burning of waste one of the key categories contributing between 10-15% of BC and $PM_{2.5}$ and nearly 30% of OC; considering anthropogenic sources. For example, waste burning could be responsible for three times more emissions of BC, OC, and $PM_{2.5}$ than agricultural waste burning or about a third of the total transport sector emissions. Current GAINS estimates of 2010 emissions from open waste burning are about 1.4, 1.3, 0.1, 0.75 Tg PM_{10} , $PM_{2.5}$, BC, OC, while Wiedinmyer et al. (2014) calculated for the same species 12, 12, 0.632, 5.1 Tg. Obviously, large uncertainties remain in activity data and actual emission factors (see discussion in section 3.8) but this activity deserves more attention in the future.

4.3 Uncertainty in emission estimates

The completeness and quality of information about emission inventories varies across the regions, sectors, and species. The underlying information about several key PM sources like residential solid fuel combustion, brick production, and residual

waste burning is often of poor quality or non-existing and that applies to both activity data and emission factors. In order to create a comprehensive emission data set, the national information is often supplemented with model estimates that rely on default parameterization; in fact, even many of the national inventories draw on the international data sets of emission factors (e.g., EEA, 2013; US EPA, 1995) owing to lack of local measurements. Finally, the level of enforcement of existing laws, as well as the real-life performance of control technology is seldom sufficiently well-known and we tend to assume rather optimistically that both deliver and work as planned which has been shown to be often false (e.g., Stoerk, 2016; Xu et al., 2009; Xu, 2011)(e.g., Stoerk, 2016; Xu, 2011; Xu et al., 2009), as, more recently, in the so-called Dieselgate affair (e.g., Lange and Domke, 2015; US EPA, 2015a, 2015b). Consequently, the level of uncertainty, or confidence, varies widely across source sectors and regions.

We have not performed a formal uncertainty analysis for emission estimates in this study, but results of analysis from other studies are helpful and indicative of the expected uncertainties for various species and regions. For example, the global BC and OC inventory developed by Bond et al. (2004) included an uncertainty analysis of total emissions providing regional 'low-high' estimates for 1996. For BC emissions from anthropogenic sources, the range was 3.1-10 Tg yr⁻¹ (-30% to +120%) and for OC 5.1-14 Tg yr⁻¹ (-40% to +130%). Estimates from the GAINS model presented in this study sit well within these ranges. ~~The TRACE-P emission inventory for Asia (Streets et al., 2003) estimated the uncertainties for emission of carbonaceous particle (BC and OC) as 95% confidence intervals to be 160-500% for the developing countries and 80-180% for developed countries in Asia (Streets et al., 2003).~~

As indicated earlier, emissions of PM, including carbonaceous aerosols, belong to the most uncertain among the air pollutants as they form usually under poor combustion conditions in small inefficient installations burning poor quality fuels, which brings variability to the emission characteristics. Additionally there is very little information globally about local emission factors. Considering local data and knowledge about emission sources and their emission factors could significantly reduce uncertainties (Zhang et al., 2009). Allocating total PM emissions into different size bins or chemical species (here BC and OC) is associated with uncertainties that for a specific source are determined by the measurement. Among others, Bond et al. (2013) discussed specific issues related to BC and OC aerosols, while for PM size distribution there exist specific analysis for particular measurement equipment (e.g., Armas et al., 2007; Coquelin et al., 2013) and most of the studies reporting measurements of size distribution estimate uncertainties for each size category. While the sum of all the PM species is constrained by the total mass, the single size distribution values rely on a large number of measurements reducing the overall uncertainty. Exceptions are source-sectors for which very few measurements exist, e.g., coke ovens, fireworks, handling of bulk materials.

In addition to the emission characteristics, the activity data also is a source of uncertainty. While for major industrial and transport sectors there are well documented and regularly updated national and international sources of activity data (e.g., IEA, 2015a, 2015b), the activities behind the major PM source categories, for example poor quality fuels in cook stoves or brick kilns, as well as local vehicle fleets, are not well known. For commercial fuels, however, the uncertainty has been estimated to vary from 2-3% for OECD countries to 5-10% for non-OECD (IPCC, 2006a).

Field Code Changed

A significant part of total aerosol emissions originate from open biomass burning, including forest fires, savannah, and agricultural residue burning (e.g., Reddington et al., 2015). Estimation of activity data and actual emission factors are bound with significant uncertainties which include, among others, amount of biomass burned and interannual variability (Chen et al., 2013; van der Werf et al., 2006; Wiedinmyer et al., 2011), drivers and impact of change in agricultural fires (Morton et al., 2008), and emission factors (Castellanos et al., 2014). The uncertainty ranges estimated by Bond et al. (2004) for BC and OC emissions from open biomass burning were 1.6 to 9.8 Tg yr⁻¹ (-45% to +185%) for BC and 31 to 58 Tg yr⁻¹ (-40% to +110%) for OC.

The uncertainties of emission estimates developed with integrated assessment models like GAINS are similar to the estimates for bottom-up inventories discussed above, at least at a regional scale. ~~In fact, they could be even lower considering that they typically rely on a harmonized data set and include a simultaneous calculation of emissions of several species using the same principal activity and technology data.~~ Additionally, the error compensation, which is especially relevant if calculated emissions are the sum of a large number of equally important source categories (and where the errors in input parameters are not correlated with each other), can lead to a further reduction of overall emission uncertainty (Schöpp et al., 2005). A careful assessment of the assumption about correlation between input parameters is essential as for example poor enforcement of legislation or measurements errors could affect several source sectors in a similar way. The GAINS model uncertainties, calculated in Schöpp et al. (2005), are consistent with the values reported by Streets et al. (2003) for developed countries. This analysis has also shown that at a finer scale the understanding of local circumstances are critically important to reduce uncertainty, and while the emission factors were estimated to be the key factor determining uncertainty in historical emissions, at least for aerosol emissions, the uncertainty in activity assumptions becomes more important for the uncertainties in projected emissions.

5 Conclusions

To our knowledge, the estimates represent the first global dataset of anthropogenic emissions where size specific mass PM calculation, including BC and OC, was performed using a uniform and consistent estimation framework including a number of previously unaccounted or often misallocated emission sources, i.e., kerosene lamps, gas flaring, diesel generators, trash burning that have been systematically evaluated for each region. Spatially, emissions were calculated for 170 regions and allocated to 0.5° x 0.5° longitude-latitude grids and are available either from the on-line GAINS model¹⁷, where assumptions and results can be displayed for 25 global regions (see section S7 in SI) or gridded emissions can be downloaded from the project website¹⁸. The ECLIPSE datasets do not include independent estimates of emissions from forest fires and savannah burning, windblown dust, and unpaved roads.

¹⁷ <http://magcat.iiasa.ac.at/gains/IAM/index.login>

¹⁸ http://www.iiasa.ac.at/web/home/research/researchPrograms/air/Global_emissions.html

We estimate that global emissions of PM have not changed much between 1990 and 2010 but there are significantly different regional trends with North America, Pacific, and Europe reducing emissions by 30 to over 50% and Asia and Africa increasing by about 30%. While these regionally varying developments are clearly visible in PM_{2.5} and PM₁₀ estimates, the BC regional changes were somewhat less dramatic, mostly because trends in power and industrial sector emissions of PM are much less relevant for total black carbon emissions. Our new global estimate of BC emissions suggests higher numbers than previously published owing primarily to inclusion of new sources. Globally, over 75% of anthropogenic PM₁₀ and PM_{2.5} originates from residential combustion, power plants and industry while for BC residential combustion and transport represent more than 75% but the importance varies across regions with Europe and North America having transport as key and rest of the world residential combustion. Our new global estimate of BC emissions suggests higher numbers than previously published owing primarily to inclusion of new sources.

We argue that this PM estimate reduces the gap in source coverage required in air quality and climate modelling studies and health impact assessments at a regional and global level as it includes both carbonaceous and non-carbonaceous constituents of primary particulate matter emissions, however, additional efforts need to be made to address several fugitive sources of anthropogenic dust, e.g., unpaved roads. The ECLIPSE emission data sets have been used in several regional and global atmospheric transport and climate model simulations (AMAP, 2015; Eckhardt et al., 2015; Gadhavi et al., 2015; Lund et al., 2014; Quennehen et al., 2016; Stohl et al., 2013, 2015; Wobus et al., 2016; Yttri et al., 2014) where various aspects of several particulate matter species were addressed. The emissions developed during ECLIPSE also served as basis for a recently published global particulate number estimates (Paasonen et al., 2016).

We envisage development of further datasets drawing on the experience of the ECLIPSE exercise. The future versions will be available via the same on-line platform where additional documentation will be placed too. As a matter of fact, the GAINS model and the ECLIPSE dataset and scenarios have been already used as a starting point to develop emission data and mitigation strategies for the recently published International Energy Agency (IEA) World Energy Outlook special report on air pollution (IEA, 2016). Furthermore, the elements of the ECLIPSE data have been part of the contribution towards improved representation of carbonaceous aerosols in the large scale integrated assessment models used in the development of the Shared Socio-economic Pathways (SSP). (O'Neill et al., 2014; Rao et al., 2017; Riahi et al., 2017). ~~(O'Neill et al., 2014; Rao et al., in press; Riahi et al., in press)~~

Acknowledgements: The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no 282688 – ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants). We acknowledge the funding received from UNEP under the Small Scale Funding Agreement (SSFA) [SLP 2294-2H73-1111-2261] which allowed to improve the resolution of the GAINS model in Latin America and the Caribbean. We would like to thank Prof Qiang Zhang from Tsinghua University (Beijing, China) for the spatial distribution of Chinese power plants in 2000, 2005, and 2010 (that information was used in developing gridded emission fields) as well as Imrich Bertok and Robert Sander from IIASA for dedicated database programming support.

Formatted: Font: 10 pt

Field Code Changed

References

[Aasestad, K.: Emissions of Black carbon and Organic carbon in Norway 1990-2011, Statistics Norway, Oslo, Norway., 2013.](#)

[Adria, O. and Bethge, J.: What users can save with energy- efficient cooking stoves and ovens Oliver Adria \(CSCP\) Jan Bethge, Wuppertal Institute for Climate, Environment and Energy, Wuppertal, Germany., 2013.](#)

[Aghalino, S.: Gas flaring, environmental pollution and abatement measures in Nigeria, 1969-2001, Journal of Sustainable Development in Africa, 11, 219–238, 2009.](#)

[Agrawal, H., Malloy, O. G. J., Welch, W. A., Wayne Miller, J. and Cocker III, D. R.: In-use gaseous and particulate matter emissions from a modern ocean going container vessel, Atmospheric Environment, 42\(21\), 5504–5510, 2008.](#)

[Agrawal, H., Welch, W. A., Henningsen, S., Miller, J. W. and III, D. R. C.: Emissions from main propulsion engine on container ship at sea, J. Geophys. Res., 115, 7 PP., doi:201010.1029/2009JD013346, 2010.](#)

[Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U. and Jimenez, J. L.: O/C and OM/OC Ratios of Primary, Secondary, and Ambient Organic Aerosols with High-Resolution Time-of-Flight Aerosol Mass Spectrometry, Environ. Sci. Technol., 42\(12\), 4478–4485, doi:10.1021/es703009q, 2008.](#)

[AIT: Small and Medium scale Industries in Asia: Energy and Environment. Brick and Ceramic Sectors, Regional Energy Resources Information Center \(RERIC\), Asian Institute of Technology \(AIT\), Pathumthani, Thailand., 2003.](#)

[Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D. and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, Atmospheric Chemistry and Physics, 11\(9\), 4039–4072, doi:10.5194/acp-11-4039-2011, 2011.](#)

[Amann, M., Bertok, I., Borken, J., Chambers, A., Cofala, J., Dentener, F., Heyes, C., Kejun, J., Klimont, Z., Makowski, M., Matur, R., Purohit, P., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F. and Winiwarter, W.: GAINS-Asia. A tool to combat air pollution and climate change simultaneously, International Institute for Applied Systems Analysis \(IIASA\), Laxenburg, Austria., 2008.](#)

[Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguven, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F. and Winiwarter, W.: Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications, Environmental Modelling & Software, 26\(2\), 1489–1501, 2011.](#)

[Amann, M., Borken-Kleefeld, J., Cofala, J., Heyes, C., Klimont, Z., Rafaj, P., Purohit, P., Schoepp, W. and Winiwarter, W.: Future emissions of air pollutants in Europe - Current legislation baseline and the scope for further reductions. TSAP Report #1, International Institute for Applied Systems Analysis, Laxenburg. \[online\] Available from: <http://gains.iiasa.ac.at/images/stories/reports/TSAP/TSAP-BASELINE-20120613.pdf> \(Accessed 13 December 2012\), 2012.](#)

[Amann, M., Klimont, Z. and Wagner, F.: Regional and Global Emissions of Air Pollutants: Recent Trends and Future Scenarios, Annual Review of Environment and Resources, 38\(1\), 31–55, doi:10.1146/annurev-environ-052912-173303, 2013.](#)

[Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Kiesewetter, G., Klimont, Z., Schöpp, W., Vellinga, N. and Winiwarter, W.: Adjusted historic emission data, projections, and optimized emission](#)

Formatted: Bibliography, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

reduction targets for 2030 – A comparison with COM data 2013, IIASA, Laxenburg, Austria. [online] Available from: http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf, 2015.

5 [AMAP: AMAP Assessment 2015: Black carbon and ozone as Arctic climate forcers, Arctic Monitoring and Assessment Programme \(AMAP\), Oslo, Norway. \[online\] Available from: <http://www.amap.no/documents/doc/amap-assessment-2015-black-carbon-and-ozone-as-arctic-climate-forcers/1299>, 2015.](#)

[Anand, M.: Diesel Pricing in India: Entangled in Policy Maze, National Institute of Public Finance and Policy \(NIPFP\), New Delhi. \[online\] Available from: <http://www.nipfp.org.in/media/medialibrary/2013/04/Diesel%20Price%20Reform.pdf>, 2012.](#)

10 [Anayochukwu, A. V., Nnene, E. A. and Onyeka, A. E.: Assessment of Environmental Impact of Power Generation in Banking Industry, Journal of Energy, Environment & Carbon Credits, 3\(1\), 13–21, 2013.](#)

[Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, Global Biogeochemical Cycles, 15\(4\), 955–966, doi:10.1029/2000GB001382, 2001.](#)

15 [Anenberg, S. C., Schwartz, J., Shindell, D., Amann, M., Faluvegi, G., Klimont, Z., Janssens-Maenhout, G., Pozzoli, L., Van Dingenen, R., Vignati, E., Emberson, L., Muller, N. Z., West, J. J., Williams, M., Demkine, V., Hicks, W. K., Kuylenstierna, J., Raes, F. and Ramanathan, V.: Global Air Quality and Health Co-benefits of Mitigating Near-Term Climate Change through Methane and Black Carbon Emission Controls, Environ Health Perspect, 120\(6\), 831–839, doi:10.1289/ehp.1104301, 2012.](#)

[Apple, J., Vicente, R., Yarberr, A., Lohse, N., Mills, E., Jacobson, A. and Poppendieck, D.: Characterization of particulate matter size distributions and indoor concentrations from kerosene and diesel lamps, Indoor Air, 20\(5\), 399–411, 2010.](#)

20 [Armas, O., Gómez, A. and Herreros, J. M.: Uncertainties in the determination of particle size distributions using a mini tunnel–SMPS system during Diesel engine testing, Measurement Science and Technology, 18\(7\), 2121, 2007.](#)

[Ban-Weiss, G. A., Lunden, M. M., Kirchstetter, T. W. and Harley, R. A.: Measurement of Black Carbon and Particle Number Emission Factors from Individual Heavy-Duty Trucks, Environmental Science & Technology, 43\(5\), 1419–1424, doi:10.1021/es8021039, 2009.](#)

25 [Boman, C., Petterson, E., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 1: Pellet Stoves, Energy Fuels, 25\(1\), 307–314, doi:10.1021/ef100774x, 2011.](#)

30 [Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H. and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, J. Geophys. Res., 109\(D14\), 1–43, doi:10.1029/2003JD003697, 2004.](#)

[Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G. and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850–2000, Global Biogeochem. Cycles, 21\(2\), 1–16, doi:10.1029/2006GB002840, 2007.](#)

35 [Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Bernsten, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G. and Zender, C. S.: Bounding the role of black carbon in the climate](#)

5
system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.

Bonjour, S., Adair-Rohani, H., Wolf, J., Bruce, N. G., Mehta, S., Pruss-Ustun, A., Lahiff, M., Rehfuess, E. A., Mishra, V. and Smith, K. R.: Solid fuel use for household cooking: country and regional estimates for, *Environ Health Perspect*, 121(7), 784–790, doi:10.1289/ehp.1205987, 2013.

Borken-Kleefeld, J. and Chen, Y.: New emission deterioration rates for gasoline cars – Results from long-term measurements, *Atmospheric Environment*, 101, 58–64, doi:10.1016/j.atmosenv.2014.11.013, 2015.

10
Brohan, P., Kennedy, J. J., Harris, I., Tett, S. F. B. and Jones, P. D.: Uncertainty estimates in regional and global observed temperature changes: A new data set from 1850, *Journal of Geophysical Research: Atmospheres*, 111(D12), n/a–n/a, doi:10.1029/2005JD006548, 2006.

Buhaug, O., Corbett, J. J., Endresen, O., Eyring, V., Faber, J., Hanayama, S., Lee, D. S., Lee, D., Lindstad, H., Markowska, A. Z., Mjelde, A., Nelissen, D., Nilsen, J., Palsson, C., Winebrake, J. J., Wu, W.-Q. and Yoshida, K.: Second IMO GHG study 2009, IMO, London, UK., 2009.

15
20
Cadle, S., Ayala, A., Black, K., Graze, R., Koupal, J., Minassian, F., Murray, H., Natarajan, M., Zhu, F., Takaoka, M., Oshita, K., Morisawa, S., Tsuno, H., Takaoka, M., Oshita, K., Morisawa, S., Tsuno, H., Kitajima, Y., Chiang, W.-F., Fang, H.-Y., Fang, H.-Y., Wu, C.-H., Huang, C.-J., Huang, C.-J., Chang, C.-Y., Chang, Y.-M., Chen, C.-L., Nielsen, A., Nielsen, L., Feilberg, A., Christensen, K., Liu, Y.-Y., Lin, T.-C., Lin, T.-C., Wang, Y.-J., Ho, W.-L., Yanowitz, J., McCormick, R., Yu, L., Jia, S., Shi, Q., Chang, T.-J., Kao, H.-M., Wu, Y.-T., Huang, W.-H., Kao, H.-M., Wu, Y.-T., Huang, W.-H., Goodman, P., Rich, D., Zeka, A., Clancy, L., Dockery, D., Lavery, T., Rogers, C., Baumgardner, R., Mishoe, K., Chen, W.-C., Lin, H.-Y., Yuan, C.-S., Hung, C.-H., He, L., Huang, G., Zeng, G. and Lu, H.: Real-World Vehicle Emissions: A Summary of the 18th Coordinating Research Council On-Road Vehicle Emissions Workshop, *Journal of the Air & Waste Management Association*, 59(2), 130–138, doi:10.3155/1047-3289.59.2.130, 2009.

25
Cadle, S. H., Mulawa, P. A., Ball, J., Donase, C., Weibel, A., Sagebiel, J. C., Knapp, K. T. and Snow, R.: Particulate Emission Rates from In-Use High-Emitting Vehicles Recruited in Orange County, California, *Environmental Science & Technology*, 31(12), 3405–3412, doi:10.1021/es9700257, 1997.

CAI-Asia: Factsheet No. 17- Roadmap for Cleaner Fuels and Vehicles in Asia, 2011.

Cames, M. and Helmers, E.: Critical evaluation of the European diesel car boom - global comparison, environmental effects and various national strategies, *Environmental Sciences Europe*, 25(1), 15–36, doi:10.1186/2190-4715-25-15, 2013.

30
Cao, G., Zhang, X. and Zheng, F.: Inventory of black carbon and organic carbon emissions from China, *Atmospheric Environment*, 40(34), 6516–6527, 2006.

Cao, G., Zhang, X., Wang, Y. and Zheng, F.: Estimation of emissions from field burning of crop straw in China, *Chinese Science Bulletin*, 53(5), 784–790, 2008.

35
CAPP: A Recommended Approach to Completing the National Pollutant Release Inventory (NPRI) for the Upstream Oil and Gas Industry, Canadian Association of Petroleum Producers (CAPP). [online] Available from: <http://www.capp.ca/library/publications/policyRegulatory/pages/pubInfo.aspx?DocId=119572>, 2007.

Cardenas, B., Maiz, P., Márquez, R. O., Munguia, J. L., Angeles, F., Baum, E. and Molina, L. T.: Determining Emissions of Black carbon, greenhouse gases and other pollutants from artisanal brick production in Mexico, *Clean Air Task Force and MCE2.*, 2012.

Carslaw, D. C., Beevers, S. D., Tate, J. E., Westmoreland, E. J. and Williams, M. L.: Recent evidence concerning higher NO_x emissions from passenger cars and light duty vehicles, *Atmospheric Environment*, 45(39), 7053–7063, doi:10.1016/j.atmosenv.2011.09.063, 2011.

Castellanos, P., Boersma, K. F. and van der Werf, G. R.: Satellite observations indicate substantial spatiotemporal variability in biomass burning NO_x emission factors for South America, *Atmospheric Chemistry and Physics*, 14(8), 3929–3943, doi:10.5194/acp-14-3929-2014, 2014.

CEPMEIP: CEPMEIP (Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance), [online] Available from: <http://www.air.sk/tno/cepmeip/>, 2002.

Chafe, Z. A., Brauer, M., Klimont, Z., Van Dingenen, R., Mehta, S., Rao, S., Riahi, K., Dentener, F. and Smith, K. R.: Household Cooking with Solid Fuels Contributes to Ambient PM_{2.5} Air Pollution and the Burden of Disease, *Environmental Health Perspectives*, 122(12), 1314–1320, 2014.

Chafe, Z. A., Brauer, M., Héroux, M.-E., Klimont, Z., Lanki, T., Salonen, R. O. and Smith, K. R.: Residential heating with wood and coal: health impacts and policy options in Europe and North America, World Health Organization. [online] Available from: <http://www.euro.who.int/en/health-topics/environment-and-health/air-quality/publications/2015/residential-heating-with-wood-and-coal-health-impacts-and-policy-options-in-europe-and-north-america>, 2015.

Chen, Y. and Borken-Kleefeld, J.: Real-driving emissions from cars and light commercial vehicles – Results from 13 years remote sensing at Zurich/CH, *Atmospheric Environment*, 88, 157–164, doi:10.1016/j.atmosenv.2014.01.040, 2014.

Chen, Y. and Borken-Kleefeld, J.: NO_x Emissions from Diesel Passenger Cars Worsen with Age, *Environmental Science & Technology*, doi:10.1021/acs.est.5b04704, 2016.

Chen, Y., Zhi, G., Feng, Y., Liu, D., Zhang, G., Li, J., Sheng, G. and Fu, J.: Measurements of Black and Organic Carbon Emission Factors for Household Coal Combustion in China: Implication for Emission Reduction, *Environ. Sci. Technol.*, 43(24), 9495–9500, doi:10.1021/es9021766, 2009.

Chen, Y., Morton, D. C., Jin, Y., Collatz, G. J., Kasibhatla, P. S., van der Werf, G. R., DeFries, R. S. and Randerson, J. T.: Long-term trends and interannual variability of forest, savanna and agricultural fires in South America, *Carbon Management*, 4(6), 617–638, doi:10.4155/cmt.13.61, 2013.

Cheung, K. L., Polidori, A., Ntziachristos, L., Tzamkiozis, T., Samaras, Z., Cassee, F. R., Gerlofs, M. and Sioutas, C.: Chemical characteristics and oxidative potential of particulate matter emissions from gasoline, diesel, and biodiesel cars, *Environmental science & technology*, 43(16), 6334–6340, 2009.

Christian, T. J., Yokelson, R. J., Cárdenas, B., Molina, L. T., Engling, G. and Hsu, S.-C.: Trace gas and particle emissions from domestic and industrial biofuel use and garbage burning in central Mexico, *Atmos. Chem. Phys.*, 10(2), 565–584, doi:10.5194/acp-10-565-2010, 2010.

Cofala, J. and Syri, S.: Sulfur emissions, abatement technologies and related costs for Europe in the RAINS model database, Interim Report, IIASA, Laxenburg, Austria. [online] Available from: <http://webarchive.iiasa.ac.at/Publications/Documents/IR-98-035.pdf> (Accessed 14 May 2013), 1998.

Cofala, J., Amann, M., Klimont, Z., Kupiainen, K. and Höglund-Isaksson, L.: Scenarios of Global Anthropogenic Emissions of Air Pollutants and Methane Until 2030, *Atmospheric Environment*, Vol 41/38, 8486–8499, 2007.

- Conrad, B. M. and Johnson, M. R.: Field Measurements of Black Carbon Yields from Gas Flaring, *Environ. Sci. Technol.*, **51**(3), 1893–1900, doi:10.1021/acs.est.6b03690, 2017.
- Coquelin, L., Fischer, N., Motzkus, C., Mace, T., Gensdarnes, F., Le Brusquet, L. and Fleury, G.: Aerosol size distribution estimation and associated uncertainty for measurement with a Scanning Mobility Particle Sizer (SMPS), *Journal of Physics: Conference Series*, **429**(1), 012018, 2013.
- Corbett, J. J. and Koehler, H. W.: Updated emissions from ocean shipping, *J. Geophys. Res.*, **108**, 15 PP., doi:200310.1029/2003JD003751, 2003.
- CPCB: Comprehensive industry document with emission standards, guidelines and stack height regulation for vertical shaft brick kilns (VSBK) vis-à-vis pollution control measures, Central Pollution Control Board (CPCB), New Delhi, India., 2007.
- Crippa, M., Janssens-Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R. and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmospheric Chemistry and Physics*, **16**(6), 3825–3841, doi:10.5194/acp-16-3825-2016, 2016.
- Dalsoren, S. B., Eide, M., Endresen, U., Mjelde, A., Gravir, G. and Isaksen, I. S. A.: Update on emissions and environmental impacts from the international fleet of ships: the contribution from major ship types and ports, *Atmospheric Chemistry and Physics*, **9**(6), 2171–2194, 2009.
- Delphi Inc.: Worldwide Emission Standards - Heavy Duty and Off-Highway Vehicles 2013/14, Delphi Inc., Troy, MICH/USA., 2013.
- Delphi Inc.: Worldwide Emission Standards - Passenger Cars and Light Duty Vehicles 2014/15, Delphi Inc., Troy, MICH/USA., 2015.
- Denier van der Gon, H. A. C., Gerlofs-Nijland, M. E., Gehrig, R., Gustafsson, M., Janssen, N., Harrison, R. M., Hulskotte, J., Johansson, C., Jozwicka, M., Keuken, M., Krijgheld, K., Ntziachristos, L., Riediker, M. and Cassee, F. R.: The Policy Relevance of Wear Emissions from Road Transport, Now and in the Future—An International Workshop Report and Consensus Statement, *Journal of the Air & Waste Management Association*, **63**(2), 136–149, doi:10.1080/10962247.2012.741055, 2013.
- Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D. and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe – revised estimates and an evaluation, *Atmospheric Chemistry and Physics*, **15**(11), 6503–6519, doi:10.5194/acp-15-6503-2015, 2015.
- Desai, S., Dubey, A., Joshi, B. L., Sen, M., Shariff, A. and Vanneman, R.: *Human Development in India: Challenges for a Society in Transition*, Oxford University Press, New Delhi, India., 2010.
- Driscoll, T., Milkovits, G. and Fredenberger, D.: *Impact and Use of Firewood in Australia*, CSIRO Sustainable Ecosystems., 2000.
- Durbin, T. D., Norbeck, J. M., Smith, M. R. and Truex, T. J.: Particulate Emission Rates from Light Duty Vehicles in the South Coast Air Quality Management District, *Environ. Sci. & Technol.*, **33**, 4401–4406, 1999.
- EAWAG: *Global waste challenge: Situation in developing countries*, 2008.
- Ebel, A., Friedrich, R. and Rodhe, H.: GENEMIS: Assessment, Improvement, and Temporal and Spatial Disaggregation of European Emission Data, in *Tropospheric Modelling and Emission Estimation: Chemical Transport and Emission Modelling*

on Regional, Global and Urban Scales, edited by A. Ebel, R. Friedrich, and H. Rodhe, pp. 181–214, Springer Berlin Heidelberg, Berlin, Heidelberg. [online] Available from: http://dx.doi.org/10.1007/978-3-662-03470-5_6, 1997.

EC: Directive 2001/80/EC of the European Parliament and of the Council of 23 October 2001 on the limitation of emissions of certain pollutants into the air from large combustion plants, 2001a.

5 EC: Directive 2001/81/EC of the European Parliament and of the Council of 23 October 2001 on national emission ceilings for certain atmospheric pollutants, European Parliament and Council, Luxembourg., 2001b.

EC: Directive 2010/75/EU of the European Parliament and of the council on industrial emissions (integrated pollution prevention and control), Commission of the European Communities, Brussels, Belgium. [online] Available from: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2010:334:0017:0119:EN:PDF>, 2010.

10 EC-JRC/PBL: Emission Database for Global Atmospheric Research (EDGAR)., European Commission (EC), Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL). [online] Available from: <http://edgar.jrc.ec.europa.eu>, 2010.

15 Eckhardt, S., Quennehen, B., Olivie, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Lund, M. T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I. E., Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J.-C., Rumbold, S. T., Schulz, M., Sharma, S., Skeie, R. B., Skov, H., Uttal, T., von Salzen, K. and Stohl, A.: Current model capabilities for simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi-model evaluation using a comprehensive measurement data set, *Atmos. Chem. Phys.*, 15(16), 9413–9433, doi:10.5194/acp-15-9413-2015, 2015.

20 Edwards, P.: Global cement emissions standards, Global Cement Magazine [online] Available from: www.GlobalCement.com, 2014.

EEA: EMEP/EEA air pollutant emission inventory guidebook, Technical report, European Environmental Agency (EEA), Copenhagen, Denmark. [online] Available from: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>, 2013.

25 Ekström, M., Sjödin, Å. and Andreasson, K.: Evaluation of the COPERT III emission model with on-road optical remote sensing measurements, *Atmospheric Environment*, 38(38), 6631–6641, doi:10.1016/j.atmosenv.2004.07.019, 2004.

Elisabeth, T.: Black Carbon Emissions from Kerosene Lamps: Potential for a new CCAC Initiative, Ecologic Institute, Berlin, Germany. [online] Available from: http://www.ecologic.eu/sites/files/publication/2014/black-carbon-and-kerosene-lamps-study_0.pdf, 2013.

30 Elvidge, C. D., Ziskin, D., Baugh, K. E., Tuttle, B. T., Ghosh, T., Pack, D. W., Erwin, E. H. and Zhizhin, M.: A Fifteen Year Record of Global Natural Gas Flaring Derived from Satellite Data, *Energies*, 2(3), 595, doi:10.3390/en20300595, 2009.

Elvidge, C. D., Baugh, K. E., Anderson, S., Ghosh, T. and Ziskin, D.: Estimation of Gas Flaring Volumes Using NASA MODIS Fire Detection Products, NOAA National Geophysical Data Center, Boulder, US. [online] Available from: http://www.ngdc.noaa.gov/dmsp/interest/gas_flares.html, 2011.

35 Elvidge, C. D., Zhizhin, M., Hsu, F.-C. and Baugh, K. E.: VIIRS Nightfire: Satellite Pyrometry at Night, *Remote Sensing*, 5(9), 4423, doi:10.3390/rs5094423, 2013.

5 [Endresen, Ø., Sørgård, E., Behrens, H. L., Brett, P. O. and Isaksen, I. S. A.: A historical reconstruction of ships' fuel consumption and emissions. *Journal of Geophysical Research: Atmospheres*, 112\(D12\), n/a–n/a, doi:10.1029/2006JD007630, 2007.](#)

ESMAP: Power Sector Reform in Africa: Assessing Impact on Poor People, Energy Sector Management Assistance Programme (ESMAP), World Bank, Washington D.C., 2005.

EUROSTAT: Statistics database, [online] Available from: http://epp.eurostat.ec.europa.eu/portal/page/portal/statistics/search_database (Accessed 5 September 2011), 2011.

10 [Eyring, V., Isaksen, I. S. A., Berntsen, T., Collins, W. J., Corbett, J. J., Endresen, O., Grainger, R. G., Moldanova, J., Schlager, H. and Stevenson, D. S.: Transport impacts on atmosphere and climate: Shipping. *Atmos. Environ.*, 44, 4735–4771, doi:10.1016/j.atmosenv.2009.04.059, 2010.](#)

FAO: Status and development issues of the brick industry in Asia, Field Document, FAO Regional Wood Energy Development Programme in Asia, Bangkok, Thailand., 1993.

Fernandes, S. D., Trautmann, N. M., Streets, D. G., Roden, C. A. and Bond, T. C.: Global biofuel use, 1850–2000, *Global Biogeochem. Cycles*, 21(2), 1–15, doi:10.1029/2006GB002836, 2007.

15 [Foell, W., Pachauri, S., Spreng, D. and Zerriffi, H.: Household cooking fuels and technologies in developing economies. *Energy Policy*, 39\(12\), 7487–7496, doi:10.1016/j.enpol.2011.08.016, 2011.](#)

Fortner, E. C., Brooks, W. A., Onasch, T. B., Canagaratna, M. R., Massoli, P., Jayne, J. T., Franklin, J. P., Knighton, W. B., Wormhoudt, J., Worsnop, D. R., Kolb, C. E. and Herndon, S. C.: Particulate Emissions Measured During the TCEQ Comprehensive Flare Emission Study, *Ind. Eng. Chem. Res.*, 51(39), 12586–12592, doi:10.1021/ie202692y, 2012.

20 [Frost & Sullivan: Indian Diesel Generator Set Market. \[online\] Available from: <http://www.frost.com/sublib/frost-content.do?sheetName=report-overview&sheetGroup=P420-01-00-00-00&viewName=virtual-brochure&repid=P420-01-00-00-00>, 2010.](#)

25 [Gadhavi, H. S., Renuka, K., Ravi Kiran, V., Jayaraman, A., Stohl, A., Klimont, Z. and Beig, G.: Evaluation of black carbon emission inventories using a Lagrangian dispersion model – a case study over southern India. *Atmos. Chem. Phys.*, 15\(3\), 1447–1461, doi:10.5194/acp-15-1447-2015, 2015.](#)

GEA: Global Energy Assessment: Toward a Sustainable Future, Cambridge University Press, UK., 2012.

Geller, M. D., Ntziachristos, L., Mamakos, A., Samaras, Z., Schmitz, D. A., Froines, J. R. and Sioutas, C.: Physicochemical and redox characteristics of particulate matter (PM) emitted from gasoline and diesel passenger cars, *Atmospheric Environment*, 40(36), 6988–7004, doi:10.1016/j.atmosenv.2006.06.018, 2006.

30 [Germain, A., Granger, F. and Gosselin, A.: A Model Municipal By-Law for regulating wood burning appliances A. Germain, F. Granger & A. Gosselin, *WIT Transactions on Ecology and the Environment*, 116, doi:10.2495/AIR080631, 2008.](#)

Gilmore, E. A., Lave, L. B. and Adams, P. J.: The costs, air quality and human health effects of meeting peak electricity demand with installed backup generators, *Environmental Science & Technology*, 40(22), 6887–6893, 2006.

Goel, R. and Guttikunda, S. K.: Role of urban growth, technology, and judicial interventions on vehicle exhaust emissions in Delhi for 1991–2014 and 2014–2030 periods, *Environmental Development*, 14, 6–21, doi:10.1016/j.envdev.2015.03.002, 2015.

GOI: 2011 Census of India: Source of Lighting, Ministry of Home Affairs, Government of India (GOI), New Delhi., 2011.

Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J.-F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J.-C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., Aardenne, J., Werf, G. R. and Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109(1–2), 163–190, doi:10.1007/s10584-011-0154-1, 2011.

Guo, H., Zhang, Q., Shi, Y. and Wang, D.: On-road remote sensing measurements and fuel-based motor vehicle emission inventory in Hangzhou, China, *Atmospheric Environment*, 41(14), 3095–3107, 2007.

Guttikunda, S. K. and Jawahar, P.: Atmospheric emissions and pollution from the coal-fired thermal power plants in India, *Atmospheric Environment*, 92, 449–460, doi:10.1016/j.atmosenv.2014.04.057, 2014.

Guttikunda, S. K., Begum, B. A. and Wadud, Z.: Particulate pollution from brick kiln clusters in the Greater Dhaka region, Bangladesh, *Air Quality, Atmosphere & Health*, 6(2), 357–365, doi:10.1007/s11869-012-0187-2, 2013.

Habib, G., Venkataraman, C., Bond, T. C. and Schauer, J. J.: Chemical, Microphysical and Optical Properties of Primary Particles from the Combustion of Biomass Fuels, *Environ. Sci. Technol.*, 42(23), 8829–8834, doi:10.1021/es800943f, 2008.

Harrison, R. M., Jones, A. M., Gietl, J., Yin, J. and Green, D. C.: Estimation of the Contributions of Brake Dust, Tire Wear, and Resuspension to Nonexhaust Traffic Particles Derived from Atmospheric Measurements, *Environmental Science & Technology*, 46(12), 6523–6529, doi:10.1021/es300894r, 2012.

Haugland, T., Saunier, S., Pederstad, A., Holm, T., Darani, H. and Kertesheva, A.: Associated Petroleum Gas Flaring Study for Russia, Kazakhstan, Turkmenistan and Azerbaijan, Final Report, Carbon Limits AS, Oslo, Norway. [online] Available from: www.carbonlimits.no, 2013.

HBEFA 3.1: HBEFA - Handbook Emission Factors for Road Transport, [online] Available from: <http://www.hbefa.net/d/index.html> (Accessed 20 March 2012), 2010.

Hegg, D. A., Livingston, J., Hobbs, P. V., Novakov, T. and Russell, P.: Chemical apportionment of aerosol column optical depth off the mid-Atlantic coast of the United States, *Journal of Geophysical Research: Atmospheres*, 102(D21), 25293–25303, doi:10.1029/97JD02293, 1997.

Heierli, U. and Maithel, S.: Brick by Brick: The Herculean Task of Cleaning of the Asian Brick Industry, Swiss Agency for Development and Cooperation (SDC), Bern., 2008.

Herzog, J. W.: Current and near-term emission control strategies for diesel powered generator sets, in *Proceedings of 24th International Telecommunications Energy Conference (INTELEC)*, pp. 394 – 399, Montréal., 2002.

Hodzic, A., Wiedinmyer, C., Salcedo, D. and Jimenez, J. L.: Impact of Trash Burning on Air Quality in Mexico City, *Environ. Sci. Technol.*, 46(9), 4950–4957, doi:10.1021/es203954r, 2012.

Hsu, Y. and Mullen, M.: Compilation of Diesel Emissions Speciation Data, Final Report for CRC., 2007.

Huang, K., Fu, J. S., Prikhodko, V. Y., Storey, J. M., Romanov, A., Hodson, E. L., Cresko, J., Morozova, I., Ignatieva, Y. and Cabaniss, J.: Russian anthropogenic black carbon: Emission reconstruction and Arctic black carbon simulation, *Journal of Geophysical Research: Atmospheres*, 120(21), 11,306–11,333, doi:10.1002/2015JD023358, 2015.

Huo, H., Zhang, Q., He, K., Yao, Z., Wang, X., Zheng, B., Streets, D. G., Wang, Q. and Ding, Y.: Modeling vehicle emissions in different types of Chinese cities: Importance of vehicle fleet and local features, *Environmental Pollution*, 159(10), 2954–2960, doi:10.1016/j.envpol.2011.04.025, 2011.

Huo, H., Lei, Y., Zhang, Q., Zhao, L. and He, K.: China's coke industry: Recent policies, technology shift, and implication for energy and the environment, *Energy Policy*, 51(0), 397–404, doi:10.1016/j.enpol.2012.08.041, 2012.

ICC & SRI: Atmospheric emissions of particulates from agriculture: a scoping study. Final report for the Ministry of Agriculture, Fisheries and Food (MAFF) Research and Development, ICCConsultants and Silsoe Research Institute, London, UK., 2000.

ICCT & Dieselnets: *Transportpolicy.net*, [online] Available from: http://transportpolicy.net/index.php?title=Main_Page (Accessed 15 March 2014), 2014.

IEA: Coal Power 2: World coal-fired power stations and their pollution control systems, International Energy Agency, London, UK., 1997.

IEA: World Energy Outlook 2007, International Energy Agency, Paris, France., 2007.

IEA: World Energy Outlook 2011, International Energy Agency, Paris, France., 2011.

IEA: Energy Technology Perspectives 2012 - Pathways to a Clean Energy System, International Energy Agency, IEA/OECD, Paris, [online] Available from: http://www.iea.org/publications/freepublications/publication/ETP2012_free.pdf, 2012a.

IEA: World Energy Outlook 2012, International Energy Agency, Paris, France., 2012b.

IEA: Energy Statistics of Non-OECD Countries 2015, International Energy Agency, IEA/OECD, Paris, France., 2015a.

IEA: Energy Statistics of OECD Countries 2015, International Energy Agency, IEA/OECD, Paris, France., 2015b.

IEA: World Energy Outlook Special Report: Energy and Air Pollution, International Energy Agency (IEA), Paris, France., 2016.

IEA CCC: Coal Power. Coal Power Database, IEA Clean Coal Centre, London, UK., 2012.

IFC/WB: Lighting Africa market assessment results: quantitative assessment – Ethiopia, International Finance Corporation (IFC) and the World Bank (WB), Washington D.C., 2008.

IIDFC: Environmental Management Framework for improving kiln efficiency in the brick making industry in Bangladesh: Hybrid Hoffman Kiln (HHK) project, Industrial and Infrastructure Development Finance Company Ltd. (IIDFC), Dhaka, Bangladesh., 2009.

IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Intergovernmental Panel on Climate Change, IGES, Japan, [online] Available from: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm>, 2006a.

IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, Waste, Intergovernmental Panel on Climate Change (IPCC), Kanagawa, Japan. [online] Available from: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm>, 2006b.

Ite, A. and Ibok, U. J.: Gas flaring and venting associated with petroleum exploration and production in the Nigeria's Niger Delta, *American Journal of Environmental Protection*, 1, 70–77, 2013.

Jacobson, A., Lam, N. L., Bond, T. C. and Hultman, N.: *Black Carbon and Kerosene Light-ing: An Opportunity for Rapid Action on Climate Change and Clean Energy for Development*, Policy Paper 2013-03, The Bookings Institution, Washington D.C., 2013.

Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B. and Li, M.: HTAP v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmospheric Chemistry and Physics*, 15(19), 11411–11432, doi:10.5194/acp-15-11411-2015, 2015.

Jiang, M., Marr, L. C., Dunlea, E. J., Herndon, S. C., Jayne, J. T., Kolb, C. E., Knighton, W. B., Rogers, T. M., Zavala, M., Molina, L. T. and Molina, M. J.: Vehicle fleet emissions of black carbon, polycyclic aromatic hydrocarbons, and other pollutants measured by a mobile laboratory in Mexico City, *Atmospheric Chemistry and Physics*, 5(12), 3377–3387, doi:10.5194/acp-5-3377-2005, 2005.

Jimenez, J. L., McRae, G. J., Nelson, D., Zahniser, M. S. and Kolb, C. E.: Remote Sensing of NO and NO₂ Emissions from Heavy-Duty Diesel Truck Using Tunable Diode Lasers, *Environ. Sci. & Technol.*, 33, 2380–2387, 2000.

Johnson, G. R., Jayaratne, E. R., Lau, J., Thomas, V., Juwono, A. M., Kitchen, B. and Morawska, L.: Remote measurement of diesel locomotive emission factors and particle size distributions, *Atmospheric Environment*, 81, 148–157, doi:10.1016/j.atmosenv.2013.09.019, 2013.

Johnson, M., Edwards, R., Alatorre Frenk, C. and Masera, O.: In-field greenhouse gas emissions from cookstoves in rural Mexican households, *Atmospheric Environment*, 42(6), 1206–1222, doi:10.1016/j.atmosenv.2007.10.034, 2008.

Johnson, M. R., Devillers, R. W. and Thomson, K. A.: Quantitative Field Measurement of Soot Emission from a Large Gas Flare Using Sky-LOSA., *Environ. Sci. Technol.*, 45(1), 345–350, 2011.

KEEI: *Yearbook of Energy Statistics 2011*, Korea Energy Economics Institute (KEEI), South Korea., 2011.

Kholod, N., Evans, M. and Kuklinski, T.: Russia's black carbon emissions: focus on diesel sources, *Atmospheric Chemistry and Physics Discussions*, 1–27, doi:10.5194/acp-2016-475, 2016.

Kinsey, J. S.: *Characterization of emissions from commercial aircraft engines during the aircraft particle emissions experiment (APEX) 1–3*, Technical report, US Environmental Protection Agency Office of Research and Development National Risk Management Research Laboratory, Washington DC, USA. [online] Available from: <http://nepis.epa.gov/Adobe/PDF/P1005KRK.pdf>, 2009.

Kirchstetter, T. W., Harley, R. A., Kreisberg, N. M., Stolzenburg, M. R. and Hering, S. V.: On-road measurement of fine particle and nitrogen oxide emissions from light- and heavy-duty motor vehicles, *Atmospheric Environment*, 33(18), 2955–2968, doi:10.1016/S1352-2310(99)00089-8, 1999.

- 5 [Klimont, Z., Hoeglund-Isaksson, L., Heyes, C., Rafaj, P., Schoepp, W., Cofala, J., Purohit, P., Borken-Kleefeld, J., Kupiainen, K., Kiesewetter, G., Winiwarter, W., Amann, M., Zhao, B., Wang, S., Bertok, I. and Sander, R.: Global scenarios of air pollutants and methane: 1990-2050, in preparation.](#)
- 5 [Klimont, Z., Streets, D. G., Gupta, S., Cofala, J., Lixin, F. and Ichikawa, Y.: Anthropogenic Emissions of Non-Methane Volatile Organic Compounds \(NMVOC\) in China, Atmospheric Environment, 36\(8\), 1309–1322, 2002a.](#)
- [Klimont, Z., Cofala, J., Bertok, I., Amann, M., Heyes, C. and Gyarmas, F.: Modelling Particulate Emissions in Europe. A Framework to Estimate Reduction Potential and Control Costs, Interim Report, International Institute for Applied Systems Analysis \(IIASA\), Laxenburg, Austria., 2002b.](#)
- 10 [Klimont, Z., Cofala, J., Xing, J., Wei, W., Zhang, C., Wang, S., Kejun, J., Bhandari, P., Mathur, R., Purohit, P., Rafaj, P., Chambers, A., Amann, M. and Hao, J.: Projections of SO₂, NO_x, and carbonaceous aerosols emissions in Asia, Tellus, 61B\(4\), 602–617, doi:10.1111/j.1600-0889.2009.00428.x, 2009.](#)
- [Klimont, Z., Smith, S. J. and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000-2011 emissions, Environ. Res. Let., 8\(1\), 014003, doi:10.1088/1748-9326/8/1/014003, 2013.](#)
- 15 [Kondo, Y., Oshima, N., Kajino, M., Mikami, R., Moteki, N., Takegawa, N., Verma, R. L., Kajii, Y., Kato, S. and Takami, A.: Emissions of black carbon in East Asia estimated from observations at a remote site in the East China Sea, Journal of Geophysical Research: Atmospheres, 116\(D16\), n/a–n/a, doi:10.1029/2011JD015637, 2011.](#)
- [Krasenbrink, A. and Dobranskyte-Niskota, A.: 2007 Technical Review of the NRM Directive 1997/68/EC as amended by Directives 2002/88/EC and 2004/26/EC, JRC, Ispra, Italy., 2008.](#)
- 20 [Kumar, S., Aggarwal, S. G., Gupta, P. K. and Kawamura, K.: Investigation of the tracers for plastic-enriched waste burning aerosols, Atmospheric Environment, 108, 49–58, doi:10.1016/j.atmosenv.2015.02.066, 2015.](#)
- [Kupiainen, K. and Klimont, Z.: Primary Emissions of Submicron and Carbonaceous Particles in Europe and the Potential for their Control, International Institute for Applied Systems Analysis \(IIASA\), Laxenburg, Austria., 2004.](#)
- [Kupiainen, K. and Klimont, Z.: Primary emissions of fine carbonaceous particles in Europe, Atmospheric Environment, 41, 2156–2170, doi:10.1016/j.atmosenv.2006.10.066, 2007.](#)
- 25 [Kupiainen, K. J. and Pirjola, L.: Vehicle non-exhaust emissions from the tyre–road interface – effect of stud properties, traction sanding and resuspension, Atmospheric Environment, 45\(25\), 4141–4146, doi:10.1016/j.atmosenv.2011.05.027, 2011.](#)
- 30 [Kupiainen, K. J., Tervahattu, H., Räisänen, M., Mäkelä, T., Aurela, M. and Hillamo, R.: Size and Composition of Airborne Particles from Pavement Wear, Tires, and Traction Sanding, Environ. Sci. Technol., 39\(3\), 699–706, doi:10.1021/es035419e, 2005.](#)
- [Lack, D., Lerner, B., Granier, C., Baynard, T., Lovejoy, E., Massoli, P., Ravishankara, A. R. and Williams, E.: Light absorbing carbon emissions from commercial shipping, Geophysical Research Letters, 35\(13\), L13815, doi:10.1029/2008GL033906, 2008.](#)
- 35 [Lack, D. A. and Corbett, J. J.: Black carbon from ships: a review of the effects of ship speed, fuel quality and exhaust gas scrubbing, Atmos. Chem. Phys., 12, 3985–4000, doi:10.5194/acp-12-3985-2012, 2012.](#)

Lack, D. A., Corbett, J. J., Onasch, T., Lerner, B., Massoli, P., Quinn, P. K., Bates, T. S., Covert, D. S., Coffman, D., Sierau, B. and others: Particulate emissions from commercial shipping: Chemical, physical, and optical properties, *J. Geophys. Res.*, 114, D00F04, 2009.

5 Lam, N. L., Chen, Y., Weyant, C., Venkataraman, C., Sadavarte, P., Johnson, M. A., Smith, K. R., Brem, B. T., Arineitwe, J., Ellis, J. E. and Bond, T. C.: Household Light Makes Global Heat: High Black Carbon Emissions From Kerosene Wick Lamps, *Environmental Science & Technology*, 46(24), 13531–13538, doi:10.1021/es302697h, 2012.

10 Lam, N. L., Pachauri, S., Cameron, C., Purohit, P. and Nagai, Y.: Characterizing Kerosene Demand for Light in India and Evaluating the Impact of Measures Affecting Access and Dependence, in *Discovering Untapped Resources*, p. 116–119., UC Berkeley. [online] Available from: <https://opus4.kobv.de/opus4-tuberlin/frontdoor/index/index/docId/5185> (Accessed 20 March 2015), 2014.

Lam, N. L., Pachauri, S., Purohit, P., Nagai, Y., Bates, M. N., Cameron, C. and Smith, K. R.: Kerosene subsidies for household lighting in India: what are the impacts?, *Environmental Research Letters*, 11(4), 044014 (1-11), doi:10.1088/1748-9326/11/4/044014, 2016.

15 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussé, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D. T., Smith, S. J., Stehfest, E., Van Aardenne, J. A., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K. and Van Vuuren, D. P. V.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application., *Atmos. Chem. Phys.*, 10(2), 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.

20 Lange, D. and Domke, F.: The exhaust emissions scandal („Dieselgate“) - Take a deep breath into pollution trickery, [online] Available from: <https://media.ccc.de/v/32c3-7331-the-exhaust-emissions-scandal-dieselgate> (Accessed 5 January 2016), 2015.

Lawson, D. R.: (On-Road) Mobile Source Emissions and Mitigation Potential, 2010.

Le, H. A. and Oanh, N. T. K.: Integrated assessment of brick kiln emission impacts on air quality Hoang Anh Le · Nguyen Thi Kim Oanh, *Environ Monit Assess.*, 171, 381–394, doi:10.1007/s10661-009-1285-y, 2010.

25 Lee, D. S., Fahey, D. W., Forster, P. M., Newton, P. J., Wit, R. C. N., Lim, L. L., Owen, B. and Sausen, R.: Aviation and global climate change in the 21st century, *Atmos. Environ.*, 43, 3520–3537, 2009.

Lee, W. J., Liu, Y. C., Mwangi, F. K., Chen, W. H., Lin, S. L., Fukushima, Y., Liao, C. N. and Wang, L. C.: Assessment of energy performance and air pollutant emissions in a diesel engine generator fueled with water-containing ethanol–biodiesel–diesel blend of fuels, *Energy*, 36(9), 5591–5599, doi:10.1016/j.energy.2011.07.012, 2011.

30 Lei, Y., Zhang, Q., He, K. B. and Streets, D. G.: Primary anthropogenic aerosol emission trends for China, 1990–2005, *Atmospheric Chemistry and Physics*, 11(3), 931–954, doi:10.5194/acp-11-931-2011, 2011.

Lewis, J. J. and Pattanayak, S. K.: Who Adopts Improved Fuels and Cookstoves? A Systematic Review, *Environmental Health Perspectives*, 120, 637–645, 2012.

35 Li, O., Li, X., Jiang, J., Duan, L., Ge, S., Zhang, O., Deng, J., Wang, S. and Hao, J.: Semi-coke briquettes: towards reducing emissions of primary PM_{2.5}, particulate carbon, and carbon monoxide from household coal combustion in China, *Scientific Reports*, 6, 19306, 2016.

Li, X. H., Wang, S. X., Duan, L., Hao, J. M. and Nie, Y. F.: Carbonaceous aerosol emissions from household biofuel combustion in China., *Environ. Sci. Technol.*, 43, 6076–6081, 2009.

Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H., AlMazroa, M. A., Amann, M., Anderson, H. R., Andrews, K. G., Arvee, M., Atkinson, C., Bacchus, L. J., Bahalim, A. N., Balakrishnan, K., Balmes, J., Barker-Collo, S., Baxter, A., Bell, M. L., Blore, J. D., Blyth, F., Bonner, C., Borges, G., Bourne, R., Boussinesq, M., Brauer, M., Brooks, P., Bruce, N. G., Brunekreef, B., Bryan-Hancock, C., Bucello, C., Buchbinder, R., Bull, F., Burnett, R. T., Byers, T. E., Calabria, B., Carapetis, J., Carnahan, E., Chafe, Z., Charlson, F., Chen, H., Chen, J. S., Cheng, A. T.-A., Child, J. C., Cohen, A., Colson, K. E., Cowie, B. C., Darby, S., Darling, S., Davis, A., Degenhardt, L., Dentener, F., Des Jarlais, D. C., Devries, K., Dherani, M., Ding, E. L., Dorsey, E. R., Driscoll, T., Edmond, K., Ali, S. E., Engell, R. E., Erwin, P. J., Fahimi, S., Falder, G., Farzadfar, F., Ferrari, A., Finucane, M. M., Flaxman, S., Fowkes, F. G. R., Freedman, G., Freeman, M. K., Gakidou, E., Ghosh, S., Giovannucci, E., Gmel, G., Graham, K., Grainger, R., Grant, B., Gunnell, D., Gutierrez, H. R., Hall, W., Hoek, H. W., Hogan, A., Hosgood, H. D., Hov, D., Hu, H., Hubbell, B. J., Hutchings, S. J., Ibeanusi, S. E., Jacklyn, G. L., Jasrasaria, R., Jonas, J. B., Kan, H., Kanis, J. A., Kassebaum, N., Kawakami, N., Khang, Y.-H., Khatibzadeh, S., Khoo, J.-P., et al.: A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010, *The Lancet*, 380(9859), 2224–2260, doi:10.1016/S0140-6736(12)61766-8, 2012.

Lin, J., Nielsen, C. P., Zhao, Y., Lei, Y., Liu, Y. and McElroy, M. B.: Recent changes in particulate air pollution over China observed from space and the ground: Effectiveness of emission control, *Environmental Science and Technology-Columbus*, 44(20), 7771, 2010.

Lin, Y. C., Lee, W. J., Chao, H. R., Wang, S. L., Tsou, T. C., Chang-Chien, G. P. and Tsai, P. J.: Approach for energy saving and pollution reducing by fueling diesel engines with emulsified biosolution/biodiesel/diesel blends, *Environmental Science and Technology*, 42(10), 3849–3855, 2008.

Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H. and He, K. B.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, *Atmospheric Chemistry and Physics*, 15(23), 13299–13317, doi:10.5194/acp-15-13299-2015, 2015.

Liu, H., He, K., Lents, J. M., Wang, Q. and Tolvet, S.: Characteristics of Diesel Truck Emission in China Based on Portable Emissions Measurement Systems, *Environmental Science & Technology*, 43(24), 9507–9511, doi:10.1021/es902044x, 2009.

Louhelainen, K., Vilhunen, P., Kangas, J. and Terho, F. O.: Dust exposure in piggeries, *European J. of Resp. Diseases*, 71(152), 80–90, 1987.

Lu, Z., Zhang, Q. and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, *Atmos. Chem. Phys.*, 11(18), 9839–9864, 2011.

Lükewille, A., Bertok, I., Amann, M., Cofala, J., Gyarmas, F., Johansson, M., Klimont, Z., Pacyna, E. and Pacyna, J.: A Module to Calculate Primary Particulate Matter Emissions and Abatement Measures in Europe, *Water, Air, and Soil Pollution*, 130(1–4), 229–234, doi:10.1023/A:1013807101561, 2001.

Lund, M. T., Berntsen, T. K., Heyes, C., Klimont, Z. and Samset, B. H.: Global and regional climate impacts of black carbon and co-emitted species from the on-road diesel sector, *Atmos Environ*, 98, 50–58, doi:10.1016/j.atmosenv.2014.08.033, 2014.

MacCarty, N., Ogle, D., Still, D., Bond, T., Roden, C. and Willson, B.: Laboratory Comparison of the Global-Warming Potential of Six Categories of Biomass Cooking Stoves, Aprovecho Research Center. [online] Available from: internal-pdf://Global warming full 9-6-07-2181006597/Global warming full 9-6-07.pdf, 2007.

- Mahapatra, S., Chanakya, H. N. and Dasappa, S.: Evaluation of various energy devices for domestic lighting in India: Technology, economics and CO₂ emissions. *Energy for Sustainable Development*, 13(4), 271–279, doi:10.1016/j.esd.2009.10.005, 2009.
- 5 Maitheh, S.: Factsheets about Brick Kilns in South and South-East Asia. [online] Available from: <http://www.ccacoalition.org/en/resources/factsheets-about-brick-kilns-south-and-south-east-asia>, 2014.
- Maitheh, S., Lalchandani, D., Malhotra, G., Bhanware, P., Uma, R., Ragavan, S., Athalye, V., Bindiya, K., Reddy, S., Bond, T. C., Weyant, C., Baum, E., Kim Thoa, V. T., Thu Phuong, N. and Kim Thanh, T.: Brick Kilns Performance Assessment: A Roadmap for Cleaner Brick Production in India, Shakti Sustainable Energy Foundation and Climate Works Foundation. [online] Available from: <http://www.catf.us/resources/publications/view/161>, 2012.
- 10 Maíz, P.: Experiencias en Medición de Emisiones en Hornos para Manufactura Artesanal de Ladrillos en Mexico. [online] Available from: <http://www.ine.gob.mx/cenica-memorias/1111-taller-ladrilleras-2012-eng> (Accessed 17 September 2012), 2012.
- Mancilla, Y., Araizaga, A. E. and Mendoza, A.: A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico. *Journal of the Air & Waste Management Association*, 62(12), 1431–1442, doi:10.1080/10962247.2012.717902, 2012.
- 15 Maricq, M. M.: Chemical characterization of particulate emissions from diesel engines: A review M. Matti Maricq*. *Aerosol Science*, 38, 1079–1118, doi:10.1016/j.jaerosci.2007.08.001, 2007.
- Mazzoleni, C., Moosmüller, H., Kuhns, H. D., Keislar, R. E., Barber, P. W., Nikolic, D., Nussbaum, N. J. and Watson, J. G.: Correlation between automotive CO, HC, NO, and PM emission factors from on-road remote sensing: implications for inspection and maintenance programs, *Transportation Research Part D: Transport and Environment*, 9(6), 477–496, 2004.
- 20 McClintock, P.: Remote Sensing Measurements of Real World High Exhaust Emitters, Coordinating Research Council, Inc., 1999.
- McClintock, P.: 2007 High Emitter Remote Sensing Project, Prepared for Southeast Michigan Council of Governments, Environmental Systems Products Inc., Tiburon, CA/USA., 2007.
- 25 McEwen, J. D. N. and Johnson, M. R.: Black Carbon Particulate Matter Emission Factors for Buoyancy Driven Associated Gas Flares. *Journal of the Air & Waste Management Association*, 63(3), 307–321, doi:10.1080/10473289.2011.650040, 2012.
- Mills, E.: Technical and Economic Performance Analysis of Kerosene Lamps and Alternative Approaches to Illumination in Developing Countries, Lawrence Berkeley National Laboratory (LBNL), Berkeley. [online] Available from: <http://evanmills.lbl.gov/pubs/pdf/offgrid-lighting.pdf>, 2003.
- 30 Mills, E.: The specter of fuel-based lighting. *Science*, 308, 1263–1264, 2005.
- Ministério do Meio Ambiente: Inventário Nacional de Emissões Atmosféricas por Veículos Automotores Rodoviários. Final Report, Ministério do Meio Ambiente, Secretaria de Mudanças Climáticas e Qualidade Ambiental (MMA), Brasília/BR., 2011.
- 35 Moldanova, J., Fridell, E., Popovichieva, O., Demirdjian, B., Tishkova, V., Faccineto, A. and Focsa, C.: Characterisation of Particulate Matter and Gaseous Emissions from a Large Ship Diesel Engine., *Atmos. Environ.*, 43, 2632–2641, 2009.

Morton, D. C., DeFries, R. S., Randerson, J. T., Giglio, L., Schroeder, W. and Van Der Werf, G. R.: Agricultural intensification increases deforestation fire activity in Amazonia, *Global Change Biology*, 14(10), 2262–2275, doi:10.1111/j.1365-2486.2008.01652.x, 2008.

5 Mukai, S., Nakata, M., Yasumoto, M., Sano, I. and Kokhanovsky, A.: A study of aerosol pollution episode due to agriculture biomass burning in the east-central China using satellite data, *Frontiers in Environmental Science*, 3(Article 57), doi:10.3389/fenvs.2015.00057, 2015.

10 Murphy, S. M., Agrawal, H., Sorooshian, A., Padro, L. T., Gates, H., Hersey, S., Welch, W. A., Jung, H., Miller, J. W., Cocker III, D. R., Nenes, A., Jonsson, H. H., Flagan, R. C. and Seinfeld, J. H.: Comprehensive Simultaneous Shipboard and Airborne Characterization of Exhaust from a Modern Container Ship at Sea, *Environ. Sci. & Technol.*, 43(13), 4626–4640, 2009.

NEA: A Year in Review: Fiscal Year 2011/2012, Nepal Electricity Authority (NEA), Kathmandu., 2012.

Neurath, C.: Open Burning of Domestic Wastes: The Single Largest Source of Dioxin to Air?, *Organohalogen Compounds*, 60–65, 2003.

15 NIELSEN: All India Study on Sectoral Demand of Diesel & Petrol, Petroleum Planning and Analysis Cell (PPAC), Ministry of Petroleum & Natural Gas, Government of India, New Delhi. [online] Available from: <http://ppac.org.in>, 2013.

Niemi, J. K.: Atmospheric Emissions from Open Biomass Burning: Development of Datasets for RAINS Model, Interim Report, International Institute for Applied Systems Analysis, Laxenburg, Austria., 2007.

20 Norris, J. O. W.: An In-Service Emissions Test for Spark Ignition (SI) Petrol Engines – PPAD 9/107/09, Phase 1 Report, Definition of an Excess Emitter and Effectiveness of Current Annual Test, AEAT Technology, Harwell, UK. [online] Available from: https://uk-air.defra.gov.uk/assets/documents/reports/cat15/0408171318_S1Phase1reportIssue3.pdf, 2001.

NSSO: Energy sources of India household for cooking and lighting, NSS 61st Round (July 2004–June 2005), National Sample Survey Organisation (NSSO), Ministry of Statistics and Programme Implementation, Government of India., New Delhi, India., 2007.

25 Ntziachristos, L., Gkatzoflias, D., Kouridis, C. and Samaras, Z.: COPERT: A European Road Transport Emission Inventory Model, in *Information Technologies in Environmental Engineering - Proceedings of the 4th International ICSC Symposium Thessaloniki, Greece, May 28-29, 2009*, pp. 491–504, Springer Berlin Heidelberg., 2009.

Oanh, N. T. K., Bich, T. L., Tipayarom, D., Manadhar, B. R., Prapat, P., Simpson, C. D. and Liu, L.-J. S.: Charazcterization of particulate matter emission from open burning of rice straw, *Atmospheric environment (Oxford, England : 1994)*, 45(2), 493–502, doi:10.1016/j.atmosenv.2010.09.023, 2011.

30 Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X. and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, *Atmos. Chem. Phys.*, 7, 4419–4444, 2007.

Ohlström, M. O., Lehtinen, K. E. J., Moisio, M. and Jokiniemi, J. K.: Fine-particle emissions of energy production in Finland, *Atmospheric Environment*, 34(22), 3701–3711, doi:10.1016/S1352-2310(00)00076-5, 2000.

35 O'Neill, B. C., Krieglger, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R. and van Vuuren, D. P.: A new scenario framework for climate change research: the concept of shared socioeconomic pathways, *Climatic Change*, 122(3), 387–400, doi:10.1007/s10584-013-0905-2, 2014.

5
OTAQ: Exhaust and Crankcase Emission Factors for Nonroad Engine Modeling-- Compression-Ignition, US Environmental Protection Agency., 2004.

Paasonen, P., Visschedijk, A., Kupiainen, K., Klimont, Z., Denier van der Gon, H. A. C., Kulmala, M. and Amann, M.: Aerosol particle number emissions and size distributions: Implementation in the GAINS model and initial results, IIASA Interim Report, International Institute for Applied Systems Analysis, Laxenburg, Austria., 2013.

Paasonen, P., Kupiainen, K., Klimont, Z., Visschedijk, A., Denier van der Gon, H. A. C. and Amann, M.: Continental anthropogenic primary particle number emissions, Atmospheric Chemistry and Physics Discussions, 2016, 1–39, doi:10.5194/acp-2015-1023, 2016.

10 Parashar, D. C., Gadi, R., Mandal, T. K. and Mitra, A. P.: Carbonaceous aerosol emissions from India, Atmospheric Environment, 39, 7861–7871, 2005.

Parikh, K. S.: Report of the expert group on a viable and sustainable system of pricing petroleum products, Government of India, New Delhi, India., 2010.

15 Pederstad, A., Smith, J. D., Jackson, R., Saunier, S. and Holm, T.: Assessment of flare strategies, techniques for reduction of flaring and associated emissions, emission factors and methods for determination of emissions to air from flaring, Carbon Limits AS, Trondheim, Norway. [online] Available from: www.miljødirektoratet.no, 2015.

Pettersson, E., Boman, C., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 2: Wood Stove, Energy Fuels, 25(1), 315–323, doi:10.1021/ef1007787, 2011.

20 Pettus, A.: Agricultural Fires and Arctic Climate Change, CATF (Clean Air Task Force), Boston, US. [online] Available from: <http://www.catf.us>, 2009.

Petzold, A., Hasselbach, J., Lauer, P., Baumann, R., Franke, K., Gurk, C., Schlager, H. and Weingartner, E.: Experimental studies on particle emissions from cruising ship, their characteristic properties, transformation and atmospheric lifetime in the marine boundary layer, Atmos. Chem. Phys., 8, 2387–2403, doi:10.5194/acp-8-2387-2008, 2008.

25 Petzold, A., Weingartner, E., Hasselbach, J., Lauer, P., Kurok, C. and Fleischer, F.: Physical Properties, Chemical Composition, and Cloud Forming Potential of Particulate Emissions from a Marine Diesel Engine at Various Load Conditions., Environ. Sci. & Technol., 44(10), 3800–3805, 2010.

PFC Energy: Using Russia's Associated Gas, PFC Energy, Washington DC, USA. [online] Available from: www.pfcenergy.com, 2007.

30 Philip, S., Martin, R. V., Snider, G., Weagle, C. L., van Donkelaar, A., Brauer, M., Henze, D. K., Klimont, Z., Venkataraman, C., Guttikunda, S. K. and Zhang, Q.: Anthropogenic fugitive, combustion and industrial dust is a significant, underrepresented fine particulate matter source in global atmospheric models, Environ. Res. Lett. [online] Available from: <http://iopscience.iop.org/10.1088/1748-9326/aa65a4>, 2017.

35 Pine, K., Edwards, R., Masera, O., Schilman, A., Marrón-Mares, A. and Riojas-Rodríguez, H.: Adoption and use of improved biomass stoves in Rural Mexico, Energy for Sustainable Development, 15(2), 176–183, doi:10.1016/j.esd.2011.04.001, 2011.

Pode, R.: Solution to enhance the acceptability of solar-powered LED lighting technology, Renewable and Sustainable Energy Reviews, 14(3), 1096–1103, doi:10.1016/j.rser.2009.10.006, 2010.

Pokhrel, A. K., Bates, M. N., Verma, S. C., Joshi, H. S., Sreeramareddy, C. T. and Smith, K. R.: Tuberculosis and indoor biomass and kerosene use in Nepal: a case-control study, *Environmental Health Perspectives*, 118(4), 558–564, doi:10.1289/ehp.0901032, 2010.

Polenske, K. R. [ed]: *The Technology-Energy-Environment-Health (TEEH) Chain in China: A Case Study of Cokemaking*, Springer Netherlands., 2006.

Purohit, P. and Michaelowa, A.: CDM potential of SPV lighting systems in India, *Mitigation and Adaptation Strategies for Global Change*, 13(1), 23–46, doi:10.1007/s11027-006-9078-x, 2008.

Purohit, P., Amann, M., Mathur, R., Gupta, I., Marwah, S., Verma, V., Bertok, I., Borken-Kleefeld, J., Chambers, A., Cofala, J., Heyes, C., Hoglund-Isaksson, L., Klimont, Z., Rafaj, P., Sandler, R., Schoepp, W., Toth, G., Wagner, F. and Winiwarter, W.: GAINS-Asia. Scenarios for cost-effective control of air pollution and greenhouse gases in India, *International Institute for Applied Systems Analysis (IIASA)*, Laxenburg, Austria., 2010.

Quennehen, B., Raut, J.-C., Law, K. S., Daskalakis, N., Ancellet, G., Clerbaux, C., Kim, S.-W., Lund, M. T., Myhre, G., Olivie, D. J. L., Safieddine, S., Skeie, R. B., Thomas, J. L., Tsyro, S., Bazureau, A., Bellouin, N., Hu, M., Kanakidou, M., Klimont, Z., Kupiainen, K., Myriokefalitakis, S., Quaas, J., Rumbold, S. T., Schulz, M., Cherian, R., Shimizu, A., Wang, J., Yoon, S.-C. and Zhu, T.: Multi-model evaluation of short-lived pollutant distributions over east Asia during summer 2008, *Atmospheric Chemistry and Physics*, 16(17), 10765–10792, doi:10.5194/acp-16-10765-2016, 2016.

Rajaratnam, U., Athalye, V., Ragavan, S., Maithel, S., Lalchandani, D., Kumar, S., Baum, E., Weyant, C. and Bond, T.: Assessment of air pollutant emissions from brick kilns, *Atmospheric Environment*, 98, 549–553, doi:10.1016/j.atmosenv.2014.08.075, 2014.

Randerson, J. T., Chen, Y., van der Werf, G. R., Rogers, B. M. and Morton, D. C.: Global burned area and biomass burning emissions from small fires, *Journal of Geophysical Research: Biogeosciences*, 117(G4), doi:10.1029/2012JG002128, 2012.

Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J. and Kasibhatla, P. S.: *Global Fire Emissions Database, Version 4, (GFEDv4)*, [online] Available from: <http://dx.doi.org/10.3334/ORNLDAAC/1293>, 2015.

Rao, S., Pachauri, S., Dentener, F., Kinney, P., Klimont, Z., Riahi, K. and Schoepp, W.: Better air for better health: Forging synergies in policies for energy access, climate change and air pollution, *Global Environmental Change*, doi:10.1016/j.gloenvcha.2013.05.003, 2013.

Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., van Vuuren, D. P., Aletuia Reis, L., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Strefler, J., van der Sluis, S. and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Global Environmental Change*, 42, 346–358, doi:10.1016/j.gloenvcha.2016.05.012, 2017.

Reddington, C. L., Butt, E. W., Ridley, D. A., Artaxo, P., Morgan, W. T., Coe, H. and Spracklen, D. V.: Air quality and human health improvements from reductions in deforestation-related fire in Brazil, *Nature Geosci.*, 8(10), 768–771, 2015.

Riahi, K., Dentener, F., Gielen, D., Grubler, A., Jewell, J., Klimont, Z., Krey, V., McCollum, D., Pachauri, S., Rao, S., van Ruijven, B., van Vuuren, D. P. and Wilson, C.: Chapter 17 - Energy Pathways for Sustainable Development, in *Global Energy Assessment - Toward a Sustainable Future*, pp. 1203–1306, Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria. [online] Available from: www.globalenergyassessment.org, 2012.

5 [Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A. and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, doi:10.1016/j.gloenvcha.2016.05.009, 2017.](#)

10 [Roden, C. A., Bond, T. C., Conway, S. and Pinel, A. B. O.: Emission Factors and Real-Time Optical Properties of Particles Emitted from Traditional Wood Burning Cookstoves, *Environ. Sci. Technol.*, 40\(21\), 6750–6757, doi:10.1021/es052080i, 2006.](#)

[Roden, C. A., Bond, T. C., Conway, S., Osorto Pinel, A. B., MacCarty, N. and Still, D.: Laboratory and field investigations of particulate and carbon monoxide emissions from traditional and improved cookstoves, *Atmospheric Environment*, 43\(6\), 1170–1181, doi:10.1016/j.atmosenv.2008.05.041, 2009.](#)

15 [Røland, T. H.: Associated Petroleum Gas in Russia: Reasons for non-utilization, Fridtjof Nansens Institute, Lysaker, Norway. \[online\] Available from: www.fni.no, 2010.](#)

[Ruiz-Mercado, I., Masera, O., Zamora, H. and Smith, K. R.: Adoption and sustained use of improved cookstoves, *Energy Policy*, 39\(12\), 7557–7566, doi:10.1016/j.enpol.2011.03.028, 2011.](#)

20 [Schaap, M., Van Der Gon, H. A. C. D., Dentener, F. J., Visschedijk, A. J. H., Van Loon, M., ten Brink, H. M., Putaud, J.-P., Guillaume, B., Liousse, C. and Builtjes, P. J. H.: Anthropogenic black carbon and fine aerosol distribution over Europe, *Journal of Geophysical Research: Atmospheres*, 109\(D18\), n/a–n/a, doi:10.1029/2003JD004330, 2004.](#)

[Schäffeler, U. and Keller, M.: Non-road fuel consumption and pollutant emissions. Study for the period from 1980 to 2020, Federal Office for the Environment, Bern/CH., 2008.](#)

[Schilderman, T. and Mason, K.: Using residues as fuel in small-scale brickmaking, in *Proceedings of the 11th International Conference on Non-conventional Materials and Technologies \(NOCMAT 2009\)*, Bath, UK., 2009.](#)

25 [Schmidl, C., Luisser, M., Padouvas, E., Lasselsberger, L., Rzaca, M., Cruz, C. -S., Handler, M., Peng, G., Bauer, H. and Puxbaum, H.: Particulate and gaseous emissions from manually and automatically fired small scale combustion systems., *Atmospheric Environment*, 45, 7443–7454, 2011.](#)

[Schmidt, C. W.: Modernizing Artisanal Brick Kilns: A Global Need, *Environmental Health Perspectives*, 121, doi:http://dx.doi.org/10.1289/ehp.121-A242, 2013.](#)

30 [Schöpp, W., Klimont, Z., Suutari, R. and Cofala, J.: Uncertainty analysis of emission estimates in the RAINS integrated assessment model, *Environmental Science & Policy*, 8, 601–613, 2005.](#)

[Schwarz, J. P., Holloway, J. S., Katich, J. M., McKeen, S., Kort, E. A., Smith, M. L., Rverson, T. B., Sweeney, C. and Peischl, J.: Black Carbon Emissions from the Bakken Oil and Gas Development Region, *Environ. Sci. Technol. Lett.*, 2\(10\), 281–285, doi:10.1021/acs.estlett.5b00225, 2015.](#)

35 [Scott, A. J.: Real-life emissions from residential wood burning appliances in New Zealand, Ministry for the Environment, Christchurch, New Zealand., 2005.](#)

Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics: from air pollution to climate change, John Wiley & Sons, New York, USA, [online] Available from: <http://books.google.com/books?hl=en&lr=&id=YH2K9eWsZOC&oi=fnd&pg=PA1991&dq=%22activities,+including+oxidation+of+anthropogenically+derived+CH4,+The%22+%22of+industrial%22+%22the+maximum+values+are+found+near+the+surface+at+northern+midlatitudes.%22+%22of+its+overwhelming+importance+in+global+climate,+we+will+delay+consideration%22+&ots=hJ9tPmbVEB&sig=J5iCU26zodWW2ugZI1T2PdhFhw> (Accessed 1 August 2013), 2012.

Shafizadeh, K., Niemeier, D. and Eisinger, D. S.: Gross Emitting Vehicles: A Review of the Literature, Caltrans Air Quality Project, US Davis., 2004.

Shah, S. D., Cocker, D. R., Miller, J. W. and Norbeck, J. M.: Emission rates of particulate matter and elemental and organic carbon from in-use diesel engines, *Environmental Science & Technology*, 38(9), 2544–2550, doi:10.1021/es0350583, 2004.

Shah, S. D., Cocker, D. R., Johnson, K. C., Lee, J. M., Soriano, B. L. and Miller, J. W.: Emissions of regulated pollutants from in-use diesel backup generators, *Atmospheric Environment*, 40(22), 4199–4209, 2006.

Shah, S. D., Cocker III, D. R., Johnson, K. C., Lee, J. M., Soriano, B. L. and Miller, J. W.: Reduction of particulate matter emissions from diesel backup generators equipped with four different exhaust after treatment devices, *Environmental Science & Technology*, 41(14), 5070–5076, doi:10.1021/es0614161, 2007.

Shi, X., Pang, X., Mu, Y., He, H., Shuai, S., Wang, J., Chen, H. and Li, R.: Emission reduction potential of using ethanol-biodiesel-diesel fuel blend on a heavy-duty diesel engine, *Atmospheric Environment*, 40(14), 2567–2574, doi:10.1016/j.atmosenv.2005.12.026, 2006.

Shindell, D., Kuylenstierna, J. C. I., Vignati, E., Van Dingenen, R., Amann, M., Klimont, Z., Anenberg, S. C., Muller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L., Kupiainen, K., Höglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh, N. T. K., Milly, G., Williams, M., Demkine, V. and Fowler, D.: Simultaneously mitigating near-term climate change and improving human health and food security, *Science*, 335(6065), 183–189, 2012.

Shrimali, G., Slaski, X., Thurber, M. C. and Zerriffi, H.: Improved stoves in India: A study of sustainable business models, *Energy Policy*, 39(12), 7543–7556, doi:10.1016/j.enpol.2011.07.031, 2011.

Silk, B. J., Sudamah, I., Patel, M. K., Were, V., Person, R., Harris, J., Otieno, R., Nygren, B., Loo, J., Eleveld, A., Quick, R. E. and Cohen, A. L.: A strategy to increase adoption of locally produced, ceramic cookstoves in rural Kenyan households Benjamin J Silk1,2*, Ibrahim Sadumah3, Minal K Patel1,4, Vincent Were3, Bobbie Person4, Julie Harris1,4, Ronald Otieno3, Benjamin Nygren4, Jennifer Loo2, Alie Eleveld5, Robert E Quick4 and Adam L Cohen2, *BMC Public Health*, 12, 359–369, 2012.

Sippula, O., Hokkinen, J., Puustinen, H., Yli-Pirila, P. and Jokiniemi, J.: Comparison of particle emissions from small heavy fuel oil and wood-fired boilers, *Atmospheric Environment*, 43, 4855–4864, 2009.

Skinder, B. M., Sheikh, A. Q., Pandit, A. K. and Ganai, B. A.: Brick kiln emissions and its environmental impact: A Review Bhat Mohd Skinder*, Afeefa Qayoom Sheikh, Ashok K. Pandit and Bashir Ahmad Ganai, *Journal of Ecology and the Natural Environment*, 6(1), 1–11, 2014.

Smit, R. and Bluett, J.: A new method to compare vehicle emissions measured by remote sensing and laboratory testing: High-emitters and potential implications for emission inventories, *Science of The Total Environment*, 409(13), 2626–2634, doi:10.1016/j.scitotenv.2011.03.026, 2011.

Smith, T. W., Jalkanen, J. P., Anderson, B. A., Corbett, J. J., Faber, J., Hanayama, S., O'Keefe, E., Parker, S., Johansson, L., Aldous, L., Raucci, C., Traut, M., Ettinger, S., Nelissen, D., Lee, D. S., Ng, S., Agrawal, A., Winebrake, J. J., Hoen, M., Chesworth, S. and Pandey, A.: Third IMO GHG Study 2014, International Maritime Organization (IMO), London, UK, [online] Available from: www.imo.org, 2015.

Stein, G., Wuenstel, E. and Travnicek-Pagaimo, W.: Reifenabrieb in Feinstaub – kein Grund zur Panik!, GAK - Gummi Fasern Kunststoffe, Jahrgang 65, 441–445, 2012.

Stettler, M. E. J., Boies, A. M., Petzold, A. and Barrett, S. R. H.: Global Civil Aviation Black Carbon Emissions, *Environmental Science & Technology*, 47(18), 10397–10404, doi:10.1021/es401356v, 2013.

Stoerk, T.: Statistical corruption in Beijing's air quality data has likely ended in 2012, *Atmospheric Environment*, 127, 365–371, doi:10.1016/j.atmosenv.2015.12.055, 2016.

Stohl, A., Berg, T., Burkhardt, J. F., Fjærraa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth, K., Treffeisen, R., Virkkunen, K. and Yttri, K. E.: Arctic smoke - record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmospheric Chemistry and Physics*, 7(2), 511–534, doi:10.5194/acp-7-511-2007, 2007.

Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V. P., Kopeikin, V. M. and Novigatsky, A. N.: Black carbon in the Arctic: the underestimated role of gas flaring and residential combustion emissions, *Atmospheric Chemistry and Physics*, 13(17), 8833–8855, doi:10.5194/acp-13-8833-2013, 2013.

Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Berntsen, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinska, M., Eckhardt, S., Fuglestedt, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U., Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Olivie, D., Quaas, J., Quennehen, B., Raut, J.-C., Rumbold, S. T., Samset, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E. and Zhu, T.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmos. Chem. Phys.*, 15(18), 10529–10566, doi:10.5194/acp-15-10529-2015, 2015.

Stratus Consulting: CCAC initiative to mitigate black carbon and other pollutants from brick production: Regional Assessment. Black Carbon Mitigation in Brick Production: A Summary for Brazil, Chile, Colombia, Mexico, Nigeria, and Peru, Climate and Clean Air Coalition (CCAC), 2014.

Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J. H. and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research*, 108(D21), 1–23, doi:10.1029/2002JD003093, 2003.

Subramanian, R., Winijkul, E., Bond, T. C., Thiansathit, W., Oanh, N. T. K., Paw-armart, I. and Duleep, K. G.: Climate-Relevant Properties of Diesel Particulate Emissions: Results from a Piggyback Study in Bangkok, Thailand, *Environmental Science & Technology*, 43(11), 4213–4218, doi:10.1021/es8032296, 2009.

Takai, H., Pedersen, S., Johnsen, J. O., Metz, J. H. M., Groot Koerkamp, P. W. G., Uenk, G. H., Phillips, V. R., Holden, M. R., Sneath, R. W., Short, J. L., White, R. P., Hartung, J., Seedorf, J., Schröder, M., Linkert, K. H. and Wathes, C. M.: Concentrations and emission of airborne dust in livestock buildings in Northern Europe, *J. Agric. Engng. Res.*, 1(70), 59–77, 1998.

Tang, N. W., Apte, J. S., Martien, P. T. and Kirchstetter, T. W.: Measurement of black carbon emissions from in-use diesel-electric passenger locomotives in California, *Atmospheric Environment*, 115, 295–303, doi:10.1016/j.atmosenv.2015.05.001, 2015.

Formatted: German (Germany)

5 Tissari, J., Hytönen, K., Lyyräinen, J. and Jokiniemi, J.: A novel field measurement method for determining fine particle and gas emissions from residential wood combustion. *Atmospheric Environment*, 41, 8330–8344, 2007.

10 Tissari, J., Lyyräinen, J., Hytonen, K., Sippula, O., Tapper, U., Frey, A., Saarnio, K., Pennanen, A. S., Hillamo, R., Salonen, R. O., Hirvonen, M.-R. and Jokiniemi, J.: Fine particle and gaseous emissions from normal and smouldering wood combustion in a conventional masonry heater. *Atmos. Environ.*, 42, 7862–7873, 2008.

15 Tissari, J., Hytonen, K., Sippula, O. and Jokiniemi, J.: The effects of operating conditions on emissions from masonry heaters and sauna stoves. *Biomass and Bioenergy*, 33, 513–520, 2009.

20 Todd, J. J.: *Wood-Smoke Handbook: Woodheaters, Firewood and Operator Practice*. Eco-Energy Options, Lindisfarne Tasmania., 2003.

25 Triple E.: Diesel power generation: Inventories, black carbon emissions, initial stakeholder meeting and data mapping for Nigeria. World Bank, Washington D.C., 2013.

30 Troncoso, K., Castillo, A., Merino, L., Lazos, E. and Masera, O. R.: Understanding an improved cookstove program in rural Mexico: An analysis from the implementers' perspective. *Energy Policy*, 39(12), 7600–7608, doi:10.1016/j.enpol.2011.04.070, 2011.

35 Tsai, J. H., Chen, S. J., Huang, K. L., Lin, Y. C., Lee, W. J., Lin, C. C. and Lin, W. Y.: PM, carbon, and PAH emissions from a diesel generator fuelled with soy-biodiesel blends. *Journal of Hazardous Materials*, 179(1–3), 237–243, doi:10.1016/j.jhazmat.2010.02.085, 2010.

40 Turn, S. Q., Jenkins, B. M., Chow, J. C., Pritchett, L. C., Campbell, D., Cahill, T. and Whalem, S. A.: Elemental Characterization of Particulate Matter Emitted from Biomass Burning: Windtunnel Derived Source Profiles for Herbaceous and Wood Fuels. *Journal of Geophysical Research*, 102(D3), 3683–3699, 1997.

45 Turpin, J. T. and Lim, H.: Species contributions to PM_{2.5} mass concentrations: revisiting common assumptions for estimating organic mass. *Aerosol Science and Technology*, 35, 602–610, 2001.

50 UBOS: Uganda national household survey 2009/2010, Uganda Bureau of Statistics (UBOS), Kampala, Uganda., 2010.

55 Uma, R., Kandpal, T. C. and Kishore, V. V. N.: Emission characteristics of an electricity generation system in diesel alone and dual fuel modes. *Biomass and Bioenergy*, 27(2), 195–203, doi:10.1016/j.biombioe.2004.01.003, 2004.

60 UNEP: Near-term Climate Protection and Clean Air Benefits: Actions for Controlling Short-Lived Climate Forcers, United Nations Environment Programme (UNEP), Nairobi, Kenya. [online] Available from: http://www.unep.org/dewa/Portals/67/pdf/Near_Term_Climate_Protection_&_Air_Benefits.pdf (Accessed 12 March 2012), 2011.

65 UNEP/WMO: Integrated Assessment of Black Carbon and Tropospheric Ozone, Nairobi, Kenya. [online] Available from: www.unep.org/dewa/Portals/67/pdf/BlackCarbon_report.pdf, 2011.

70 UN-Habitat: Global Reports on Human Settlements 2005, United Nations Human Settlements Programme (UN-Habitat), Nairobi, Kenya., 2005.

75 US EPA: *Compilation of Air Pollutant Emission Factors - Volume I: Stationary Point and Area Sources*. AP-42 5th Edition., U.S. Environmental Protection Agency (US EPA), Springfield, VA., 1995.

US EPA: SPECIATE3.2 US EPA's repository of Total Organic Compound (TOC) and Particulate Matter (PM) speciated profiles for a variety of sources for use in source apportionment studies. [online] Available from: <http://www.epa.gov/ttn/chief/software/speciate/>. 2002.

US EPA: Notice of Violation. [online] Available from: <http://www3.epa.gov/otaq/cert/documents/vw-nov-2015-11-02.pdf>, 2015a.

US EPA: Notice of Violation, 2015b.

US EPA. (first): Emission Factors for Uncontrolled Industrial Diesel Engines, Section 3.3 Small Engines, United States Environmental Protection Agency (US EPA), Washington D.C., 1996.

US-EPA OTAQ: Development of Emission Rates for Light-Duty Vehicles in the Motor Vehicle Emissions Simulator (MOVES2010), Final Report, US Environmental Protection Agency, Washington, DC, USA. [online] Available from: <http://www.epa.gov/otaq/models/moves/documents/420r11011.pdf> (Accessed 5 October 2013), 2011.

Van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Peters, J. A. H. W. and Ganzeveld, L. N.: The EDGAR 3.2 Fast Track 2000 dataset (32FT2000). [online] Available from: [http://themasites.pbl.nl/images/Description_of_EDGAR_32FT2000\(v8\)_tcm61-46462.pdf](http://themasites.pbl.nl/images/Description_of_EDGAR_32FT2000(v8)_tcm61-46462.pdf), 2005.

Van Vuuren, D. P., Edmonds, J. A., Kainuma, M., Riahi, K., Thomson, A. M., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J. and Rose, S.: The representative concentration pathways: an overview, Climatic Change, 109, 5–31, doi:10.1007/s10584-011-0148-z, 2011.

Venkataraman, C., Habib, G., Eiguren-Fernandez, A., Miguel, A. H. and Friedlander, S. K.: Residential Biofuels in South Asia: Carbonaceous Aerosol Emissions and Climate Impacts, Science, 307, 1454–1456, 2005.

Venkataraman, C., Habib, G., Kadamba, D., Shrivastava, M., Leon, J. F., Crouzille, B., Boucher, O. and Streets, D. G.: Emissions from open biomass burning in India: Integrating the inventory approach with high-resolution Moderate Resolution Imaging Spectroradiometer (MODIS) active-fire and land cover data, Global Biogeochem. Cycles, 20(2), 1–12, doi:10.1029/2005GB002547, 2006.

Venkataraman, C., Sagar, A. D., Habib, G. and Smith, K. R.: The Indian National Initiative for Advanced Biomass Cookstoves: The benefits of clean combustion, Energy for Sustainable Development, 14, 63–72, 2010.

Vongmahadlek, C., Thao, P. T. B., Satayopas, B. and Thongboonchu, N.: An Inventory of Primary Gaseous Emissions from Thailand with Spatial and Temporal Allocation Profile, in Proceedings of the 3rd IASME/WSEAS International Conference on Energy & Environment, pp. 313–318, World Scientific and Engineering Academy and Society (WSEAS), Stevens Point, Wisconsin, USA. [online] Available from: <http://dl.acm.org/citation.cfm?id=1576758.1576812>, 2008.

Vongmahadlek, C., Pham, T. B. T., Satayopas, B. and Thongboonchoo, N.: A compilation and development of spatial and temporal profiles of high-resolution emissions inventory over Thailand., J Air Waste Manag Assoc, 59(7), 845–856, 2009.

Wang, S. X., Zhao, B., Cai, S. Y., Klimont, Z., Nielsen, C. P., Morikawa, T., Woo, J. H., Kim, Y., Fu, X., Xu, J. Y., Hao, J. M. and He, K. B.: Emission trends and mitigation options for air pollutants in East Asia, Atmos. Chem. Phys., 14(13), 6571–6603, doi:10.5194/acp-14-6571-2014, 2014.

Wang, S.-X., Zhao, X.-J., Li, X.-H., Wei, W. and Hao, J.-M.: Emission characteristics of fine particles from grate firing boilers, Environmental Science, 30(4), 963–968, 2009.

Weitkamp, E. A., Lipsky, E. M., Pancras, P. J., Ondov, J. M., Polidori, A., Turpin, B. J. and Robinson, A. L.: Fine particle emission profile for a large coke production facility based on highly time-resolved fence line measurements, *Atmospheric Environment*, 39(36), 6719–6733, doi:10.1016/j.atmosenv.2005.06.028, 2005.

5 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S. and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6(11), 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y. and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric Chemistry and Physics*, 10(23), 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

10 Weyant, C., Athalve, V., Ragavan, S., Rajarathnam, U., Lalchandani, D., Maithe, S., Baum, E. and Bond, T. C.: Emissions from South Asian Brick Production, *Environ. Sci. Technol.*, 48(11), 6477–6483, doi:10.1021/es500186g, 2014.

Weyant, C. L., Shepson, P. B., Subramanian, R., Cambaliza, M. O. L., Heimbürger, A., McCabe, D., Baum, E., Stirm, B. H. and Bond, T. C.: Black Carbon Emissions from Associated Natural Gas Flaring, *Environ. Sci. Technol.*, 50(4), 2075–2081, doi:10.1021/acs.est.5b04712, 2016.

15 WHO: Health aspects of air pollution: Results from the WHO project “Systematic review of Health aspects of air pollution in Europe.” Technical report, Copenhagen, Denmark. [online] Available from: <http://www.aerohabitat.org/link/2006/12-06-2006%20-%20WHO.%20Health%20aspects%20of%20air%20pollution.pdf> (Accessed 2 December 2013), 2004.

Wickramasinghe, A.: Energy access and transition to cleaner cooking fuels and technologies in Sri Lanka: Issues and policy limitations, *Energy Policy*, 39(12), 7567–7574, doi:10.1016/j.enpol.2011.07.032, 2011.

20 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J. and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geoscientific Model Development*, 4(3), 625–641, doi:10.5194/gmd-4-625-2011, 2011.

25 Wiedinmyer, C., Yokelson, R. J. and Gullett, B. K.: Global Emissions of Trace Gases, Particulate Matter, and Hazardous Air Pollutants from Open Burning of Domestic Waste, *Environ. Sci. Technol.*, 48(16), 9523–9530, doi:10.1021/es502250z, 2014.

Wobus, C., Flanner, M., Sarofim, M. C., Moura, M. C. P. and Smith, S. J.: Future Arctic temperature change resulting from a range of aerosol emissions scenarios, *Earth’s Future*, 4, doi:10.1002/2016EF000361, 2016.

World Bank: Diesel Power Generation: Inventories and Black Carbon Emissions in Kathmandu Valley, Nepal, The World Bank, Washington DC, USA., 2014a.

30 World Bank: Diesel Power Generation: Inventories and Black Carbon Emissions in Nigeria, The World Bank, Washington DC, USA., 2014b.

World Bank and ICCI: On Thin Ice: How cutting pollution can slow warming and save lives, The World Bank, The International Cryosphere Climate Initiative, Washington D.C., U.S.A., 2013.

35 Xu, Y.: Improvements in the operation of SO₂ scrubbers in China’s coal power plants, *Environ. Sci. Technol.* 45(2), 380–385, 2011.

Xu, Y., Williams, R. H. and Socolow, R. H.: China's rapid deployment of SO₂ scrubbers, *Energy Environ. Sci.*, 2(5), 459–465, doi:10.1039/B901357C, 2009.

Yan, F., Winijkul, E., Jung, S., Bond, T. C. and Streets, D. G.: Global emission projections of particulate matter (PM): I. Exhaust emissions from on-road vehicles, *Atmospheric Environment*, 45(28), 4830–4844, doi:10.1016/j.atmosenv.2011.06.018, 2011.

Yan, F., Winijkul, E., Streets, D. G., Lu, Z., Bond, T. C. and Zhang, Y.: Global emission projections for the transportation sector using dynamic technology modeling, *Atmospheric Chemistry and Physics*, 14(11), 5709–5733, doi:10.5194/acp-14-5709-2014, 2014.

Yanowitz, J., McCormick, R. L. and Graboski, M. S.: In-Use Emissions from Heavy-Duty Diesel Vehicles, *Environmental Science & Technology*, 34(5), 729–740, doi:10.1021/es990903w, 2000.

Yelverton, T. L. B., Holder, A. L. and Pavlovic, J.: Emissions removal efficiency from diesel gensets using aftermarket PM controls, *Clean Technologies and Environmental Policy*, 17(7), 1861–1871, doi:10.1007/s10098-015-0900-6, 2016.

Yim, S. H. L., Lee, G. L., Lee, I. H., Allroggen, F., Ashok, A., Caiazzo, F., Eastham, S. D., Malina, R. and Barrett, S. R. H.: Global, regional and local health impacts of civil aviation emissions, *Environmental Research Letters*, 10(3), 034001, 2015.

Yttri, K. E., Lund Myhre, C., Eckhardt, S., Fiebig, M., Dye, C., Hirdman, D., Ström, J., Klimont, Z. and Stohl, A.: Quantifying black carbon from biomass burning by means of levoglucosan – a one-year time series at the Arctic observatory Zeppelin, *Atmos. Chem. Phys.*, 14(12), 6427–6442, doi:10.5194/acp-14-6427-2014, 2014.

Zhang, Q., Klimont, Z., Streets, D., Huo, H. and He, K.: An anthropogenic PM emission model for China and emission inventory for the year 2001, *Progress in Natural Science*, 16(2), 223–231 (in Chinese), 2006.

Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T. and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, 2009.

Zhang, W., Wang, J., Zhang, B., Bi, J. and Jiang, H.: Can China Comply with Its 12th Five-Year Plan on Industrial Emissions Control: A Structural Decomposition Analysis, *Environ. Sci. Technol.*, 49(8), 4816–4824, doi:10.1021/es504529x, 2015.

Zhao, Y., Zhang, J. and Nielsen, C. P.: The effects of recent control policies on trends in emissions of anthropogenic atmospheric pollutants and CO₂ in China, *Atmos. Chem. Phys.*, 13, 487–513, 2013.

Zhi, G., Chen, Y., Feng, Y., Xiong, S., Li, J., Zhang, G., Sheng, G. and Fu, J.: Emission Characteristics of Carbonaceous Particles from Various Residential Coal-Stoves in China, *Environ. Sci. Technol.*, 42(9), 3310–3315, doi:10.1021/es702247q, 2008.

Zhi, G., Peng, C., Chen, Y., Liu, D., Sheng, G. and Fu, J.: Deployment of Coal Briquettes and Improved Stoves: Possibly an Option for both Environment and Climate, *Environ. Sci. Technol.*, 43(15), 5586–5591, doi:10.1021/es802955d, 2009.

Aasestad, K.: Emissions of Black carbon and Organic carbon in Norway 1990–2011, Statistics Norway, Oslo, Norway., 2013.

Adria, O. and Bethge, J.: What users can save with energy efficient cooking stoves and ovens Oliver Adria (CSCP) Jan Bethge, Wuppertal Institute for Climate, Environment and Energy, Wuppertal, Germany., 2013.

Formatted: German (Germany)

- Aghalino, S.: Gas flaring, environmental pollution and abatement measures in Nigeria, 1969-2001, *Journal of Sustainable Development in Africa*, 11, 219–238, 2009.
- Agrawal, H., Malloy, Q.-G. J., Welch, W. A., Wayne Miller, J. and Cocker III, D. R.: In-use gaseous and particulate matter emissions from a modern ocean going container vessel, *Atmospheric Environment*, 42(21), 5504–5510, 2008.
- 5 Agrawal, H., Welch, W. A., Henningsen, S., Miller, J. W. and III, D. R. C.: Emissions from main propulsion engine on container ship at sea, *J. Geophys. Res.*, 115, 7 PP., doi:201010.1029/2009JD013346, 2010.
- Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U. and Jimenez, J. L.: O/C and OM/OC Ratios of Primary, Secondary, and Ambient Organic Aerosols with High Resolution Time-of-Flight Aerosol Mass Spectrometry, *Environ. Sci. Technol.*, 42(12), 4478–4485, doi:10.1021/es703009q, 2008.
- 10 AIT: Small and Medium scale Industries in Asia: Energy and Environment. Brick and Ceramic Sectors, Regional Energy Resources Information Center (RERIC), Asian Institute of Technology (AIT), Pathumthani, Thailand., 2003.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crouse, J. D. and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmospheric Chemistry and Physics*, 11(9), 4039–4072, doi:10.5194/acp-11-4039-2011, 2011.
- 15 Amann, M., Bertok, I., Borken, J., Chambers, A., Cofala, J., Dentener, F., Heyes, C., Kejun, J., Klimont, Z., Makowski, M., Matur, R., Purohit, P., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F. and Winiwarter, W.: GAINS Asia. A tool to combat air pollution and climate change simultaneously, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria., 2008.
- 20 Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F. and Winiwarter, W.: Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications, *Environmental Modelling & Software*, 26(2), 1489–1501, 2011.
- 25 Amann, M., Borken-Kleefeld, J., Cofala, J., Heyes, C., Klimont, Z., Rafaj, P., Purohit, P., Schoepp, W. and Winiwarter, W.: Future emissions of air pollutants in Europe—Current legislation baseline and the scope for further reductions. TSAP Report #1, International Institute for Applied Systems Analysis, Laxenburg. [online] Available from: <http://gains.iiasa.ac.at/images/stories/reports/TSAP/TSAP-BASELINE-20120613.pdf> (Accessed 13-December 2012), 2012.
- 30 Amann, M., Klimont, Z. and Wagner, F.: Regional and Global Emissions of Air Pollutants: Recent Trends and Future Scenarios, *Annual Review of Environment and Resources*, 38(1), 31–55, doi:10.1146/annurev-environ-052912-173303, 2013.
- Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Kiesewetter, G., Klimont, Z., Schöpp, W., Vellinga, N. and Winiwarter, W.: Adjusted historic emission data, projections, and optimized emission reduction targets for 2030—A comparison with COM data 2013, IIASA, Laxenburg, Austria. [online] Available from: http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf, 2015.
- 35 AMAP: AMAP Assessment 2015: Black carbon and ozone as Arctic climate foreers, Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway. [online] Available from: <http://www.amap.no/documents/doc/amap-assessment-2015-black-carbon-and-ozone-as-arctic-climate-foreers/1299>, 2015.

Anand, M.: Diesel Pricing in India: Entangled in Policy Maze, National Institute of Public Finance and Policy (NIPFP), New Delhi. [online] Available from: <http://www.nipfp.org.in/media/medialibrary/2013/04/Diesel%20Price%20Reform.pdf>; 2012.

5 Anayoehukwu, A. V., Naene, E. A. and Onyeka, A. E.: Assessment of Environmental Impact of Power Generation in Banking Industry, *Journal of Energy, Environment & Carbon Credits*, 3(1), 13–21, 2013.

Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochemical Cycles*, 15(4), 955–966, doi:10.1029/2000GB001382, 2001.

10 Anenberg, S. C., Schwartz, J., Shindell, D., Amann, M., Faluvegi, G., Klimont, Z., Janssens-Maenhout, G., Pozzoli, L., Van Dingenen, R., Vignati, E., Emberson, L., Müller, N. Z., West, J. J., Williams, M., Demkine, V., Hicks, W. K., Kuylensstierna, J., Raes, F. and Ramanathan, V.: Global Air Quality and Health Co-benefits of Mitigating Near-Term Climate Change through Methane and Black Carbon Emission Controls, *Environ Health Perspect*, 120(6), 831–839, doi:10.1289/ehp.1104301, 2012.

Apple, J., Vicente, R., Yarberry, A., Lohse, N., Mills, E., Jacobson, A. and Poppendieck, D.: Characterization of particulate matter size distributions and indoor concentrations from kerosene and diesel lamps, *Indoor Air*, 20(5), 399–411, 2010.

15 Ban-Weiss, G. A., Lunden, M. M., Kirchstetter, T. W. and Harley, R. A.: Measurement of Black Carbon and Particle Number Emission Factors from Individual Heavy Duty Trucks, *Environmental Science & Technology*, 43(5), 1419–1424, doi:10.1021/es8021039, 2009.

20 Boman, C., Pettersson, E., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 1: Pellet Stoves, *Energy Fuels*, 25(1), 307–314, doi:10.1021/ef100774x, 2011.

Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H. and Klimont, Z.: A technology based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109(D14), 1–43, doi:10.1029/2003JD003697, 2004.

25 Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G. and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy related combustion, 1850–2000, *Global Biogeochem. Cycles*, 21(2), 1–16, doi:10.1029/2006GB002840, 2007.

30 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Bernsten, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G. and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.

35 Bonjour, S., Adair-Rohani, H., Wolf, J., Bruce, N. G., Mehta, S., Pruss-Ustun, A., Lahiff, M., Rehfuess, E. A., Mishra, V. and Smith, K. R.: Solid fuel use for household cooking: country and regional estimates for, *Environ Health Perspect*, 121(7), 784–790, doi:10.1289/ehp.1205987, 2013.

Borken-Kleefeld, J. and Chen, Y.: New emission deterioration rates for gasoline cars—Results from long term measurements, *Atmospheric Environment*, 101, 58–64, doi:10.1016/j.atmosenv.2014.11.013, 2015.

Brohan, P., Kennedy, J. J., Harris, I., Tett, S. F. B. and Jones, P. D.: Uncertainty estimates in regional and global observed temperature changes: A new data set from 1850, *Journal of Geophysical Research: Atmospheres*, 111(D12), n/a–n/a, doi:10.1029/2005JD006548, 2006.

Buhaus, O., Corbett, J. J., Endresen, O., Eyring, V., Faber, J., Hanayama, S., Lee, D. S., Lee, D., Lindstad, H., Markowska, A. Z., Mjelde, A., Nelissen, D., Nilsen, J., Palsson, C., Winebrake, J. J., Wu, W. Q. and Yoshida, K.: Second IMO GHG study 2009, IMO, London, UK., 2009.

Cadle, S., Ayala, A., Black, K., Graze, R., Koupal, J., Minassian, F., Murray, H., Natarajan, M., Zhu, F., Takaoka, M., Oshita, K., Morisawa, S., Tsuno, H., Takaoka, M., Oshita, K., Morisawa, S., Tsuno, H., Kitajima, Y., Chiang, W. F., Fang, H. Y., Fang, H. Y., Wu, C. H., Huang, C. J., Huang, C. J., Chang, C. Y., Chang, Y. M., Chen, C. L., Nielsen, A., Nielsen, L., Feilberg, A., Christensen, K., Liu, Y. Y., Lin, T. C., Lin, T. C., Wang, Y. J., Ho, W. L., Yanowitz, J., McCormick, R., Yu, L., Jia, S., Shi, Q., Chang, T. J., Kao, H. M., Wu, Y. T., Huang, W. H., Kao, H. M., Wu, Y. T., Huang, W. H., Goodman, P., Rich, D., Zeka, A., Clancy, L., Doekery, D., Lavery, T., Rogers, C., Baumgardner, R., Mishoe, K., Chen, W. C., Lin, H. Y., Yuan, C. S., Hung, C. H., He, L., Huang, G., Zeng, G. and Lu, H.: Real World Vehicle Emissions: A Summary of the 18th Coordinating Research Council On Road Vehicle Emissions Workshop, *Journal of the Air & Waste Management Association*, 59(2), 130–138, doi:10.3155/1047-3289.59.2.130, 2009.

Cadle, S. H., Mulawa, P. A., Ball, J., Donase, C., Weibel, A., Sagebiel, J. C., Knapp, K. T. and Snow, R.: Particulate Emission Rates from In-Use High-Emitting Vehicles Recruited in Orange County, California, *Environmental Science & Technology*, 31(12), 3405–3412, doi:10.1021/es9700257, 1997.

CAI Asia: Factsheet No. 17 Roadmap for Cleaner Fuels and Vehicles in Asia, 2011.

Cames, M. and Helmers, E.: Critical evaluation of the European diesel car boom—global comparison, environmental effects and various national strategies, *Environmental Sciences Europe*, 25(1), 15–36, doi:10.1186/2190-4715-25-15, 2013.

Cao, G., Zhang, X. and Zheng, F.: Inventory of black carbon and organic carbon emissions from China, *Atmospheric Environment*, 40(34), 6516–6527, 2006.

Cao, G., Zhang, X., Wang, Y. and Zheng, F.: Estimation of emissions from field burning of crop straw in China, *Chinese Science Bulletin*, 53(5), 784–790, 2008.

CAPP: A Recommended Approach to Completing the National Pollutant Release Inventory (NPRI) for the Upstream Oil and Gas Industry, Canadian Association of Petroleum Producers (CAPP). [online] Available from: <http://www.capp.ca/library/publications/policyRegulatory/pages/pubInfo.aspx?DoelD=119572>, 2007.

Cardenas, B., Maiz, P., Márquez, R. O., Munguia, J. L., Angeles, F., Baum, E. and Molina, L. T.: Determining Emissions of Black carbon, greenhouse gases and other pollutants from artisanal brick production in Mexico, *Clean Air Task Force and MCE2*, 2012.

Carslaw, D. C., Beevers, S. D., Tate, J. E., Westmoreland, E. J. and Williams, M. L.: Recent evidence concerning higher NO_x emissions from passenger cars and light duty vehicles, *Atmospheric Environment*, 45(39), 7053–7063, doi:10.1016/j.atmosenv.2011.09.063, 2011.

Castellanos, P., Boersma, K. F. and van der Werf, G. R.: Satellite observations indicate substantial spatiotemporal variability in biomass burning NO_x emission factors for South America, *Atmospheric Chemistry and Physics*, 14(8), 3929–3943, doi:10.5194/acp-14-3929-2014, 2014.

CEPMEIP: CEPMEIP (Co-ordinated European Programme on Particulate Matter Emission Inventories, Projections and Guidance), [online]. Available from: <http://www.air.sk/tno/cepmcip/>, 2002.

Chafe, Z. A., Brauer, M., Klimont, Z., Van Dingenen, R., Mehta, S., Rao, S., Riahi, K., Dentener, F. and Smith, K. R.: Household Cooking with Solid Fuels Contributes to Ambient PM_{2.5} Air Pollution and the Burden of Disease, *Environmental Health Perspectives*, 122(12), 1314–1320, 2014.

Chafe, Z. A., Brauer, M., Héroux, M.-E., Klimont, Z., Lanki, T., Salonen, R. O. and Smith, K. R.: Residential heating with wood and coal: health impacts and policy options in Europe and North America, World Health Organization. [online]. Available from: <http://www.euro.who.int/en/health-topics/environment-and-health/air-quality/publications/2015/residential-heating-with-wood-and-coal-health-impacts-and-policy-options-in-europe-and-north-america>, 2015.

Chen, Y. and Borken-Kleeefeld, J.: Real driving emissions from cars and light commercial vehicles — Results from 13 years remote sensing at Zurich/CH, *Atmospheric Environment*, 88, 157–164, doi:10.1016/j.atmosenv.2014.01.040, 2014.

Chen, Y. and Borken-Kleeefeld, J.: NO_x Emissions from Diesel Passenger Cars Worsen with Age, *Environmental Science & Technology*, doi:10.1021/acs.est.5b04704, 2016.

Chen, Y., Zhi, G., Feng, Y., Liu, D., Zhang, G., Li, J., Sheng, G. and Fu, J.: Measurements of Black and Organic Carbon Emission Factors for Household Coal Combustion in China: Implication for Emission Reduction, *Environ. Sci. Technol.*, 43(24), 9495–9500, doi:10.1021/es9021766, 2009.

Chen, Y., Morton, D. C., Jin, Y., Collatz, G. J., Kasibhatla, P. S., van der Werf, G. R., DeFries, R. S. and Randerson, J. T.: Long-term trends and interannual variability of forest, savanna and agricultural fires in South America, *Carbon Management*, 4(6), 617–638, doi:10.4155/cmt.13.61, 2013.

Cheung, K. L., Polidori, A., Ntziachristos, L., Tzamkiozis, T., Samaras, Z., Cassee, F. R., Gerlofs, M. and Sioutas, C.: Chemical characteristics and oxidative potential of particulate matter emissions from gasoline, diesel, and biodiesel cars, *Environmental science & technology*, 43(16), 6334–6340, 2009.

Christian, T. J., Yokelson, R. J., Cárdenas, B., Molina, L. T., Engling, G. and Hsu, S. C.: Trace gas and particle emissions from domestic and industrial biofuel use and garbage burning in central Mexico, *Atmos. Chem. Phys.*, 10(2), 565–584, doi:10.5194/acp-10-565-2010, 2010.

Cofala, J. and Syri, S.: Sulfur emissions, abatement technologies and related costs for Europe in the RAINS model database, Interim Report, IIASA, Laxenburg, Austria. [online]. Available from: <http://web.archive.iiasa.ac.at/Publications/Documents/IR-98-035.pdf> (Accessed 14 May 2013), 1998.

Cofala, J., Amann, M., Klimont, Z., Kupiainen, K. and Höglund-Isaksson, L.: Scenarios of Global Anthropogenic Emissions of Air Pollutants and Methane Until 2030, *Atmospheric Environment*, Vol 41/38, 8486–8499, 2007.

Conrad, B. and Johnson, M.: Field Measurements of Black Carbon Yields from Gas Flaring, *Environ. Sci. Technol.*, in review.

Corbett, J. J. and Koehler, H. W.: Updated emissions from ocean shipping, *J. Geophys. Res.*, 108, 15 PP., doi:10.1029/2003JD003751, 2003.

CPCB: Comprehensive industry document with emission standards, guidelines and stack height regulation for vertical shaft brick kilns (VSBK) vis-à-vis pollution control measures, Central Pollution Control Board (CPCB), New Delhi, India., 2007.

Crippa, M., Janssens Maenhout, G., Dentener, F., Guizzardi, D., Sindelarova, K., Muntean, M., Van Dingenen, R. and Granier, C.: Forty years of improvements in European air quality: regional policy-industry interactions with global impacts, *Atmospheric Chemistry and Physics*, 16(6), 3825–3841, doi:10.5194/acp-16-3825-2016, 2016.

5 Dalsoren, S. B., Eide, M., Endresen, U., Mjelde, A., Gravir, G. and Isaksen, I. S. A.: Update on emissions and environmental impacts from the international fleet of ships: the contribution from major ship types and ports, *Atmospheric Chemistry and Physics*, 9(6), 2171–2194, 2009.

Delphi Inc.: *Worldwide Emission Standards—Heavy Duty and Off-Highway Vehicles 2013/14*, Delphi Inc., Troy, MICH/USA., 2013.

10 Delphi Inc.: *Worldwide Emission Standards—Passenger Cars and Light Duty Vehicles 2014/15*, Delphi Inc., Troy, MICH/USA., 2015.

Denier van der Gon, H. A. C., Gerlofs Nijland, M. E., Gehrig, R., Gustafsson, M., Janssen, N., Harrison, R. M., Hulskotte, J., Johansson, C., Jozwička, M., Keuken, M., Krijgheld, K., Ntziachristos, L., Riediker, M. and Cassee, F. R.: The Policy Relevance of Wear Emissions from Road Transport, Now and in the Future—An International Workshop Report and Consensus Statement, *Journal of the Air & Waste Management Association*, 63(2), 136–149, doi:10.1080/10962247.2012.741055, 2013.

15 Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D. and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe—revised estimates and an evaluation, *Atmospheric Chemistry and Physics*, 15(11), 6503–6519, doi:10.5194/acp-15-6503-2015, 2015.

20 Desai, S., Dubey, A., Joshi, B. L., Sen, M., Shariff, A. and Vanneman, R.: *Human Development in India: Challenges for a Society in Transition*, Oxford University Press, New Delhi, India., 2010.

Driscoll, T., Milkovits, G. and Fredenberger, D.: *Impact and Use of Firewood in Australia*, CSIRO Sustainable Ecosystems., 2000.

Durbin, T. D., Norbeck, J. M., Smith, M. R. and Truex, T. J.: Particulate Emission Rates from Light Duty Vehicles in the South Coast Air Quality Management District, *Environ. Sci. & Technol.*, 33, 4401–4406, 1999.

25 EAWAG: *Global waste challenge: Situation in developing countries*, 2008.

Ebel, A., Friedrich, R. and Rodhe, H.: GENEMIS: Assessment, Improvement, and Temporal and Spatial Disaggregation of European Emission Data, in *Tropospheric Modelling and Emission Estimation: Chemical Transport and Emission Modelling on Regional, Global and Urban Scales*, edited by A. Ebel, R. Friedrich, and H. Rodhe, pp. 181–214, Springer Berlin Heidelberg, Berlin, Heidelberg. [online] Available from: http://dx.doi.org/10.1007/978-3-662-03470-5_6, 1997.

30 EC: Directive 2001/80/EC of the European Parliament and of the Council of 23 October 2001 on the limitation of emissions of certain pollutants into the air from large combustion plants, 2001a.

EC: Directive 2001/81/EC of the European Parliament and of the Council of 23 October 2001 on national emission ceilings for certain atmospheric pollutants, European Parliament and Council, Luxembourg., 2001b.

35 EC: Directive 2010/75/EU of the European Parliament and of the council on industrial emissions (integrated pollution prevention and control), Commission of the European Communities, Brussels, Belgium. [online] Available from: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2010:334:0017:0119:EN:PDF>, 2010.

EC JRC/PBL: Emission Database for Global Atmospheric Research (EDGAR), European Commission (EC), Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL). [online] Available from: <http://edgar.jrc.ec.europa.eu>; 2010.

5 Eckhardt, S., Quennehen, B., Ollivié, D. J. L., Berntsen, T. K., Cherian, R., Christensen, J. H., Collins, W., Crepinsek, S., Daskalakis, N., Flanner, M., Herber, A., Heyes, C., Hodnebrog, Ø., Huang, L., Kanakidou, M., Klimont, Z., Langner, J., Law, K. S., Lund, M. T., Mahmood, R., Massling, A., Myriokefalitakis, S., Nielsen, I. E., Nøjgaard, J. K., Quaas, J., Quinn, P. K., Raut, J. C., Rumbold, S. T., Schulz, M., Sharma, S., Skeie, R. B., Skov, H., Uttal, T., von Salzen, K. and Stohl, A.: Current model capabilities for simulating black carbon and sulfate concentrations in the Arctic atmosphere: a multi model evaluation using a comprehensive measurement data set, *Atmos. Chem. Phys.*, 15(16), 9413–9433, doi:10.5194/acp-15-9413-2015, 2015.

10 Edwards, P.: Global cement emissions standards, *Global Cement Magazine* [online] Available from: www.GlobalCement.com, 2014.

EEA: EMEP/EEA air pollutant emission inventory guidebook, Technical report, European Environmental Agency (EEA), Copenhagen, Denmark. [online] Available from: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>, 2013.

15 Ekström, M., Sjödin, Å. and Andreasson, K.: Evaluation of the COPERT III emission model with on-road optical remote sensing measurements, *Atmospheric Environment*, 38(38), 6631–6641, doi:10.1016/j.atmosenv.2004.07.019, 2004.

Elisabeth, T.: Black Carbon Emissions from Kerosene Lamps: Potential for a new CCAC Initiative, Ecologic Institute, Berlin, Germany. [online] Available from: http://www.ecologic.eu/sites/files/publication/2014/black-carbon-and-kerosene-lamps-study_0.pdf, 2013.

20 Elvidge, C. D., Ziskin, D., Baugh, K. E., Tuttle, B. T., Ghosh, T., Paek, D. W., Erwin, E. H. and Zhizhin, M.: A Fifteen Year Record of Global Natural Gas Flaring Derived from Satellite Data, *Energies*, 2(3), 595, doi:10.3390/en20300595, 2009.

Elvidge, C. D., Baugh, K. E., Anderson, S., Ghosh, T. and Ziskin, D.: Estimation of Gas Flaring Volumes Using NASA MODIS Fire Detection Products, NOAA National Geophysical Data Center, Boulder, US. [online] Available from: http://www.ngdc.noaa.gov/dmsp/interest/gas_flares.html, 2011.

25 Elvidge, C. D., Zhizhin, M., Hsu, F. C. and Baugh, K. E.: VIIRS Nightfire: Satellite Pyrometry at Night, *Remote Sensing*, 5(9), 4423, doi:10.3390/rs5094423, 2013.

Endresen, Ø., Sorgård, E., Behrens, H. L., Brett, P. O. and Isaksen, I. S. A.: A historical reconstruction of ships' fuel consumption and emissions, *Journal of Geophysical Research: Atmospheres*, 112(D12), n/a–n/a, doi:10.1029/2006JD007630, 2007.

30 ESMAP: Power Sector Reform in Africa: Assessing Impact on Poor People, Energy Sector Management Assistance Programme (ESMAP), World Bank, Washington D.C., 2005.

EUROSTAT: Statistics database, [online] Available from: http://epp.eurostat.ec.europa.eu/portal/page/portal/statistics/search_database (Accessed 5 September 2011), 2011.

35 Eyring, V., Isaksen, I. S. A., Berntsen, T., Collins, W. J., Corbett, J. J., Endresen, Ø., Grainger, R. G., Moldanova, J., Schlager, H. and Stevenson, D. S.: Transport impacts on atmosphere and climate: Shipping, *Atmos. Environ.*, 44, 4735–4771, doi:10.1016/j.atmosenv.2009.04.059, 2010.

- 5
10
15
20
25
30
35
- FAO: Status and development issues of the brick industry in Asia, Field Document, FAO Regional Wood Energy Development Programme in Asia, Bangkok, Thailand., 1993.
- Fernandes, S. D., Trautmann, N. M., Streets, D. G., Roden, C. A. and Bond, T. C.: Global biofuel use, 1850–2000, *Global Biogeochem. Cycles*, 21(2), 1–15, doi:10.1029/2006GB002836, 2007.
- Foell, W., Pachauri, S., Spreng, D. and Zerriffi, H.: Household cooking fuels and technologies in developing economies, *Energy Policy*, 39(12), 7487–7496, doi:10.1016/j.enpol.2011.08.016, 2011.
- Fortner, E. C., Brooks, W. A., Onasch, T. B., Canagaratna, M. R., Massoli, P., Jayne, J. T., Franklin, J. P., Knighton, W. B., Wormhoudt, J., Worsnop, D. R., Kolb, C. E. and Herndon, S. C.: Particulate Emissions Measured During the TCEQ Comprehensive Flare Emission Study, *Ind. Eng. Chem. Res.*, 51(39), 12586–12592, doi:10.1021/ie202692y, 2012.
- Frost & Sullivan: Indian Diesel Generator Set Market. [online] Available from: <http://www.frost.com/sublib/frost-content.do?sheetName=report-overview&sheetGroup=P420-01-00-00-00&viewName=virtual-brochure&repid=P420-01-00-00-00>, 2010.
- Gadhavi, H. S., Renuka, K., Ravi Kiran, V., Jayaraman, A., Stohl, A., Klimont, Z. and Beig, G.: Evaluation of black carbon emission inventories using a Lagrangian dispersion model—a case study over southern India, *Atmos. Chem. Phys.*, 15(3), 1447–1461, doi:10.5194/acp-15-1447-2015, 2015.
- GEA: Global Energy Assessment: Toward a Sustainable Future, Cambridge University Press, UK., 2012.
- Geller, M. D., Ntziachristos, L., Mamakos, A., Samaras, Z., Schmitz, D. A., Froines, J. R. and Sioutas, C.: Physicochemical and redox characteristics of particulate matter (PM) emitted from gasoline and diesel passenger cars, *Atmospheric Environment*, 40(36), 6988–7004, doi:10.1016/j.atmosenv.2006.06.018, 2006.
- Germain, A., Granger, F. and Gosselin, A.: A Model Municipal By-Law for regulating wood burning appliances A. Germain, F. Granger & A. Gosselin, *WIT Transactions on Ecology and the Environment*, 116, doi:10.2495/AIR080631, 2008.
- Gilmore, E. A., Lave, L. B. and Adams, P. J.: The costs, air quality and human health effects of meeting peak electricity demand with installed backup generators, *Environmental Science & Technology*, 40(22), 6887–6893, 2006.
- Goel, R. and Guttikunda, S. K.: Role of urban growth, technology, and judicial interventions on vehicle exhaust emissions in Delhi for 1991–2014 and 2014–2030 periods, *Environmental Development*, 14, 6–21, doi:10.1016/j.envdev.2015.03.002, 2015.
- GOI: 2011 Census of India: Source of Lighting, Ministry of Home Affairs, Government of India (GOI), New Delhi., 2011.
- Granier, C., Bessagnet, B., Bond, T., D’Angiola, A., Denier van der Gon, H., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J. F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J. C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., Aardenne, J., Werf, G. R. and Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109(1–2), 163–190, doi:10.1007/s10584-011-0154-1, 2011.
- Guo, H., Zhang, Q., Shi, Y. and Wang, D.: On-road remote sensing measurements and fuel-based motor vehicle emission inventory in Hangzhou, China, *Atmospheric Environment*, 41(14), 3095–3107, 2007.

- Guttikunda, S. K. and Jawahar, P.: Atmospheric emissions and pollution from the coal fired thermal power plants in India, *Atmospheric Environment*, 92, 449–460, doi:10.1016/j.atmosenv.2014.04.057, 2014.
- Guttikunda, S. K., Begum, B. A. and Wadud, Z.: Particulate pollution from brick kiln clusters in the Greater Dhaka region, Bangladesh, *Air Quality, Atmosphere & Health*, 6(2), 357–365, doi:10.1007/s11869-012-0187-2, 2013.
- 5 Habib, G., Venkataraman, C., Bond, T. C. and Schauer, J. J.: Chemical, Microphysical and Optical Properties of Primary Particles from the Combustion of Biomass Fuels, *Environ. Sci. Technol.*, 42(23), 8829–8834, doi:10.1021/es800943f, 2008.
- Harrison, R. M., Jones, A. M., Gietl, J., Yin, J. and Green, D. C.: Estimation of the Contributions of Brake Dust, Tire Wear, and Resuspension to Nonexhaust Traffic Particles Derived from Atmospheric Measurements, *Environmental Science & Technology*, 46(12), 6523–6529, doi:10.1021/es300894r, 2012.
- 10 Haugland, T., Saunier, S., Pederstad, A., Holm, T., Darani, H. and Kertesheva, A.: Associated Petroleum Gas Flaring Study for Russia, Kazakhstan, Turkmenistan and Azerbaijan, Final Report, Carbon Limits AS, Oslo, Norway. [online] Available from: www.carbonlimits.no, 2013.
- HBEFA 3.1: HBEFA Handbook Emission Factors for Road Transport, [online] Available from: <http://www.hbefa.net/d/index.html> (Accessed 20 March 2012), 2010.
- 15 Hegg, D. A., Livingston, J., Hobbs, P. V., Novakov, T. and Russell, P.: Chemical apportionment of aerosol column optical depth off the mid-Atlantic coast of the United States, *Journal of Geophysical Research: Atmospheres*, 102(D21), 25293–25303, doi:10.1029/97JD02293, 1997.
- Heierli, U. and Maithel, S.: Brick by Brick: The Herculean Task of Cleaning of the Asian Brick Industry, Swiss Agency for Development and Cooperation (SDC), Bern., 2008.
- 20 Herzog, J. W.: Current and near-term emission control strategies for diesel powered generator sets, in *Proceedings of 24th International Telecommunications Energy Conference (INTELEC)*, pp. 394–399, Montréal., 2002.
- Hodzie, A., Wiedinmyer, C., Salcedo, D. and Jimenez, J. L.: Impact of Trash Burning on Air Quality in Mexico City, *Environ. Sci. Technol.*, 46(9), 4950–4957, doi:10.1021/es203954r, 2012.
- Hsu, Y. and Mullen, M.: Compilation of Diesel Emissions Speciation Data, Final Report for CRC., 2007.
- 25 Huang, K., Fu, J. S., Prikhodko, V. Y., Storey, J. M., Romanov, A., Hodson, E. L., Cresko, J., Morozova, I., Ignatieva, Y. and Cabaniss, J.: Russian anthropogenic black carbon: Emission reconstruction and Arctic black carbon simulation, *Journal of Geophysical Research: Atmospheres*, 120(21), 11,306–11,333, doi:10.1002/2015JD023358, 2015.
- Huo, H., Zhang, Q., He, K., Yao, Z., Wang, X., Zheng, B., Streets, D. G., Wang, Q. and Ding, Y.: Modeling vehicle emissions in different types of Chinese cities: Importance of vehicle fleet and local features, *Environmental Pollution*, 159(10), 2954–2960, doi:10.1016/j.envpol.2011.04.025, 2011.
- 30 Huo, H., Lei, Y., Zhang, Q., Zhao, L. and He, K.: China's coke industry: Recent policies, technology shift, and implication for energy and the environment, *Energy Policy*, 51(0), 397–404, doi:10.1016/j.enpol.2012.08.041, 2012.
- 35 ICC & SRI: Atmospheric emissions of particulates from agriculture: a scoping study, Final report for the Ministry of Agriculture, Fisheries and Food (MAFF) Research and Development, ICCConsultants and Silsoe Research Institute, London, UK., 2000.

- ICCT & Dieselnet: Transportpolicy.net, [online] Available from: http://transportpolicy.net/index.php?title=Main_Page (Accessed 15 March 2014), 2014.
- IEA: Coal Power 2: World coal fired power stations and their pollution control systems, International Energy Agency, London, UK., 1997.
- 5 IEA: World Energy Outlook 2007, International Energy Agency, Paris, France., 2007.
- IEA: World Energy Outlook 2011, International Energy Agency, Paris, France., 2011.
- IEA: World Energy Outlook 2012, International Energy Agency, Paris, France., 2012.
- IEA: Energy Statistics of Non-OECD Countries 2015, International Energy Agency, IEA/OECD, Paris, France., 2015a.
- IEA: Energy Statistics of OECD Countries 2015, International Energy Agency, IEA/OECD, Paris, France., 2015b.
- 10 IEA: World Energy Outlook Special Report: Energy and Air Pollution, International Energy Agency (IEA), Paris, France., 2016.
- IEA CCC: Coal Power. Coal Power Database, IEA Clean Coal Centre, London, UK., 2012.
- IEA/ETP: Energy Technology Perspectives 2012, International Energy Agency (IEA), Paris., 2012.
- 15 IFC/WB: Lighting Africa market assessment results: quantitative assessment—Ethiopia, International Finance Corporation (IFC) and the World Bank (WB), Washington D.C., 2008.
- HDFC: Environmental Management Framework for improving kiln efficiency in the brick making industry in Bangladesh; Hybrid Hoffman Kiln (HHK) project, Industrial and Infrastructure Development Finance Company Ltd. (HDFC), Dhaka, Bangladesh., 2009.
- 20 IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Intergovernmental Panel on Climate Change, IGES, Japan. [online] Available from: <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm>, 2006a.
- IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, Waste, Intergovernmental Panel on Climate Change (IPCC), Kanagawa, Japan. [online] Available from: <http://www.ipee-nggip.iges.or.jp/public/2006gl/index.htm>, 2006b.
- 25 Ite, A. and Ibok, U. J.: Gas flaring and venting associated with petroleum exploration and production in the Nigeria's Niger Delta, American Journal of Environmental Protection, 1, 70–77, 2013.
- Jacobson, A., Lam, N. L., Bond, T. C. and Hultman, N.: Black Carbon and Kerosene Light ing: An Opportunity for Rapid Action on Climate Change and Clean Energy for Development, Policy Paper 2013-03, The Brookings Institution, Washington D.C., 2013.
- 30 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B. and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, Atmospheric Chemistry and Physics, 15(19), 11411–11432, doi:10.5194/acp-15-11411-2015, 2015.

Jiang, M., Marr, L. C., Dunlea, E. J., Herndon, S. C., Jayne, J. T., Kolb, C. E., Knighton, W. B., Rogers, T. M., Zavala, M., Molina, L. T. and Molina, M. J.: Vehicle fleet emissions of black carbon, polycyclic aromatic hydrocarbons, and other pollutants measured by a mobile laboratory in Mexico City, *Atmospheric Chemistry and Physics*, 5(12), 3377–3387, doi:10.5194/acp-5-3377-2005, 2005.

5 Jimenez, J. L., McRae, G. J., Nelson, D., Zahniser, M. S. and Kolb, C. E.: Remote Sensing of NO and NO₂ Emissions from Heavy-Duty Diesel Truck Using Tunable Diode Lasers, *Environ. Sci. & Technol.*, 33, 2380–2387, 2000.

Johnson, G. R., Jayaratne, E. R., Lau, J., Thomas, V., Juwono, A. M., Kitchen, B. and Morawska, L.: Remote measurement of diesel locomotive emission factors and particle size distributions, *Atmospheric Environment*, 81, 148–157, doi:10.1016/j.atmosenv.2013.09.019, 2013.

10 Johnson, M., Edwards, R., Alatorre-Frenk, C. and Masera, O.: In field greenhouse gas emissions from cookstoves in rural Mexican households, *Atmospheric Environment*, 42(6), 1206–1222, doi:10.1016/j.atmosenv.2007.10.034, 2008.

Johnson, M. R., Devillers, R. W. and Thomson, K. A.: Quantitative Field Measurement of Soot Emission from a Large Gas Flare Using Sky LOSA., *Environ. Sci. Technol.*, 45(1), 345–350, 2011.

KEEL: Yearbook of Energy Statistics 2011, Korea Energy Economics Institute (KEEI), South Korea., 2011.

15 Kholod, N., Evans, M. and Kuklinski, T.: Russia's black carbon emissions: focus on diesel sources, *Atmospheric Chemistry and Physics Discussions*, 1–27, doi:10.5194/acp-2016-475, 2016.

Kinsey, J. S.: Characterization of emissions from commercial aircraft engines during the aircraft particle emissions experiment (APEX) 1–3, Technical report, US Environmental Protection Agency Office of Research and Development National Risk Management Research Laboratory, Washington DC, USA. [online] Available from: <http://nepis.epa.gov/Adobe/PDF/P1005KRK.pdf>, 2009.

20 Kirchstetter, T. W., Harley, R. A., Kreisberg, N. M., Stolzenburg, M. R. and Hering, S. V.: On road measurement of fine particle and nitrogen oxide emissions from light and heavy duty motor vehicles, *Atmospheric Environment*, 33(18), 2955–2968, doi:10.1016/S1352-2310(99)00089-8, 1999.

25 Klimont, Z., Hoeglund-Isaksson, L., Heyes, C., Rafaj, P., Schoepp, W., Cofala, J., Purohit, P., Borken-Kleefeld, J., Kupiainen, K., Kiesewetter, G., Winiwarter, W., Amann, M., Zhao, B., Wang, S., Bertok, I. and Sander, R.: Global scenarios of air pollutants and methane: 1990–2050, in preparation.

Klimont, Z., Streets, D. G., Gupta, S., Cofala, J., Lixin, F. and Ichikawa, Y.: Anthropogenic Emissions of Non Methane Volatile Organic Compounds (NMVOC) in China, *Atmospheric Environment*, 36(8), 1309–1322, 2002a.

30 Klimont, Z., Cofala, J., Bertok, I., Amann, M., Heyes, C. and Gyarmas, F.: Modelling Particulate Emissions in Europe. A Framework to Estimate Reduction Potential and Control Costs, Interim Report, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria., 2002b.

Klimont, Z., Cofala, J., Xing, J., Wei, W., Zhang, C., Wang, S., Kejun, J., Bhandari, P., Mathur, R., Purohit, P., Rafaj, P., Chambers, A., Amann, M. and Hao, J.: Projections of SO₂, NO_x, and carbonaceous aerosols emissions in Asia, *Tellus*, 61B(4), 602–617, doi:10.1111/j.1600-0889.2009.00428.x, 2009.

35 Klimont, Z., Smith, S. J. and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions, *Environ. Res. Lett.*, 8(1), 14003, doi:10.1088/1748-9326/8/1/014003, 2013.

- Kondo, Y., Oshima, N., Kajino, M., Mikami, R., Moteki, N., Takegawa, N., Verma, R. L., Kajii, Y., Kato, S. and Takami, A.: Emissions of black carbon in East Asia estimated from observations at a remote site in the East China Sea, *Journal of Geophysical Research: Atmospheres*, 116(D16), n/a–n/a, doi:10.1029/2011JD015637, 2011.
- 5 Krasenbrink, A. and Dobranskyte Niskota, A.: 2007 Technical Review of the NRMM Directive 1997/68/EC as amended by Directives 2002/88/EC and 2004/26/EC, JRC, Ispra, Italy., 2008.
- Kumar, S., Aggarwal, S. G., Gupta, P. K. and Kawamura, K.: Investigation of the tracers for plastic-enriched waste-burning aerosols, *Atmospheric Environment*, 108, 49–58, doi:10.1016/j.atmosenv.2015.02.066, 2015.
- Kupiainen, K. and Klimont, Z.: Primary Emissions of Submicron and Carbonaceous Particles in Europe and the Potential for their Control, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria., 2004.
- 10 Kupiainen, K. and Klimont, Z.: Primary emissions of fine carbonaceous particles in Europe, *Atmospheric Environment*, 41, 2156–2170, doi:10.1016/j.atmosenv.2006.10.066, 2007.
- Kupiainen, K. J. and Pirjola, L.: Vehicle non-exhaust emissions from the tyre-road interface—effect of stud properties, traction sanding and resuspension, *Atmospheric Environment*, 45(25), 4141–4146, doi:10.1016/j.atmosenv.2011.05.027, 2011.
- 15 Kupiainen, K. J., Tervahattu, H., Räisänen, M., Mäkelä, T., Aurela, M. and Hillamo, R.: Size and Composition of Airborne Particles from Pavement Wear, Tires, and Traction Sanding, *Environ. Sci. Technol.*, 39(3), 699–706, doi:10.1021/es035419e, 2005.
- Lack, D., Lerner, B., Granier, C., Baynard, T., Lovejoy, E., Massoli, P., Ravishankara, A. R. and Williams, E.: Light absorbing carbon emissions from commercial shipping, *Geophysical Research Letters*, 35(13), L13815, doi:10.1029/2008GL033906, 2008.
- 20 Lack, D. A. and Corbett, J. J.: Black carbon from ships: a review of the effects of ship speed, fuel quality and exhaust gas scrubbing, *Atmos. Chem. Phys.*, 12, 3985–4000, doi:10.5194/acp-12-3985-2012, 2012.
- Lack, D. A., Corbett, J. J., Onasch, T., Lerner, B., Massoli, P., Quinn, P. K., Bates, T. S., Covert, D. S., Coffman, D., Sierau, B. and others: Particulate emissions from commercial shipping: Chemical, physical, and optical properties, *J. Geophys. Res.*, 114, D00F04, 2009.
- 25 Lam, N. L., Chen, Y., Weyant, C., Venkataraman, C., Sadavarte, P., Johnson, M. A., Smith, K. R., Brem, B. T., Arinaitwe, J., Ellis, J. E. and Bond, T. C.: Household Light Makes Global Heat: High Black Carbon Emissions From Kerosene Wick Lamps, *Environmental Science & Technology*, 46(24), 13531–13538, doi:10.1021/es302697h, 2012.
- Lam, N. L., Pachauri, S., Cameron, C., Purohit, P. and Nagai, Y.: Characterizing Kerosene Demand for Light in India and Evaluating the Impact of Measures Affecting Access and Dependence, in *Discovering Untapped Resources*, p. 116–119., UC Berkeley, [online] Available from: https://opus4.kobv.de/opus4_tuberlin/frontdoor/index/index/docId/5185 (Accessed 20 March 2015), 2014.
- 30 Lam, N. L., Pachauri, S., Purohit, P., Nagai, Y., Bates, M. N., Cameron, C. and Smith, K. R.: Kerosene subsidies for household lighting in India: what are the impacts?, *Environmental Research Letters*, 11(4), 044014 (1–11), doi:10.1088/1748-9326/11/4/044014, 2016.
- 35 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Lioussé, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D. T., Smith, S. J., Stehfest, E., Van Aardenne, J. A., Cooper, O. R., Kainuma, M., Mahowald, N.,

- McConnell, J. R., Naik, V., Riahi, K. and Van Vuuren, D. P. V.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application., *Atmos. Chem. Phys.*, 10(2), 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Lange, D. and Domke, F.: The exhaust emissions scandal („Dieselgate“) – Take a deep breath into pollution trickery, [online] Available from: https://media.ecc.de/v/32e3-7331-the_exhaust_emissions_scandal_dieselgate (Accessed 5 January 2016), 2015.
- Lawson, D. R.: (On-Road) Mobile Source Emissions and Mitigation Potential, 2010.
- Le, H. A. and Oanh, N. T. K.: Integrated assessment of brick kiln emission impacts on air quality Hoang Anh Le – Nguyen Thi Kim Oanh, *Environ Monit Assess.*, 171, 381–394, doi:10.1007/s10661-009-1285-y, 2010.
- Lee, D. S., Fahey, D. W., Forster, P. M., Newton, P. J., Wit, R. C. N., Lim, L. L., Owen, B. and Sausen, R.: Aviation and global climate change in the 21st century, *Atmos. Environ.*, 43, 3520–3537, 2009.
- Lee, W. J., Liu, Y. C., Mwangi, F. K., Chen, W. H., Lin, S. L., Fukushima, Y., Liao, C. N. and Wang, L. C.: Assessment of energy performance and air pollutant emissions in a diesel engine generator fueled with water containing ethanol biodiesel–diesel blend of fuels, *Energy*, 36(9), 5591–5599, doi:10.1016/j.energy.2011.07.012, 2011.
- Lei, Y., Zhang, Q., He, K. B. and Streets, D. G.: Primary anthropogenic aerosol emission trends for China, 1990–2005, *Atmospheric Chemistry and Physics*, 11(3), 931–954, doi:10.5194/acp-11-931-2011, 2011.
- Lewis, J. J. and Pattanayak, S. K.: Who Adopts Improved Fuels and Cookstoves? A Systematic Review, *Environmental Health Perspectives*, 120, 637–645, 2012.
- Li, Q., Li, X., Jiang, J., Duan, L., Ge, S., Zhang, Q., Deng, J., Wang, S. and Hao, J.: Semi-coke briquettes: towards reducing emissions of primary PM_{2.5}, particulate carbon, and carbon monoxide from household coal combustion in China, *Scientific Reports*, 6, 19306, 2016.
- Li, X. H., Wang, S. X., Duan, L., Hao, J. M. and Nie, Y. F.: Carbonaceous aerosol emissions from household biofuel combustion in China., *Environ. Sci. Technol.*, 43, 6076–6081, 2009.
- Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair Rohani, H., AlMazroa, M. A., Amann, M., Anderson, H. R., Andrews, K. G., Aryee, M., Atkinson, C., Bacchus, L. J., Bahalim, A. N., Balakrishnan, K., Balmes, J., Barker Collo, S., Baxter, A., Bell, M. L., Blore, J. D., Blyth, F., Bonner, C., Borges, G., Bourne, R., Boussinesq, M., Brauer, M., Brooks, P., Bruce, N. G., Brunekreef, B., Bryan Hancock, C., Bucello, C., Buehler, R., Bull, F., Burnett, R. T., Byers, T. E., Calabria, B., Carapetis, J., Carnahan, E., Chafe, Z., Charlson, F., Chen, H., Chen, J. S., Cheng, A. T. A., Child, J. C., Cohen, A., Colson, K. E., Cowie, B. C., Darby, S., Darling, S., Davis, A., Degenhardt, L., Dentener, F., Des Jarlais, D. C., Devries, K., Dherani, M., Ding, E. L., Dorsey, E. R., Driscoll, T., Edmond, K., Ali, S. E., Engell, R. E., Erwin, P. J., Fahimi, S., Falder, G., Farzadfar, F., Ferrari, A., Finucane, M. M., Flaxman, S., Fowkes, F. G. R., Freedman, G., Freeman, M. K., Gakidou, E., Ghosh, S., Giovannucci, E., Gmel, G., Graham, K., Grainger, R., Grant, B., Gunnell, D., Gutierrez, H. R., Hall, W., Hoek, H. W., Hogan, A., Hosgood, H. D., Hoy, D., Hu, H., Hubbell, B. J., Hutchings, S. J., Ibeanusi, S. E., Jacklyn, G. L., Jasrasaria, R., Jonas, J. B., Kan, H., Kanis, J. A., Kassebaum, N., Kawakami, N., Khang, Y. H., Khatibzadeh, S., Khoo, J. P., et al.: A comparative risk assessment of burden and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010, *The Lancet*, 380(9859), 2224–2260, doi:10.1016/S0140-6736(12)61766-8, 2012.

- Lin, J., Nielsen, C. P., Zhao, Y., Lei, Y., Liu, Y. and McElroy, M. B.: Recent changes in particulate air pollution over China observed from space and the ground: Effectiveness of emission control, *Environmental Science and Technology-Columbus*, 44(20), 7771, 2010.
- 5 Lin, Y.-C., Lee, W. J., Chao, H. R., Wang, S. L., Tsou, T. C., Chang-Chien, G. P. and Tsai, P. J.: Approach for energy saving and pollution reducing by fueling diesel engines with emulsified biosolution/biodiesel/diesel blends, *Environmental Science and Technology*, 42(10), 3849–3855, 2008.
- Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H. and He, K. B.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, *Atmospheric Chemistry and Physics*, 15(23), 13299–13317, doi:10.5194/acp-15-13299-2015, 2015.
- 10 Liu, H., He, K., Lents, J. M., Wang, Q. and Tolvett, S.: Characteristics of Diesel Truck Emission in China Based on Portable Emissions Measurement Systems, *Environmental Science & Technology*, 43(24), 9507–9511, doi:10.1021/es902044x, 2009.
- Louhelainen, K., Vilhunen, P., Kangas, J. and Terho, F. O.: Dust exposure in piggeries, *European J. of Resp. Diseases*, 71(152), 80–90, 1987.
- 15 Lu, Z., Zhang, Q. and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, *Atmos. Chem. Phys.*, 11(18), 9839–9864, 2011.
- Lükewille, A., Bertok, I., Amann, M., Cofala, J., Gyarmas, F., Johansson, M., Klimont, Z., Pacyna, E. and Pacyna, J.: A Module to Calculate Primary Particulate Matter Emissions and Abatement Measures in Europe, *Water, Air, and Soil Pollution*, 130(1–4), 229–234, doi:10.1023/A:1013807101561, 2001.
- 20 Lund, M. T., Berntsen, T. K., Heyes, C., Klimont, Z. and Samset, B. H.: Global and regional climate impacts of black carbon and co-emitted species from the on-road diesel sector, *Atmos. Environ.*, 98, 50–58, doi:10.1016/j.atmosenv.2014.08.033, 2014.
- MacCarty, N., Ogle, D., Still, D., Bond, T., Roden, C. and Willson, B.: Laboratory Comparison of the Global Warming Potential of Six Categories of Biomass Cooking Stoves, Aprovecho Research Center. [online] Available from: internal-pdf://Global_warming_full_9-6-07-2181006597/Global_warming_full_9-6-07.pdf, 2007.
- 25 Mahapatra, S., Chanakya, H. N. and Dasappa, S.: Evaluation of various energy devices for domestic lighting in India: Technology, economics and CO₂ emissions, *Energy for Sustainable Development*, 13(4), 271–279, doi:10.1016/j.esd.2009.10.005, 2009.
- Maithel, S.: Factsheets about Brick Kilns in South and South-East Asia, [online] Available from: <http://www.ceacoalition.org/en/resources/factsheets-about-brick-kilns-south-and-south-east-asia>, 2014.
- 30 Maithel, S., Lalchandani, D., Malhotra, G., Bhanware, P., Uma, R., Ragavan, S., Athalye, V., Bindiya, K., Reddy, S., Bond, T. C., Weyant, C., Baum, E., Kim Thoa, V. T., Thu Phuong, N. and Kim Thanh, T.: Brick Kilns Performance Assessment: A Roadmap for Cleaner Brick Production in India, Shakti Sustainable Energy Foundation and Climate Works Foundation. [online] Available from: <http://www.catf.us/resources/publications/view/161>, 2012.
- 35 Maíz, P.: Experiencias en Medición de Emisiones en Hornos para Manufactura Artesanal de Ladrillos en Mexico, [online] Available from: http://www.ine.gob.mx/ceinea/memorias/1111_taller_ladrilleras_2012_eng (Accessed 17 September 2012), 2012.

Mancilla, Y., Araizaga, A. E. and Mendoza, A.: A tunnel study to estimate emission factors from mobile sources in Monterrey, Mexico, *Journal of the Air & Waste Management Association*, 62(12), 1431-1442, doi:10.1080/10962247.2012.717902, 2012.

5 Marieq, M. M.: Chemical characterization of particulate emissions from diesel engines: A review M. Matti Marieq*, *Aerosol Science*, 38, 1079-1118, doi:10.1016/j.jaerosci.2007.08.001, 2007.

Mazzoleni, C., Moosmüller, H., Kuhns, H. D., Keislar, R. E., Barber, P. W., Nikolic, D., Nussbaum, N. J. and Watson, J. G.: Correlation between automotive CO, HC, NO, and PM emission factors from on-road remote sensing: implications for inspection and maintenance programs, *Transportation Research Part D: Transport and Environment*, 9(6), 477-496, 2004.

10 McClintock, P.: Remote Sensing Measurements of Real-World High-Exhaust Emitters, Coordinating Research Council, Inc., 1999.

McClintock, P.: 2007 High-Emitter Remote Sensing Project, Prepared for Southeast Michigan Council of Governments, Environmental Systems Products Inc., Tiburon, CA/USA., 2007.

15 McEwen, J. D. N. and Johnson, M. R.: Black Carbon Particulate Matter Emission Factors for Buoyancy Driven Associated Gas Flares, *Journal of the Air & Waste Management Association*, 63(3), 307-321, doi:10.1080/10473289.2011.650040, 2012.

Mills, E.: Technical and Economic Performance Analysis of Kerosene Lamps and Alternative Approaches to Illumination in Developing Countries, Lawrence Berkeley National Laboratory (LBNL), Berkeley. [online] Available from: <http://evanmills.lbl.gov/pubs/pdf/offgrid-lighting.pdf>, 2003.

Mills, E.: The specter of fuel-based lighting, *Science*, 308, 1263-1264, 2005.

20 Ministério do Meio Ambiente: Inventário Nacional de Emissões Atmosféricas por Veículos Automotores Rodoviários, Final Report, Ministério do Meio Ambiente, Secretaria de Mudanças Climáticas e Qualidade Ambiental (MMA), Brasília/BR., 2011.

Moldanova, J., Fridell, E., Popovichieva, O., Demirdjian, B., Tishkova, V., Faccinotto, A. and Foesa, C.: Characterisation of Particulate Matter and Gaseous Emissions from a Large Ship Diesel Engine., *Atmos. Environ.*, 43, 2632-2641, 2009.

25 Morton, D. C., DeFries, R. S., Randerson, J. T., Giglio, L., Schroeder, W. and Van Der Werf, G. R.: Agricultural intensification increases deforestation fire activity in Amazonia, *Global Change Biology*, 14(10), 2262-2275, doi:10.1111/j.1365-2486.2008.01652.x, 2008.

30 Mukai, S., Nakata, M., Yasumoto, M., Sano, I. and Kokhanovsky, A.: A study of aerosol pollution episode due to agriculture biomass burning in the east central China using satellite data, *Frontiers in Environmental Science*, 3(Article 57), doi:10.3389/fenvs.2015.00057, 2015.

Murphy, S. M., Agrawal, H., Sorooshian, A., Padro, L. T., Gates, H., Hersey, S., Welch, W. A., Jung, H., Miller, J. W., Cocker III, D. R., Nenes, A., Jonsson, H. H., Flagan, R. C. and Seinfeld, J. H.: Comprehensive Simultaneous Shipboard and Airborne Characterization of Exhaust from a Modern Container Ship at Sea., *Environ. Sci. & Technol.*, 43(13), 4626-4640, 2009.

35 NEA: A Year in Review: Fiscal Year 2011/2012, Nepal Electricity Authority (NEA), Kathmandu., 2012.

Neurath, C.: Open Burning of Domestic Wastes: The Single Largest Source of Dioxin to Air?, *Organohalogen Compounds*, 60–65, 2003.

NIELSEN: All India Study on Sectoral Demand of Diesel & Petrol, Petroleum Planning and Analysis Cell (PPAC), Ministry of Petroleum & Natural Gas, Government of India, New Delhi. [online] Available from: <http://ppac.org.in>, 2013.

5 Niemi, J. K.: Atmospheric Emissions from Open Biomass Burning; Development of Datasets for RAINS Model, Interim Report, International Institute for Applied Systems Analysis, Laxenburg, Austria., 2007.

Norris, J. O. W.: An In-Service Emissions Test for Spark Ignition (SI) Petrol Engines — PPAD 9/107/09. Phase 1 Report. Definition of an Excess Emitter and Effectiveness of Current Annual Test, AEAT Technology, Harwell, UK. [online] Available from: https://uk-air.defra.gov.uk/assets/documents/reports/eat15/0408171318_S1Phase1reportIssue3.pdf, 2001.

10 NSSO: Energy sources of India household for cooking and lighting, NSS 61st Round (July 2004–June 2005), National Sample Survey Organisation (NSSO), Ministry of Statistics and Programme Implementation, Government of India., New Delhi, India., 2007.

15 Ntziachristos, L., Gkatzoflias, D., Kouridis, C. and Samaras, Z.: COPERT: A European Road Transport Emission Inventory Model, in *Information Technologies in Environmental Engineering—Proceedings of the 4th International ICSC Symposium Thessaloniki, Greece, May 28–29, 2009*, pp. 491–504, Springer Berlin Heidelberg., 2009.

Oanh, N. T. K., Bieh, T. L., Tipayarom, D., Manadhar, B. R., Prapat, P., Simpson, C. D. and Liu, L. J. S.: Characterization of particulate matter emission from open burning of rice straw, *Atmospheric environment (Oxford, England : 1994)*, 45(2), 493–502, doi:10.1016/j.atmosenv.2010.09.023, 2011.

20 Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X. and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, *Atmos. Chem. Phys.*, 7, 4419–4444, 2007.

Ohlström, M. O., Lehtinen, K. E. J., Moisio, M. and Jokiniemi, J. K.: Fine particle emissions of energy production in Finland, *Atmospheric Environment*, 34(22), 3701–3711, doi:10.1016/S1352-2310(00)00076-5, 2000.

25 O'Neill, B. C., Krieglner, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R. and van Vuuren, D. P.: A new scenario framework for climate change research: the concept of shared socioeconomic pathways, *Climatic Change*, 122(3), 387–400, doi:10.1007/s10584-013-0905-2, 2014.

OTAQ: Exhaust and Crankcase Emission Factors for Nonroad Engine Modeling—Compression Ignition, US Environmental Protection Agency., 2004.

30 Paasonen, P., Visschedijk, A., Kupiainen, K., Klimont, Z., Denier van der Gon, H. A. C., Kulmala, M. and Amann, M.: Aerosol particle number emissions and size distributions: Implementation in the GAINS model and initial results, IIASA Interim Report, International Institute for Applied Systems Analysis, Laxenburg, Austria., 2013.

Paasonen, P., Kupiainen, K., Klimont, Z., Visschedijk, A., Denier van der Gon, H. A. C. and Amann, M.: Continental anthropogenic primary particle number emissions, *Atmospheric Chemistry and Physics Discussions*, 2016, 1–39, doi:10.5194/acp-2015-1023, 2016.

35 Parashar, D. C., Gadi, R., Mandal, T. K. and Mitra, A. P.: Carbonaceous aerosol emissions from India, *Atmospheric Environment*, 39, 7861–7871, 2005.

- Parikh, K. S.: Report of the expert group on a viable and sustainable system of pricing petroleum products, Government of India, New Delhi, India., 2010.
- Pederstad, A., Smith, J. D., Jackson, R., Saunier, S. and Holm, T.: Assessment of flare strategies, techniques for reduction of flaring and associated emissions, emission factors and methods for determination of emissions to air from flaring, Carbon Limits AS, Trondheim, Norway. [online] Available from: www.miljødirektoratet.no, 2015.
- Pettersson, E., Boman, C., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 2: Wood Stove, *Energy Fuels*, 25(1), 315–323, doi:10.1021/ef1007787, 2011.
- Pettus, A.: *Agricultural Fires and Arctic Climate Change*, CATF (Clean Air Task Force), Boston, US. [online] Available from: <http://www.catf.us>, 2009.
- Petzold, A., Hasselbach, J., Lauer, P., Baumann, R., Franke, K., Gurk, C., Schlager, H. and Weingartner, E.: Experimental studies on particle emissions from cruising ship, their characteristic properties, transformation and atmospheric lifetime in the marine boundary layer, *Atmos. Chem. Phys.*, 8, 2387–2403, doi:10.5194/acp-8-2387-2008, 2008.
- Petzold, A., Weingartner, E., Hasselbach, J., Lauer, P., Kurok, C. and Fleischer, F.: Physical Properties, Chemical Composition, and Cloud-Forming Potential of Particulate Emissions from a Marine Diesel Engine at Various Load Conditions, *Environ. Sci. & Technol.*, 44(10), 3800–3805, 2010.
- PFC Energy: *Using Russia's Associated Gas*, PFC Energy, Washington DC, USA. [online] Available from: www.pfcenergy.com, 2007.
- Pine, K., Edwards, R., Masera, O., Schilman, A., Marrón-Mares, A. and Riojas-Rodríguez, H.: Adoption and use of improved biomass stoves in Rural Mexico, *Energy for Sustainable Development*, 15(2), 176–183, doi:10.1016/j.esd.2011.04.001, 2011.
- Pode, R.: Solution to enhance the acceptability of solar powered LED lighting technology, *Renewable and Sustainable Energy Reviews*, 14(3), 1096–1103, doi:10.1016/j.rser.2009.10.006, 2010.
- Pokhrel, A. K., Bates, M. N., Verma, S. C., Joshi, H. S., Sreeramareddy, C. T. and Smith, K. R.: Tuberculosis and indoor biomass and kerosene use in Nepal: a case control study, *Environmental Health Perspectives*, 118(4), 558–564, doi:10.1289/ehp.0901032, 2010.
- Polenske, K. R. [ed]: *The Technology Energy Environment Health (TEEH) Chain in China: A Case Study of Cokemaking*, Springer Netherlands., 2006.
- Purohit, P. and Michaelowa, A.: CDM potential of SPV lighting systems in India, *Mitigation and Adaptation Strategies for Global Change*, 13(1), 23–46, doi:10.1007/s11027-006-9078-x, 2008.
- Purohit, P., Amann, M. and Mathur, R.: *GAINS Asia: Scenarios for Cost Effective Control of Air Pollution and Greenhouse Gases in India*. [online] Available from: <http://gains.iiasa.ac.at/gains/reports/GAINS-Asia-Methodology-20081205.pdf> (Accessed 9 August 2012), 2010.
- Quennehen, B., Raut, J.-C., Law, K. S., Daskalakis, N., Anechet, G., Clerbaux, C., Kim, S.-W., Lund, M. T., Myhre, G., Olivie, D. J. L., Safieddine, S., Skeie, R. B., Thomas, J. L., Tsyro, S., Bazureau, A., Bellouin, N., Hu, M., Kanakidou, M., Klimont, Z., Kupiainen, K., Myriokefalitakis, S., Quaa, J., Rumbold, S. T., Schulz, M., Cherian, R., Shimizu, A., Wang, J.,

Yoon, S.-C. and Zhu, T.: Multi-model evaluation of short lived pollutant distributions over east Asia during summer 2008, *Atmospheric Chemistry and Physics*, 16(17), 10765–10792, doi:10.5194/acp-16-10765-2016, 2016.

Rajaratnam, U., Athalye, V., Ragavan, S., Maithel, S., Lalchandani, D., Kumar, S., Baum, E., Weyant, C. and Bond, T.: Assessment of air pollutant emissions from brick kilns, *Atmospheric Environment*, 98, 549–553, doi:10.1016/j.atmosenv.2014.08.075, 2014.

Randerson, J. T., Chen, Y., van der Werf, G. R., Rogers, B. M. and Morton, D. C.: Global burned area and biomass-burning emissions from small fires, *Journal of Geophysical Research: Biogeosciences*, 117(G4), doi:10.1029/2012JG002128, 2012.

Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J. and Kasibhatla, P. S.: Global Fire Emissions Database, Version 4, (GFEDv4), [online] Available from: <http://dx.doi.org/10.3334/ORNLDAAC/1293>, 2015.

Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., van Vuuren, D. P., Aleluia Reis, L., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Streffler, J., van der Sluis, S. and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Global Environmental Change*, doi:10.1016/j.gloenvcha.2016.05.012, in press.

Rao, S., Pachauri, S., Dentener, F., Kinney, P., Klimont, Z., Riahi, K. and Schoepp, W.: Better air for better health: Forging synergies in policies for energy access, climate change and air pollution, *Global Environmental Change*, doi:10.1016/j.gloenvcha.2013.05.003, 2013.

Reddington, C. L., Butt, E. W., Ridley, D. A., Artaxo, P., Morgan, W. T., Coe, H. and Spracklen, D. V.: Air quality and human health improvements from reductions in deforestation-related fire in Brazil, *Nature Geosci*, 8(10), 768–771, 2015.

Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Streffler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A. and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, *Global Environmental Change*, doi:10.1016/j.gloenvcha.2016.05.009, in press.

Riahi, K., Dentener, F., Gielen, D., Grubler, A., Jewell, J., Klimont, Z., Krey, V., McCollum, D., Pachauri, S., Rao, S., van Ruijven, B., van Vuuren, D. P. and Wilson, C.: Chapter 17 Energy Pathways for Sustainable Development, in *Global Energy Assessment – Toward a Sustainable Future*, pp. 1203–1306, Cambridge University Press, Cambridge, UK and New York, NY, USA and the International Institute for Applied Systems Analysis, Laxenburg, Austria. [online] Available from: www.globalenergyassessment.org, 2012.

Roden, C. A., Bond, T. C., Conway, S. and Pinel, A. B. O.: Emission Factors and Real-Time Optical Properties of Particles Emitted from Traditional Wood Burning Cookstoves, *Environ. Sci. Technol.*, 40(21), 6750–6757, doi:10.1021/es052080i, 2006.

Roden, C. A., Bond, T. C., Conway, S., Osorto Pinel, A. B., MacCarty, N. and Still, D.: Laboratory and field investigations of particulate and carbon monoxide emissions from traditional and improved cookstoves, *Atmospheric Environment*, 43(6), 1170–1181, doi:10.1016/j.atmosenv.2008.05.041, 2009.

Røland, T. H.: Associated Petroleum Gas in Russia: Reasons for non utilization, Fridtjof Nansens Institute, Lysaker, Norway. [online] Available from: www.fni.no, 2010.

- Ruiz Mercado, I., Masera, O., Zamora, H. and Smith, K. R.: Adoption and sustained use of improved cookstoves, *Energy Policy*, 39(12), 7557–7566, doi:10.1016/j.enpol.2011.03.028, 2011.
- Schaap, M., Van Der Gon, H. A. C. D., Dentener, F. J., Visschedijk, A. J. H., Van Loon, M., ten Brink, H. M., Putaud, J. P., Guillaume, B., Liousse, C. and Buitjes, P. J. H.: Anthropogenic black carbon and fine aerosol distribution over Europe, *Journal of Geophysical Research: Atmospheres*, 109(D18), n/a–n/a, doi:10.1029/2003JD004330, 2004.
- Schäffeler, U. and Keller, M.: Non-road fuel consumption and pollutant emissions. Study for the period from 1980 to 2020, Federal Office for the Environment, Bern/CH., 2008.
- Schilderman, T. and Mason, K.: Using residues as fuel in small scale brickmaking, in *Proceedings of the 11th International Conference on Non-conventional Materials and Technologies (NOCMAT 2009)*, Bath, UK., 2009.
- Schmidl, C., Luisser, M., Padouvas, E., Lasselsberger, L., Rzaca, M., Cruz, C. S., Handler, M., Peng, G., Bauer, H. and Puxbaum, H.: Particulate and gaseous emissions from manually and automatically fired small scale combustion systems, *Atmospheric Environment*, 45, 7443–7454, 2011.
- Schmidt, C. W.: Modernizing Artisanal Brick Kilns: A Global Need, *Environmental Health Perspectives*, 121, doi:http://dx.doi.org/10.1289/ehp.121.A242, 2013.
- Schöpp, W., Klimont, Z., Suutari, R. and Cofala, J.: Uncertainty analysis of emission estimates in the RAINS integrated assessment model, *Environmental Science & Policy*, 8, 601–613, 2005.
- Schwarz, J. P., Holloway, J. S., Katich, J. M., McKeen, S., Kort, E. A., Smith, M. L., Ryerson, T. B., Sweeney, C. and Peischl, J.: Black Carbon Emissions from the Bakken Oil and Gas Development Region, *Environ. Sci. Technol. Lett.*, 2(10), 281–285, doi:10.1021/acs.estlett.5b00225, 2015.
- Scott, A. J.: Real-life emissions from residential wood burning appliances in New Zealand, Ministry for the Environment, Christchurch, New Zealand., 2005.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, New York, USA. [online] Available from: http://books.google.com/books?hl=en&lr=&id=YH2K9eWsZOeC&oi=fnd&pg=PA1991&dq=%22activities,+including+oxidation+of+anthropogenically+derived+CH4.+The%22+%22of+industrial%22+%22the+maximum+values+are+found+near+the+surface+at+northern+midlatitudes.%22+%22of+its+overwhelming+importance+in+global+climate,+we+will+delay+eonsideration%22+&ots=hJ9tPmbVEB&sig=_J5iCU26zodWW2ugZH1T2PdhFhw (Accessed 1 August 2013), 2012.
- Shafizadeh, K., Niemeier, D. and Eisinger, D. S.: *Gross Emitting Vehicles: A Review of the Literature*, Caltrans Air Quality Project, US Davis., 2004.
- Shah, S. D., Cocker, D. R., Miller, J. W. and Norbeck, J. M.: Emission rates of particulate matter and elemental and organic carbon from in-use diesel engines, *Environmental Science & Technology*, 38(9), 2544–2550, doi:10.1021/es0350583, 2004.
- Shah, S. D., Cocker, D. R., Johnson, K. C., Lee, J. M., Soriano, B. L. and Miller, J. W.: Emissions of regulated pollutants from in-use diesel backup generators, *Atmospheric Environment*, 40(22), 4199–4209, 2006.
- Shah, S. D., Cocker III, D. R., Johnson, K. C., Lee, J. M., Soriano, B. L. and Miller, J. W.: Reduction of particulate matter emissions from diesel backup generators equipped with four different exhaust after-treatment devices, *Environmental Science & Technology*, 41(14), 5070–5076, doi:10.1021/es0614161, 2007.

Shi, X., Pang, X., Mu, Y., He, H., Shuai, S., Wang, J., Chen, H. and Li, R.: Emission reduction potential of using ethanol-biodiesel diesel fuel blend on a heavy-duty diesel engine, *Atmospheric Environment*, 40(14), 2567–2574, doi:10.1016/j.atmosenv.2005.12.026, 2006.

5 Shindell, D., Kuylenstierna, J. C. I., Vignati, E., Van Dingenen, R., Amann, M., Klimont, Z., Anenberg, S. C., Müller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L., Kupiainen, K., Höglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh, N. T. K., Milly, G., Williams, M., Demkine, V. and Fowler, D.: Simultaneously mitigating near term climate change and improving human health and food security, *Science*, 335(6065), 183–189, 2012.

10 Shrimali, G., Slaski, X., Thurber, M. C. and Zerriffi, H.: Improved stoves in India: A study of sustainable business models, *Energy Policy*, 39(12), 7543–7556, doi:10.1016/j.enpol.2011.07.031, 2011.

15 Silk, B. J., Sudamah, I., Patel, M. K., Were, V., Person, R., Harris, J., Otieno, R., Nygren, B., Loo, J., Eleveld, A., Quick, R. E. and Cohen, A. L.: A strategy to increase adoption of locally produced, ceramic cookstoves in rural Kenyan households Benjamin J Silk1,2*, Ibrahim Sadumah3, Minal K Patel1,4, Vincent Were3, Bobbie Person4, Julie Harris1,4, Ronald Otieno3, Benjamin Nygren4, Jennifer Loo2, Alie Eleveld5, Robert E Quick4 and Adam L Cohen2, *BMC Public Health*, 12, 359–369, 2012.

Sippula, O., Hokkinen, J., Puustinen, H., Yli-Pirila, P. and Jokiniemi, J.: Comparison of particulate emissions from small heavy fuel-oil and wood-fired boilers, *Atmospheric Environment*, 43, 4855–4864, 2009.

20 Skinder, B. M., Sheikh, A. Q., Pandit, A. K. and Ganai, B. A.: Brick kiln emissions and its environmental impact: A Review Bhat Mohd Skinder*, Afeefa Qayoom Sheikh, Ashok K. Pandit and Bashir Ahmad Ganai, *Journal of Ecology and the Natural Environment*, 6(1), 1–11, 2014.

Smit, R. and Bluett, J.: A new method to compare vehicle emissions measured by remote sensing and laboratory testing: High emitters and potential implications for emission inventories, *Science of The Total Environment*, 409(13), 2626–2634, doi:10.1016/j.scitotenv.2011.03.026, 2011.

25 Stein, G., Wuenstel, E. and Travnicek Pagaimo, W.: Reifenabrieb in Feinstaub – kein Grund zur Panik!, *GAK – Gummi Fasern Kunststoffe*, Jahrgang 65, 441–445, 2012.

Stettler, M. E. J., Boies, A. M., Petzold, A. and Barrett, S. R. H.: Global Civil Aviation Black Carbon Emissions, *Environmental Science & Technology*, 47(18), 10397–10404, doi:10.1021/es401356v, 2013.

30 Stoerk, T.: Statistical corruption in Beijing's air quality data has likely ended in 2012, *Atmospheric Environment*, 127, 365–371, doi:10.1016/j.atmosenv.2015.12.055, 2016.

Stohl, A., Berg, T., Burkhardt, J. F., Fjårraa, A. M., Forster, C., Herber, A., Hov, Ø., Lunder, C., McMillan, W. W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Ström, J., Tørseth, K., Treffeisen, R., Virkkunen, K. and Yttri, K. E.: Arctic smoke – record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006, *Atmospheric Chemistry and Physics*, 7(2), 511–534, doi:10.5194/acp-7-511-2007, 2007.

35 Stohl, A., Klimont, Z., Eckhardt, S., Kupiainen, K., Shevchenko, V. P., Kopeikin, V. M. and Novigatsky, A. N.: Black carbon in the Arctic: the underestimated role of gas flaring and residential combustion emissions, *Atmospheric Chemistry and Physics*, 13(17), 8833–8855, doi:10.5194/acp-13-8833-2013, 2013.

Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Berntsen, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinska, M., Eckhardt, S., Fuglestad, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U.,

Formatted: German (Germany)

Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Olivici, D., Quaas, J., Quennehen, B., Raut, J. C., Rumbold, S. T., Samsat, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E. and Zhu, T.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmos. Chem. Phys.*, 15(18), 10529–10566, doi:10.5194/acp-15-10529-2015, 2015.

5 Stratus Consulting: CCAC initiative to mitigate black carbon and other pollutants from brick production: Regional Assessment. Black Carbon Mitigation in Brick Production: A Summary for Brazil, Chile, Colombia, Mexico, Nigeria, and Peru, Climate and Clean Air Coalition (CCAC), 2014.

10 Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J. H. and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research*, 108(D21), 1–23, doi:10.1029/2002JD003093, 2003.

Subramanian, R., Winijkul, E., Bond, T. C., Thiansathit, W., Oanh, N. T. K., Pawarmart, I. and Duleep, K. G.: Climate-Relevant Properties of Diesel Particulate Emissions: Results from a Piggyback Study in Bangkok, Thailand, *Environmental Science & Technology*, 43(11), 4213–4218, doi:10.1021/es8032296, 2009.

15 Takai, H., Pedersen, S., Johnsen, J. O., Metz, J. H. M., Groot Koerkamp, P. W. G., Uenk, G. H., Phillips, V. R., Holden, M. R., Sneath, R. W., Short, J. L., White, R. P., Hartung, J., Seedorf, J., Schröder, M., Linkert, K. H. and Wathes, C. M.: Concentrations and emission of airborne dust in livestock buildings in Northern Europe, *J. Agric. Engng. Res.*, 1(70), 59–77, 1998.

20 Tang, N. W., Apte, J. S., Martien, P. T. and Kirchstetter, T. W.: Measurement of black carbon emissions from in-use diesel-electric passenger locomotives in California, *Atmospheric Environment*, 115, 295–303, doi:10.1016/j.atmosenv.2015.05.001, 2015.

Tissari, J., Hytönen, K., Lyyrinen, J. and Jokiniemi, J.: A novel field measurement method for determining fine particle and gas emissions from residential wood combustion, *Atmospheric Environment*, 41, 8330–8344, 2007.

25 Tissari, J., Lyyrinen, J., Hytonen, K., Sippula, O., Tapper, U., Frey, A., Saarnio, K., Pennanen, A. S., Hillamo, R., Salonen, R. O., Hirvonen, M. R. and Jokiniemi, J.: Fine particle and gaseous emissions from normal and smouldering wood combustion in a conventional masonry heater, *Atmos. Environ.*, 42, 7862–7873, 2008.

Tissari, J., Hytonen, K., Sippula, O. and Jokiniemi, J.: The effects of operating conditions on emissions from masonry heaters and sauna stoves, *Biomass and Bioenergy*, 33, 513–520, 2009.

Todd, J. J.: *Wood Smoke Handbook: Woodheaters, Firewood and Operator Practice*, Eco Energy Options, Lindisfarne Tasmania, 2003.

30 Triple E.: Diesel power generation: Inventories, black carbon emissions, initial stakeholder meeting and data mapping for Nigeria, World Bank, Washington D.C., 2013.

Troncoso, K., Castillo, A., Merino, L., Lazos, E. and Masera, O. R.: Understanding an improved cookstove program in rural Mexico: An analysis from the implementers' perspective, *Energy Policy*, 39(12), 7600–7608, doi:10.1016/j.enpol.2011.04.070, 2011.

35 Tsai, J. H., Chen, S. J., Huang, K. L., Lin, Y. C., Lee, W. J., Lin, C. C. and Lin, W. Y.: PM, carbon, and PAH emissions from a diesel generator fuelled with soy biodiesel blends, *Journal of Hazardous Materials*, 179(1–3), 237–243, doi:10.1016/j.jhazmat.2010.02.085, 2010.

Turn, S. Q., Jenkins, B. M., Chow, J. C., Pritchett, L. C., Campbell, D., Cahill, T. and Whalem, S. A.: Elemental Characterization of Particulate Matter Emitted from Biomass Burning: Windtunnel-Derived Source Profiles for Herbaceous and Wood Fuels, *Journal of Geophysical Research*, 102(D3), 3683–3699, 1997.

5 Turpin, J. T. and Lim, H.: Species contributions to PM_{2.5} mass concentrations: revisiting common assumptions for estimating organic mass, *Aerosol Science and Technology*, 35, 602–610, 2001.

UBOS: Uganda national household survey 2009/2010, Uganda Bureau of Statistics (UBOS), Kampala, Uganda., 2010.

Uma, R., Kandpal, T. C. and Kishore, V. V. N.: Emission characteristics of an electricity generation system in diesel-alone and dual-fuel modes, *Biomass and Bioenergy*, 27(2), 195–203, doi:10.1016/j.biombioe.2004.01.003, 2004.

10 UNEP: Near-term Climate Protection and Clean Air Benefits: Actions for Controlling Short-Lived Climate Forcers, United Nations Environment Programme (UNEP), Nairobi, Kenya. [online] Available from: http://www.unep.org/dewa/Portals/67/pdf/Near_Term_Climate_Protection_&_Air_Benefits.pdf (Accessed 12 March 2012), 2011.

UNEP/AWMO: Integrated Assessment of Black Carbon and Tropospheric Ozone, Nairobi, Kenya. [online] Available from: www.unep.org/dewa/Portals/67/pdf/BlackCarbon_report.pdf, 2011.

15 UN-Habitat: Global Reports on Human Settlements 2005, United Nations Human Settlements Programme (UN-Habitat), Nairobi, Kenya., 2005.

US EPA: Compilation of Air Pollutant Emission Factors—Volume I: Stationary Point and Area Sources. AP-42 5th Edition., U.S. Environmental Protection Agency (US EPA), Springfield, VA., 1995.

20 US EPA: SPECIATE3.2 US EPA's repository of Total Organic Compound (TOC) and Particulate Matter (PM) speciated profiles for a variety of sources for use in source apportionment studies, [online] Available from: <http://www.epa.gov/ttn/chief/software/speciate/>, 2002.

US EPA: Notice of Violation, [online] Available from: <http://www3.epa.gov/otaq/cert/documents/vw-nov-2015-11-02.pdf>, 2015a.

US EPA: Notice of Violation, 2015b.

25 US EPA, (first): Emission Factors for Uncontrolled Industrial Diesel Engines, Section 3.3 Small Engines, United States Environmental Protection Agency (US EPA), Washington D.C., 1996.

US EPA-OTAQ: Development of Emission Rates for Light-Duty Vehicles in the Motor Vehicle Emissions Simulator (MOVES2010), Final Report, US Environmental Protection Agency, Washington, DC, USA. [online] Available from: <http://www.epa.gov/otaq/models/moves/documents/420r11011.pdf> (Accessed 5 October 2013), 2011.

30 Van Aardenne, J. A., Dentener, F. J., Olivier, J. G. J., Peters, J. A. H. W. and Ganzeveld, L. N.: The EDGAR 3.2 Fast Track 2000 dataset (32FT2000), [online] Available from: [http://themasites.pbl.nl/images/Description_of_EDGAR_32FT2000\(v8\)_tem61-46462.pdf](http://themasites.pbl.nl/images/Description_of_EDGAR_32FT2000(v8)_tem61-46462.pdf), 2005.

35 Van Vuuren, D. P., Edmonds, J. A., Kainuma, M., Riahi, K., Thomson, A. M., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J. F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J. and Rose, S.: The representative concentration pathways: an overview, *Climatic Change*, 109, 5–31, doi:10.1007/s10584-011-0148-z, 2011.

Venkataraman, C., Habib, G., Eiguren Fernandez, A., Miguel, A. H. and Friedlander, S. K.: Residential Biofuels in South Asia: Carbonaceous Aerosol Emissions and Climate Impacts, *Science*, 307, 1454–1456, 2005.

Venkataraman, C., Habib, G., Kadamba, D., Shrivastava, M., Leon, J. F., Crouzille, B., Boucher, O. and Streets, D. G.: Emissions from open biomass burning in India: Integrating the inventory approach with high resolution Moderate Resolution Imaging Spectroradiometer (MODIS) active fire and land cover data, *Global Biogeochem. Cycles*, 20(2), 1–12, doi:10.1029/2005GB002547, 2006.

Venkataraman, C., Sagar, A. D., Habib, G. and Smith, K. R.: The Indian National Initiative for Advanced Biomass Cookstoves: The benefits of clean combustion, *Energy for Sustainable Development*, 14, 63–72, 2010.

Vongmahadlek, C., Thao, P. T. B., Satayopas, B. and Thongboonchu, N.: An Inventory of Primary Gaseous Emissions from Thailand with Spatial and Temporal Allocation Profile, in Proceedings of the 3rd IASME/WSEAS International Conference on Energy & Environment, pp. 313–318, World Scientific and Engineering Academy and Society (WSEAS), Stevens Point, Wisconsin, USA. [online] Available from: <http://dl.acm.org/citation.cfm?id=1576758.1576812>, 2008.

Vongmahadlek, C., Pham, T. B. T., Satayopas, B. and Thongboonchoo, N.: A compilation and development of spatial and temporal profiles of high-resolution emissions inventory over Thailand., *J Air Waste Manag Assoc.* 59(7), 845–856, 2009.

Wang, S. X., Zhao, B., Cai, S. Y., Klimont, Z., Nielsen, C. P., Morikawa, T., Woo, J. H., Kim, Y., Fu, X., Xu, J. Y., Hao, J. M. and He, K. B.: Emission trends and mitigation options for air pollutants in East Asia, *Atmos. Chem. Phys.*, 14(13), 6571–6603, doi:10.5194/acp-14-6571-2014, 2014.

Wang, S. X., Zhao, X. J., Li, X. H., Wei, W. and Hao, J. M.: Emission characteristics of fine particles from grate firing boilers, *Environmental Science*, 30(4), 963–968, 2009.

Weitkamp, E. A., Lipsky, E. M., Paneras, P. J., Ondov, J. M., Polidori, A., Turpin, B. J. and Robinson, A. L.: Fine particle emission profile for a large coke production facility based on highly time resolved fence line measurements, *Atmospheric Environment*, 39(36), 6719–6733, doi:10.1016/j.atmosenv.2005.06.028, 2005.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S. and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6(11), 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y. and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric Chemistry and Physics*, 10(23), 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Weyant, C., Athalye, V., Ragavan, S., Rajarathnam, U., Lalchandani, D., Maithel, S., Baum, E. and Bond, T. C.: Emissions from South Asian Brick Production, *Environ. Sci. Technol.*, 48(11), 6477–6483, doi:10.1021/es500186g, 2014.

Weyant, C. L., Shepson, P. B., Subramanian, R., Cambaliza, M. O. L., Heimburger, A., McCabe, D., Baum, E., Stirm, B. H. and Bond, T. C.: Black Carbon Emissions from Associated Natural Gas Flaring, *Environ. Sci. Technol.*, 50(4), 2075–2081, doi:10.1021/acs.est.5b04712, 2016.

WHO: Health aspects of air pollution; Results from the WHO project “Systematic review of Health aspects of air pollution in Europe,” Technical report, Copenhagen, Denmark. [online] Available from: <http://www.aerohabitat.org/link/2006/12-06-2006%20-%20WHO,%20Health%20aspects%20of%20air%20pollution.pdf> (Accessed 2 December 2013), 2004.

- Wickramasinghe, A.: Energy access and transition to cleaner cooking fuels and technologies in Sri Lanka: Issues and policy limitations, *Energy Policy*, 39(12), 7567–7574, doi:10.1016/j.enpol.2011.07.032, 2011.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al Saadi, J. A., Orlando, J. J. and Soja, A. J.: The Fire INventory from NCAR (FINN): a high-resolution global model to estimate the emissions from open burning, *Geoscientific Model Development*, 4(3), 625–641, doi:10.5194/gmd-4-625-2011, 2011.
- Wiedinmyer, C., Yokelson, R. J. and Gullett, B. K.: Global Emissions of Trace Gases, Particulate Matter, and Hazardous Air Pollutants from Open Burning of Domestic Waste, *Environ. Sci. Technol.*, 48(16), 9523–9530, doi:10.1021/es502250z, 2014.
- Wobus, C., Planner, M., Sarofim, M. C., Moura, M. C. P. and Smith, S. J.: Future Arctic temperature change resulting from a range of aerosol emissions scenarios, *Earth's Future*, 4, doi:10.1002/2016EF000361, 2016.
- World Bank: Diesel Power Generation: Inventories and Black Carbon Emissions in Kathmandu Valley, Nepal, The World Bank, Washington DC, USA., 2014a.
- World Bank: Diesel Power Generation: Inventories and Black Carbon Emissions in Nigeria, The World Bank, Washington DC, USA., 2014b.
- World Bank and ICCI: On Thin Ice: How cutting pollution can slow warming and save lives, The World Bank, The International Cryosphere Climate Initiative, Washington D.C., U.S.A., 2013.
- Xu, Y.: Improvements in the operation of SO₂ scrubbers in China's coal power plants, *Environ. Sci. Technol.*, 45(2), 380–385, 2011.
- Xu, Y., Williams, R. H. and Socolow, R. H.: China's rapid deployment of SO₂ scrubbers, *Energy Environ. Sci.*, 2(5), 459–465, doi:10.1039/B901357C, 2009.
- Yan, F., Winijkul, E., Jung, S., Bond, T. C. and Streets, D. G.: Global emission projections of particulate matter (PM): I. Exhaust emissions from on road vehicles, *Atmospheric Environment*, 45(28), 4830–4844, doi:10.1016/j.atmosenv.2011.06.018, 2011.
- Yan, F., Winijkul, E., Streets, D. G., Lu, Z., Bond, T. C. and Zhang, Y.: Global emission projections for the transportation sector using dynamic technology modeling, *Atmospheric Chemistry and Physics*, 14(11), 5709–5733, doi:10.5194/acp-14-5709-2014, 2014.
- Yanowitz, J., McCormick, R. L. and Graboski, M. S.: In Use Emissions from Heavy Duty Diesel Vehicles, *Environmental Science & Technology*, 34(5), 729–740, doi:10.1021/es990903w, 2000.
- Yelverton, T. L. B., Holder, A. L. and Pavlovic, J.: Emissions removal efficiency from diesel gensets using aftermarket PM controls, *Clean Technologies and Environmental Policy*, 17(7), 1861–1871, doi:10.1007/s10098-015-0900-6, 2016.
- Yim, S. H. L., Lee, G. L., Lee, I. H., Allroggen, F., Ashok, A., Caiazzo, F., Eastham, S. D., Malina, R. and Barrett, S. R. H.: Global, regional and local health impacts of civil aviation emissions, *Environmental Research Letters*, 10(3), 34001, 2015.
- Yttri, K. E., Lund Myhre, C., Eckhardt, S., Fiebig, M., Dye, C., Hirdman, D., Ström, J., Klimont, Z. and Stohl, A.: Quantifying black carbon from biomass burning by means of levoglucosan — a one-year time series at the Arctic observatory Zeppelin, *Atmos. Chem. Phys.*, 14(12), 6427–6442, doi:10.5194/acp-14-6427-2014, 2014.

Zhang, Q., Klimont, Z., Streets, D., Huo, H. and He, K.: An anthropogenic PM emission model for China and emission inventory for the year 2001, *Progress in Natural Science*, 16(2), 223–231 (in Chinese), 2006.

Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T. and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, 2009.

Zhang, W., Wang, J., Zhang, B., Bi, J. and Jiang, H.: Can China Comply with Its 12th Five-Year Plan on Industrial Emissions Control: A Structural Decomposition Analysis, *Environ. Sci. Technol.*, 49(8), 4816–4824, doi:10.1021/es504529x, 2015.

Zhao, Y., Zhang, J. and Nielsen, C. P.: The effects of recent control policies on trends in emissions of anthropogenic atmospheric pollutants and CO₂ in China, *Atmos. Chem. Phys.*, 13, 487–513, 2013.

Zhi, G., Chen, Y., Feng, Y., Xiong, S., Li, J., Zhang, G., Sheng, G. and Fu, J.: Emission Characteristics of Carbonaceous Particles from Various Residential Coal Stoves in China, *Environ. Sci. Technol.*, 42(9), 3310–3315, doi:10.1021/es702247q, 2008.

Zhi, G., Peng, C., Chen, Y., Liu, D., Sheng, G. and Fu, J.: Deployment of Coal Briquettes and Improved Stoves: Possibly an Option for both Environment and Climate, *Environ. Sci. Technol.*, 43(15), 5586–5591, doi:10.1021/es802955d, 2009.

Formatted: German (Germany)

Table 1. Overview of the ECLIPSE emission data sets available to date

Version	Release date	Period covered	Comments; key features
V3	Nov 2013	2005, 2008, 2009, 2010	Estimates for 2008 and 2009 based on activity proxies and trends in internationally reported emissions; activity data for 2010 based on the IEA World Energy Outlook 2011 (IEA, 2011)
V4a	Jan 2014	2005, 2010, 2030, 2050	Major updates of EU-28 data (Amann et al., 2012)
V5	Apr 2014	1990-2010 ^a , 2015-30 ^a , 2040, 2050	IEA and FAO statistical data reimported for the period 1990-2010, international shipping included
V5a	Jul 2015	1990-2010 ^a , 2015-30 ^a , 2040, 2050	China 12 th 5-year plan included, improved regional resolution for Latin America, update of: global cement legislation, gas flaring, OC/OM ratios for residential combustion in Asia, Africa, Latin America, EU-28 update (Amann et al., 2015)

^a Estimated in 5-year intervals

5

Table 2. Residential-commercial sector fuel and source structure in GAINS

Fuels	Non-specific	Lighting	Three-stone	Fireplace	Stove ^a	Household boiler		Medium boiler	
						Manual	Auto	Manual	Auto
Gaseous fuels	•								
Liquid fuels	•	•							
Charcoal	•								
Coal					•	•	•	•	•
Biomass									
- Fuelwood			•	•	•	•	•	•	•
- Agricultural residue			•		•		•		•
- Dung cake			•		•				

^a distinguishing cooking and heating stoves as separate categories

Table 3. Mitigation measures distinguished in the residential-commercial sector in GAINS

Control option	Non-specific	Lighting	Three-stone	Fireplace	Stove		Household boiler		Medium boiler	
					Cooking	Heating	Manual	Auto	Manual	Auto
Improved	•			•	•	•	•			
New				•	•	•	•			
Fan stove					•					
Coal briquettes					•	•				
Hurricane lamp		•								
LED ^a lamp		•								
Pellets						•	•	•	•	•
Cyclone									•	•
ESP ^b						•	•	•		•

^a Light Emitting Diode

^b Electrostatic precipitator

5

Table 4. Brick sector technology structure assumed in GAINS for different regions

Kiln type	East Asia ^a	South-East Asia ^b	Central Asia	Africa	Latin America and Caribbean	Other
Traditional clamp	•	•	•	•	•	
Downdraft		•	•	•	•	
Moving chimney Bull's trench		•				
Fixed chimney Bull's trench		•				
Zig-Zag		•		•		
Vertical shaft brick kiln	•	•		•	•	
Marques kiln					•	
Hoffman kiln	•	•	•	•	•	
Tunnel kiln (coal)	•	•	•	•	•	•
Tunnel kiln (gas, oil)	•	•	•	•	•	•

^a Excluding OECD countries which are included in 'Other'

^b Including Middle East

Table 5. Overview of sectoral layers included in the gridded ECLIPSE emissions of PM

Sector layer	Included activities
Energy ^a	Power plants, energy production/conversion, fossil fuel distribution
Industry	Industrial combustion and processes
Residential	Residential and commercial combustion sources
Transport ^b	Road and non-road transport sources; including tyre and brake wear, road abrasion
Waste	Waste disposal, including trash burning
Agriculture	Livestock and arable land operations (ploughing, harvesting)
Agriculture (open burning) ^c	Open burning of agricultural residues (excluding forest and savannah burning)
Total	The sum of the above sectors
Shipping ^d	International shipping; available in version V5 and V5a

^a Includes associated petroleum gas flaring which is also available as a separate gridded layer

^b Does not include resuspension and international air and shipping; for the latter recommendation to use the RCP datasets, except for version V5 and V5a where international shipping was also included

^c The gridding proxy has been acquired from the GFED3.1 (van der Werf et al., 2010)

^d Available as a separate file where all pollutants' emissions are included; the resolution of this layer is 1°x1°

Table 6. Particulate matter amplification factors for high emitting light- and heavy-duty diesel and gasoline vehicles used in the GAINS model

	Light duty		Heavy duty	
	diesel	gasoline	diesel	gasoline
No control	3	6	3	4
Euro 1/I	3	6	3	4
Euro 2/II	5	6	5	10
Euro 3/III	5	10	5	10
Euro 4/IV	5	10	5	-
Euro 5/V	10	10	10	-
Euro 6/VI	10	10	10	-

Table 7. Regional emissions of particulate matter in 2010, ECLIPSE V5a, Gg year⁻¹

	PM ₁₀	PM _{2.5}	PM ₁	BC	OC	OM
Africa	9161	7973	6959	1347	3023	5207
East Asia	27172	20241	15291	2622	4974	7996
Europe and Russia	6027	4105	2781	660	897	1399
Latin and Central America	3736	2947	2358	508	994	1617
North America	1964	1268	917	249	382	594
Pacific	609	347	220	62	75	115
South-West and Central Asia	11982	9174	7654	1686	2796	4667
International shipping	1856	1758	1612	120	398	517
International aviation ^a	30	30	28	10	10	13
Global anthropogenic	62537	47843	37819	7264	13548	22125
Forest and savannah fires ^b	48207	33014	33014	2268	19489	31363
Global total	110744	80858	70834	9532	33037	53489

^a Values are middle of the range estimates referring to the ranges reported in Settler et al. (2013), Yim et al. (2015), and based on global fuel consumption and ranges of emission factors from Kinsey (2009)

^b GFED3.1 without agricultural waste burning; PM₁₀ value based on TPM (total particulate matter); PM₁ not available in GFED – here assumed equal PM_{2.5}

Field Code Changed

Table 8. Sectoral emissions of particulate matter in 2010, ECLIPSE V5a, Gg year⁻¹

	PM ₁₀	PM _{2.5}	PM ₁	BC	OC	OM
Agriculture	6555	3848	2883	337	1313	2364
Residential combustion	23078	21857	20742	4163	8852	15329
Industrial processes	12162	8340	4135	462	633	823
Large scale combustion	11561	6420	3812	136	164	248
Oil & gas, mining	1706	571	412	226	93	120
Transport – road	3339	2925	2524	1349	1116	1451
Transport – non-road	861	823	795	363	217	283
Waste	1388	1272	876	97	751	977
International shipping	1856	1758	1612	120	398	517
International aviation ^a	30	30	28	10	10	13
Global anthropogenic	62537	47843	37819	7264	13548	22125
Forest and savannah fires ^b	48207	33014	33014	2268	19489	31363
Global total	110744	80858	70834	9532	33037	53489

^a Values are middle of the range estimates based on the ranges reported in Settler et al. (2013), Yim et al. (2015), and based on global fuel consumption and ranges of emission factors from Kinsey (2009)

^b GFED3.1 without agricultural waste burning that is included based on GAINS estimates in category 'Agriculture'; PM₁₀ value based on TPM (total particulate matter); PM₁ not available in GFED – here assumed equal PM_{2.5}

Field Code Changed

Table 9. Comparison of global anthropogenic emissions of BC by sector, Gg year⁻¹

	1995		2000	
	Bond et al. (2004) ^a	This study (V5a)	Bond et al. (2013)	This study (V5a)
Diesel engines – road	792	872	840	980
Diesel engines – off-road	579	415	470	432
Residential combustion	2046	3703	1880	3891
<i>of which:</i>				
<i>Biomass cooking</i>		1660	1290	1711
<i>Biomass heating</i>	1481	411	260	392
<i>Residential coal</i>	480	710	330	908
<i>Other</i> ^b	85	922	^c	880
Agricultural burning	328	323	330	326
Industrial coal ^d	642	282	740	315
Other ^e	610	612	600	649
Global anthropogenic	4997	6206	4870	6594

^a Estimates for 1996

^b GAINS includes oil appliances and kerosene lamps – the latter are estimated in GAINS at 750 and 692 Gg BC in 1995 and 2000

^c Other residential sources (oil) included in category 'Other'

^d Includes coke and brick production, coal boilers and furnaces

^e Includes power plants, gas flaring, waste, gasoline engines in transport; for Bond et al. also oil use in residential sector




Figure 1. Change in implied $PM_{2.5}$ and BC emission factors for residential wood heating in selected countries and world regions; changes relative to 1990 in ECLIPSE V5a dataset.




Figure 2. Change in implied BC emission factors for road diesel vehicles in selected countries and world regions; changes relative to 1990 in ECLIPSE V5a dataset.




Figure 3. Change in implied PM_{2.5} emission factors for cement production and coal power plants in selected countries and world regions: changes relative to 1990 in ECLIPSE V5a dataset.





Figure 4. Global distribution of grids (0.5°x0.5°) for which flaring of associated petroleum gas emissions were calculated; derived from the 2009 data from Elvidge et al. (2011).



PM emission factors
(mg MJ⁻¹)

	Sulphur content (%S)				
	0.1	0.5	1	2	3
PM ₁₀	38	57	76	124	176
PM _{2.5}	37	54	73	120	167
BC	8.8	8.8	8.8	6	4
OC	17	20	24	35	37

Figure 52. Particulate matter emission factors for shipping used in the GAINS model




Figure 6.3. Global and regional emissions of PM species [Tg] and global energy consumption [EJ] in the period 1990-2010, ECLIPSE V5a.




Figure 74. Global and sectoral emissions of PM species [Tg] in the period 1990-2010, ECLIPSE V5a.





Figure 85. Global distribution of emissions of PM_{2.5} (left) and BC (right) in 2010 [Gg year⁻¹ per grid] from land-based sources, ECLIPSE V5a; the scale is the same across sectors but there is a factor ten between PM_{2.5} and BC



5 | **Figure 96.** Source-sector distribution of global anthropogenic emission of BC estimated with the GAINS model (ECLIPSE V5a) for the year 2000. Tg year⁻¹




Figure 107. Comparison of black carbon emission in this work (ECLIPSE V5a) with Lamarque et al. (2010) and Granier et al. (2011). The black star (*) symbols show emissions reported in global and regional studies listed in Table 8.1 in the SI.

Supplement of

Global anthropogenic emissions of particulate matter including black carbon


5 Zbigniew Klimont¹ et al.

¹International Institute for Applied Systems Analysis (IIASA), Laxenburg, 2361, Austria

Correspondence to: Zbigniew Klimont (klimont@iiasa.ac.at)

S1 Comparison of temporal distribution patterns

Fig S1.1 shows a comparison of the temporal patterns (it is an aggregate as the actual patterns are grid specific) for residential combustion sector, applied in the ECLIPSE project, with other data for selected countries.



5 **Figure S1.1.** Comparison of monthly distribution of emissions used in ECLIPSE with profiles from EDGAR (EC-JRC/PBL, 2010), EMEP (<http://emep.int/mscw/>), national Finish model FRES (Karvosenoja, 2008), and US EPA.

S2 Particulate matter emission factors for residential combustion

The GAINS model distinguishes three principal solid fuel stove categories: *traditional*, *improved* and *new stoves*. *Traditional heating stoves* using wood or coal as fuel have simple grate based firebox designs with usually only primary air supply and no heat storing components. Consequently there is restricted availability of air for combustion and poor mixing of air and pyrolysis gases. *Traditional stoves* in general have very high PM emission factors compared with more advanced technologies, but within this category the variability in the emission factors is also large. For example highest emission factors for traditional wood stoves have been measured in situations with restricted combustion air supply that leads to lower burn rate (Jordan and Seen, 2005). Such conditions might prevail when the user wants a lower heat supply to the room. *Improved stoves* have secondary air supply and heat storing components in the firebox construction that improve the combustion performance and reduce emissions of PM compared with the *traditional stoves*. *New stoves* represent the most advanced stove models on the market that have firebox, construction and airflow characteristics that optimize combustion efficiency. Additionally, an electrostatic precipitator (ESP) can be fitted into the latest stoves, which further improve the PM emission performance. GAINS distinguishes also *wood pellet stoves*. Pellets are a very homogenous fuel and combustion is more optimized than batch fired wood log stoves and thus also the PM emissions are lower than with wood log stoves.

A stove heats the surrounding room, but a boiler heats water to be circulated through a piping system to heat an entire house (Johansson et al., 2004). In *old-type wood log boilers* up-draught combustion is commonly used, which resembles the combustion in a stove; *modern wood boilers*, however, use downdraught combustion and often have an isolated burn-out zone (Johansson et al., 2004). In contrast to stoves, wood boilers can be connected to a water tank to store heat, which allows the boiler to be run at a regular heat output and to certain extent optimizing the combustion conditions. Storage tanks are common in modern wood boilers and also old boilers may be equipped with them, leading to lower emissions and higher efficiencies (Johansson et al., 2004). The single family house boilers are typically smaller than 50 kW_{th}, the larger residential boilers are allocated to a category *medium size boilers* where manual and automatic boilers are distinguished (Klimont et al., 2002; Kupiainen and Klimont, 2004, 2007). Such boilers might be an important emission source, especially when many of them are fired with coal, but there are not a of lot measurements available. The GAINS model relies on studies discussed previously (EEA, 2013; Klimont et al., 2002; Kupiainen and Klimont, 2004, 2007) but for a number of countries in Europe updates were made drawing on national information provided within EU consultations (Amann et al., 2015) and recent measurements in China where 100,000s of such installations are used in both residential as well as industrial sector (Wang et al., 2009).

GAINS distinguishes also open fireplaces as a separate category which is of relevance mostly in North America and some European countries, even though in Europe less than 5% of fuelwood would be used in such installations (Klimont et al., 2002; Kupiainen and Klimont, 2004, 2007).

Field Code Changed

Here we summarize the published measurements of emission factors for cooking and heating stoves boilers and compare them to the current ranges of region- and technology-specific GAINS values. The focus is on studies that appeared after the original development of the GAINS particulate matter module (Klimont et al., 2002; Kupiainen and Klimont, 2004, 2007).

5 **Table S2.1:** Summary of PM emission factors for residential wood boilers.

Emission factors (mg/MJ)				Shares (%)			References
PM	TC ^a	BC	OC	TC ^a	BC	OC	
wood log							
1300 (350-2200) ^b	715			55			(Boman et al., 2008) old, no accumulator, large fuel charge
120 (73-260) ^b	60			50			(Boman et al., 2008) old, no accumulator, adjusted fuel charge
95 (87-100) ^b	48			50			(Boman et al., 2008) old, with accumulator
44 (11-450) ^b	18			42			(Boman et al., 2008) modern, with accumulator
37 70-700	27	12	16	75	32	43	(Gaegauf et al., 2005), 35 kW apartment house GAINS ^c , >50 kW, uncontrolled boiler
230-1300		75-200	75-600				GAINS ^c , <50 kW, old uncontrolled boiler
80-520		32-50	22-230				GAINS ^c , <50 kW, improved
40-260		13-37	12-100				GAINS ^c , <50 kW, new/modern
wood chip							
				44	23	21	(Schmidl et al., 2011) 40 kW moving grate, start-up
				5	1	4	(Schmidl et al., 2011) 40 kW moving grate, full load
				35	33	2	(Schmidl et al., 2011) 40 kW moving grate, part load
85	8	2	6	9	2	7	(Gaegauf et al., 2005) 70 kW, institute building
wood pellet							
20		0.1	0.9	0.5	5		(Lamberg et al., 2011a) efficient combustion
12 (3-29) ^b		0.8 (0-14) ^b	0.3 (0-3) ^b	6 (0-51) ^b	2 (2-11) ^b		(Lamberg et al., 2011b), 25 kW, nominal load
16		1	0.1				(Tissari et al., 2008), 20 kW, nominal load
24		3	0.2				(Tissari et al., 2008), 20 kW, partial load
49	35	24	11	72	49	23	(Gaegauf et al., 2005) 10-32 kW, apartment house
8-25		0.8-1	0.4-1				GAINS ^c , >50 kW
20-68		5	2.5-10				GAINS ^c , <50 kW

^a Total Carbon (TC)

^b (min-max)

^c PM value refers to PM2.5

Table S2.2: Summary of PM emission factors for residential heating wood stoves.

Emission factors (mg/MJ)			Shares (%)		Reference
PM	BC	OC	BC	OC	
traditional					
673-1373	24-72	263-623	2-7	39-53	(Alves et al., 2011)
300-1400	-	-	2-9	35-50	(Gonçalves et al., 2011) incl. cold start
90-900	-	-	2-9	35-48	(Gonçalves et al., 2011) incl. hot start
750-1060	-	-	-	-	(Jordan and Seen, 2005), full airflow
1560-1700	-	-	-	-	(Jordan and Seen, 2005), half airflow
1870-3000	-	-	-	-	(Jordan and Seen, 2005), closed airflow
128-400	20	157	8	64	(McDonald et al., 2000)
-	39-43	70-390	5-14	47-67	Studies in Kupiainen& Klimont (2007)
150 ^a - 930 (2400) ^b	32 ^a - 100	60 ^a - 435 (1200) ^b	4-22	41-50	GAINS; the PM value represents PM2.5
improved					
22-180	-	-	-	-	(Boman et al., 2008)
86-105	9-11	52-58	-	-	(Fine et al., 2004)
130	88	39	68	30	(Gaegauf et al., 2005)
60-160	-	-	11-37	20-43	(Gonçalves et al., 2010)
75-97	15-28	17-35	24-32	27-39	(Schmidl et al., 2011)
38-350	-	-	-	-	(Pettersson et al., 2011)
-	56-79	11-16	-	-	Studies in Kupiainen& Klimont (2007)
55 ^a - 372	30 ^a - 95	11 ^a -133	25-55	19-35	GAINS; the PM value represents PM2.5
new					
67-122	13-15	43-67	-	-	(Fine et al., 2004), catalytic
72-89	21-33	16-32	30-37	22-36	(Schmidl et al., 2011)
30 ^a - 186	9 ^a - 30	8 ^a - 67	18-30	28-35	GAINS; the PM value represents PM2.5
pellet					
10-66	-	-	-	-	(Boman et al., 2008)
15-47	-	-	-	-	(Boman et al., 2011)
17	0.7	-	4	-	(Frey et al., 2014)
20	0.1	0.9	0.5	5	(Lamberg et al., 2011b)
3-29	0-14	0.1-3	0-51	2-11	(Lamberg et al., 2011a)
-	-	-	14	11	(Schmidl et al., 2011)
47-129	0.5-1.3	0.3-5.2	1-2	1-9	(Sippula et al., 2007)
10 ^a - 47	1.3 ^a - 4	2 ^a - 7	10-17	12-17	GAINS; the PM value represents PM2.5

^a The lowest values represent Swiss data

^b Norwegian wood stove

Table S2.3: Summary of PM emission factors for cookstoves using biofuels.

Emission factors (mg/MJ)			References
PM	BC	OC	
traditional			
530	44	250	(Just et al., 2013)
106	50	44	(Roden et al., 2009), 3-stone, lab measurements
515 (300-1000) ^a	83 (10-210) ^a	254 (90-660) ^a	(Roden et al., 2009), Honduras, field measurements
510 (280-510) ^b	65-75 (40-75) ^b	229 (125-229) ^b	GAINS ^c
improved			
150	80	20	(Just et al., 2013), rocket stove
270 (100-500) ^a			(Li et al., 2009), improved stoves, PM2.5
394 (120-700) ^a	102 (6-325) ^a	208 (60-460) ^a	(Roden et al., 2009), improved no chimney, field measurements
205 (105-270) ^b	50-75 (27-75) ^b	63 (31-68) ^b	GAINS ^c
new			
255 (40-720) ^a	116 (6-660) ^a	93 (33-370) ^a	(Roden et al., 2009), improved with chimney, field measurements
56-102	11-21	19-34	GAINS ^c
fan assisted			
86 (25-125) ^a	33 (6-100) ^a	38 (4-71) ^a	(Roden et al., 2009), fan assisted, lab measurements
54	33	14	(Just et al., 2013), gasifier with fan
17	4	9	GAINS ^c

^a (min-max)

^b central value for fuelwood and in brackets the whole range including also dung and agricultural residues

^c the PM value represents PM2.5

Table S2.4: Summary of PM emission factors for coal cooking and heating stoves

Emission factors (mg/MJ)			References
PM	BC	OC	
traditional			
805 (214-1360) ^a	250 (11-540) ^a	400 (116-710) ^a	(Zhi et al., 2009), portable stove, bituminous coals (Chen et al., 2009), simple low-efficiency stove without chimney, bituminous coals
	332 (10-610) ^a	472 (129-822) ^a	
351	135	108	GAINS ^b (cooking)
315-495	90-220	160-200	GAINS ^b (heating)
improved			
	466 (6-1377) ^a	248 (35-551) ^a	(Chen et al., 2009), high-efficiency stove with chimney
492	183	200	(Zhang et al., 2008), steel stove, brown coal
36	1	16	(Zhang et al., 2008), steel stove, bituminous coal
408 (155-685) ^a	40 (2-140) ^a	230 (78-470) ^a	(Zhi et al., 2009), bituminous coals
246	132	60	GAINS ^b (cooking)
315-350	82-200	88-112	GAINS ^b (heating)
new			
270	23	96	(Li et al., 2016), average for bituminous coals
176	108	32	GAINS ^b (cooking)
158-248	73-176	48-60	GAINS ^b (heating)
briquettes			
	16 (2-33) ^a	329 (71-668) ^a	(Chen et al., 2009), simple low-efficiency, no chimney
	4 (0.5-9) ^a	219 (27-423) ^a	(Chen et al., 2009), high-efficiency, with chimney
184	3	80	(Zhang et al., 2008), steel stove
440 (98-930) ^a	12 (2-23) ^a	233 (67-460) ^a	(Zhi et al., 2009), traditional portable stove
202 (90-346) ^a	2 (0.5-6) ^a	124 (36-217) ^a	(Zhi et al., 2009), improved stove with chimney
17	0.4	6.5	(Li et al., 2016), semi-coke briquettes
23-135	0.3-1	9-55	GAINS ^b

^a (min-max)^b the PM value represents PM2.5

S3 Summary of particulate matter emissions factors for diesel generators

Note that the ranges presented for GAINS represent the spread across GAINS regions or technologies (if a category refers to an aggregate across several measures) defined in the GAINS model.

5 **Table S3.1:** Summary of PM emission factors for diesel generator sets

Emission factors (mg/MJ)			Shares (%)		Reference
PM	BC	OC	BC	OC	
69-189					Uma et al. (2004), 10 kW (higher value), 40 kW (lower value)
139			66%		Bond et al. (2004)
13/22					Gilmore et al. (2006), ICE 10 kW, with/without DPF
		116-585			Watson et al. (2006) ^a
59-190	12-54	30-120	31%	51%	Shah et al. (2007) ^b 300 kW 1985 Detroit Diesel V92, 2-str
45-219	30-145	8-56	67%	21%	Shah et al. (2007) ^b 350 kW 2000 Cat 3406C, 4-str
22-143	10-80	6-37	53%	25%	Shah et al. (2007) ^b 300 kW 1985 Detroit Diesel V92, 2-str, DOC
59-203	28-145	4-16	67%	8%	Shah et al. (2007) ^b 350 kW 2000 Cat 3406C, 4-str, DOC
23-190	9-96	10-81	49%	36%	Shah et al. (2007) ^b 300 kW 1985 Detroit Diesel V92, 2-str, DOC+FBC
4-26	2.5-19	1-3	76%	15%	Shah et al. (2007) ^b 350 kW 2000 Cat 3406C, 4-str, passive-DPF
1-3	0.8-2	1-6	67%	49%	Shah et al. (2007) ^b 350 kW 2000 Cat 3406C, 4-str, active-DPF
				20-70%	Watson et al. (2008)
14-42					Zhu et al. (2009) ^c
174-433					Tsai et al. (2010) ^d
55					Anayochukwu et al. (2013)
GAINS emission factors; the PM value represents PM2.5					
96	40	28	41%	29%	No control
48-64	20-26	14-19	41%	29%	Controlled, no DPF
<1-3	0.5-2	0.3-0.8			Controlled, with DPF

^a Higher value with 10% load and lower value with 100% load for a 100 kW DG set

^b Lower value with 100% load and higher value with 10% load, share of BC/OC is average of all loads

^c Average of 14 military diesel generators with rated capacities of 10, 30, 60, and 100 kW under different load conditions. The fleet average EFs are 1.2+/-0.6 g/kg for PM.

^d Higher value with no load and lower value with 10 kW

10

15

S4 Transport sector

Note that the ranges presented for GAINS represent the spread across GAINS regions or technologies (if a category refers to an aggregate across several measures) defined in the GAINS model.

5

Table S4.1: Comparison of selected measured emissions factors and ranges used in the GAINS model for diesel and gasoline cars and light duty vehicles.

	Emission factors (mg/MJ)			Reference
	PM	BC	OC	
Diesel				
Pre-/early regulation	44-67	9-17	13-34	(Subramanian et al., 2009)
Euro 1	67	17	13	(Subramanian et al., 2009)
Euro 2	30-33	7-16	8-12	(Cheung et al., 2009; Subramanian et al., 2009)
Euro 3	10-29			(Graham, 2005)
Euro 4	6-11	3-8	1-2	(Cheung et al., 2009; Geller et al., 2006)
Euro 4 with DPF	0.2-0.3	0.02-0.1	0.02-0.06	(Dwyer et al., 2010; Louis et al., 2016)
Pre-/early regulation	56-133	38-76	21-51	GAINS ^a
Euro 1	22-50	16-35	5-11	GAINS ^a ; for developing countries the values only marginally lower than pre/early regulation
Euro 2	15-40	12-32	3-6	GAINS ^a ; for developing countries the values only marginally lower than pre/early regulation
Euro 3	11-29	10-22	1-2	GAINS ^a
Euro 4	5-20	4-17	0.5-1.6	GAINS ^a
Euro 4 with DPF	0.5-1	0.1-0.3	0.1-0.6	GAINS ^a
Gasoline				
Pre-/early regulation	4-10	0.5-2	2-10	see studies in Kupiainen and Klimont (2004, 2007)
Euro 1, 2	1-4	0.6-1.5	0.3-1.6	see studies in Kupiainen and Klimont (2004, 2007)
Euro 3	0.2-2	0.01-0.2	0.2-0.6	(Cheung et al., 2009; Geller et al., 2006; Graham, 2005)
Euro 4		0.001-0.4		(Louis et al., 2016)
Pre-/early regulation	6	1	3-4	GAINS ^a
Euro 1, 2	1-4	0.2-1	0.3-1.7	GAINS ^a
Euro 3, 4	0.3-1.1	0.05-0.5	0.1-0.4	GAINS ^a

^a the PM value represents PM2.5

Table S4.2: Comparison of selected measured emissions factors and ranges used in the GAINS model for diesel heavy duty vehicles

	Emission factors (mg/MJ)			Reference
	PM	BC	OC	
Diesel heavy duty trucks				
Pre-/early regulation	28-33			(Herner et al., 2009; Yanowitz et al., 2000)
	44-244	4-50	15-122	(Subramanian et al., 2009), Bangkok, Thailand
	30-50			(Liu et al., 2009), on-road measurements in China
Euro I	11			(Yanowitz et al., 2000)
	22	4	9	(Subramanian et al., 2009), Bangkok, Thailand
Euro II	10-20			(Liu et al., 2009), on-road measurements in China
	22-44	2-9	7-22	(Subramanian et al., 2009), Bangkok, Thailand
Euro III	7-17	16		(Liu et al., 2009; Wang et al., 2011), on-road measurements in China
	3-7	9		(Liu et al., 2009; Wang et al., 2011), on-road measurements in China
Euro IV		4		(Wang et al., 2011), on-road measurements in China
Pre-/early regulation	34-107	17-53	10-37	GAINS ^a
Euro I	21-71	17-53	6-19	GAINS ^a
Euro II	11-44	7-30	2-10	GAINS ^a
Euro III	10-27	8-25	2-7	GAINS ^a
Euro IV, V	2-7	2-5	0.3-1	GAINS ^a
Euro VI	0.1-0.4	0.01-0.06	0.06-0.15	GAINS ^a

^a the PM value represents PM2.5

Table S4.3: Comparison of selected measured emissions factors and ranges used in the GAINS model for non-road machinery.

	Emission factors (mg/MJ)			Reference
	PM	BC	OC	
Diesel locomotives				
Pre-/early regulation	49-67			(Dincer and Elbir, 2007; Johnson et al., 2013; Tang et al., 2015)
Regulated	20-40	20		(Dincer and Elbir, 2007; Galvis et al., 2013; Johnson et al., 2013; Tang et al., 2015)
	30	14		(Galvis et al., 2013)
	20	15		(Jaffe et al., 2014)
	37	21		(Krasowsky et al., 2015)
pre-regulated	49-98	24-45	12-25	GAINS ^a
regulated (stage I)	26-49	11-22	6-12	GAINS ^a
Agriculture				
Pre-regulation	141	58	41	(Kupiainen and Klimont, 2007)
	89	49		(EEA, 2013)
Stage I	20-39	16-21		(EEA, 2013)
Stage II	15	11.5		(EEA, 2013)
Pre-regulation	100-170	41-70	29-50	GAINS ^a
Stage I	57-96	23-40	16-27	GAINS ^a
Stage II, III	27-43	10-19	8-12	GAINS ^a
Stage IV,V	6-10	0.7-1.2	0.5-0.8	GAINS ^a
Construction				
Pre-regulation	140	65	30	(Kupiainen and Klimont, 2007)
	103	56		(EEA, 2013)
Stage I	85	47		(EEA, 2013)
Pre-regulation	95-140	46-68	21-31	GAINS ^a
Stage I	57-76	26-39	12-18	GAINS ^a
Stage II, III	24-36	12-17	5-8	GAINS ^a
Stage IV,V	6-8	0.8-1.2	0.4-0.6	GAINS ^a

^a the PM value represents PM2.5

Table S4.4: Comparison of selected measured emissions factors and ranges used in the GAINS model for 2-wheelers.

	Emission factors (mg/MJ)			References
	PM	BC	OC	
2-stroke				
Euro 0 mopeds	250 (198-295)			(Spezzano et al., 2008), hot start
	160 (121-878)			(Spezzano et al., 2008), cold start
Euro 1 mopeds	169 (102-235)			(Spezzano et al., 2008), hot start
	42 (26-71)			(Spezzano et al., 2008), cold start
Euro 2 mopeds	147-217			(Spezzano et al., 2008), hot start
	13-215			(Spezzano et al., 2008), cold start
CNG rickshaw, Delhi, India	124-160			(Grieshop et al., 2012)
Euro 0 mopeds	132-1400	10-75	90-1015	GAINS ^a
Euro 1 mopeds	12-450	7-49	40-300	GAINS ^a
Euro 2 mopeds	37-280	6-45	23-172	GAINS ^a
Euro 3 mopeds	14-112	3-30	8-61	GAINS ^a
4-stroke				
Motorcycles	2.6-3.7			(Yang et al., 2005), cold start
Euro 0 motorcycles	4			(Spezzano et al., 2007)
Euro 1 motorcycles	2			(Spezzano et al., 2007)
Rickshaw, Delhi, India	30-45			(Grieshop et al., 2012)
CNG rickshaw, Delhi, India	12-13			(Grieshop et al., 2012)
Euro 0 motorcycles	6-14	1-2	3-9	GAINS ^a
Euro 1 motorcycles	5-12	1-2	2-7	GAINS ^a
Euro 2 motorcycles	3-5	0.5-0.8	0.4-1.7	GAINS ^a
Euro 3 motorcycles	2-3	0.5-0.75	0.3-1.4	GAINS ^a

^a the PM value represents PM2.5

Table S4.5: Summary of PM emission factor ranges used in the GAINS model for non-exhaust transport sources

	Emission factors (mg/km)			
	PM ₁₀	PM _{2.5}	BC	OC
Brake wear				
Cars	3.5 – 12	2.5 – 5	0.05 – 0.12	0.8 – 2.2
Light duty vehicles	3.5 – 19	2.5 – 8	0.05 – 0.2	0.8 – 3.5
Heavy duty vehicles	21 – 53	13 – 21	0.25 – 0.5	5 – 17
Tyre wear				
Cars	1.5 – 9	0.15 – 0.7	0.2 – 1	0.5 – 2.4
Light duty vehicles	2.5 – 7	0.2 – 0.7	0.35 – 1	0.85 – 2.4
Heavy duty vehicles	40 – 47	4.2 – 4.7	6 – 7	15 – 17
Road abrasion				
Cars & Light duty vehicles	7 – 10	3 – 5	0.15 – 0.6	0.7 – 1
	30 – 140 ^a	20 – 80 ^a	0.2 – 1.5 ^a	4 – 14 ^a
Heavy duty vehicles	38 – 50	18 – 27	0.7 – 1	3 – 5

^a vehicles with studded tires; variation between estimates for Scandinavian and alpine countries

S5 Industry

GAINS model PM emission factors (as used for the ECLIPSE V5a) for brick making compared with values used in GAINS previously (UNEP/WMO, 2011) and recent set of measurements on typical kilns used in South Asia (Weyant et al., 2014).

5 **Table S5.1:** Comparison of emissions factors used in the GAINS model for brick kilns with selected other studies.

	Emission factors (g kg ⁻¹ brick)			References
	PM2.5	BC	OC	
Clamp kiln				
	1.6	0.35	0.3	(UNEP/WMO, 2011) ^a
	1	0.3	0.1	GAINS (Asia)
	1	0.35	0.15	GAINS (Latin America and Africa)
Downdraft kiln				
	0.49	0.19	0.07	(Weyant et al., 2014)
	0.97	0.29	0.09	GAINS (all regions)
Bull's trench kiln (BTK)				
	1.31	0.27	0.24	(UNEP/WMO, 2011) ^a
	0.19 (0.08-0.33)	0.15 (0.09-0.27)	0.007	(Weyant et al., 2014) ^b
	0.18/0.8	0.13/0.25	0.01/0.07	GAINS (Asia); fixed /moving chimney
Vertical shaft brick kiln (VSBK)				
	0.77	0.175	0.15	(UNEP/WMO, 2011) ^a
	0.07 (0.005-0.009)	0.0015 (0.001-0.002)	0.014	(Weyant et al., 2014) ^b
	0.093	0.001-0.004	0.002-0.059	GAINS (Asia)
	0.093	0.002	0.059	GAINS (Latin America and Africa)
Zig-zag kiln				
	0.06 (0.03-0.06)	0.01 (0.014-0.03)	0.005	(Weyant et al., 2014) ^b
	0.13	0.04	0.02	GAINS (Asia)
Tunnel kiln (coal)				
	0.28	0.0035	0.003	(UNEP/WMO, 2011) ^a
	0.24	0.001	<0.00	(Weyant et al., 2014)
	0.18	0.002	0.0035	GAINS (all regions)
Hoffman kiln				
	0.08	0.003	0.005	GAINS (all regions)
Marquez kiln (MK)				
	0.15	0.06	0.02	GAINS (Latin America)

^a Previous version of the GAINS model was used

^b Central value and ranges of average values; all measurement data provided in the original study

Brick sector production structure in Asia has been analysed in a number of studies addressing either the whole region where selected countries, typically key producers including China, India, Pakistan, Bangladesh, Vietnam, are discussed (AIT, 2003; BASIN, 1999; FAO, 1993; Heierli and Maithel, 2008; Maithel, 2014) or focusing on particular countries like China (Zhang, 1997), India (BASIN, 1998; Maithel et al., 2012; Verma and Uppal, 2013), Bangladesh (Croitoru and Sarraf, 2012; Guttikunda et al., 2013; World Bank, 2011), Cambodia (Rozemuller, 1999), Afghanistan (Samuel Hall Consulting, 2011), Nepal (Heierli et al., 2007). More recently, a number of development programs and local air pollution studies focused on this sector in the Latin America and Caribbean regions, including some where information about kiln structure was collected (Bellprat, 2009; EELA, 2011; Erbe, 2011; PRAL, 2012; Stratus Consulting, 2014; SwissContact, 2014a). Fewer assessments exist for Africa (Scott, 2013; SwissContact, 2014c). The updated and country specific data for Latin America and Caribbean (LAC) is included only in version V5a of ECLIPSE since the previous versions included just five regions for the whole LAC; Argentina, Brazil, Chile, Mexico, other LAC.


GAINS activity data has been built on the basis of several regional studies where production, energy efficiency, and sector structure were discussed, i.e., Asia (AIT, 2003; Co et al., 2009; Croitoru and Sarraf, 2012; FAO, 1993; Guttikunda et al., 2013; Heierli et al., 2007; Heierli and Maithel, 2008; Maithel, 2014; Maithel et al., 2012; Samuel Hall Consulting, 2011; Subrahmanya, 2006; Verma and Uppal, 2013; World Bank, 2011; Zhang, 1997), Africa (Alam, 2006; Scott, 2013; SwissContact, 2014c), Latin America and Caribbean (Bellprat, 2009; EELA, 2011; PRAL, 2012; Stratus Consulting, 2014; SwissContact, 2014b). For several countries where we found no regional analysis, the United Nations data on ‘building bricks, made of clay’ was used (<http://unstats.un.org/unsd/industry/commoditylist2.asp>). There are some differences between different versions of the ECLIPSE datasets; specifically during the development of the V5a version, the data for all countries in Latin America and Caribbean was revisited and updated, and a new version of the UN statistics was downloaded.

Table S5.2: Brick production in key regions; GAINS model assumptions - ECLIPSE V5a, Tg bricks year⁻¹

	1990	1995	2000	2005	2010
Global	1542	2357	2688	3022	3574
Asia	1314	2130	2530	2819	3320
<i>of which:</i>					
<i>China</i>	1050	1800	2106	2204	2508
<i>India</i>	131	178	254	406	553
<i>Vietnam</i>	20	20	27	46	65
<i>Bangladesh</i>	9	15	18	17	25
<i>Pakistan</i>	32	41	50	59	74
<i>Other Asia</i>	71	75	76	87	95
Africa	18	18	15	17	22
Europe	158	156	72	82	79
Latin America and Caribbean	29	30	43	75	127
Other	23	23	27	29	25

S6 Emissions of PM species over time in ECLIPSE datasets

The Fig S6.1 shows emissions of PM₁₀, PM_{2.5}, BC, and OC calculated with the GAINS model within different versions of the ECLIPSE dataset. These have been created between 2013 and 2015 and include a number of updates to activity data and emission factors; the methodology remained the same. The changes for PM₁₀ and PM_{2.5} are similar, driven by updates of activity data, i.e., the energy statistics from IEA were reimported for the whole time series for the version V5 and V5a and for China the regional coal statistics were used. Control strategies have been updated continuously considering more up to date information available over time. Additionally, in version V5a Latin America and Caribbean were revised since higher spatial resolution was introduced in the GAINS model. Several of the above mentioned updates affected also emissions of BC and OC but the largest impact on the BC emissions was due to introduction of emissions from kerosene lamps which were not specifically distinguished in V4a; this represents the key component of the higher emissions in V5, V5a. For OC the change is in the opposite direction and V5a has significantly lower emissions than previous versions which is due to update of the OC emission factor for residential cooking in Asia and Africa.



15 **Figure S6.1.** Global emissions of PM (excluding international shipping and open biomass burning) in the period 1990-2010 in different ECLIPSE scenarios; unit [Tg year⁻¹]

Table S6.2: Global anthropogenic (excluding international shipping & aviation) emissions of PM10 in ECLIPSE V5a; [Gg year⁻¹]

Region	1990	1995	2000	2005	2010
1 Canada	333	315	345	337	334
2 USA	2416	2158	1954	1920	1630
3 Mexico	643	621	653	574	572
4 Rest Central America	454	455	479	498	516
5 Brazil	1228	1295	1250	1385	1456
6 Rest South America	1018	1155	1138	1131	1192
7 Northern Africa	1022	1152	1355	1144	1194
8 Other Africa	4393	4993	5831	6425	7150
10 South Africa	682	738	747	848	818
11 Western Europe	3294	2458	2031	1747	1577
12 Central Europe	2944	1608	1236	1046	1038
13 Turkey	1007	756	525	477	571
14 Ukraine+	1854	856	679	707	680
15 Asia-Stan	836	325	303	314	392
16 Russia+	5833	2434	2314	2316	2161
17 Middle East	836	954	1055	962	996
18 India	7828	8785	8654	7952	8061
19 Korea	1227	913	844	816	768
20 China+	14057	17612	18205	21230	21976
21 Southeastern Asia	2291	2855	2783	2451	2526
22 Indonesia+	1383	1576	1673	1768	1902
23 Japan	545	435	354	319	267
24 Oceania	295	303	354	354	342
25 Rest South Asia	1695	1894	2211	2349	2533
Global	58112	56646	56974	59071	60651

Table S6.3: Global anthropogenic (excluding international shipping & aviation) emissions of PM2.5 in ECLIPSE V5a; [Gg year⁻¹]

Region	1990	1995	2000	2005	2010
1 Canada	252	244	250	242	241
2 USA	1629	1482	1296	1275	1027
3 Mexico	495	498	526	459	454
4 Rest Central America	395	394	416	428	446
5 Brazil	938	974	933	1054	1098
6 Rest South America	825	933	909	901	949
7 Northern Africa	762	852	982	847	909
8 Other Africa	4056	4606	5308	5887	6575
10 South Africa	408	444	431	501	490
11 Western Europe	2125	1700	1360	1157	1037
12 Central Europe	1610	1020	843	752	775
13 Turkey	585	480	388	356	425
14 Ukraine+	1072	531	464	483	455
15 Asia-Stan	562	222	211	222	283
16 Russia+	3702	1614	1530	1495	1413
17 Middle East	686	778	845	784	794
18 India	5768	6453	6472	5957	6032
19 Korea	784	600	547	565	529
20 China+	10863	13072	13633	15673	16096
21 Southeastern Asia	1878	2257	2198	1974	2012
22 Indonesia+	1230	1371	1447	1510	1604
23 Japan	337	295	236	203	160
24 Oceania	188	193	210	201	188
25 Rest South Asia	1455	1629	1859	1962	2065
Global	42606	42640	43294	44888	46055

Table S6.4: Global anthropogenic (excluding international shipping & aviation) emissions of PM1 in ECLIPSE V5a; [Gg year⁻¹]

Region	1990	1995	2000	2005	2010
1 Canada	184	195	196	187	190
2 USA	1163	1095	949	930	727
3 Mexico	375	378	395	361	357
4 Rest Central America	329	331	353	366	390
5 Brazil	706	720	718	819	846
6 Rest South America	657	732	708	712	764
7 Northern Africa	447	476	514	485	542
8 Other Africa	3724	4213	4838	5416	6064
10 South Africa	285	309	307	354	354
11 Western Europe	1397	1171	966	834	751
12 Central Europe	894	667	619	579	607
13 Turkey	386	341	286	263	311
14 Ukraine+	565	325	279	278	261
15 Asia-Stan	292	154	146	154	198
16 Russia+	1988	1078	1011	936	852
17 Middle East	501	562	596	614	615
18 India	4500	4992	5016	4700	5031
19 Korea	635	510	450	464	429
20 China+	9153	11251	11731	12473	11606
21 Southeastern Asia	1800	2204	2093	1791	1803
22 Indonesia+	1135	1254	1315	1373	1453
23 Japan	258	229	157	126	87
24 Oceania	140	146	155	143	133
25 Rest South Asia	1303	1445	1625	1714	1811
Global	32816	34780	35422	36073	36180

Table S6.5: Global anthropogenic (excluding international shipping & aviation) emissions of BC in ECLIPSE V5a; [Gg year⁻¹]

Region	1990	1995	2000	2005	2010
1 Canada	44	49	51	49	49
2 USA	311	291	281	279	201
3 Mexico	76	77	82	84	88
4 Rest Central America	52	54	61	65	71
5 Brazil	143	148	160	171	179
6 Rest South America	115	135	140	150	169
7 Northern Africa	127	120	117	121	140
8 Other Africa	752	836	942	1030	1135
10 South Africa	57	59	57	74	72
11 Western Europe	331	335	307	287	246
12 Central Europe	126	112	112	121	134
13 Turkey	60	59	53	51	67
14 Ukraine+	88	59	45	41	36
15 Asia-Stan	50	28	33	38	55
16 Russia+	439	251	238	226	177
17 Middle East	174	183	210	243	262
18 India	853	931	884	908	1022
19 Korea	135	84	71	84	74
20 China+	1348	1347	1655	1823	1924
21 Southeastern Asia	300	299	304	328	333
22 Indonesia+	243	260	275	279	290
23 Japan	67	74	66	50	29
24 Oceania	30	32	35	35	33
25 Rest South Asia	288	304	325	337	348
Global	6210	6129	6505	6872	7134

Table S6.6: Global anthropogenic (excluding international shipping & aviation) emissions of OC in ECLIPSE V5a; [Gg year⁻¹]

Region	1990	1995	2000	2005	2010
1 Canada	72	77	77	72	74
2 USA	448	434	388	379	308
3 Mexico	162	162	164	158	155
4 Rest Central America	144	149	159	169	181
5 Brazil	251	258	275	311	314
6 Rest South America	297	329	315	324	344
7 Northern Africa	145	150	155	166	192
8 Other Africa	1627	1842	2124	2408	2701
10 South Africa	101	108	110	129	130
11 Western Europe	495	422	343	284	253
12 Central Europe	224	201	217	220	234
13 Turkey	114	108	95	88	107
14 Ukraine+	149	102	82	77	72
15 Asia-Stan	90	66	62	64	86
16 Russia+	509	332	304	256	231
17 Middle East	190	217	220	237	229
18 India	1530	1623	1596	1630	1755
19 Korea	200	157	147	157	148
20 China+	3147	3264	3500	3564	3599
21 Southeastern Asia	526	548	567	598	632
22 Indonesia+	431	473	514	551	595
23 Japan	51	54	49	40	29
24 Oceania	52	55	57	51	46
25 Rest South Asia	502	562	628	680	726
Global	11456	11695	12150	12610	13140

The following charts in this section show emissions of PM species by key sectors for seven global regions defined as in Figure 3 in the main paper. These charts are the same as Figure 4 in the paper where global emissions are shown. Note that the scale varies across sectors and also across regions. For all regions, except Pacific, the units are million tons [Tg year⁻¹]; for Pacific they are [Gg year⁻¹].




Figure S6.2. Emissions of PM species by key sectors in *East Asia* in the period 1990-2010 [Tg year^{-1}]; ECLIPSE V5a




Figure S6.3. Emissions of PM species by key sectors in *Other Asia* in the period 1990-2010 [Tg year^{-1}]; ECLIPSE V5a




Figure S6.4. Emissions of PM species by key sectors in *Africa* in the period 1990-2010 [Tg year^{-1}]; ECLIPSE V5a




Figure S6.5. Emissions of PM species by key sectors in *Latin America and Caribbean* in the period 1990-2010 [Tg year^{-1}]; ECLIPSE V5a




Figure S6.6. Emissions of PM species by key sectors in *Europe and Russia* in the period 1990-2010 [Tg year^{-1}]; ECLIPSE V5a




Figure S6.7. Emissions of PM species by key sectors in *North America* in the period 1990-2010 [Tg year^{-1}]; ECLIPSE V5a





Figure S6.8. Emissions of PM species by key sectors in Pacific in the period 1990-2010 [Gg year⁻¹]; ECLIPSE V5a

S7 Regional resolution

The spatial resolution of the GAINS model is discussed section 2.4 of the paper and the list of all 170 regions can be obtained from the online model. In principle, GAINS distinguishes single countries in Europe (exception in Russia for which European and Asian part is included separately) North America, Australia and New Zealand, for Asia several larger countries are divided into provinces or states (larger administrative units in, e.g., China, India, Indonesia, Japan, etc.) while Middle East represented as one region or (most recent versions) distinguishes Iran, Saudi Arabia, Israel, and the rest of Middle East. Africa is divided into four regions: South Africa, Egypt, North Africa, and other Africa. Latin America and Caribbean includes now 13 regions with all larger countries treated separately while Central America as well as Caribbean states are grouped in two regions. While such resolution of 170 regions is used for the calculation of emissions, the presentation of data and results differs between the on-line models available for specific world regions, e.g., for Europe and Asia the full resolution is available, while in the global model application (<http://magcat.iiasa.ac.at/gains/IAM/index.login>) the data and results are presented for 25 regions (Fig. S7.1). This follows closely the IMAGE model¹ resolution; often used or compatible with several global integrated assessment models.



15 **Figure S7.1.** Regions distinguished in the global GAINS online application.

¹ http://themasites.pbl.nl/models/image/index.php/Region_classification_map

S8 Sectoral resolution

Table S8.1: Source sector resolution in the GAINS model for calculation of PM emissions

Key source category	Source sectors	Fuel category or activity type
<i>Energy sector</i>	Power plants (distinguishing small, large, old, new plants); Diesel generators;	Coal, oil, gas, biomass, waste
	Extraction and distribution of solid and liquid fuels (fugitive as well as combustion from gas flaring)	Coal, oil
	Briquette production	Production
<i>Residential combustion</i>	Cooking stoves; Heating (distinguishing fireplaces, stoves, house boilers, mid-size residential boilers)	Coal, fuelwood, dung, oil, gas, agricultural residues, charcoal
	Kerosene lighting	Kerosene
	Waste (trash) burning	Waste
<i>Industrial combustion</i>	Iron and Steel; Pulp and Paper; Chemical; Non-ferrous metals; Non-metallic minerals (excl. Bricks); Other	Coal, oil, gas, biomass, waste
<i>Industrial processes</i>	Iron and steel industry divided into: Pig iron; Coke ovens; Agglomeration plants – pellets; Agglomeration plants – sinter; Open hearth; Electric Arc; Basic oxygen; Rolling mills; Cast Iron	Production
	Non-ferrous metals (copper and nickel smelters); Primary aluminium; Secondary aluminium; Cement; Lime; Carbon black production; Glass production; Mineral fertilizer production; Brick manufacturing; Pulp and paper	Production
	Refineries	Crude oil throughput
	Handling and storage of bulk industrial and agricultural products (fugitive)	Million tons of products
<i>Road transport</i>	Passenger cars and vans; Light duty vehicles; Heavy duty vehicles; Busses; Motorcycles (4-stroke); Mopeds (2-stroke)	Gasoline, diesel, CNG, LPG, km driven (for calculation of non-exhaust emissions)
<i>Non-road transport</i>	Agricultural and forestry; Construction and mining; Railways; Inland navigation; Coastal shipping; Aviation (landing and take-off); 2-stroke engines (e.g., in household, forestry, etc.); Other land based machinery	Diesel, gasoline, CNG, jet fuel and kerosene, heavy fuel oil, coal
<i>Agriculture</i>	Arable land operations	Arable land area
	Livestock housing	Cattle, pigs, poultry
	Open burning of agricultural waste	Waste burned
<i>Other</i>	Fireworks; Cigarette smoking; Barbeques; Cremation	Population
	Construction (fugitive)	Constructed area

S8 Comparison of regional estimates with selected studies

The table S8.1 provides ECLIPSE V5a PM estimates for selected regions and years (from the period 2000-2010) and compares them with selected regional peer-reviewed studies.

5 **Table S8.1:** Comparison of regional estimates for anthropogenic ^a emissions of PM species, Gg year⁻¹

Region – (Source) – Year	PM10	PM2.5	PM1	BC	OC
Global					
<i>This study – 1995</i>	57830	43762	35902	6206	11949
(Bond et al., 2004) - 1996				4997	10481
<i>This study - 2000</i>	58366	44613	36741	6595	12449
(Bond et al., 2013) - 2000				4870	
<i>This study - 2010</i>	62537	47843	37819	7264	13548
HTAP_v2 (Janssens-Maenhout et al., 2015) - 2010	50292	32761		5525	13581
China					
<i>This study - 2000</i>	18061	13554	11685	1646	3487
(Cao et al., 2006) - 2000				1496	4211
(Streets et al., 2003) - 2000				1049	3385
(Klimont et al., 2009) - 2000				1345	3205
(Lu et al., 2011) - 2000				1244	2823
(Ohara et al., 2007) - 2000				1093	2563
(Bond et al., 2013) - 2000				1200 ^b	2800 ^b
(Zhang et al., 2006) - 2001	17120	12100			
<i>This study - 2005</i>	21087	15593	12428	1813	3552
(Zhang et al., 2009) - 2006	18223	13266		1811	3217
(Klimont et al., 2009) - 2005				1366	2812
<i>This study - 2010</i>	21827	16019	11564	1915	3589
(Lu et al., 2011) - 2010				1838	3907
(Kurokawa et al., 2013) - 2008	21606	14514		1589	3081
(Guan et al., 2014) - 2010		12100			
HTAP_v2 (Janssens-Maenhout et al., 2015) - 2010	16615	12199		1764	3384
(Kondo et al., 2011) - 2008				1940	
India					
<i>This study - 2000</i>	8654	6472	5016	884	1596
(Streets et al., 2003) - 2000				600	2837
(Ohara et al., 2007) - 2000				795	3268

Region – (Source) – Year	PM10	PM2.5	PM1	BC	OC
(Klimont et al., 2009) - 2000				842	1887
(Lu et al., 2011) - 2000				736	1990
(Bond et al., 2013) - /2000				500 ^b	1600 ^b
(Reddy and Venkataraman, 2002a, 2002b) - 1998-99		4300		380	1250
<i>This study</i> - 2005	7952	5957	4700	908	1630
(Zhang et al., 2009) - 2006	4002	3111		344	888
(Klimont et al., 2009) - 2005				1029	2132
<i>This study</i> - 2010	8061	6032	5091	1022	1755
(Lu et al., 2011) - 2010				996	2582
HTAP_v2 (Janssens-Maenhout et al., 2015) - 2010	8280	6230		1019	2530
(Kurokawa et al., 2013) - 2008	6651	4884		713	2286
Europe^c					
<i>This study</i> - 1995	6905	4584	3071	675	1021
(Kupiainen and Klimont, 2007) - 1995				717	1053
(Schaap et al., 2004) - 1995				760	
(Bond et al., 2004) - 1996				678	947
<i>This study</i> - 2000	5579	3843	2668	618	910
(Kupiainen and Klimont, 2007) - 2000				680	996
(Kupiainen and Klimont, 2004) - 2000			2772	672	988
<i>This study</i> - 2010	5008	3471	2393	562	806
<u>TNO-MACCII (Kuenen et al., 2014) - 2009</u>	<u>4694</u>	<u>3199</u>		<u>548</u>	<u>906</u>
HTAP_v2 (Janssens-Maenhout et al., 2015) ^d - 2010	2951	2133		382	638
LRTAP reporting (www.ceip.at) - 2010	4784	3250			
Russian Federation					
<i>This study</i> - 2010	2108	1368	815	170	213
HTAP_v2 (Janssens-Maenhout et al., 2015) - 2010	562	313		60	42
(Huang et al., 2015) - 2010				224	
Russian Federation – European part only					
<i>This study</i> - 2010	1090	734	427	71	122
LRTAP reporting (www.ceip.at) - 2010	569	367			
US					
<i>This study</i> - 2000	1954	1296	949	289	388
(Battye et al., 2002) - 1999				430	
(Reff et al., 2009) - 2000				440	960
(Bond et al., 2013) - 2000				350 ^b	500 ^b

Field Code Changed

Formatted: German (Germany)

Formatted: German (Germany)

Formatted: German (Germany)

Field Code Changed

Region – (Source) – Year	PM10	PM2.5	PM1	BC	OC
<i>This study - 2010</i>	1630	1027	727	201	308
HTAP_v2 (Janssens-Maenhout et al., 2015) - 2010	1973	1640		295	471
<u>(US EPA, 2016)^c - 2011</u>	<u>2847</u>	<u>1909</u>			
<u>(US EPA, 2016)^c - 2014</u>	<u>2830</u>	<u>1875</u>		<u>280</u>	<u>602</u>

^a Based on the information available in the quoted studies, all presented estimates exclude forest fires but include agricultural burning, unless stated otherwise; ^b Excluding agricultural burning; ^c Includes European part of Russian Federation (except HTAP_v2);

^d Excluding any territories of Russian Federation; ^e ~~Including wildfires and prescribed burning;~~ ^f Excluding wildfires and prescribed burning, unpaved roads, and construction dust

References

[AIT: Small and Medium scale Industries in Asia: Energy and Environment. Brick and Ceramic Sectors, Regional Energy Resources Information Center \(RERIC\), Asian Institute of Technology \(AIT\), Pathumthani, Thailand., 2003.](#)

[Alam, S. A.: Use of biomass fuels in the brick-making industries of Sudan: Implications for deforestation and greenhouse gas emission, M. Sc., University of Helsinki, Helsinki, Finland, April., 2006.](#)

[Alves, C., Gonçalves, C., Fernandes, A. P., Tarelho, L. and Pio, C.: Fireplace and woodstove fine particle emissions from combustion of western Mediterranean wood types, Atmospheric Research, 101\(3\), 692–700, doi:10.1016/j.atmosres.2011.04.015, 2011.](#)

[Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Hoglund-Isaksson, L., Kiesewetter, G., Klimont, Z., Schöpp, W., Vellinga, N. and Winiwarter, W.: Adjusted historic emission data, projections, and optimized emission reduction targets for 2030 – A comparison with COM data 2013, IIASA, Laxenburg, Austria. \[online\] Available from: \[http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf\]\(http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf\), 2015.](#)

[Anayochukwu, A. V., Nnene, E. A. and Onyeka, A. E.: Assessment of Environmental Impact of Power Generation in Banking Industry, Journal of Energy, Environment & Carbon Credits, 3\(1\), 13–21, 2013.](#)

[BASIN: Vertical Shaft Brick Kiln - Technology transfer Indian experience - 2, Case Study, Building Advisory Aervice and Information Network \(BASIN\), Eschborn, Germany. \[online\] Available from: <http://www.gtz.de/basin>, 1998.](#)

[BASIN: VSBK-Newsletter 3, Newsletter, Building Advisory Aervice and Information Network \(BASIN\), Eschborn, Germany. \[online\] Available from: <http://www.gtz.de/basin>, 1999.](#)

[Battye, W., Boyer, K. and Pace, T. G.: Methods for improving global inventories of black carbon and organic carbon particulates, EC/R Inc., Chapel Hill, North Carolina, US., 2002.](#)

[Bellprat, O.: Brick Kiln Evaluation Study in the Bajio Region GTO, Mexico., INE, ETH., 2009.](#)

[Boman, C., Nordin, A., Pettersson, E. and Wiinikka, H.: State-of-the-art of small-scale biomass combustion with respect to fine particle emissions – country report from Sweden, Graz, Austria., 2008.](#)

[Boman, C., Pettersson, E., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 1: Pellet Stoves, Energy Fuels, 25\(1\), 307–314, doi:10.1021/ef100774x, 2011.](#)

[Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H. and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, J. Geophys. Res., 109\(D14\), 1–43, doi:10.1029/2003JD003697, 2004.](#)

[Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Bernsten, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G. and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, Journal of Geophysical Research: Atmospheres, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.](#)

Formatted: Bibliography, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Field Code Changed

5 Cao, G., Zhang, X. and Zheng, F.: Inventory of black carbon and organic carbon emissions from China, *Atmospheric Environment*, 40(34), 6516–6527, doi:10.1016/j.atmosenv.2006.05.070, 2006.

Chen, Y., Zhi, G., Feng, Y., Liu, D., Zhang, G., Li, J., Sheng, G. and Fu, J.: Measurements of Black and Organic Carbon Emission Factors for Household Coal Combustion in China: Implication for Emission Reduction, *Environ. Sci. Technol.*, 43(24), 9495–9500, doi:10.1021/es9021766, 2009.

Cheung, K. L., Polidori, A., Ntziachristos, L., Tzankiozis, T., Samaras, Z., Cassee, F. R., Gerlofs, M. and Sioutas, C.: Chemical characteristics and oxidative potential of particulate matter emissions from gasoline, diesel, and biodiesel cars, *Environmental science & technology*, 43(16), 6334–6340, 2009.

10 Co, H. X., Dung, N. T., Le, H. A., An, D. D., van Chinh, K. and Oanh, N. T. K.: Integrated management strategies for brick kiln emission reduction in Vietnam: A case study, *International Journal of Environmental Studies*, 66(1), 113–124, 2009.

Croituru, L. and Sarraf, M.: Benefits and Costs of the Informal Sector: The Case of Brick Kilns in Bangladesh, *Journal of Environmental Protection*, 3, 476–484, doi:http://dx.doi.org/10.4236/jep.2012.36058, 2012.

Dincer, F. and Elbir, T.: Estimating national exhaust emissions from railway vehicles in Turkey, *Science of The Total Environment*, 374(1), 127–134, doi:10.1016/j.scitotenv.2006.12.043, 2007.

15 Dwyer, H., Ayala, A., Zhang, S., Collins, J., Huai, T., Herner, J. and Chau, W.: A study of emissions from a Euro 4 light duty diesel vehicle with the European particulate measurement programme, *Atmospheric Environment*, 44(29), 3469–3476, doi:10.1016/j.atmosenv.2010.06.028, 2010.

20 EC-JRC/PBL: Emission Database for Global Atmospheric Research (EDGAR), European Commission (EC), Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL), [online] Available from: <http://edgar.jrc.ec.europa.eu>, 2010.

EEA: EMEP/EEA air pollutant emission inventory guidebook, Technical report, European Environmental Agency (EEA), Copenhagen, Denmark, [online] Available from: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>, 2013.

EELA: Caracterización de los hornos usados en la industria ladrillera - Proyecto Colombia, Programa Eficiencia Energética en Ladrilleras Artesanales (EELA), 2011.

25 Erbe, S. O.: Technical, economical and organization analysis of informal brick production in Tercera Chica, SLP, Mexico, Master of Science, Universidad Autonoma de San Luis Potosi, San Luis Potosi, Mexico, February., 2011.

FAO: Status and development issues of the brick industry in Asia, Field Document, FAO Regional Wood Energy Development Programme in Asia, Bangkok, Thailand., 1993.

30 Fine, P. M., Cass, G. R. and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the wood stove combustion of prevalent United States tree species, *Environ. Eng. Sci.*, 21(6), 705–721, 2004.

Frey, A. K., Saarnio, K., Lamberg, H., Mylläri, F., Karjalainen, P., Teinilä, K., Carbone, S., Tissari, J., Niemelä, V., Häyrynen, A., Rautiainen, J., Kytömäki, J., Artaxo, P., Virkkula, A., Pirjola, L., Rönkkö, T., Keskinen, J., Jokiniemi, J. and Hillamo, R.: Optical and Chemical Characterization of Aerosols Emitted from Coal, Heavy and Light Fuel Oil, and Small-Scale Wood Combustion, *Environ. Sci. Technol.*, 48(1), 827–836, doi:10.1021/es4028698, 2014.

5 Gaegauf, C. K., Schmid, M. R. and Güntert, P.: Elemental and Organic Carbon in Flue Gas Particles of Various Wood Combustion Systems, Ökozentrum Langenbruck, Langenbruck, Switzerland. [online] Available from: www.oekozentrum.ch, 2005.

Galvis, B., Bergin, M. and Russell, A.: Fuel-based fine particulate and black carbon emission factors from a railyard area in Atlanta, *Journal of the Air & Waste Management Association*, 63(6), 648–658, doi:10.1080/10962247.2013.776507, 2013.

Geller, M. D., Ntziachristos, L., Mamakos, A., Samaras, Z., Schmitz, D. A., Froines, J. R. and Sioutas, C.: Physicochemical and redox characteristics of particulate matter (PM) emitted from gasoline and diesel passenger cars, *Atmospheric Environment*, 40(36), 6988–7004, doi:10.1016/j.atmosenv.2006.06.018, 2006.

10 Gilmore, E. A., Lave, L. B. and Adams, P. J.: The costs, air quality and human health effects of meeting peak electricity demand with installed backup generators, *Environmental Science & Technology*, 40(22), 6887–6893, 2006.

Gonçalves, C., Alves, C., Evtuyugina, M., Mirante, F., Pio, C., Caseiro, A., Schmidl, C., Bauer, H. and Carvalho, F.: Characterisation of PM10 emissions from woodstove combustion of common woods grown in Portugal, *Atmospheric Environment*, 44(35), 4474–4480, doi:10.1016/j.atmosenv.2010.07.026, 2010.

15 Gonçalves, C., Evtuyugina, M., Alves, C., Monteiro, C., Pio, C. and Tomé, M.: Organic particulate emissions from field burning of garden and agriculture residues, *Atmospheric Research*, 101(3), 666–680, doi:10.1016/j.atmosres.2011.04.017, 2011.

Graham, L.: Chemical characterization of emissions from advanced technology light-duty vehicles, *Atmospheric Environment*, 39(13), 2385–2398, doi:10.1016/j.atmosenv.2004.10.049, 2005.

20 Grieshop, A. P., Boland, D., Reynolds, C. C. O., Gouge, B., Apte, J. S., Rogak, S. N. and Kandlikar, M.: Modeling air pollutant emissions from Indian auto-rickshaws: Model development and implications for fleet emission rate estimates, *Atmospheric Environment*, 50, 148–156, doi:10.1016/j.atmosenv.2011.12.046, 2012.

Guan, D., Su, X., Zhang, Q., Peters, G. P., Liu, Z., Lei, Y. and He, K.: The socioeconomic drivers of China's primary PM 2.5 emissions, *Environmental Research Letters*, 9(2), 024010, 2014.

25 Guttikunda, S. K., Begum, B. A. and Wadud, Z.: Particulate pollution from brick kiln clusters in the Greater Dhaka region, Bangladesh, *Air Quality, Atmosphere & Health*, 6(2), 357–365, doi:10.1007/s11869-012-0187-2, 2013.

Heierli, U. and Maithe, S.: Brick by Brick: The Herculean Task of Cleaning of the Asian Brick Industry, Swiss Agency for Development and Cooperation (SDC), Bern., 2008.

Heierli, U., Jha, H. B. and Gurung, Y.: VSBK & CESEF Nepal; External Review, 2007.

30 Herner, J. D., Hu, S., Robertson, W. H., Huai, T., Collins, J. F., Dwyer, H. and Ayala, A.: Effect of Advanced Aftertreatment for PM and NOx Control on Heavy-Duty Diesel Truck Emissions, *Environ. Sci. Technol.*, 43(15), 5928–5933, doi:10.1021/es9008294, 2009.

Huang, K., Fu, J. S., Prikhodko, V. Y., Storey, J. M., Romanov, A., Hodson, E. L., Cresko, J., Morozova, I., Ignatieva, Y. and Cabaniss, J.: Russian anthropogenic black carbon: Emission reconstruction and Arctic black carbon simulation, *Journal of Geophysical Research: Atmospheres*, 120(21), 11,306–11,333, doi:10.1002/2015JD023358, 2015.

Jaffe, D. A., Hof, G., Malashanka, S., Putz, J., Thayer, J., Fry, J. L., Ayres, B. and Pierce, J. R.: Diesel particulate matter emission factors and air quality implications from in-service rail in Washington State, USA, *Atmospheric Pollution Research*, 5(2), 344–351, doi:10.5094/APR.2014.040, 2014.

Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B. and Li, M.: HTAP v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmospheric Chemistry and Physics*, 15(19), 11411–11432, doi:10.5194/acp-15-11411-2015, 2015.

Johansson, L. S., Leckner, B., Gustavsson, L., Cooper, D., Tullin, C. and Potter, A.: Emission characteristics of modern and old-type residential boilers fired with wood logs and wood pellets, *Atmospheric Pollution*, 38, 4183–4195, 2004.

Johnson, G. R., Javaratne, E. R., Lau, J., Thomas, V., Juwono, A. M., Kitchen, B. and Morawska, L.: Remote measurement of diesel locomotive emission factors and particle size distributions, *Atmospheric Environment*, 81, 148–157, doi:10.1016/j.atmosenv.2013.09.019, 2013.

Jordan, T. B. and Seen, A. J.: Effect of airflow setting on the organic composition of woodheater emissions, *Environ. Sci. & Technol.*, 39(10), 3601–3610, 2005.

Just, B., Rogak, S. and Kandlikar, M.: Characterization of Ultrafine Particulate Matter from Traditional and Improved Biomass Cookstoves, *Environ. Sci. Technol.*, 47(7), 3506–3512, doi:10.1021/es304351p, 2013.

Karvosenoja, N.: Emission scenario model for regional air pollution, Monographs of the Boreal Environment Research, 32 [online] Available from: www.environment.fi/publications, 2008.

Klimont, Z., Cofala, J., Bertok, I., Amann, M., Heyes, C. and Gyarfas, F.: Modelling Particulate Emissions in Europe. A Framework to Estimate Reduction Potential and Control Costs, Interim Report, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria., 2002.

Klimont, Z., Cofala, J., Xing, J., Wei, W., Zhang, C., Wang, S., Kejun, J., Bhandari, P., Mathur, R., Purohit, P., Rafaj, P., Chambers, A., Amann, M. and Hao, J.: Projections of SO₂, NO_x, and carbonaceous aerosols emissions in Asia, *Tellus*, 61B(4), 602–617, doi:10.1111/j.1600-0889.2009.00428.x, 2009.

Kondo, Y., Oshima, N., Kajino, M., Mikami, R., Moteki, N., Takegawa, N., Verma, R. L., Kajii, Y., Kato, S. and Takami, A.: Emissions of black carbon in East Asia estimated from observations at a remote site in the East China Sea, *Journal of Geophysical Research: Atmospheres*, 116(D16), n/a–n/a, doi:10.1029/2011JD015637, 2011.

Krasowsky, T., Daher, N., Sioutas, C. and Ban-Weiss, G.: Measurement of particulate matter emissions from in-use locomotives, *Atmospheric Environment*, 113, 187–196, doi:10.1016/j.atmosenv.2015.04.046, 2015.

Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M. and Denier van der Gon, H. A. C.: TNO-MACC II emission inventory: a multi-year (2003 - 2009) consistent high-resolution European emission inventory for air quality modelling, *Atmospheric Chemistry and Physics*, 14(20), 10963–10976, doi:10.5194/acp-14-10963-2014, 2014.

Kupiainen, K. and Klimont, Z.: Primary Emissions of Submicron and Carbonaceous Particles in Europe and the Potential for their Control, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria., 2004.

Kupiainen, K. and Klimont, Z.: Primary emissions of fine carbonaceous particles in Europe, *Atmospheric Environment*, 41, 2156–2170, doi:10.1016/j.atmosenv.2006.10.066, 2007.

5 [Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K. and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia \(REAS\) version 2, Atmospheric Chemistry and Physics, 13\(21\), 11019–11058, doi:10.5194/acp-13-11019-2013, 2013.](#)

[Lamberg, H., Sippula, O., Tissari, J. and Jokiniemi, J.: Effects of Air Staging and Load on Fine-Particle and Gaseous Emissions from a Small-Scale Pellet Boiler, Energy Fuels, 25\(11\), 4952–4960, doi:10.1021/ef2010578, 2011a.](#)

[Lamberg, H., Nuutinen, K., Tissari, J., Ruusunen, J., Yli-Pirilla, P., Sippula, O., Tapanainen, M., Jalava, P., Makkonen, U. and Teinilla, K.: Physicochemical Characterization of Fine Particles from Small-Scale Wood Combustion, Atmospheric Environment, 45, 7635–7643, doi:10.1016/j.atmosenv.2011.02.072, 2011b.](#)

10 [Li, Q., Li, X., Jiang, J., Duan, L., Ge, S., Zhang, Q., Deng, J., Wang, S. and Hao, J.: Semi-coke briquettes: towards reducing emissions of primary PM_{2.5}, particulate carbon, and carbon monoxide from household coal combustion in China, Scientific Reports, 6, 19306, 2016.](#)

[Li, X. H., Wang, S. X., Duan, L., Hao, J. M. and Nie, Y. F.: Carbonaceous aerosol emissions from household biofuel combustion in China., Environ. Sci. Technol., 43, 6076–6081, 2009.](#)

15 [Liu, H., He, K., Lents, J. M., Wang, Q. and Tolvet, S.: Characteristics of Diesel Truck Emission in China Based on Portable Emissions Measurement Systems, Environmental Science & Technology, 43\(24\), 9507–9511, doi:10.1021/es902044x, 2009.](#)

[Louis, C., Liu, Y., Tassel, P., Perret, P., Chaumont, A. and André, M.: PAH, BTEX, carbonyl compound, black-carbon, NO₂ and ultrafine particle dynamometer bench emissions for Euro 4 and Euro 5 diesel and gasoline passenger cars, Atmospheric Environment, 141, 80–95, doi:10.1016/j.atmosenv.2016.06.055, 2016.](#)

20 [Lu, Z., Zhang, Q. and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, Atmos. Chem. Phys., 11\(18\), 9839–9864, 2011.](#)

[Maithe, S.: Factsheets about Brick Kilns in South and South-East Asia, \[online\] Available from: <http://www.ccacoalition.org/en/resources/factsheets-about-brick-kilns-south-and-south-east-asia>, 2014.](#)

25 [Maithe, S., Lalchandani, D., Malhotra, G., Bhanware, P., Uma, R., Ragavan, S., Athalye, V., Bindiya, K., Reddy, S., Bond, T. C., Weyant, C., Baum, E., Kim Thoa, V. T., Thu Phuong, N. and Kim Thanh, T.: Brick Kilns Performance Assessment: A Roadmap for Cleaner Brick Production in India, Shakti Sustainable Energy Foundation and Climate Works Foundation, \[online\] Available from: <http://www.catf.us/resources/publications/view/161>, 2012.](#)

[McDonald, J. D., Zielinska, B., Fujita, E. M., Sagebiel, J. C., Chow, J. C. and Watson, J. G.: Fine Particle and Gaseous Emission Rates from Residential Wood Combustion., Environ. Sci. & Technol., 34, 2080–2091, 2000.](#)

30 [Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X. and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, Atmos. Chem. Phys., 7, 4419–4444, 2007.](#)

[Pettersson, E., Boman, C., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 2: Wood Stove, Energy Fuels, 25\(1\), 315–323, doi:10.1021/ef1007787, 2011.](#)

35 [PRAL: Case study, Behind the bricks: An experience in the integrated management of the informal sector, PRAL \(Regional Clean Air Program\), Lima, Peru., 2012.](#)

Reddy, M. S. and Venkataraman, C.: Inventory of aerosol and sulphur dioxide emissions from India. Part II - biomass combustion, *Atmospheric Environment*, 36(4), 699–712. doi:[http://dx.doi.org/10.1016/S1352-2310\(01\)00464-2](http://dx.doi.org/10.1016/S1352-2310(01)00464-2), 2002a.

Reddy, S. M. and Venkataraman, C.: Inventory of aerosol and sulphur dioxide emissions from India. Part I - Fossil fuel combustion, *Atmospheric Environment*, 36, 677–697, 2002b.

Reff, A., Bhave, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D. and Houyoux, M.: Emissions Inventory of PM_{2.5} Trace Elements across the United States, *Environ. Sci. Technol.*, 43(15), 5790–5796. doi:10.1021/es802930x, 2009.

Roden, C. A., Bond, T. C., Conway, S., Osorto Pinel, A. B., MacCarty, N. and Still, D.: Laboratory and field investigations of particulate and carbon monoxide emissions from traditional and improved cookstoves, *Atmospheric Environment*, 43(6), 1170–1181. doi:10.1016/j.atmosenv.2008.05.041, 2009.

Rozemuller, B.: *An Overview of the Brick and Tile Manufacturing Industry in North West Cambodia*, Center for Advanced Studies, UNDP/CARERE, Phnom Penh, Kingdom of Cambodia., 1999.

Samuel Hall Consulting: *A Rapid Assessment of Bonded Labour in Brick Kilns in Afghanistan*, International Labour Organisation (ILO), Kabul, Afghanistan., 2011.

Schaap, M., Van Der Gon, H. A. C. D., Dentener, F. J., Visschedijk, A. J. H., Van Loon, M., ten Brink, H. M., Putaud, J.-P., Guillaume, B., Liousse, C. and Builtjes, P. J. H.: Anthropogenic black carbon and fine aerosol distribution over Europe, *Journal of Geophysical Research: Atmospheres*, 109(D18), n/a–n/a. doi:10.1029/2003JD004330, 2004.

Schmidl, C., Luisser, M., Padouvas, E., Lasselsberger, L., Rzaca, M., Cruz, C. -S., Handler, M., Peng, G., Bauer, H. and Puxbaum, H.: Particulate and gaseous emissions from manually and automatically fired small scale combustion systems., *Atmospheric Environment*, 45, 7443–7454, 2011.

Scott, G.: *Baseline Study for Clamp Kilns in South Africa and Possible Regulatory Approaches*, 2013.

Shah, S. D., Cocker III, D. R., Johnson, K. C., Lee, J. M., Soriano, B. L. and Miller, J. W.: Reduction of particulate matter emissions from diesel backup generators equipped with four different exhaust after treatment devices, *Environmental Science & Technology*, 41(14), 5070–5076. doi:10.1021/es0614161, 2007.

Sippula, O., Hytönen, K., Tissari, J., Raunemaa, T. and Jokiniemi, J.: Effect of Wood Fuel on the Emissions from a Top-Feed Pellet Stove, *Energy Fuels*, 21(2), 1151–1160. doi:10.1021/ef060286e, 2007.

Spezzano, P., Picini, P., Cataldi, D., Messale, F., Manni, C. and Santino, D.: Particle-phase Polycyclic Aromatic Hydrocarbon Emissions from Non-catalysed, In-use Four-stroke Scooters, *Environmental Monitoring and Assessment*, 133(1), 105–117. doi:10.1007/s10661-006-9564-3, 2007.

Spezzano, P., Picini, P., Cataldi, D., Messale, F. and Manni, C.: Particle- and gas-phase emissions of polycyclic aromatic hydrocarbons from two-stroke, 50-cm³ mopeds, *Atmospheric Environment*, 42(18), 4332–4344. doi:10.1016/j.atmosenv.2008.01.008, 2008.

Stratus Consulting: *CCAC initiative to mitigate black carbon and other pollutants from brick production: Regional Assessment. Black Carbon Mitigation in Brick Production: A Summary for Brazil, Chile, Colombia, Mexico, Nigeria, and Peru*, Climate and Clean Air Coalition (CCAC), 2014.

Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J. H. and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research*, 108(D21), 1–23, doi:10.1029/2002JD003093, 2003.

Subrahmanya, M. H. B.: Labour productivity, energy intensity and economic performance in small enterprises: A study of brick enterprises cluster in India, *Energy Conversion and Management*, 47, 763–777, doi:10.1016/j.enconman.2005.05.021, 2006.

Subramanian, R., Winijkul, E., Bond, T. C., Thiansathit, W., Oanh, N. T. K., Paw-armart, I. and Duleep, K. G.: Climate-Relevant Properties of Diesel Particulate Emissions: Results from a Piggyback Study in Bangkok, Thailand, *Environmental Science & Technology*, 43(11), 4213–4218, doi:10.1021/es8032296, 2009.

SwissContact: Report on comparative portfolio of technologies and practices in Asia and LAC; Assessment of Gaps., CCAC: Brick Production Initiative., 2014a.

SwissContact: Report on identified past work on effective policies., CCAC: Brick Production Initiative., 2014b.

SwissContact: Survey of the Status of the Informal Clay Brick Making Sector: Eastern Cape, South Africa, 2014, 2014c.

Tang, N. W., Apte, J. S., Martien, P. T. and Kirchstetter, T. W.: Measurement of black carbon emissions from in-use diesel-electric passenger locomotives in California, *Atmospheric Environment*, 115, 295–303, doi:10.1016/j.atmosenv.2015.05.001, 2015.

Tissari, J., Sippula, O., Kouki, J., Vuorio, K. and Jokiniemi, J.: Fine Particle and Gas Emissions from the Combustion of Agricultural Fuels Fired in a 20 kW Burner, *Energy Fuels*, 22(3), 2033–2042, doi:10.1021/ef700766y, 2008.

Tsai, J. H., Chen, S. J., Huang, K. L., Lin, Y. C., Lee, W. J., Lin, C. C. and Lin, W. Y.: PM, carbon, and PAH emissions from a diesel generator fuelled with soy-biodiesel blends, *Journal of Hazardous Materials*, 179(1–3), 237–243, doi:10.1016/j.jhazmat.2010.02.085, 2010.

Uma, R., Kandpal, T. C. and Kishore, V. V. N.: Emission characteristics of an electricity generation system in diesel alone and dual fuel modes, *Biomass and Bioenergy*, 27(2), 195–203, doi:10.1016/j.biombioe.2004.01.003, 2004.

UNEP/WMO: Integrated Assessment of Black Carbon and Tropospheric Ozone, Nairobi, Kenya. [online] Available from: www.unep.org/dewa/Portals/67/pdf/BlackCarbon_report.pdf, 2011.

US EPA: National Emissions Inventory (NEI), [online] Available from: <https://www.epa.gov/air-emissions-inventories/national-emissions-inventory-nei> (Accessed 2 March 2017), 2016.

Verma, S. and Uppal, J.: Use of biomass in brick kilns, *Ministry of New and Renewable Energy, Government of India*, 6(4), 24–27, 2013.

Wang, A., Ge, Y., Tan, J., Fu, M., Shah, A. N., Ding, Y., Zhao, H. and Liang, B.: On-road pollutant emission and fuel consumption characteristics of buses in Beijing, *J. Env. Sci.*, 23(3), 419–426, doi:10.1016/S1001-0742(10)60426-3, 2011.

Wang, S.-X., Zhao, X.-J., Li, X.-H., Wei, W. and Hao, J.-M.: Emission characteristics of fine particles from grate firing boilers, *Environmental Science*, 30(4), 963–968, 2009.

Watson, J. G., Miller, W., Moosmuller, H., Kuhns, H. D., Chow, J. C., Zhu, D. and Cocker, D. R.: Characterization of Off-Road Diesel Emissions of Criteria Pollutants, CY2006 Annual Report, WP-1336, Desert Research Institute, Reno, NV., 2006.

5 Watson, J. G., Miller, J. W., Moosmüller, H., Kuhns, H. D., Chow, J. C., Chang, O., Nussbaum, N., Mazzoleni, C., Zhu, D., Barber, P. W., Kemme, M. R. and Cocker, D. R.: Characterization of off-road diesel emissions of criteria pollutants, Desert Research Institute, Reno, NV. [online] Available from: <https://www.dri.edu/images/stories/editors/cafeditor/Watsonetal2008SERDPFinalReportDieselWP1366.pdf>, 2008.

Weyant, C., Athalye, V., Ragavan, S., Rajarathnam, U., Lalchandani, D., Maithel, S., Baum, E. and Bond, T. C.: Emissions from South Asian Brick Production, *Environ. Sci. Technol.*, 48(11), 6477–6483, doi:10.1021/es500186g, 2014.

10 World Bank: Introducing energy-efficient clean technologies in the brick sector of Bangladesh, World Bank, Washington DC, USA. [online] Available from: <http://documents.worldbank.org/curated/en/2011/06/15492357/introducing-energy-efficient-clean-technologies-brick-sector-bangladesh>, 2011.

Yang, H.-H., Hsieh, L.-T., Liu, H.-C. and Mi, H.-H.: Polycyclic aromatic hydrocarbon emissions from motorcycles, *Atmospheric Environment*, 39(1), 17–25, doi:10.1016/j.atmosenv.2004.09.054, 2005.

15 Yanowitz, J., McCormick, R. L. and Graboski, M. S.: In-Use Emissions from Heavy-Duty Diesel Vehicles, *Environmental Science & Technology*, 34(5), 729–740, doi:10.1021/es990903w, 2000.

Zhang, Q., Klimont, Z., Streets, D., Huo, H. and He, K.: An anthropogenic PM emission model for China and emission inventory for the year 2001, *Progress in Natural Science*, 16(2), 223–231 (in Chinese), 2006.

20 Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T. and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, 2009.

Zhang, Y., Schauer, J. J., Zhang, Y., Zeng, L., Wei, Y., Liu, Y. and Shao, M.: Characteristics of Particulate Carbon Emissions from Real-World Chinese Coal Combustion, *Environ. Sci. Technol.*, 42(14), 5068–5073, doi:10.1021/es702257g, 2008.

25 Zhang, Z.: Energy efficiency and environmental pollution of brickmaking in China, *Energy*, 22(1), 33–42, 1997.

Zhi, G., Peng, C., Chen, Y., Liu, D., Sheng, G. and Fu, J.: Deployment of Coal Briquettes and Improved Stoves: Possibly an Option for both Environment and Climate, *Environ. Sci. Technol.*, 43(15), 5586–5591, doi:10.1021/es802955d, 2009.

30 Zhu, D., Nussbaum, N. J., Kuhns, H. D., Chang, M. C. O., Sodeman, D., Uppapalli, S., Moosmüller, H., Chow, J. C. and Watson, J. G.: In-Plume Emission Test Stand 2: Emission Factors for 10- to 100-kW U.S. Military Generators, *Journal of the Air & Waste Management Association*, 59(12), 1446–1457, doi:10.3155/1047-3289.59.12.1446, 2009.

AIT: Small and Medium scale Industries in Asia: Energy and Environment. Brick and Ceramic Sectors, Regional Energy Resources Information Center (RERIC), Asian Institute of Technology (AIT), Pathumthani, Thailand., 2003.

Alam, S. A.: Use of biomass fuels in the brick-making industries of Sudan: Implications for deforestation and greenhouse gas emission, M. Sc., University of Helsinki, Helsinki, Finland, April., 2006.

Alves, C., Gonçalves, C., Fernandes, A. P., Tarelho, L. and Pio, C.: Fireplace and woodstove fine particle emissions from combustion of western Mediterranean wood types, *Atmospheric Research*, 101(3), 692–700, doi:10.1016/j.atmosres.2011.04.015, 2011.

5 Amann, M., Bertok, I., Borken-Kleefeld, J., Cofala, J., Heyes, C., Hoglund-Isaksson, L., Kiesewetter, G., Klimont, Z., Schöpp, W., Vellinga, N. and Winiwarter, W.: Adjusted historic emission data, projections, and optimized emission reduction targets for 2030—A comparison with COM data 2013, HASA, Laxenburg, Austria. [online] Available from: http://ec.europa.eu/environment/air/pdf/review/TSAP_16a.pdf, 2015.

Anayochukwu, A. V., Naene, E. A. and Onyeka, A. E.: Assessment of Environmental Impact of Power Generation in Banking Industry, *Journal of Energy, Environment & Carbon Credits*, 3(1), 13–21, 2013.

10 BASIN: Vertical Shaft Brick Kiln—Technology transfer Indian experience—2, Case Study, Building Advisory Service and Information Network (BASIN), Eschborn, Germany. [online] Available from: <http://www.gtz.de/basin>, 1998.

BASIN: VSBK Newsletter 3, Newsletter, Building Advisory Service and Information Network (BASIN), Eschborn, Germany. [online] Available from: <http://www.gtz.de/basin>, 1999.

15 Battye, W., Boyer, K. and Pace, T. G.: Methods for improving global inventories of black carbon and organic carbon particulates, EC/R Inc., Chapel Hill, North Carolina, US., 2002.

Bellprat, O.: Brick Kiln Evaluation Study in the Bajío Region GTO, Mexico., INE, ETH., 2009.

Boman, C., Nordin, A., Pettersson, E. and Wiinikka, H.: State-of-the-art of small-scale biomass combustion with respect to fine particle emissions—country report from Sweden, Graz, Austria., 2008.

20 Boman, C., Pettersson, E., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 1: Pellet Stoves, *Energy Fuels*, 25(1), 307–314, doi:10.1021/ef100774x, 2011.

Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H. and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109(D14), 1–43, doi:10.1029/2003JD003697, 2004.

25 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Bernsten, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schult, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G. and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.

30 Cao, G., Zhang, X. and Zheng, F.: Inventory of black carbon and organic carbon emissions from China, *Atmospheric Environment*, 40(34), 6516–6527, doi:10.1016/j.atmosenv.2006.05.070, 2006.

35 Chen, Y., Zhi, G., Feng, Y., Liu, D., Zhang, G., Li, J., Sheng, G. and Fu, J.: Measurements of Black and Organic Carbon Emission Factors for Household Coal Combustion in China: Implication for Emission Reduction, *Environ. Sci. Technol.*, 43(24), 9495–9500, doi:10.1021/es9021766, 2009.

- Cheung, K. L., Polidori, A., Ntziachristos, L., Tzankiozis, T., Samaras, Z., Cassee, F. R., Gerlofs, M. and Sioutas, C.: Chemical characteristics and oxidative potential of particulate matter emissions from gasoline, diesel, and biodiesel cars, *Environmental science & technology*, 43(16), 6334–6340, 2009.
- 5 Co, H. X., Dung, N. T., Le, H. A., An, D. D., van Chinh, K. and Oanh, N. T. K.: Integrated management strategies for brick kiln emission reduction in Vietnam: A case study, *International Journal of Environmental Studies*, 66(1), 113–124, 2009.
- Croitoru, L. and Sarraf, M.: Benefits and Costs of the Informal Sector: The Case of Brick Kilns in Bangladesh, *Journal of Environmental Protection*, 3, 476–484, doi:<http://dx.doi.org/10.4236/jep.2012.36058>, 2012.
- Dincer, F. and Elbir, T.: Estimating national exhaust emissions from railway vehicles in Turkey, *Science of The Total Environment*, 374(1), 127–134, doi:[10.1016/j.scitotenv.2006.12.043](https://doi.org/10.1016/j.scitotenv.2006.12.043), 2007.
- 10 Dwyer, H., Ayala, A., Zhang, S., Collins, J., Huai, T., Herner, J. and Chau, W.: A study of emissions from a Euro 4 light duty diesel vehicle with the European particulate measurement programme, *Atmospheric Environment*, 44(29), 3469–3476, doi:[10.1016/j.atmosenv.2010.06.028](https://doi.org/10.1016/j.atmosenv.2010.06.028), 2010.
- EC JRC/PBL: Emission Database for Global Atmospheric Research (EDGAR), European Commission (EC), Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL). [online] Available from: <http://edgar.jrc.ec.europa.eu>, 2010.
- 15 EEA: EMEP/EEA air pollutant emission inventory guidebook, Technical report, European Environmental Agency (EEA), Copenhagen, Denmark. [online] Available from: <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>, 2013.
- EELA: Caracterización de los hornos usados en la industria ladrillera — Proyecto Colombia, Programa Eficiencia Energética en Ladrillos Artesanales (EELA), 2011.
- 20 Erbe, S. O.: Technical, economical and organization analysis of informal brick production in Tercera Chica, SLP, Mexico; Master of Science, Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico, February, 2011.
- FAO: Status and development issues of the brick industry in Asia, Field Document, FAO Regional Wood Energy Development Programme in Asia, Bangkok, Thailand., 1993.
- 25 Fine, P. M., Cass, G. R. and Simoneit, B. R. T.: Chemical characterization of fine particle emissions from the wood stove combustion of prevalent United States tree species, *Environ. Eng. Sci.*, 21(6), 705–721, 2004.
- Frey, A. K., Saarnio, K., Lamberg, H., Mylläri, F., Karjalainen, P., Teinilä, K., Carbone, S., Tissari, J., Niemelä, V., Häyrynen, A., Rautiainen, J., Kytömäki, J., Artaxo, P., Virkkula, A., Pirjola, L., Rönkkö, T., Keskinen, J., Jokiniemi, J. and Hillamo, R.: Optical and Chemical Characterization of Aerosols Emitted from Coal, Heavy and Light Fuel Oil, and Small-Scale Wood Combustion, *Environ. Sci. Technol.*, 48(1), 827–836, doi:[10.1021/es4028698](https://doi.org/10.1021/es4028698), 2014.
- 30 Gaegauf, C. K., Schmid, M. R. and Güntert, P.: Elemental and Organic Carbon in Flue Gas Particles of Various Wood Combustion Systems, Ökozentrum Langenbruck, Langenbruck, Switzerland. [online] Available from: www.oekozentrum.ch, 2005.
- Galvis, B., Bergin, M. and Russell, A.: Fuel-based fine particulate and black carbon emission factors from a railyard area in Atlanta, *Journal of the Air & Waste Management Association*, 63(6), 648–658, doi:[10.1080/10962247.2013.776507](https://doi.org/10.1080/10962247.2013.776507), 2013.

- Geller, M. D., Ntziachristos, L., Mamakos, A., Samaras, Z., Schmitz, D. A., Froines, J. R. and Sioutas, C.: Physicochemical and redox characteristics of particulate matter (PM) emitted from gasoline and diesel passenger cars, *Atmospheric Environment*, 40(36), 6988–7004, doi:10.1016/j.atmosenv.2006.06.018, 2006.
- 5 Gilmore, E. A., Lave, L. B. and Adams, P. J.: The costs, air quality and human health effects of meeting peak electricity demand with installed backup generators, *Environmental Science & Technology*, 40(22), 6887–6893, 2006.
- Gonçalves, C., Alves, C., Evtuygina, M., Mirante, F., Pio, C., Caseiro, A., Schmidl, C., Bauer, H. and Carvalho, F.: Characterisation of PM10 emissions from woodstove combustion of common woods grown in Portugal, *Atmospheric Environment*, 44(35), 4474–4480, doi:10.1016/j.atmosenv.2010.07.026, 2010.
- 10 Gonçalves, C., Evtuygina, M., Alves, C., Monteiro, C., Pio, C. and Tomé, M.: Organic particulate emissions from field burning of garden and agriculture residues, *Atmospheric Research*, 101(3), 666–680, doi:10.1016/j.atmosres.2011.04.017, 2011.
- Graham, L.: Chemical characterization of emissions from advanced technology light duty vehicles, *Atmospheric Environment*, 39(13), 2385–2398, doi:10.1016/j.atmosenv.2004.10.049, 2005.
- 15 Grieshop, A. P., Boland, D., Reynolds, C. C. O., Gouge, B., Apte, J. S., Rogak, S. N. and Kandlikar, M.: Modeling air pollutant emissions from Indian auto-rickshaws: Model development and implications for fleet emission rate estimates, *Atmospheric Environment*, 50, 148–156, doi:10.1016/j.atmosenv.2011.12.046, 2012.
- Guan, D., Su, X., Zhang, Q., Peters, G. P., Liu, Z., Lei, Y. and He, K.: The socioeconomic drivers of China's primary PM 2.5 emissions, *Environmental Research Letters*, 9(2), 24010, 2014.
- 20 Guttikunda, S. K., Begum, B. A. and Wadud, Z.: Particulate pollution from brick kiln clusters in the Greater Dhaka region, Bangladesh, *Air Quality, Atmosphere & Health*, 6(2), 357–365, doi:10.1007/s11869-012-0187-2, 2013.
- Heierli, U. and Maithel, S.: Brick by Brick: The Herculean Task of Cleaning of the Asian Brick Industry, Swiss Agency for Development and Cooperation (SDC), Bern., 2008.
- Heierli, U., Jha, H. B. and Gurung, Y.: VSBK & CESEF Nepal; External Review, 2007.
- 25 Hemer, J. D., Hu, S., Robertson, W. H., Huai, T., Collins, J. F., Dwyer, H. and Ayala, A.: Effect of Advanced Aftertreatment for PM and NOx Control on Heavy Duty Diesel Truck Emissions, *Environ. Sci. Technol.*, 43(15), 5928–5933, doi:10.1021/es9008294, 2009.
- Huang, K., Fu, J. S., Prikhodko, V. Y., Storey, J. M., Romanov, A., Hodson, E. L., Cresko, J., Morozova, I., Ignatieva, Y. and Cabaniss, J.: Russian anthropogenic black carbon: Emission reconstruction and Arctic black carbon simulation, *Journal of Geophysical Research: Atmospheres*, 120(21), 11,306–11,333, doi:10.1002/2015JD023358, 2015.
- 30 Jaffe, D. A., Hof, G., Malashanka, S., Putz, J., Thayer, J., Fry, J. L., Ayres, B. and Pierce, J. R.: Diesel particulate matter emission factors and air quality implications from in-service rail in Washington State, USA, *Atmospheric Pollution Research*, 5(2), 344–351, doi:10.5094/APR.2014.040, 2014.
- 35 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B. and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmospheric Chemistry and Physics*, 15(19), 11411–11432, doi:10.5194/acp-15-11411-2015, 2015.

Johansson, L. S., Leckner, B., Gustavsson, L., Cooper, D., Tullin, C. and Potter, A.: Emission characteristics of modern and old-type residential boilers fired with wood logs and wood pellets, *Atmospheric Pollution*, 38, 4183–4195, 2004.

Johnson, G. R., Jayaratne, E. R., Lau, J., Thomas, V., Juwono, A. M., Kitchen, B. and Morawska, L.: Remote measurement of diesel locomotive emission factors and particle size distributions, *Atmospheric Environment*, 81, 148–157, doi:10.1016/j.atmosenv.2013.09.019, 2013.

Jordan, T. B. and Seen, A. J.: Effect of airflow setting on the organic composition of woodheater emissions, *Environ. Sci. & Technol.*, 39(10), 3601–3610, 2005.

Just, B., Rogak, S. and Kandlikar, M.: Characterization of Ultrafine Particulate Matter from Traditional and Improved Biomass Cookstoves, *Environ. Sci. Technol.*, 47(7), 3506–3512, doi:10.1021/es304351p, 2013.

Karvosenoja, N.: Emission scenario model for regional air pollution, *Monographs of the Boreal Environment Research*, 32 [online] Available from: www.enviroment.fi/publications, 2008.

Klimont, Z., Cofala, J., Bertok, I., Amann, M., Heyes, C. and Gyarfas, F.: Modelling Particulate Emissions in Europe. A Framework to Estimate Reduction Potential and Control Costs, *Interim Report, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria*, 2002.

Klimont, Z., Cofala, J., Xing, J., Wei, W., Zhang, C., Wang, S., Kejun, J., Bhandari, P., Mathur, R., Purohit, P., Rafaj, P., Chambers, A., Amann, M. and Hao, J.: Projections of SO₂, NO_x, and carbonaceous aerosols emissions in Asia, *Tellus*, 61B(4), 602–617, doi:10.1111/j.1600-0889.2009.00428.x, 2009.

Kondo, Y., Oshima, N., Kajino, M., Mikami, R., Moteki, N., Takegawa, N., Verma, R. L., Kajii, Y., Kato, S. and Takami, A.: Emissions of black carbon in East Asia estimated from observations at a remote site in the East China Sea, *Journal of Geophysical Research: Atmospheres*, 116(D16), n/a–n/a, doi:10.1029/2011JD015637, 2011.

Krasowsky, T., Daher, N., Sioutas, C. and Ban-Weiss, G.: Measurement of particulate matter emissions from in-use locomotives, *Atmospheric Environment*, 113, 187–196, doi:10.1016/j.atmosenv.2015.04.046, 2015.

Kupiainen, K. and Klimont, Z.: Primary Emissions of Submicron and Carbonaceous Particles in Europe and the Potential for their Control, *International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria*, 2004.

Kupiainen, K. and Klimont, Z.: Primary emissions of fine carbonaceous particles in Europe, *Atmospheric Environment*, 41, 2156–2170, doi:10.1016/j.atmosenv.2006.10.066, 2007.

Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens Maenhout, G., Fukui, T., Kawashima, K. and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia (REAS)-version 2, *Atmospheric Chemistry and Physics*, 13(21), 11019–11058, doi:10.5194/acp-13-11019-2013, 2013.

Lamberg, H., Sippula, O., Tissari, J. and Jokiniemi, J.: Effects of Air Staging and Load on Fine Particle and Gaseous Emissions from a Small Scale Pellet Boiler, *Energy Fuels*, 25(11), 4952–4960, doi:10.1021/ef2010578, 2011a.

Lamberg, H., Nuutinen, K., Tissari, J., Ruusunen, J., Yli-Pirilla, P., Sippula, O., Tapanainen, M., Jalava, P., Makkonen, U. and Teinilla, K.: Physicochemical Characterization of Fine Particles from Small-Scale Wood Combustion, *Atmospheric Environment*, 45, 7635–7643, doi:10.1016/j.atmosenv.2011.02.072, 2011b.

Li, Q., Li, X., Jiang, J., Duan, L., Ge, S., Zhang, Q., Deng, J., Wang, S., and Hao, J.: Semi coke briquettes: towards reducing emissions of primary PM_{2.5}, particulate carbon, and carbon monoxide from household coal combustion in China, *Scientific Reports*, 6, 19306, 2016.

Li, X. H., Wang, S. X., Duan, L., Hao, J. M. and Nie, Y. F.: Carbonaceous aerosol emissions from household biofuel combustion in China., *Environ. Sci. Technol.*, 43, 6076–6081, 2009.

Liu, H., He, K., Lents, J. M., Wang, Q. and Tolvett, S.: Characteristics of Diesel Truck Emission in China Based on Portable Emissions Measurement Systems, *Environmental Science & Technology*, 43(24), 9507–9511, doi:10.1021/es902044x, 2009.

Louis, C., Liu, Y., Tassel, P., Perret, P., Chaumond, A. and André, M.: PAH, BTEX, carbonyl compound, black carbon, NO₂ and ultrafine particle dynamometer bench emissions for Euro 4 and Euro 5 diesel and gasoline passenger cars, *Atmospheric Environment*, 141, 80–95, doi:10.1016/j.atmosenv.2016.06.055, 2016.

Lu, Z., Zhang, Q. and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, *Atmos. Chem. Phys.*, 11(18), 9839–9864, 2011.

Maithel, S.: Factsheets about Brick Kilns in South and South East Asia, [online] Available from: <http://www.ceacoalition.org/en/resources/factsheets-about-brick-kilns-south-and-south-east-asia>, 2014.

Maithel, S., Lalchandani, D., Malhotra, G., Bhanware, P., Uma, R., Ragavan, S., Athalye, V., Bindiya, K., Reddy, S., Bond, T. C., Weyant, C., Baum, E., Kim Thoa, V. T., Thu Phuong, N. and Kim Thanh, T.: Brick Kilns Performance Assessment: A Roadmap for Cleaner Brick Production in India, Shakti Sustainable Energy Foundation and Climate Works Foundation. [online] Available from: <http://www.catf.us/resources/publications/view/161>, 2012.

McDonald, J. D., Zielinska, B., Fujita, E. M., Sagebiel, J. C., Chow, J. C. and Watson, J. G.: Fine Particle and Gaseous Emission Rates from Residential Wood Combustion., *Environ. Sci. & Technol.*, 34, 2080–2091, 2000.

Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X. and Hayasaka, T.: An Asian emission inventory of anthropogenic emission sources for the period 1980–2020, *Atmos. Chem. Phys.*, 7, 4419–4444, 2007.

Pettersson, E., Boman, C., Westerholm, R., Boström, D. and Nordin, A.: Stove Performance and Emission Characteristics in Residential Wood Log and Pellet Combustion, Part 2: Wood Stove, *Energy Fuels*, 25(1), 315–323, doi:10.1021/ef1007787, 2011.

PRAL: Case study. Behind the bricks: An experience in the integrated management of the informal sector, PRAL (Regional Clean Air Program), Lima, Peru., 2012.

Reddy, S. M. and Venkataraman, C.: Inventory of aerosol and sulphur dioxide emissions from India. Part I—Fossil fuel combustion, *Atmospheric Environment*, 36, 677–697, 2002a.

Reddy, S. M. and Venkataraman, C.: Inventory of aerosol and sulphur dioxide emissions from India. Part II—biomass combustion, *Atmospheric Environment*, 36, 699–712, 2002b.

Reff, A., Bhawe, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D. and Houyoux, M.: Emissions Inventory of PM_{2.5} Trace Elements across the United States, *Environ. Sci. Technol.*, 43(15), 5790–5796, doi:10.1021/es802930x, 2009.

Roden, C. A., Bond, T. C., Conway, S., Osorto Pinel, A. B., MacCarty, N. and Still, D.: Laboratory and field investigations of particulate and carbon monoxide emissions from traditional and improved cookstoves, *Atmospheric Environment*, 43(6), 1170–1181, doi:10.1016/j.atmosenv.2008.05.041, 2009.

- Rozemuller, B.: An Overview of the Brick and Tile Manufacturing Industry in North West Cambodia, Center for Advanced Studies, UNDP/CARERE, Phnom Penh, Kingdom of Cambodia., 1999.
- Samuel Hall Consulting: A Rapid Assessment of Bonded Labour in Brick Kilns in Afghanistan, International Labour Organisation (ILO), Kabul, Afghanistan., 2011.
- 5 Schaap, M., Van Der Gon, H. A. C. D., Dentener, F. J., Visschedijk, A. J. H., Van Loon, M., ten Brink, H. M., Putaud, J. P., Guillaume, B., Liousse, C. and Builtjes, P. J. H.: Anthropogenic black carbon and fine aerosol distribution over Europe, *Journal of Geophysical Research: Atmospheres*, 109(D18), n/a–n/a, doi:10.1029/2003JD004330, 2004.
- 10 Schmidl, C., Luisser, M., Padouvas, E., Lasselsberger, L., Rzaea, M., Cruz, C. S., Handler, M., Peng, G., Bauer, H. and Puxbaum, H.: Particulate and gaseous emissions from manually and automatically fired small scale combustion systems, *Atmospheric Environment*, 45, 7443–7454, 2011.
- Scott, G.: Baseline Study for Clamp Kilns in South Africa and Possible Regulatory Approaches, 2013.
- Shah, S. D., Cocker III, D. R., Johnson, K. C., Lee, J. M., Soriano, B. L. and Miller, J. W.: Reduction of particulate matter emissions from diesel backup generators equipped with four different exhaust after treatment devices, *Environmental Science & Technology*, 41(14), 5070–5076, doi:10.1021/es0614161, 2007.
- 15 Sippula, O., Hytönen, K., Tissari, J., Raunemaa, T. and Jokiniemi, J.: Effect of Wood Fuel on the Emissions from a Top-Feed Pellet Stove, *Energy Fuels*, 21(2), 1151–1160, doi:10.1021/ef060286e, 2007.
- Spezzano, P., Picini, P., Cataldi, D., Messale, F., Manni, C. and Santino, D.: Particle phase Polycyclic Aromatic Hydrocarbon Emissions from Non catalysed, In use Four stroke Scooters, *Environmental Monitoring and Assessment*, 133(1), 105–117, doi:10.1007/s10661-006-9564-3, 2007.
- 20 Spezzano, P., Picini, P., Cataldi, D., Messale, F. and Manni, C.: Particle and gas phase emissions of polycyclic aromatic hydrocarbons from two stroke, 50 cm³ mopeds, *Atmospheric Environment*, 42(18), 4332–4344, doi:10.1016/j.atmosenv.2008.01.008, 2008.
- 25 Stratus Consulting: CCAC initiative to mitigate black carbon and other pollutants from brick production: Regional Assessment. Black Carbon Mitigation in Brick Production: A Summary for Brazil, Chile, Colombia, Mexico, Nigeria, and Peru, Climate and Clean Air Coalition (CCAC), 2014.
- Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J. H. and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research*, 108(D21), 1–23, doi:10.1029/2002JD003093, 2003.
- 30 Subrahmanya, M. H. B.: Labour productivity, energy intensity and economic performance in small enterprises: A study of brick enterprises cluster in India, *Energy Conversion and Management*, 47, 763–777, doi:10.1016/j.enconman.2005.05.021, 2006.
- Subramanian, R., Winijkul, E., Bond, T. C., Thiansathit, W., Oanh, N. T. K., Pawarmart, I. and Duleep, K. G.: Climate-Relevant Properties of Diesel Particulate Emissions: Results from a Piggyback Study in Bangkok, Thailand, *Environmental Science & Technology*, 43(11), 4213–4218, doi:10.1021/es8032296, 2009.
- 35 SwissContact: Report on comparative portfolio of technologies and practices in Asia and LAC; Assessment of Gaps; CCAC; Brick Production Initiative., 2014a.

SwissContact: Report on identified past work on effective policies., CCAC; Brick Production Initiative., 2014b.

SwissContact: Survey of the Status of the Informal Clay Brick Making Sector; Eastern Cape, South Africa, 2014, 2014c.

5 Tang, N. W., Apte, J. S., Martien, P. T. and Kirchstetter, T. W.: Measurement of black carbon emissions from in-use diesel electric passenger locomotives in California, *Atmospheric Environment*, 115, 295–303, doi:10.1016/j.atmosenv.2015.05.001, 2015.

Tissari, J., Sippula, O., Kouki, J., Vuorio, K. and Jokiniemi, J.: Fine Particulate and Gas Emissions from the Combustion of Agricultural Fuels Fired in a 20 kW Burner, *Energy Fuels*, 22(3), 2033–2042, doi:10.1021/ef700766y, 2008.

10 Tsai, J. H., Chen, S. J., Huang, K. L., Lin, Y. C., Lee, W. J., Lin, C. C. and Lin, W. Y.: PM, carbon, and PAH emissions from a diesel generator fuelled with soy biodiesel blends, *Journal of Hazardous Materials*, 179(1–3), 237–243, doi:10.1016/j.jhazmat.2010.02.085, 2010.

Uma, R., Kandpal, T. C. and Kishore, V. V. N.: Emission characteristics of an electricity generation system in diesel alone and dual fuel modes, *Biomass and Bioenergy*, 27(2), 195–203, doi:10.1016/j.biombioe.2004.01.003, 2004.

UNEP/WMO: Integrated Assessment of Black Carbon and Tropospheric Ozone, Nairobi, Kenya. [online] Available from: www.unep.org/dewa/Portals/67/pdf/BlackCarbon_report.pdf, 2011.

15 US EPA: 2011 Version 6 Air Emissions Modeling Platforms, [online] Available from: <https://www.epa.gov/air-emissions-modeling/2011-version-6-air-emissions-modeling-platforms>, 2011.

Verma, S. and Uppal, J.: Use of biomass in brick kilns, Ministry of New and Renewable Energy, Government of India, 6(4), 24–27, 2013.

20 Wang, A., Ge, Y., Tan, J., Fu, M., Shah, A. N., Ding, Y., Zhao, H. and Liang, B.: On-road pollutant emission and fuel consumption characteristics of buses in Beijing, *J. Env. Sci.*, 23(3), 419–426, doi:10.1016/S1001-0742(10)60426-3, 2011.

Wang, S. X., Zhao, X. J., Li, X. H., Wei, W. and Hao, J. M.: Emission characteristics of fine particles from grate firing boilers, *Environmental Science*, 30(4), 963–968, 2009.

25 Watson, J. G., Miller, W., Moosmuller, H., Kuhns, H. D., Chow, J. C., Zhu, D. and Cocker, D. R.: Characterization of Off-Road Diesel Emissions of Criteria Pollutants, CY2006 Annual Report, WP 1336, Desert Research Institute, Reno, NV., 2006.

Watson, J. G., Miller, J. W., Moosmüller, H., Kuhns, H. D., Chow, J. C., Chang, O., Nussbaum, N., Mazzoleni, C., Zhu, D., Barber, P. W., Kemmle, M. R. and Cocker, D. R.: Characterization of off-road diesel emissions of criteria pollutants, Desert Research Institute, Reno, NV. [online] Available from: <https://www.dri.edu/images/stories/editors/cafeditor/Watsonetal2008SERDPFinalReportDieselWP1366.pdf>, 2008.

30 Weyant, C., Athalye, V., Ragavan, S., Rajarathnam, U., Lalchandani, D., Maithel, S., Baum, E. and Bond, T. C.: Emissions from South Asian Brick Production, *Environ. Sci. Technol.*, 48(11), 6477–6483, doi:10.1021/es500186g, 2014.

World Bank: Introducing energy efficient clean technologies in the brick sector of Bangladesh, World Bank, Washington DC, USA. [online] Available from: <http://documents.worldbank.org/curated/en/2011/06/15492357/introducing-energy-efficient-clean-technologies-brick-sector-bangladesh>, 2011.

Yang, H. H., Hsieh, L. T., Liu, H. C. and Mi, H. H.: Polycyclic aromatic hydrocarbon emissions from motoreycles, *Atmospheric Environment*, 39(1), 17–25, doi:10.1016/j.atmosenv.2004.09.054, 2005.

Yanowitz, J., McCormick, R. L. and Graboski, M. S.: In Use Emissions from Heavy Duty Diesel Vehicles, *Environmental Science & Technology*, 34(5), 729–740, doi:10.1021/es990903w, 2000.

5 Zhang, Q., Klimont, Z., Streets, D., Huo, H. and He, K.: An anthropogenic PM emission model for China and emission inventory for the year 2001, *Progress in Natural Science*, 16(2), 223–231 (in Chinese), 2006.

Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T. and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX B mission, *Atmos. Chem. Phys.*, 9, 5131–5153, 2009.

10 Zhang, Y., Schauer, J. J., Zhang, Y., Zeng, L., Wei, Y., Liu, Y. and Shao, M.: Characteristics of Particulate Carbon Emissions from Real-World Chinese Coal Combustion, *Environ. Sci. Technol.*, 42(14), 5068–5073, doi:10.1021/es7022576, 2008.

Zhang, Z.: Energy efficiency and environmental pollution of brickmaking in China, *Energy*, 22(1), 33–42, 1997.

15 Zhi, G., Peng, C., Chen, Y., Liu, D., Sheng, G. and Fu, J.: Deployment of Coal Briquettes and Improved Stoves: Possibly an Option for both Environment and Climate, *Environ. Sci. Technol.*, 43(15), 5586–5591, doi:10.1021/es802955d, 2009.

Zhu, D., Nussbaum, N. J., Kuhns, H. D., Chang, M. C. O., Sodeman, D., Uppapalli, S., Moosmüller, H., Chow, J. C. and Watson, J. G.: In Plume Emission Test Stand 2: Emission Factors for 10–100 kW U.S. Military Generators, *Journal of the Air & Waste Management Association*, 59(12), 1446–1457, doi:10.3155/1047-3289.59.12.1446, 2009.