

Case study of wave breaking with high-resolution turbulence measurements with LITOS and WRF simulations

Andreas Schneider^{1,3}, Johannes Wagner², Jens Söder¹, Michael Gerding¹, and Franz-Josef Lübken¹

¹Leibniz Institute of Atmospheric Physics at the University of Rostock (IAP), Kühlungsborn, Germany

²German Aerospace Center (DLR), Institute of Atmospheric Physics (IPA), Wessling, Germany

³Now at: SRON Netherlands Institute for Space Research, Utrecht, the Netherlands

Correspondence to: Andreas Schneider (a.schneider@sron.nl)

Abstract. Measurements of turbulent energy dissipation rates obtained from wind fluctuations observed with the balloon-borne instrument LITOS (Leibniz-Institute Turbulence Observations in the Stratosphere) are combined with simulations with the Weather Research and Forecasting (WRF) model to study the breakdown of waves into turbulence. One flight from Kiruna (68° N, 21° E) and two flights from Kühlungsborn (54° N, 12° E) are analysed. Dissipation rates are in the order of 0.1 mW kg^{-1} ($\sim 0.01 \text{ K d}^{-1}$) in the troposphere and in the stratosphere below 15 km, increasing in distinct layers by about two orders of magnitude. For one flight covering the stratosphere up to $\sim 28 \text{ km}$, the measurement shows nearly no turbulence at all above 15 km. Another flight features a patch with highly increased dissipation directly below the tropopause, collocated with strong wind shear and wave filtering conditions. In general, small or even negative Richardson numbers are affirmed being a sufficient condition for increased dissipation. On the other side, significant turbulence has also been observed in the lower stratosphere under stable conditions. Observed energy dissipation rates are related to wave patterns visible in the modelled vertical winds. In particular, the drop in turbulent fraction at 15 km mentioned above coincides with a drop in amplitude in the wave patterns visible in WRF. This indicates wave saturation being visible in the LITOS turbulence data.

1 Introduction

Gravity waves transport energy and momentum and are thus an important factor in the atmospheric energetics. Typically, they are excited in the troposphere and propagate upwards and horizontally. Due to decreasing density, the amplitudes

increase with altitude in the absence of damping. Eventually, the waves become unstable and break, producing turbulence and dissipation, and thereby depose their energy and momentum. This mechanism has been suggested by Hodges (1967) to explain turbulence in the mesosphere. There are two variants of wave breaking (e.g. Hocking, 2011, Section 9): First catastrophic wave breaking, where the wave is completely annihilated (e.g. Andreassen et al., 1994), and second wave saturation, where a wave loses energy to turbulence so that the amplitude does not increase further, i.e. the wave breaks only partially (e.g. Lindzen, 1981). Hines (1991) defines saturation to imply that the wave amplitude is at a maximum and the excess energy is shed by physical processes to prevent further growth. There are several theories for saturation (Fritts and Alexander, 2003, Section 6.3), and the phenomenon has been observed as well. For example, using a balloon-borne instrument Cot and Barat (1986) measured a gravity wave in winds and temperature with vertical wavelength of $\sim 1 \text{ km}$ and nearly constant amplitude over $\sim 5 \text{ km}$ height. Simultaneously they observed several turbulent patches collocated with negative temperature gradient and Richardson numbers between 0.3 and 6. They concluded that clear air turbulence is related to a long-period wave via shear instability, and that the energy budget of the wave-turbulence interaction is in an order of magnitude that the wave amplitude would not change much. Franke and Collins (2003) observed gravity waves in the mesosphere with Na lidar and found upwards propagating waves still present (with less amplitude) above an overturning region. Catastrophic wave breaking has been observed, e.g., in the lowermost stratosphere by Worthington (1998) and Pavelin et al. (2001) with radar and radiosonde. Model studies of breaking gravity waves have, e.g., been carried out by Achatz (2005) and by

Fritts and Wang (2013), Fritts et al. (2016), who performed direct numerical simulations (DNS) of a gravity wave superposed by fine-scale shear.

Regarding turbulence measurements, there are two aspects of importance: first, its energy dissipation, and secondly its diffusive properties. We will concentrate on the former. Large-scale diffusion in the stratosphere is a complex process due to the intermittent nature of the turbulence there, as summarised in some detail by Osman et al. (2016), among others. A relatively extensive dataset exists for the troposphere and tropopause region (e. g. Lilly et al., 1974; Hauf, 1993; Cho et al., 2003), but in the middle stratosphere observations are sparse. Remote sensing is mainly performed by radars in the troposphere and lower stratosphere as well as in the mesosphere (see Wilson, 2004, for an overview), and with satellites in the upper stratosphere (e. g. Gavrilov, 2013). In situ observations in the middle stratosphere have been carried out with balloon-borne instruments. Pioneering work has been done by Barat (1982) and Dalaudier et al. (1994). An instrument with a similar anemometer has been developed by Yamanaka et al. (1985). Indirect measurements using the Thorpe method were performed by Luce et al. (2002); Clayson and Kantha (2008) and others, mainly using standard radiosondes. A recent high-resolution balloon-borne instrument for the direct measurement of turbulent wind fluctuations is Leibniz Institute Turbulence Observations in the Stratosphere (LITOS) (Theuerkauf et al., 2011), which can resolve the inner scale of turbulence in the stratosphere for the first time. This state of the art instrument is used for this study.

To study wave breaking into turbulence, a wide range of scales from kilometres (the wavelength of GWs) to millimetres (the viscous subrange of turbulence) has to be resolved. This cannot be performed by a single instrument. Thus several techniques have to be combined. In this study, LITOS is used for the turbulence part and radiosonde observations from the same gondola for local atmospheric background conditions. To put the observations into a geophysical context and to obtain information about waves, regional model simulations with WRF (Weather Research and Forecasting model) driven by reanalysis data are applied. Three flights are analysed, comprising one from Kiruna (northern Sweden, 67.9° N, 21.1° E) and two from Kühlungsborn (northern Germany, 54.1° N, 11.8° E).

This paper is structured as follows: Section 2 gives an overview of the instrument LITOS and the data retrieval (Section 2.1) as well as the WRF model setup (Section 2.2). The results for three different flights are presented in Section 3. These are interrelated and discussed in Section 4, and finally conclusions are drawn in Section 5.

2 Instrumentation and model

2.1 Balloon-borne measurements

LITOS (Leibniz-Institute Turbulence Observations in the Stratosphere) is a balloon-borne instrument to observe small-scale fluctuations in the stratospheric wind field (Theuerkauf et al., 2011). The wind measurements are performed with a constant temperature anemometer (CTA) which has a precision of a few cm s^{-1} . It is sampled with 8 kHz yielding a sub-millimetre vertical resolution at 5 m s^{-1} ascent rate. Thus the inner scale of turbulence is typically covered. A standard meteorological radiosonde (Vaisala RS92 or RS41) is used to record atmospheric background parameters. LITOS was launched three times as part of a $\sim 120 \text{ kg}$ payload from Kiruna (67.9° N, 21.1° E) within Balloon Experiments for University Students (BEXUS) 6, 8 and 12 in 2008, 2009 and 2011, respectively (Theuerkauf et al., 2011; Haack et al., 2014; Schneider et al., 2015). The second generation of the small version of the instrument is an improvement of the one described by Theuerkauf et al. (2011) and consists of a spherical payload of $\sim 3 \text{ kg}$ weight. It is suspended $\sim 180 \text{ m}$ below a meteorological rubber balloon. Two CTA sensors are mounted on booms protruding at the top of the gondola. The instrument was launched several times from IAP's site at Kühlungsborn (54.1° N, 11.8° E), e. g. at 27 Mar 2014, 06 Jun 2014, and 12 Jul 2015.

In this paper, only flights are taken into account where data from more than one CTA sensor on the same gondola are available. Summarised, the data analysis is performed in three steps. First, the dissipation rate is retrieved similar as described by Theuerkauf et al. (2011). Then the ϵ values from both sensors are compared to detect sections where one sensor is possibly affected by the wake of ropes. Finally, the remaining spectra are manually inspected to sort out cases where both sensors potentially have been affected. Another source of artificial turbulence is the wake of the balloon (Barat et al., 1984). Typically, the wake influences both sensors similarly and cannot be detected by the above methods. Therefore, we limit our analysis to flights and altitude regions, where wake effects do not play a role due to sufficient wind shear that brings the payload out of the balloon's wake.

The details of the retrieval are as follows: The data of the ascent is split into windows with depths of 5 m altitude with 50 % overlap. In each window, the mean value is subtracted, and the periodogram is computed, which is an estimation of the power spectral density (PSD). The periodogram is smoothed with a Gaussian-weighted running average. The instrumental noise level is detected and subtracted. Initially, turbulence is assumed in each window and thus the Heisenberg (1948) model for fully developed turbulence in the form given by Lübken and Hillert (1992) and Theuerkauf et al. (2011) is tried to fit to the observed spectrum (cf. Equation (A3) in Appendix A). If the fit succeeds, the inner scale

l_0 is obtained. This leads to the energy dissipation rate ε given by

$$\varepsilon = c_{l_0}^4 \frac{v^3}{l_0^4}, \quad (1)$$

where v is the kinematic viscosity (known from the radiosonde measurement) and c_{l_0} is a constant depending on the type of sensor. The determination of c_{l_0} for our sensor configurations is described Appendix A. Non-turbulent (or disturbed) spectra manifest in bad fits which are sorted out with the following set of criteria:

- The noise level detection fails, which usually means that the noise is not white, i. e. the periodogram is disturbed at small scales.
- The mean logarithmic difference between data and fit exceeds a given threshold. This condition captures cases where the fit does not describe the data well, e. g. when no turbulence is present so that the periodogram does not follow form of the turbulence model.
- The inner scale l_0 lies outside the fit range. This means that the bend in the spectrum is not within the fit range and thus the fit is not meaningful, allowing no useful retrieval of ε . That can occur when the spectrum does not have the expected form of the turbulence model, when the inner scale lies at very small scales where the periodogram is dominated by noise, or when the periodogram is disturbed.
- The fit width is smaller than a threshold; in this case the fit is determined by too few data points.
- The value of the periodogram at l_0 is too close to the value of the noise level, i. e. too small a part of the viscous subrange is resolved.
- The slope of the fit function at the small-scale end is less than a given threshold (less steep than m^{-4} , where m is the vertical wave number). This indicates that the bend in the spectrum is not well covered by the fit and the data.

If one of the above conditions applies, the spectrum does not follow the form for fully developed turbulence, thus ε is set to zero. Requiring the spectrum to follow Heisenberg's turbulence model may exclude turbulence that is not fully developed. However, it is not feasible to retrieve ε in cases where the periodogram does not follow the turbulence model.

Sometimes a sensor has been located in the wake of a rope supporting the gondola and the other sensor not, causing the ε values of both sensors to differ by up to 5 orders of magnitude. To sort out such sections, altitude bins where the dissipation rate from both sensors deviates by more than a factor of 15 are discarded. For the flights with the small payload, the remaining spectra have been inspected manually for

sections where both sensors have been affected by the rope wake, and those that look suspicious have been taken out. A spectrum is regarded as wake-affected if it has a plateau in PSD near 10 cm spatial scale, which is estimated to be the extent of a Kármán vortex street originating from the lines supporting the gondola. This problem of wake effects from the ropes does not occur for the BEXUS flights, where the sensors were placed further away from the supporting lines. For all other altitude bins the average of both sensors is taken.

On the other hand, the BEXUS flight had a comparatively small distance between balloon and payload of 50 m. Thus, during considerable times the payload flew through the wake of the balloon. Therefore, only limited altitude sections with large wind shears are considered for this flight.

To quantify the stability of the atmosphere, the gradient Richardson number $Ri = N^2/S^2$ is used, which is the ratio of the squared Brunt-Väisälä frequency N^2 and the square of the vertical shear of the horizontal wind S^2 . The Brunt-Väisälä frequency can be written as $N^2 = \frac{g}{\Theta} \frac{d\Theta}{dz}$, where Θ is the potential temperature and g the acceleration due to gravity. The wind shear is defined as $S^2 = \left(\frac{du}{dz}\right)^2 + \left(\frac{dv}{dz}\right)^2$, where u and v are the zonal and meridional wind components, respectively. The Richardson number represents the ratio of buoyancy forces (which suppress turbulence) to shear forces (which generate turbulence). According to a theory for plane-parallel flow established by Miles (1961) and Howard (1961), turbulence occurs below a critical Richardson number of $Ri_c = 1/4$. The general applicability of that criterion was recently questioned based on measurements (e. g. Balsley et al., 2008) and model simulations (e. g. Achatz, 2005). Often the shear is not strictly horizontal so that the theory by Miles (1961) and Howard (1961) is not applicable, as pointed out by Achatz (2005). To take into account slanted shear, Hines (1988) proposed a concept of slantwise instability. However, the Richardson number is still useful as an estimation of stability. The Richardson number also depends on the scale on which it is computed (Balsley et al., 2008; Haack et al., 2014). Usually, computing Ri on a smaller scale yields locally smaller numbers, since for a computation on larger scales an average over regions with small and large Ri is obtained. In this study Ri is retrieved from the radiosonde measurements. In order not to dominate the derivatives by instrumental noise, the potential temperatures and winds are smoothed with a Hann-weighted running average over 150 m prior to differentiation with central finite differences.

2.2 Model simulations

Mesoscale numerical simulations are performed with the Weather Research and Forecasting (WRF) model, version 3.7 (Skamarock et al., 2008). Two nested domains with horizontal resolutions of 6 km and 2 km and time step 15 s and 5 s, respectively, are applied. In the vertical direction 138 terrain following levels with stretched level distances of 80 m near the surface and 300 m in the stratosphere are used and

the model top is set to 2 hPa (about 40 km altitude) for the BEXUS flights and 5 hPa (about 32 km altitude) for the flights from K hlungsborn. At the model top a 7 km thick Rayleigh damping layer is applied to prevent wave reflections (Klemp et al., 2008), i. e. the top of the damping layer is the model top. Physical parametrisations contain the Rapid Radiative Transfer Model longwave scheme (Mlawer et al., 1997), the Goddard shortwave scheme (Chou and Suarez, 1994), the Mellor-Yamada-Nakanishi-Niino boundary layer scheme (Nakanishi and Niino, 2009), the Noah land surface model (Chen and Dudhia, 2001), the WRF single-moment 6-class microphysics scheme (WSM6; Hong and Lim, 2006) and the Kain-Fritsch cumulus parametrisation scheme (Kain and Fritsch, 1990). The initial and boundary conditions are supplied by ECMWF (European Centre for Medium-Range Weather Forecasts) operational analyses on 137 model levels with a temporal resolution of 6 hours. In WRF a temporal output interval of 1 hour is used, data interpolated along the flight track are output with an interval of 5 minutes. Simulations are initialised 5 to 6 hours before the launch time of the balloon. The computation of turbulent kinetic energy (TKE) is done by the boundary layer scheme and described in Nakanishi and Niino (2009). It is based on a prognostic equation which is solved additionally to the equations of motion and which includes transport, shear production, buoyancy production and dissipation terms. Shear and buoyancy terms include deformation and stability effects of the resolved flow and are related to turbulent motions by the horizontal and vertical eddy viscosities. The equation operates on the scale of the grid size.

In this paper WRF simulations are used to get an overview of the meteorological situation. Ehard et al. (2016) showed that regions of GW breaking can be simulated by WRF simulations with horizontal grid distances of 2 km and a similar model set-up by means of convective overturning and reduced Richardson numbers. Here, the TKE output from the model is also used to identify regions of intensified turbulent mixing in the atmosphere along the balloon flight tracks. This can be a hint that observed turbulence was caused by large-scale GW breaking. It is not intended to quantitatively compare observed dissipation rates with simulated regions of enhanced TKE values.

3 Results

3.1 The BEXUS 12 flight (27 September 2011)

The BEXUS 12 flight was launched from Kiruna on 27 Sep 2011 at 17:36 UT. The two left panels of Figure 1 show atmospheric conditions as observed by the radiosonde on board the payload. Temperatures decreased up to the tropopause at 10.3 km, excepting some small inversion layers. Above there was a sharp increase in temperature known as tropopause inversion layer (TIL) (Birner et al., 2002;

Birner, 2006). Higher up, temperatures slightly decreased. Winds came from north-west near the surface and reversed between ~ 6 km and 10 km. The reversal caused nearly opposite wind direction at 9 km altitude compared to 5 km, and a change of sign in both wind components. It further entailed strong wind shear below the tropopause, causing low Richardson numbers (below the critical number of $1/4$). Above the tropopause the wind field showed signatures of gravity wave activity with short wavelengths and no obvious altitude-dependent structure. In the stratosphere, Richardson numbers were generally larger than in the troposphere.

The right panel of Figure 1 depicts observed dissipation rates. Each blue cross corresponds to an altitude bin classified as turbulent (as described in Section 2.1). The orange curve depicts a Hann-weighted running average over 500 m. Please note that large sections in the troposphere and stratosphere are subject to wake influence (marked grey) due to the small distance of only 50 m between the payload and the balloon. These sections are generally not discussed here. Between 9 km and 10 km there was a thick layer with high dissipation. As described above, this altitude region featured low Richardson numbers caused by high wind shears. Thus turbulence was presumably induced by dynamic instability. Additionally, at this altitude a wind reversal was observed which caused filtering of gravity waves with phase velocities equal to the background winds (if present). Most probably, these high dissipation rates are not caused by wake because calculations show that the gondola was outside the wake in this altitude section due to the large wind shear. Furthermore, the dissipation rates are even larger than typical wake turbulence.

WRF model simulations were performed for the time and place of the flight. To show that these produced reasonable results, model winds and temperatures interpolated along the flight trajectory are plotted in the left panel of Figure 1 along with the radiosonde profiles. Observed and modelled results compare very well, the only difference is that the radiosonde data contain signatures from small-scale gravity waves which WRF cannot resolve. In Figure 2, model snapshots at the middle of the ascent are shown. The upper left panel depicts horizontal winds at 850 hPa. Westerly winds flowed over the Scandinavian mountains which are expected to have excited mountain waves. Another potential source of gravity waves is geostrophic adjustment. Bending stream lines are visible, e.g., over the Scandinavian mountains, west of the flight track. The upper right panel presents a vertical section of horizontal winds and potential temperatures. It demonstrates that the jet (~ 7 km to 10 km altitude) had a local structure and involved strong wind shears.

With a grid resolution of 2 km WRF can resolve waves with horizontal wavelengths larger than about 10 km. These waves can be seen, e.g., in the vertical winds, which are used as a proxy. This quantity is plotted in the lower left panel of Figure 2. Strong wave-like patterns are visible especially over the Scandinavian mountains, which correspond to the mountain wave excitation mentioned above. Weaker




Figure 1. Observations during the BEXUS 12 flight. Left: Zonal winds u (blue), meridional winds v (green) and temperatures T (red) from the radiosonde. The light blue, light green, and orange curves show the corresponding results from the WRF model interpolated along the balloon trajectory. Centre left: Wind direction (blue) and horizontal wind speed (green) from the radiosonde. Centre right: Richardson number Ri computed from the radiosonde data, using a smoothing over 150 m prior to numerical differentiation. The Ri axis is split at 1 into a linear and a logarithmic part. The red line shows the critical Richardson number $1/4$. Right: Energy dissipation rates ϵ observed by LITOS. The blue crosses mark single turbulent spectra computed on a 5 m grid with 50 % overlap, the orange curve shows a Hann-weighted running average over 500 m (non-turbulent bins count as zero in the average). The top axis gives the heating rate due to turbulent dissipation, $dT/dt = \epsilon/c_p$. The grey areas mark the regions with likely wake influence. The horizontal black line in all four panels marks the tropopause.

wave patterns are visible near the flight trajectory, downstream of the mountains. Between roughly $x = 400$ km and $x = 550$ km, the wave patterns change at tropopause height (approximately 10 km altitude): Above there is less amplitude than below. This is ascribed to the wave breaking and filtering mentioned before. Filtering means catastrophic breaking of waves, i. e. a wave that is filtered is annihilated. Further upwards the amplitude increases slowly.

Waves can propagate over considerable distances and times. Therefore it is not sufficient to look at potential sources in the vicinity of the flight track. Even if sources are found, the waves may have propagated to other places (away from the point of interest), while waves from sources outside the domain may have propagated to the location of observation. For resolved waves the model takes care of these issues. Waves seen in WRF at the location of the flight may have travelled from remote places, yet the important information is not their origin, but that they were present during the measurement.

To trigger turbulence, wave *breaking* is necessary. Such events are triggered by dynamic or convective instabilities or by wave-wave interactions (e. g. Fritts and Alexander, 2003). In WRF, the break-down to turbulence is parametrised by solving a prognostic equation for turbulent kinetic energy

(TKE), which is based on production terms due to shear and buoyancy obtained from the resolved flow. TKE is plotted in the lower right panel of Figure 2. It peaks near 10 km height at the location of the flight. This corresponds nicely to the intense turbulent layer observed by LITOS. It is reproduced in WRF due to the shear instability on scales resolved by the model. That highlights the geophysical significance of that layer.

3.2 The 27 March 2014 flight

A small LITOS payload of second generation was launched from Kühlungsborn on 27 Mar 2014 at 10:10 UT. It was carried by a comparatively small (3000 g) balloon and a 60 m dereeler.

The left panel of Figure 3 shows temperatures smoothed over 15 data points (~ 150 m) as well as zonal and meridional winds. The smoothing is necessary because for this flight the temperature measurement is perturbed by radiation effects as the radiosonde was incorporated in the main payload; these effects get worse with increasing altitude. Temperatures decreased up to the tropopause at 9 km. Between 9 km and ~ 30 km altitude they stayed nearly constant and started to increase further upwards. Winds were easterly and turned northerly above ~ 20 km altitude. A strong southeast-




Figure 2. Map of horizontal winds at 850 hPa (upper left), vertical section of horizontal winds (upper right), vertical section of vertical winds (lower left), and vertical section of turbulent kinetic energy (TKE) (lower right) from WRF simulations for 27 Sep 2011, 18:00 UT. The black curves visualise the trajectory of the BEXUS 12 flight. In the upper left panel, the blue streamlines show the wind direction, the white lines visualise coastlines and a latitude/longitude grid, and the black line indicates the location of the vertical sections. In the upper right panel, the white isolines show potential temperature.




Figure 3. Same as Figure 1, but for the flight from K hlungsborn at 27 Mar 2014. Due to disturbances of the temperature data, temperatures are smoothed in the plot in the left panel, and Richardson numbers are shown only for altitudes lower than 9.4 km. The dissipation profile excludes the lowermost 650 m due to disturbances from the launch procedure (dereeling of the payload suspension), and the part above 9.4 km altitude due to potential wake effects from the balloon.

erly jet was present between ~ 6 km and 10 km height. Superposed are signatures of small-scale gravity waves. Wind shears originating from the jet may have excited turbulence and/or waves. The effect of the shear is visible as a layer with enhanced dissipation at this altitude (see below). Richardson numbers are shown for altitudes below 9.4 km only because they involve derivatives of the temperature profile which was disturbed by radiation effects as described above.

Dissipation rates are presented in the right panel of Figure 3. The data below 650 m altitude are affected by the unwinding of the dereelers while the data above the tropopause are subject to wake influence. Therefore, these are discarded and not shown in the plot. Dissipation rates varied over several orders of magnitude within only small altitude ranges (typically a few 10 m). The running average shows some structure in the troposphere, e. g. a few layers that are standing out with larger rates. Most prominently this can be seen near 8 km. That is in the same altitude as the wind shear due to the jet, which speaks for shear-induced turbulence. Precisely, there were two turbulent layers from 7.5 km to 7.9 km and from 8.1 km to 8.3 km height; within both, Richardson numbers were below 1 and partly below 1/4. Other sheets with large dissipation were detected, e. g., near 6.1 km and around 3.0 km altitude.

To validate the corresponding WRF simulations, winds and temperatures interpolated to the flight track are plotted in the left panel of Figure 3. They agree very well to the radiosonde data. Figure 4 depicts WRF results for the time

of the flight. The upper left panel shows horizontal winds at 850 hPa, which were easterly or south-easterly. In the upper right panel horizontal winds are depicted as altitude section, showing that the strong jet had not much structure in horizontal direction, while the sharp vertical structure is reproduced as observed by the radiosonde. The lower left panel shows a vertical profile of vertical winds. Wave patterns are visible, which stretch over the whole altitude range. Particularly, a superposition of a wave with long vertical wavelength ($\lambda_z \approx 8$ km) and nearly horizontal phase fronts and waves with short horizontal wavelength (10 km to 20 km) and phase fronts in the vertical can be seen. The lower right panel of Figure 4 shows the TKE. Outside the boundary layer there is an enhancement near 7.5 km altitude. It corresponds nicely to a thick, strong turbulent layer in the measurement by LITOS between ~ 7 km and 8.5 km height. Within this observed turbulent layer, which in fact consists of several layers, Richardson numbers are smaller than 1 almost everywhere and at times smaller than 1/4.

3.3 The 11/12 July 2015 flight

A night-time flight with LITOS was performed on 11/12 Jul 2015 from K hlungsborn, launched at midnight local time (22:01 UT on 11 Jul). A dereeler of 180 m (with a 3000 g balloon) was used for payload suspension, making balloon wake effects negligible for this flight. The radiosonde was positioned 60 m below the main payload to




Figure 4. Same as Figure 2, but for WRF simulations for 27 Mar 2014, 11:00 UT.




Figure 5. Same as Figure 1, but for the flight from Kühlungsborn at 11/12 Jul 2015. The dissipation profile excludes the lowermost 550 m due to disturbances from the launch procedure (dereeling of the payload suspension).

avoid disturbances of the temperature sounding. The observed background parameters are depicted in the two left panels of Figure 5. Westerly winds prevailed up to ~ 19 km altitude, whereas above winds came from the east. This change in direction was not associated with a significant wind shear because velocities were small in that altitude region. A jet is visible at about 10 km height. Superposed on the winds are signatures of small-scale gravity waves. Above the tropopause at 11.3 km altitude there was a small tropopause inversion layer. Higher up temperatures remained rather constant up to ~ 20 km, where they started to increase.

Richardson numbers were typically lower than for the other flights, indicating less stability. There are several layers where the Richardson number is below the critical limit of Ri_c ($1/4$). These layers are relatively thin.

Energy dissipation rates (data below 550 m are excluded due to disturbances from the launch procedure) showed a strong patchy structure, with enhanced dissipation at, e.g., ~ 2.0 km, 3.8 km, 7.2 km, 8.9 km, 11.0 km, 12.1 km, and 14.3 km. These layers of intense turbulence mostly corresponded to Richardson numbers smaller than $Ri_c = 1/4$, or at least to $Ri < 1$. But particularly in the lower stratosphere between 11 km and 15 km, turbulence occurred also for high Richardson numbers. It should be kept in mind that the Richardson number depends on the scale on which it is computed (e.g. Balsley et al., 2008; Haack et al., 2014). A higher resolution (i.e. computing Ri on smaller scales) may result in locally smaller Ri numbers, because the computation on large scales yields a kind of average. Similarly, Paoli et al. (2014) found in Large Eddy Simulations larger Richardson numbers

for smaller model resolutions (i.e. larger scales). Here, due to measurement noise a smoothing over 150 m has been applied before computing Ri , determining the resolution. However, this issue cannot explain the whole discrepancy. In simulations of gravity waves, Achatz (2005) found instabilities and onset of turbulence for Richardson numbers both smaller and larger than $1/4$. He noted that the theory by Miles (1961) and Howard (1961) is not applicable to his simulations because the gravity wave phase propagation and thus the wave-induced shear is slanted. In the real atmosphere waves usually propagate at a tilt (i.e. the shear is not orthogonal to the altitude axis). Already Hines (1988) discussed slantwise static instabilities created by gravity waves. He developed a wave period criterion for turbulence by comparing the e-folding time of the (slantwise) instability with the period of the wave. Turbulence is more likely to occur for slantwise static instability than for vertical static instability. In the light of these comments, the violation of the Richardson criterion for the LITOS measurements is comprehensible.

Above ~ 15 km altitude, hardly any turbulence was detected; only a few thin turbulent layers were observed. Thus above 15 km the average dissipation rate (for which no turbulence is counted as zero) was only 0.01 mW kg^{-1} , while below 15 km it was 0.64 mW kg^{-1} .

Results from corresponding WRF simulations are depicted in Figure 6. Horizontal winds at the 850 hPa level were mainly westerly. The altitude section shows that the strong jet did not have much variation in the horizontal direction. Vertical winds reveal wave patterns that are particularly intense around the tropopause and gradually become weaker




Figure 6. Same as Figure 2, but for WRF simulations for 11 Jul 2015, 23:00 UT

near ~ 15 km, with less amplitude above. This drop in wave amplitude is at the same altitude as the drop in observed dissipation. The TKE has enlarged values around 3 km altitude and near the tropopause, however the enhancement is small at the flight path. Correspondingly, the thickness of the strong turbulent layers detected by LITOS is relatively small; that means that these dissipative layers are potentially not resolved in the model.

4 Discussion

A comparison of the observed dissipation profiles and the wave patterns in the model vertical winds for the different flights suggests that more turbulence observed by LITOS comes along with stronger wave patterns visible in WRF, and vice versa. Particularly, this can be seen at 11/12 July 2015 at the drop in dissipation and wave amplitude at ~ 15 km altitude. A similar feature has been observed during another flight at 06 Jun 2014 (not shown): Likewise, LITOS data exhibit a sharp drop in turbulence at ~ 15 km, and the corresponding WRF simulation shows strong wave patterns below ~ 15 km and very weak ones above. For the troposphere, vertical winds in WRF show similar gravity wave amplitudes for both Kühlungsborn soundings, even if the wave structures are different. Accordingly, dissipation rates are generally similar, showing up as a highly structured profile that is partly related to shear instabilities measured by the radiosonde. This reflects also in the WRF turbulent kinetic energy, attesting that the structures are sufficiently large to be resolved in WRF. The same is true for the turbulent layer below the tropopause observed during BEXUS 12.

The relation between waves and turbulence can also be seen in averages over altitude regions. For 12 Jul 2015 the most significant drop in mean dissipation does not happen at the tropopause where the stability increases due to the changing temperature gradient, but at ~ 15 km where the wave activity decreases. Mean energy dissipation rates are 0.64 mW kg^{-1} below 15 km altitude and 0.01 mW kg^{-1} above. Consistently, the average absolute vertical flux calculated from WRF data as a measure for wave activity is 64 mW m^{-2} below 15 km and 6.9 mW m^{-2} above.

We interpret this behaviour as the effect of wave saturation. As described in the introduction, a saturated wave loses part of its energy to turbulence so that the amplitude does not grow further. Such effects have already been observed, e.g., by Cot and Barat (1986), who measured a gravity wave with almost constant amplitude over an altitude range of 5 km and collocated isolated turbulent patches with a dissipation rate approximately accounting for the energy loss of the wave. Franke and Collins (2003) found regions of strong overturning, and upwards propagating waves present below as well as (with less amplitude) above the overturning region. They argue that, depending on the amplitude, a breaking wave is not always completely annihilated, but the ampli-

tude may be modulated in a highly non-linear event. Nappo (2002, p. 125) states that “gravity wave and turbulence are often observed to exist simultaneously.” Via the process of wave saturation, the occurrence of waves is connected to the intensity of turbulence. Pavelin et al. (2001) observed intense turbulence in the lowermost stratosphere during a period of maximal wave intensity using radar at Aberystwyth (52.4° N , 4.0° W), which supports the above hypothesis.

Saturation theories proposed several mechanisms, e.g. linear instability dynamics due to large wave amplitudes, non-linear damping, or non-linear wave-wave interactions (Fritts and Alexander, 2003, Section 6.3). The present study cannot answer that debate, yet the relatively large Richardson numbers hint that non-linear interactions may play a role.

Mean dissipation rates observed by LITOS are in the order of $10^{-4} \text{ W kg}^{-1}$ (roughly 0.01 K d^{-1}). This is two orders of magnitude below typical solar or chemical heating rates which are in the order of 1 K d^{-1} (Brasseur and Solomon, 1986, Fig. 4.19b). However, within thin layers rates of $10^{-2} \text{ W kg}^{-1}$ to $10^{-1} \text{ W kg}^{-1}$ ($\sim 1 \text{ K d}^{-1}$ to 10 K d^{-1}) are observed, which is larger than solar heating. The low mean energy dissipation rates are not explicitly contained even in high-resolution models, which cannot describe the large intermittency. Only large layers with highly increased dissipation as encountered, e.g., during BEXUS 12 are captured.

Observed dissipation rates are partly larger than those reported by other publications using different methods. Barat (1982) obtained values between $1.4 \times 10^{-5} \text{ W kg}^{-1}$ and $3.9 \times 10^{-5} \text{ W kg}^{-1}$ from balloon measurements. Wilson et al. (2014) found ε values between $3 \times 10^{-5} \text{ W kg}^{-1}$ and $6 \times 10^{-4} \text{ W kg}^{-1}$ in the upper troposphere from radar measurements. These are lower rates than the averages in this work, but within the range of the variability. Lilly et al. (1974) observed stratospheric dissipation rates between $7 \times 10^{-4} \text{ W kg}^{-1}$ and $2 \times 10^{-3} \text{ W kg}^{-1}$, depending on the underlying terrain, with aircraft. These results are in similar order of magnitude as the averages in this study. Haack et al. (2014) reported mean dissipation rates between $2 \times 10^{-2} \text{ W kg}^{-1}$ and $5 \times 10^{-3} \text{ W kg}^{-1}$ for the altitude range 7 km to 26.5 km, using a different retrieval and potentially including wake effects.

5 Conclusions

In this paper high-resolution turbulence observations with LITOS are complemented by model simulations with WRF to study the relation between turbulence, waves, and background conditions. Three flights are selected where in each case data from two wind sensors are available; this allows a high quality assurance. Furthermore, any data that is possibly influenced by the balloon’s wake has been removed for this study.

Enhanced energy dissipation rates were observed where pronounced instabilities were detected by the radiosonde.

Moreover, measured shear instabilities and associated enhancements in dissipation on scales resolved by WRF also coincide with enlarged model turbulent kinetic energies (TKE). For instance, during the BEXUS 12 flight (27 Sep 2011), a wind reversal was observed which caused a large shear instability (indicated by Richardson numbers smaller than 1/4) as well as potential wave filtering. The resulting turbulence was detected by LITOS as a region with large dissipation rates. The model turbulent kinetic energy (TKE) peaks in this region, highlighting the significance of that layer. Similar effects are observed for some strong layers of the 27 Mar 2014 and 11/12 Jul 2015 flights. Thus, in these cases the geophysical causes of the observed turbulent layers are clearly visible. The large scale instabilities are resolved by the radiosondes and the model. On the other hand, many other (less intense) turbulent layers observed by LITOS are obviously too thin to be related to the much coarser data of the radiosonde or the WRF results.

Another relation between turbulence detected by LITOS and the presence of wave-like structures in WRF is noted: For the available summer flights at 06 Jun 2014 (not shown) and 12 Jul 2015, a drop in turbulence occurrence at approximately 15 km altitude with hardly any turbulence above was observed. In the associated model simulations, wave signatures become weaker around 15 km. Altogether, observed dissipation is weaker during lower wave activity (as seen in WRF), and larger where larger wave amplitudes are seen. These findings can be explained by wave saturation, while a change in, e. g., static stability is less prominent.

Turbulence has been observed for Richardson numbers below as well as above the critical number of 1/4, partly even for values much larger than 1. Such a violation of the classical theory by Miles (1961) and Howard (1961) has already been described by several researchers, e. g. Achatz (2005); Galperin et al. (2007); Balsley et al. (2008). Hines (1988) recognised the limitation of considering only vertical instability (as done when using the Richardson number) and proposed a concept of slantwise instabilities as created by gravity waves. He showed that turbulence is more likely to develop via slanted instability compared to vertical instability. Thus turbulence for $Ri > 1/4$ is comprehensible.

The results are based on the limited dataset from a few flights. More flights at selected meteorological situations are planned to further study the relation between waves and turbulence. A redesign of the instrumental setup shall eliminate the wake effects of balloon and ropes. Moreover, a direct measurement of gravity wave activity in combination to the turbulence observations is preferable.

Appendix A: Derivation of the constant c_{l_0} in Equation (1)

To retrieve energy dissipation rates from observed spectra, relation (1) between inner scale l_0 and dissipation rate ε ,

$\varepsilon = c_{l_0}^4 v^3 / l_0^4$, and especially the value of the constant c_{l_0} is important. To obtain correct values, care has to be taken of which component(s) of the spectral tensor are observed. In the following, the derivation of the constant c_{l_0} is summarised.

In the inertial subrange, the longitudinal component, transversal component, and trace of the structure function tensor for velocity fluctuations have the form

$$D_{xx}(r) = C_{xx} r^{2/3}, \quad (A1)$$

where xx is a placeholder for rr (longitudinal), tt (transversal), or ii (trace), and the structure constant has the form $C_{xx} = b_{xx} a_v^2 \varepsilon^{2/3}$ with $b_{rr} = 1$, $b_{tt} = \frac{4}{3}$, $b_{ii} = b_{rr} + 2b_{tt} = \frac{11}{3}$ (Tatarskii, 1971, p. 54ff) and the empirical constant $a_v^2 = 2.0$ (e. g. Pope, 2000, p. 193f). In the viscous subrange, the structure function is

$$D_{xx}(r) = \tilde{C}_{xx} r^2 \quad (A2)$$

with $\tilde{C}_{xx} = c_{xx} \frac{\varepsilon}{v}$ and the factors $c_{rr} = \frac{1}{15}$, $c_{tt} = \frac{2}{15}$, $c_{ii} = c_{rr} + 2c_{tt} = \frac{1}{3}$ (Tatarskii, 1971, p. 49).

Based on Heisenberg (1948, (28)), Lübken and Hillert (1992, (4)) gave a form of the temporal spectrum in the inertial and viscous subranges, which reads for velocity fluctuations

$$W(\omega) = \frac{\Gamma(\frac{5}{3}) \sin(\frac{\pi}{3})}{2\pi u_b} C_{xx} \frac{(\omega/u_b)^{-5/3}}{(1 + (\omega/u_b)^{8/3})^2} \quad (A3)$$

where u_b is the ascent velocity of the balloon, $\Gamma(z) := \int_0^\infty t^{z-1} e^{-t} dt$ is the Gamma function, and k_0 denotes the breakpoint between inertial and viscous subrange. The normalisation is obtained by considering the limit $k \ll k_0$ for the inertial subrange. Using the relation $\Phi(k) = -\frac{u_b^2}{2\pi k} \frac{dW}{d\omega}(ku_b)$ between temporal and spatial spectrum (Tatarskii, 1971, (6.14)), the corresponding three-dimensional spectrum is

$$\Phi_{xx}(k) = \frac{1}{6\pi} \frac{\Gamma(\frac{5}{3}) \sin(\frac{\pi}{3})}{2\pi} C_{xx} k^{-11/3} \frac{5 + 21(\frac{k}{k_0})^{8/3}}{(1 + (\frac{k}{k_0})^{8/3})^3}. \quad (A4)$$

The constant c_{l_0} in (1) can be computed from the condition of the structure function at the origin

$$\frac{d^2 D_{xx}}{dr^2}(0) = \frac{8\pi}{3} \int_0^\infty \Phi_{xx}(k) k^4 dk \quad (A5)$$

(Tatarskii, 1971, p. 49f). Inserting the structure function (A2) and the spectrum (A4) into condition (A5), integrating and solving for $1/k_0$ yields

$$l_0 = \frac{2\pi}{k_0} = 2\pi \underbrace{\left(\frac{3}{16} \Gamma(5/3) \sin(\pi/3) \frac{b_{xx}}{c_{xx}} a_v^2 \right)^{3/4}}_{=c_{l_0}} \left(\frac{v^3}{\varepsilon} \right)^{1/4}.$$

(A6)

CTA wire probes are sensitive perpendicular to the wire axis but insensitive parallel to the wire axis. For the earlier flights, the wires of the CTA sensors were oriented vertically so that they are sensitive in both horizontal directions and insensitive in the vertical direction, i.e. for an ascending balloon both transversal components are measured. Thus $b_{xx} = 4/3 + 4/3 = 8/3$ and $c_{xx} = 2/15 + 2/15 = 4/15$, which leads to $c_{l_0} = 14.1$. For the flight at 12 Jul 2015, one sensor with the wire oriented horizontally was flown, which is sensitive in the vertical and one horizontal direction yet insensitive in the other horizontal direction (parallel to the wire). In this case $b_{xx} = 1 + 4/3 = 7/3$ and $c_{xx} = 1/15 + 2/15 = 3/15$ so that $c_{l_0} = 15.8$.

Haack et al. (2014, Section 4) used different components of the structure function constant yielding $c_{l_0} = 5.7$. Since in (1) the constant occurs with $c_{l_0}^4$, this results in a difference in ε of a factor of ~ 50 for the same l_0 .

Acknowledgements. The BEXUS programme was financed by the German Aerospace Center (DLR) and the Swedish National Space Board (SNSB). We are grateful for the support by the “International Leibniz Graduate School for Gravity Waves and Turbulence in the Atmosphere and Ocean” (ILWAO) funded by the Leibniz Association (WGL). This study was partly funded by the German Federal Ministry for Education and Research (BMBF) research initiative “Role of the Middle Atmosphere In Climate” (ROMIC) under project numbers 01LG1206A and 01LG1218A (METROSI), and by the German Research Foundation (DFG) under project numbers LU 1174 (PACOG) and FOR 1898 (MS-GWaves). We thank Wayne K. Hocking and two anonymous reviewers for their valuable comments leading to the improvement of this article. The publication of this article was funded by the Open Access Fund of the Leibniz Association.

References

- Achatz, U.: On the role of optimal perturbations in the instability of monochromatic gravity waves, *Phys. Fluids*, 17, doi:10.1063/1.2046709, 2005.
- Andreassen, O., Wasberg, C. E., Fritts, D. C., and Isler, J. R.: Gravity wave breaking in two and three dimensions: 1. Model description and comparison of two-dimensional evolutions, *J. Geophys. Res.*, 99, 8095–8108, doi:10.1029/93JD03435, 1994.
- Balsley, B. B., Svensson, G., and Tjernström, M.: On the Scale-dependence of the Gradient Richardson Number in the Residual Layer, *Bound.-Layer Meteor.*, 127, 57–72, doi:10.1007/s10546-007-9251-0, 2008.
- Barat, J.: Some characteristics of clear-air turbulence in the middle stratosphere, *J. Atmos. Sci.*, 39, 2553–2564, doi:10.1175/1520-0469(1982)039<2553:SCOCAT>2.0.CO;2, 1982.
- Barat, J., Cot, C., and Sidi, C.: On the measurement of turbulence dissipation rate from rising balloons, *J. Atmos. Oceanic Technol.*, 1, 270–275, 1984.
- Birner, T.: Fine-scale structure of the extratropical tropopause region, *J. Geophys. Res.*, 111, D04104, doi:10.1029/2005JD006301, 2006.
- Birner, T., Dörnbrack, A., and Schumann, U.: How sharp is the tropopause at midlatitudes?, *Geophys. Res. Lett.*, 29, 45–1–45–4, doi:10.1029/2002GL015142, 2002.
- Brasseur, G. and Solomon, S.: *Aeronomy of the middle atmosphere: chemistry and physics of the stratosphere and mesosphere*, Atmospheric sciences library, Reidel, Dordrecht, 2nd edn., 1986.
- Chen, F. and Dudhia, J.: Coupling an Advanced Land Surface–Hydrology Model with the Penn State-NCAR MM5 Modeling System. Part I: Model Implementation and Sensitivity, *Mon. Wea. Rev.*, 129, 569–585, doi:10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2, 2001.
- Cho, J. Y. N., Newell, R. E., Anderson, B. E., Barrick, J. D. W., and Thornhill, K. L.: Characterizations of tropospheric turbulence and stability layers from aircraft observations, *J. Geophys. Res.*, 108, doi:10.1029/2002JD002820, 8784, 2003.
- Chou, M. D. and Suarez, M. J.: An efficient thermal infrared radiation parameterization for use in general circulation models., NASA Tech. Memo., 104606, 85pp., 1994.
- Clayson, C. A. and Kantha, L.: On Turbulence and Mixing in the Free Atmosphere Inferred from High-Resolution Soundings, *J. Atmos. Oceanic Technol.*, 25, 833–852, doi:10.1175/2007JTECHA992.1, 2008.
- Cot, C. and Barat, J.: Wave-turbulence interaction in the stratosphere: A case study, *J. Geophys. Res.*, 91, 2749–2756, doi:10.1029/JD091iD02p02749, 1986.
- Dalaudier, F., Sidi, C., Crochet, M., and Vernin, J.: Direct Evidence of “Sheets” in the Atmospheric Temperature Field, *J. Atmos. Sci.*, 51, 237–248, doi:10.1175/1520-0469(1994)051<0237:DEOITA>2.0.CO;2, 1994.
- Ehard, B., Achtert, P., Dörnbrack, A., Gisinger, S., Gumbel, J., Khamplanov, M., Rapp, M., and Wagner, J. S.: Combination of lidar and model data for studying deep gravity wave propagation, *Mon. Wea. Rev.*, 144, 77–98, doi:10.1175/MWR-D-14-00405.1, 2016.
- Franke, P. M. and Collins, R. L.: Evidence of gravity wave breaking in lidar data from the mesopause region, *Geophys. Res. Lett.*, 30, doi:10.1029/2001GL014477, 1155, 2003.
- Fritts, D. C. and Alexander, M. J.: Gravity wave dynamics and effects in the middle atmosphere, *Rev. Geophys.*, 41, doi:10.1029/2001RG000106, 2003.
- Fritts, D. C. and Wang, L.: Gravity Wave–Fine Structure Interactions. Part II: Energy Dissipation Evolutions, Statistics, and Implications, *J. Atmos. Sci.*, 70, 3735–3755, doi:10.1175/JAS-D-13-059.1, 2013.
- Fritts, D. C., Wang, L., Geller, M. A., Lawrence, D. A., Werne, J., and Balsley, B. B.: Numerical Modeling of Multiscale Dynamics at a High Reynolds Number: Instabilities, Turbulence, and an Assessment of Ozmidov and Thorpe Scales, *J. Atmos. Sci.*, 73, 555–578, doi:10.1175/JAS-D-14-0343.1, 2016.
- Galperin, B., Sukoriansky, S., and Anderson, P. S.: On the critical Richardson number in stably stratified turbulence, *Atmos. Sci. Lett.*, 8, 65–69, doi:10.1002/asl.153, 2007.
- Gavrilov, N. M.: Estimates of turbulent diffusivities and energy dissipation rates from satellite measurements of spectra of stratospheric refractivity perturbations, *Atmos. Chem. Phys.*, 13, 12 107–12 116, doi:10.5194/acp-13-12107-2013, 2013.

- Haack, A., Gerding, M., and Lübken, F.-J.: Characteristics of stratospheric turbulent layers measured by LITOS and their relation to the Richardson number, *J. Geophys. Res.*, 119, 10 605–10 618, doi:10.1002/2013JD021008, 2014.
- Hauf, T.: Aircraft Observation of Convection Waves over Southern Germany—A Case Study, *Mon. Wea. Rev.*, 121, 3282–3290, doi:10.1175/1520-0493(1993)121<3282:AOCWO>2.0.CO;2, 1993.
- Heisenberg, W.: Zur statistischen Theorie der Turbulenz, *Z. Phys.*, 124, 628–657, doi:10.1007/BF01668899, 1948.
- Hines, C. O.: Generation of Turbulence by Atmospheric Gravity Waves, *J. Atmos. Sci.*, 45, 1269–1278, doi:10.1175/1520-0469(1988)045<1269:GOTBAG>2.0.CO;2, 1988.
- Hines, C. O.: The Saturation of Gravity Waves in the Middle Atmosphere. Part I: Critique of Linear-Instability Theory, *J. Atmos. Sci.*, 48, 1348–1360, doi:10.1175/1520-0469(1991)048<1348:TSOGWI>2.0.CO;2, 1991.
- Hocking, W. K.: A review of Mesosphere–Stratosphere–Troposphere (MST) radar developments and studies, circa 1997–2008, *Journal of Atmospheric and Solar-Terrestrial Physics*, 73, 848–882, doi:http://dx.doi.org/10.1016/j.jastp.2010.12.009, 2011.
- Hodges, R. R.: Generation of turbulence in the upper atmosphere by internal gravity waves, *J. Geophys. Res.*, 72, 3455–3458, doi:10.1029/JZ072i013p03455, 1967.
- Hong, S.-Y. and Lim, J.-O. J.: The WRF single-moment 6-class microphysics scheme (WSM6), *J. Korean Meteor. Soc.*, 42, 129–151, 2006.
- Howard, L. N.: Note on a paper of John W. Miles, *J. Fluid Mech.*, 10, 509–512, doi:10.1017/S0022112061000317, 1961.
- Kain, J. S. and Fritsch, J. M.: A One-Dimensional Entrainment/Detraining Plume Model and Its Application in Convective Parameterization, *J. Atmos. Sci.*, 47, 2784–2802, doi:10.1175/1520-0469(1990)047<2784:AODEPM>2.0.CO;2, 1990.
- Klemp, J. B., Dudhia, J., and Hassiotis, A. D.: An Upper Gravity-Wave Absorbing Layer for NWP Applications, *Mon. Wea. Rev.*, 136, 3987–4004, doi:10.1175/2008MWR2596.1, 2008.
- Lilly, D. K., Waco, D. E., and Adelfang, S. I.: Stratospheric Mixing Estimated from High-Altitude Turbulence Measurements, *J. Appl. Meteor.*, 13, 488–493, doi:10.1175/1520-0450(1974)013<0488:SMEFHA>2.0.CO;2, 1974.
- Lindzen, R. S.: Turbulence and stress owing to gravity wave and tidal breakdown, *J. Geophys. Res.*, 86, 9707–9714, doi:10.1029/JC086iC10p09707, 1981.
- Lübken, F.-J. and Hillert, W.: Measurements of turbulent energy dissipation rates applying spectral models, in: *Coupling Processes in the Lower and Middle Atmosphere*, pp. 345–351, NATO Advanced Research Workshop, Kluwer Press, Loen, Norway, 1992.
- Luce, H., Fukao, S., Dalaudier, F., and Crochet, M.: Strong Mixing Events Observed near the Tropopause with the MU Radar and High-Resolution Balloon Techniques, *J. Atmos. Sci.*, 59, 2885–2896, doi:10.1175/1520-0469(2002)059<2885:SMEONT>2.0.CO;2, 2002.
- Miles, J. W.: On the stability of heterogeneous shear flows, *J. Fluid Mech.*, 10, 496–508, doi:10.1017/S0022112061000305, 1961.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.*, 102, 16 663–16 682, doi:10.1029/97JD00237, 1997.
- Nakanishi, M. and Niino, H.: Development of an Improved Turbulence Closure Model for the Atmospheric Boundary Layer, *J. Meteor. Soc. Japan*, 87, 895–912, http://ci.nii.ac.jp/naid/110007465760/en/, 2009.
- Nappo, C. J.: An Introduction to Atmospheric Gravity Waves, vol. 85 of *International Geophysics Series*, Academic Press, San Diego, 2002.
- Osman, M., Hocking, W., and Tarasick, D.: Parameterization of large-scale turbulent diffusion in the presence of both well-mixed and weakly mixed patchy layers, *Journal of Atmospheric and Solar-Terrestrial Physics*, 143–144, 14–36, doi:http://dx.doi.org/10.1016/j.jastp.2016.02.025, http://www.sciencedirect.com/science/article/pii/S1364682616300645, 2016.
- Paoli, R., Thouren, O., Escobar, J., Picot, J., and Cariolle, D.: High-resolution large-eddy simulations of stably stratified flows: application to subkilometer-scale turbulence in the upper troposphere–lower stratosphere, *Atmos. Chem. Phys.*, 14, 5037–5055, doi:10.5194/acp-14-5037-2014, 2014.
- Pavelin, E., Whiteway, J. A., and Vaughan, G.: Observation of gravity wave generation and breaking in the lowermost stratosphere, *J. Geophys. Res.*, 106, 5173–5179, doi:10.1029/2000JD900480, 2001.
- Pope, S. B.: *Turbulent Flows*, Cambridge University Press, Cambridge, 2000.
- Schneider, A., Gerding, M., and Lübken, F.-J.: Comparing turbulent parameters obtained from LITOS and radiosonde measurements, *Atmos. Chem. Phys.*, 15, 2159–2166, doi:10.5194/acp-15-2159-2015, 2015.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Powers, J. G.: A description of the Advanced Research WRF Version 3, NCAR technical note, Mesoscale and Microscale Meteorology Division, National Center for Atmospheric Research, Boulder, Colorado, USA, http://www2.mmm.ucar.edu/wrf/users/docs/arw_v3.pdf, 2008.
- Tatarskii, V. I.: The effects of the turbulent atmosphere on wave propagation, Israel Program for Scientific Translations, Jerusalem, translated from Russian, 1971.
- Theuerkauf, A., Gerding, M., and Lübken, F.-J.: LITOS – a new balloon-borne instrument for fine-scale turbulence soundings in the stratosphere, *Atmos. Meas. Tech.*, 4, 55–66, doi:10.5194/amt-4-55-2011, 2011.
- Wilson, R.: Turbulent diffusivity in the free atmosphere inferred from MST radar measurements: a review, *Ann. Geophys.*, 22, 3869–3887, doi:10.5194/angeo-22-3869-2004, 2004.
- Wilson, R., Luce, H., Hashiguchi, H., Nishi, N., and Yabuki, Y.: Energetics of persistent turbulent layers underneath mid-level clouds estimated from concurrent radar and radiosonde data, *J. Atmos. Sol.-Terr. Phys.*, 118, Part A, 78–89, doi:10.1016/j.jastp.2014.01.005, 2014.
- Worthington, R. M.: Tropopausal turbulence caused by the breaking of mountain waves, *J. Atmos. Sol.-Terr. Phys.*, 60, 1543–1547, doi:10.1016/S1364-6826(98)00105-9, 1998.
- Yamanaka, M. D., Tanaka, H., Hirose, H., Matsuzaka, Y., Yamagami, T., and Nishimura, J.: Measurement of Stratospheric Turbulence by Balloon-Borne “Glow-Discharge” Anemometer,

J. Meteor. Soc. Japan Ser. II, 63, 483–489, https://www.jstage.jst.go.jp/article/jmsj1965/63/3/63_3_483/_article, 1985.