



## Resolution dependence of uncertainties in gridded emission inventories: a case study in Hebei, China

Bo Zheng<sup>1</sup>, Qiang Zhang<sup>2</sup>, Dan Tong<sup>2</sup>, Chuchu Chen<sup>2</sup>, Chaopeng Hong<sup>2</sup>, Meng Li<sup>2</sup>, Guannan Geng<sup>2</sup>, Yu Lei<sup>3</sup>, Hong Huo<sup>4</sup>, Kebin He<sup>1,5</sup>

5 <sup>1</sup>State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing 100084, China

<sup>2</sup>Ministry of Education Key Laboratory for Earth System Modeling, Center for Earth System Science, Tsinghua University, Beijing 100084, China

<sup>3</sup>The Atmospheric Environment Department, Chinese Academy for Environmental Planning, Beijing 100012, China

10 <sup>4</sup>Institute of Energy, Environment and Economy, Tsinghua University, Beijing 100084, China

<sup>5</sup>State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex, Beijing 100084, China

*Correspondence to:* Qiang Zhang (qiangzhang@tsinghua.edu.cn) and Kebin He (hekb@tsinghua.edu.cn)

**Abstract.** Gridded emission inventories are essential inputs for chemical transport models and climate models. Spatial proxies are applied to allocate emissions from regional totals to spatially resolved grids when the exact locations of emissions are absent, with additional uncertainties arising due to the spatial mismatch between the locations of emissions and spatial proxies. In this study, we investigate the impact of spatial proxies on the accuracy of gridded emission inventories at different spatial resolutions by comparing gridded emissions developed from different spatial proxies (proxy-based inventory) with a highly spatially disaggregated bottom-up emission inventory developed from the extensive use of locations of emitting facilities (bottom-up inventory) in Hebei Province, China. We find that proxy-based inventories are generally comparable to bottom-up inventories for grid sizes larger than 0.25 °because spatial errors are largely diminished at coarse resolutions. However, for gridded emissions with finer resolutions, large positive biases in urban centers and negative biases in suburban and rural regions are identified in proxy-based inventories and are then propagated into significant biases in urban-scale chemical transport modeling. Compared to bottom-up inventories, the use of proxy-based emissions exhibits similar modeling results, with biases varying from 3%–13% in predicting surface concentrations of different pollutants at 36 km resolution and an additional 8%–73% at 4 km resolution. The resolution dependence of uncertainties in proxy-based gridded inventories can be explained by the decoupling of emission facility locations from spatial surrogates, especially because industry facilities tend to be located away from urban centers. This distance results in a divergence between emission distributions and the allocation of proxies on smaller grids. The decoupling effects are weakened when the grid size increases to cover both urban and rural regions. We conclude that proxy-based inventories are of sufficient quality to support regional and global models (larger than 0.25 ° in this case study); however, to support urban-scale models with accurate emission inputs, bottom-up inventories incorporating the exact locations of emitting facilities should be developed instead of proxy-based inventories.

15  
20  
25  
30



## 1 Introduction

35 Gridded emission inventories have emerged as a critical component of atmospheric chemistry and climate models. The importance of these inventories has been driven by the advent of regional chemical transport models at different spatial scales. As the key inputs, spatial representations of emission estimates containing errors can be propagated into modeled concentrations, affecting subsequent studies based on those chemical transport models.

Gridded emission inventories can be developed using two methods: bottom-up estimates and proxy-based estimates with downscaling techniques. Bottom-up estimates rely on massive spatial information such as point sources, detailed censuses, and traffic statistics. For large point sources, emissions can be estimated for each individual facility and then mapped at high 40 resolutions (e.g., Zhao et al., 2008; Liu et al., 2015). For on-road emissions, high-resolution mapping can be achieved using road-specific traffic count data at spatial scales ranging from 500 m to 10 km (e.g., Gately et al., 2013, 2015; McDonald et al., 2014). The above bottom-up inventory methods rely on mass data inputs, which are accurate but difficult to extend through all sectors and regions. For example, Gurney et al. (2009) developed the Vulcan inventory to quantify fossil fuel 45 CO<sub>2</sub> emissions for the contiguous U.S. at a resolution of 10 km. This data product was built upon the best available data sources through all sectors throughout the U.S., including point, nonpoint and airport datasets with additional emission monitoring data from individual facilities.

The proxy-based method relies on spatial proxies to build emission inventories in gridded form. Because they require relatively less data, proxy-based approaches are widely used for developing gridded emission inventories. Proxies such as 50 population and nighttime lights have been used to derive gridded emissions (e.g., Raupach et al., 2010; Oda et al., 2011; Wang et al., 2013), which are implicitly assumed to be resolution-independent at sizes ranging from district level to grid cells. However, this correlation is likely sensitive to fine-scale spatial resolutions, which introduce uncertainties in high-resolution emission mapping. Gurney et al. (2009) highlighted the spatial biases inherent in a population-based gridded emission inventory due to the decoupling of emissions and population at 0.1 °. Oda et al. (2011) presented the uncertainties in 55 a nighttime light-based inventory due to the saturation errors in nighttime light data at 1 km. Gately et al. (2013, 2015) found that the spatial correlation between the per-mile CO<sub>2</sub> emissions of motor vehicles and population density changed from positive to negative when population density increased (cut-off point of approximately 2000 persons km<sup>-2</sup>). The above studies suggest that nonlinearities exist between emissions and spatial proxies at fine scales. The downscaling method with fixed correlation can involve large uncertainties in proxy-based gridded emissions, especially at high resolutions.

60 The uncertainties in proxy-based gridded emission inventories are sensitive to spatial resolution. Recent efforts suggest that the spatial errors of proxy-based CO<sub>2</sub> emission inventories tend to increase as spatial resolutions rise (e.g., Gurney et al., 2009; Rayner et al., 2010; Oda et al., 2011; Wang et al., 2013; Asefi-Najafabady et al., 2014) because spatial proxies are decoupled from emissions at fine resolutions; however, the influence of uncertainties in gridded emission inventories of air pollutants and their propagation in atmospheric chemistry models are not considered. This shortcoming hampers high-



65 resolution air pollution modeling when proxy-based inventories are used because the grid size suitable for constraining the uncertainties is unknown.

This paper aims to investigate how resolutions influence uncertainties in gridded emission inventories of air pollutants and the subsequent atmospheric chemistry modeling. The spatial resolution dependence of uncertainties in gridded emission inventory is quantified using multi-resolution emission inventories, a chemical transport model, and *in situ* measurements.

70 We use Hebei Province in China, where a detailed bottom-up emission inventory is available. We first develop gridded emission inventories for Hebei at the resolution of 1 km using proxy-based and bottom-up methods and then compare the two datasets at multiple aggregated spatial resolutions. The Community Multi-scale Air Quality model (CMAQ) is driven by the two gridded emission datasets to explore how uncertainties affect the performance in predicting the surface concentrations of different air pollutants.

## 75 2 Methodology

### 2.1 Gridded emission inventories

#### 2.1.1 Bottom-up inventory

We used the bottom-up method to develop a high-resolution emission inventory for Hebei Province in 2013 (denoted by HB-EI). The emission sources were subdivided into more than 700 sector/fuel (or product)/technology combinations of source categories and aggregated into five sectors (power, industry, residential, transportation, and agriculture). Each source category was classified as point, nonpoint or mobile, with a different emission accounting method applied to each category. Point sources were stationary emitting sources inventoried at a facility level. The power and industry sectors were treated as point sources, from which the emissions were calculated using the following:

$$Emis_{i,s} = A_i \times EF_{i,s} \times \prod_n (1 - \eta_{i,n,s}), \quad (1)$$

85 where  $i$  represents the emitting facility,  $s$  represents air pollutants (i.e., SO<sub>2</sub>, NO<sub>x</sub>, VOCs, NH<sub>3</sub>, CO, CO<sub>2</sub>, PM<sub>2.5</sub>, BC, OC, PM<sub>10</sub> and TSP (particulate matter with aerodynamic diameter of 100 μm or less)),  $A$  is the activity rate,  $EF$  is the unabated emission factor,  $n$  represents an air pollution control device, and  $\eta_{i,n,s}$  is the removal efficiency of pollutant  $s$  by control device  $n$  installed at facility  $i$ . The point sources were located according to their latitudinal and longitudinal coordinates. The locations of large point sources were checked and corrected by visual inspection in Google Maps. Large point sources included power and heating plants, large industrial boilers ( $\geq 24.5$  MW), and manufacturing factories of coke, iron, steel, cement and flat glass. These sites constituted 90% of the energy demand from all the point sources. For the other point sources, the coordinates of registered addresses were used directly.



Nonpoint or area sources were stationary emitting sources inventoried at a province level, from which emissions exit from diffuse sources without identifiable stacks. The residential, non-road transportation and agriculture sectors were estimated as nonpoint sources using the following:

$$Emis_{j,k,m,s} = A_{j,k} \times X_{j,k,m} \times EF_{j,k,m,s} \times \sum_n (C_{j,k,m,n} \times (1 - \eta_{n,s})), \quad (2)$$

where  $j$  represents sector,  $k$  represents fuel or product,  $m$  represents technology,  $X$  is the fraction of activity rates contributed by a specific technology,  $C$  is the penetration of a specific pollution-control technology, and the other parameters are the same as in Eq. (1). The nonpoint sources were allocated to a  $30'' \times 30''$  grid in two steps. First, the provincial emission totals were distributed to each county based on county-level activity statistics. For example, we used the residential coal and biofuel use of each county to allocate residential emissions. Second, county emissions were allocated to grids based on spatial proxies, such as urban or rural extents (Schneider et al., 2009) and population (Oak Ridge National Laboratory, 2013). The parameters used in nonpoint source emission distributions are summarized in Table 1.

Mobile sources referred to the on-road transportation sector, which were estimated using the method established by Zheng et al. (2014). The county-specific vehicle activity and emission factors were simulated and multiplied to calculate county-level vehicle emissions. County emissions were downscaled to a  $30'' \times 30''$  grid using a GIS road atlas and traffic flow statistics specific to different vehicle and road types (Zheng et al., 2014).

### 2.1.2 Proxy-based inventory

We used the data from the Multi-resolution Emission Inventory for China (MEIC) as the proxy-based estimate. MEIC is a technology-based emission model framework developed by Tsinghua University (<http://www.meicmodel.org>). This model was built on the foundation laid by the same group responsible for the present study (e.g., Zhang et al., 2007, 2009; Lei et al., 2011), with high-resolution mapping of emissions for power plants (Liu et al., 2015) and on-road vehicles (Zheng et al., 2014).

The emission source categorization in MEIC is the same as that in HB-EI. Emissions of power plants were estimated for each unit using Eq. (1). The on-road transportation sector was estimated following the method established by Zheng et al. (2014) as the bottom-up inventory. The industrial, residential, non-road transportation and agricultural sectors were estimated as nonpoint sources using Eq. (2) with activity rates and emission factors at the provincial level. Emission totals from nonpoint sources were allocated to grid cells using different spatial proxies. The proxies used for MEIC are also presented in Table 1 and compared with HB-EI. For example, in MEIC, the residential sector used county population to split provincial emissions by county instead of residential energy consumption used in HB-EI.

In this work, we used Hebei emissions from 2013 from MEIC v1.2 and scaled the emission magnitude to match the HB-EI inventory by source category. This approach can support the magnitude-independent comparison to reflect the discrepancies in the spatial distribution of gridded emissions. Multi-resolution gridded emissions aggregated from the  $30'' \times 30''$  grids were used to assess the resolution dependence of uncertainties in the gridded emission inventory.



## 125 2.2 Chemical transport model

### 2.2.1 Model configuration

The WRF/CMAQ system was used to evaluate emission inventories and assess how uncertainties in gridded emission inventory influenced atmospheric chemical modeling. CMAQ v5.0.1 (<http://www.cmascenter.org/cmaq/>) was applied in this work, which was driven by assimilated meteorological fields from the Weather Research and Forecasting model v3.5.1 (<http://www.wrf-model.org/>). The model configurations were determined following the method established in our previous work (Zheng et al., 2015). We used the updated carbon bond gas-phase mechanism with updated toluene mechanism (Whitten et al., 2010), aerosol module 6 (AERO6) and ISORROPIA v2.1 inorganic chemistry (Fountoukis and Nenes, 2007). The aqueous-phase chemistry method used in this study was the updated mechanism of the Regional Acid Deposition Model (Walcek and Taylor, 1986; Chang et al., 1987). Photolytic rates were calculated in line using simulated aerosol and ozone concentrations. The ACM methodology was adopted in the cloud module to compute convective mixing for aerosols. The above configurations were evaluated in Zheng et al. (2015).

Anthropogenic emissions outside Hebei Province were taken from MEIC for China (<http://www.meicmodel.org>) and MIX for the other Asian countries (Li et al., 2015). Emission inputs were processed on simulation domains from their native resolutions ( $30\text{ s} \times 30\text{ s}$  for MEIC and  $0.25^\circ \times 0.25^\circ$  for MIX). Other emissions, such as biomass burning, sea salt and biogenic VOCs, were taken from various models and datasets following the method established by Zheng et al. (2015). The boundary and initial conditions were processed from the GEOS-Chem model output (Bey et al., 2001) using the tool developed by Henderson et al. (2014).

### 2.2.2 Simulation design

Two full-year simulations of 2013 were conducted using MEIC (denoted by S1) and HB-EI (denoted by S2), both at triple-nested domains (36, 12 and 4 km), with the finest resolution focusing on Hebei Province (Fig. 1). We also conducted a sensitivity simulation (denoted by S3) for Jan, Apr, Jul and Oct using the adjusted HB-EI emissions by changing ~20% point sources into nonpoint source estimates to assess the influence of highly spatially resolved emission sources on atmospheric chemistry modeling. We aggregated the emissions from small industrial boilers ( $\leq 24.5\text{ MW}$ ) into provincial totals and then distributed the totals onto grid cells using the same spatial proxies as those used in the industrial sector in MEIC. The model performances were evaluated against ground-based measurements. We compared the annual daily mean of simulation and observation for the criteria air pollutants and calculated mean bias (MB) and normalized mean bias (NMB) to evaluate modeling results.

## 2.3 *In situ* measurements

The ground-based observations in Hebei were acquired from the China National Environmental Monitoring Center (<http://106.37.208.233:20035/>), which published hourly concentrations of  $\text{SO}_2$ ,  $\text{NO}_2$ ,  $\text{CO}$ ,  $\text{O}_3$ ,  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  from 53



monitoring stations over Hebei in 2013. This dataset was built and maintained by the Ministry of Environmental Protection in China and is used as the official data for national air-quality management. The ground-based stations are primarily located in urban centers because they are designed to assess population exposures in the densest areas.

### 3 Results

#### 160 3.1 Emission inventories: bottom-up versus proxy-based

In the HB-EI inventory, we estimated the anthropogenic emissions of Hebei Province in 2013 as follows: 1.4 Tg SO<sub>2</sub>, 2.0 Tg NO<sub>x</sub>, 1.5 Tg VOCs, 0.5 Tg NH<sub>3</sub>, 16.8 Tg CO, 827.2 Tg CO<sub>2</sub>, 2.7 Tg TSP, 1.3 Tg PM<sub>10</sub>, 0.9 Tg PM<sub>2.5</sub>, 0.1 Tg BC and 0.2 Tg OC. The magnitudes of emissions in MEIC were scaled to match HB-EI by emission source; therefore, the discrepancies between MEIC and HB-EI were primarily attributed to the differences in spatial distributions. The MEIC inventory  
165 represented a mixture of data sources from downscaled province-level emissions and point sources (i.e., power plants), in which 25% SO<sub>2</sub>, NO<sub>x</sub> and CO<sub>2</sub> emissions were identified as point sources (Fig. 2a). Conversely, more than 70% emissions in the HB-EI inventory were inventoried as point sources (Fig. 2b), and the remaining were constrained by road- and county-level activities.

The discrepancy in emission shares of point sources varied by pollutant between MEIC and HB-EI. For pollutants that  
170 dominated emissions by industrial combustion and production (e.g., CO<sub>2</sub>, SO<sub>2</sub>, and NO<sub>x</sub>), HB-EI presented a much larger share of point source emissions than MEIC, while for the pollutants mainly emitted from diffuse sources (e.g., VOCs, NH<sub>3</sub>, BC, and OC), the two datasets presented similar contributions from point sources. This finding suggested that the nonlinearities in CO<sub>2</sub> emissions and spatial proxy covariance identified in early studies (e.g., Gurney et al., 2009; Rayner et al., 2010; Oda et al., 2011; Wang et al., 2013; Asefi-Najafabady et al., 2014) may have diverged for different air pollutants.

#### 175 3.2 Resolution dependence of uncertainties in the gridded emission inventory

Table 2 shows the comparison between the gridded emissions of the bottom-up inventory (HB-EI) and the proxy-based inventory (MEIC) over the Hebei region at different resolutions. Following the method established by Rayner et al. (2010), three metrics were adopted for comparison: (1) spatial correlation (R), which quantifies the correspondence of spatial patterns; (2) summed absolute difference (SAD), which is the sum of the absolute difference for the whole domain; and (3)  
180 relative summed absolute difference (RSAD), which is calculated as the SAD divided by total emissions over the domain. For the pollutants of which emissions are dominated from point sources, the two gridded inventories agreed well at coarse resolutions (grid size larger than 0.25 °), while the differences tended to increase when grid size decreased, indicating more spatial biases involved in proxy-based inventories at high spatial resolutions. For example, the resolutions of 0.05 ° and 0.1 ° produced normalized bias (RSAD) as large as 80%–100%, results that were much higher than the uncertainties in total  
185 emissions (e.g., 20%–40% for SO<sub>2</sub> and NO<sub>x</sub>). For the pollutants of which emissions are dominated by non-point sources (e.g., NH<sub>3</sub>), the two inventories agreed well regardless of spatial resolution because they shared the same spatial proxies.



Figure 3 compares the spatial distributions of gridded emissions using  $\text{NO}_x$  as an example. Figure 3(a–d) presents  $\text{NO}_x$  emission distributions in HB-EI at different spatial resolutions, and Fig. 3(e–h) presents the differences between the two inventories. At high resolutions ( $0.05^\circ$  and  $0.1^\circ$ ), MEIC tended to overestimate emissions in urban centers but underestimate emissions in rural areas, leading to unrealistically higher urban-rural emission gradients. Following economic development and air quality control progress in China, polluted industries have tended to move away from urban centers in large cities (see Fig. 4 for the largest two cities in Hebei), resulting in the divergence polluting industries from dense population distributions. Therefore, the use of population distribution to allocate emissions tended to overestimate the urban-rural gradients of emissions. At coarse resolutions of  $0.5^\circ$  and  $1.0^\circ$ , relative differences between the two inventories were smaller because the large grid covered both urban and rural regions and smoothed emission distributions.

To elucidate the variations in urban and rural areas, we analyzed the relationships between emission fluxes of  $\text{NO}_x$  and demographic trends at multiple resolutions, as shown in Fig. 5. In HB-EI, 32% of emissions were attributed to the top 25% of the populated dense grids, as shown by the line of HB-EI- $\text{NO}_x$ . Emission fluxes gradually declined, moving from the dense urban grids to the less dense suburban and rural grids, while emission fluxes in MEIC declined relatively sharply compared to HB-EI. This trend occurred because the proxy-based downscaling method tended to over-allocate industrial emissions to urban areas, such that 42% of  $\text{NO}_x$  emissions were distributed to populated grids in the upper quartile, 10% higher than HB-EI, resulting in larger urban-rural gradients. This pattern was evident at the finer scales given that the  $\text{NO}_x$  emission fluxes from the 10% most densely populated grids were 46%–140% higher than those of HB-EI at  $0.05^\circ$  and  $0.1^\circ$ . At coarse resolutions, the urban and rural areas were aggregated with evenly distributed emissions. Therefore, both HB-EI- $\text{NO}_x$  and MEIC- $\text{NO}_x$  approached the 1:1 line at  $0.5^\circ$  and  $1.0^\circ$  and exhibited similar emission flux patterns that were not influenced by spatial allocation biases.

Urban-scale models have emerged in response to a critical need for fine-scale modeling. However, the proxy-based downscaling method over-allocates emissions to urban centers, producing artificial emission hotspots that can result in biases in high-resolution models. When estimated as population-weighted exposures, the health risks induced by air pollution may also be overestimated due to the collocation of gridded emissions with dense populations. We evaluate how uncertainties in fine-scale gridded emissions influence air pollution modeling in the next section.

### 3.3 Resolution dependence of biases in air pollution modeling

The uncertainties in gridded emission inventory induced by the proxy-based method can affect the biases of chemical transport modeling. The bias propagation was quantified by comparing performance of the WRF-CMAQ model with the MEIC (S1) and HB-EI (S2) emission input. The comparison was conducted at 36 km, 12 km and 4 km resolutions, which roughly corresponded to the  $0.5^\circ$ – $0.05^\circ$  grid sizes discussed above.

For the densest urban areas, the finer-scale simulations predicted higher concentrations of air pollutants due to concentrated emissions (Fig. 6). This enhancement led to better agreement with the observations, as demonstrated by the S2 simulation moving from 36 km to 4 km. Higher concentrations modeled at finer scales were also observed in the S1 simulation, but S1





220 tended to introduce higher positive biases rather than better performances. This tendency was likely due to the over-allocated urban emissions produced by the proxy-based downscaling method. Compared to bottom-up inventories, the use of proxy-based emissions produced similar modeling results, with biases of 3%–13% in predicting surface concentrations of different pollutants at 36 km resolution. At 12 km and 4 km, the major pollutants modeled by proxy-based inventories were overestimated by 28%–114% and O<sub>3</sub> was underestimated by 17% due to the enhanced titration effect caused by concentrated

225 NO<sub>x</sub> emissions. The finer-scale modeling using proxy-based emissions introduced additional 8%–73% biases compared with bottom-up inventories. This finding suggested that urban-scale modeling efforts at ~10 km cannot achieve corresponding accuracies until factory-level inventories are used. Regarding the emission spatial biases (see Sect. 3.2), the proxy-based downscaled emissions were appropriate for modeling at the global and regional scales (e.g., 0.25 °–0.5 ° or 36 km in this case study), while they could cause larger biases for finer resolutions at the urban scale.

230 Figure S1 presents the spatial distribution of modeled surface concentrations for primary pollutants. The S1 simulation predicted higher NO<sub>2</sub> and SO<sub>2</sub> in the south-central province but lower concentrations in the southwest. The densest cities in Hebei are located in the former region, and the industrial district along Taihang Mountain is located in the latter. The proxy-based downscaling method allocated more emissions to the populated regions, resulting in higher estimates of air pollution levels in the densest cities. Compared to bottom-up emissions, the modeling surface concentrations of air pollutants in urban

235 areas were approximately 20%–50% higher for the 36 km domain and were much higher at 12 km and 4 km (see Fig. 7). The bias level was influenced by the size of cities. The cities with large populations and industries tended to experience rapid urbanization, with people gathering in the urban centers and polluted industries moving outward. The modeling discrepancies of the two largest cities in Hebei (i.e., Tangshan and Shijiazhuang) were as large as 100%–200%. The small cities presented scattered distributions for both populations and industries and exhibited smaller differences when using

240 population to distribute emissions.

Figure S2 presents the spatial distribution of modeled PM<sub>2.5</sub> and O<sub>3</sub> concentrations. The S1 simulation tended to predict an 18%–55% higher PM<sub>2.5</sub> in the densely populated regions but 30%–110% lower in the less dense regions. The differences were slightly smaller than those found in the primary pollutants because of the nonlinear particle formation processes and long-range regional transport. For O<sub>3</sub>, the enhanced titration by NO due to concentrated emissions resulted in negative biases.

245 The S1 simulation predicted 12%–30% lower O<sub>3</sub> in the dense regions but 16%–35% higher in the less dense regions. When estimated as population-weighted exposures and averaged within the areas with population density over 600 per km<sup>-2</sup> (see Fig. 7), the PM<sub>2.5</sub> exposures tended to be 10%–20% higher, whereas the O<sub>3</sub> exposures tended to be 10%–15% lower. In the urban centers, the population exposures to PM<sub>2.5</sub> and O<sub>3</sub> were biased at rates as high as 50%–100%. The overestimated exposures of PM<sub>2.5</sub> associated with the underestimated exposures of O<sub>3</sub> suggested that more effort is needed to evaluate air pollution

250 health risks at the urban scale. The uncertainties in gridded emissions comprised important uncertainties in health risk assessment.





## 4 Discussion

### 4.1 Sensitivity to different proxies

The proxy-based spatial allocation method assumed linear correlations between emission intensities and spatial proxy densities within a given district. To investigate the sensitivity of the assumption to spatial proxies, we evaluated the spatial correlations between the gridded emissions of HB-EI and different spatial proxies (see Fig. 8). We concluded that the spatial correlation was enhanced significantly with the increasing of grid size, which was not sensitive to spatial proxy type. The grid size of  $0.25^\circ$  roughly corresponded to the correlation coefficient of 0.5. At coarse resolutions (grid size larger than  $0.25^\circ$ ), good spatial correlations were found for different proxies across all pollutants. However, the correlation worsened when the spatial resolutions grew, except for pollutants dominated by nonpoint sources that were correlated with their allocating proxies (e.g.,  $\text{NH}_3$  vs. rural population). Due to the discrepancies of emission source compositions, the air pollutants presented different distribution patterns (see Fig. 8). Compared to nonpoint source emissions (e.g.,  $\text{NH}_3$ , VOCs, OC), the pollutants dominated by point sources (e.g., TSP,  $\text{CO}_2$ ,  $\text{SO}_2$ ) showed relatively poor correlations with spatial proxies. This result was not very sensitive to spatial proxy type. For example, changing allocators caused the correlation coefficient to vary only between 0.07 and 0.18 for TSP at  $0.05^\circ$ . This pattern arose because point sources have been increasingly sited away from urban areas, a phenomenon that was difficult to represent by spatial proxy distribution. In contrast, the pollutant distributions of nonpoint sources were sensitive to spatial proxies at resolutions finer than  $0.25^\circ$  because the pollutants allocated with one proxy were clearly correlated with this proxy but uncorrelated with other proxies at fine scales. However, all the proxies tended to present similar spatial distributions at large scales, demonstrated by the correlation coefficient of almost 1 at resolutions coarser than  $0.25^\circ$ . The above findings suggest that the underlying assumption inherent in the proxy-based emission allocation process is potentially valid at a coarse scale but is highly suspect at scales finer than  $0.25^\circ$ .

We evaluated the spatial correlations of HB-EI with different proxies by sector, as shown in Fig. 9.  $\text{NO}_x$  is shown as an example (the results for other pollutants are similar). Because the locations of point sources were decoupled from spatial proxies at fine resolutions, power plant and industrial emissions presented poor correlations ( $R^2 < 0.2$ ) with various spatial proxies when grid size was smaller than  $0.5^\circ$ . The decoupling effect was weakened when grids were aggregated to  $0.5^\circ$  or higher, and consistent spatial patterns were found between emissions from point sources and different spatial proxies. The residential and transportation emissions were sensitive to proxy selections at finer scales, which were affected by the allocating proxies used by the two sectors. When grids were aggregated to  $0.25^\circ$  or higher, these nonpoint source emissions showed consistent spatial distributions regardless of spatial proxies. For total emissions, the variation of spatial correlation between gridded emissions and proxies was narrowed for grids larger than  $0.25^\circ$ . Therefore, the proxy-based downscaling method tended to introduce large errors at scales finer than  $0.25^\circ$ , where local patterns in the distributed point sources dominated over diffuse nonpoint sources, which have not yet been reproduced by any spatial proxy at finer scales.



285 The above findings imply that spatial proxies should be used with caution. When building regional and global gridded emissions (typically larger than  $0.25^\circ$ ), the two proxies of total population and nighttime light performed slightly better than the other proxies did. These two proxies correlated well with gridded emissions, while other proxies such as urban population, rural population and road network had several limitations. Urban population tended to over-allocate emissions to urban areas. Rural population had poor spatial correlation with gridded emissions. The spatial distribution patterns of road network were not suitable for emission allocation except those from vehicles. When mapping emissions at higher resolutions  
290 (e.g., finer than  $0.25^\circ$ ), the proxies of total population and nighttime light were poorly correlated with gridded emissions dominated by point sources, and the spatial distributions of nonpoint sources emissions were very sensitive to spatial proxies. In this case, the bottom-up method must be used instead of the proxy-based method to improve the spatial representation of emission distributions.

#### 4.2 Sensitivity to point sources

295 The bottom-up method estimated emissions for each individual facility as a point source to improve the accuracy of inventory. The point source estimates constituted the main difference between MEIC and HB-EI, demonstrated by the discrepancy in point source shares (Fig. 2). Compared to MEIC, the distinct improvement of HB-EI was to treat all facilities from the industry sector as point sources in addition to a minor improvement in province-to-county emission allocation. These changes produced better agreements with *in situ* observations when used for urban-scale modeling. At 4 km resolution,  
300 the bottom-up inventory of HB-EI reduced biases by 8%–73% in predicting the surface concentrations of different air pollutants. To evaluate the sensitivity of fine-scale modeling to point source estimates, we conducted a sensitivity analysis by slightly reducing the contributions of point sources, an analysis denoted by S3 (see Sect. 2.2.2). The extent to which fine-scale modeling influenced the modeling performance highlighted the implications for point source estimates and the role high-resolution emission mapping played in fine-scale modeling.

305 For S3, we aggregated the emissions from small industrial boilers ( $\leq 24.5$  MW) into provincial totals and then distributed them onto grid cells using spatial proxies. The emission shares of point sources were reduced by ~20% for almost all pollutants (Table 3). The modeling biases that used this adjusted inventory (S3) fell between those from the MEIC-based simulation (S1) and HB-EI-based simulation (S2). The normalized mean biases of the criteria pollutants were approximately 5%–15% higher than those produced by S2 due to the inclusion of ~20% fewer point sources, whereas they were 18%–90%  
310 lower than those produced by S1 due to the inclusion of ~50% more point sources. This difference suggested that the spatial biases tended to decline significantly as more point sources were included. The improvement of modeling biases roughly corresponded to the increased contributions of point sources to emissions.



## 5 Concluding remarks

In this study, we assessed the resolution dependence of uncertainties in gridded emission inventories, using Hebei, China as  
315 a case. The inherent uncertainties involved in emission distributions caused systematic biases in both the emission flux  
patterns and subsequent chemical transport modeling. In the case of Hebei, China, the proxy-based downscaling method  
tended to over-allocate emissions to the urban center. For example, the NO<sub>x</sub> emission fluxes from the 10% most densely  
populated grids tended to be overestimated by 46%–140% at 0.1 ° and 0.05 ° in this case study. This effect was demonstrated  
by the modeling performance of the CMAQ model, in which the modeling biases of different pollutants using proxy-based  
320 inventory were 8%–73% higher than using bottom-up inventories at fine scale (12 km and 4 km). The modeling biases  
induced by uncertainties in gridded emissions caused population exposures of PM<sub>2.5</sub> to be overestimated and O<sub>3</sub> to be  
underestimated in urban areas.

In the proxy-based inventory, the inherent assumption of spatial correlation between emissions and allocating proxies was  
highly suspect at an urban scale (grid sizes smaller than 0.25 ° in this case study). This lack of validity was caused by  
325 polluted industries increasingly moving away from urban centers, a phenomenon that resulted in a decoupling of emissions  
from spatial proxies at finer scales. The gridded emissions at a coarse scale tended to aggregate urban, suburban and rural  
areas; smooth emission distributions; and weaken the decoupling effect. Extensive use of point sources could improve the  
accuracy of gridded emissions, demonstrated by the improvement of modeling performance as the contribution of point  
sources increased. We concluded that proxy-based inventories are capable of supporting regional and global models (larger  
330 than 0.25 ° in this case study); however, to support urban-scale models with accurate emission inputs, bottom-up inventories  
with exact locations of emitting facilities should be developed instead.

## Acknowledgements

This work was supported by China's National Basic Research Program (2014CB441301), the National Science Foundation  
of China (41222036 and 41571130032), the National Key Technology R&D Program (2014BAC16B03 and  
335 2014BAC21B02), and the public welfare program of China's Ministry of Environmental Protection (201509014).

## References

Amann, M., Derwent, D., Forsberg, B., Hänninen, O., Hurley, F., Krzyzanowski, M., Leeuw, F., Liu, S., Mandin, C.,  
Schneider, J., Schwarze, P., and Simpson, D.: Health risks of ozone from long-range transboundary air pollution, WHO  
Regional Office for Europe, [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0005/78647/E91843.pdf](http://www.euro.who.int/__data/assets/pdf_file/0005/78647/E91843.pdf) (last access: March,  
340 2016), 2008.



- Asefi-Najafabady, S., Rayner, P. J., Gurney, K. R., McRobert, A., Song, Y., Coltin, K., Huang, J., Elvidge, C., and Baugh, K.: A multiyear, global gridded fossil fuel CO<sub>2</sub> emission data product: Evaluation and analysis of results, *J. Geophys. Res-Atmos.*, 119, 10213-10231, 10.1002/2013JD021296, 2014.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res-Atmos.*, 106, 23073-23095, 10.1029/2001JD000807, 2001.
- Boersma, K. F., Eskes, H. J., Dirksen, R. J., van der A, R. J., Veefkind, J. P., Stammes, P., Huijnen, V., Kleipool, Q. L., Sneep, M., Claas, J., Leitão, J., Richter, A., Zhou, Y., and Brunner, D.: An improved tropospheric NO<sub>2</sub> column retrieval algorithm for the Ozone Monitoring Instrument, *Atmos. Meas. Tech.*, 4, 1905-1928, 10.5194/amt-4-1905-2011, 2011.
- 345 Chang, J. S., Brost, R. A., Isaksen, I. S. A., Madronich, S., Middleton, P., Stockwell, W. R., and Walcek, C. J.: A three-dimensional Eulerian acid deposition model: Physical concepts and formulation, *J. Geophys. Res-Atmos.*, 92, 14681-14700, 10.1029/JD092iD12p14681, 1987.
- Fountoukis, C., and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K<sup>+</sup>-Ca<sup>2+</sup>-Mg<sup>2+</sup>-NH<sub>4</sub><sup>+</sup>-Na<sup>+</sup>-SO<sub>4</sub><sup>2-</sup>-NO<sub>3</sub><sup>-</sup>-Cl<sup>-</sup>-H<sub>2</sub>O aerosols, *Atmos. Chem. Phys.*, 7, 4639-4659, 10.5194/acp-7-4639-2007, 2007.
- 355 Gately, C. K., Hutyra, L. R., Wing, I. S., and Brondfield, M. N.: A Bottom up Approach to on-Road CO<sub>2</sub> Emissions Estimates: Improved Spatial Accuracy and Applications for Regional Planning, *Environ. Sci. Technol.*, 47, 2423-2430, 10.1021/es304238v, 2013.
- Gately, C. K., Hutyra, L. R., and Sue Wing, I.: Cities, traffic, and CO<sub>2</sub>: A multidecadal assessment of trends, drivers, and scaling relationships, *P. Natl. Acad. Sci.*, 112, 4999-5004, 10.1073/pnas.1421723112, 2015.
- 360 Gurney, K. R., Mendoza, D. L., Zhou, Y., Fischer, M. L., Miller, C. C., Geethakumar, S., and de la Rue du Can, S.: High Resolution Fossil Fuel Combustion CO<sub>2</sub> Emission Fluxes for the United States, *Environ. Sci. Technol.*, 43, 5535-5541, 10.1021/es900806c, 2009.
- Henderson, B. H., Akhtar, F., Pye, H. O. T., Napelenok, S. L., and Hutzell, W. T.: A database and tool for boundary conditions for regional air quality modeling: description and evaluation, *Geosci. Model Dev.*, 7, 339-360, 10.5194/gmd-7-339-2014, 2014.
- 365 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058, 10.5194/acp-13-11019-2013, 2013.
- Lei, Y., Zhang, Q., He, K. B., and Streets, D. G.: Primary anthropogenic aerosol emission trends for China, 1990–2005, *Atmos. Chem. Phys.*, 11, 931-954, 10.5194/acp-11-931-2011, 2011.
- 370 Li, M., Zhang, Q., Streets, D. G., He, K. B., Cheng, Y. F., Emmons, L. K., Huo, H., Kang, S. C., Lu, Z., Shao, M., Su, H., Yu, X., and Zhang, Y.: Mapping Asian anthropogenic emissions of non-methane volatile organic compounds to multiple chemical mechanisms, *Atmos. Chem. Phys.*, 14, 5617-5638, 10.5194/acp-14-5617-2014, 2014.



- Li, M., Zhang, Q., Kurokawa, J., Woo, J. H., He, K. B., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng,  
375 Y. F., Hong, C. P., Huo, H., Jiang, X. J., Kang, S. C., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic  
emission inventory for the MICS-Asia and the HTAP projects, *Atmos. Chem. Phys. Discuss.*, 15, 34813-34869,  
10.5194/acpd-15-34813-2015, 2015.
- Liu, F., Beirle, S., Zhang, Q., Dörner, S., He, K. B., and Wagner, T.: NO<sub>x</sub> lifetimes and emissions of hotspots in polluted  
380 background estimated by satellite observations, *Atmos. Chem. Phys. Discuss.*, 15, 24179-24215, 10.5194/acpd-15-24179-  
2015, 2015.
- Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., and He, K. B.: High-resolution inventory of technologies, activities,  
and emissions of coal-fired power plants in China from 1990 to 2010, *Atmos. Chem. Phys.*, 15, 13299-13317, 10.5194/acp-  
15-13299-2015, 2015.
- McDonald, B. C., McBride, Z. C., Martin, E. W., and Harley, R. A.: High-resolution mapping of motor vehicle carbon  
385 dioxide emissions, *J. Geophys. Res-Atmos.*, 119, 5283-5298, 10.1002/2013JD021219, 2014.
- National Bureau of Statistics: China Statistical Yearbook for Regional Economy 2014, China Statistics Press, Beijing, 2014.
- Oda, T., and Maksyutov, S.: A very high-resolution (1 km×1 km) global fossil fuel CO<sub>2</sub> emission inventory derived using a  
point source database and satellite observations of nighttime lights, *Atmos. Chem. Phys.*, 11, 543-556, 10.5194/acp-11-543-  
2011, 2011.
- 390 Oak Ridge National Laboratory (ORNL): LandScan Global Population Database, Oak Ridge National Laboratory, Oak  
Ridge, Tenn., USA, 2013.
- Raupach, M. R., Rayner, P. J., and Paget, M.: Regional variations in spatial structure of nightlights, population density and  
fossil-fuel CO<sub>2</sub> emissions, *Energ. Policy*, 38, 4756-4764, 10.1016/j.enpol.2009.08.021, 2010.
- Rayner, P. J., Raupach, M. R., Paget, M., Peylin, P., and Koffi, E.: A new global gridded data set of CO<sub>2</sub> emissions from  
395 fossil fuel combustion: Methodology and evaluation, *J. Geophys. Res-Atmos.*, 115, D19306, 10.1029/2009JD013439, 2010.
- Schneider, A., Friedl, M. A., and Potere, D.: A new map of global urban extent from MODIS satellite data, *Environ. Res.  
Lett.*, 4, 044003, 2009.
- Walcek, C. J., and Taylor, G. R.: A Theoretical Method for Computing Vertical Distributions of Acidity and Sulfate  
Production within Cumulus Clouds, *J. Atmos. Sci.*, 43, 339-355, 10.1175/1520-0469(1986)043<0339:ATMFCV>2.0.CO;2,  
400 1986.
- Wang, R., Tao, S., Ciais, P., Shen, H. Z., Huang, Y., Chen, H., Shen, G. F., Wang, B., Li, W., Zhang, Y. Y., Lu, Y., Zhu, D.,  
Chen, Y. C., Liu, X. P., Wang, W. T., Wang, X. L., Liu, W. X., Li, B. G., and Piao, S. L.: High-resolution mapping of  
combustion processes and implications for CO<sub>2</sub> emissions, *Atmos. Chem. Phys.*, 13, 5189-5203, 10.5194/acp-13-5189-2013,  
2013.
- 405 Whitten, G. Z., Heo, G., Kimura, Y., McDonald-Buller, E., Allen, D. T., Carter, W. P. L., and Yarwood, G.: A new  
condensed toluene mechanism for Carbon Bond: CB05-TU, *Atmos. Environ.*, 44, 5346-5355,  
10.1016/j.atmosenv.2009.12.029, 2010.



- 410 Zhang, Q., Streets, D. G., He, K., Wang, Y., Richter, A., Burrows, J. P., Uno, I., Jang, C. J., Chen, D., Yao, Z., and Lei, Y.:  
NO<sub>x</sub> emission trends for China, 1995–2004: The view from the ground and the view from space, *J. Geophys. Res-Atmos.*,  
112, D22306, 10.1029/2007JD008684, 2007.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S.,  
Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos.*  
*Chem. Phys.*, 9, 5131-5153, 10.5194/acp-9-5131-2009, 2009.
- 415 Zhao, Y., Wang, S., Duan, L., Lei, Y., Cao, P., and Hao, J.: Primary air pollutant emissions of coal-fired power plants in  
China: Current status and future prediction, *Atmos. Environ.*, 42, 8442-8452, 10.1016/j.atmosenv.2008.08.021, 2008.
- Zheng, B., Huo, H., Zhang, Q., Yao, Z. L., Wang, X. T., Yang, X. F., Liu, H., and He, K. B.: High-resolution mapping of  
vehicle emissions in China in 2008, *Atmos. Chem. Phys.*, 14, 9787-9805, 10.5194/acp-14-9787-2014, 2014.
- Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L., and Kimoto, T.: Heterogeneous  
420 haze episode in North China, *Atmos. Chem. Phys.*, 15, 2031-2049, 10.5194/acp-15-2031-2015, 2015.





**Table 1. Proxies used for spatial distributions of emissions in HB-EI and MEIC <sup>a</sup>**

Sector	Subsector	HB-EI		MEIC	
		Province to County	County to Grid	Province to County	County to Grid
Power		point source		point source	
Industry		point source		industrial GDP <sup>b</sup>	urban population <sup>d</sup>
Residential	urban	residential coal use <sup>e</sup>	urban population <sup>d</sup>	urban population <sup>b</sup>	urban population <sup>d</sup>
	rural	residential coal/biofuel use <sup>e</sup>	rural population <sup>d</sup>	rural population <sup>b</sup>	rural population <sup>d</sup>
Transportation	on-road <sup>e</sup>	vehicle numbers	road network; traffic flow data	vehicle numbers	road network; traffic flow data
	non-road: agriculture	machine power <sup>b</sup>	rural population <sup>d</sup>	machine power <sup>b</sup>	rural population <sup>d</sup>
	non-road: construction	construction area <sup>c</sup>	urban population <sup>d</sup>	total GDP <sup>b</sup>	urban population <sup>d</sup>
	non-road: other source	total population <sup>b</sup>	total population <sup>d</sup>	total population <sup>b</sup>	total population <sup>d</sup>
Agriculture	fertilizer	fertilizer use <sup>b</sup>	rural population <sup>d</sup>	fertilizer use <sup>b</sup>	rural population <sup>d</sup>
	livestock	livestock amount <sup>c</sup>	rural population <sup>d</sup>	meat consumption <sup>b</sup>	rural population <sup>d</sup>

<sup>a</sup> The proxies in light blue are used in HB-EI and are different from those used in MEIC.

<sup>b</sup> Data source: National Bureau of Statistics, 2014.

<sup>c</sup> Data source: statistics from local agencies.

425 <sup>d</sup> Data source: population data (Oak Ridge National Laboratory, 2013), urban/rural extents (Schneider et al., 2009).

<sup>e</sup> Data source: Zheng et al., 2014.

**Table 2. Comparison of gridded emissions from MEIC and HB-EI at different resolutions <sup>a</sup>**

	R					SAD (Tg)					RSAD (%)				
	0.05°	0.1°	0.25°	0.5°	1.0°	0.05°	0.1°	0.25°	0.5°	1.0°	0.05°	0.1°	0.25°	0.5°	1.0°
<b>TSP</b>	0.25	0.44	0.71	0.84	0.84	3.56	2.92	2.09	1.56	1.28	136	111	80	60	49
<b>CO<sub>2</sub></b>	0.59	0.66	0.76	0.91	0.92	783	628	439	287	200	96	77	54	35	24
<b>SO<sub>2</sub></b>	0.54	0.68	0.81	0.9	0.89	1.33	1.06	0.75	0.57	0.47	99	79	56	42	35
<b>PM<sub>10</sub></b>	0.31	0.49	0.73	0.87	0.86	1.38	1.13	0.83	0.64	0.52	111	91	67	51	42
<b>CO</b>	0.23	0.4	0.69	0.79	0.77	18.66	16.34	12.63	10.63	9.24	114	100	77	65	56
<b>PM<sub>2.5</sub></b>	0.31	0.49	0.73	0.86	0.85	0.91	0.76	0.57	0.45	0.38	103	86	64	51	42
<b>NO<sub>x</sub></b>	0.78	0.83	0.87	0.94	0.95	1.34	1.04	0.77	0.5	0.39	68	52	39	25	20
<b>BC</b>	0.33	0.49	0.71	0.88	0.9	0.11	0.09	0.07	0.05	0.04	77	65	51	39	32
<b>VOCs</b>	0.72	0.84	0.91	0.96	0.97	0.8	0.65	0.5	0.37	0.28	53	44	34	25	19
<b>OC</b>	0.42	0.59	0.79	0.91	0.93	0.12	0.1	0.08	0.06	0.05	56	48	38	30	24
<b>NH<sub>3</sub></b>	0.93	0.97	0.99	1	1	0.04	0.03	0.03	0.02	0.02	7	6	5	4	3

<sup>a</sup>R: spatial correlation coefficient; SAD: summed absolute difference; RSAD: relative summed absolute difference.

**Table 3. Sensitivity analysis results**

<i>Emission contributions of point sources</i>											
Emissions	TSP	CO <sub>2</sub>	SO <sub>2</sub>	PM <sub>10</sub>	CO	PM <sub>2.5</sub>	NO <sub>x</sub>	BC	VOCs	OC	NH <sub>3</sub>
MEIC	2%	22%	17%	4%	2%	4%	23%	0%	0%	0%	0%
HB-EI	86%	82%	80%	73%	69%	67%	64%	47%	36%	34%	3%
Adjusted <sup>a</sup>	60%	58%	59%	53%	59%	50%	34%	29%	17%	23%	2%


  

<i>Model evaluation<sup>b</sup> of Domain 3 for S1, S2 and S3<sup>c</sup></i>											
Pollutants	Obs (μg m <sup>-3</sup> )	Sim (μg m <sup>-3</sup> )			MB (μg m <sup>-3</sup> )			NMB (%)			
	/	S1	S2	S3	S1	S2	S3	S1	S2	S3	
SO <sub>2</sub>	69.2	149.6	106	107.3	80.3	36.7	38	147.8	53	54.9	
NO <sub>2</sub>	52.5	86.6	75.4	82.2	34.1	22.9	29.7	80.5	43.6	56.6	
CO	1.9	2.4	1.5	1.4	0.5	-0.4	-0.5	18.1	-19.9	-24.2	
O <sub>3</sub>	60.8	47.7	53.3	48.9	-13.2	-7.7	-12.1	-21.6	-12.6	-19.8	
PM <sub>2.5</sub>	105.4	141	107.9	112.8	35.6	2.5	7.4	34	2.4	7	
PM <sub>10</sub>	202.1	174.2	125.8	132.7	-28	-76.3	-69.4	-16	-37.7	-34.3	

430 <sup>a</sup> Decrease point source shares by ~20% on the basis of the HB-EI inventory.

<sup>b</sup> The modeling evaluation is based on simulations from Jan, Apr, Jul and Oct.

<sup>c</sup> S1 used MEIC emissions; S2 used HB-EI emissions; S3 used adjusted emissions.



435 **Figure 1. Triple-nested domains of the CMAQ simulation: (a) spatial extent of triple-nested domains with OMI-derived NO<sub>2</sub> column densities (Boersma et al., 2011) in background; (b) domain 3 with NO<sub>x</sub> emissions from point sources in HB-EI inventory.**





Figure 2. Emission percentages contributed by point, nonpoint and mobile sources in Hebei Province: (a) the MEIC inventory; (b) the HB-EI inventory.



440 **Figure 3.** Comparison of the spatial distributions of gridded emissions from HB-EI and MEIC at multiple resolutions. (a–d) present  $\text{NO}_x$  emission distributions in HB-EI, and (e–h) present the differences between the two inventories (MEIC subtracted from HB-EI).







Figure 4. The spatial distributions of polluted industries and population in (a) Shijiazhuang and (b) Tangshan.



445

**Figure 5. Cumulative ratio of emissions by population density spanning the resolutions of (a) 0.05°, (b) 0.1°, (c) 0.5° and (d) 1.0°. The population is sorted according to descending numbers of people along the x axis.**






Figure 6. Evaluations against in situ measurements for atmospheric modeling using MEIC and HB-EI inventories at 36 km, 12 km and 4 km. The air pollutants used for evaluation include (a) SO<sub>2</sub>, (b) NO<sub>2</sub>, (c) CO, (d) O<sub>3</sub>, (e) PM<sub>2.5</sub>, and (f) PM<sub>10</sub>.

450




**Figure 7** Air pollution distributions and exposures by population density at 36 km, 12 km and 4 km: (a) NO<sub>2</sub>, (b) SO<sub>2</sub>, (c) population-weighted PM<sub>2.5</sub> and (d) population-weighted SOMO35. Note: the SOMO35 is used to evaluate the health risk of O<sub>3</sub> according to Amann et al. (2008).



455

**Figure 8. Spatial correlations between gridded emissions of HB-EI and various spatial proxies at the resolutions of (a) 0.05 °, (b) 0.1 °, (c) 0.25 °, (d) 0.5 ° and (e) 1.0 °. The pollutants are sorted according to descending contribution of point-source emissions from left to right. NL: nighttime light (<http://ngdc.noaa.gov/eog/dmsp/downloadV4composites.html>); TP: total population; UP: urban population; RP: rural population (population data (Oak Ridge National Laboratory, 2013) with urban/rural extents (Schneider et al., 2009); RN: road network (Zheng et al, 2014).**

460



**Figure 9** Sector-specific spatial correlation between NO<sub>x</sub> emissions from HB-EI and allocating proxies spanning resolutions from 30'' to 1.0°: (a) total emissions; (b) power sector; (c) industry sector; (d) residential sector; (e) transportation sector.