

# Current estimates of biogenic emissions from Eucalypts uncertain for Southeast Australia

## Supplementary Material

K.M. Emmerson<sup>1</sup>, I.E. Galbally<sup>1</sup>, A.B. Guenther<sup>2</sup>, C. Paton-Walsh<sup>3</sup>, E-A. Guerette<sup>3</sup>, M.E. Cope<sup>1</sup>, M.D.

5 Keywood<sup>1</sup>, S.J. Lawson<sup>1</sup>, S.B. Molloy<sup>1</sup>, E. Dunne<sup>1</sup>, M. Thatcher<sup>1</sup>, T. Karl<sup>4</sup>, S.D. Maleknia<sup>5</sup>

1 CSIRO Oceans & Atmosphere, PMB1, Aspendale, VIC, Australia.

2 Department of Earth System Science, University of California, Irvine, USA.

3 Centre for Atmospheric Chemistry, School of Chemistry, University of Wollongong, Wollongong, NSW, Australia.

4 Institute of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria.

10 5 Centre for Ecosystem Science, School of Biological, Earth and Environmental Sciences, University of New South Wales, Australia.

*Correspondence to:* K.M. Emmerson (kathryn.emmerson@csiro.au)

### Construction of high resolution PFT map for Australia

15 A new PFT dataset has been constructed for this work, at 3km resolution from the International Geosphere Biosphere Project (IGBP) dataset. Table S1 gives details of how the IGBP landcover dataset was split into the NCAR 16 PFTs suitable for MEGAN.

20

25

30

Table S1 Relationship of IGBP landcover types to NCAR PFTs, sorted according to boreal, temperate and tropical biomes. Not all landcover types are present in Australia.

IGBP Landcover type		NCAR Plant Functional Types			
		% Bare Ground	Boreal ≤19 °C	Temperate 19 °C ≤ X ≤ 15.5 °C	Tropical ≥15.5 °C
<b>0</b>	Ocean	100%			
<b>1</b>	Evergreen Needleleaf Forest	25%	75% Nt Eg B	75% Nt Eg T	
<b>2</b>	Evergreen Broadleaf Forest	5%		95% Bt Eg T	95% Bt Eg Tr
<b>3</b>	Deciduous Needleleaf Forest	50%	50% Nt Dc B		
<b>4</b>	Deciduous Broadleaf Forest	25%	75% Bt Dc B	75% Bt Dc T	75% Bt Dc Tr
<b>5</b>	Mixed Forest	26%	55.5% Nt Eg B 18.5% Bt Dc B	55.5% Bt Eg T 18.5% Bt Dc T	55.5% Bt Eg Tr 18.5% Bt Dc Tr
<b>6</b>	Closed Shrubland	20%	64% Sb Dc B 16% Gs C3 Cd	32% Sb Eg T 32% Sb Dc T 16% Gs C3 Cl	16% Gs C3 W
<b>7</b>	Open Shrubland	50%	10% Sb Dc B 40% Gs C3 Cd	5% Sb Eg T 5% Sb Dc T 40% Gs C3 Cl	40% Gs C3 W
<b>8</b>	Woody Savannah	-	30% Nt Eg B 10% Bt Dc B 60% Gs C3 Cd	30% Bt Eg T 10% Bt Dc T 60% Gs C3 Cl	30% Bt Eg Tr 10% Bt Dc Tr 60% Gs C3 W
<b>9</b>	Savannahs	-	7.5% Nt Eg B 2.5% Bt Dc B 90% Gs C3 Cd	7.5% Bt Eg T 2.5% Bt Dc T 90% Gs C3 Cl	7.5% Bt Eg Tr 2.5% Bt Dc Tr 90% Gs C3 W
<b>10</b>	Grasslands	20%	80% Gs C3 Cd	80% Gs C3 Cl	80% Gs C3 W
<b>11</b>	Permanent Wetland	100%			
<b>12</b>	Cropland	15%	85% Crop	76.5% Crop 8.5% Corn	59.5% Crop 25.5% Corn
<b>13</b>	Urban and Built-Up	70%	7.5% Nt Eg B 7.5% Nt Dc B 5% Sb Dc B 10% Gs C3 Cd	7.5% Bt Eg T 7.5% Bt Dc T 5% Sb Eg T 10% Gs C3 Cl	7.5% Bt Eg Tr 7.5% Bt Dc Tr 10% Gs C3 W
<b>14</b>	Cropland/Natural Vegetation Mosaic	15%	85% Crop	85% Crop	85% Crop
<b>15</b>	Snow and Ice	100%			
<b>16</b>	Barren or Sparsely Vegetated	100%			
<b>17</b>	Water Bodies	100%			

5 Notes: B = Boreal, T = Temperate, Tr = Tropical.

Nt Eg = needleleaf evergreen tree, Nt Dc = needleleaf deciduous tree, Bt Eg = broadleaf evergreen tree, Bt Dc = broadleaf deciduous tree, Sb Eg = evergreen shrub, Sb Dc = deciduous shrub, Gs C3 = grass (can be Cd = cold, Cl = cool or W = warm depending on climatic zone).

## Comparisons with air quality data

We have tested modelled meteorological, chemical and dilution variables and conclude that there are no confounding influences that distort the model response to BVOCs. Figure S1 shows diurnal cycles of temperature, ozone and NO<sub>x</sub> where data are available. These comparisons give a broader impression of how the model predicts airshed chemistry, as both ozone and NO<sub>x</sub> are influenced by urban emissions. Temperature is a major driver of MEGAN and many compound classes exhibit a temperature dependence. The average modelled profiles compare to within 1 or 2 degrees C of the observations. The modelled ozone profiles are high (<5ppb) for SPS1 and MUMBA, and low for SPS2. Reducing the isoprene and increasing the monoterpene emission rates has reduced the peak SPS1 modelled ozone concentration by up to 2 ppb more than the peak MUMBA ozone concentration, bringing them more into line with the observations. The shape of the NO<sub>x</sub> diurnal profiles show that the timing of the morning and evening traffic peaks is correct. However, there is a high bias in modelled NO<sub>x</sub> during SPS1 which is not present in either SPS2 or MUMBA. The anthropogenic emissions inventory is the same for each of the sites, thus differences are likely due to modelled dilution and mixing heights during SPS1. Modelled NO<sub>x</sub> in SPS2 shows a high peak during the late afternoon, which is attributed to a collapsing boundary layer occurring too early in the day in the model compared with the observations. The increase/reduction in the biogenic VOCs from the emission factor sensitivity test has not changed the diurnal profiles of NO<sub>x</sub> much, although there are small differences at nighttime during SPS1 and SPS2.

There is uncertainty in the NO<sub>x</sub> emissions which affects the airshed chemistry of the modelled domains. A sensitivity test is conducted whereby the NO<sub>x</sub> emissions associated with motor vehicles, industrial processes and fire sources are scaled by  $\pm 20\%$ . Results in

Table S2 show the effects of the NO<sub>x</sub> emissions increase/decrease (and emission factor sensitivity test) on average concentrations of NO<sub>x</sub>, ozone, xylene and the biogenic species. Larger percentage changes are seen in locations close to anthropogenic sources, i.e. Sydney. Tumbarumba is rural and located far from the influences of urban NO<sub>x</sub> emissions, thus no effects are observed. Overall a  $\pm 20\%$  change in NO<sub>x</sub> causes 1-5% change in isoprene and monoterpene concentrations. Xylene is chosen to represent an anthropogenic primary species, whose concentration is dependent on the availability of oxidant species. The results indicate that changes in the biogenic species and NO<sub>x</sub> impact on xylene by 1%. There are some unexpected consequences of the non-linearities in the chemical system, i.e. a decrease in ozone should cause an increase in isoprene concentrations at MUMBA. Horizontal advection is also a factor as is the time taken for chemical production of ozone.

The model has been tested against observed concentrations and meteorological data. Sensitivity studies varying NO<sub>x</sub> and VOC emissions have been conducted to test the robustness of the modelling system. We conclude that there are no confounding influences that distort the model response to BVOCs.

5




Figure S1 Campaign average diurnal cycles for temperature, ozone and NO<sub>x</sub>. S1=SPS1, S2=SPS2, M=MUMBA. F2= percentage within a factor of 2 between observations and base run.

10

15

Table S2 Change in domain average concentrations for the duration of each field campaign. The percentage change from the base run is indicated number in brackets, for the emission factor sensitivity test and changes in anthropogenic NOx emissions.

		<b>Ozone</b>	<b>NOx</b>	<b>Isoprene</b>	<b>Monoterpenes</b>	<b>Xylene</b>
SPS1	Base	24.5	2.5	4.0	0.3	0.1
	EF test	23.9(-2%)	2.6 (2%)	1.1(-73%)	1.0(207%)	0.1(<1%)
	NOx +20%	24.8(1%)	3.0 (22%)	3.9(-2%)	0.3(-2%)	0.1(<1%)
	NOx -20%	24.2(-1%)	2.0(-21%)	4.1(3%)	0.3(2%)	0.1(<1%)
SPS2	Base	19.6	3.8	0.9	0.2	0.1
	EF test	19.2(-2%)	3.9(2%)	0.3(-70%)	0.7(224%)	0.1(-1%)
	NOx +20%	19.3(-2%)	4.6(22%)	0.9(-1%)	0.2(-1%)	0.1(2%)
	NOx -20%	19.9(2%)	3.0(-21%)	0.9(2%)	0.2(1%)	0.1(<1%)
Mumba	Base	26.2	2.9	1.8	0.1	0.03
	EF test	25.3(3%)	3.0(2%)	0.5(-72%)	0.4(211%)	0.03(<1%)
	NOx +20%	26.1(<1%)	3.6(21%)	1.7(-2%)	0.1(-3%)	0.03(2%)
	NOx -20%	26.2(<1%)	2.3(-21%)	1.8(3%)	0.1(1%)	0.03(<1%)
Tumba.	Base	18.4	0.4	0.7	0.2	0.009
	EF test	18.3(<1%)	0.3(-4%)	0.2(-71%)	0.6(211%)	0.009(<1%)
	NOx +20%	18.4(<1%)	0.4(<1%)	0.7(<1%)	0.2 (<1%)	0.009(<1%)
	NOx -20%	18.4(<1%)	0.4(<1%)	0.7 (<1%)	0.2 (<1%)	0.009(<1%)