

Interactive comment on “Impact of Saharan dust on North Atlantic marine stratocumulus clouds: Importance of the semi-direct effect” by Anahita Amiri-Farahani et al.

Dr. Kishcha

pavel@cyclone.tau.ac.il

Received and published: 16 November 2016

Large uncertainty of the estimated dust radiative effect in winter and the contribution of non-dusty aerosols.

Based on satellite-retrieved parameters of cloud properties, the authors (Amiri-Farahani et al., 2016) concluded that, in the winter season, the dust – cloud radiative effect is “weakly positive $0.92 \pm 2.86 \text{ W/m}^2$ ”. However, in fact, their estimate indicates that, in winter, the dust – cloud radiative effect could be either positive or negative, because of the large uncertainty of their estimate. The presence of non-dusty aerosols could also be a causal factor for the above-mentioned large uncertainty. In our papers (Kishcha et al., 2014, 2015), using NASA MERRAero reanalysis, we showed that, in

Printer-friendly version

Discussion paper



winter, Saharan dust is not the predominant aerosol species over the tropical North Atlantic, including the area 45W – 15E; 0N – 35N under the study by Amiri-Farahani et al. (2016). Apart from dust, non-dusty aerosols, such as carbonates (organic and black carbon), sea salt and sulfates also significantly contribute to the total AOD over the tropical North Atlantic. As shown in Fig. 1 (below), in contrast to July, in January dust dominates other aerosol species only near the African coast. The non-dusty aerosol species could affect cloudiness in a different manner. In the winter season, absorbing aerosols, such as organic and black carbon, produce mainly a positive semi-direct radiative effect, similar to the dust effect. Sulfates and sea salt, non-absorbing aerosols, produce a negative indirect radiative effect, acting as effective CCN. Thus, non-dusty aerosols, producing either positive or negative radiative effects, significantly contribute to the large uncertainty of the aerosol-cloud radiative effect in the winter season.

References: Kishcha P., da Silva A., Starobinets B., Long C.N., Kalashnikova O., Alpert P. (2014). Meridional distribution of aerosol optical thickness over the tropical Atlantic Ocean. *Atmospheric Chemistry and Physics Discussion* 14, 23309-23339, doi:10.5194/acpd-14-23309-2014.

Kishcha P., da Silva A., Starobinets B., Long C.N., Kalashnikova O., Alpert P. (2015). Saharan dust as a causal factor of hemispheric asymmetry in aerosols and cloud cover over the tropical Atlantic Ocean. *International Journal of Remote Sensing* 36, 3423-3445, doi: 10.1080/01431161.2015.1060646.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2016-933, 2016.

[Printer-friendly version](#)[Discussion paper](#)




Fig. 1. The ratio of dust aerosol optical depth (AOD) to AOD of all other aerosol species, based on the 10-year MERRAero data (July 2002 - June 2012).