

Observations of Atmospheric Chemical Deposition to High Arctic Snow – Second Response to Anonymous Referee #1

Referee comments received and published: 14 Mar 2017 (quoted below in blue text)

We would like to thank Referee #1 for their detailed comments and discussion. We greatly appreciate the care with which the referee has reviewed this manuscript and the improvements gained through their insight.

Response to Referee Discussion

Referee Comment: This draft of the manuscript is significantly improved from the original submission. I am still far from convinced that the “effective deposition velocity” is a useful construct, but it is obvious that the authors feel that it is. I also feel that the further attempt to explain the very high EVD for Ca^{2+} is not convincing. It is troubling that the detection limit (reported in Supplemental Table 6) is so much higher (more than a factor of 7) than all other ions, suggesting very large variability in the blanks, or a serious analytical problem. (Note this is not true for the aerosol composition detection limits in Supplemental Table 8). The authors assert that qualitative agreement with ICPMS measurements of Ca gives confidence in both data sets, but only about half of all possible samples were above detection limits on both instruments. I find it hard to believe that so many samples were below ICPMS detection limits, raising concern about that data set as well. Given that the authors can not really explain the high EVD, they might be better served leaving Ca^{2+} out of the story due to analytical issues causing large uncertainties.

Response: The high method detection limit (MDL) of Ca^{2+} , as pointed out by the referee, is a result of variation in the shipping/handling blanks (Hblks). There were five of these blanks collected over the course of the campaign, roughly one per month. These bottles were opened at Alert, but with no snow collected, and then transported and analysed alongside the other samples to capture any background associated with the full handling process. Of these five Hblks, three were below the Ca^{2+} instrument detection limit (IDL) of 18 ppb, and two were above IDL at 50 and 79 ppb. The MDL for each compound was calculated as the maximum of the following:

- IDL
- Three standard deviations of the Hblks
- Three standard deviation of the preparation blanks

Thus, for Ca^{2+} , the MDL was set to three standard deviation of the five Hblks. This is a conservative approach since the three Hblks below IDL were very imprecise, but reflects the full variability in the potential shipping/handling background. While all but one Ca^{2+} samples were well above the IDL, only 63% were above the calculated MDL. The other ions measured by IC did not show such a large variability in HBlks, leading to their lower MDL values.

To estimate the impact of this potential background we have used the MDL to get a “background effective deposition velocity”. Dividing the MDL by the maximum snow depth and the maximum atmospheric concentration give a conservative background of about 0.45 cm/s (note, this background is over exaggerated since these maxima do not coincide). Subtracting this exaggerated background from the calculated Ca^{2+} effective deposition velocities, the warm month Ca^{2+} median would still be very high at about 2.2 cm/s and the cold month median would be more similar to the other particle-dominated analytes at about 0.4 cm/s. Thus, we believe the observation of high Ca^{2+} deposition rates in the warm months to be reasonable, even if we do not at present have an explanation for the anomaly. Figures 2, 3, and 5 in the manuscript show the associated uncertainty about the Ca^{2+}

deposition velocity to intersect zero for several cold months; thus we believe it is clear that the cold-month deposition velocity is not significantly different from those of other analytes.

We have also considered the Ca^{2+} values relative to other crustal-related analytes. Figure 5 shows that the maximum monthly deposition velocities for K^+ and Mg^{2+} are similar to or greater than the maximum Ca^{2+} monthly velocity. In fact, if the average velocity is considered instead of the median, K^+ has the highest warm-month value at 3.2 cm/s. Furthermore, the ratios of Ca^{2+} to Mg^{2+} in this study was found to be typical compared to those seen by others (as listed in supplemental Table S5: Ross and Granat, 1986; Li and Winchester, 1993; Osada et al., 1996; Toom-Sauntry and Barrie, 2002; Dibb et al., 2007).

While we agree that uncertain measurements should not be included in the analysis, we believe the Ca^{2+} to be sufficiently substantiated to keep in the paper with the provided information on data uncertainty. Although we do not have an explanation for the discrepancy between Ca^{2+} and Mg^{2+} we hope that a reasonable hypothesis may be proposed in the future and perhaps spur future research in this area.

Referee Comment: One thing that the authors may want to expand on a bit more is the suggestion that the study period may have been unusual due to the active Icelandic volcano (mentioned in just one sentence near the end of discussion; pg 17 lines 2-5). Provision of the detailed aerosol data in Supplement Table 7 really makes it clear that the canonical “Arctic Haze” pattern for sulfate (and other pollution and dust derived constituents that usually covary with sulfate) with elevated concentrations in late winter into spring, generally peaking DJF, was not obvious. The really striking observation to me is that sulfate deposition to/with snow is so weak in the winter time, even in a year that may have had the usual haze peak in the atmosphere plus additional enhancement from the volcano. Cliff Davidson suggested very weak scavenging of sulfate by winter snow near Dye 3 way back (several papers before DGASP), based on snowpits that showed winter enhancements that were not proportional to increase in aerosol sulfate at Barrow, Alert, NyAesund during the haze season. DGASP showed that Haze really does not impact sites high on the Greenland ice sheet, so the inferred weak scavenging was not confirmed over Greenland. (Cliff actually claimed marked seasonality of sulfate scavenging, stronger in summer, since he did have aerosol measurements in summer that showed low sulfate compared to what he assumed it was in winter/spring.) Now it may be that you are confirming weak removal of sulfate during winter for Alert, and perhaps other parts of the Arctic basin, but this probably means untangling any impacts on this winter from the volcano.

Response: We agree with the referee that this is an interesting observation. The unusual fall peak in SO_4^{2-} is also mentioned earlier in the text while comparing the observed snow mixing ratios to previous studies (revised page 9 /lines 9-10). To emphasize that the observations of SO_4^{2-} may not be typical of Arctic Haze the following line has been added to the manuscript: *“Thus, the SO_4^{2-} observations of this campaign, especially the peak snow mixing ratio seen in the fall, may not reflect a seasonal trend for typical Arctic Haze.”* (16/5-6)

We believe the Davidson study referred to by the referee is the 1987 study listed below. This Davidson study was referenced in the manuscript in regards to the typical range of dry deposition velocities but not for the temporal variability in deposition. The following line has been added to the manuscript to correct this oversight: *“This aligns with seasonal trend in SO_4^{2-} deposition observed by Davidson et al. (1985b and 1987) and the suggested link to deposition mechanism.”* (14/29-30)

Davidson, C. I., Honrath, R. E., Kadane, J. B., Tsay, R. S., Mayewski, P. A., Lyons, W. B., and Heidam, N. Z.: The Scavenging of atmospheric sulfate by Arctic snow, *Atmos. Environ. A-Gen.*, 21 (4), 871–882, doi:10.1016/0004-6981(87)90083-7, 1987.

Davidson, C. I., Santhanam, Fortmann, S. R. C., and Marvin, P. O.: Atmospheric transport and deposition of trace elements onto the Greenland ice sheet, *Atmos. Environ.*, 19 (12), 2065-2081, 1985b.

Referee Comment: As noted last time, the authors really need to look at Dibb et al., 2007 (Seasonal variations in the soluble ion content of snow at Summit, Greenland: Constraints from three years of daily surface snow samples, *Atmos. Environ.*, 41, 5007-5019). As suggested by the title, this paper reports on 3 years, through the winters, of daily snow sampling at Summit, including ion concentrations and fluxes, and specifically separates freshly fallen from surface snow aged days to weeks. It probably needs to be considered in relation to the summary of previous work at top of page 2, and contradicts the statement made in lines 4-5 on page 3. Also, a main point of Dibb et al., 2007 was to compare seasonality of ions in snow falling at Summit to the seasonality inferred from snowpit studies, so it would seem an additional valid source for the comparisons made at the end of section 3.1 in this manuscript (page 9, lines 5-15).

Response: We apologize for missing the Dibb et al. (2007) study in our last revision. This reference has been added to the discussion in section 3.1 and Table S5.

Dibb, J. E., Whitlow, S. I., and Arsenault, M.: Seasonal variations in the soluble ion content of snow at Summit. Greenland: Constraints from three years of daily surface snow samples, *Atmos. Environ.*, 41, 5007-5019, doi:10.1016/j.atmosenv.2006.12.010, 2007.

Response to Detailed Comments

Referenced to Page/Line #(s) in the original manuscript:

3/30

Referee Comment: is unavoidable and natural within natural snowpack systems. → is unavoidable within natural snowpacks and still expected to smaller extent on the snow table.

Response: Line corrected in text. (revised manuscript page/line 3/32)

3/30-32

Referee Comment: It is actually not necessary to know the depth of a sampled surface stratigraphic layer to calculate fluxes, in fact the depth can, and often does, vary horizontally. All that is needed is the area sampled and the mass of snow in that area (e.g., g snow/cm²). Given concentrations in mass or mole per g of melted snow, conc x g snow/cm² give the amount of chemical in the layer per cm². The harder challenge is to only sample from single stratigraphic layer, but that can be done with care.

Original Line: *Also, the collection of samples from a snow table eliminated the need to estimate the depth of fresh snow, a source of uncertainty for traditional surface snow sampling.*

Response: This line was meant to touch on the second challenge described by the referee: the difficulty in sampling from a single stratigraphic layer. The line has been revised to clarify this point as follows: *Also, the collection of samples from a snow table eliminated the difficulty in distinguishing the fresh stratigraphic snow layer from aged layers below, a source of uncertainty for traditional surface snow sampling.* (4/1-2)

4/3-7

Referee Comment: Possible underestimate of snowfall up to a factor of 10 seems too large to just pass over. Here in the text it is stated that large discrepancies between operator reports and a met station separated by 6 km justify ignoring the station data, referring to section 4.2 in the supplement for details. However, there are no details on the discrepancies in the supplement, just a rephrased version of the same statement.

Response: The discussion of meteorological data in the supplemental has been expanded. We agree with the referee that the difference in snow depth may indicate under-catch by the snow table; however, observed significant differences in weather conditions between the collection site and meteorological station during the campaign as well as operator notes from previous years make us weary to simply adopt the meteorological station data for this analysis. Some dates with recorded precipitation at the collection site showed none at the meteorological station, and vice-versa. To provide better confidence in the analysis we have included a revised version of Figure 3 in the supplemental (revised Figure S3) which uses meteorological station snow depths instead of those measured at the snow table. The results show a change in absolute value of the effective deposition velocities, though largely within the uncertainty shown in the original plot. Yet, interpretation of the data remains largely unchanged as the relative values between analytes are unaffected.

4/30

Referee Comment: Be more specific what you mean by underestimate of BC mass due to size cut-off. First point is that 0.02-50 fg is not directly a size (need to assume shape factor and density). Also not clear whether you are worried about truncation on the small end or the large end of the distribution (both could be possible). It also seems that the cited mass range is based on the SP2 detection principle, probably also need to think about any issues related to transmission efficiency of the nebulizer.

Original Line: *Observed BC mass distributions did not suggest significant underestimation of the total BC mass due to this size cut-off.*

Response: Details on the SP2 analysis are provided in the supplemental section S2.1.1. This section provides the density and shape assumptions and a description of the cut-off concern. As stated in the text, the mass outside the detectable range was found to be negligible, both above and below.

7/3-6 and Table 1

Referee Comment: It is a little confusing that the list in the text does not overlap very well with the elements in the table (e.g. Ba, Ti, K, Cd, Ca, Cr, and Na are in text but not table while V, Se, Sb, and Tl are in the table but not listed in the text). Could either make the list in text agree with this table, or expand the list to include all of the elements in supplemental tables S3 and S4 and make it clear you are referring to those tables and not Table 1.

Response: We apologise for this oversight. The line in question has been corrected to discuss only those metals included in Table 1. The metals Se, Sb, and Tl have been left out of this list due to insufficient filtrate measurements above detection limit. Since Table 1 only described the insoluble portion of these metals their inclusion in the table is considered acceptable.

Corrected Line: *The metal measurements can be roughly classified into three categories: predominantly insoluble analytes Fe and Al (>50% insoluble over full campaign); variably soluble/insoluble analytes Co, V, As, Cu, Pb, Mn, K, and Mg; and predominantly soluble analytes Ca and Na (<50% insoluble) (in order from least to greatest average soluble fraction), excluding analytes with insufficient soluble or insoluble measurements above MDL. (7/5-9)*

Table 1 Footnote

Referee Comment: MSA, ACE, PRP, and FOR are abbreviations, so OK to define, but $C_2O_4^{2-}$ and NH_4^+ are not. If you need to define these ion formulae, you probably need to define all of them. Better choice would seem to stop after the first 4.

Related comment, MSA usually signifies methylsulfonic acid, not the ion methylsulfonate that you correctly indicate was measured by IC. I think that MS^- would be better shorthand to indicate methylsulfonate (here and everywhere else in the MS).

Response: The names for all abbreviations and all compounds have been provided in the text at their first appearance to provide clarity to the reader. The other abbreviations/compounds in Table 1 (BC , NO_3^- , and SO_4^{2-}) were named at their first appearance in section 1, but the chemical species listed above appear in the manuscript for the first time in Table 1. However, we agree that naming compounds is not strictly necessary and that Tables and the abbreviations therein should be stand-alone if possible. Thus, all abbreviations with Table 1 have been defined in the revised footnote and compound names have only been provided at their first appearance in the body of the text.

As per the referee's suggestion, the acronym MSA has been replaced with MS throughout the manuscript and supplemental.

Section 3.2.2, 3.2.3, and 4

Referee Comment: Need to be consistent and show the valences for all of the ions everytime they appear in the text. Right now only a few are shown. This is especially important since you also discuss results of elemental analyses for Na, K, Mg and Ca. Throughout these 3 sections the references are all to the ions, not elements.

Response: We agree that adding valences for all chemical species measured by IC would provide better consistency throughout the discussion. We had hoped to avoid readers misinterpreting the inclusion of valences as an indication of the expected form in the environment; however, if the exclusion of valence causes greater confusion then we are happy to add them back in. The exception is when chemical species are being discussed in general, not specifically as measurements (for example “...Cl partitioned to the gas phase...” or “... Ca-rich mineral dust...”). This exception may lead to a lack of consistency but we believe that incorrect inclusion of valence in such a discussion would lead to greater confusion. We have previously added the following line in hope to discourage readers from misinterpreting any included valences as an indication of the expected in-situ form: “It should be noted that although IC measurements are provided as the measured ions throughout the discussion, these analytes may not necessarily exist in the dissociated ionic form in the environment.”

12/6

Referee Comment: Not so clear that NH_4^+ is always dominant compared to NH_3 . Maybe in the Arctic, and during cold season, but this reads like a global declaration.

Original Line: *First, the measured chemical species differ in terms of dominant phase: BC, NH_4^+ , SO_4^{2-} , Na, K, Mg, $\text{C}_2\text{O}_4^{2-}$, and Ca are typically observed predominantly in the particle phase, while MSA, Br, Cl, and NO_3^- and their associated precursors can have appreciable gas-phase portions (Barrie and Hoff, 1985).*

Response: This line has been revised to clarify that the predominant phases discussed are in reference to within the Arctic.

Revised Line: *First, the measured chemical species differ in terms of dominant phase: BC, NH_4^+ (ammonium), SO_4^{2-} , Na^+ , K^+ , Mg^{2+} , $\text{C}_2\text{O}_4^{2+}$ (oxalate), and Ca^{2+} are typically observed predominantly in the particle phase within the Arctic, while MSA, Br, Cl, and NO_3^- and their associated precursors can have appreciable gas-phase portions (Barrie and Hoff, 1985). (12/5-8)*

12/8-9

Referee Comment: As noted earlier, Ca^{2+} is a striking exception to this statement, for no plausible reason. Do you really think this is real?

Response: See response to comment above.

13/8

Referee Comment: Seems “in-cloud scavenging” should be “below-cloud”

Original Line: *Particles containing these analytes, as well as NH_4^+ and SO_4^{2-} , have been previously suggested to act as better precipitation nuclei than BC (e.g., Zobrist et al., 2006; Hoose and Möhler, 2012; Farmer, Cappa, and Kreidenweis, 2015); thus, their enhanced velocities might be attributed to either enhanced scavenging via cloud formation or in-cloud scavenging.*

Revised Line: *Particles containing these analytes, as well as NH_4^+ and SO_4^{2-} , have been previously suggested to act as better precipitation nuclei than BC (e.g., Zobrist et al., 2006; Hoose and Möhler, 2012; Farmer, Cappa, and Kreidenweis, 2015); thus, their enhanced velocities might be attributed to either in-cloud scavenging. (13/7-10)*

Observations of Atmospheric Chemical Deposition to High Arctic Snow

Katrina M. Macdonald¹, Sangeeta Sharma², Desiree Toom², Alina Chivulescu², Sarah Hanna³, Allan K. Bertram³, Andrew Platt², Mike Elsasser², Lin Huang², David Tarasick⁴, Nathan Chellman⁵, Joseph R. McConnell⁵, Heiko Bozem⁶, Daniel Kunkel⁶, Ying Duan Lei¹, Greg J. Evans¹, Jonathan P. D. Abbatt⁷

¹ Department of Chemical Engineering and Applied Chemistry, University of Toronto, M5S 3E5, Canada

² Climate Research Divisions, Environment and Climate Change Canada, Toronto, M3H 5T4, Canada

³ Department of Chemistry, University of British Columbia, Vancouver, V6T 1Z1, Canada

⁴ Air Quality Research Divisions, Environment and Climate Change Canada, Toronto, M3H 5T4, Canada

10 ⁵ Desert Research Institute, Reno, 89512, United States of America

⁶ Institute for Atmospheric Physics, Johannes Gutenberg University Mainz, 55128, Germany

⁷ Department of Chemistry, University of Toronto, M5S 3H6, Canada

Correspondence to: Jonathan Abbatt (jabbatt@chem.utoronto.ca)

Abstract. Rapidly rising temperatures and loss of snow and ice cover have demonstrated the unique vulnerability of the high Arctic to climate change. There are major uncertainties in modelling the chemical depositional and scavenging processes of Arctic snow. To that end, fresh snow samples collected on average every four days at Alert, Nunavut, from September 2014 to June 2015 were analyzed for black carbon, major ions, and metals, and their concentrations and fluxes reported. Comparison with simultaneous measurements of atmospheric aerosol mass loadings yields effective deposition velocities that encompass all processes by which the atmospheric species are transferred to the snow. It is inferred from these values that dry deposition is the dominant removal mechanism for several compounds over the winter while wet deposition increased in importance in the fall and spring, possibly due to enhanced scavenging by mixed-phase clouds. Black carbon aerosol was the least efficiently deposited species to the snow.

1 Introduction and Background

In recent decades drastic changes have been observed within the Arctic, including a rapid increase in surface temperatures and loss of sea ice and snow cover (Rigor et al., 2000; Stroeve et al., 2005; Hartmann et al., 2013). Not only have these changes had adverse consequences for local populations and ecosystems, it has been suggested that their impacts may be significant at the global scale (Law and Stohl, 2007; AMAP, 2011). Light-absorbing compounds, the most widely-studied of which being black carbon (BC) particles, can have a particularly significant impact on the Arctic atmosphere and snow systems through the absorption of solar radiation and subsequent warming and snow melt (Bond et al., 2013). While the Arctic atmosphere has been previously explored spatially, temporally, and compositionally (e.g., Hartmann et al., 2013), Arctic snow and the mechanisms linking snow to the atmosphere have been the subject of only a relatively small number of

studies (AMAP, 2011), despite the enormous amount of research conducted on the Arctic Haze phenomenon (Quinn et al., 2007). Seasonal observations of fresh snow samples are particularly uncommon (e.g., Davidson et al., 1993; Toom-Saunry and Barrie, 2002; Hagler et al., 2007) and previous explorations of snow deposition and scavenging mechanisms have been largely reliant on short-term or aged snowpack sampling (e.g., Bergin et al., 1995), ice cores (e.g., Legrand and De Angelis 5 1995), modelling, and laboratory tests.

Aerosols entering the Arctic atmosphere, either generated locally or transported from elsewhere, can be removed by atmospheric transport or deposition. Deposition of particles follows two mechanisms: dry deposition, whereby particles are deposited to the ground by impaction, gravitational settling, and Brownian motion; and wet deposition, whereby particles are scavenged by hydrometeors and deposited through precipitation. Wet deposition is further split into two scavenging 10 mechanisms: in-cloud scavenging which removes particles from the cloud layer during precipitation formation, and below-cloud scavenging which removes particles from the atmospheric column through which precipitation falls. Gaseous compounds also undergo similar scavenging processes (Seinfeld and Pandis, 2006).

The rate of dry deposition is dependent on the properties of the depositing particle, the surface onto which deposition occurs, and the air-surface boundary layer (Sehmel, 1980; Zhang and Vet, 2006). Dry deposition velocities of accumulation mode 15 particles, the dominant mass-weighted mode of particles observed in the non-summer Arctic (Sharma et al., 2013), to snow have been modelled and observed over a range of 0.01 to 0.60 cm/s, typically within 0.02 to 0.10 cm/s; gaseous deposition velocities to a snow surface show a similar range, with observations from 0.05 to 0.50 cm/s and a typical velocity of approximately 0.10 cm/s (McMahon and Denison, 1979; Sehmel, 1980; [Davidson et al., 1985b](#); Davidson et al., 1987; Hillamo et al., 1993; Bergin et al., 1995; Petroff and Zhang, 2010; Liu et al., 2011). Wet deposition is dependent on the 20 properties of the depositing aerosol and the atmospheric conditions. In-cloud scavenging is largely controlled by a particle's size and composition which dictate its ability to nucleate hydrometeors and to be scavenged by cloud droplets. Particles can act as cloud condensation nuclei (CCN) that nucleate water droplets or ice nuclei (IN) that nucleate ice crystals. Common CCN components include sea salt, sulphate (SO_4^{2-}), and nitrate (NO_3^-), while mineral dust and bioaerosols are common IN (Hoose and Möhler, 2012; Farmer, Cappa, and Kreidenweis, 2015). Typically, BC is considered to be an ineffective CCN or 25 IN (Hoose and Möhler, 2012; Farmer, Cappa, and Kreidenweis, 2015). Although liquid water clouds are not expected during the Arctic winter, mixed-phase clouds, which contain both liquid water and ice, have been observed in this region at temperatures well below 0 °C, in unusual cases down to -40 °C but more commonly between -20 °C and -10 °C (Morrison et al., 2005; Shupe et al., 2006). Below-cloud deposition is dependent on snow type and meteorological conditions, which dictate the volume of air scavenged per snowfall (Zhang and Vet, 2006). Particle size also affects below-cloud scavenging with higher scavenging efficiencies for particles above 2.5 µm diameter relative to accumulation mode particles (Zhang and 30 Vet, 2006). The total wet deposition is a function of the volume of precipitation. Hence, wet deposition is not typically described via a deposition velocity.

The goals of this paper are to present a new dataset in which the chemical composition of freshly fallen snow was measured through a fall-winter-spring period at a high Arctic field station. By combining these data with simultaneous measurements

of ambient aerosol, the efficiency of deposition of individual species from the atmosphere to the snow can be evaluated under a set of broad assumptions. While this paper presents the measurement data set in detail and focuses on the depositional and scavenging mechanisms that can be inferred from it, a subsequent publication will identify potential pollutant sources based on the snow compositional data. To our knowledge, this is the first time that the composition and flux of freshly fallen snow has been analyzed at high temporal frequency throughout an entire cold season in the high Arctic.

[All data from this study will be available upon conclusion of the NETCARE project via the Government of Canada Open Data Portal.](#)

2 Methodology

2.1 Snow Sample Collection

Snow samples were collected at Environment and Climate Change Canada's (ECCC) Dr. Neil Trivett Global Atmosphere Watch Observatory at Alert, Nunavut from September 14th, 2014 to June 1st, 2015 as part of the Network on Climate and Aerosols Research (NETCARE) initiative to create a temporally-refined and broadly speciated dataset of high Arctic snow measurements. Alert is a remote outpost in the Canadian high Arctic, at the northern coast of Ellesmere Island (82°27' N, 62°30' W), with a small transient population of research and military personnel (location details provided in supplemental Sect. S4). Snow samples were collected from two Teflon-surfaced snow tables (about 1 m² by 1 m above ground level, shown in supplemental Figure S2) located in an open-air minimal traffic site, about 6 km SSW of the Alert base camp, 201 m above sea level. Freshly fallen snow was collected from the tables using a Teflon scraper and scoop by dividing the table into rectangular portions for replicate sample collection. Four replicate samples were collected for this study and the table area cleared to fill each bottle was recorded. Prior to their first use and between snow sample collections, both snow tables were fully cleared of all remaining snow and cleaned with methanol. Samples were collected as soon after the end of each snowfall as feasible, conditions allowing. From September 14th, 2014 to June 1st, 2015, 59 sets of snow samples were collected. When insufficient snow volume was available for complete collection, a subset of the replicate samples was collected as listed in supplemental Table S1. The interval between collections varied based on snowfall frequency, ranging from 1 to 19 days with an average of 4 days. The table area and collection period length associated with each sample allowed the measured concentration of each analyte to be converted to a flux. New sample bottles were used for the collection campaign and each bottle was thoroughly cleaned prior to use. Bottles and lids were soaked in 5% nitric acid, 1% detergent in water (Alconox), and deionized 18.2 MΩ water (DIW), allowing 10 to 14 hours for each soak. Each dried bottle was then sealed in a protective plastic sleeve until use. At Alert, plastic outer gloves and lab coats were used to minimize contamination during collection, and the scraper and scoop were cleaned with DIW prior to each collection. The collection of fresh snow samples reduces the impact of snow sublimation and/or melt as well as the movement of chemical species between snow and air, which can be a concern for snowpack sampling; however, some bi-directional exchange between snow and atmosphere is unavoidable ~~and natural~~ within natural snowpack [and still expected to smaller](#)

[extent on the snow table systems](#). Also, the collection of samples from a snow table eliminated the [difficulty in distinguishing the fresh stratigraphic snow layer from aged layers below](#)~~need to estimate the depth of fresh snow~~, a source of uncertainty for traditional surface snow sampling. This ability to assign a well-defined deposition area and time period to each sample was an advantage over traditional sampling campaigns of aged snowpack. However, both this and traditional snow collection techniques are prone to the uncertainty introduced by the redistribution of snow by winds. Measurements of snowfall accumulation were not available for the collection site. Snow depths measured at the Alert ECCC station indicate that the snow collected on the tables may have underestimated the total snowfall volume by a factor of approximately 1 to 10; however, the meteorological station and collection site were separated by over 6 km with a 50 m difference in elevation, and there was significant disagreement between operator records of weather and that indicated by the meteorological station (see supplemental S4.2 for details). Thus, it was unclear whether this disagreement was the result of snow loss from the snow table or the natural spatial variability in precipitation, and no correction was applied to the collected snow depth. Furthermore, it should be noted that dry deposition via the filtration of air as it is pumped through the snowpack (as described in Harder et al., 1996) may differ between snow on a snow table and that on the surface.

2.2 Campaign Meteorological Conditions

Alert station operators recorded the collection conditions for each sample. Atypical snowfall events were noted: diamond dust events, small crystalline snowfalls; and blowing snow events, periods when high winds potentially resuspended snow from the ground. Operators also made note of any unusual weather conditions such as fog or blizzard conditions. Local ground-level meteorological conditions were monitored by the Alert ECCC stations, approximately 6 km NNE of the collection site (station IDs 2400306, 2400305, and 2400302; retrieved Nov. 2015 from climate.weather.gc.ca). In addition to ground-level meteorological information, vertical profiles were monitored via 6 to 12-hour radiosondes. The radiosonde data were used to estimate mixing height and cloud height over the campaign. Mixing height was taken as the lowest altitude corresponding to an inflection point in the potential temperature. If the potential temperature gradient did not change from negative to positive within the lowest 3 km, no mixing height was found. The vertical humidity profiles were used to identify cloud height as the lowest altitude, within 3 km of the surface, with 100% relative humidity. When 100% humidity was not reached, this criterion was relaxed to 95%. Details of meteorology data are provided in the supplemental section S4.2.

2.3 Snow Sample Preparation and Analysis

All snow samples were kept frozen prior to analysis, throughout storage and shipping. A broad suite of analytes was quantified using replicate snow samples from each snowfall: BC, major ions, and metals. Detailed procedures are provided in the supplemental S2.

Refractory BC quantification was completed via single-particle soot photometry (SP2) as per McConnell et al. (2007). Briefly, melted and sonicated snow samples were atomized via Apex-Q nebulizer and dried particles with 0.02 to 50 fg BC

were quantified via SP2. Observed BC mass distributions did not suggest significant underestimation of the total BC mass due to this size cut-off. A quality control standard and analysis blank were analyzed for every batch of 17 samples.

Major ions were measured via ion chromatography (IC) at ECCC, as per Toom-Saunty and Barrie (2002). Briefly, melted samples were quantified using a Dionex IC: DX600 for anions and cations, ICS2000 for organic acids. Aliquots of these samples were also used for pH analysis (Denver pH analyzer). Equipment was calibrated daily and quality control runs completed every ten samples.

Metals analysis was completed via inductively coupled plasma mass spectrometry (ICP-MS) at the University of Toronto. Briefly, melted samples were filtered to separate insoluble and soluble metals (considered as that which was retained or passed through a 0.45 μm cellulose acetate filter, respectively). Both filtrate and filter were digested using 70% nitric acid, ultra-trace grade (SCP Science PlasmaPure), and filter digestion was augmented using a microwave digester (CEM MARS 6). Centrifuged samples were then quantified via ICP-MS (Thermo Scientific iCAP Q). A performance test and calibration (SCP Science PlasmaCAL QC Std 4) were completed prior to each run, and quality control checks were completed every ten samples. Also, an internal standard was included to quantify and correct for any instrument drift or inter-sample variability (SCP Science Int. Std. Mix 1). All sample preparation was completed in a class 100 vertical laminar flow cabinet (AirClean Systems AC 632).

Quality assurance is of the upmost importance in the analysis of dilute Arctic samples. Instrument accuracy was confirmed through the analysis of certified reference materials. The uncertainty of each measurement was estimated based on analysis detection limits and reproducibility; details are provided in supplemental Sect. S2 (as per Reff et al., 2007; Norris et al., 2014). Also, the signal-to-noise (S/N) of each analyte was calculated to indicate the strength of each measurement, with a S/N value over one considered to be strong (Norris et al., 2014). Regular analysis of blanks was used for background subtraction and to define method detection limits (MDL) as three standard deviations of the blank levels. Beyond typical preparation blanks, which used DIW in the place of snow melt water, field blanks were also analyzed. Once per month, a set of empty sample bottles was taken to the snow table, opened, and resealed without collection. These field blank bottles were stored and shipped with the regular samples and rinsed with DIW to quantify any contamination throughout the sampling process. Any influence from the local Alert base camp was identified using local wind records and the activity logs of the base camp personnel. The only analytes that showed a potential influence from base camp winds were crustal metals, with Pearson's correlation coefficients (R) of 0.4 to 0.6 (p-value 0.0001 to 0.002) between snow mixing ratios and periods of base camp winds. Base camp combustion activity logs showed no significant impact on the samples.

2.4 Atmospheric Monitoring

Ground-level atmospheric monitoring data from the Alert Global Atmospheric Watch Observatory were provided by ECCC (see supplemental S3 for details). Atmospheric BC was monitored hourly by SP2 (Droplet Measurement Technology) (as per Schroder et al., 2015) and major ions by IC of 6 to 8-day high-volume filters of total suspended particles (Hi-Vol) (as per Sirois and Barrie, 1999). Both the SP2 and Hi-Vol were operational throughout the campaign with coverages of 92% and

94%, respectively. As in the snow SP2 analysis, the influence of BC particle size cut-off limits was considered but found to have minimal impact, especially when considered at a monthly scale.

2.5 Transport Modelling

The Lagrangian particle dispersion model FLEXPART (Stohl et al., 2005) was used to determine the source region of air masses that were measured over Alert. This model has previously been shown to be an effective tool for the prediction of transport pathways into and within the Arctic (e.g., Paris et al., 2009). The simulations were driven using meteorological analysis data from the European Centre for Medium-Range Weather Forecasts with a horizontal grid spacing of 0.25° in longitude and latitude and 137 levels in the vertical. For each 5 day period during the measurement period we released virtual tracers over Alert in four different altitude levels, 100 m, 500 m, 1000 m, and 2000 m above sea level, to distinguish between the levels which may be scavenged by snow. The tracers were then followed for ten days backward to obtain the source region for the particular time period. The FLEXPART results were used to explore the dominant source regions associated with each sample. As a simple quantification of the variability in source region, FLEXPART trajectories were summarized by the observed southern limit of transport. This southern limit was calculated as the latitude which encircles 98% of the 10-day transport source area. Values were calculated using each of the four initialization altitudes.

15 3 Results and Discussion

3.1 Total Deposition of Arctic Snowfall Events

Each sample for this study was collected fresh after a known time and over a known area. Given that the snow tables were exposed to the ambient atmosphere for the entirety of each collection period, the measured deposition is considered to represent the total deposition (wet and dry) for said period; however, it is known that surrogate surfaces do not provide an exact proxy for the deposition which would be seen to a natural snow surface (Ibrahim, Barrie, and Fanaki, 1983; Davidson et al., 1985a; Hicks, 1986). There are two additional caveats to this assumption. Firstly, dry deposition at the beginning of each period would fall directly on the exposed clean table rather than onto previously deposited snow. It is unknown what impact these different surface characteristics could have had on the initial deposition rate and collection efficiency. Thus, there is additional uncertainty in the capture of initial dry deposition to the bare table. Secondly, strong winds can disturb and redistribute the snowpack and cause snow to be blown off and/or onto the snow table. Alert operators recorded four occasions when the snowpack was observed to be resuspended due to high winds and these were excluded from the presented results. The dates of these blowing snow events are noted in Table S1 as are missed collections.

The observed snow mixing ratios and fluxes are summarized in Table 1 and Figure 1 for measured analytes with a strong S/N. Mixing ratio is reported as parts per billion by mass (ppb) with the exception of pH. Flux is reported on a per day basis to take into account the differing collection period lengths; however, it should be noted that this length corresponds to the entire collection period (i.e., the number of days between clearing the snow table), not just the length of time when snow was

actually falling. A full record of the measured deposition over the campaign is provided in the supplemental (Table S1-6) along with the associated uncertainties and notes of atypical collection conditions. It should be noted that although IC measurements are provided as the measured ions throughout the discussion, these analytes may not necessarily exist in the dissociated ionic form in the environment. Also, the metal measurements provided in Table 1 are total values, insoluble and soluble. The soluble fractions differed by analyte and by date and are provided in the supplemental (Table S4-6). The metal measurements can be roughly classified into three categories: predominantly insoluble analytes Fe and Al and ~~Fe~~ (>50% insoluble over full campaign); variably soluble/insoluble analytes Co, V, As, Cu, Pb, Cu, Ba, Ti, Mn, K, Cd, and Mg; and predominantly soluble analytes Ca, Cr, Co, and Na (<50% insoluble) (in order from least to greatest average soluble fraction), excluding analytes with insufficient soluble or insoluble measurements above MDL.

10 **Table 1: Overview of fresh snow composition and inferred fluxes over the 2014-15 winter season.**

Analysis	Analyte	Snow Mixing Ratio (ppb)			Snow Flux ($\mu\text{g}/\text{m}^2/\text{d}$)		
		25 th Percentile	50 th Percentile	75 th Percentile	25 th Percentile	50 th Percentile	75 th Percentile
SP2	BC	1.3	2.3	4.1	0.24	0.42	0.86
IC	MS A	<1.9	<1.9	2.5	<0.1	<0.1	<0.1
	ACE	9.6	19.9	27.3	1.9	3.5	7.5
	PRP	<1.5	2.2	5.3	<0.10	0.62	2.06
	FOR	8.4	11.0	14.8	1.32	2.66	4.72
	Cl ⁻	132	249	605	35.6	59.2	122.5
	Br ⁻	<5.0	<5.0	12.1	<0.3	<0.3	2.0
	NO ₃ ⁻	85.4	152.5	265.8	10.8	23.8	50.2
	SO ₄ ²⁻	204	297	554	32.4	69.9	132.5
	C ₂ O ₄ ²⁻	<18.0	<18.0	20.8	<1.2	0.2	2.7
	Na ⁺	55.4	110.7	237.9	10.9	20.1	52.4
	NH ₄ ⁺	10.6	12.4	16.6	1.3	2.5	5.9
	K ⁺	8.0	15.4	23.5	0.8	2.0	3.9
	Mg ²⁺	22.3	43.3	77.4	2.2	7.6	13.3
Ca ²⁺	<133.1	193.1	409.2	<9.1	14.6	51.5	
pH Analyzer	H ⁺ (pH)	1.39 (5.16)	4.25 (5.37)	6.97 (5.86)	0.25 n/a	0.88 n/a	1.87 n/a
ICP-MS	Mg	18.2	28.6	67.6	3.0	6.5	14.5
	Al	<3.2	7.2	19.2	<0.2	1.4	3.3
	V	0.006	0.012	0.086	0.002	0.003	0.014
	Mn	0.23	0.64	1.14	0.06	0.10	0.20
	Fe	3.6	10.8	29.1	0.5	2.0	3.9
	Co	<0.002	0.004	0.011	<0.0002	0.0007	0.0015
	Cu	<0.02	0.05	0.28	0.001	0.010	0.053
	As	0.007	0.044	0.071	0.002	0.006	0.013
	Se	0.010	0.024	0.058	0.002	0.004	0.010
	Sb	0.004	0.010	0.018	0.001	0.002	0.004
	Tl	<0.0001	0.0001	0.0004	<7.2×10 ⁻⁶	15.6×10 ⁻⁶	54.0×10 ⁻⁶
	Pb	0.05	0.25	0.41	0.012	0.039	0.086

Notes: BC = black carbon; MSA = methanesulphonate; ACE = acetate; PRP = propionate; FOR = formate; $C_2O_4^{2-}$ = oxalate; NH_4^+ = ammonium.
< # indicates measurement is below MDL.




Figure 1: Measured snow mixing ratio (line) and uncertainty (shaded area) of key analytes over 2014 to 2015 campaign.

A review of existing Arctic snow measurements found the measured median mixing ratios to fall within expected ranges (see supplemental Table S57 for details); however, it should be noted that the referred data represent a variety of collection and analysis techniques. In general, measurements of this campaign showed salt species and non-crustal metals to be at the lower end of the typical range while SO_4^{2-} and NO_3^- were at the higher end. A limited number of seasonal snow collection campaigns were available for comparison of the observed seasonal trend in analytes. The winter peak in BC was similar to that observed by Davidson et al. (1993); however, spring values observed in this campaign were higher than previously seen. The observed seasonal trend in major ions was generally consistent with existing literature (Davidson et al. 1993; Toom-Sauntry and Barrie, 2002; Dibb et al., 2007). Specifically, a winter peak in sea salt, fall/spring peaks in MSA , and a winter peak in NO_3^- are all typical. However, the fall peak observed in the NSS-SO_4^{2-} mixing ratio and spring peak in NO_3^- were unlike seasonal trends observed previously (Davidson et al. 1993; Toom-Sauntry and Barrie, 2002). Unusual weather events, as noted by the operators, are highlighted in Figure 1; however, no obvious relationship was observed with snow measurements, with the exception of blizzard and high wind conditions in January and February which were associated with elevated mixing ratios for several chemical species. Atmospheric measurements are provided in the supplemental Table S7. As observed for snow, most atmospheric analytes experienced a winter high. The fall/spring peaks in MSA and notable fall peak in SO_4^{2-} persisted in both snow and atmospheric measurements.

3.2 Factors Influencing Snow Scavenging and Deposition

3.2.1 Effective Deposition Velocity

As discussed above, the collected snow samples provide information on the total deposition of material to the surface over a given time and area. In order to elucidate the mechanisms controlling this bulk deposition, a simplistic model for flux, Eq. (1), was adopted to describe the measured deposition:

$$F_{S,total}^{ij} = C_A^{ij} v_{d,eff}^{ij}, \quad (1)$$

where $F_{S,total}^{ij}$ is the flux deposited to snow of the j^{th} analyte over the i^{th} period; C_A^{ij} is the arithmetic average atmospheric concentration; and $v_{d,eff}^{ij}$ is the effective deposition velocity.

The measured snow flux ($F_{S,total}$) represents the total deposition by wet and dry mechanisms. Thus, the effective deposition velocity ($v_{d,eff}$) encompasses the variety of aerosol and meteorological properties controlling deposition efficiency and the relative importance of each deposition pathway: dry, wet in-cloud, and wet below-cloud scavenging. This effective velocity is akin to a typical deposition velocity; however, it encapsulates the bulk movement of material by combining the dry deposition velocity and the wet deposition efficiency as an equivalent velocity. Thus, this parameter can be used to provide a holistic view of Arctic deposition. Use of the effective deposition velocity also avoids the uncertainties of estimating the split between dry and wet deposited mass. A caveat to this analysis is that the three deposition mechanisms relate to different atmospheric concentrations, a gradient which is not necessarily captured when the ground-level atmospheric concentration


(C_A) is used to calculate the effective deposition velocity: dry deposition affects the lower atmosphere, in-cloud scavenging the cloud layer, and below-cloud scavenging the full below-cloud atmospheric column. Previous observations of vertical profiles in the Arctic have shown notable variability with altitude (Hansen and Rosen, 1984; Leaitch et al., 1989; Spackman et al., 2010; Brock et al., 2011; Sharma et al., 2013). So, the calculated effective velocity includes an intrinsic variability
5 dependent on the vertical atmospheric profile of each analyte.

Effective deposition velocities were calculated for chemical species measured in both snow (SP2 and IC) and atmospheric (SP2 and Hi-Vol) samples. Figure 2 shows effective deposition velocities calculated as the ratio of total summed snow flux and average atmospheric concentration measured over the same period. Both a six-day resolution, as per the Hi-Vol sampling frequency, and monthly resolution are provided. The calculated effective deposition velocities ranged from 0.001
10 to 10 cm/s (0 to 16 cm/s with uncertainty). Episodic and monthly peaks are observed in Figure 2 for each analyte. The variance in deposition observed by composition and temporally is discussed below.

3.2.2 Variation in Deposition by Composition

Monthly effective deposition velocities were used to contrast deposition mechanisms by aerosol composition. A monthly resolution provides insight into the general deposition regime of each analyte, highlighting the impact of bulk deposition
15 characteristics rather than event-specific variability. The variability between aerosol of different composition and the influences of seasonal changes within the Arctic system are simpler to identify without the interference of variability between event-specific conditions. A monthly analysis also facilitates future comparison with modelled results which may not replicate individual events. January and February, 2015, were excluded from the monthly analysis because blizzard and high wind conditions were believed to have caused significant losses of snow from the snow tables during these months
20 (based on operator reports), which would lead to underestimation of these snow flux values. The effective deposition velocity is best suited to analysis across periods of equal length and precipitation volume, since both of these parameters are inherently included when the wet deposition efficiency is converted to an equivalent deposition velocity. With the exception of January and February, the total monthly snow precipitation over the campaign was relatively constant, with a relative standard deviation of 20%.

25 Figure 3 shows that the typical deposition characteristics varied by analyte, with median effective deposition velocities ranging from 0.03 to 1.1 cm/s; however, error bars describing the combined uncertainty of the snow and atmospheric measurements show that the range of calculated effective deposition velocities of each analyte have considerable overlap. The calculated monthly effective deposition velocities showed a relative standard deviation of 60 to 150% across the measured analytes, while measurement-based uncertainty was estimated as only $\pm 35\%$, indicating a significant impact of
30 aerosol state or properties on deposition.



2014 to 2015 Winter Season

Figure 2: Effective deposition velocity at monthly (solid) and approximately 6-day (dashed) frequencies with 6-day uncertainties (shaded area). Missing values indicate periods with snow and/or atmospheric measurements below detection limits.




Figure 3: Monthly effective deposition velocities by composition (points, excluding January and February). The median of each analyte (bar) and full range with uncertainty (error bar) are also shown. Also shown is the typical range of dry deposition velocity for accumulation mode particles to snow by others (Davidson et al., 1987; Petroff and Zhang, 2010).

The variability observed between analytes may be the result of variations in aerosol properties. First, the measured chemical species differ in terms of dominant phase: BC, NH₄⁺ (ammonium), SO₄²⁻, Na⁺, K⁺, Mg²⁺, C₂O₄²⁻ (oxalate), and Ca²⁺ are typically observed predominantly in the particle phase within the Arctic, while MSA, Br⁻, Cl⁻, and NO₃⁻ and their associated precursors can have appreciable gas-phase portions (Barrie and Hoff, 1985). In general, analytes considered to be dominantly particle-based were seen to exhibit lower deposition velocities than those which may have existed as gases. This observation is supported by a detailed study of sea salt species. The Cl in salt particles has been previously observed to partition to the gas phase over winter months (e.g., Barrie and Hoff, 1985; Toom-Sauntry and Barrie, 2002; Quinn et al., 2009); thus, Arctic salt particles often show a Cl deficit while gaseous Cl shows enhancement. Such a process was corroborated by comparison of the atmospheric Hi-Vol and snow IC measurements for Na⁺ and Cl⁻. Both atmospheric and snow measurements showed high correlation of Na⁺ and Cl⁻ analytes, 0.92 and 0.99, respectively. Yet, the ratio of Cl⁻/Na⁺ differed between measurement mediums with a higher proportion of Cl⁻ observed in snow (Cl⁻/Na⁺=2.49 mass/mass) than in atmosphere (Cl⁻/Na⁺=1.36). Compared to the typical marine ratio (Cl⁻/Na⁺=1.795; Pytkowicz and Kester, 1971), the atmosphere showed a deficit in Cl⁻ while snow showed enhanced Cl⁻ (see supplemental Figure S4). It is expected that the Hi-Vol measurement technique would collect predominantly atmospheric particles, while snow would scavenge both gaseous and particulate aerosol. Thus, the enhanced Cl⁻ in snow above that of sea salt indicates that Cl partitioned to the gas phase, that both phases deposited to snow, and that gas-phase Cl was scavenged to snow preferentially. Assuming all Na⁺ and Cl⁻ originated as sea salt particles, the Cl mass in from each phase can be estimated in the atmosphere and snow. Based on these estimated proportions in the snow and atmosphere, median effective deposition velocities were calculated as 0.16 and 0.40

cm/s for the particle and gas phases, respectively. Comparison of these velocities indicates an 86% enhancement in gas-phase deposition relative to that of the particle phase. Thus, gaseous scavenging may contribute to the enhanced bulk deposition observed in MSA, Br⁻, Cl⁻, and NO₃⁻.

Particle nucleation affinity may also be a significant contributor to the observed differences in bulk deposition. The lowest velocities were observed for BC, NH₄⁺, and SO₄²⁻, which largely fell within the typical range of deposition velocities seen for dry deposition: 0.02 to 0.10 cm/s for typical accumulation mode Arctic particles (Davidson et al., 1987; Petroff and Zhang, 2010), as highlighted in Figure 3. In contrast, the monthly velocities of other particle-dominated chemical species, Na⁺, K⁺, Mg²⁺, C₂O₄²⁻, and Ca²⁺, all fell above the typical dry deposition range. Particles containing these analytes, as well as NH₄⁺ and SO₄²⁻, have been previously suggested to act as better precipitation nuclei than BC (e.g., Zobrist et al., 2006; Hoose and Möhler, 2012; Farmer, Cappa, and Kreidenweis, 2015); thus, their enhanced velocities might be attributed to ~~either enhanced scavenging via cloud formation or~~ in-cloud scavenging. Thus, dry deposition may be the dominant deposition mechanism in some measurements while others show enhanced deposition which likely reflects an increased contribution of wet deposition processes. The observed difference in deposition velocities also implies that BC/SO₄²⁻/NH₄⁺-containing particles are to a significant extent externally mixed from these other constituents.

Furthermore, salt and crustal particles may experience enhanced deposition since they typically consist of coarser particles than BC, SO₄²⁻, and NH₄⁺. Specifically, coarse mode particles can exhibit dry deposition velocities to snow up to 0.6 cm/s and below-cloud scavenging efficiencies enhanced over accumulation mode particles by a factor of 10 (Zhang and Vet, 2006; Petroff and Zhang, 2010). The enhancement of Ca²⁺ above other crustal-related analytes is unexpected. This cannot be satisfactorily explained without further study; however, the phenomenon may be connected to the dissimilar sources of Ca-rich mineral dust to that of other crustal species suggested by other studies (Banta et al., 2008). In addition to aerosol phase, nucleation affinity and size as discussed above, particle coating could also impact the scavenging and deposition process. However, the potential influence of coatings on the observed velocities cannot be addressed with the available information.

3.2.3 Temporal Variability of Deposition

Shared temporal trends in effective deposition velocity can be observed in Figure 2. A general trend of heightened deposition in the fall and spring can be observed across all analytes. In particular, BC, Na⁺, Mg²⁺, and Ca²⁺ can be seen to share a similar seasonal and episodic trend, with Pearson's correlation coefficients of 0.7 to 0.9 (comparing six-day resolution, excluding January and February, 2015, p-values <0.001). Episodic peaks in Mg²⁺ and Ca²⁺ also show some similarity to SO₄²⁻ (correlation of 0.70-0.85, p-value <0.001). The trend of NH₄⁺ is more difficult to distinguish as September measurements were below detection limit; however fall and spring peaks are suggested. A more pronounced fall peak was observed in SO₄²⁻, Na⁺, K⁺, and Cl⁻, though their episodic peaks differ. Although MSA is distinguished by a spring peak, MSA and SO₄²⁻ share similar episodic peaks with a six-day correlation of 0.86 (p-value <0.001). The seasonal trends of Br⁻, C₂O₄²⁻, and NO₃⁻ are more distinct: Br⁻ exhibits a broad spring peak, C₂O₄²⁻ a short October peak, and NO₃⁻ a broad fall/winter peak. Episodic C₂O₄²⁻ showed moderate correlation with BC, Na⁺, Mg²⁺, and Ca²⁺ (coefficient=0.5-0.7, p-

value < 0.02) and NO_3^- episodes showed correlation with K^+ (coefficient = 0.83, p-value < 0.001). There are some peak events shared across several analytes, for example, an early October peak is observed in most chemical species. While the differing magnitudes of effective deposition velocities observed across analytes implies separate scavenging and deposition, these shared temporal trends indicate that the externally mixed particles and gaseous chemical species are subject to similar temporal influences controlling deposition.

Several factors controlling deposition experience seasonal variations, which may have contributed to the observed inter-monthly variability. Six properties of the Arctic system with seasonal trends were considered as possible influences on the observed velocity trend: precipitation, temperature, mixing height, cloud height, dominant aerosol source region, and sunlight availability, as shown in Figure 4. The precipitated snow-water equivalent depth was calculated from the snow mass and table area of each sample. Temperature was monitored at local ground-level meteorological stations over the campaign (supplemental Table S9) and sunlight estimated from location and time of year. The dominant aerosol source of each month was described using the southern limit to transport and mixing/cloud heights were estimated from radiosonde data, as described above.

Temperature, transport, and sunlight can be seen to follow similar seasonal trends with fall/spring peaks. Precipitation, mixing height, and cloud height exhibit episodic peaks and a less significant seasonal trend (relative standard deviation intra-monthly was a factor of 1.5 to 2 times higher than inter-monthly, whereas these values were approximately equal for temperature, transport, and sunlight). When compared to effective deposition velocities, BC, SO_4^{2-} , Na^+ , Mg^{2+} , NO_3^- , and Ca^{2+} show better correlation with temperature than the other meteorological conditions shown in Figure 4, with Pearson's correlation coefficients above 0.5 (excluding January and February, 2015, p-value < 0.02). Specifically, SO_4^{2-} , K^+ , Mg^{2+} , and Ca^{2+} had high correlation coefficients with temperature of 0.6-0.7 (p-values < 0.002). Medium to high correlations were observed between mixing height and SO_4^{2-} , Na^+ , K^+ , Mg^{2+} , and NO_3^- , with coefficients of 0.4-0.7 (highest correlation for K^+ at 0.76) and p-values below 0.05. Bromide deposition showed high correlation with sunlight (coefficient: 0.6, p-value: 0.001). In contrast, NH_4^+ and MSA only showed weak correlation with the described meteorological parameters (maximum coefficients of about 0.4 with transport and mixing height, respectively). In contrast, $\text{C}_2\text{O}_4^{2-}$ and Cl^- did not show strong correlations with any of the described meteorological parameters (coefficients below 0.2). Overall, temperature showed the best correlation with deposition velocities for most analytes, followed by mixing height. Thus, changes in temperature may be linked to seasonal changes in the dominant scavenging mechanism of several analyzed species. Specifically, it is hypothesized that the increased presence of mixed-phase clouds with warmer temperatures may have enhanced wet deposition via in-cloud scavenging due to CCN activity for those analytes expected to exist predominantly in the particle phase: BC, NH_4^+ , SO_4^{2-} , Na^+ , K^+ , Mg^{2+} , NO_3^- , and Ca^{2+} . [This aligns with seasonal trend in \$\text{SO}_4^{2-}\$ deposition observed by Davidson et al. \(1985b and 1987\) and the suggested link to deposition mechanism.](#)

The effective deposition velocities for warmer and colder months were separated using a -20 °C cut-off between ice-cloud dominated periods, November to April (N/D/Mr/A), and months with a significant potential for mixed-phase clouds, September, October, and May (S/O/My). Radiosonde observations show temperatures at typical cloud heights usually above




Figure 4: Seasonal variation in precipitation (a), temperature (b), mixing height (c), cloud height (d), transport (e), and sunlight (f). Precipitation and sunlight are presented as the average per snow sample collection period. Temperature, mixing/cloud height, and southern transport limit are presented as the average for each collection period (line) along with the full range (shaded area).

5

-20 °C in S/O/My and below -20 °C in N/D/Mr/A, supporting the mixed-phase cloud hypothesis. Comparison of the effective deposition velocities delineated by these periods minimized the influence of precipitation volume as it was relatively constant: an average of 12 mm/month in S/O/My and 18 mm/month in N/D/Mr/A. However, the influence of aerosol source may differ by period given their distinct source profiles: long-range transport dominated in N/D/Mr/A, and local transport dominated in S/O/My. Figure 5 depicts the range of effective deposition velocities calculated for each analyte

10

over these two periods (again excluding January and February). Bromide was excluded from this analysis since it was below detection limit in the snow and/or atmospheric measurements from September to November.




Figure 5: Effective deposition velocities split by season. The effective deposition velocities are separated into two time periods: warmer months S/O/My = September/October/May, and colder months N/D/Mr/A = November/December/March/April.

With the exception of C₂O₄²⁻, all analytes showed a larger median effective deposition velocity for the warmer S/O/My months than the colder N/D/Mr/A months. Although insufficient data are available to confirm the statistical significance for each individual analyte, the combined normalized dataset of velocities showed significant enhancement during S/O/My using the ANOVA test (p-value 4.9×10⁻⁸). Specifically, marked enhancement was seen for BC, NH₄⁺, SO₄²⁻, Na⁺, K⁺, and Mg²⁺. The cold-month deposition of these analytes can largely be described by dry deposition alone with velocities below 0.1 cm/s for analytes expected to be dominated by accumulation mode particles and 0.6 cm/s for those expected to include significant coarse-mode mass; however, their warm-month effective deposition velocities were a factor of 2 to 12 higher.

The unexpected discrepancy in the deposition of Ca²⁺ and Mg²⁺ given their typically similar origins, as discussed above, was also observed when comparing median warm and cold month periods. Both Mg²⁺ and Ca²⁺ are common to sea salt and crustal origins but their non-sea salt (NSS) portions can be estimated based on typical sea salt ratios with Na⁺ (Pytkowicz and Kester, 1971). The NSS-Mg²⁺ and NSS-Ca²⁺ showed similar behaviour over the warm months, with average velocities of 2.4 and 2.6 cm/s, respectively, but a larger difference in the cold months, with average velocities of 0.4 and 1.0 cm/s, respectively. No explanation of this discrepancy can be supported without further study; however, it appears that Ca²⁺ and Mg²⁺ exist as externally mixed Ca-rich and Mg-rich crustal particles which are subject to differing deposition processes, particularly in dry deposition-dominated colder months.

Thus, the observed effective deposition velocities suggest that most analytes and particularly those expected to exist primarily as particle phase were preferentially scavenged during the warmer S/O/My months, possibly due to the presence of

5 mixed-phase clouds and the associated CCN activation of these chemical species or enhanced below-cloud deposition of those compounds typically associated with larger particles. However, the change in source profile typically experienced during these months along with other seasonal changes in aerosol processing and altitudinal distribution might have also contributed to the observed S/O/My enhancement. In particular, records of volcanic activity show that the Icelandic volcano Bárðarbunga was active August, 2014 through February, 2015 (Global Volcanism Program, retrieved March 2016 from <http://volcano.si.edu/>), which likely contributed to a shift in the dominant source and scavenging-related properties of SO_4^{2-} over the campaign that would not be representative of a typical year. [Thus, the \$\text{SO}_4^{2-}\$ observations of this campaign, especially the peak snow mixing ratio seen in the fall, may not reflect a seasonal trend for typical Arctic Haze.](#) The overlapping warm/cold-month ranges observed for Cl^- , MSA, $\text{C}_2\text{O}_4^{2-}$, and NO_3^- suggest that the deposition of these chemical species was more strongly driven by factors other than nucleation, such as gas-phase partitioning or other aerosol aging processes. For example, enhanced deposition of MSA and Br^- was observed as early as March and April, which could imply that their deposition was impacted by changes in the atmospheric processing of these chemical species during polar sunrise.

4 Conclusions

15 To help characterize the chemical state of the rapidly changing high Arctic, an intensive campaign of fresh snow sampling at Alert, Nunavut, was completed and snow quantified for a broad suite of analytes. Comparison of these snow measurements with coincident atmospheric measurements allowed estimation of monthly effective deposition velocities describing the total dry and wet deposition in the range of about 0.02 to 8 cm/s. The calculated effective deposition velocities for several measured chemical species resemble those expected for dry deposition alone, suggesting that dry deposition may be the dominant removal mechanism, especially for winter scavenging of BC, NH_4^+ , SO_4^{2-} , Na^\pm , and K^\pm . Enhanced deposition during September, October, and May suggests that wet deposition may increase in importance during these warmer months, possibly due to the presence of mixed-phase clouds and the associated scavenging of crustal, salt, and SO_4^{2-} species as CCN; however, other factors such as changes in the dominant aerosol source profile may also contribute to the observed trend. Comparison of salt species measurements in the Arctic snow and atmosphere suggested that Cl experiences significant gas-phase partitioning and that this gas phase may be preferentially scavenged. Such gas-phase scavenging may contribute to the enhanced deposition of MSA, Br^- , Cl^- , and NO_3^- observed relative to BC, in conjunction with other aerosol processing differences. The low deposition velocity of BC-containing particles is consistent with those particles being externally mixed from more soluble species and having a low cloud nucleation efficiency. Given the rarity of temporally-refined and broadly speciated Arctic snow sampling campaigns, measured deposition magnitudes and insights on deposition mechanisms such as these are valuable for future model validation.

Author Contribution

Organization of the snow collection campaign was led by S. Sharma with the assistance of A. Platt and sample collection by M. Elsasser. Snow SP2 analysis was completed by J. McConnell and N. Chellman, snow IC analysis lead by D. Toom with the assistance of A. Chivulescu, and snow ICP-MS by K. Macdonald with the assistance of Y. Lei. Analysis of radiosonde data was completed by D. Tarasick. Ambient atmospheric monitoring of inorganic aerosols was completed by D. Toom and monitoring of BC by S. Hanna with the assistance of A. Bertram. FLEXPART simulations were completed by H. Bozem and D. Kunkel with data analysis assisted by K. Macdonald. Data interpretation was led by K. Macdonald with input and comments by all authors. G. Evans and J. Abbatt provided oversight for the project, including input on the manuscript.

Competing interests

10 The authors declare that they have no conflict of interest.

Acknowledgements

Funding of this study was provided as part of the Network on Climate and Aerosols Research (NETCARE), Natural Science and Engineering Research Council of Canada (NSERC), the government of Ontario through the Ontario Graduate Scholarship (OGS), and Environment and Climate Change Canada. This project would not have been possible without the collaboration of many skilled individuals: Richard Leaitch at Environment Canada; and Catherine Philips-Smith and Cheol-Heon Jeong at the University of Toronto.

References

- AMAP: Snow, water, ice and permafrost in the Arctic: Climate change and the cryosphere, Arctic Monitoring and Assessment Programme, Oslo, Norway, 2011.
- 20 Banta, J. R., McConnell, J. R., Edwards, R., and Engelbrecht, J. P.: Delineation of carbonate dust, aluminous dust, and sea salt deposition in a Greenland glaciochemical array using positive matrix factorization, *Geochem. Geophys. Geos.*, 9 (7), 1–19, doi:10.1029/2007GC001908, 2008.
- Barrie, L. A., and Hoff, R.M.: Five years of air chemistry observations in the Canadian Arctic, *Atmos. Environ.*, 19 (12), 1995–2010, doi:10.1016/0004-6981(85)90108-8, 1985.
- 25 Bergin, M. H., Jaffrezo, J.-L., Davidson, C. I., Dibb, J. E., Pandis, S. N., Hillamo, R., Maenhaut, W., Kuhns, H. D., and Makela, T.: The contributions of snow, fog, and dry deposition to the summer flux of anions and cations at Summit, Greenland, *J. Geophys. Res.-Atmos.*, 100, D8, 16,275–16,288A, doi:10.1029/95JD01267, 1995.

- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Bernsten, T., Deangelo, B. J., Flanner, M. G., et al.: Bounding the role of black carbon in the climate system: A Scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.
- Brock C. A., Cozic, J., Bahreini, R., Froyd, K. D., Middlebrook, A. M., McComiskey, A., Brioude, J., et al.: Characteristics, sources, and transport of aerosols measured in spring 2008 during the aerosol, radiation, and cloud processes affecting Arctic Climate (ARCPAC) Project; *Atmos. Chem. Phys.*, 11, 2423–2453, doi:10.5194/acp-11-2423-2011, 2011.
- Davidson, C. I., Lindberg, S. E., Schmidt, J. A., Cartwright, L. G., and Landis, L. R.: Dry deposition of sulfate onto surrogate surfaces, *J. Geophys. Res.*, 90 (D1), 2123–30, doi:10.1029/JD090iD01p02123, 1985a.
- [Davidson, C. I., Santhanam, S. R. C., and Marvin, P. O.: Atmospheric transport and deposition of trace elements onto the Greenland ice sheet, *Atmos. Environ.*, 19 \(12\), 2065-2081, 1985b.](#)
- Davidson, C. I., Honrath, R. E., Kadane, J. B., Tsay, R. S., Mayewski, P. A., Lyons, W. B., and Heidam, N. Z.: The Scavenging of atmospheric sulfate by Arctic snow, *Atmos. Environ. A-Gen.*, 21 (4), 871–882, doi:10.1016/0004-6981(87)90083-7, 1987.
- Davidson, C. I., Jaffrezo, J.-L., Mosher, B. W., Dibb, J. E., Borys, R. D., Bodhaine, B. A., Rasmussen, R. A., et al.: Chemical constituents in the air and snow at Dye 3, Greenland: I. Seasonal variations, *Atmos. Environ. A-Gen.*, 27 (17), 2709–22, doi:10.1016/0960-1686(93)90304-H, 1993.
- [Dibb, J. E., Whitlow, S. I., and Arsenault, M.: Seasonal variations in the soluble ion content of snow at Summit, Greenland: Constraints from three years of daily surface snow samples, *Atmos. Environ.*, 41, 5007-5019, doi:10.1016/j.atmosenv.2006.12.010, 2007.](#)
- Farmer, D. K., Cappa, C. D., and Kreidenweis, S. M.: Atmospheric processes and their controlling influence on cloud condensation nuclei activity. *Chem. Rev.*, 115 (10), 4199–4217, doi:10.1021/cr5006292, 2015.
- Hagler, G. S. W., Bergin, M. H., Smith, E. A., and Dibb, J. E.: A Summer time series of particulate carbon in the air and snow at Summit, Greenland, *J. Geophys. Res.-Atmos.*, 112 (August), 1–12, doi:10.1029/2007JD008993, 2007.
- Hansen, A. D. A., and Rosen, H.: Vertical distributions of particulate carbon, sulfur, and bromine in the Arctic haze and comparison with ground-level measurements at Barrow, Alaska, *Geophys. Res. Lett.*, 11 (5), 381–84, doi:10.1029/GL011i005p00381, 1984.
- Harder, S. L., Warren, S. G., Charlson, R. J., and Covert, D. S.: Filtering of air through snow as a mechanism for aerosol deposition to the Antarctic ice sheet, *J. Geophys. Res.*, 101(D13), 18729–18743, doi:10.1029/96JD01174, 1996.
- Hartmann, D. J., Klein Tank, A. M. G., Rusticucci, M., Alexander, L. V., Brönnimann, S., Charabi, Y. A.-R., Dentener, F. J., et al.: Observations: Atmosphere and surface, *Climate change 2013: The Physical science basis, Contribution of working group I to the fifth assessment report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)], Cambridge University Press, Cambridge and New York, NY, doi:10.1017/CBO9781107415324.008, 2013.

- Hicks, B. B.: Measuring dry deposition: A Re-assessment of the state of the art, *Water Air Soil Poll.*, 30, 75–90, doi:10.1007/BF00305177, 1986.
- Hillamo, R.E., Kerminen, V.-M., Maenhaut, W., Jaffrezo, J.-L., Balachandran, S., and Davidson, C.I.: Size distributions of atmospheric trace elements at Dye 3, Greenland: I. Distribution characteristics and dry deposition velocities, *Atmos. Environ. A-Gen.*, 27 (17), 2787–2802, doi:10.1016/0960-1686(93)90311-L, 1993.
- 5 Hoose, C. and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: A Review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12 (20), 9817–9854, doi:10.5194/acp-12-9817-2012, 2012.
- Ibrahim, M., Barrie, L. A., and Fanaki, F.: An Experimental and theoretical investigation of the dry deposition of particles to snow, pine trees and artificial collectors, *Atmos. Env.*, 17 (4), 781–88, doi:10.1016/0004-6981(83)90427-4. 1983.
- 10 Law, K. S. and Stohl, A.: Arctic air pollution: Origins and impacts, *Science*, 315, 1537–1540, doi:10.1126/science.1137695, 2007.
- Leaitch, W. R., Hoff, R. M., and MacPherson, J. I.: Airborne and lidar measurements of aerosol and cloud particles in the troposphere over Alert Canada in April 1986, *J. Atmos. Chem.*, 9, 187–211, doi:10.1007/BF00052832, 1989.
- Legrand, M. and De Angelis, M.: Origins and variations of light carboxylic acids in polar precipitation, *J Geophys. Res.*, 100 (Di), 1445–1462, doi:10.1029/94jd02614, 1995.
- 15 Liu, J., Fan, S., Horowitz, L. W., and Levy, H.: Evaluation of factors controlling long-range transport of black carbon to the Arctic, *J. Geophys. Res.*, 116, D04307, doi:10.1029/2010JD015145, 2011.
- McConnell, J. R., Edwards, R., Kok, G. L., Flanner, M. G., Zender, C. S., Saltzman, E. S., Banta, J. R., et al.: 20th-Century industrial black carbon emissions altered Arctic climate forcing, *Science*, 317, 1381–1384, doi:10.1126/science.1144856, 2007.
- 20 McMahon, T. A. and Denison, P. J.: Empirical atmospheric deposition parameters: A Survey, *Atmos. Environ.* (1967), 13 (5), 571–585, doi:10.1016/0004-6981(79)90186-0, 1979.
- Morrison, H., Shupe, M. D., Pinto, J. O., and Curry, J.: Possible roles of ice nucleation mode and ice nuclei depletion in the extended lifetime of Arctic mixed-phase clouds, *Geophys. Res. Lett.*, 32 (18), 1–5, doi:10.1029/2005GL023614, 2005.
- 25 Nguyen, Q. T., Skov, H., Sørensen, L. L., Jensen, B. J., Grube, A. G., Massling, A., Glasius, M., and Nøjgaard, J. K.: Source apportionment of particles at Station Nord, north east Greenland during 2008-2010 using COPREM and PMF analysis, *Atmos. Chem. Phys.*, 13, 35–49, doi:10.5194/acp-13-35-2013, 2013.
- Norris, G., Duvall, R., Brown, S., and Bai, S.: EPA Positive matrix factorization (PMF) 5.0 fundamentals and user guide, U.S. Environmental Protection Agency, 2014.
- 30 Paris, J.-D., Stohl, A., Nédélec, P., Arshinov, M. Y., Panchenko, M. V., Shmargunov, V. P., Law, K. S., et al.: Wildfire smoke in the Siberian Arctic in summer: Source characterization and plume evolution from airborne measurements, *Atmos. Chem. Phys. Discussions*, 9, 18201–18233, doi:10.5194/acpd-9-18201-2009, 2009.
- Petroff, A. and Zhang, L.: Development and validation of a size-resolved particle dry deposition scheme for application in aerosol transport models, *Geoscientific Model Development*, 3 (2), 753–769, doi:10.5194/gmd-3-753-2010, 2010.

- Pytkowicz, R. M. and Kester, D. R.: The Physical chemistry of sea water, *Oceanogr. Mar. Biol.*, 9, 11–60, doi:10.1029/WR001i002p00263, 1971.
- Quinn, P. K., Shaw, G., Andrews, E., Dutton, E. G., Ruoho-Airola, T., and Gong, S. L.: Arctic haze – Current trends and knowledge gaps, *Tellus B.*, 59 (1), 99–114, doi:10.1111/j.1600-0889.2006.00238.x, 2007.
- 5 Quinn, P. K., Bates, T. S., Schulz, K., and Shaw, G. E.: Decadal trends in aerosol chemical composition at Barrow, AK: 1976–2008, *Atmos. Chem. Phys.*, 9, 18727–43, doi:10.5194/acpd-9-18727-2009, 2009.
- Reff, A., Eberly, S. I., and Bhave, P. V.: Receptor modeling of ambient particulate matter data using positive matrix factorization: Review of existing methods, *Journal of the Air and Waste Management Association* (1995), 57 (August 2014), 146–154, doi:10.1080/10473289.2007.10465319, 2007.
- 10 Rigor, I. G., Colony, R. L., and Martin, S.: Variations in surface air temperature observations in the Arctic: 1979–97, *J. Climate*, 13 (5), 896–914, doi:10.1175/1520-0442(2000)013<0896:VISATO>2.0.CO;2, 2000.
- Schroder, J. C., Hanna, S. J., Modini, R. L., Corrigan, A. L., Kreidenwies, S. M., MacDonald, A. M., Noone, K. J., et al.: Size-resolved observations of refractory black carbon particles in cloud droplets at a marine boundary layer site, *Atmos. Chem. Phys.*, 15 (3), 1367–1383, doi:10.5194/acp-15-1367-2015, 2015.
- 15 Sehmel, G. A.: Particle and gas dry deposition: A Review, *Atmos. Environ.* (1967), 14 (9), 983–1011, doi:10.1016/0004-6981(80)90031-1, 1980.
- Seinfeld J. H., and Pandis, S. N.: *Atmospheric chemistry and physics – From Air pollution to climate change*, Wiley, Michigan, USA, 2006.
- Sharma, S., Ishizawa, M., Chan, D., Lavoué, D., Andrews, E., Eleftheriadis, K., and Maksyutov, S.: 16-Year simulation of Arctic black carbon: Transport, source contribution, and sensitivity analysis on deposition, *J. Geophys. Res-Atmos.*, 118, 943–964, doi:10.1029/2012JD017774, 2013.
- 20 Shupe, M. D., Matrosov, S. Y., and Uttal, T.: Arctic mixed-phase cloud properties derived from surface-based sensors at SHEBA, *J. Atmos. Sci.*, 63 (2), 697–711, doi:10.1175/JAS3659.1, 2006.
- Sirois, A. and Barrie, L. A.: Arctic lower tropospheric aerosol trends and composition at Alert, Canada: 1980–1995, *J. Geophys. Res.*, 104 (D9), 11599–11618, doi:10.1029/1999JD900077, 1999.
- 25 Spackman, J. R., Gao, R. S., Neff, W. D., Schwarz, J. P., Watts, L. A., Fahey, D. W., Holloway, J. S., et al.: Aircraft observations of enhancement and depletion of black carbon mass in the springtime Arctic, *Atmos. Chem. Phys.*, 10, 9667–9680, doi:10.5194/acp-10-9667-2010, 2010.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note – The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5(9), 2461–2474, doi:10.5194/acp-5-2461-2005, 2005.
- 30 Stroeve, J. C., Serreze, M. C., Fetterer, F., Arbetter, T., Meier, W., Maslanik, J., and Knowles, K.: Tracking the Arctic’s shrinking ice cover: another extreme September minimum in 2004, *Geophys. Res. Lett.*, 32 (4), 1–4, doi:10.1029/2004GL021810, 2005.

Toom-Saunty, D. and Barrie, L. A.: Chemical composition of snowfall in the high Arctic: 1990 – 1994, *Atmos. Environ.*, 36, 2683–2693, doi:10.1016/S1352-2310(02)00115-2 , 2002.

Zhang, L. and Vet, R.: A Review of current knowledge concerning size-dependent aerosol removal, *China Part.*, 4 (6), 272–282, doi:10.1016/S1672-2515(07)60276-0, 2006.

- 5 Zobrist, B., Marcolli, C., Koop, T., Luo, B. P., Murphy, D. M., Lohmann, U., Zardini, A., et al.: Oxalic acid as a heterogeneous ice nucleus in the upper troposphere and its indirect aerosol effect, *Atmos. Chem. Phys.*, 6, 3115–3129, doi:10.5194/acpd-6-3571-2006, 2006.