

Response to reviewers

Alison Ming, Amanda Maycock, Peter Hitchcock and Peter Haynes

March 6, 2017

We would like to thank both reviewers for their detailed comments which have helped us to improve the paper significantly in revision. We have implemented many of the suggestions and provided detailed responses when we have not. In light of the comments from both reviewers about the length of the manuscript, we have revised the text to make the paper more concise.

Page and line numbers quoted in the responses refer to the updated text.

In response to the comments of Reviewer 1 concerning the differences between SEFDH and dynamical calculations we have reexamined the IGCM calculation and modified it slightly to minimise any differences between it and the SEFDH calculation arising from implementation rather than dynamical adjustment.

Below we indicate how we have dealt with each individual comment.

Reviewer 1

What was assumed in this study regarding clouds? Are the results sensitive to clouds, both in the region from 100-130 mbar and below that level?

All calculations in the paper assume clear-sky conditions as stated on page 4, line 15. While it is possible that cloud effects could modify the TTL temperature responses to ozone and water vapour, and as such a careful exploration of this effect would be a valuable contribution, we note that our calculations accounting for the effects of both clear-sky radiative and dynamical heating reproduce well the observed annual cycle in TTL temperature, suggesting that the net effects of clouds (other than those included indirectly in the dynamical heating) are small, justifying this choice of focus.

Nevertheless, in revising the paper we have thought carefully about cloud effects and have performed additional SEFDH calculations to give an indication of the order of magnitude of cloud effects. These calculations show the effect of including annual mean climatological cloud cover is to decrease the peak-to-peak annual cycle temperature change due to ozone at 70 hPa by 5-10% at all latitudes between 20° N and 20° S. The effect on the water vapour annual cycle at the same level is negligible. For this estimate, we used the high, medium and low annual mean cloud fractions from the ISCCP dataset (1983 to 2006). The clouds lead primarily to a reduction in the amount of upwelling longwave radiation reaching 70 hPa of about 0.05 K day^{-1} which in turn decreases the ozone temperature response. Clouds also have an annual cycle in radiative heating. One estimate, at a single location in the tropics, from Fueglistaler and Fu (2006) shows that the annual cycle in radiative heating from clouds is about 0.05 K day^{-1} (peak to peak) at 70 hPa but these values are at present poorly constrained. On the basis of these rough quantitative estimates we consider it unlikely that including cloud effects will invalidate our main arguments and conclusions.

However, we do not feel that it is practical or appropriate to present these estimates in the revised version of the paper – the problem is sufficiently complicated that a brief explanation could not do it justice and the scientific uncertainties are large.

b) How sensitive are the conclusions regarding the role of water vapor in the 100-130 mbar region to the background values of water vapor adopted, both in that region and below? The paper should discuss sensitivities to both the assumed water vapor background values and to cloudiness. If either or both are important, then what does that imply for the robustness of the results presented?

The results are not sensitive to the background concentrations of water vapour within the ranges of values in the SWOOSH dataset (page 16 Line 1-3). The main sensitivity is to background concentrations of ozone which we have now discussed in more detail with supporting calculations added in Appendix C.

2) The paper is quite long. It would greatly benefit from shortening with a particular eye to focusing on what is new here, and how robust it is. Results that are not robust to uncertainties should be edited, including where they are used to highlight differences relative to other studies. For example, the heavy emphasis on differences in ozone responses compared to Fueglistaler et al. on page 9 isn't warranted given the strong sensitivity to ozone climatology later stated on page 10. If the results are that sensitive, this doesn't merit a page of discussion. This occurs in several other places, and the paper would benefit from tightening throughout.

The paper has been considerably shortened. The main text (excluding references and appendices) has been reduced from 28 pages to 24 pages during revision. In particular, the review of current literature in the introduction has been made more concise and the discussion of the differences compared to Fueglistaler et al. (2011) is now shorter.

3) In many places, the review of past literature that is not new here could usefully be shortened. To give one example among many, potential reasons advanced in prior work as to why ozone varies between hemispheres don't need to be discussed in detail. This work is not about the reasons for ozone variations. Its focus is on their radiative effects.

This point has been addressed together with the comment above whilst shortening the paper.

4) Differences in the “smoothness” between SEFDH calculations and a 2-D dynamical model are heavily emphasized, but the computed changes are mainly in a few limited regions. It is useful and very helpful that the authors show that there is essentially no difference in the tropical mean. But its also true that over much of the region from 20N- 20S, Figures 10a,b and 11 show differences of less than 20% between the SEFDH and 2-D dynamical model calculations; i.e., Figures 10 and 11 show that the SEFDH and 2-D model agree quite well in more places and times than they disagree. Highlighting local differences that occur only in limited places doesnt provide a balanced representation of the findings. Language should be changed throughout the paper to avoid over-emphasizing spatially limited changes, and a more balanced account should be provided.

We have added Figure 10(c) which shows explicitly the difference between the SEFDH and dynamical temperature responses.

The validity of assuming fixed dynamical heating within the tropics has been questioned by a number of important studies on this topic. Fels et al. (1980): “Supporting auxiliary calculations using purely radiative models are also presented. One of these, in which the thermal sensitivity is computed using the assumption that heating by dynamical processes is unaffected by changed composition, gives results that are generally in excellent agreement with the GCM. Exceptions to this occur in the ozone reduction experiment at the tropical tropopause...” – i.e., Fels et al. (1980) are saying that (SE)FDH may give a poor estimate of the temperature response in the tropical tropopause region to changes in ozone in this region. Garcia (1987): “This implies that tropical heating will tend to be balanced by a mean meridional circulation rather than by radiative relaxation, in agreement with the conclusions of Fels et al. (1980).” If this statement by Garcia (1987) was taken as a rule of thumb then it would imply that a zeroth-order approach to considering the effect of changing ozone in the tropical tropopause region should be to assume that all the anomalous heating is balanced by a change in dynamical heating – i.e. a kind of anti-SEFDH. In practice, the extent to which the temperature response to an ozone perturbation is correctly predicted by SEFDH comes down to details, in particular the latitudinal structure in the ozone perturbation. The heating anomalies associated with broad features (more than 20 degrees of latitude for this problem, as argued in the paper) give rise to a broad temperature change as predicted by SEFDH. The heating anomalies associated with narrow features are primary balanced by changes in dynamical heating and the temperature response is small. So simply giving the size of local differences does not seem an effective description of how things work.

We believe that this characterization in terms of latitudinal scale is the most useful and general way to describe the effectiveness (or otherwise) of the SEFDH calculation, rather than focusing on whether or not numerical values are in agreement in particular locations. Nonetheless, in revision we have taken account of this comment from the reviewer by trying to ensure that our description is as clear and quantitative as possible (see also Point 6 below). As such, we have explicitly shown the differences between the SEFDH and dynamical adjustment calculations.

5) a) Are your statements about differences in the SEFDH and 2D calculations robust to uncertainties in the adopted constituent distributions? Given the strong sensitivity of the results to the background climatologies in ozone, how might errors in the SWOOSH datasets background climatologies of ozone as a function of latitude affect your results on this point? What about water vapor gradients? I would expect dynamical responses to a radiative perturbation to depend upon the background climatological gradients and was surprised that there was no discussion of that.

As explained in the paper and noted above, the differences in SEFDH and 2D dynamical calculations are largest when latitudinal gradients in the anomalous heating implied by the change in constituents are large. Therefore, broadly speaking, errors in the SWOOSH dataset on small horizontal scales will have a greater effect on the differences (and will be manifested in small horizontal scales), while errors on larger scales will have a less significant effect on the difference.

In a similar vein, the effect of errors in the background climatologies will largely be determined by how those errors modulate the anomalous heating associated with the annual variations in ozone and water vapour. If errors in the climatologies tend to reduce the small horizontal scale features in that anomalous heating, then the corresponding SEFDH vs. dynamical differences will be smaller. If background errors tend to increase the small horizontal scale features, then the differences will be larger. There does not seem to be any particular reason why errors in the background field should systematically reduce or increase the small horizontal scale features in the anomalous heating. Furthermore, irrespective of the effect of such errors, the same broad principle applies – large horizontal scale features in SEFDH calculated temperature response are likely to be robust, small horizontal scale features are not.

6) The paper does not present a clear case for what causes the decrease in 'smoothness' for the SEFDH versus IGCM, which is the central key point in the paper. The comparison of changes in vertical velocity in Figure 10c and the difference between the SEFDH and IGCM calculations suggests as many mismatches as it does matches, so this on its own doesn't serve to convince the reader. It is suggested in the text on the bottom of page 22, top of page 23 that there is a balance involving Q_{rad} , the time rate of change of temperature, and dynamical heating, but the paper doesn't demonstrate a balance. To be publishable, the paper needs to show exactly how these (or other) factors change in the model to produce the results shown. It would be appropriate to do that for a few of the key places where there are larger local differences between the SEFDH and the IGCM, and a few of the places where there are no such large changes.

In addressing this important comment (some aspects of which are addressed by our replies above to 4 and 5), we have first made some minor modifications to the implementation of the radiative code in the IGCM calculation to ensure that the SEFDH and IGCM calculations are directly comparable. These changes have only a minor effect in the updated Figure 10 (see below), but this verifies that differences between the SEFDH and dynamical adjustment calculations do not arise simply from differences in implementation.

The following text has been added:




Figure 10: (a) Monthly temperature changes showing the annual cycle at 70 hPa calculated using the idealised dynamical model (IGCM) with an annual cycle in ozone. (b) Figure 3(b) is reproduced here for comparison and shows the corresponding SEFDH calculation at 70 hPa. (c) Difference in temperature change (K) between SEFDH calculation, (b), and the IGCM calculation, (a). (d) Change in upwelling in idealised dynamical model. (e) Temperature change at 70 hPa calculated by imposing the term $\Delta(\bar{w}\bar{S})$ from the dynamical model as a perturbation to the SEFDH calculation. See main text for more details.

“Figures 10(a) and (b) compare the temperature change at 70 hPa caused by the annual cycle in ozone in the dynamical model and in the SEFDH calculation, respectively (Fig. 10(b) is identical to Fig. 3(b), but is included here for ease of comparison). Fig. 10(c) shows the difference between the two. The figures show the importance of including the dynamical adjustment, which tends to broaden the temperature response in latitude in the tropical region, making it more symmetric about the Equator. Note, in particular, the effect on the off-equatorial maximum at about 10° N in the SEFDH calculation, which is no longer a distinct isolated feature in the dynamical calculation.”

We have further added Figure 10(e), which shows the temperature response induced by the dynamical heating from the change in upwelling shown in Figure 10(d). The temperature response

in Figure 10(e) closely matches the SEFDH - dynamical adjustment temperature differences in Figure 10(c), thus demonstrating quantitatively that the temperature differences are explained by the dynamical response. We then include a discussion of the criterion that determines the range of latitudes over which dynamical heating response will dominate.

7) It would be helpful to have more information here on how the 2-D dynamical model performs. 2-D models obviously have many limitations, and the references cited generally focus more on broad dynamical phenomena than on quantitative performance. Does the 2-D model generate accurate seasonal and latitudinal climatologies of temperature, winds, and circulation from apriori information? Is the model mean circulation or background temperature distribution tuned? How much confidence is there in the models ability to simulate the strength of the Brewer-Dobson circulation (critical here), and how has it been tested? More discussion of confidence in the models quantitative performance is needed, since the papers key findings rest on a robust quantitative simulation of meridional circulation perturbations from a model of reduced dimensionality.

The dynamical model is being used to calculate responses to the changes in constituents, not calculate ‘background quantities such as the time-average Brewer-Dobson circulation. There is no sense in which the dynamical model is tuned. It simply contains the basic ingredients of the 2-D dynamical equations, plus radiative relaxation as represented by a radiative calculation considering the temperature response as an anomaly relative to a specified background temperature field. We noted in the original text that tests had shown that the calculated response was insensitive to plausible changes in that background temperature field – e.g., changing it from annual average to solstice distributions.

We have made it clear that the dynamical model does not include the effects of change in eddy forcing and we have carefully discussed the limitations of that.

8) The paper doesnt state what years were used to define the background climatologies for H₂O and O₃ against which seasonal anomalies were evaluated from among the range of 1984-2015 available in SWOOSH. This is particularly concerning for ozone, where it is clear that there have been long-term trends in the tropics. Decadal variability in tropical H₂O has also been established in the scientific literature. What years were used? How much do the years chosen matter, both in terms of background climatologies and amplitudes of the responses you are interested in, both for ozone and water vapor?

This information is given on page 5 line 13. The seasonal cycle is provided by SWOOSH using data from 1984 to 2015. The interannual variability is incorporated into the uncertainty estimates for both ozone and water vapour (see Appendix B).

Detailed comments: 1, Lines 8-9. Given the high sensitivity to the adopted background ozone values, as you show in the text, it isn't clear that your results being high deserves such emphasis here.

We were intending here to put our results into the context of other literature on this topic. However, taking our results together with those of Gilford and Solomon (2017) and Chae and Sherwood (2007), all of which show a larger amplitude for the ozone annual cycle effect than Fueglistaler et al. (2011), we have removed this clause from the abstract. A discussion of the quantitative magnitudes and how these compare to other literature is still given in the main text (Pages 6 to 9).

1, Lines 12-13. Is the non-local result robust to assumptions about background water vapor and background and seasonality of cloudiness? Clarify or delete. tropical upwelling?

The non-local radiative result arises from the properties of the radiative transfer for water vapour in the TTL and is robust to background water vapour [tested and reported in the paper, page 11, line 5] and tropospheric cloudiness [tested and not explicitly reported in the paper]. Neither the effects of background cirrus nor the seasonality in cloudiness have been considered, but this is clearly stated as outside the scope of the paper.

4, Lines 10-13. Should tropical gravity waves be explicitly noted as a possible factor in tropical upwelling?

Tropical gravity waves are possibly relevant but whilst condensing the review of past literature, as recommended by the referee, this paragraph has been modified to

“Many studies have focused on the role of wave-induced forces in driving the annual cycle in temperature through their effects on upwelling in the TTL and hence on $\overline{Q}_{\text{dyn}}$ in Eq. 1, although uncertainty remains about what types of waves are the most important (Randel and Jensen, 2013, and references therein).”

6, Lines 18-19. What is meant by “a three-point Gaussian is used to account for the diurnal variation in solar zenith angle”? Please clarify what you assume regarding diurnal changes in heating rates.

The Morcrette/Zhong and Haigh radiation code incorporates a common method of calculating a diurnal average which accounts for the variation in solar zenith angle over the day. Note that how we compute these diurnal averages does not affect the results in this paper so we have revised this sentence to “Shortwave heating rates are calculated as diurnal averages”.

7, Lines 1-2. Are you sure that the temperature changes would necessarily be zonally uniform if, for example, there are zonal asymmetries in potentially thick clouds?

This is a statement about our calculation, in which all quantities are assumed to be independent of longitude, rather than a statement that ‘if TTL temperatures and constituents are zonally uniform then the temperature response will be zonally uniform’. We have amended the text to clarify this as well as to emphasise the fact that the zonal mean adjustment would smooth regional changes in temperature that would be predicted by an SEFDH calculation.

9, Lines 21-23. Why is it necessary to presume this rather than determining whether this is actually true in your calculations?

This sentence has been edited to remove the word “presumably”. This claim is essentially verified by Fig. 3; the sentence has accordingly been strengthened.

Page 10, line 13. The claim of a 10% accuracy in the SWOOSH dataset is remarkable. Is this 2-sigma? Does it apply for local values across the full range of latitudes of interest here, all the way down to 130 mbar? What is this claim based on? I would not expect 2-sigma absolute uncertainties in tropical ozone in SWOOSH to be better than 20 or 30%, at best.

A consideration of the uncertainties as far as they have been characterized for the SWOOSH dataset (see Davis et al., 2016) suggests that 10% for climatological values is reasonable at 70 hPa

although this uncertainty increases to 30-40% by 130 hPa. We emphasise to the referee that these calculations are included in the manuscript to highlight the sensitivity of the calculations to the background ozone profile, and as such the 10% values quoted are intended to represent plausible ranges of ozone values rather than a precise assessment of uncertainty, for which a more detailed explanation for the SWOOSH dataset is given in Davis et al. (2016).

We have also added an Appendix C with additional calculations to describe the effect of varying background ozone values which is the primary source of sensitivity in our results. These calculations are intended as illustrative calculations rather than a precise assessment of uncertainty.

Page 11, line 17. Why ‘perhaps’? Please provide a clear statement based on your quantitative results as to whether it is or isn’t.

The word ‘perhaps’ has been removed.

Page 13, line 8. Garbled sentence.

This sentence was too long and has been revised to:

“In each case, the latitude of the maximum response is further north than the latitude of the maximum amplitude in the water vapour mixing ratios at that level. The fact that there is no simple relation between the latitude-time structure of the SEFDH-predicted annual cycle in temperature at a given level and the latitude-time structure of the water vapour mixing ratios at that level is further evidence for important non-local contributions in the vertical from water vapour to the temperature variations.”

Page 14, line 18. A secondary

It is not clear from this comment what the reviewer’s concern here was; however the relevant text has been modified in the course of shortening the paper.

Page 20, line 2. Why ‘maximum possible relevance ... to the real atmosphere’? A full 3-D model, and a complete line-by-line radiative code would have more relevance to the real world, so this statement should be deleted.

The text has been amended.

Sections 5.2, 5.3, 5.4, and 6 require revision to deal with the above comments. I will not repeat those remarks here on a line-by-line basis but they occur in many places.

These Sections have been revised. See responses to previous comments.

Page 25, lines 2-4. Gilford and Solomons paper has been accepted in J. Climate. Your statement that your consideration of different vertical layers, and water vapor, is different from that paper is not correct. Gilford and Solomon did consider water vapor, as well as concentration perturbations in different layers. Please revise to accurately quote what Gilford and Solomon did.

We have modified our discussion of Gilford and Solomon (2017) accordingly.

Page 28, lines 15-21. 'should not be taken too seriously' is not clear and its not scientific language. Please provide a quantitative statement that is specific, and balanced across regions of agreement and disagreement and considers robustness of your results as noted above in the major comments section.

The text has been amended to improve clarity and exploit the new Fig. 10. As noted in earlier replies, the key and robust point about the difference between SEFDH temperature response and dynamical temperature response is that this is largest at small horizontal scales. As can be seen in the revised Fig. 10(c), local differences are zero for some latitudes and dates but for others, these are up to about 1.5 K, i.e., about 50% of the maximum SEFDH 'signal'. Again, simply giving the size of local differences does not seem an effective description of how things work.

Reviewer 2

Abstract: I think the abstract could be somewhat condensed, and (i) more quantitative information could be given; (ii) the text switches back and forth between amplitude, phase, ozone, water and temperature (e.g. around Lines 6-12), which is confusing.

The abstract has been revised to discuss the ozone annual cycle first then the water vapour.

P1/L7,8: "... and out of phase with the observed annual cycle of the tracer mixing ratio."

P1/L8: Suggest: "The ozone contribution calculated here is ..."

P1/L9: Suggest to replace "This difference" with "The difference"; but see general comment on Abstract above.

These suggestions have been taken into account in the revised text.

P2/L20: Suggest (the tropopause temperature is important for stratospheric water, not dehydration per se.) "Therefore the temperatures at 90hPa and 100hPa shown in Figs 1(c/d) are more directly relevant to water entering the stratosphere than the temperature at 70hPa."

This is a good point and the sentence has been changed.

P4/L26: Quantitative details for H₂O missing in previous work. Fueglistaler et al. state that their calculations "show that the temperature adjustment for the annual cycle in water vapour around 70 hPa is about an order of magnitude smaller than that for ozone." At the time of writing Fueglistaler et al., this seemed to be sufficient information. However, in the context of your ozone effect being substantially larger than theirs, it is interesting to note that their water vapor impact (around 0.2K) is also smaller than your water vapor impact (your Figure 5b gives 0.6K). I think it would be great if your paper could resolve these

quantitative differences better, such that your paper can be cited as the best quantification of the radiative effects. The Fueglistaler et al. numbers are taken from the calculations with the Edwards-Slingo code, but results with the Fu-Liou code were rather similar (see Fueglistaler et al., page 3702; last paragraph). The calculations using the Fu-Liou code gave slightly larger amplitudes, but the differences were much smaller than the differences to your calculations; at the time, we decided to go with the Edwards-Slingo results as they were obtained from a software configuration that has been used in previous studies by Piers Foster. It would be great if you could resolve the questions you discuss on page 10/11 regarding the role of the radiative transfer code versus trace input data -the ozone and water vapor files used by Fueglistaler et al. for the calculations with the Edwards-Slingo and Fu-Liou code can be made available.

We have performed a number of SEFDH calculations that try to narrow down the causes of the differences between our calculation and Fueglistaler et al. and whilst we have not been able to explain the differences, the main effect is likely to be differing ozone climatologies rather than radiation code. The temperature change was more sensitive to the background ozone field than any other changes we tested. For instance, an SEFDH calculation with the RRTM radiation code and SWOOSH ozone obtained similar results to those in this manuscript.

We suggest that future studies, as far as possible, make the actual fields of ozone and water vapour used in the radiative calculation publicly available, to make comparison more straightforward.

Finally, can you quantify the impact on the global average temperature? An important point of Fueglistaler et al. is that one should not assume that the global average temperature remains constant (as e.g. implied in Yulaeva et al. 1994); it would be interesting to see the numbers from your calculation. You do discuss the role of extratropical heating rate variations on the tropical amplitude, and it appears to me it that would be fairly straightforward to also show the annual mean cycle of global average temperatures.

The plot below shows the global averaged temperature in the IGCM model calculation and ERA-Interim. This supports the point that the global average temperature is not constant with a peak to peak amplitude of about 1.4 K in the case of the ozone and water vapour perturbation. The 20° N-S averaged for this calculation was about 2.8 K.

We have considered including a mention of the global average but having significantly reduced the length of the paper, there did not seem an appropriate location for the detailed discussion that this topic would require.




Figure 11: Globally averaged temperature at 70 hPa in the IGCM model calculations with ozone and water vapour perturbations. The ERA-Interim plot is also shown.

Bibliography

- Chae, J. H., and S. C. Sherwood, 2007: Annual temperature cycle of the tropical tropopause: A simple model study. *Journal of Geophysical Research*, **112** (D19), D19 111, doi:10.1029/2006JD007956, URL <http://doi.wiley.com/10.1029/2006JD007956>.
- Fels, S. B., J. D. Mahlman, M. D. Schwarzkopf, and R. W. Sinclair, 1980: Stratospheric Sensitivity to Perturbations in Ozone and Carbon Dioxide: Radiative and Dynamical Response. *Journal of the Atmospheric Sciences*, **37** (10), 2265–2297, doi:10.1175/1520-0469(1980)037<2265:SSTPIO>2.0.CO;2.
- Fueglistaler, S., and Q. Fu, 2006: Impact of clouds on radiative heating rates in the tropical lower stratosphere. *Journal of Geophysical Research Atmospheres*, **111** (23), 1–13, doi:10.1029/2006JD007273.
- Fueglistaler, S., P. H. Haynes, and P. M. Forster, 2011: The annual cycle in lower stratospheric temperatures revisited. *Atmospheric Chemistry and Physics*, **11** (8), 3701–3711, doi:10.5194/acp-11-3701-2011, URL <http://www.atmos-chem-phys.net/11/3701/2011/>.
- Garcia, R. R., 1987: On the Mean Meridional Circulation of the Middle Atmosphere. *Journal of Atmospheric Sciences*, **44** (24), 3599–3609, doi:10.1175/1520-0469(1987)044<3599:OTMMCO>2.0.CO;2.
- Gilford, D. M., and S. Solomon, 2017: Radiative effects of stratospheric seasonal cycles in the tropical upper troposphere and lower stratosphere. *Journal of Climate*, JCLI-D-16-0633.1, doi:10.1175/JCLI-D-16-0633.1, URL <http://journals.ametsoc.org/doi/10.1175/JCLI-D-16-0633.1>.
- Randel, W. J., and E. J. Jensen, 2013: Physical processes in the tropical tropopause layer and their roles in a changing climate. *Nature Geoscience*, **6** (3), 169–176, doi:10.1038/ngeo1733, URL <http://www.nature.com/doifinder/10.1038/ngeo1733>.

The radiative role of ozone and water vapour in the temperature annual cycle in the tropical tropopause layer

Alison Ming¹, Amanda C. Maycock², Peter Hitchcock³, and Peter Haynes¹

¹Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge, UK

²School of Earth and Environment, University of Leeds, Leeds, UK

³National Center for Atmospheric Research, Boulder, Colorado, USA

Correspondence to: Alison Ming (A.Ming@damtp.cam.ac.uk)

Abstract. The structure and amplitude of the radiative contributions of the annual cycles in ozone and water vapour to the prominent annual cycle in temperatures in the tropical tropopause layer (TTL) are considered. This is done initially through a seasonally evolving fixed dynamical heating (SEFDH) calculation. The annual cycle in ozone is found to drive significant temperature changes predominantly locally (in the vertical) and roughly in phase with the observed TTL annual cycle. In contrast, temperature changes driven by the annual cycle in water vapour are out of phase with the latter. The effects are weaker than those of ozone, but still quantitatively significant, particularly near the cold point (100 to 90 hPa) where there are significant non-local effects from variations in water vapour at lower layers of the TTL. The combined radiative heating effect of the annual cycles in ozone and water vapour maximizes above the cold point, and is one factor contributing to the vertical structure of the amplitude of the annual cycle in lower stratospheric temperatures, which has a relatively localized maximum around 70 hPa. Other important factors are identified here: radiative damping time scales, which are shown to maximize over a deep layer centred on the cold point; the vertical structure of the dynamical heating; and non-radiative processes in the upper troposphere that are inferred to impose a strong constraint on tropical temperature perturbations below 130 hPa. The latitudinal structure of the radiative contributions to the annual cycle in temperatures is found to be significantly modified when the SEFDH assumption is relaxed and the dynamical response, as represented by a zonally symmetric calculation, is taken into account. The effect of the dynamical response is to reduce the strong latitudinal gradients and inter-hemispheric asymmetry seen in the purely radiative SEFDH temperature response, while leaving the 20° N–S average response relatively unchanged. The net contribution of the ozone and water vapour annual cycles to the peak-to-peak amplitude in the annual cycle of TTL temperatures is found to be around 35% of the observed 8 K at 70 hPa, 40% of 6 K at 90 hPa and 45% of 3 K at 100 hPa. The primary sensitivity of the calculated magnitude of the temperature response is identified as the assumed annual mean ozone mixing ratio in the TTL.

1 Introduction

The tropical tropopause layer (TTL), spanning from 150 to 70 hPa or 14 to 18.5 km, is the main entry region for air into the stratosphere from the troposphere (e.g. Fueglistaler et al., 2009). The properties of this region are influenced by the presence of

a prominent annual cycle in temperatures which is clear in, for example, radiosonde measurements Reed and Vlcek (1969) and

5 GPS radio occultation measurements (Randel et al., 2003). Figure 1 shows the structure of the temperature annual cycle in a

month-by-month climatology from the ERA-Interim reanalysis dataset (Dee et al., 2011), constructed using data from 1991 to

2010. Consistent with earlier studies, the annual cycle in temperatures is coherent over the layer from 130 to 40 hPa (Fig. 1a),

with relatively cold temperatures in northern hemisphere (NH) winter, relatively warm temperatures in NH summer and early

autumn, and with weak latitudinal gradients over the tropics (20° N–S, Fig. 1(b)–(d)). The maximum peak-to-peak amplitude

10 of the annual cycle is 8 K near 70 hPa, decreasing to 6 K at 90 hPa and 3 K at 100 hPa, below which the amplitude reduces very

rapidly (Fig. 1(a)). Above about 30 hPa (not shown), temperature variations are dominated by the semi-annual oscillation.




Figure 1. Climatology of the temperature annual cycle in ERA-Interim constructed by averaging from 1991 to 2010 and (a) between 20° N–S, (b) at 70 hPa, (c) at 90 hPa and (d) at 100 hPa. The tick marks for the months indicate the first day of each month. The values of the temperature are shown by solid contours with contour values labelled explicitly. The coloured contours are shown at intervals equal to half of the solid contour interval. These conventions are followed in all of the figures in the paper.

The temperature variations at the tropical cold point near 100 to 90 hPa regulate the water vapour entering the stratosphere on annual and inter-annual time scales by modulating the freeze drying of upwelling air (e.g., Fueglistaler and Haynes, 2005; Fueglistaler et al., 2005; Randel and Jensen, 2013). In particular, the regular annual cycle in temperatures shown in Fig. 1(c)

at 90hPa, and (d) at 100hPa leads to a significant annual cycle in water vapour. The additional effect of upward transport in the lower stratospheric Brewer-Dobson circulation creates the well known water vapour tape recorder signal (e.g., Mote et al., 1996; Randel et al., 2001). Note that the larger temperature annual cycle at 70hPa (Fig. 1(b)) does not have a direct effect on water vapour because overall temperatures are higher than at the cold point below.

Despite the potential significance of the annual cycle in TTL temperatures, both in its role in determining stratospheric water vapour mixing ratios and also simply as a conspicuous and persistent aspect of temperature variation, the mechanisms responsible for the cycle are not yet completely clear. Furthermore, state-of-the-art climate models, e.g., within the CMIP5 dataset, still exhibit large inter-model differences in the amplitude of the annual cycle at 100hPa with peak-to-peak amplitudes ranging from ~ 1 K to ~ 5 K compared to ~ 4 K for the 15° N–S average in ERA-Interim (Kim et al., 2013). This indicates that many current climate models do not capture correctly the processes that drive the TTL temperature annual cycle. Moreover the lack of understanding of the quantitative impact of different physical mechanisms on TTL temperatures precludes development of a set of general principles for improving models (Hardiman et al., 2015).

To consider the annual cycle further, it is useful to begin by introducing the thermodynamic equation in the Transformed Eulerian Mean framework (Andrews et al., 1987), neglecting eddy terms,

$$\partial_t \bar{T} = \bar{Q}_{\text{rad}} - \bar{w}^* \bar{S} - \bar{v}^* \partial_y \bar{T} = \bar{Q}_{\text{rad}} + \bar{Q}_{\text{dyn}}, \quad (1)$$

which predicts the rate of change of zonal mean temperature, \bar{T} , with time, t where $\bar{(\cdot)}$ represents a zonal mean. The dynamical heating, \bar{Q}_{dyn} , is defined by the second equality in Eq. (1). \bar{v}^* and \bar{w}^* are the horizontal and vertical components of the mean residual velocity respectively. y is the meridional coordinate. $\bar{S} = \partial \bar{T} / \partial z + \kappa \bar{T} / H$ is a measure of the static stability where z is the log-pressure height. $z = -H \log(\sigma)$ where H is a scale height taken to be 7 km and $\sigma = p/p_0$ where p is pressure and $p_0 = 1000$ hPa. $\kappa = R/c_p \simeq 2/7$ where R is the gas constant for dry air and c_p is the specific heat at constant pressure. The radiative heating, \bar{Q}_{rad} , depends in general on the temperature and the distributions of various radiatively active components including clouds, aerosols and trace gases.

Combined with the equations for zonal wind, continuity, and thermal wind balance, Eq. (1) determines the zonally symmetric response to imposed heatings and mechanical forcing (Andrews et al., 1987, Chapter 3). This is particularly important in considering the response to an imposed forcing on the right-hand side of Eq. (1), or indeed an imposed forcing term in one of the other equations.

Many studies have focused on the role of wave-induced forces in driving the annual cycle in temperature through their effects on upwelling in the TTL and hence on \bar{Q}_{dyn} in Eq. (1), although uncertainty remains about what types of waves are the most important (Randel and Jensen, 2013, and references therein). However, radiative contributions to the annual cycle have also been suggested, principally in connection with the strong annual cycle in TTL ozone mixing ratios (Folkins et al., 2006; Randel et al., 2007). The quantitative effect of ozone on TTL temperatures was first investigated by Chae and Sherwood (2007) who used a one-dimensional radiative convective model representing a tropical average profile. They concluded that at 70 hPa, about 3 K of the observed 8 K peak-to-peak variation in temperature might be caused by the radiative effects of the

annual cycle in ozone, reducing to about 1 K of the observed 3 K peak-to-peak variation at 100 hPa. Fueglistaler et al. (2011) used a seasonally evolving fixed dynamical heating (SEFDH; see Sect. 2) calculation and found a slightly smaller contribution from ozone of 2 K at 70 hPa.

Both Chae and Sherwood (2007) and Fueglistaler et al. (2011) asserted that annual variations in TTL water vapour mixing ratios have only a small role in determining the annual cycle in temperatures; quantitative details, however, were omitted. There has been recent significant interest in the radiative effect of variations in stratospheric water vapour, both the effect on the radiative balance of the troposphere (e.g., Forster and Shine, 2002; Solomon et al., 2010), but also the effect on the lower stratosphere. For example, Maycock et al. (2011), used a set of radiative calculations to show that a uniform increase in stratospheric water vapour gives rise to a cooling that is largest in the lower stratosphere at all latitudes.

In this work, we investigate, first using the SEFDH approach, the individual and combined radiative effects of the annual cycles in ozone and water vapour on TTL temperatures, including at 70 hPa where the amplitude of the annual cycle is at a maximum, and at 90 hPa near the cold point which is crucial for determining stratospheric water vapour mixing ratios. The radiative calculations required for this investigation also allow us to examine carefully how vertical structure in the background radiative environment combines with the variations in radiative and dynamical heating to determine the vertical structure of the annual cycle in temperatures. All the calculations are done using clear-sky assumptions since a full analysis of the cloud effects is beyond the scope of this work.

For a more complete assessment of the effect of seasonal variations of ozone and water vapour on the annual cycle in TTL temperatures, which goes beyond the simplifying assumptions of the SEFDH approach, it is necessary to take account of dynamical changes. The seasonal cycle in radiative heating induced by ozone and water vapour variations will in part be balanced by a change in the meridional circulation (e.g., Plumb, 1982; Garcia, 1987; Haynes et al., 1991). This is shown in Section 5 below to modify strongly the latitudinal structure of the temperature response.

The structure of the paper is as follows. Section 2 describes the data and the radiative calculations. Section 3 describes the results of SEFDH calculations to quantify the effect of annual variations in ozone and water vapour on the annual cycle of temperature in the TTL. This section includes a detailed discussion of the radiative effects of water vapour variations omitted by previous authors. These calculations are complemented by a set of illustrative fixed dynamical heating (FDH) radiative calculations in Appendix A. The SEFDH temperature changes are also sensitive to the background ozone mixing ratios and a set of further calculations is presented in Appendix C. (The results presented in Section 3 are in broad agreement with those from similar work by (Gilford and Solomon, 2017), which we became aware of during the review process.) Section 4 discusses the vertical structure of the annual cycle in temperature, distinguishing the role of the background radiative environment from that of the radiative and dynamical heating in determining this structure. Details regarding the estimates of uncertainty associated with the calculations in Sect. 3 and 4 are given in Appendix B. Section 5 then goes beyond the SEFDH calculation reported in Sect. 3 to consider how the temperature response to variations in ozone and water vapour is modified by the zonally symmetric dynamical response to the radiative heating. The final section discusses the results and their implications and reviews the various simplifying assumptions that have been made.

2 Data and Radiative method

We use temperature and dynamical fields from the ERA-Interim reanalysis dataset covering the period 1991 to 2010, using data at a horizontal resolution of 1° , at 6-hourly analysis time intervals (0000, 0006, 0012 and 0018 UTC) and on 60 model levels. The mean residual vertical velocity in the Transformed Eulerian Mean framework, calculated using the same method as Seviour et al. (2012), and the dynamical heating used in Sect. 4 are both computed on the original grid from the ERA-Interim data and then smoothed by linearly interpolating the monthly averages to daily values. The temperatures are also linearly interpolated to the grids relevant for the calculations described below and in Sect. 5.

Ozone and water vapour mixing ratios are obtained from the Stratospheric Water and OzOne Satellite Homogenized dataset (SWOOSH; Tummon et al., 2015; Davis et al., 2016). This record is formed from a combination of measurements from various limb and solar occultation satellite instruments from 1984 to 2015, namely: SAGE-II/III, UARS HALOE Harries et al. (1996); Bruhl et al. (1996), UARS MLS, and Aura MLS Lambert et al. (2007) instruments. The measurements are homogenized by applying corrections that are calculated from data taken during time periods of instrument overlap. Using the same data, SWOOSH also provides combined monthly climatologies of ozone and water vapour which we make use of in this work. SWOOSH is chosen for this study because it provides a homogenized record useful for climate studies and has been used previously to study both stratospheric water vapour (Maycock et al., 2014) and stratospheric ozone (Harris et al., 2015) variability. The pressure at the lowest altitude level in SWOOSH is 316 hPa. The results presented in this paper are not sensitive to mixing ratios of water vapour and ozone below 316 hPa (within plausible limits) and for convenience the vertical profiles below 316 hPa were simply defined by linear interpolation between the SWOOSH values at 316 hPa and the surface values taken from ERA-Interim.

The radiative calculations were performed using a modified version of the Morcrette (1991) radiation scheme, which includes updates to the longwave absorption properties of water vapour (Zhong and Haigh, 1995). All calculations were performed on zonal mean data at 5° intervals in latitude and on 100 pressure levels (which are the same as those listed in Appendix A for the FDH calculations). Shortwave heating rates are calculated as diurnal averages and the surface albedo is taken from ERA-Interim data. Carbon dioxide is assumed to be well mixed and the volume mixing ratio is set to 360 ppmv. All calculations in this study assume clear-sky conditions (i.e., neglecting radiative effects of clouds). While cloud radiative effects may also contribute to the TTL annual cycle in temperature, investigating their effects is beyond the scope of this work. However, as we will show below, our calculation accounting for the effects of both radiative and dynamical heating is able to reproduce most of the observed annual cycle in TTL temperature, suggesting that any effects of clouds (other than those included indirectly in the dynamical heating) are small.

To study the radiative contributions of seasonal variations in ozone and water vapour to the annual cycle in TTL temperatures, we make use of the seasonally evolving fixed dynamical heating calculation (Forster et al., 1997). This method calculates the time varying temperature change due to a specified radiative perturbation (e.g., a change in a trace gas) and takes into account the specified time dependence of temperature and trace gas concentration profiles in a background state, to which the perturbation is applied.

Given time-varying background profiles (at a specified latitude) of temperatures, \bar{T}^0 , and mixing ratios of trace gases, $\bar{\chi}_{\text{O}_3}^0$ and $\bar{\chi}_{\text{H}_2\text{O}}^0$ (where $(\cdot)^0$ denotes the background state), the dynamical heating, \bar{Q}_{dyn}^0 , is first calculated by assuming the balance in Eq. (1), i.e.,

$$\partial_t \bar{T}^0 = \bar{Q}_{\text{rad}}(\bar{T}^0, \bar{\chi}_{\text{O}_3}^0, \bar{\chi}_{\text{H}_2\text{O}}^0) + \bar{Q}_{\text{dyn}}^0. \quad (2)$$

A perturbation is applied to the trace gas mixing ratios ($\Delta\bar{\chi}_{\text{O}_3}$, $\Delta\bar{\chi}_{\text{H}_2\text{O}}$) and the new time evolving equilibrium temperature state, $\bar{T}^0 + \Delta\bar{T}$, is obtained from

$$\partial_t (\bar{T}^0 + \Delta\bar{T}) = \bar{Q}_{\text{rad}}(\bar{T}^0 + \Delta\bar{T}, \bar{\chi}_{\text{O}_3}^0 + \Delta\bar{\chi}_{\text{O}_3}, \bar{\chi}_{\text{H}_2\text{O}}^0 + \Delta\bar{\chi}_{\text{H}_2\text{O}}) + \bar{Q}_{\text{dyn}}^0. \quad (3)$$

Eq. (3) is integrated forward in time with a daily time step until the perturbed temperature field, $\bar{T}^0 + \Delta\bar{T}$, is also annually repeating. Five years is found to be sufficient for accurate convergence. The radiative transfer calculation couples vertical levels, but not latitudes, so that each calculation is local in latitude. Following a similar method to Forster et al. (1997), Eq. (3) is applied to update the temperature only above a certain level taken here to be 130 hPa, on the basis that there are distinct processes determining temperature variations in the troposphere below. The choice of this level is further justified in Sect. 4. In setting up the calculation we verified that the Forster et al. (1997) results could be reproduced. The SEFDH technique reduces to the more standard and widely used FDH technique if the imposed background temperature and species mixing ratios are constant in time and Eq. (3) is integrated to a steady state (Appendix A).

The background state is taken to be the annual average ERA-Interim temperature (which implies $\partial_t \bar{T}^0 = 0$) and the annual mean SWOOSH constituent mixing ratios. The latter are then perturbed to their annually varying climatologies. One could alternatively use the annually varying temperature climatology as the base state (e.g., Fueglistaler et al., 2011), but this was found to have a negligible impact on the simulated temperature response. The use of a time independent background state was also easier to implement in the dynamical calculations reported in Sect. 5.

3 SEFDH calculations of temperature response

3.1 Temperature response due to ozone annual cycle

Figure 2 shows differences in ozone mixing ratios from the annual mean over the tropics. The annual cycle in tropical lower stratospheric ozone mixing ratios, and in particular the large amplitude of the annual cycle relative to annual mean values, is well known on the basis of ozonesondes (e.g., Logan, 1999) and satellite data (e.g., Randel et al., 2007; Tegtmeier et al., 2013). The height and latitude structure of the ozone annual cycle at low latitudes from SWOOSH is shown in Fig. 2(a), and the corresponding latitudinal structure at 70 hPa is shown in Fig. 2(b). Ozone mixing ratios are lowest across the tropics in NH winter/spring and highest in NH summer/autumn. The cycle has a broad latitudinal structure, but the amplitude is significantly larger in the NH subtropics than in the SH subtropics (Stolarski et al., 2014). Whilst the amplitude of the annual cycle in ozone mixing ratio increases with height (Fig. 2(a)), the amplitude as a proportion of the annual mean mixing ratio (i.e., the ‘relative amplitude’) is largest at about 80 hPa and decreases upward above that level (e.g., Randel et al., 2007, their Fig. 3).




Figure 2. Ozone volume mixing ratio (ppmv) from the SWOOSH dataset plotted as a difference from the annual mean (a) averaged between 20° N-S and (b) at 70 hPa. (c) The total change in heating rate (longwave and shortwave) due to the ozone annual cycle assuming temperatures are fixed at the annual average values.

The factors that determine the temperature response to a change in ozone mixing ratio in the TTL are explained in detail in Appendix A1. The main effect of a reduction in ozone in a particular shallow layer is to decrease heating in that layer, through both decreased shortwave absorption and decreased absorption of upwelling longwave radiation, with the latter being the dominant effect in the TTL. The decreased opacity of the perturbed layer also leads to increased longwave heating in overlying layers. As an illustration, Fig. 2(c) shows the change in heating rate at 70 hPa due to the ozone annual cycle assuming temperatures are fixed at the annual average values. (The quantity plotted is $\overline{Q}_{\text{rad}}(\overline{T}^0, \overline{\chi}_{O_3}^0 + \Delta \overline{\chi}_{O_3}, \overline{\chi}_{H_2O}^0) - \overline{Q}_{\text{rad}}(\overline{T}^0, \overline{\chi}_{O_3}^0, \overline{\chi}_{H_2O}^0)$.)

The temperature response associated with the ozone anomalies in Fig. 2 predicted by the SEFDH calculation are shown in Fig. 3. A significant annual cycle in temperature is simulated across the tropics (averaged between 20° N-S), with cooler temperatures when ozone mixing ratios are relatively low, in NH winter, and warmer temperatures when ozone mixing ratios are relatively high, in NH summer.

In the vertical, the temperature response to ozone is largest between 90 to 70 hPa with a peak-to-peak amplitude over the annual cycle of about 3.5 ± 0.4 K at 70 hPa and about 3.3 ± 0.5 K at 90 hPa (Fig. 3(a); values are quoted with 95% confidence intervals, see Appendix B for details). The simulated temperature response has a lag of about 1.5 months compared to the annual cycle in ozone. The response essentially has the same sign at all levels, because the change in ozone mixing ratios occurs over a relatively deep layer so that, at a given level, any effects of the reduction in upwelling radiation by increased ozone in the levels below are dominated by the increased absorption by ozone at that level. The latitudinal structure of the simulated temperature response at 70 hPa is shown in Fig. 3(b). Within the tropics, the latitudinal structure closely matches that of the ozone variations shown in Fig. 2(b). Both are stronger in the NH subtropics than in the SH subtropics.

The temporal and latitudinal structure of the temperature response to ozone at 70 hPa are similar to those presented by Fueglistaler et al. (2011) who used ozone from the HALOE dataset and the Edwards and Slingo radiation code. However, the peak-to-peak amplitude we obtain, 3.5 ± 0.4 K, is significantly larger than they report (~ 2 K). The 3 K amplitude found by




Figure 3. SEFDH temperature change (K) due to the annual cycle in ozone. All other trace gases are kept at their annual mean values. (a) Temperature change averaged between 20° N-S. (b) Temperature change at 70 hPa. (c) Temperature changes at 70 hPa and averaged between 20° N-S calculated with the annual cycle in ozone imposed within different pressure ranges. Outside of each range and including the pressure level at the lower bound (in terms of height) of the range, the ozone mixing ratio is kept at the annual mean value.

Chae and Sherwood (2007) is closer to our result and provides useful comparison since the assumptions underlying the time-dependent 1D radiative-convective calculation from which it was obtained are very similar to those in the SEFDH approach. The recent study of Gilford and Solomon (2017), which used the same SEFDH approach with a different radiation code and ozone climatology, finds an amplitude of 3.1 K, which is more consistent with the present results.

There are several possible causes for the quantitative differences between results, including differences in the satellite ozone datasets employed (which reflect real observational uncertainties) and differences in radiation schemes. Our quoted uncertainties for the magnitude of the annual cycle in temperature at different levels (refer previous text and/or Appendix B) are intended to estimate the effect of the observational uncertainties. While we have not been able to isolate the specific reason for the different results, it is clear from further sensitivity tests, reported in Appendix C, that the quantitative temperature response to the ozone annual cycle has significant sensitivity to the annual mean background ozone concentrations. For example, if the background ozone concentration is reduced at each level by about 10% (which corresponds to about two standard deviations of the estimated uncertainty in the annual mean ozone mixing ratio), the peak-to-peak amplitude of the annual cycle is reduced by about 5%.

To investigate the role of ozone variations in different layers of the TTL for the observed annual cycle, the SEFDH calculation was repeated with the annual cycle in ozone imposed only within a set of sub-layers: 1000 to 90 hPa, 90 to 50 hPa, 50 to 30 hPa, and 30 to 1 hPa (see Fig. 3(c)). Outside the given layer, and including the pressure level at the lower bound of the range (in height), ozone is left at the annual mean value. The annual cycle in ozone in the region 90 to 50 hPa accounts for about 80% of the temperature annual cycle at 70 hPa. A similar result is found at 90 hPa, where about 60% of the temperature variation is driven by ozone variations in the 100 to 80 hPa layer and about 30% by those in the 80 to 50 hPa layer (not shown). The relation between ozone variations and the resulting temperature variations in the TTL region is therefore primarily local in the vertical.

To summarise the results of this subsection, we have shown using an SEFDH calculation that the ozone annual cycle can
5 account for 3.5 ± 0.4 K of the 8.2 ± 0.3 K observed peak-to-peak amplitude of the annual cycle in tropical averaged temperature
at 70 hPa. The response amounts to an even larger fraction of the observed annual cycle near the cold point, accounting for
 3.3 ± 0.5 K of the 5.8 ± 0.2 K amplitude at 90 hPa, and 2.6 ± 0.2 K of the 3.4 ± 0.1 K amplitude at 100 hPa.

3.2 Temperature response due to water vapour annual cycle

Figure 4 shows the annual cycle in tropical water vapour mixing ratio anomalies from the SWOOSH dataset. The tropical
10 average water vapour mixing ratios show a clear tape recorder signal of tilted bands of positive and negative anomalies in the
vertical (Fig. 4(a)). Above the tropopause, the amplitude of the water vapour annual cycle is largest at around 90 hPa where
temperatures are coldest, consistent with the fact that water vapour is directly controlled by temperature. The amplitude of the
annual cycle increases significantly below 150 hPa.

Since the vertical structure of the water vapour annual cycle is quite complex relative to that of ozone, we show the latitudinal
15 structure at several different levels, 70, 90 and 100 hPa (Fig. 4(b)–(d)). Some hemispheric differences are apparent, especially
at 100 hPa. The amplitude in the annual cycle in water vapour is greater in the NH, with the largest values near the cold point
in September.

The radiative factors that determine the temperature response to a change in water vapour in the TTL are described in detail
in Appendix A2. The main effect of a reduction in water vapour within a particular shallow layer is cooling below the layer
20 and heating within and above it. The reduction in water vapour implies less local emission of longwave radiation and therefore
reduced absorption above and below (hence the cooling), together with less absorption of upwelling radiation within the layer
and increased absorption above. Within the layer the effect of reduced local emission is stronger, so the net effect is heating.
Above the layer the effects of increased absorption of upwelling radiation dominate, leading to net heating.

Figure 5(a) shows the temperature response from the SEFDH calculation for the water vapour changes in Fig. 4. The
25 temperature response peaks near 90 hPa, i.e., at a lower altitude than the maximum response to ozone (see Fig. 3(a)). In
contrast to the ozone response, the water vapour response has a phase lag of about one month between 90 hPa and 70 hPa.
Note the phase lag of about 2 months between the annual cycle in water vapour mixing ratios at these levels (Fig. 4(a)). These
phase lags result because of the non-locality of the radiative response and the fact the effect of one layer on another is being
communicated in part by changes in temperature, and hence changes in radiation, in the intermediate layers. The temperature
30 response in the range 100 hPa to 70 hPa is, broadly speaking, opposite in phase to the observed annual cycle in temperatures
(Fig. 1(a)) and the response to ozone (Fig. 3(a)).

Figures 5(b)–(d) show that the water vapour response is largest in the NH subtropics at all three levels 70, 90 and 100 hPa.
In each case, the latitude of the maximum response is further north than the latitude of the maximum amplitude in the water
vapour mixing ratios at that level. The fact that there is no simple relation between the latitude-time structure of the SEFDH-
predicted annual cycle in temperature at a given level and the latitude-time structure of the water vapour mixing ratios at that
level is further evidence for important non-local contributions in the vertical from water vapour to the temperature variations.




Figure 4. Water vapour volume mass mixing ratio (ppmv) from SWOOSH plotted as a difference from the annual mean (a) averaged over the region 20° N–S, (b) at 70 hPa, (c) at 90 hPa and (d) at 100 hPa.

As in the previous section, we examine these non-local contributions further by imposing the water vapour changes only within a set of sub-layers: 1000 to 200 hPa, 200 to 130 hPa, 130 to 100 hPa, 100 to 80 hPa, 80 to 60 hPa, and 60 to 1 hPa. Typical results are illustrated by Fig. 6(a), which shows the response at 90 hPa for each calculation. The total peak-to-peak amplitude is 1.1 ± 0.1 K, which consists of a local contribution from the 80 to 100 hPa layer of 0.7 K, and a significant non-local contribution of 0.4 K from the 100 to 130 hPa layer. Contributions from above 80 hPa and from below 130 hPa are small. The net contribution from the 130 to 200 hPa layer is small in the 20° N–S average as a result of cancellations between the Northern and Southern Hemisphere temperature changes. There is also a large meridional gradient in water vapour resulting in a larger temperature change in the Northern Hemisphere which is about 15% of the temperature change at 90 hPa and 20° N (not shown).

Further illustration is given in Fig. 6(b), which shows the time evolution of the temperature response at all levels when the water vapour perturbation is confined to 130 to 100 hPa. In this layer, the water vapour anomaly is at a minimum in February–March and at a maximum in September–October. The features of the response described above are all visible, except there is no cooling below 130 hPa due to the SEFDH temperature constraint. Figure 6(c) shows the same temperature response plotted at 90 hPa. Comparing to Fig. 5(c) (note the different contour interval), the 100 to 130 hPa region contributes to about 35% of




Figure 5. Temperature change (K) due to the water vapour annual cycle in an SEFDH calculation (a) averaged between 20° N–S, (b) at 70 hPa, (c) at 90 hPa and (d) at 100 hPa.

the total response at 90 hPa. The peak response is centred at around 25° N, demonstrating that the maximum response at 90 hPa (Fig. 5(c)) is shifted northwards by non-local effects.

- 5 Gilford and Solomon (2017) find a response to water vapour changes with a peak-to-peak amplitude of 0.6 K at 70 hPa, consistent with the present results. Further sensitivity tests show that, unlike the case with ozone, the temperature response is not very sensitive to changes in the background value of water vapour (considering changes typical of interannual variations within the range of years covered by the SWOOSH dataset).

3.3 Temperature response to annual cycle in both constituents and dynamical heating

- 10 Figure 7(a), for 70 hPa, and Fig. 7(b), for 90 hPa, show the combined effects of the ozone and water vapour annual cycles on temperature in an SEFDH calculation. These figures also show the observed annual cycle in temperature, the estimated annual cycle in temperature due to the annual cycle in dynamical heating (based on ERA-Interim data, see Sect. 4 for further details) and the estimated annual cycle due to the combined effects of ozone, water vapour and dynamical heating. To a good




Figure 6. Annual cycle temperature changes (K) at 90 hPa calculated using SEFDH with the annual cycle in water vapour imposed within different pressure ranges. Outside of this range and on the pressure level at the lower bound (in terms of height) of the range, the water vapour mixing ratio is kept at the annual mean value. The plots are averaged between 20° N–S. The contributions from each layer add linearly to reproduce the total change (not shown). The temperature change (K) for the case in (a) where the water vapour annual cycle is imposed only from 100 to 130 hPa is shown in (b) averaged between 20° N–S and in (c) at 90 hPa.

approximation the combined effect of ozone and water vapour is simply the sum of the individual effects discussed in Sect. 3.1 and 3.2.

At 70 hPa (Fig. 7(a)), ozone and water vapour together can account for an annual cycle in temperature of about 2.8 ± 0.3 K peak-to-peak, i.e., about 35% of the observed annual cycle in temperature. The cancellation between the effects of ozone and water vapour on temperature is strongest at 90 hPa (Fig. 7(b)) with the combined amplitude being about 2.3 ± 0.4 K peak-to-peak, i.e., again about 40% of the observed annual cycle. At 100 hPa, the combined amplitude is about 1.5 ± 0.4 K peak-to-peak, or about 45% of the observed annual cycle (not shown). Thus, while the estimated contribution of dynamical heating to the annual cycle in temperatures is substantially smaller than the observed annual cycle (Fig. 7(a)–(c)), when the contributions from dynamical heating, ozone and water vapour are combined the result is in remarkably good agreement with the observed annual cycle, both in amplitude and in phase.

In summary, the combined effects of ozone and water vapour variations exert a substantial radiative influence on the annual cycle in TTL temperatures and the lower stratosphere above. The estimated radiative effect of ozone and water vapour and the observed annual cycle both peak in amplitude at 70 hPa. The fractional effect of ozone and water vapour relative to the annual cycle is substantial throughout the TTL, including at the cold point, where temperatures control the entry values of stratospheric water vapour.

4 Vertical structure of the temperature annual cycle

The annual cycle in tropical lower stratospheric temperature is largest over a shallow layer from 100 to 50 hPa, with a maximum amplitude at 70 hPa (Fig. 1(c)). This vertical structure has been attributed by Randel et al. (2002) to the presence of long




Figure 7. Temperature changes (K) calculated using SEFDH method with annual cycles in ozone and water vapour and from the modified SEFDH method (Sect. 4) for the dynamical heating. The plots are averaged between 20° N–S at (a) 70 hPa and (b) 90 hPa. The ERA-Interim temperature annual cycle is also shown. Note that the vertical axes are different in (a) and (b). (c) The peak-to-peak amplitude of the temperature change averaged between 20° N–S. Note that there is a phase difference between the temperature from the water vapour and ozone annual cycles. Shadings show 95 % confidence intervals arising from uncertainties in the datasets (see Appendix B for more details).

radiative time scales in this region. In this section, we reconsider the question of whether the location of the maximum variation in tropical temperatures over the annual cycle is due to the structure of the major radiative and dynamical forcings, and/or to the structure of the background radiative environment.

Figure 8(a) shows the variation of tropical averaged $\overline{w^*}$, in height and time, and reveals a systematic decrease in amplitude with increasing height from 150 to 50 hPa. Figure 8(b) shows the full dynamical heating term, $\overline{w^*S}$, and Fig. 8(c) shows the same quantity with the annual mean, $\langle \overline{w^*S} \rangle$, removed. The annual cycle in dynamical heating is larger above 100 hPa compared to below and rather uniform in amplitude over a deep layer that extends from 90 hPa up to about 40 hPa (Fig. 8(c)). This behaviour is strongly influenced by the annual cycle in upwelling (Fig. 8(d)), and in the region 120 to 90 hPa, is further modified by the annual cycle in static stability (Fig. 8(e)), which causes a reduction in the annual cycle in dynamical heating around 100 hPa. The peak-to-peak amplitude of the annual cycle in dynamical heating in the lower stratosphere is around 0.15 to 0.2 K day⁻¹. This amplitude does not decrease below the tropopause as rapidly as does the observed peak-to-peak amplitude of temperature (Fig. 1(a)).

To probe the effects of these heating structures on the temperature annual cycle, we now consider a set of SEFDH-like calculations forced by a set of specified dynamical heating structures. This is achieved by imposing an additional dynamical heating, $\Delta \overline{Q}_{\text{dyn}}$, on the right hand side of Eq. (1). To determine whether the localization of temperature variation is a result of the structure of the radiative environment, we first consider an idealized dynamical heating perturbation with no vertical structure, $\Delta \overline{Q}_{\text{dyn}} = -0.1 \cos(2\pi t/365) \text{ K day}^{-1}$. For this and the next few calculations, we remove the constraint on temperatures below 130 hPa, permitting them to evolve freely in response to the radiative perturbations. Any vertical dependence in the response to this heating will therefore be solely determined by vertical structure in the temperature-dependent part of the radiative heating.




Figure 8. Data from ERA-Interim averaged between 1991 to 2010 and 20° N-S. Monthly averages are interpolated to daily values to smooth out the noise in the upwelling field. (a) Mean residual vertical velocity, \bar{w}^* . (b) Dynamical heating term $\bar{w}^* \bar{S} (= -\bar{Q}_{\text{dyn}})$. (c) Same as (b) but with the annual mean removed. (d) $(\bar{w}^* - \langle \bar{w}^* \rangle) \langle \bar{S} \rangle$ component of the dynamical heating. (e) $\langle \bar{w}^* \rangle (\bar{S} - \langle \bar{S} \rangle)$ component of the dynamical heating.

5 The resulting temperature change for this case is shown in Fig. 9(a). The amplitude of the response is largest in a layer centred on 100hPa. The phase lag with respect to the imposed heating is also largest in this layer and equal to about 60 days. This is consistent with a Newtonian cooling model in which the radiative relaxation time scale was a maximum of about 60 days at 100hPa, roughly at the cold point, consistent with theoretical expectations (Bresser et al., 1995; Fels, 1982). The implied radiative time scales peak over a broader height range than those found by Randel et al. (2002) and in particular do not

10 show such a strong reduction below 100hPa. One reason for the different results likely arises from the fact that Randel et al. (2002) inferred time scales by looking at the cross-correlation between the annual components of \bar{T} and \bar{w}^* and that includes the effects of the extra non-radiative physics operating in the tropical upper troposphere. We note, however, that radiative time scales in this region depend strongly on the vertical scale of the imposed temperature perturbation (Fels, 1982).

The response to the annual cycle in $\bar{w}^* \bar{S}$ from ERA-Interim (Fig. 8(c)) is now considered. For convenience, we set the dynamical heating below 450hPa to have the same value as at 450hPa. This does not affect the main conclusions of this

5 calculation. The corresponding temperature response is shown in Fig. 9(b). The vertical structure in the dynamical heating

significantly modifies the vertical structure in the temperature response. In particular, the fact that the dynamical heating is larger at 70hPa than at 100hPa leads to a larger temperature response at 70hPa than at 100hPa, in contrast to the response to the uniform dynamical heating shown in Fig. 9(a). Therefore the vertical structure in amplitude of the temperature annual cycle driven by dynamical heating is determined by both the background radiative environment and by the vertical structure of the dynamical heating itself.

To further illustrate this, Fig. 9(c) shows the temperature response to the ERA-Interim dynamical heating assuming a constant radiative relaxation time scale of 40 days. The response is a good approximation to that in Fig. 9(b) around 70hPa, suggesting that the radiative time scale appropriate for the dynamical heating perturbation is around 40 days, somewhat shorter than that inferred from Fig. 9(a) and consistent with the smaller vertical length scale of the imposed perturbation. However, there remains below 100hPa a peak-to-peak amplitude in the temperature that is significantly larger than is observed. If we assume that this calculation of the dynamical heating provides a reasonable estimate of the magnitude of the dominant terms in the thermodynamic budget of the upper troposphere, this suggests that upper tropospheric processes provide a stronger constraint on temperature perturbations than do clear-sky radiative processes. Further calculations (results not shown) suggest (subject to the preceding assumption) that the effective time scale of this constraint is approximately 10 days.

To illustrate the implications of the observed tropospheric constraint on temperatures, we re-introduce the clamp on the temperatures below 130hPa in the SEFDH calculation as a simple representation of these processes. Figure 9(e) shows the resulting temperature response to the same dynamical heating perturbation imposed in Fig. 9(b). The tropospheric constraint causes the maximum amplitude of the response to shift upwards from around 80hPa to about 70hPa and reduces the magnitude of the peak response compared to Fig. 9(b). It is clear, therefore, that this upper tropospheric constraint has a significant radiative effect on the region above.

We now add the radiative heating perturbations from the ozone and water vapour annual cycles to the ERA-Interim dynamical heating to produce Fig. 9(f). The net effect of the ozone and water vapour annual cycles, as shown in Sect. 3.3, is to increase the amplitude of the temperature response. This produces an annual cycle with a structure that is in better agreement than that in Fig. 9(e) with the ERA-Interim annual cycle, Fig. 9(d), with a more pronounced peak at 70hPa.

In summary, the calculations reported in this section suggest that the vertical structure of the peak-to-peak amplitude in the annual cycle of temperatures arises from a combination of several effects. In the absence of the implied upper tropospheric constraint, we find that clear-sky radiative processes produce long radiative time scales over a deep layer centred around 100hPa and would, in the absence of other effects, imply a similarly deep structure in the amplitude of the annual cycle. The vertical structure in the dynamical heating and radiative heating from constituent changes, both of which exhibit a peak in the region around 80 to 70hPa, combined with the tropospheric constraint, lead to a shallower vertical structure with a stronger response at 70hPa than at 100hPa.




Figure 9. Temperature change from the annual mean averaged between 20°N – S . (a) An SEFDH like calculation with a perturbation of $-0.1 \cos(2\pi t/365) \text{ K day}^{-1}$ is added to the dynamical heating with annual mean ozone and water vapour. (b) The temperature change from the annual cycle in the ERA-Interim dynamical heating, $\overline{w^*S} - \langle \overline{w^*S} \rangle$, shown in Figure 8(c). (c) Temperature change due to the annual cycle in the ERA-Interim dynamical but assuming a constant radiative relaxation rate of $1/40 \text{ days}^{-1}$. (d) ERA-Interim annual mean temperature averaged between 1991 to 2010 (same as Figure 1(a)). (e) Similar to (b) but with temperature held fixed at the annual mean below 130 hPa. (f) Similar to (e) but with the additional perturbation from the annual cycle in ozone and water vapour included.

5 The effect of zonally symmetric dynamical adjustment

We will now consider the temperature response to annual cycles in ozone and water vapour, relaxing the SEFDH assumption to include zonally symmetric dynamical adjustment. This approach assumes no change in the zonally averaged wave force, which might well be a significant part of the full dynamical response in a three-dimensional atmosphere, even if the imposed annual cycles in ozone and water vapour are zonally symmetric. We discuss the implications of this simplifying assumption in Sect. 6 below. The zonally symmetric dynamical response problem has been considered in many previous papers (e.g., Plumb, 1982; Garcia, 1987; Haynes et al., 1991). The expectation from this previous work is that the response to the heating implied by an imposed change in constituents will occur in part through dynamical heating, modifying the vertical and latitudinal structure of the temperature response. One important difference from these previous studies is that, rather than approximating the temperature-dependent part of the radiative heating by Newtonian cooling, we continue to use the modified Morcrette/Zhong and Haigh radiation code.

5.1 Model description

For the dynamical calculations, we use the University of Reading IGCM 3.1 (de F. Forster et al., 2000) which is a hydrostatic primitive equation model based on the original Hoskins and Simmons (1975) spectral dynamical model. Only the coefficients of the zonally symmetric spherical harmonics are retained, up to the total wavenumber 42, resulting in an approximate latitudinal resolution of 3° . There are 60 levels equally spaced in log-pressure coordinates in the vertical with the model top at 50 km. The velocities in the layer near the surface $\sigma > 0.7$ are linearly damped as described in Held and Suarez (1994).

The temperature tendency in the model is given by:

$$\partial_t \bar{T} = [\dots] + (1 - G(\phi, \sigma))(\bar{Q}_{\text{rad}}(\bar{T}(t), \bar{\chi}(t)) - \bar{Q}_{\text{rad}}(\bar{T}^0, \bar{\chi}^0)) - G(\phi, \sigma) \alpha (\bar{T} - \bar{T}^0) \quad (4)$$

where the $[\dots]$ represents other advective processes in the model and $G(\phi, \sigma) = 0.5(1 + \tanh(50(\sigma - \sigma_{\text{trop}}(\phi))))$. The notation $\bar{Q}_{\text{rad}}(\bar{T}(t), \bar{\chi}(t))$ is used for brevity, with the second argument indicating schematically the mixing ratio of any radiatively active species (i.e., both ozone and water vapour in this case). Radiative transfer is imposed above 130 hPa by setting $\sigma_{\text{trop}}(\phi) = 0.13$.

This leads to a smooth transition between the troposphere, which is relaxed to ERA-Interim annual mean temperatures on a time scale $1/\alpha = 10$ days, and the stratosphere where the radiative heating rates are determined explicitly by the radiation code. The radiative calculation is implemented in exactly the same way as in the SEFDH calculations in Sect. 3 and 4. We have verified that the standalone radiation code and the version in the model produce consistent longwave and shortwave heating rates.

The additional term $\bar{Q}_{\text{rad}}(\bar{T}^0, \bar{\chi}^0)$ in Eq. (4) is imposed in the model such that, with the annual mean ozone and water vapour fields, $\bar{\chi}^0$, the temperature, \bar{T}^0 , remains close to the ERA-Interim annual mean. The control run evolves away from this state through the effect of dissipative dynamical processes such as surface drag and model hyperdiffusion, but differences from the ERA-Interim annual mean are small (less than 2 K in the tropical stratosphere) and do not affect the results presented below.

Perturbation experiments are carried out by imposing an annual cycle in ozone and/or water vapour. Mixing ratios are read in daily and heating rates are linearly interpolated to the model time step. Both the control and perturbed runs are five years

5 long, of which four years are the spin-up period. The results shown in the remainder of Sect. 5 are differences between the perturbed and control runs from the last year of the run. Results in this section are shown for the nearest model levels to 70, 90 and 100 hPa, which are 68.8, 87.3 and 98.3 hPa respectively.

5.2 Temperature response to ozone annual cycle




Figure 10. (a) Monthly temperature changes showing the annual cycle at 70 hPa calculated using the idealised dynamical model (IGCM) with an annual cycle in ozone. (b) Figure 2(b) is reproduced here for comparison and shows the corresponding SEFDH calculation at 70 hPa. (c) Difference in temperature change (K) between SEFDH calculation, (b), and the IGCM calculation, (a). (d) Change in upwelling in idealised dynamical model. (e) Temperature change at 70 hPa calculated by imposing the term $\Delta(\bar{w}\bar{S})$ from the dynamical model as a perturbation to the SEFDH calculation. See main text for more details.

Figures 10(a) and (b) compare the temperature change at 70 hPa caused by the annual cycle in ozone in the dynamical model and in the SEFDH calculation, respectively (Fig. 10(b) is identical to Fig. 3(b), but is included here for ease of comparison). Fig. 10(c) shows the difference between the two. The figures show the importance of including the dynamical adjustment, which tends to broaden the temperature response in latitude in the tropical region, making it more symmetric about the Equator. Note, in particular, the effect on the off-equatorial maximum at about 10° N in the SEFDH calculation, which is no longer a distinct isolated feature in the dynamical calculation.

This difference between the dynamical and SEFDH calculations is as expected from the previously cited theoretical work on the zonally symmetric dynamical response adjustment problem. In the dynamical calculation there is a change in vertical velocity, \overline{w}^* , and in consequence, the applied heating is balanced in part by $\partial_t \overline{T}$ and $\overline{Q}_{\text{rad}}$ (the ‘temperature part’ of the response), and in part by a response in dynamical heating (principally $\overline{w}^* \overline{S}$). In considering the annual cycle in the TTL, time scales are comparable to or somewhat larger than the radiative damping time, implying that the change in the temperature-dependent part of $\overline{Q}_{\text{rad}}$ is significant (but not necessarily dominant) in the ‘temperature part’ of the response. On the basis of simple scaling arguments which follow as a corollary to those presented, for example in Garcia (1987) or Haynes (2005), the dynamical heating response is then expected to dominate over the ‘temperature part’ of the response when the latitudinal scale, L , is less than $(ND/(2\Omega \sin \phi))(\omega_a/\alpha)^{1/2}$ where N is the buoyancy frequency, D is the vertical scale of the heating, Ω is the rotation rate and ω_a is the annual frequency. This condition holds when the latitudinal scale L is sufficiently small or at sufficiently low latitudes. Close to the Equator, this criterion is modified to L being less than $(NDa/(2\Omega))^{1/2}(\omega_a/\alpha)^{1/4}$, where a the radius of the Earth. (Note that this condition can be re-written in terms of $\beta = 2\Omega/a$, the gradient of the Coriolis parameter at the Equator.) Since the ratio of ω_a/α is close to 1 (recall that in Sect. 3, the relevant value of the radiative relaxation time was deduced to be about 40 days), it follows from the latter expression, assuming a vertical scale, D , of 4 km, that the dynamical heating response will dominate on latitudinal scales of less than about 2000 km or 20° .

The \overline{w}^* response at 70 hPa to the ozone annual cycle variations is shown in Fig. 10(d). Consistent with the dynamical scaling argument, the \overline{w}^* field tends to emphasise the smaller latitudinal-scale features in the heating field shown in Fig. 2(c), e.g., the two regions of strong cooling near 30° S and 10° N in January and February, and the regions of strong heating at about 20° S in September and October and at about 10° N in August and September. On the other hand, between these regions there tends to be an opposite signed dynamical response. We have verified consistency by applying the dynamical heating corresponding to the vertical velocity field shown in Fig 10(d), extracted from the IGCM calculation, as a perturbation heating in an SEFDH calculation using the same procedure described in Sect. 3.3 and 4. The resulting temperature response shown in Fig. 10(e) is a very good match to the difference in temperature in Fig. 10(c). The temperature difference at 70 hPa between the SEFDH and IGCM calculations (Fig. 10(c)) is, therefore, that forced by a heating anomaly equal to the dynamical heating response. The overall effect of the dynamical adjustment is to smooth the SEFDH predicted temperature response in latitude, eliminating features of latitudinal scale (in this particular problem) less than about 20° . The amplitude of \overline{w}^* between 20° N–S is typically about 20% of the amplitude of the annual cycle in upwelling in ERA-Interim, implying that the ozone heating plays a non-negligible role in determining the latitudinal structure of the overall annual cycle in \overline{w}^* .

5.3 Temperature response to water vapour annual cycle

The temperature response of the dynamical model to a perturbation from annual average water vapour to annually varying water vapour is now considered in a similar way to the ozone perturbation just discussed. Given the significant radiative interactions in the water vapour response between different vertical layers, the temperature responses at each of the levels 70 hPa, 90 hPa and 100 hPa are displayed respectively in Fig. 11(a)–(c). The corresponding SEFDH temperature responses at 70 hPa, 90 hPa and 100 hPa are shown respectively in Fig. 11(d)–(f). As was the case for ozone, the temperature responses in the dynamical

model are broader and smoother than the corresponding SEFDH temperature responses. The prominent maxima in heating in March and April at 20° N at 70 hPa and about 25° N at 90 hPa and 100 hPa, and in cooling in September to November at the same locations, are reduced in magnitude, but over the Equator and extending into the SH there is increased heating in March and April and increased cooling in September to November. The resulting structure in the tropics is much more symmetric across the Equator than the SEFDH temperature response.




Figure 11. Temperature changes (K) calculated using the idealised dynamical model (IGCM) with an annual cycle in water vapour at (a) 70 hPa, (b) 90 hPa and (c) 100 hPa. Temperature change from the SEFDH calculation (same as Fig. 5(b)–(d)) at (d) 70 hPa, (e) 90 hPa and (f) 100 hPa for comparison.

5.4 Temperature response to ozone and water vapour annual cycles

5 The combined effect of the ozone and water vapour annual cycles in the dynamical calculation is now considered. Their effects, to very good approximation, add linearly. The latitudinal structure of the combined response is shown for 70 hPa (Fig. 12(a)) and for 90 hPa (Fig. 12(b)). Figures 12(c) and (d) show the temperature responses in the dynamical model averaged between 20° N–S, to ozone and water vapour individually and their combined response, at 70 hPa and at 90 hPa respectively. Also shown on these figures are the SEFDH results for comparison (same as Fig. 7(a) and (b)). By this tropical average measure,




Figure 12. Temperature changes (K) calculated using the idealised dynamical model (IGCM) with annual cycles in both ozone and water vapour shown at (a) 70hPa and at (b) 90hPa. Temperature changes averaged between 20° N–S at (c) 70hPa and (d) 90hPa and showing the effects of ozone and water vapour in the dynamical model (thick lines) as well as the corresponding SEFDH temperature changes from Fig. 7(a) and (b) (thin lines).

there is virtually no change in the peak-to-peak amplitudes of the individual and combined temperature responses to ozone and water vapour variations between the SEFDH and dynamical calculations. Any reduction in local latitudinal maxima in the temperatures response is offset by the broadening effect, leaving the tropical average essentially the same. However, we reiterate that important changes in the structure of the temperature responses across the tropics occur as a result of including the zonally symmetric dynamical adjustment.

The non-locality in latitude in the dynamical problem means that the temperature response in the tropics, shown in Fig. 12(a) and (b), is potentially determined in part by the change in trace gases in the extratropics. To quantify this effect, we restricted both the ozone and water vapour annual cycle perturbations to the tropical region between 30° N–S. The net effect of the annual variation in trace gases in the extratropics is to increase the amplitude of the temperature response in the tropics from 2.6 K to

2.8 K peak-to-peak (not shown). The dominant contribution to the annual cycle change in temperature in the tropics is therefore due to ozone and water vapour variations in the tropics.

5 6 Discussion

We have analysed radiative aspects of the prominent annual cycle in temperature in the TTL/tropical lower stratosphere, which has a maximum peak-to-peak amplitude at 70 hPa of ~ 8 K. Building on previous work (Chae and Sherwood, 2007; Fueglistaler et al., 2011), we have applied the seasonally evolving fixed dynamical heating (SEFDH) method to calculate the temperature response to the annual cycle variations in zonal mean ozone and water vapour, derived here from the SWOOSH satellite dataset (Davis et al., 2016). We extend the previous work by presenting explicit results for the effects of water vapour variations and by paying particular attention to the vertical structure of the temperature response and the role of variations in the trace gas mixing ratios in different vertical layers. In our first approach, we have used a SEFDH calculation in which the temperature response to annual variations in a trace gas is calculated independently at each latitude, assuming that the dynamical heating at each height is unchanged from its value in a control state in which the trace gas mixing ratios are constant (and equal to their annual mean values).

We find substantial contributions to the peak-to-peak amplitude of the tropical average (20° N-S) annual cycle in temperatures from ozone: 3.5 ± 0.4 K at 70 hPa, 3.3 ± 0.5 K at 90 hPa and 2.6 ± 0.2 K at 100 hPa, and from water vapour: 0.9 ± 0.1 K at 70 hPa, 1.1 ± 0.1 K at 90 hPa and 0.99 ± 0.03 K at 100 hPa. Whilst the ozone contribution maximises around 70 hPa and is roughly in phase with the observed temperature annual cycle, the water vapour contribution maximises around 90 hPa and is of the opposite phase. Despite the cancellation, the net effect of variations in ozone and water vapour together is substantial and amounts to about 35% of the observed annual cycle at both 70 hPa and 90 hPa and about 45% at 100 hPa (Fig. 7). Our results are broadly consistent with the recent independent work of Gilford and Solomon (2017).

Further SEFDH calculations showed that the ozone-induced temperature variation at 70 hPa is caused primarily (80%) by local ozone variations (Fig. 3). In contrast, the water vapour induced temperature variation is caused by both local and non-local water vapour variations, e.g., 60% of the water vapour induced temperature variation at 90 hPa comes from water vapour variation in the region 100 to 80 hPa and 40% from the region 130 to 100 hPa (Fig. 5). This upward non-local radiative effect is seen throughout the lower stratosphere and has important implications for cold point temperatures. For example, if the amplitude of the annual cycle in water vapour below the cold point was to increase, then the radiative effect would reduce the amplitude in the annual cycle in cold point temperatures and hence reduce the amplitude of the annual cycle in water vapour at and above the cold point.

We also examined the factors controlling the vertical structure of the amplitude of the annual cycle in temperatures. The observed maximum centred on 70 hPa and largely restricted to the 50 to 100 hPa layer arises from a combination of several factors. The vertical structure cannot be explained by clear-sky radiative damping time scales alone, which maximize over a deep region, centred near the cold point at 100 hPa. However, both the dynamical and radiative forcings maximize above the

cold point, and in combination with an inferred upper-tropospheric constraint active below 130 hPa, these lead to the observed maximum at 70 hPa.

Finally, we investigated the effect on the temperature response of relaxing the SEFDH assumption, thereby going beyond the work of Fueglistaler et al. (2011) and Gilford and Solomon (2017). We do this by incorporating the radiative code used for the SEFDH calculations within a 2D (height-latitude) dynamical model. Consistent with dynamical expectations, part of the heating associated with annual cycle variations in both ozone and water vapour drives an annual cycle in the upwelling and that may play a non-negligible role in determining the latitudinal structure of the observed annual cycle in upwelling. This has the effect of reducing latitudinal gradients in the SEFDH-predicted temperature response, particularly across the tropics. However, this modification of the response leaves the tropical (20° N-S) average temperature response essentially unchanged. Therefore the conclusion that the net effect of ozone and water vapour contributes about 35% of the annual cycle peak-to-peak amplitude at 70 hPa and 90 hPa from the SEFDH calculations is robust to including the dynamical adjustment. The detailed latitudinal structure predicted by the SEFDH calculation, however, is not robust to this adjustment.

As explicitly illustrated by Fig. 10(d) (for ozone), the differences between the temperature responses to ozone and water vapour calculated through the SEFDH approach and those calculated using the 2D dynamical model demonstrate that low-latitude temperature features with small latitudinal scales predicted by SEFDH calculations are unlikely to be reproducible when the SEFDH assumption is relaxed, because these features will be smoothed out by the dynamical response. (When considering annual variations in the TTL ‘small latitudinal scales’ means less than about 20° of latitude.) This applies to Fig. 3(b) and 5(b)–(d) in this paper, to previous SEFDH calculations of the temperature response to annual variations in ozone (Fueglistaler et al., 2011, their Fig. 5(b)) and to similar calculations of the effect of recent interannual variations in ozone and water vapour (Gilford et al., 2016, their Fig. 6).

Within the 2D zonally symmetric dynamical formalism presented here we do not take account of changes in wave-induced forces. This effect has been discussed by several authors over the last 30 years or so, including Fels et al. (1980) and Garcia (1987), usually making the assumption that the wave force can be represented by Rayleigh friction (so that local wave force is proportional and opposite to the local zonal velocity). However, it is generally accepted that Rayleigh friction is a poor representation of the wave forces that operate in the upper troposphere and stratosphere. Ming et al. (2016) analyse the effect of the change in wave force in the response to imposed steady localised zonally symmetric heating in a simple 3D model (where the waves are resolved and no Rayleigh friction assumption is necessary) and argue that the effect is to broaden the temperature response, particularly at low latitudes. Latitudinal structure in the imposed heating tends to be balanced by the dynamical heating associated with the meridional velocity response and the change in wave force provides the necessary angular momentum balance. A similar effect is seen in the zonally symmetric problem with Rayleigh friction (e.g., Garcia, 1987, their Fig. 6). There is an analogous effect in the time-dependent zonally symmetric response problem, without any change in wave force or Rayleigh friction, considered in Sect. 5, with the angular momentum balance including the zonal acceleration. Therefore the effect of including the change in wave force in the dynamical problem is, broadly speaking, expected to be similar and in addition to that already seen in the time-dependent zonally symmetric problem, that at low latitudes the dynamical adjustment will smooth the temperature response to latitudinally varying heating. If the change in wave force is

weak then the additional effect will be small. If the change in wave force is strong then the result will be that the smoothing is over a larger range of latitudes. The fact that the observed annual cycle in temperature is coherent over the latitude range 20° N–S, but no more than that (Fig. 1(b)–(d)) suggests that the wave force effect cannot be too strong. Therefore, we expect that our conclusions from the zonally symmetric dynamical problem studied here would not be changed too much if the change in wave force was included. Furthermore, we expect that similar dynamical principles will allow extension to the fully three-dimensional case, implying that an SEFDH calculation will have limited ability to predict geographical (i.e., latitudinal and longitudinal) variations in temperature resulting from geographical latitudinal variations in radiatively active gases or in other relevant quantities such as clouds or aerosol.

Current comprehensive global (chemistry-)climate models show a large spread in the amplitude of the TTL annual cycle in temperature (e.g., Kim et al., 2013), but the quantitative causes of these differences are not well understood. The results of this study show that an erroneous representation of the climatology of ozone and water vapour, as is commonplace amongst such models (e.g., Gettelman et al., 2010), is likely to be a major contributor to poor model performance for capturing the TTL temperature annual cycle. Similar conclusions are likely to apply to interannual variations, e.g., in the 2010–2013 period investigated by Gilford et al. (2016) using SEFDH calculations. Progress in improving the representation of the TTL in comprehensive global models therefore requires consideration of the coupling through transport and radiative effects between dynamics, ozone and water vapour in the TTL. Specific aspects highlighted by our results include a strong sensitivity of ozone radiative effects to mean ozone mixing ratios in the 90 to 70 hPa region, for which models with interactive chemistry simulate a range of values (Gettelman et al., 2010) and for which a range of observation-based gridded datasets exist for climate models that do not include chemistry (Cionni et al., 2011; Bodeker et al., 2013). Furthermore, because of the importance shown here of non-local radiative effects for water vapour in the TTL, modelled cold point temperatures are also likely to be sensitive to the representation of water vapour mixing ratios in the upper tropical troposphere.

Appendix A: FDH calculations

A first order estimate of the effect of specified perturbations to radiative trace gases on temperatures in the TTL and the stratosphere can be made using a fixed dynamical heating (FDH) calculation where it is assumed that the dynamical heating remains constant from the unperturbed to the perturbed state, i.e., that no changes in circulation occur as a result of the perturbation (Fels et al., 1980; Ramanathan and Dickinson, 1979). The time scale for stratospheric adjustment to the perturbation is essentially the stratospheric radiative damping time. This is about 40 days in the tropical lower stratosphere and less than a week near the stratopause, although different techniques estimate different values and furthermore the time scale is dependent on the vertical scale of the heating perturbation (e.g., Dickinson, 1973; Mlynczak et al., 1999; Hitchcock et al., 2010). These stratospheric time scales are relatively short compared to that required for tropospheric temperatures to adjust to the perturbation, because these are strongly constrained to surface temperatures, which particularly in oceanic regions, will evolve only on time scales of months or years. Hence, in FDH calculations, temperatures are held fixed below some level, often corresponding to the

(radiative) tropopause. We choose this level to be 130 hPa, consistent with previous calculations. The reasons for this choice are justified in Sect. 3.

The FDH calculation is a simplified version of the SEFDH calculations and the equations below can be compared to Eq. (2) and (3). Given the background profiles of temperatures and mixing ratios of trace gases ($\bar{T}_0, \bar{\chi}_{\text{O}_3}^0, \bar{\chi}_{\text{H}_2\text{O}}^0$), the dynamical heating,

5 \bar{Q}_{dyn}^0 , is first calculated by assuming the balance

$$\bar{Q}_{\text{rad}}(\bar{T}^0, \bar{\chi}_{\text{O}_3}^0, \bar{\chi}_{\text{H}_2\text{O}}^0) + \bar{Q}_{\text{dyn}}^0 = 0. \quad (\text{A1})$$

The dynamical heating is not a function of time unlike in the SEFDH calculations. A perturbation is then applied to trace gas mixing ratios ($\Delta\bar{\chi}_{\text{O}_3}, \Delta\bar{\chi}_{\text{H}_2\text{O}}$) and the equilibrium temperature state, $\bar{T}^0 + \Delta\bar{T}$, is obtained from

$$\bar{Q}_{\text{rad}}(\bar{T}^0 + \Delta\bar{T}, \bar{\chi}_{\text{O}_3}^0 + \Delta\bar{\chi}_{\text{O}_3}, \bar{\chi}_{\text{H}_2\text{O}}^0 + \Delta\bar{\chi}_{\text{H}_2\text{O}}) + \bar{Q}_{\text{dyn}}^0 = 0. \quad (\text{A2})$$

10 Time averaged profiles of ozone and water vapour from the SWOOSH dataset and the annual mean temperature from ERA-Interim at the Equator are used as the base profile and the trace gases are then perturbed. The calculation is done at the Equator on January 1 and the albedo is set to 0.085. The 100 pressure levels used in all radiative calculations are: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 25, 27, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 93, 95, 97, 100, 103, 105, 107, 110, 113, 115, 117, 120, 123, 125, 127, 130, 133, 135, 137, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190,
15 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, 265, 270, 275, 280, 285, 290, 295, 300, 320, 330, 340, 350, 370, 400, 450, 500, 600, 700, 800, 900, 1000 hPa.

Numerically, the FDH calculation is done by iterating the temperatures forward with a time step of one day using the long-wave heating rates to find the new equilibrium temperature. The values are considered to have converged when the temperature change and the fluxes between pressure levels after consecutive time steps falls below $5 \times 10^{-4} \text{ K}$ and $1 \times 10^{-7} \text{ K m}^{-1}$ respectively. In practice, these thresholds are reached after about 500 days which is much larger than any radiative time scales in the stratosphere, hence, ensuring that the temperatures in the stratosphere have converged.

In Sect. A1 and A2 below, we describe in detail the temperature response to example perturbations in ozone and in water vapour. These provide helpful background for understanding the response to the annual cycle in these two gases reported in Sect. 3.

25 A1 Ozone perturbation

The example perturbation applied to ozone mixing ratios is a reduction in mixing ratios in the lower stratosphere (solid line in Fig. 13(a) left). This is a simple representation of lower stratospheric mixing ratios in NH winter, relative to the annual mean. The perturbation is a Gaussian of the form $A_0 \exp[-0.5((z - 18.6)/2)^2]$ where $A_0 = -0.07 \text{ (ppmv)}$ and $z = -7 \log(p/1 \times 10^5) \text{ km}$. Removing ozone in the lower stratosphere leads to an instantaneous local decrease in the longwave and shortwave heating (Fig. 13(a) right) and results in a local decrease in the temperature in an FDH calculation, Fig. 13(b) (where ‘local’ refers to the vertical region in which the perturbation in mixing ratios is applied).

The time evolution of various components of the longwave radiative heating after the perturbation is applied are shown in Fig. 13(b)–(e), respectively total longwave heating and then the individual contributions from ozone, carbon dioxide and




Figure 13. (a) Left: ozone reference profile (solid) and perturbed profile (dashed) used in the FDH calculation. Right: instantaneous change in heating rate from perturbation. (b) Temperature change resulting from the ozone perturbation. The calculation is done at the Equator on January 1. The lines correspond to perturbations of $A_0 = -0.2, -0.1, -0.07$ (black line), $-0.05, 0.05, 0.07, 0.1$ and 0.2 ppmv from left to right at the maximum temperature change (see text for more details). For the perturbation of -0.7 ppmv, (c) total longwave heating rate from all the constituents, (d) longwave heating rate due to ozone, (e) longwave heating rate due to carbon dioxide and (f) longwave heating rate due to water vapour.

water vapour. The instantaneous effect of the reduction in ozone mixing ratios is to cause a decrease in both the shortwave heating, and the local longwave heating, because of reduction in local longwave absorption. The shortwave change, which has peak amplitude -1.1×10^{-2} K day $^{-1}$, occurs because of reduced shortwave absorption and is essentially proportional to the local change in mixing ratio. The instantaneous longwave change is significantly larger, with peak amplitude -4×10^{-2} K day $^{-1}$, and, in addition to the local decrease, there is an increase, with similar peak amplitude in the region above the mixing ratio perturbation (see Fig. 13(d)). The explanation for this vertical structure is that, because ozone mixing ratios are small in the troposphere, in the lower stratosphere there is a substantial upwelling flux of longwave radiation of wavelength relevant to ozone ($9.6 \mu\text{m}$ -band), and the imposed perturbation in ozone mixing ratios leads to less local absorption of this upwelling radiation with, correspondingly increased absorption above the perturbation. Note that another potential effect of the perturbation to ozone mixing ratios is reduced local emission which would imply a local heating. Figure 13(d) shows that any effect of change in emission is dominated by the changed absorption of upwelling radiation.

In the response to the instantaneous change in heating just described, the temperature and hence the longwave fluxes change, with both carbon dioxide (Fig. 13(e)) and to a lesser extent water vapour (Fig. 13(f)) contributing significantly. Note that changes in the ozone longwave heating, after the instantaneous change resulting from the perturbation to ozone mixing ratios, are weak, suggesting that it plays little role in the temperature adjustment. An equilibrium is reached where the net longwave heating (Fig. A1(c)) balances the reduction in shortwave heating. The equilibrium temperature change is dominated by a local decrease centred on 70 hPa (i.e., the centre of the region where ozone mixing ratios were perturbed). Several time scales are involved in the adjustment process and Fig. 13(c) shows that the heating rates and hence the temperature are still evolving after 100 days. This justifies the use of an SEFDH rather than an FDH calculation when studying the annual cycle in temperatures.

Further experiments show that the FDH temperature response varies approximately linearly with the peak value of the Gaussian perturbation in the range -0.1 to 0.1 ppmv (thin grey lines in Fig. 13(b)), so that the detailed time evolution described above continues to hold if heating and temperature anomalies are multiplied by the appropriate factor. In particular a modest increase in ozone mixing ratios will lead to a local temperature increase, in which the net (negative) change in longwave heating balances an increase in shortwave heating. For mixing ratio anomalies with peak values of ± 0.2 ppmv, significant non-linear effects appear.

15 A2 Water vapour perturbation

Following the approach in Appendix A1 above, a corresponding calculation is now described in which water vapour is perturbed by removing a Gaussian of the form $B_0 \exp[-0.5((z - 16.9)/1.5)^2]$ where $B_0 = 1.0$ (ppmv) (Fig. 14(a) left) which leads to an instantaneous local decrease in the shortwave and a local increase in the longwave radiation (Fig. 14(a) right). This is also a very simple representation of lower stratospheric mixing ratios in NH winter, relative to the annual mean. As in Appendix A1, Fig. 14(c)–(f) respectively show the total longwave heating and then the individual contributions from ozone, carbon dioxide and water vapour, during the evolution in response to the water vapour perturbation.

The abundance of water vapour in the troposphere means it is relatively opaque to upwelling longwave radiation in the main water vapour absorption bands. This means that, in contrast to ozone, the dominant instantaneous effect in the longwave of locally reducing the water vapour in the lower stratosphere, is to cause less local emission, i.e., local heating, and, correspondingly, less non-local absorption in neighbouring regions, i.e., non-local cooling, rather than any effect on absorption of upwelling radiation. This can be seen in the water vapour longwave heating shown in Fig. 14(f). Note that the change in non-local absorption is seen primarily in the upper troposphere below the region where the mixing ratios are reduced, because background water vapour mixing ratios are relatively large there compared to those in the stratosphere. The reduction in water vapour mixing ratio also leads to a reduction in shortwave absorption, as was the case for ozone, but the magnitude ($-0.3 \times 10^{-2} \text{ K day}^{-1}$) is smaller than the corresponding change in longwave heating ($4.3 \times 10^{-2} \text{ K day}^{-1}$).

In the evolution following the initial instantaneous change in heating the longwave heating contributions to due to carbon dioxide, water vapour and ozone all play a role to limit the temperature response and redistribute it in the vertical. (Fig. 14(d)–(f)). In particular the initial local increase in temperatures is transmitted in the vertical through longwave fluxes in the carbon dioxide bands to give subsequent temperature increases significantly above the layer in which water vapour mixing ratios were

perturbed. This sort of behaviour is not captured by a local Newtonian cooling approximation. As was the case for ozone, the longwave heating (and hence the temperatures) continue to evolve beyond 100 days. This suggests that a sequence of quasi-steady FDH calculations would be inadequate for studying the annual cycle in temperatures and again justifies the use of the SEFDH approach.

- 5 Experiments with different amplitudes of perturbation to water vapour mixing ratio (Figure 19(b)) show that the response is linear for peak values up to ± 1.0 ppmv, with non-linear effects visible at ± 2.0 ppmv. Note that a similar amplitude and shape of perturbation as the ozone perturbation with $A_0 = 0.2$ ppmv is shown for comparison as a dashed grey line in Fig. 14(b) and the magnitude of the temperature change is small (0.14 K at 70 K) compared to that for the equivalent ozone perturbation (2.8 K at 70 K).




Figure 14. Similar to the plots in Figure 13 but for water vapour. (a) Left: water vapour profile (solid) and perturbation (dashed) used in the FDH calculation. Right: instantaneous change in heating rate from perturbation. (b) Temperature change resulting from the water vapour perturbation for $B_0 = 2.0, 1.0, 0.5, 0.2, -0.2, -0.5, -1.0$ (thick black line) and -2.0 ppmv from left to right at the maximum temperature change for the solid lines. For comparison, a perturbation in water vapour of a similar form to the ozone perturbation with $A_0 = 0.2$ ppmv is also included (dashed grey line) (see text for more details). For the perturbation $B_0 = -1.0$ ppmv, (c) is total longwave heating rate from all the constituents, (d) is longwave heating rate due to ozone, (e) is longwave heating rate due to carbon dioxide and (f) is longwave heating rate due to water vapour.

Appendix B: Statistical methods

Estimates of the 95% confidence intervals are shown for the SEFDH calculations in Fig. 7. For ozone and water vapour in the SWOOSH dataset, a combined uncertainty arising from the uncertainties in the various instruments and a standard deviation arising from interannual variability can be obtained. These two quantities are provided as part of the SWOOSH dataset and are of similar magnitude in the region of interest. A 95% confidence interval is obtained for each month by summing these two uncertainties in quadrature and assuming that each year in the dataset is independent. This assumption has been checked and is adequate. The SEFDH calculation for each constituent is then repeated to give bounds for the temperature change given the uncertainty in that constituent only. For example, the water vapour uncertainty in Fig. 7 is small and only reflects that coming from the water vapour dataset and not from differences in ozone which will also affect the temperature change from water vapour. However, the combined effect of both uncertainties is present in the calculation of the temperature change from both ozone and water vapour. When calculating the peak-to-peak amplitude, the uncertainties at the maximum amplitude and minimum amplitude are added in quadrature.

The residual mean vertical velocity in reanalysis datasets has a large interannual variability and this is the only source of uncertainty taken into account in the calculation in Fig. 7. Again, in estimating this quantity, we assume that each year of the dataset is independent. This leads to a peak-to-peak amplitude from the dynamical heating averaged over 20° N–S at 70 hPa of 5.6 ± 0.6 K and of 1.5 ± 0.6 K at 90 hPa. In addition, there are other large discrepancies in estimates of the dynamical heating which are not taken into account in this calculation. For example, the difference between calculating the dynamical heating directly from $\overline{w^* \overline{S}}$ and from the thermodynamic equation can be as high as about 40% in certain months. A full treatment of all the sources of uncertainty in this calculation is beyond the scope of this work.

Appendix C: Background ozone mixing ratio

The greatest sensitivity of the temperature changes calculated in the SEFDH calculations is to the background value of ozone. A set of illustrative SEFDH calculations are presented below to show how this affects the temperature change for the ozone annual cycle. Fig. 15(a) shows an illustrative perturbation (third line from the left, solid grey) to the annual mean ozone profile (middle line, solid black) used in the SEFDH calculations in Sect. 3.1. (This perturbation is a decrease of twice the standard deviation of the annual mean values in the SWOOSH dataset adjusted for time correlations between the years.) This decrease in the annual mean ozone leads to an increase in the peak-to-peak amplitude of the temperature change due to ozone at 70 hPa by about 0.16 K, Fig 15(b). Similarly, an increase in the annual mean ozone leads to a smaller amplitude ozone annual cycle. Further experiments show that the change in the peak-to-peak amplitude varies roughly linearly with the change in the background ozone mixing ratio within the range of values shown in Fig. 15(a). These values are of a magnitude comparable to those seen in other ozone datasets. For instance, Tummon et al. (2015) quote a spread of about $\pm 10\%$ in the annual mean ozone in the lower stratosphere between seven newly available merged satellite ozone profile data sets. The spread in individual satellite instruments is larger with differences of up to $\pm 20\%$ from the multi-instrument mean (Tegtmeier et al., 2013).

Fig. 15(c) shows the contribution from different pressure ranges to the change in the temperature annual cycle from the illustrative perturbation. A decrease in the annual mean ozone increases the upwelling longwave radiation reaching 70hPa leading to a larger temperature annual cycle response at 70hPa. This can be seen from the largest contributions coming from the regions 90 to 70hPa and below 90hPa.




Figure 15. Annual mean ozone profile averaged between 20° N–S (middle solid black line). Solid dark grey line to the left of the middle line shows an illustrative perturbation to the annual mean profile where the ozone background value is decreased. Solid grey lines represent negative perturbations two times and three times this perturbation. Corresponding positive perturbations are shown as dashed lines. (b) Difference to the SEFDH temperature change at 70hPa for the ozone annual cycle due to the different annual mean ozone values in (a). The darker solid grey line shows the temperature change for the illustrative perturbation. The grey lines and dotted lines correspond to the increasingly larger negative and positive perturbations respectively as shown in (a). (c) The contribution of different ranges of pressure levels to the temperature change for the illustrative perturbation.

- 5 *Acknowledgements.* The authors wish to thank Stephan Fueglistaler for helpful discussions and Manoj Joshi for assistance with the IGCM3.1 model and acknowledge helpful comments from the two referees. AM, ACM and PHi were supported by an ERC ACCI grant (project no. 267760). In addition, ACM was supported by an AXA Postdoctoral Research Fellowship, and a NERC Research Fellowship (grant NE/M018199/1) and PHi was supported by a NSERC postdoctoral fellowship. PHa acknowledges support from the IDEX Chaires d'Attractivité programme of l'Université Fédérale de Toulouse, Midi-Pyrénées.

References

- Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle atmosphere dynamics, Academic Press, Orlando, 1987.
- Bodeker, G. E., Hassler, B., Young, P. J., and Portmann, R. W.: A vertically resolved, global, gap-free ozone database for assessing or constraining global climate model simulations, *Earth System Science Data*, 5, 31–43, doi:10.5194/essd-5-31-2013, 2013.
- 5 Bresser, G., Manning, A. J. L., Pawson, S., and Rodgers, C. D.: A New Parameterization of Scale-Dependent Radiative Rates in the Stratosphere, *Journal of the Atmospheric Sciences*, 52, 4429–4447, doi:10.1175/1520-0469(1995)052<4429:ANPOSD>2.0.CO;2, 1995.
- Bruhl, C., Drayson, S. R., Russell, J. M., Crutzen, P. J., McInerney, J. M., Purcell, P. N., Claude, H., Gernandt, H., McGee, T. J., McDerimid, I. S., and Gunson, M. R.: Halogen occultation experiment ozone channel validation, *J. Geophys. Res.*, 101, 10 217–10 240, doi:10.1029/95JD02031, 1996.
- 10 Chae, J. H. and Sherwood, S. C.: Annual temperature cycle of the tropical tropopause: A simple model study, *Journal of Geophysical Research*, 112, D19 111, doi:10.1029/2006JD007956, 2007.
- Cionni, I., Eyring, V., Lamarque, J. F., Randel, W. J., Stevenson, D. S., Wu, F., Bodeker, G. E., Shepherd, T. G., Shindell, D. T., and Waugh, D. W.: Ozone database in support of CMIP5 simulations: results and corresponding radiative forcing, *Atmospheric Chemistry and Physics*, 11, 11 267–11 292, doi:10.5194/acp-11-11267-2011, 2011.
- 15 Davis, S. M., Rosenlof, K. H., Hassler, B., Hurst, D. F., Read, W. G., Vömel, H., Selkirk, H., Fujiwara, M., and Damadeo, R.: The Stratospheric Water and Ozone Satellite Homogenized (SWOOSH) database: A long-term database for climate studies, *Earth System Science Data Discussions*, pp. 1–59, doi:10.5194/essd-2016-16, 2016.
- de F. Forster, P. M., Blackburn, M., Glover, R., and Shine, K. P.: An examination of climate sensitivity for idealised climate change experiments in an intermediate general circulation model, *Climate Dynamics*, 16, 833–849, doi:10.1007/s003820000083, 2000.
- 20 Dee, D. P., Uppala, S. M., Simmons, a. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. a., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, a. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, a. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, a. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, doi:10.1002/qj.828, 2011.
- 25 Dickinson, R. E.: Method of parameterization for infrared cooling between altitudes of 30 and 70 kilometers, *Journal of Geophysical Research*, 78, 4451–4457, doi:10.1029/JC078i021p04451, 1973.
- Fels, S. B.: A Parameterization of Scale-Dependent Radiative Damping Rates in the Middle Atmosphere, *Journal of the Atmospheric Sciences*, 39, 1141–1152, 1982.
- 30 Fels, S. B., Mahlman, J. D., Schwarzkopf, M. D., and Sinclair, R. W.: Stratospheric Sensitivity to Perturbations in Ozone and Carbon Dioxide: Radiative and Dynamical Response, *Journal of the Atmospheric Sciences*, 37, 2265–2297, doi:10.1175/1520-0469(1980)037<2265:SSTPIO>2.0.CO;2, 1980.
- Folkens, I., Bernath, P., Boone, C., Lesins, G., Livesey, N., Thompson, A. M., Walker, K., and Witte, J. C.: Seasonal cycles of O₃, CO, and convective outflow at the tropical tropopause, *Geophysical Research Letters*, 33, 1–5, doi:10.1029/2006GL026602, 2006.
- 35 Forster, P. M. d. F. and Shine, K. P.: Assessing the climate impact of trends in stratospheric water vapor, *Geophysical Research Letters*, 29, 1086, doi:10.1029/2001GL013909, 2002.

- Forster, P. M. F., Freckleton, R. S., and Shine, K. P.: On aspects of the concept of radiative forcing, *Climate Dynamics*, 13, 547–560, doi:10.1007/s003820050182, 1997.
- Fueglistaler, S. and Haynes, P. H.: Control of interannual and longer-term variability of stratospheric water vapor, *Journal of Geophysical Research Atmospheres*, 110, 1–14, doi:10.1029/2005JD006019, 2005.
- 5 Fueglistaler, S., Bonazzola, M., Haynes, P., and Peter, T.: Stratospheric water vapor predicted from the Lagrangian temperature history of air entering the stratosphere in the tropics, *Journal of Geophysical Research*, 110, D08 107, doi:10.1029/2004JD005516, 2005.
- Fueglistaler, S., Dessler, a. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Reviews of Geophysics*, 47, 1–31, doi:10.1029/2008RG000267, 2009.
- Fueglistaler, S., Haynes, P. H., and Forster, P. M.: The annual cycle in lower stratospheric temperatures revisited, *Atmospheric Chemistry and Physics*, 11, 3701–3711, doi:10.5194/acp-11-3701-2011, 2011.
- 10 Garcia, R. R.: On the Mean Meridional Circulation of the Middle Atmosphere, *Journal of Atmospheric Sciences*, 44, 3599–3609, doi:10.1175/1520-0469(1987)044<3599:OTMMCO>2.0.CO;2, 1987.
- Gettelman, A., Hegglin, M. I., Son, S.-W., Kim, J., Fujiwara, M., Birner, T., Kremser, S., Rex, M., Añel, J. A., Akiyoshi, H., Austin, J., Bekki, S., Braesike, P., Brühl, C., Butchart, N., Chipperfield, M., Dameris, M., Dhomse, S., Garny, H., Hardiman, S. C., Jöckel, P., Kinnison, D. E., Lamarque, J. F., Mancini, E., Marchand, M., Michou, M., Morgenstern, O., Pawson, S., Pitari, G., Plummer, D., Pyle, J. A., Rozanov, E., Scinocca, J., Shepherd, T. G., Shibata, K., Smale, D., Teyssèdre, H., and Tian, W.: Multimodel assessment of the upper troposphere and lower stratosphere: Tropics and global trends, *Journal of Geophysical Research*, 115, 1–22, doi:10.1029/2009JD013638, 2010.
- 15 Gilford, D. M. and Solomon, S.: Radiative effects of stratospheric seasonal cycles in the tropical upper troposphere and lower stratosphere, *Journal of Climate*, pp. JCLI–D–16–0633.1, doi:10.1175/JCLI-D-16-0633.1, 2017.
- 20 Gilford, D. M., Solomon, S., and Portmann, R. W.: Radiative Impacts of the 2011 Abrupt Drops in Water Vapor and Ozone in the Tropical Tropopause Layer, *Journal of Climate*, 29, 595—612, doi:10.1175/JCLI-D-15-0167.1, 2016.
- Hardiman, S. C., Boutle, I. A., Bushell, A. C., Butchart, N., Cullen, M. J. P., Field, P. R., Furtado, K., Manners, J. C., Milton, S. F., Morcrette, C., O’Connor, F. M., Shipway, B. J., Smith, C., Walters, D. N., Willett, M. R., Williams, K. D., Wood, N., Abraham, L. N., Keeble, J., Maycock, A. C., Thuburn, J., and Woodhouse, M. T.: Processes controlling tropical tropopause temperature and stratospheric water vapor in climate models, *Journal of Climate*, 28, 6516–6535, doi:10.1175/JCLI-D-15-0075.1, 2015.
- 25 Harries, J. E., Iii, J. M. R., Tuck, A. F., Gordley, L. L., Purcell, P., Stone, K., Bevilacqua, R. M., and Traub, W. A.: Validation of measurements of water vapour from the Halogen Occultation Experiment (HALOE), *Journal of Geophysical Research*, 101, 10 205–10 216, 1996.
- Harris, N. R. P., Hassler, B., Tummon, F., Bodeker, G. E., Hubert, D., Petropavlovskikh, I., Steinbrecht, W., Anderson, J., Bhartia, P. K., Boone, C. D., Bourassa, A., Davis, S. M., Degenstein, D., Delcloo, A., Frith, S. M., Froidevaux, L., Godin-Beekmann, S., Jones, N., Kurylo, M. J., Kyrölä, E., Laine, M., Leblanc, S. T., Lambert, J. C., Liley, B., Mahieu, E., Maycock, A., De Mazière, M., Parrish, A., Querel, R., Rosenlof, K. H., Roth, C., Sioris, C., Staehelin, J., Stolarski, R. S., Stübi, R., Tamminen, J., Vigouroux, C., Walker, K. A., Wang, H. J., Wild, J., and Zawodny, J. M.: Past changes in the vertical distribution of ozone - Part 3: Analysis and interpretation of trends, *Atmospheric Chemistry and Physics*, 15, 9965–9982, doi:10.5194/acp-15-9965-2015, 2015.
- 30 Haynes, P.: Stratospheric dynamics, *Annual Review of Fluid Mechanics*, 37, 263–293, doi:10.1146/annurev.fluid.37.061903.175710, 2005.
- Haynes, P., McIntyre, M., Shepherd, T., Marks, C., and Shine, K.: On the ‘Downward Control’ of Extratropical Diabatic Circulations by Eddy-Induced Mean Zonal Forces., *Journal of Atmospheric Sciences*, 48, 651–680, doi:10.1175/1520-0469(1991)048<0651:OTCOED>2.0.CO;2, 1991.

- Held, I. M. and Suarez, M. J.: A Proposal for the Intercomparison of the Dynamical Cores of Atmospheric General Circulation Models, *Bulletin of the American Meteorological Society*, 75, 1825–1830, doi:10.1175/1520-0477(1994)075<1825:APFTIO>2.0.CO;2, 1994.
- Hitchcock, P., Shepherd, T. G., and Yoden, S.: On the Approximation of Local and Linear Radiative Damping in the Middle Atmosphere, *Journal of the Atmospheric Sciences*, 67, 2070–2085, doi:10.1175/2009JAS3286.1, 2010.
- 5 Hoskins, B. J. and Simmons, A. J.: A multi-layer spectral model and the semi-implicit method, *Quarterly Journal of the Royal Meteorological Society*, 101, 637–655, doi:10.1002/qj.49710142918, 1975.
- Kim, J., Grise, K. M., and Son, S.-W. W.: Thermal characteristics of the cold-point tropopause region in CMIP5 models, *Journal of Geophysical Research: Atmospheres*, 118, 8827–8841, doi:10.1002/jgrd.50649, 2013.
- Lambert, A., Read, W. G., Livesey, N. J., Santee, M. L., Manney, G. L., Froidevaux, L., Wu, D. L., Schwartz, M. J., Pumphrey, H. C.,
10 Jimenez, C., Nedoluha, G. E., Cofield, R. E., Cuddy, D. T., Daffer, W. H., Drouin, B. J., Fuller, R. a., Jarnot, R. F., Knosp, B. W., Pickett, H. M., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., Wagner, P. a., Waters, J. W., Jucks, K. W., Toon, G. C., Stachnik, R. a., Bernath, P. F., Boone, C. D., Walker, K. a., Urban, J., Murtagh, D., Elkins, J. W., and Atlas, E.: Validation of the Aura Microwave Limb Sounder middle atmosphere water vapor and nitrous oxide measurements, *Journal of Geophysical Research: Atmospheres*, 112, 1–24, doi:10.1029/2007JD008724, 2007.
- 15 Logan, J. A.: An analysis of ozonesonde data for the troposphere: Recommendations for testing 3-D models and development of a gridded climatology for tropospheric ozone, *Journal of Geophysical Research: Atmospheres*, 104, 16 115–16 149, doi:10.1029/1998JD100096, 1999.
- Maycock, A. C., Shine, K. P., and Joshi, M. M.: The temperature response to stratospheric water vapour changes, *Quarterly Journal of the Royal Meteorological Society*, 137, 1070–1082, doi:10.1002/qj.822, 2011.
- 20 Maycock, A. C., Joshi, M. M., Shine, K. P., Davis, S. M., and Rosenlof, K. H.: The potential impact of changes in lower stratospheric water vapour on stratospheric temperatures over the past 30 years, *Quarterly Journal of the Royal Meteorological Society*, 140, 2176–2185, doi:10.1002/qj.2287, 2014.
- Ming, A., Hitchcock, P., and Haynes, P.: The Response of the Lower Stratosphere to Zonally Symmetric Thermal and Mechanical Forcing, *Journal of the Atmospheric Sciences*, 73, 1903–1922, doi:10.1175/JAS-D-15-0294.1, 2016.
- 25 Mlynczak, M. G., Mertens, C. J., Garcia, R. R., and Portmann, R. W.: A detailed evaluation of the stratospheric heat budget: 2. Global radiation balance and diabatic circulations, *Journal of Geophysical Research: Atmospheres*, 104, 6039–6066, doi:10.1029/1998JD200099, 1999.
- Morcrette, J.-J.: Radiation and cloud radiative properties in the European Centre for Medium Range Weather Forecasts forecasting system, *Journal of Geophysical Research: Atmospheres*, 96, 9121–9132, doi:10.1029/89JD01597, 1991.
- 30 Mote, P. W., Rosenlof, K. H., McIntyre, E., Carr, E. S., Gille, J. C., Holton, R., Kinnersley, S., Pumphrey, H. C., Russell III, J. M., and Waters, J. W.: An atmospheric tape recorder: The imprint of tropical tropopause temperatures on stratospheric water vapor, *Journal of Geophysical Research*, 101, 3989–4006, 1996.
- Plumb, R. A.: Zonally-symmetric Hough modes and meridional circulations in the middle atmosphere., *Journal of Atmospheric Sciences*, 39, 983–991, 1982.
- 35 Ramanathan, V. and Dickinson, R. E.: The Role of Stratospheric Ozone in the Zonal and Seasonal Radiative Energy Balance of the Earth-Troposphere System, *Journal of the Atmospheric Sciences*, 36, 1084–1104, doi:10.1175/1520-0469(1979)036<1084:TROSOI>2.0.CO;2, 1979.

- Randel, W. J. and Jensen, E. J.: Physical processes in the tropical tropopause layer and their roles in a changing climate, *Nature Geoscience*, 6, 169–176, doi:10.1038/ngeo1733, 2013.
- Randel, W. J., Wu, F., Gettelman, A., Russell III, J. M., Zawodny, J. M., and Oltmans, S. J.: Seasonal variation of water vapor in the lower stratosphere observed in Halogen Occultation Experiment data, *Journal of Geophysical Research*, 106, 14 313–14 325, doi:10.1029/2001JD900048, 2001.
- Randel, W. J., Garcia, R. R., and Wu, F.: Time-Dependent Upwelling in the Tropical Lower Stratosphere Estimated from the Zonal-Mean Momentum Budget, *Journal of the Atmospheric Sciences*, 59, 2141–2152, doi:10.1175/1520-0469(2002)059<2141:TDUITT>2.0.CO;2, 2002.
- Randel, W. J., Wu, F., and Rios, W.: Thermal variability of the tropical tropopause region derived from GPS / MET observations, *Journal of Geophysical Research*, 108, 1–12, doi:10.1029/2002JD002595, 2003.
- Randel, W. J., Park, M., Wu, F., and Livesey, N.: A Large Annual Cycle in Ozone above the Tropical Tropopause Linked to the Brewer–Dobson Circulation, *Journal of the Atmospheric Sciences*, 64, 4479–4488, doi:10.1175/2007JAS2409.1, 2007.
- Reed, R. J. and Vlcek, C. L.: The Annual Temperature Variation in the Lower Tropical Stratosphere, *Journal of Atmospheric Sciences*, 26, 163–167, doi:10.1175/1520-0469(1969)026<0163:TATVIT>2.0.CO;2, 1969.
- Seviour, W. J. M., Butchart, N., and Hardiman, S. C.: The Brewer-Dobson circulation inferred from ERA-Interim, *Quarterly Journal of the Royal Meteorological Society*, 138, 878–888, doi:10.1002/qj.966, 2012.
- Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G.-K.: Contributions of stratospheric water vapor to decadal changes in the rate of global warming., *Science*, 327, 1219–1223, doi:10.1126/science.1182488, 2010.
- Stolarski, R. S., Waugh, D. W., Wang, L., Oman, L. D., Douglass, A. R., and Newman, P. A.: Seasonal variation of ozone in the tropical lower stratosphere: Southern tropics are different from northern tropics, *Journal of Geophysical Research: Atmospheres*, 119, 6196–6206, doi:10.1002/2013JD021294, 2014.
- Tegtmeier, S., Hegglin, M. I., Anderson, J., Bourassa, A., Brohede, S., Degenstein, D., Froidevaux, L., Fuller, R., Funke, B., Gille, J., Jones, A., Kasai, Y., Krüger, K., Kyrölä, E., Lingenfelser, G., Lumpe, J., Nardi, B., Neu, J., Pendlebury, D., Remsberg, E., Rozanov, A., Smith, L., Toohey, M., Urban, J., Von Clarmann, T., Walker, K. A., and Wang, R. H. J.: SPARC Data Initiative: A comparison of ozone climatologies from international satellite limb sounders, *Journal of Geophysical Research: Atmospheres*, 118, 12 229–12 247, doi:10.1002/2013JD019877, 2013.
- Tummon, F., Hassler, B., Harris, N. R. P., Staehelin, J., Steinbrecht, W., Anderson, J., Bodeker, G. E., Bourassa, A., Davis, S. M., Degenstein, D., Frith, S. M., Froidevaux, L., Kyrölä, E., Laine, M., Long, C., Penckwitt, A. A., Sioris, C. E., Rosenlof, K. H., Roth, C., Wang, H. J., and Wild, J.: Intercomparison of vertically resolved merged satellite ozone data sets: Interannual variability and long-term trends, *Atmospheric Chemistry and Physics*, 15, 3021–3043, doi:10.5194/acp-15-3021-2015, 2015.
- Zhong, W. and Haigh, J. D.: Improved Broadband Emissivity Parameterization for Water Vapor Cooling Rate Calculations, *Journal of Atmospheric Sciences*, 52, 124–138, 1995.