

# Supplementary Information for: Hygroscopic Properties of Amminium Sulphate Aerosols

Grazia Rovelli<sup>1,2</sup>, Rachael E.H. Miles<sup>1</sup>, Jonathan P. Reid<sup>1</sup>, Simon L. Clegg<sup>3</sup>

<sup>1</sup> School of Chemistry, University of Bristol, Bristol, BS8 1TS, UK

5 <sup>2</sup> Department of Earth and Environmental Sciences, University of Milano-Bicocca, 20124 Milan, Italy

<sup>3</sup> School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, UK

Correspondence to: Jonathan P. Reid ([j.p.reid@bristol.ac.uk](mailto:j.p.reid@bristol.ac.uk))

10

## Supplementary Information outline:

- **Table S1:** List and description of the uncertainties taken into account when calculating the hygroscopicity properties of aerosols from comparative kinetics experiments in a CK-EDB.
- 15 - **Table S2:** Measured densities and refractive indices of amminium sulphates solutions.
- **Table S3:** 3<sup>rd</sup> order polynomial fitted parameters for densities, calculated melt densities and melt refractive indices calculated with molar refraction mixing rule for the six considered amminium sulphates.
- **Table S4:** Comparison of pure melt densities from this work and from Clegg et al. (2013) and solid densities from Qiu and Zhang (2012).
- 20 - **Table S5:** Coefficients of the fitted equations of the measured apparent molar volumes.
- **Table S6:** Extended ZSR fitted coefficients at two fixed water activities ( $a_w$ , 0.80 and 0.925) for mixtures of amminium sulphate salts and sulphuric acid.
  
- **Figure S1:** Measured densities of amminium sulphates solutions and 3<sup>rd</sup> order polynomial fittings of data and measured refractive indices as a function of mass fraction and the corresponding molar refraction mixing rule fitting.
- 25 - **Figure S2:** Comparison of measured  $GF_r$  from CK-EDB experiments and  $GF_r$  data from H-TDMA measurements by Qiu and Zhang (2012)
- **Figure S3:** Measured and fitted apparent molar volumes of amminium sulphates in aqueous solutions from this work and from Clegg et al. (2013).

**Table S1:** List and description of the uncertainties taken into account when calculating the hygroscopicity properties of aerosols from comparative kinetics experiments in a CK-EDB.

Quantity	Symbol/Formula	Uncertainties
Water activity	$a_w$	<ul style="list-style-type: none"> <li>- <math>a_w</math>: uncertainty from RH determination in the EDB measurements (Davies et al., 2013):</li> <li>- RH&gt;80% (pure water used as a probe):  <math display="block">RH = RH_w^{+(0.169RH_w^2 - 0.364RH_w + 0.194) - (-0.020RH_w + 0.021)}</math> </li> <li>- RH&lt;80% (NaCl used as a probe):  <math display="block">RH = RH_{eq}^{+(-0.0175RH_{eq}^2 - 0.0005RH_{eq} + 0.017) - (-0.0266RH_{eq}^2 + 0.0086RH_{eq} + 0.017)}</math> </li> </ul>
Radial growth factor	$GF_r = \frac{a}{a_{dry}}$	<ul style="list-style-type: none"> <li>- <math>a</math> (measured): <math>\pm 100</math> nm (Davies, 2014), &lt;1% for droplets bigger than 10 <math>\mu\text{m}</math>.<sup>(a)</sup></li> <li>- <math>a_{dry}</math>: uncertainty on the initial radius extrapolation at <math>t_0</math> (<math>a_{dry}^{+150\text{ nm}}_{-100\text{ nm}}</math>); uncertainty on the initial solution concentration.</li> </ul>
Mass fraction	$mfs = \frac{m_{solute}}{m_{tot}}$	<ul style="list-style-type: none"> <li>- <math>m_{solute}</math> (dry mass from the initial concentration): <math>a_{dry}^{+150\text{ nm}}_{-100\text{ nm}}</math>; uncertainty on the initial solution concentration.<sup>(b)</sup></li> <li>- <math>m_{tot}</math>: uncertainty from radius determination.</li> </ul>
Mole curves	$\frac{n_{water}}{n_{solute}}$	<ul style="list-style-type: none"> <li>- <math>n_{water}</math>, calculated from <math>m_{water} = m_{tot} - m_{solute}</math> (see uncertainties indicated above for <math>mfs</math>).</li> </ul>
Stoichiometric osmotic coefficient	$\phi_{st} = \frac{-\ln(a_w)}{(m_{st} \cdot M_w / 1000)}$	<ul style="list-style-type: none"> <li>- <math>m_{st}</math> (stoichiometric molality): uncertainties on <math>n_{solute}</math> and <math>m_{water}</math> (see uncertainties indicated for <math>mfs</math> and for <math>n_{water}/n_{solute}</math>).</li> </ul>

5 Symbols:  $a_w$ ; water activity;  $GF_r$  – radial growth factor;  $a$  – droplet radius;  $a_{dry}$  – dry particle radius;  $mfs$  – mass fraction of solute;  $m_{solute}$  – mass of solute in the particle;  $m_{tot}$  – total particle mass;  $n_{water}$  – moles of water;  $n_{solute}$  – moles of solute;  $\phi_{st}$  – stoichiometric osmotic coefficients;  $m_{st}$  – stoichiometric molality;  $M_w$  – molecular mass of water.

Notes:

- 5 (a) The optical phase function measurements in the CK-EDB used to estimate the size of a levitated droplet are taken at a wavelength of 532 nm; however, the wavelength dependence of refractive index was considered negligible (e.g. for pure water  $m=1.333$  at 589 nm and  $m=1.335$  at 532 nm) and introduced negligible error in the determination of droplet size when compared with other experimental uncertainties that must be accounted for (Rovelli et al., 2016).
- 10 (b) No uncertainties on the applied density parameterisations are taken into account. However, the sensitivity of the hygroscopic properties retrieved by means of CK-EDB measurements to potential uncertainties in the parameterisation of solution density have been evaluated in Section III.d of the main manuscript.

**Table S2:** Measured densities and refractive indices of aminium sulphates solutions.

<b>MMAS</b>						<b>MEAS</b>					
<i>mfs</i>		$\rho$ (kg m <sup>-3</sup> )		<i>m</i>		<i>mfs</i>		$\rho$ (kg m <sup>-3</sup> )		<i>m</i>	
		$\sigma$	$\sigma$	$\sigma$	$\sigma$			$\sigma$	$\sigma$		
0	± 0	998.1	0	1.333	-	0	± 0	998.2	0.1	1.333	-
0.0021	± 8.90E-06	999.0	0.1	1.3334	-	0.0073	± 3.00E-05	1000.6	0.1	1.3342	-
0.0070	± 3.00E-05	1001.4	0.1	1.3342	-	0.0129	± 5.30E-05	1002.8	0.1	1.3350	-
0.0178	± 0.0001	1006.3	0.1	1.3359	-	0.0345	± 0.0001	1010.6	0.1	1.3382	5.77E-05
0.0406	± 0.0002	1016.8	0	1.3394	-	0.0776	± 0.0003	1026.1	0	1.3444	-
0.0675	± 0.0003	1028.8	0.1	1.3436	-	0.1393	± 0.0006	1047.3	0.2	1.3531	-
0.1061	± 0.0005	1045.8	0.2	1.3494	-	0.2087	± 0.0009	1071.1	0.4	1.3632	5.77E-05
0.1487	± 0.0006	1064.8	0.1	1.3561	-	0.3097	± 0.0013	1107.1	0.1	1.3778	-
0.1995	± 0.0009	1087	0	1.3638	-	0.4178	± 0.0017	1144.2	0.2	1.3934	1.15E-04
0.2579	± 0.0011	1111.5	0.4	1.3724	-	0.5295	± 0.0022	1180.6	0.2	1.4091	5.77E-05
0.3002	± 0.0013	1127.6	0.1	1.3778	5.77E-05	0.7687	± 0.0031	1247.6	1.2	1.4413	2.00E-04
0.3657	± 0.0016	1154.7	0.1	1.3872	5.77E-05						
0.4411	± 0.0019	1186.8	1.2	1.3981	-						
0.5482	± 0.0023	1231.3	0.6	1.4145	1.53E-04						

<b>DMAS</b>						<b>DEAS</b>					
<i>mfs</i>		$\rho$ (kg m <sup>-3</sup> )		<i>m</i>		<i>mfs</i>		$\rho$ (kg m <sup>-3</sup> )		<i>m</i>	
		$\sigma$	$\sigma$	$\sigma$	$\sigma$			$\sigma$	$\sigma$		
0	± 0	998.5	0.1	1.333	-	0	± 0	998.1	0.1	1.333	-
0.0108	± 0.0001	1001.8	0.1	1.3345	-	0.0024	± 4.70E-05	998.3	0.1	1.3335	5.77E-05
0.0643	± 0.0005	1019.1	0.1	1.3421	-	0.0124	± 2.50E-04	1000.5	0.1	1.3347	5.77E-05
0.1232	± 0.0010	1037.6	0.1	1.3502	-	0.0291	± 0.0006	1004.1	0.1	1.3371	-
0.1742	± 0.0015	1052.9	0.6	1.3573	-	0.0690	± 0.0014	1012.7	0.1	1.3426	-
0.2351	± 0.0020	1071.9	0.1	1.3657	1.53E-04	0.1126	± 0.0022	1026.7	0.1	1.3518	-
0.2956	± 0.0025	1088.9	1.2	1.3737	2.65E-04	0.2094	± 0.0041	1043.3	0.1	1.3629	1.15E-04
0.3557	± 0.0031	1104.3	4.7	1.3823	2.00E-04	0.3061	± 0.0061	1064.9	0.1	1.3773	5.77E-05
0.4082	± 0.0035	1124.5	0.4	1.3898	5.77E-05	0.4292	± 0.0085	1092.2	0.1	1.3962	1.15E-04
0.4699	± 0.0040	1143.4	0.4	1.3985	5.77E-05	0.5622	± 0.0111	1117.3	1.4	1.4152	9.87E-04
0.5330	± 0.0045	1156.0	1.1	1.4067	1.15E-04	0.6667	± 0.0132	1134.5	0.2	1.4309	4.16E-04

TMAS					TEAS						
<i>mfs</i>	$\rho$ (kg m <sup>-3</sup> )		<i>m</i>		<i>mfs</i>	$\rho$ (kg m <sup>-3</sup> )		<i>m</i>			
	$\pm$	$\sigma$	$\sigma$	$\sigma$		$\pm$	$\sigma$	$\sigma$	$\sigma$		
0	$\pm$ 0	998.0	0.1	1.333	0	$\pm$ 0	998.0	0.2	1.333		
0.0017	$\pm$ 1.70E-05	998.5	0	1.3334	-	0.0025	$\pm$ 3.20E-04	998.5	0.1	1.3336	-
0.0039	$\pm$ 3.70E-05	999.2	0.1	1.3336	-	0.0129	$\pm$ 7.30E-04	1001.3	0.1	1.3354	5.77E-05
0.0109	$\pm$ 0.0001	1001.2	0.1	1.3346	-	0.0335	$\pm$ 0.0012	1006.9	0.2	1.3392	-
0.0197	$\pm$ 0.0002	1003.5	0.1	1.3359	-	0.0637	$\pm$ 0.0016	1014.9	0.2	1.3447	-
0.0331	$\pm$ 0.0003	1007.3	0.1	1.3378	-	0.1236	$\pm$ 0.0023	1030.8	0.1	1.3558	-
0.0515	$\pm$ 0.0005	1012.1	0.2	1.3404	-	0.1884	$\pm$ 0.0028	1048.9	0.1	1.3683	1.53E-04
0.0786	$\pm$ 0.0007	1019.4	0.1	1.3444	-	0.2747	$\pm$ 0.0034	1075.1	0.1	1.3864	-
0.1027	$\pm$ 0.0010	1025.7	0.1	1.3477	5.77E-05	0.3807	$\pm$ 0.0040	1104.8	1.2	1.4082	1.00E-04
0.1303	$\pm$ 0.0012	1032.9	0.1	1.3517	-	0.4999	$\pm$ 0.0046	1135.6	1.6	1.4298	1.18E-03
0.1450	$\pm$ 0.0014	1036.8	0.1	1.3539	5.77E-05	0.6008	$\pm$ 0.0050	1158.2	1.8	1.4480	4.54E-03
0.1828	$\pm$ 0.0017	1046.6	0.2	1.3593	5.77E-05						
0.2104	$\pm$ 0.0020	1053.4	0.1	1.3634	-						
0.2557	$\pm$ 0.0024	1065.3	0.1	1.3701	-						
0.3074	$\pm$ 0.0029	1078.6	0.2	1.3778	5.77E-05						
0.3638	$\pm$ 0.0035	1094.0	0.2	1.3864	-						
0.4580	$\pm$ 0.0044	1118.1	1.6	1.4006	1.53E-04						

Symbols: *mfs* – mass fraction of solute;  $\rho$  – measured density of solution; *m* – measured refractive index of solution;  $\sigma$  – standard deviation arising from the averaging of three repeated measurements.

**Table S3:** 3<sup>rd</sup> order polynomial fitted parameters for densities ( $\rho$ ), calculated melt densities ( $\rho_{melt}$ ) and melt refractive indices ( $m_{melt}$ ) calculated with molar refraction mixing rule for the six considered aminium sulphates.

	$a_0$	$a_1$	$a_2$	$a_3$	$\rho_{melt}$ ( $kg\ m^{-3}$ )	$m_{melt}$
MMAS	998.10	5.12	447.11	-40.19	1410.1	1.4771
DMAS	998.47	-2.55	341.03	-51.91	1285.0	1.4725
TMAS	998.03	4.64	252.17	4.82	1259.7	1.4921
MEAS	998.17	-19.25	459.22	-127.00	1311.1	1.4708
DEAS	998.10	-10.09	285.60	-82.47	1191.1	1.4810
TEAS	998.50	-25.09	378.58	-100.86	1251.1	1.5357

### Parameterisations used:

3<sup>rd</sup> order polynomial parameterization for densities ( $\rho$ ) as a function of the square-rooted mass fraction ( $mfs$ ) of solute:

$$\rho = a_0 + a_1(mfs^{0.5}) + a_2(mfs) + a_3(mfs^{1.5}) \quad \text{Eq. S1}$$

Molar refraction mixing rule for the representation of the refractive index ( $m$ ) of a solution. The molar refraction ( $R$ ) of a species is defined as (Liu and Daum, 2008):

$$R = \frac{(m^2-1)M}{(m^2+2)\rho} \quad \text{Eq. S2}$$

where  $M$  is the compound's molecular mass and  $\rho$  is its pure density (melt density in case of a solute, pure liquid density in the case of the solvent, i.e. water). The molar refraction for a mixture of different components is defined as the sum of the molar refractions of each compound ( $i$ ) weighted by their mole fractions ( $x_i$ ):

$$R = \sum_i x_i R_i \quad \text{Eq. S3}$$

Therefore, in the case of a binary solution composed of water and a generic solute:

$$\frac{(m_{solution}^2-1)M_{solution}}{(m_{solution}^2+2)\rho_{solution}} = x_{H_2O} \frac{(m_{H_2O}^2-1)M_{H_2O}}{(m_{H_2O}^2+2)\rho_{H_2O}} + x_{solute} \frac{(m_{melt}^2-1)M_{solute}}{(m_{melt}^2+2)\rho_{melt}} \quad \text{Eq. S4}$$

where  $m_{solution}$  and  $\rho_{solution}$  are measured,  $M_{solution}$  is calculated as  $M_{solution} = \sum_i x_i M_i$  (Liu and Daum, 2008), the mole fractions of water and solute are known from the concentration of the prepared solution, density and refractive index of pure water are known,  $\rho_{melt}$  is calculated from the density parameterisation at  $mfs=1$ . Therefore, the only unknown is  $m_{melt}$  and a least squares fit of the experimental refractive index data is performed with Eq. S4 to calculate it.

When correcting the obtained radius vs. time data from CK-EDB experiments, the iterative procedure proposed in a previous work (Davies et al., 2012) is applied, but the variation in refracting index with the variation of the solution concentration is calculated with the molar refraction mixing rule.

**Table S4:** Comparison of pure melt densities ( $\rho_{melt}$ ) from this work and from Clegg et al. (2013) and solid densities ( $\rho_{solid}$ ) from Qiu and Zhang (2012).

<b>Compound</b>	<b><math>\rho_{melt}</math> (this work)</b>	<b><math>\rho_{melt}</math></b>	<b><math>\rho_{solid}</math></b>
	(g cm <sup>-3</sup> )	(Clegg et al., 2013) (g cm <sup>-3</sup> )	(Qiu and Zhang, 2012) (g cm <sup>-3</sup> )
MMAS	1.410	1.451 ( $\pm 0.0029$ )	1.485 $\pm$ 0.089
DMAS	1.285	1.278 ( $\pm 0.0022$ )	1.408 $\pm$ 0.084
TMAS	1.259	1.289 ( $\pm 0.0031$ )	1.342 $\pm$ 0.081
MEAS	1.311	-	-
DEAS	1.191	1.212 ( $\pm 0.0030$ )	1.268 $\pm$ 0.076
TEAS	1.251	1.190 ( $\pm 0.0018$ )	1.247 $\pm$ 0.075

**Table S5:** Values of the coefficients of the fitted equations of measured apparent molar volumes ( $V^\phi$ ) of aminium sulphates in aqueous solutions (displayed in Figure S3). The used fitting equation is the same used by Clegg et al. (2013):

$$V^\phi = V^{\phi\infty} + a_1 \cdot (wt\%)^{0.5} + a_2 \cdot (wt\%)^{0.75} + a_3 \cdot (wt\%) + a_4 \cdot (wt\%)^{1.5}$$

where  $V^{\phi\infty}$  is the apparent molar volume of the aminium salt taken from literature (as in Clegg et al.) and  $wt\%$  is the weight percentage of salt in solution. Note that the fitted equation is constrained at  $wt\% = 0$  to the  $V^{\phi\infty}$  literature value.


	$V^{\phi\infty}$	$a_1$	$a_2$	$a_3$	$a_4$
MMAS	86.27	-0.542 ± 0.276	0	0.792 ± 0.099	-0.0507 ± 0.00882
DMAS	123.83	2.039 ± 0.150	0	0	0.00266 ± 0.00367
TMAS	159.47	-8.051 ± 0.962	6.866 ± 0.883	-1.318 ± 0.201	0
MEAS	120.8	-1.043 ± 0.238	1.062 ± 0.088	0	0
DEAS	191.91	0	0	0.115 ± 0.009	0
TEAS	256.16	-22.759 ± 1.264	7.080 ± 0.495	0	0




**Table S6:** Extended ZSR fitted coefficients (Eq. 3 in Sauerwein et al. (2015)) at two fixed water activities ( $a_w$ , 0.80 and 0.925) for mixtures of aminium sulphate salts and sulphuric acid, as in Figure 7 in the main manuscript. The coefficients refer to the ZSR fitting of data from Sauerwein et al. (2016) together with the CK-EDB datapoint (red dashed lines in Figure 7 in the main manuscript). Symbols:  $w^o_a$  – water associated to  $H_2SO_4$  in an aqueous solution at the same water activity as the mixture;  $w^o_{as}$  – water associated to the aminium salt in an aqueous solution at the same water activity as the mixture;  $A^o$  – fitted empirical parameter.

$a_w = 0.80$						
	$w^o_a$	$w^o_{as}$			$A^o$	
MMAS	0.2682	0.2501	±	0.0092	-0.3218	± 0.0447
DMAS	0.2682	0.3072	±	0.0047	-0.4272	± 0.0240
TMAS	0.2682	0.3782	±	0.0112	-0.6147	± 0.0388
MEAS	0.2682	0.2747	±	0.0076	-0.3733	± 0.0256
DEAS	0.2682	0.3299	±	0.0195	-0.4655	± 0.0882
$a_w = 0.925$						
	$w^o_a$	$w^o_{as}$			$A^o$	
MMAS	0.5641	0.6035	±	0.0194	-0.6155	± 0.0934
DMAS	0.5641	0.6997	±	0.0248	-0.7095	± 0.1273
TMAS	0.5641	0.8143	±	0.0215	-1.0166	± 0.0749
MEAS	0.5641	0.6587	±	0.0140	-0.6166	± 0.0476
DEAS	0.5641	0.8472	±	0.0325	-1.0603	± 0.1233


**Figure S1:** Measured densities of aminium sulphates solutions and 3<sup>rd</sup> order polynomial fittings of data (Panel A). Measured refractive indices as a function of mass fraction and the corresponding molar refraction mixing rule fitting (Panel B).



**Figure S2:** Comparison of measured  $GF_r$  from CK-EDB experiments and  $GF_r$  data from H-TDMA measurements by Qiu and Zhang (2012).



**Figure S3:** Apparent molar volumes ( $V^\phi$ ) of aminium sulphates in aqueous solutions as a function of the square root of the solute weight percentage ( $wt\%$ ). Symbols: black diamonds – measured values, this work; black line – fitted equations (coefficients in Table S5); grey dots – measured values by Clegg et al. (2013); grey lines – fitted equations of measured data in Clegg et al. (2013); dashed lines – uncertainty on apparent molar volumes when a  $\pm 0.001 \text{ g cm}^{-3}$  uncertainty on the measured density is considered; open circles – apparent molar volumes at infinite dilutions from literature (as in Fig. 1 in Clegg et al. (2013)).



## References

- Clegg, S. L., Qiu, C. and Zhang, R.: The deliquescence behaviour, solubilities, and densities of aqueous solutions of five methyl- and ethyl-aminium sulphate salts, *Atmos. Environ.*, 73, 145–158, doi:10.1016/j.atmosenv.2013.02.036, 2013.
- 5 Davies, J. F.: *Characterising The Impact Of Bulk, Surface And Gas-Phase Limitations On Mass Transport In Aerosol.*, 2014.
- Davies, J. F., Haddrell, A. E. and Reid, J. P.: Time-Resolved Measurements of the Evaporation of Volatile Components from Single Aerosol Droplets, *Aerosol Sci. Technol.*, 46(6), 666–677, doi:10.1080/02786826.2011.652750, 2012.
- Davies, J. F., Haddrell, A. E., Rickards, A. M. J. and Reid, J. P.: Simultaneous analysis of the equilibrium hygroscopicity and water transport kinetics of liquid aerosol, *Anal. Chem.*, 85(12), 5819–5826, doi:10.1021/ac4005502, 2013.
- 10 Liu, Y. and Daum, P. H.: Relationship of refractive index to mass density and self-consistency of mixing rules for multicomponent mixtures like ambient aerosols, *J. Aerosol Sci.*, 39(11), 974–986, doi:10.1016/j.jaerosci.2008.06.006, 2008.
- Qiu, C. and Zhang, R.: Physiochemical properties of alkylaminium sulfates: Hygroscopicity, thermostability, and density, *Environ. Sci. Technol.*, 46(8), 4474–4480, doi:10.1021/es3004377, 2012.
- 15 Rovelli, G., Miles, R. E. H., Reid, J. P. and Clegg, S. L.: Accurate Measurements of Aerosol Hygroscopic Growth Over a Wide Range in Relative Humidity, *J. Phys. Chem. A*, 120, 4376–4388, doi:10.1021/acs.jpca.6b04194, 2016.
- Sauerwein, M., Clegg, S. L. and Chan, C. K.: Water Activities and Osmotic Coefficients of Aqueous Solutions of Five Alkylaminium Sulfates and Their Mixtures with H<sub>2</sub>SO<sub>4</sub> at 25°C, *Aerosol Sci. Technol.*, 49(8), 566–579, doi:10.1080/02786826.2015.1043045, 2015.