

Interactive comment on “NO₂ pollution over India observed from space – the impact of rapid economic growth, and a recent decline” by Andreas Hilboll et al.

Anonymous Referee #3

Received and published: 5 April 2017

Hilboll et al. presents an analysis of NO₂ pollution changes over different regions of India and their socioeconomic drivers, by combining multiple satellite products and official socioeconomic data. I have a few suggestions as follows.

Multiple satellite products are used. Although some consistency is found in trend results (Fig. 1 and Fig. S1), there are clear quantitative differences among these products, especially after 2012. For example, the trend of OMI NO₂ is clearly different from those in GOME2-a and GOME2-b (for all regions in Fig. 1 except North Indian Plain). For North Indian Plain, the OMI trend (Fig.1) is different from the DOMINO NO₂ trend in Fig. S1. The large uncertainties in these satellite datasets make it difficult to conduct further linkage to socioeconomics. Is it possible to focus the analysis on regions

C1

that multiple satellite products show quantitatively consistent trends? What are the uncertainties in trends from individual products considering retrieval and representative errors?

The NO₂ growth rates are quantitatively significantly different from those in socioeconomic data, often by a factor of 10 (Table 1). It appears that chemistry, meteorology and/or other factors play major roles here. Can these factors be better accounted for in linking NO₂ trends to emission trends? How are the roles of chemistry/meteorology in NO₂ trends over India compared to the roles over other countries?

A statistical model is used to calculate the NO₂ trends. Please discuss the model here briefly. Also, the model does not account for shift in seasonality when the pollution grew, which is important for fast changing pollution regions. Please discuss the caveat of this model.

Many regions are discussed here. A map defining all these regions will be extremely helpful for a general reader to understand the analyses.

The explanations in the last three paragraphs of Sect. 3.1 could be further improved. North Indian Plain also have large emissions from non-traffic sources such as power generation and industry. Can other factors be ruled out? The explanation for Chhattisgarh, Jharkand and Odisha is focused on power generation, how about other factors? Also, it is not clear why and how the monsoon signal is clear for South India but not for other regions. Can the changes over these regions be also found in OMI NO₂ data? Overall, a region-specific analysis of major socioeconomic factors before discussing the causes of NO₂ trends in these regions will much help the causation analysis.

The OMI NO₂ data should be analyzed more intensively (e.g., in Figs 3,5,6 and Table 1), given its long temporal coverage (2004-present), different time of day (to help discuss the role of chemistry), and a higher resolution (to help reveal the hotspots). Comparing OMI with morning-time instruments will also help reveal the satellite uncertainty.

C2

Fig.1 and S1 – starting the y-axis from a higher value (e.g., 15) than zero will help visualize the NO₂ changes.

Fig. 3 – can you show results from OMI and quantitatively compare to GOME-2 results?

Fig. 6 – NO₂ was flat (or even declined) from 2008 to 2011 while electricity and GSDP grew clearly. Why? How about the OMI NO₂ data?

Table 1 – how about the NO₂ trends from OMI?

Sect. 3.4 – where are the numbers ($3e14$ – $24e14$ molec cm⁻²) from?

Conclusion – “This may imply that changes in meteorology or up to now not understood changes in tropospheric chemistry are also of significance.” – given the uncertainty (especially after 2012), similar sentences addressing the roles of non-emission factors should be highlighted in the abstract.

Interactive comment on Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2017-101, 2017.