

## Which processes drive observed variations of HCHO columns over India?

Luke Surl<sup>1,2</sup>, Paul I. Palmer<sup>1,2</sup>, and Gonzalo González Abad<sup>3</sup>

<sup>1</sup>National Centre for Earth Observation, University of Edinburgh, Edinburgh, UK.

<sup>2</sup>School of GeoSciences, University of Edinburgh, Edinburgh, UK.

<sup>3</sup>Atomic and Molecular Physics Division, Harvard–Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA

*Correspondence to:* Paul I. Palmer (paul.palmer@ed.ac.uk)

**Abstract.** We interpret HCHO column variations observed by the Ozone Monitoring Instrument (OMI), aboard the NASA Aura satellite, over India during 2014 using the GEOS-Chem atmospheric chemistry and transport model. We use a nested version of the model with a horizontal resolution of approximately 25 km. HCHO columns are related to local emissions of volatile organic compounds (VOCs) with a spatial smearing that increases with the VOC lifetime. Over India, HCHO has biogenic, pyrogenic, and anthropogenic VOC sources. Using a 0-D photochemistry model, we find that isoprene has the largest molar yield of HCHO that is typically realized within a few hours. We find that forested regions that neighbour major urban conurbations are exposed to high levels of nitrogen oxides. This results in depleted hydroxyl radical concentrations and a delay in the production of HCHO from isoprene oxidation. We find that propene is the only anthropogenic VOC emitted in major Indian cities that produces HCHO at a comparable (but slower) rate to isoprene. The GEOS-Chem model reproduces the broad-scale annual mean HCHO column distribution observed by OMI ( $r=0.6$ ), which is dominated by a distinctive meridional gradient in the northern half of the country, and by localized regions of high columns that coincide with forests. Major discrepancies are over the Indo-Gangetic Plain and Delhi. We find that the model has more skill at reproducing observations during winter (JF) and pre-monsoon (MAM) months with Pearson correlations  $r>0.5$  but with a positive model bias of  $\simeq 1 \times 10^{15}$  molec/cm<sup>2</sup>. During the monsoon season (JJAS) we reproduce only a diffuse version of the observed meridional gradient ( $r=0.4$ ). We find that on a continental scale most of the HCHO column seasonal cycle is explained by monthly variations in surface temperature ( $r=0.9$ ), suggesting a role for biogenic VOCs, in agreement with the 0-D and GEOS-Chem model calculations. We also find that the seasonal cycle during 2014 is not significantly different from the 2008–2015 mean seasonal variation. There are two main loci for biomass burning (states of Punjab

and Haryana, and northeastern India), which we find contributes a significant contribution (up to  $1 \times 10^{15}$  molec/cm<sup>2</sup>) to observed HCHO columns only during March and April over northeastern India. The slow production of HCHO from propene oxidation results in a smeared hotspot over Delhi that we resolve only on an annual mean timescale by using a temporal oversampling method. Using a linear regression model to relate GEOS-Chem isoprene emissions to HCHO columns we infer seasonal isoprene emissions over two key forest regions from the OMI HCHO column data. We find that the *a posteriori* emissions are typically lower than the *a priori* emissions, with a much stronger reduction of emissions during the monsoon season. We find that this reduction in emissions during monsoon months coincides with a large drop in satellite observations of leaf phenology that recovers in post monsoon months. This may signal a forest-scale response to monsoon conditions.

## 1 Introduction

Formaldehyde (HCHO) is an important source of the hydroperoxyl radical (Volkamer et al., 2010; Whalley et al., 2010), and a source of upper tropospheric hydroxyl radical (OH) (Jaeglé et al., 1998). It therefore plays a role in determining the oxidizing capacity of the global troposphere. The principal source of HCHO is from the oxidation of methane (CH<sub>4</sub>), which provides a global ambient background. Shorter-lived non-methane volatile organic compounds (NMVOCs) elevate HCHO concentrations over continental atmospheres. Minor direct HCHO sources include biomass burning, industry, agriculture, automobiles, shipping, and vegetation. The atmospheric lifetime of HCHO, determined by OH and photolysis, is typically several hours. Building on our past studies (Palmer et al., 2003, 2006; Barkley et al., 2008; Gonzi et al., 2011), we interpret HCHO column distributions over India observed by the Ozone Monitoring Instrument (OMI) aboard the NASA Aura spacecraft (Levelt et al., 2006). We interpret spatial and temporal variations in terms of biogenic, pyrogenic, and anthropogenic VOC sources.

India has the sixth largest economy and the second largest population of any country. It also has one of the largest increases in mortality rates due to chronic exposure to elevated levels of surface ozone and particulate matter (Cohen et al., 2017). Figure 1 shows that India has a rich landscape that includes the Thar desert over northwestern India, major forests over the southwestern coast and over the east and northeast, and five megacities (New Delhi, Mumbai, Kolkata, Bengalura, and Chennai). The Indo-Gangetic Plain (IGP) stretches from eastern Pakistan, across the northern edge of India (bounded by the Himalayas), to Bangladesh. The IGP represents more than a quarter of a million acres of fertile land, which is used primarily to grow rice and wheat, but also maize, sugarcane, and cotton. The southwest monsoon represents the main source of water to the IGP, with contributions also from rivers flowing from the Himalayas. The southwest monsoon begins in June and subsides in September. High temperatures over the Thar desert cause a region of low pressure that helps to establish a large-scale land-sea breeze with the Indian Ocean. This results in warm, moisture-

laden air from the Indian Ocean travelling inland. Eventually, this air meets the Himalayas where it is forced to rise. As the air rises, cooler temperatures result in precipitation. Some areas of India  
60 receive ten metres of rain annually, mostly during the monsoon season.

Satellite columns observations of HCHO were originally developed using observed UV spectra from the Global Ozone Monitoring Experiment (Thomas et al., 1998; Chance et al., 2000). HCHO column data are now available from a range of satellite instruments, but here we focus on data from OMI aboard the NASA Aura spacecraft. Generally, slant HCHO columns are retrieved by directly  
65 fitting to observed spectra in a narrow UV window (Chance et al., 2000; De Smedt et al., 2008; González Abad et al., 2015). Vertical columns are determined by scaling these slant columns by scene-dependent air-mass factors (AMFs), taking into account clouds and aerosol scattering (Palmer et al., 2001). Past work has shown that HCHO columns over India have increased nationwide on average between 1.6%/yr (1997–2009, De Smedt et al. (2010)) and 1.5%/yr (1995–2013, Mahajan  
70 et al. (2015)). Mahajan et al. (2015) also showed using coincident satellite measurements of HCHO and NO<sub>2</sub> that over much of India O<sub>3</sub> production is limited by the availability by nitrogen oxides but over urban regions it is limited by the availability of VOCs, supported by detailed modelling studies over Delhi (Sharma et al., 2016). Here, we employ a high-resolution ( $\simeq 25$  km) model of atmospheric chemistry that is closely aligned with the resolution of the satellite data, allowing us to  
75 take advantage of the richness of these data.

HCHO columns are related to their parent VOC emissions with a smearing spatial scale that is related to the production rate and molar yield of HCHO (Palmer et al., 2003). Past studies have used this relationship to infer isoprene emissions from major forests (Palmer et al., 2003; Abbot et al., 2003; Shim et al., 2005; Palmer et al., 2006, 2007; Barkley et al., 2008; Millet et al., 2008;  
80 Stavrakou et al., 2009; Curci et al., 2010; Marais et al., 2012; Barkley et al., 2013), biomass burning emissions (Young and Paton-Walsh, 2010; Gonzi et al., 2011; Stavrakou et al., 2016), and anthropogenic emissions (Fu et al., 2007; Stavrakou et al., 2009). Detailed photochemical calculations that link VOCs and the time-dependent production of HCHO lay the groundwork for interpreting the HCHO column data. Many past studies have inferred VOC emissions from the HCHO columns using  
85 a linear regression model between these two variables (e.g. Palmer et al. (2003); Millet et al. (2008)), but others have adopted a more rigorous Bayesian inverse model approach (e.g. Shim et al. (2005)). For central Africa, Marais et al. (2012) employed an inversion method that accounted for the NO<sub>x</sub> dependence of the isoprene-HCHO relationship. Given the large uncertainties associated with VOC emissions (e.g. Guenther et al. (2012)) and the production of HCHO in the low-NO<sub>x</sub>  
90 regime (Wolfe et al., 2016) both approaches provide useful insights. Our study is focused on India where there are significant sources of biogenic, pyrogenic, and anthropogenic VOCs.

The next section describes the OMI HCHO column data, the detailed box-model used to study the time-dependent production of HCHO from VOC oxidation, and the GEOS-Chem atmospheric chemistry transport model focused on India. Section 3 reports the results from our analysis of the

95 OMI HCHO column data over India, the associated model interpretation of these data, and the iso-  
prene emissions we infer from the HCHO column data collected over two major forest regions. We  
conclude in Section 4.

## 2 Data and Methods

Our data are focused on India, as defined by the database of Global Administrative Areas ([www.gadm.org](http://www.gadm.org)). We adopt climatological definitions of seasons from the Indian Meteorological Department that are determined by the onset of the regional monsoon system: winter includes January and February; the pre-monsoon season is from March to May; the monsoon season is from June to September; and the post-monsoon season is from October to December. Our focus is on OMI HCHO column variations during 2014, but we put this year into a longer temporal context by comparing it  
105 with data collected from 2008 to 2015.

### 2.1 Ozone Monitoring Instrument HCHO Columns

For our analysis we use HCHO vertical columns from OMI, a nadir-viewing UV/Vis spectrometer aboard the NASA Aura satellite that was launched in 2004. Aura is in a sun-synchronous orbit with a local equatorial crossing time of 13:38, achieving daily global coverage subject to cloud coverage.

110 OMI uses two imaging grating spectrometers each with a CCD detector to collect solar backscattered radiation in the spectral range 270–500 nm using three channels: UV-1 (264–311 nm), UV-2 (307–383 nm), and Vis (349–504 nm). OMI has an across-track swath width of 2,600 km, which in global mode is described by 60 scenes that have ground footprints from  $13 \times 24 \text{ km}^2$  at nadir to  $28 \times 160 \text{ km}^2$  at the swath edges.

115 Determination of HCHO vertical columns is a two-step procedure, which we describe below. First, slant columns are retrieved from the observed spectra. Second, an air-mass factor (AMF) is used to transform these slant columns into geophysical vertical columns that can be compared more easily with models.

We use the NASA OMHCHOv003 data product (González Abad et al., 2015) from the NASA  
120 Data and Information Services Center, which fits HCHO slant columns in the 328.5–356.5 nm window and accounts for competing absorbers, the Ring effect, and undersampling. Typical slant columns range from  $4 \times 10^{15}$  to  $6 \times 10^{16} \text{ molec/cm}^2$  with associated fitting uncertainties ranging from 30% for larger columns to more than 100% for smaller columns (González Abad et al., 2015).

We adopt a conservative approach to filtering data, based on previous studies (e.g. De Smedt et al. (2015)). We filter slant column data using the main data quality flag (removing suspect, bad or missing scenes); scenes that have been flagged as being affected by the row anomaly corresponding to problem with a row of CCD detectors on OMI; scenes with slant column values greater than  $15 \times 10^{16} \text{ molec/cm}^2$  that we attribute also to potential row anomalies; scenes with solar zenith angles

less than  $0^\circ$  or greater than  $70^\circ$ ; scenes with cloud fractions  $> 0.4$ ; and scenes from the last two rows  
130 at the edge of the swath, which we are too wide to be useful in our analysis. We do not anticipate that  
these data filters will introduce bias in our analysis, with the exception of removing cloudy scenes that  
introduces a clear-sky bias. To evaluate the clear-sky bias, we compared the corresponding model  
HCHO columns (described below) with and without these cloudy scenes and find that removing  
them results in a monthly mean positive bias of 13%, consistent with previous work (Palmer et al.,  
135 2001).

To transform each observed slant column to a vertical column we calculate an AMF that accounts  
for temporal and spatial variations from scattering due to clouds and aerosols, and for the vertical  
distribution of HCHO. This approach is described in detail by Palmer et al. (2001); Martin et al.  
(2003), and has since been evaluated in a large number of studies (e.g., Palmer et al. (2003, 2006);  
140 Fu et al. (2007); Millet et al. (2008); Barkley et al. (2008); Curci et al. (2010); Gonzi et al. (2011);  
Marais et al. (2012)). We use the OMI OMCL02 cloud data product, and the nested GEOS-Chem  
model to provide information about aerosols and HCHO vertical distributions, described below.

Finally, as a post-processing step, we remove any large-scale biases using a reference sector nor-  
malization procedure (Palmer et al., 2003; Martin et al., 2003). Over the remote Pacific we expect  
145 observed HCHO columns to be determined exclusively by the oxidation of methane. By anchoring  
observed values over the remote Pacific ( $0\text{--}40^\circ\text{N}$ ,  $160\text{--}180^\circ\text{W}$ ) to the corresponding GEOS-Chem  
model values we determine the monthly bias that we subsequently subtract from the data within the  
study domain.

Previous work using aircraft observations over Southeastern U.S. has reported that the OMH-  
150 CHOv0003 OMI HCHO column data product has a negative bias (37%), and that the GEOS-Chem  
model (v9-02) has a 10% negative bias (Zhu et al., 2016). In the absence of similar data over In-  
dia, we cannot determine with certainty the generality of these biases, and have chosen to implicitly  
assume that the OMI data are not significantly impacted by systematic error

## 2.2 Models of Tropospheric Chemistry

155 We use the CAABA box model, including a comprehensive description of atmospheric chemistry,  
to understand the time-dependent yield of HCHO from the oxidation of VOCs in Indian forest and  
urban environments. These calculations help determine which VOC emissions are responsible for  
observed HCHO column variations. We also use a nested version of the GEOS-Chem 3-D atmo-  
spheric chemistry model to interpret the satellite data and to understand chemistry on a regional  
160 spatial scale. We use this model to establish a relationship between emissions of VOCs and HCHO  
columns that is used to infer emissions of VOCs that correspond to observed HCHO columns.

The chemical mechanisms from the CAABA box model and the GEOS-Chem model are not  
identical. CAABA includes a more explicit chemical mechanism and a reduced version of the Mainz  
Isoprene Mechanism 2 (MIM2, Taraborrelli et al. (2009)). The version of GEOS-Chem we use here

165 is described by Eastham et al. (2014). The latest version of the GEOS-Chem chemical mechanism,  
released after the majority of our calculations had been completed, results in an increase of the  
HCHO yield from the oxidation of isoprene. We find using preliminary calculations with v11-01  
that the revised mechanism does not systematically change the results shown here. Previous studies  
have evaluated the performance of GEOS-Chem (Marvin et al., 2017) and MIM2 (Taraborrelli et al.,  
170 2009) against the Master Chemical Mechanism. The GEOS-Chem v10-01 mechanism slightly un-  
derestimates HCHO production in high-NO<sub>x</sub> conditions and the MIM2 mechanism shows similar  
HCHO production across a wide range of NO<sub>x</sub> values.

### Box modelling

We use v3.0 of the CAABA/MECCA (0-D) box model (Sander et al., 2011), but without halo-  
175 gen, sulphur, and mercury chemistry, to estimate the time-dependent production of HCHO from the  
chemical oxidation of different VOCs in forest and urban photochemical environments (Table 1).

We set up the model to describe a well-mixed summertime boundary layer, with photochemistry  
driven by a diurnal cycle in sunlight. For each environment, the photolysis model JVAL (Sander  
et al., 2014) is used to calculate photolysis rates assuming clear-sky conditions given latitude, lon-  
180 gitude, and day of year. For each study location, we assume constant values for humidity, pressure,  
temperature and boundary layer height, taken from colocated GEOS-FP meteorology (see below).  
For forested environments we also prescribe fixed mixing ratios of O<sub>3</sub>, NO<sub>2</sub> and CO from the GEOS-  
Chem model, described below. For the Delhi urban environment we use daytime average surface air  
pollutant values in 2014 reported by Tyagi et al. (2016): O<sub>3</sub> (37 ppbv), CO (2.3 ppmv), and NO<sub>2</sub>  
185 (19 ppbv), Table 1. Figure 1 shows the three forested regions (denoted by E, NE, and S) we chose to  
explore the range of photochemical environments. For each calculation, we spun-up the model from  
initial conditions for 48 hours before running the model for a further seven days. We use a model  
timestep of two minutes.

To evaluate the time-dependent HCHO yield of anthropogenic and biogenic VOCs we run paired  
190 calculations: 1) a control run and 2) a run in which we perturb a VOC by approximately  $5 \times$   
 $10^{14}$  molec cm<sup>-2</sup> over a 15 minute period at 0900 local time on the first day after the two-day spin-up  
period. The perturbed amount is sufficiently small that we can assume that the chemistry response is  
approximately linear so that we can compare the perturbed run with the control run. We can then use  
control minus perturbed model calculations to determine HCHO per-carbon yield from the oxidation  
195 of the perturbed VOC.

### GEOS-Chem 3-D Modelling

We use v10-01 of the GEOS-Chem global model of atmospheric chemistry and transport ([www.geos-chem.org](http://www.geos-chem.org)), driven by GEOS-FP analyzed meteorological fields, provided by the Global Mod-  
eling and Assimilation Office (GMAO) at NASA Goddard Space Flight Center. The native spatial

200 resolution of these data is  $0.25^\circ$  (latitude)  $\times$   $0.3125^\circ$  (longitude), and includes 47 vertical terrain-  
following sigma-levels that describe the atmosphere from the surface to 0.01 hPa of which about  
30 are typically below the dynamic tropopause. The 3-D meteorological data are updated hourly,  
and 2-D fields and surface fields are updated every 3 hours. The chemical mechanism used is the  
“tropchem” mechanism (Eastham et al., 2014). The atmospheric chemistry and transport timesteps  
205 are five and ten minutes, respectively.

We use the nesting capability of the model to focus on India, defined here as  $0\text{--}40^\circ\text{N}$ ,  $65\text{--}100^\circ\text{E}$ ,  
using the native resolution of the meteorological data. We use time-dependent lateral boundary con-  
ditions archived from a self-consistent,  $4^\circ \times 5^\circ$  version of the global full-chemistry model, which is  
initialized in January 2013 to minimize the influence of initial conditions. Here, we describe emis-  
210 sion inventories relevant to our nested India simulation and direct the reader to the GEOS-Chem  
website (<http://geos-chem.org>) for a more comprehensive description of inventories.

Monthly anthropogenic emissions for India are taken from the mosaic MIX inventory (Li et al.,  
2017), including NO, CO, SO<sub>2</sub> and NMVOCs that are available on a spatial resolution of  $0.25 \times$   
 $0.25^\circ$ . The emissions for India are a mosaic of regional emission inventories, including the Regional  
215 Emission inventory in ASia (REAS) (Kurokawa et al., 2013) and those developed by the Argonne  
National Laboratory (Lu et al., 2011; Lu and Streets, 2012). Indian NMVOC emissions represent  
by mass 17 Tg per year, of which 43% is from the residential sector, 36% is from transport, 20% is  
from industry, and a small amount is attributed to the power sector.

Approximately 20% of India’s geographical area is described as forest. To describe biogenic VOC  
220 emissions of isoprene, monoterpenes, alkenes, and acetone we use the MEGAN v2.1 emission model  
(Model of Emissions of Gases and Aerosols from Nature, Guenther et al. (2012)). Figure 2 shows  
seasonal isoprene emissions over India. Monoterpenes emission have a similar distribution to iso-  
prene but are a factor of five smaller in magnitude. Given their relatively small influence on HCHO  
column variability (Palmer et al., 2003) we do not discuss them further. Within GEOS-Chem, we  
225 adjust the MEGAN model with time-dependent local environment conditions (e.g., surface temper-  
ature, photosynthetic active radiation, and leaf area index (LAI) from MODIS).

Monthly pyrogenic emissions are taken from the Global Fire Emissions Database (GFED) version  
4 (van der Werf et al., 2017), which are available with a spatial resolution of  $0.25 \times 0.25^\circ$ .

To interpret OMI HCHO column data we sample the model at the local time averaged between  
230 1300 and 1500, corresponding to the overpass time of OMI, and the location of each observed scene.  
To put 2014 into a broader temporal context, we interpreted OMI HCHO columns for years 2008–  
2015. We find that the AMF plays only a minor role in determining OMI vertical column variations  
so we do not expect our approach to significantly influence our study of year to changes in the HCHO  
columns.

### 235 3 Results

First, we report results from the box-model calculation to provide some insights into the VOCs that determine the observed HCHO column variations. We then report the annual and seasonal spatial distributions of OMI and GEOS-Chem HCHO columns. Using correlative space-borne data we explore the role of biogenic, pyrogenic, and anthropogenic emissions in determining the spatial distributions of HCHO. Finally, using a model relationship between isoprene emissions and HCHO columns we infer isoprene emissions that are consistent with the OMI vertical columns.

#### 3.1 HCHO Yield from VOC Oxidation in Urban and Forest Environments

We use the CAABA/MECCA box model to determine the time-dependent production of HCHO from the VOCs that we expect to be emitted from urban and forest environments over India. This represents important groundwork for interpreting the satellite observations of HCHO. Table 1 provides an overview of the box model calculations we describe below.

Relating HCHO column variations to emissions of its parent VOC requires that the VOC 1) has a high yield of HCHO so that eventual concentrations are elevated above the global background, determined mainly by the oxidation of CH<sub>4</sub>; and 2) produces HCHO rapidly so that most of the HCHO is produced close to the emission source and not smeared over long spatial scales. Isoprene is the dominant HCHO precursor over many northern midlatitude and tropical forest ecosystems (Palmer et al., 2003, 2006; Barkley et al., 2008; Curci et al., 2010). Over tropical latitudes biomass burning emissions of VOCs also play a role in HCHO column variations (Fu et al., 2007; Barkley et al., 2008; Gonzi et al., 2011; Marais et al., 2012). Fu et al. (2007) also showed that reactive anthropogenic VOCs played a role in HCHO column variations over China.

#### Biogenic VOCs from Forest Environments

We explore three contrasting forest regions throughout 2014 (Figure 1), characterized by latitude-dependent levels of photosynthetically active radiation (PAR) and by their proximity to urban emissions. We focus on the yield of HCHO from the emission of isoprene (Palmer et al., 2003).

For each calculation we report in Table 2 the resulting lifetime of the injected isoprene (section 2.2), and the cumulative per-C HCHO yield. We find that the duration of the calculation is sufficiently long that the peak HCHO had time to diminish to a negligible amount.

The atmospheric lifetime of isoprene against OH is typically much shorter than an hour, with shortest values (<15 mins) during summer months. The most southern site in the state of Kerala (Figure 1) is where isoprene has the longest lifetime. This is due to OH being suppressed by high ambient NO<sub>2</sub> concentrations (in excess of 8 ppbv) originating from the coastal conurbation in that state. We find that levels of NO<sub>x</sub> are sufficiently high (averages generally around or above 1 ppbv) in all forested regions such that isoprene peroxy radicals preferentially react with NO to rapidly

produce HCHO. Peak values for HCHO production are typically reached within 90 minutes of the  
270 isoprene oxidation. The OH suppression in Kerala is primarily responsible for the slow production  
of HCHO. We find that the peak HCHO signal is typically reached within 30-90 minutes, corre-  
sponding to a smearing length scale of 9–27 km assuming an example wind speed of 5 m/s. For  
the Kerala region where the time taken to reach the peak HCHO signal is typically 2–3 hours the  
corresponding smearing length scale is 36–54 km. Even in Kerala the smearing length is compara-  
275 ble to the longitudinal extent of a single OMI scene and well within the swath width (section 2).  
Figure 3 shows that the corresponding values of the per-C HCHO yield ranges from 0.38 to 0.66.  
Higher values are generally associated with higher values of  $\text{NO}_x$  (c.f.  $\text{NO}_2$  mixing ratios in Table 1  
and yields reported in Table 2), consistent with previous studies (Palmer et al., 2006; Barkley et al.,  
2013). The simulations reported in this work are specific to the Indian scenarios discussed.

### 280 **Anthropogenic VOC in urban environment**

To study the role of anthropogenic non-methane VOCs (NMVOCs) in determining HCHO columns  
we simulate boundary layer chemistry over Delhi using emissions from the MIX emission inventory  
(Li et al., 2017). Over major cities, NMVOC emissions originate mainly from stationary combus-  
tion and the transport sector. The species in the MIX inventory are defined following the SAPRC-  
285 99 chemical mechanism (Carter, 2000). From a comparison of the correlation coefficients we find  
that over Delhi, according to the MIX inventory, NMVOC emissions include propane, propene and  
ALK4, representing 13%, 32%, and 55% of NMVOC C emissions. ALK4 denotes  $>\text{C}_4$  alkanes.  
The longest chain alkane in the 0-D chemical mechanism is n-butane. We denote this urban model  
scenario as “D05” (Table 1). We calculate individual HCHO yields from the oxidation of propane,  
290 propene, and n-butane.

Propene ( $\text{CH}_3\text{CH}=\text{CH}_2$ ) has an atmospheric lifetime of several hours, determined by OH addition  
to its double bond. The major intermediate products include HCHO and acetaldehyde. The ultimate  
HCHO yield from the oxidation of propene (close to 0.5) is comparable with the value from isoprene  
oxidation (Figure 3) but it only reaches 50% of that final value within 12 hours, consistent with the  
295 longer lifetime of propene. Propane and n-butane have atmospheric lifetimes of 10 and 5 days,  
respectively, determined by OH. Long-lived intermediate oxidation products include acetone that  
further delay the production of HCHO (Figure 3).

Assuming an average wind speed of 3 m/s, taken from wind measurements at Indira Gandhi airport  
in Delhi, only HCHO production from the oxidation of propene (out of all the gases that represent a  
300 major contribution to the regional emission inventory) would produce a signal that could potentially  
be distinguishable above the ambient concentration. We show later there is some evidence that we  
can observe a Delhi HCHO hotspot but only after temporally oversampling the data.

## 3.2 OMI and GEOS-Chem Model HCHO Column Distributions

### Data Filtering and AMF Statistics

305 Using the data quality criteria, as defined above, we remove 58% of all OMI HCHO measurements collected during 2014. The proportion of data removed per month varies during the year, with most scenes removed during the cloudy monsoon season in July (71%) and the least number of scenes removed during March (45%).

Figure 4 shows the annual mean distribution of AMFs across India. AMF values range from 0.7 to 310 1.5 with a median value of 1. AMF values are highest in the Himalayas at the extreme north of India, which are often covered by snow, and over the Rann of Kutch salt marsh. In both cases, elevated AMF values are due to high surface albedos.

To determine the influence of the AMFs on the spatial distribution of HCHO vertical columns we compare fitted OMI slant HCHO columns and the corresponding vertical columns to the GEOS-315 Chem model. We find that the vertical columns reproduce 11% more of the model spatial distribution than the observed slant columns, consistent with past studies (Palmer et al., 2001; Millet et al., 2008) that show that the AMF plays only a minor role in the observed spatial distribution of vertical HCHO columns.

To investigate the role of model resolution in our calculation of vertical columns we repeated the 320 annual mean analysis using the  $2^\circ \times 2.5^\circ$  version of the GEOS-Chem driven by the same inventories as described above. Figure 5 shows that the coarser resolution model fails to reproduce smaller-scale variations, as expected, that define much of the India western coast, and misses variations over northeastern India. While the model via the AMF calculation only contributes a small amount to the distribution of HCHO vertical columns it does affect the magnitude of the columns. We find that 325 there is an overwhelming argument to justify the use of the higher-resolution model.

### Annual Mean Spatial Distribution

Figure 5 shows the annual mean model and observed HCHO columns for 2014. The lowest observed and model columns are in the north of the study region coinciding with the Himalayas and (to a lesser extent) in the Thar desert in the Northwest. The observations show three main regions associated 330 with elevated HCHO columns: 1) the non-Himalayan part of the Northeast region of India, east of Bangladesh; 2) territory at the south of the country, around the southern part of the Western Ghats mountain range, roughly following the borders of the state of Kerala; and 3) a broad area over the east of the country, roughly outlined by the states of Chattisgarh, Odisha, and the northern part of Anhdra Pradesh.

335 The nested GEOS-Chem model reproduces the observed magnitude and broad-scale spatial distribution of OMI HCHO columns over India (Figure 5), but is much smoother. The model has a small but positive bias of 14% (12%) for the mean (median) annual column amounts. We find that

the model captures 33% ( $r=0.58$ ) of the observed annual mean spatial variation of HCHO columns. The major discrepancy between the model and observed annual mean HCHO distributions is over the IGP and over Delhi. Observed HCHO columns are elevated over the IGP but not to the values shown in the model, which we discuss below. A HCHO hotspot over Delhi is apparently absent from the OMI columns, but as we show later there is evidence that the hotspot exists in the observations.

Based on the box modelling, the largest source of HCHO column variability is expected to be isoprene emissions. Figure 1 shows that the broad-scale annual distribution of HCHO over India is consistent with the annual mean distribution of Normalized Diffusive Vegetation Index (NDVI) with a Pearson correlation of  $r = 0.49$ . Below, we use seasonal variations of HCHO columns to improve understanding of the drivers.

### 3.3 Seasonal Spatial Distributions

Figure 6 shows OMI and GEOS-Chem model HCHO columns for seasons during 2014. There are distinct seasonal cycles to the HCHO columns over three broad regions: northeastern and southwestern India, and over the IGP. Over (north)eastern India HCHO columns peak in pre-monsoon and monsoon months. Columns over southwestern coast of India peak in winter and pre-monsoon months and are very low in other seasons. Columns over the IGP are elevated above those elsewhere in northern Indian during pre-monsoon months and peak in monsoon months. Figure 7 shows that the timing of these elevated columns coincide with warmer surface temperatures. This suggests a role for biogenic VOC emissions, consistent with the strong, local relationship between isoprene emissions and HCHO production that we demonstrated above.

The GEOS-Chem model captures these broad-scale observed distributions of HCHO but is much smoother, as expected. We find the model has some skill at reproducing the observed spatial distribution with a Pearson correlation  $r$  of  $>0.4$ , with larger values in winter and pre-monsoon months ( $r=0.50$  and  $0.62$ , respectively) and smaller values in monsoon and post-monsoon months ( $r=0.39$  and  $0.44$ , respectively). On an annual timescale the model has a positive bias of  $1.0 \times 10^{15}$  molec/cm<sup>2</sup> (33%), which is skewed due to a large bias during the monsoon months ( $1.9 \times 10^{15}$  molec/cm<sup>2</sup>, 17%). The model has a more defined peak over Delhi, peaking in monsoon months, but there is only a small, diffuse peak in the observations. We discuss this discrepancy below.

Figure 7 shows that on a continental scale the seasonal distribution of HCHO slant columns over India in 2014 is not significantly different from observed distributions from 2008 to 2015. In general, HCHO columns follow a similar cycle in the winter and pre-monsoon months, but there are substantial year to year variations during the monsoon and post-monsoon months suggesting a role for large-scale variation in meteorology associated with the monsoon. Figure 8 shows the corresponding differences in the spatial distribution of HCHO columns from 2014 compared to the mean 2008–2015 distribution. Median HCHO columns for 2014 are slightly higher than the eight-year average by  $0.4 \times 10^{15}$  molec/cm<sup>2</sup>, but show a similar spatial distribution taking into account year-to-

year variability (Figure 7). Mahajan et al. (2015) reports an analysis of the drivers of year-to-year  
375 variations in HCHO columns over India.

### Biogenic VOCs

Figure 7 shows that the observed continental-scale HCHO seasonal cycle is reproduced by the model  
but with a positive model bias typically within  $1 \times 10^{15}$  molec/cm<sup>2</sup>. On this spatial scale, most of the  
seasonal cycle can be explained by variations in surface temperature, suggesting a larger than ex-  
380 pected role for biogenic emissions. However, the model fails to capture the elevated monthly column  
during October 2014. We find this is driven mostly by observed variations in HCHO columns over  
tropical India, coinciding with the withdrawal of the monsoon from India between late September  
and mid-October (Pai and Bhan, 2015). We suggest this variation represents a response from the  
vegetation that is missing in the model. Model error associated with cloud coverage and consequent  
385 errors associated with the partitioning between direct and diffuse PAR could result in large-scale  
changes in biogenic emissions.

The only forested region that is the exception to the continental-scale picture is over Kerala (de-  
noted by S in Figure 1). Kerala has a tropical maritime climate with little seasonal variation in  
temperature. The forested region in Kerala neighbours an urban conurbation associated with a high  
390 level of NO<sub>x</sub> (Figure 9), which influence the HCHO yields from the oxidation of biogenic VOCs, as  
described in Section 3.1. Figure 10 shows that the observed seasonal distribution of HCHO columns  
at this site, broadly reproduced by the model, peaks during the winter and is lowest during pre-  
monsoon and monsoon months. The size of the seasonal variation in HCHO columns is not fully  
explained by the small seasonal variation in surface temperature, suggesting a role for an additional  
395 driver. Figure 10 shows that over this region, the seasonal distribution is driven mostly by changes in  
leaf phenology rather than temperature or PAR. Satellite observations of LAI drop from  $\simeq 4$  m<sup>2</sup>/m<sup>2</sup>  
to  $\simeq 1$  m<sup>2</sup>/m<sup>2</sup> during the monsoon months and recover in the post-monsoon months. We find sim-  
ilar behaviour over the east and northeastern forests with large reductions in LAI during monsoon  
months (not shown). This behaviour is consistent with vegetation taking advantage of the decreased  
400 temperatures and higher precipitation rates during the monsoon season to regulate leaf flushing.

Past work has found a contrasting relationship between leaf phenology and satellite observations  
of HCHO columns over the Amazon basin (Barkley et al., 2009) in which HCHO columns and  
LAI values dropped in the transition period between the wet and dry seasons, and recovered soon  
afterwards. Thermotolerance is one hypothesis that describes why leaves emit isoprene (Singaas  
405 et al., 1997). After a few weeks of emerging, the emission capacity of leaves peaks and subsequently  
declines with age. To explain the variation in HCHO columns, Barkley et al. (2009) proposed wide-  
scale leaf flushing that allowed vegetation to maximize their protection against the light-rich envi-  
ronment of the dry season. A demographic model of leaf phenology based on the hypothesis that  
trees seek an optimal LAI as a function of available light and soil water (Caldararu et al., 2012,

410 2014) explained the observed increase in LAI over the Amazon basin during the dry season as a net addition of leaves in response to increased solar radiation.

### Pyrogenic VOCs

The main loci for biomass burning are: 1) a region approximately encompassed by the state boundaries of Punjab and Haryana (Figure 1) and 2) Northeastern India. The states of Punjab and Haryana  
415 have two growing seasons: May–September (rice) and November–April (wheat). Paddy stubble burning in May and October/November represents agriculture burning of wheat and rice residue, respectively. For our purposes Northeastern India includes the Seven Sister states (Arunachal Pradesh, Assam, Meghalaya, Manipur, Mizoram, Nagaland, and Tripura), where there are significant forest fires particularly during March and April. This is mainly due to deforestation to convert forests to  
420 agricultural land (Santenda and Kaushik, 2014). These two geographical regions account for more than 70% of these emissions during 2014.

To illustrate the impact of fires on HCHO column variations we use MODIS firecount data (Justice et al., 2002) to identify when and where fires occur over the states of Punjab and Haryana, and Northeastern India. We calculate monthly mean HCHO vertical columns and firecounts. We then  
425 determine which  $0.25^\circ \times 0.3125^\circ$  grid cells are most affected by fires by selecting these cells in the top 20th percentile of cumulative fires.

Figure 11 shows that the highest number of firecounts generally correspond to when there is the largest difference between all grid cells and those most affected by fire. This difference is relatively small over the states of Punjab and Haryana with a peak value of  $1.5 \times 10^{15}$  molec/cm<sup>2</sup> during May.  
430 There is a large difference during March and April over the Northeastern India where fires contribute up to  $5 \times 10^{15}$  molec/cm<sup>2</sup> to the monthly mean. We find that this contribution to HCHO columns is localized in time and geography.

### Anthropogenic Hotspots

Guided by *a priori* emissions and our box modelling, we anticipate that propene is the only anthropogenic VOC likely to produce HCHO sufficiently rapidly that we can relate elevated HCHO  
435 columns to emissions. However, we find little evidence that seasonally-averaged OMI HCHO columns are elevated over Indian megacities due to limits in the signal to noise, in agreement with previous work (Mahajan et al., 2015).

We use a temporal oversampling approach, following Zhu et al. (2014), to improve the spatial  
440 resolution of HCHO columns over Delhi and the surrounding region. Oversampling increases the signal-to-noise ratio and allows for inspection of finer spatial features, at the expense of the temporal information. We focus on Delhi because the National Capital Region has a population of approximately 17 million people (Perianayagam and Goli, 2012) over a geographical area of approximately 2000 km<sup>2</sup>. Based on the MIX emissions inventory, which is indicative of values from 2010, (Figure

445 6) we expect to see an elevated signal from this region. This bottom-up emission inventory likely overestimates emissions from the transport sector, which has seen the biggest change from 2010 to 2014 (per. comm.: Jun-ichi Kurokawa, Japan Environmental Sanitation Center, October 2017).

First, our area of focus is divided into a very high resolution grid ( $0.02^\circ \times 0.02^\circ$ ). The temporally-averaged column for each point in this grid is the average of the OMI observational vertical columns  
450 collected during 2014 with the centre point within 43 km in both the latitudinal and longitudinal directions. This effectively smears out these observations over 43 km squares. Here, we average over  $43 \times 43 \text{ km}^2$  squares rather than the 24 km radius circles), sampling all the 2014 vertical HCHO columns from the observational dataset for the area around Delhi.

Figure 12 shows that the oversampling method results in distinct elevated HCHO columns over  
455 New Delhi and along major roadways, although the gradients are still noisy. The magnitude of this elevation is  $\mathcal{O}(10^{15})$  molec/cm<sup>2</sup>. Elevated areas to the East and Southeast of the city may represent HCHO produced from VOC transported downwind from Delhi. Based on our results we find that anthropogenic emissions do not appear to play a large role in the observed column variations of HCHO over India.

460 However, our use of HCHO columns from the OMI instrument, which has a local overpass time of 1330, may be hindering our ability to observe the anthropogenic contribution to HCHO. Biogenic emissions and to a lesser extent biomass burning emissions peak in early afternoon hours, which is ideal for OMI. Emissions from the transportation sector, a major source of anthropogenic VOCs, peak during early morning and late afternoon associated with commuter traffic. We argue that the  
465 early morning 0930 overpass of the Global Ozone Monitoring Experiment (GOME-2) aboard the MetOp satellite is better suited to capture these anthropogenic emissions. Secondary production of HCHO is generally larger than direct emissions of HCHO, and will occur a few hours after the peak commuter time (e.g. Lin et al. (2012); Wang et al. (2017)). The early morning rush hour in Delhi starts after 0700 so we expect a 0930 overpass to also capture some fraction of the secondary HCHO  
470 production. Using data collected from morning and afternoon overpass times to describe diurnal variations of HCHO was presented by De Smedt et al. (2015), but they did not discuss the relative importance of different VOC emission sources at these different times.

Figure 13 shows the annual mean HCHO columns observed by GOME-2 and OMI. For our preliminary argument we are interested only in the distribution of HCHO columns. Here, we have  
475 standardized these data for the whole country so that they have a mean of zero and a unit standard deviation using:

$$z_i = \frac{x_i - \bar{x}}{s}, \quad (1)$$

where  $x_i$  is a data point,  $\bar{x}$  is the sample mean, and  $s$  is the sample standard deviation. This allows us to compare the two data without worrying about bias. The resulting z-scores represent the number  
480 of standard deviations from the population mean. We find that GOME-2 data has higher columns over the IGP, while OMI more clearly emphasizes the forested regions that we can identify indepen-

dently through LAI or NDVI measurements (Figure 1). This qualitative test appears to support our hypothesis and is an early demonstration of how datasets with continuous measurements of a region (such as those from a geostationary satellite) could help capture the temporal variability of HCHO. Taking advantage of the complementary information from multiple sensors that have different local overpass time requires a sophisticated inverse model approach.

### 3.4 Inferring Isoprene Emissions from OMI HCHO Columns

Based on our analysis of the HCHO yields from Indian VOC sources, and the distribution of observed HCHO columns we conclude that the majority of the observed HCHO column variation is due to biogenic VOC emissions. Here, we adopt a simple inversion methodology based on linear regression to infer isoprene emissions from OMI HCHO columns (Palmer et al., 2006).

First, we filter HCHO column data over India to minimize any interference from pyrogenic and anthropogenic sources. We focus on two relatively remote areas of India: “East” defined approximately as the area spanning 16–25°N and 76–87°E, and “Northeast” defined as the region of India east of 90°E (Figure 1). We remove scenes with MODIS land cover classifications (Friedl et al., 2010) corresponding to croplands (including cropland mosaics), urban/built-up, snow/ice, barren/sparsely vegetated, and water bodies. We also filter out potential fire-affected data by identifying for each day the cells of the  $0.25^\circ \times 0.3125^\circ$  grid in which fires are reported in the MODIS active fire product. The data from these, and the adjacent cells, are then removed for that day as well as the preceding and succeeding days, following Barkley et al. (2013).

Second, we determine the model relationship between local isoprene emissions  $E$  (molec/cm<sup>2</sup>/s), as calculated by MEGAN (Guenther et al., 2012), and HCHO columns  $\Omega$  (molec/cm<sup>2</sup>):

$$\Omega = SE_{\text{VOC}} + \Omega_b, \quad (2)$$

where the slope  $S$  represents the production of HCHO column per emission of isoprene, and the intercept  $\Omega_b$  represent the HCHO column contributions from longer-lived VOCs mainly from the oxidation of methane. We resolve seasonal *a priori* emissions of isoprene from observed HCHO columns by transposing the model linear relationship between isoprene emissions and HCHO columns.

For both study regions, we find a statistically significant linear relationship between those variables (Table 3), where Pearson correlation coefficients  $r$  range from 0.52 to 0.82, with a typical value in excess of 0.70. The slope values ( $10^3$ s) vary with region and season. The offsets that represent the background HCHO column is higher during the pre-monsoon and monsoon summer months when we expect larger HCHO production from a range of longer-lived VOC (including CH<sub>4</sub>) due to higher values of OH.

Our *a posteriori* emission estimates are generally lower than *a priori* values, reflecting the positive model HCHO column bias. This is most pronounced over the Northeast during the monsoon season, where *a posteriori* isoprene emissions are 88% lower than *a priori* estimate, due to the model not

capturing the sharp observed decline in HCHO that appears to be linked with monsoon conditions. We acknowledge that the bias between model and OMI HCHO columns could also reflect a bias in the OMI data (Zhu et al., 2016) but without independent measurements over this region we have  
520 chosen to assign these biases exclusively to the model.

#### 4 Concluding Remarks

We used models of atmospheric chemistry to interpret HCHO column distributions during 2014 observed by the Ozone Monitoring Instrument (OMI) satellite instrument over India. The annual mean OMI distribution of clear-sky HCHO columns is dominated by a distinctive meridional gradient in the northern half of the country, and by localized regions of high columns that coincide with  
525 forests. We found that the nested GEOS-Chem atmospheric chemistry model (spatially resolved at  $\simeq 25$  km) reproduces these broad-scale observed features with a positive model bias, particularly over the Indo-Gangetic Plain and Delhi.

Over India, HCHO has biogenic, pyrogenic, and anthropogenic sources of volatile organic compounds (VOCs), some of which are spatially and temporally disaggregated. Using the CAABA 0-D photochemistry model, we explored a range of forest and urban photochemical environments found over India and their subsequent influence on HCHO concentrations. HCHO columns are related to local VOC emissions with a spatial smearing that increase with the VOC lifetime. We found that isoprene has the largest molar yield of HCHO that is typically realized within a few hours in the  
535 presence of moderate levels of nitrogen oxides ( $\simeq 1$  ppbv), in agreement with previous studies. However, we also found that forested regions that neighbour major urban conurbations (e.g. in the state of Kerala) are exposed to much higher levels of nitrogen oxides ( $\simeq 8$  ppbv). This results in depleted hydroxyl radical concentrations and a delay in the production of HCHO from isoprene oxidation. Informed by a regional bottom-up emission inventory for India, we found that propene is the only  
540 major component of anthropogenic VOCs that produces HCHO at comparable (but slower) rate to isoprene.

We found that the GEOS-Chem model reproduces observed spatial distributions during winter (JF) and pre-monsoon months (MAM) better than during monsoon (JJAS) and post-monsoon (OND) months. We attributed these differences in model skill to the response of the natural biosphere to  
545 changes in the meteorological and photochemical environments associated with the onset and retreat of the monsoon. We found that on a continental scale much of the seasonal cycle in observed HCHO columns can be explained by monthly variations in surface temperature. This observation together with the strong local relationship we found between isoprene emissions and HCHO production suggests a role for biogenic VOCs, in agreement with the GEOS-Chem model calculation. We also  
550 found that the seasonal cycle during 2014 is not significantly different from the 2008-2015 mean seasonal variation but there are large year to year variations. There are two main loci for biomass

burning (states of Punjab and Haryana, and northeastern India), which we found contributes a significant contribution to observed columns only during March to April over northeastern India. The slow production of HCHO from propene oxidation results in a smeared hotspot over Delhi that we resolved only by using a temporal oversampling method. Based on comparing GOME-2 and OMI HCHO column distributions, we argue that the early morning overpass time is better for quantifying anthropogenic emissions soon after the rush hour and before biogenic emissions are at their early afternoon peak.

Using a linear regression model to relate GEOS-Chem isoprene emissions to HCHO columns we inferred seasonal isoprene emissions over two key forest regions from the OMI HCHO column data. We found that the *a posteriori* emissions are typically lower than the *a priori* emissions, with a much stronger reduction of emissions during the monsoon season. This reduction in emissions during monsoon months coincided with a large drop in satellite observations of leaf phenology. Large-scale differences in observed and model HCHO columns during monsoon months may highlight errors in seasonal variations in basal emission rates and/or model errors associated with the underlying meteorological environments, e.g. partitioning of direct and diffuse photosynthetically active radiation.

The next logical step to this analysis is to estimate simultaneous estimates of anthropogenic, pyrogenic, and biogenic VOC emissions by using data collecting data from morning and afternoon local overpass times. In the case of biogenic VOC emissions, information from HCHO columns together with leaf phenology (e.g. leaf area index) and land surface parameters (e.g., soil moisture), can be integrated to develop a new satellite data-driven isoprene emission inventory. A self-consistent pan-tropical emission inventory for isoprene, for example, would help to improve understanding of tropospheric O<sub>3</sub> and organic aerosol that represent some of the largest uncertainties associated with the Earth system. Our ability to achieve this capability is improved by the launch of TROPOMI aboard Sentinel-5P that will result in daily maps of HCHO columns and complementary trace gases at a spatial resolution of 7 km, which dramatically increases the number of clear-sky scenes available for the analysis.

## 5 Data availability

The OMHCHOv003 OMI HCHO column data are publicly available through NASA's Mirador website. Model data are archived at the Edinburgh Data Share <http://dx.doi.org/10.7488/ds/2305>.

*Acknowledgements.* L.S. was funded by the NERC National Centre for Earth Observation (NCEO020005), and P.I.P. gratefully acknowledges his Royal Society Wolfson Research Merit Award. L.S. also acknowledges funding from the British Council Newton Fund (215829867), administered by the University of Birmingham. We are grateful to Kelly Chance from the Harvard-Smithsonian Center for Astrophysics, the Harvard University

GEOS-Chem group who maintains the model, and to Rolf Sander for maintaining the CAABA/MECCA box model.

*Author contributions.* L.S. and P.I.P designed the computational experiments; P.I.P. and L.S. wrote the paper; G.G.A. provided input on the paper regarding the OMI data analysis.

## 590 References

- Abbot, D. S., Palmer, P. I., Martin, R. V., Chance, K. V., Jacob, D. J., and Guenther, A.: Seasonal and interannual variability of North American isoprene emissions as determined by formaldehyde column measurements from space, *Geophysical Research Letters*, 30, doi:10.1029/2003GL017336, <http://dx.doi.org/10.1029/2003GL017336>, 1886, 2003.
- 595 Barkley, M. P., Palmer, P. I., Kuhn, U., Kesselmeier, J., Chance, K., Kurosu, T. P., Martin, R. V., Helmig, D., and Guenther, A.: Net ecosystem fluxes of isoprene over tropical South America inferred from Global Ozone Monitoring Experiment (GOME) observations of HCHO columns, *Journal of Geophysical Research: Atmospheres*, 113, doi:10.1029/2008JD009863, <http://dx.doi.org/10.1029/2008JD009863>, d20304, 2008.
- Barkley, M. P., Palmer, P. I., De Smedt, I., Karl, T., Guenther, A., and Van Roozendaal, M.: Regulated large-scale annual shutdown of Amazonian isoprene emissions?, *Geophysical Research Letters*, 36, doi:10.1029/2008GL036843, 2009.
- 600 Barkley, M. P., De Smedt, I., Van Roozendaal, M., Kurosu, T. P., Chance, K., Arneeth, A., Hagberg, D., Guenther, A., Paulot, F., Marais, E., and Mao, J.: Top-down isoprene emissions over tropical South America inferred from SCIAMACHY and OMI formaldehyde columns, *Journal of Geophysical Research: Atmospheres*, 118, 6849–6868, doi:10.1002/jgrd.50552, 2013.
- 605 Caldararu, S., Palmer, P. I., and Purves, D. W.: Inferring Amazon leaf demography from satellite observations of leaf area index, *Biogeosciences*, 9, 1389–1404, doi:10.5194/bg-9-1389-2012, <https://www.biogeosciences.net/9/1389/2012/>, 2012.
- Caldararu, S., Purves, D. W., and Palmer, P. I.: Phenology as a strategy for carbon optimality: a global model, *Biogeosciences*, 11, 763–778, doi:10.5194/bg-11-763-2014, <https://www.biogeosciences.net/11/763/2014/>, 2014.
- 610 Carter, W.: Documentation of the SAPRC-99 Chemical Mechanism for VOC Reactivity Assessment, 2000.
- Chance, K., Palmer, P. I., Spurr, R. J. D., Martin, R. V., Kurosu, T. P., and Jacob, D. J.: Satellite observations of formaldehyde over North America from GOME, *Geophysical Research Letters*, 27, 3461–3464, doi:10.1029/2000GL011857, <http://dx.doi.org/10.1029/2000GL011857>, 2000.
- 615 Cohen, A. J., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope, C. A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C. J. L., and Forouzanfar, M. H.: Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015, *The Lancet*, 389, 1907 – 1918, doi:[https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6), <http://www.sciencedirect.com/science/article/pii/S0140673617305056>, 2017.
- 620 Curci, G., Palmer, P. I., Kurosu, T. P., Chance, K., and Visconti, G.: Estimating European volatile organic compound emissions using satellite observations of formaldehyde from the Ozone Monitoring Instrument, *Atmospheric Chemistry and Physics*, 10, 11 501–11 517, doi:10.5194/acp-10-11501-2010, <https://www.atmos-chem-phys.net/10/11501/2010/>, 2010.
- 625 De Smedt, I., Müller, J.-F., Stavrou, T., van der A, R., Eskes, H., and Van Roozendaal, M.: Twelve years of global observations of formaldehyde in the troposphere using GOME and SCIAMACHY sen-

- sors, *Atmospheric Chemistry and Physics*, 8, 4947–4963, doi:10.5194/acp-8-4947-2008, <https://www.atmos-chem-phys.net/8/4947/2008/>, 2008.
- 630 De Smedt, I., Stavrou, T., Müller, J.-F., van der A, R. J., and Van Roozendael, M.: Trend detection in satellite observations of formaldehyde tropospheric columns, *Geophysical Research Letters*, 37, doi:10.1029/2010GL044245, <http://dx.doi.org/10.1029/2010GL044245>, 118808, 2010.
- De Smedt, I., Stavrou, T., Hendrick, F., Danckaert, T., Vlemmix, T., Pinardi, G., Theys, N., Lerot, C., Gielen, C., Vigouroux, C., Hermans, C., Fayt, C., Veeffkind, P., Müller, J.-F., and Van Roozendael, M.: Diurnal, seasonal and long-term variations of global formaldehyde columns inferred from combined OMI and GOME-2 observations, *Atmospheric Chemistry and Physics*, 15, 12 519–12 545, doi:10.5194/acp-15-12519-2015, <http://www.atmos-chem-phys.net/15/12519/2015/>, 2015.
- 635 Eastham, S. D., Weisenstein, D. K., and Barrett, S. R.: Development and evaluation of the unified tropospheric–stratospheric chemistry extension (UCX) for the global chemistry-transport model GEOS-Chem, *Atmospheric Environment*, 89, 52 – 63, doi:10.1016/j.atmosenv.2014.02.001, 2014.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X.: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sensing of Environment*, 114, 168–182, doi:10.1016/j.rse.2009.08.016, 2010.
- 645 Fu, T.-M., Jacob, D. J., Palmer, P. I., Chance, K., Wang, Y. X., Barletta, B., Blake, D. R., Stanton, J. C., and Pilling, M. J.: Space-based formaldehyde measurements as constraints on volatile organic compound emissions in east and south Asia and implications for ozone, *Journal of Geophysical Research: Atmospheres*, 112, doi:10.1029/2006JD007853, <http://dx.doi.org/10.1029/2006JD007853>, d06312, 2007.
- González Abad, G., Liu, X., Chance, K., Wang, H., Kurosu, T. P., and Suleiman, R.: Updated Smithsonian Astrophysical Observatory Ozone Monitoring Instrument (SAO OMI) formaldehyde retrieval, *Atmospheric Measurement Techniques*, 8, 19–32, doi:10.5194/amt-8-19-2015, 2015.
- 650 Gonzi, S., Palmer, P. I., Barkley, M. P., De Smedt, I., and Van Roozendael, M.: Biomass burning emission estimates inferred from satellite column measurements of HCHO: Sensitivity to co-emitted aerosol and injection height, *Geophysical Research Letters*, 38, doi:10.1029/2011GL047890, <http://dx.doi.org/10.1029/2011GL047890>, 114807, 2011.
- 655 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geoscientific Model Development*, 5, 1471–1492, doi:10.5194/gmd-5-1471-2012, 2012.
- 660 Jaeglé, L., Jacob, D. J., Wang, Y., Weinheimer, A. J., Ridley, B. A., Campos, T. L., Sachse, G. W., and Hagen, D. E.: Sources and chemistry of NO<sub>x</sub> in the upper troposphere over the United States, *Geophysical Research Letters*, 25, 1705–1708, doi:10.1029/97GL03591, <http://dx.doi.org/10.1029/97GL03591>, 1998.
- Justice, C., Giglio, L., Korontzi, S., Owens, J., Morisette, J., Roy, D., Descloitres, J., Alleaume, S., Petitcolin, F., and Kaufman, Y.: The MODIS fire products, *Remote Sensing of Environment*, 83, 244 – 262, doi:[https://doi.org/10.1016/S0034-4257\(02\)00076-7](https://doi.org/10.1016/S0034-4257(02)00076-7), <http://www.sciencedirect.com/science/article/pii/S0034425702000767>, the Moderate Resolution Imaging Spectroradiometer (MODIS): a new generation of Land Surface Monitoring, 2002.

- Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission inventory in ASia (REAS) version 2, *Atmospheric Chemistry and Physics*, 13, 11 019–11 058, doi:10.5194/acp-13-11019-2013, 2013.
- Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Mälkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J. O. V., and Saari, H.: The Ozone Monitoring Instrument, *IEEE Transactions on Geoscience and Remote Sensing*, 44, 1093–1101, 2006.
- 675 Li, M., Zhang, Q., Kurokawa, J.-I., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmospheric Chemistry and Physics*, 17, 935–963, doi:10.5194/acp-17-935-2017, 2017.
- Lin, Y. C., Schwab, J. J., Demerjian, K. L., Bae, M.-S., Chen, W.-N., Sun, Y., Zhang, Q., Hung, H.-M., and Perry, J.: Summertime formaldehyde observations in New York City: Ambient levels, sources and its contribution to HO<sub>x</sub> radicals, *Journal of Geophysical Research: Atmospheres*, 117, doi:10.1029/2011JD016504, <http://dx.doi.org/10.1029/2011JD016504>, d08305, 2012.
- 680 Lu, Z. and Streets, D. G.: Increase in NO<sub>x</sub> Emissions from Indian Thermal Power Plants during 1996–2010: Unit-Based Inventories and Multisatellite Observations, *Environmental Science & Technology*, 46, 7463–7470, doi:10.1021/es300831w, 2012.
- Lu, Z., Zhang, Q., and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996-2010, *Atmospheric Chemistry and Physics*, 11, 9839–9864, doi:10.5194/acp-11-9839-2011, 2011.
- Mahajan, A. S., Smedt, I. D., Biswas, M. S., Ghude, S., Fadnavis, S., Roy, C., and van Roozendaal, M.: Inter-annual variations in satellite observations of nitrogen dioxide and formaldehyde over India, *Atmospheric Environment*, 116, 194 – 201, doi:10.1016/j.atmosenv.2015.06.004, 2015.
- 690 Marais, E. A., Jacob, D. J., Kurosu, T. P., Chance, K., Murphy, J. G., Reeves, C., Mills, G., Casadio, S., Millet, D. B., Barkley, M. P., Paulot, F., and Mao, J.: Isoprene emissions in Africa inferred from OMI observations of formaldehyde columns, *Atmospheric Chemistry and Physics*, 12, 6219–6235, doi:10.5194/acp-12-6219-2012, 2012.
- 695 Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., and Evans, M. J.: Global inventory of nitrogen oxide emissions constrained by space-based observations of NO<sub>2</sub> columns, *Journal of Geophysical Research: Atmospheres*, 108, doi:10.1029/2003JD003453, 2003.
- Marvin, M. R., Wolfe, G. M., Salawitch, R. J., Canty, T. P., Roberts, S. J., Travis, K. R., Aikin, K. C., de Gouw, J. A., Graus, M., Hanisco, T. F., Holloway, J. S., Hübler, G., Kaiser, J., Keutsch, F. N., Peischl, J., Pollack, I. B., Roberts, J. M., Ryerson, T. B., Veres, P. R., and Warneke, C.: Impact of evolving isoprene mechanisms on simulated formaldehyde: An inter-comparison supported by in situ observations from SENEX, *Atmospheric Environment*, 164, 325 – 336, doi:10.1016/j.atmosenv.2017.05.049, 2017.
- 700 Millet, D. B., Jacob, D. J., Boersma, K. F., Fu, T.-M., Kurosu, T. P., Chance, K., Heald, C. L., and Guenther, A.: Spatial distribution of isoprene emissions from North America derived from formaldehyde column measurements by the OMI satellite sensor, *Journal of Geophysical Research: Atmospheres*, 113, doi:10.1029/2007JD008950, <http://dx.doi.org/10.1029/2007JD008950>, d02307, 2008.
- 705

- Pai, D. S. and Bhan, S. C.: Monsoon 2014 A Report, Tech. Rep. ESSO/IMD/Synoptic Met./01(2015)/17, India Meteorological Department, 2015.
- 710 Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A., and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *Journal of Geophysical Research: Atmospheres*, 106, 14 539–14 550, doi:10.1029/2000JD900772, 2001.
- Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over North America using formaldehyde column observations from space, *Journal of Geophysical Research: Atmospheres*, 108, doi:10.1029/2002JD002153, 2003.
- 715 Palmer, P. I., Abbot, D. S., Fu, T.-M., Jacob, D. J., Chance, K., Kurosu, T. P., Guenther, A., Wiedinmyer, C., Stanton, J. C., Pilling, M. J., Pressley, S. N., Lamb, B., and Sumner, A. L.: Quantifying the seasonal and interannual variability of North American isoprene emissions using satellite observations of the formaldehyde column, *Journal of Geophysical Research: Atmospheres*, 111, doi:10.1029/2005JD006689, 2006.
- 720 Palmer, P. I., Barkley, M. P., Kurosu, T. P., Lewis, A. C., Saxton, J. E., Chance, K., and Gatti, L. V.: Interpreting satellite column observations of formaldehyde over tropical South America, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 365, 1741–1751, doi:10.1098/rsta.2007.2042, <http://rsta.royalsocietypublishing.org/content/365/1856/1741>, 2007.
- 725 Perianayagam, A. and Goli, S.: Provisional results of the 2011 Census of India: Slowdown in growth, ascent in literacy, but more missing girls, *International Journal of Social Economics*, 39, 785–801, doi:10.1108/03068291211253395, 2012.
- Sander, R., Baumgaertner, A., Gromov, S., Harder, H., Jöckel, P., Kerkweg, A., Kubistin, D., Regelin, E., Riede, H., Sandu, A., Taraborrelli, D., Tost, H., and Xie, Z.-Q.: The atmospheric chemistry box model CAABA/MECCA-3.0, *Geoscientific Model Development*, 4, 373–380, doi:10.5194/gmd-4-373-2011, 2011.
- 730 Sander, R., Jöckel, P., Kirner, O., Kunert, A. T., Landgraf, J., and Pozzer, A.: The photolysis module JVAL-14, compatible with the MESSy standard, and the JVal PreProcessor (JVPP), *Geoscientific Model Development*, 7, 2653–2662, doi:10.5194/gmd-7-2653-2014, <https://www.geosci-model-dev.net/7/2653/2014/>, 2014.
- Santenda and Kaushik, A. D.: Forest Fire Disaster Management, Tech. rep., Institute of Disaster Management, Ministry of Home Affairs, New Delhi, ISBN-978-93-82571-09-4, 2014.
- 735 Sharma, S., Chatani, S., Mahtta, R., Goel, A., and Kumar, A.: Sensitivity analysis of ground level ozone in India using WRF-CMAQ models, *Atmospheric Environment*, 131, 29 – 40, doi:<https://doi.org/10.1016/j.atmosenv.2016.01.036>, <http://www.sciencedirect.com/science/article/pii/S1352231016300504>, 2016.
- 740 Shim, C., Wang, Y., Choi, Y., Palmer, P. I., Abbot, D. S., and Chance, K.: Constraining global isoprene emissions with Global Ozone Monitoring Experiment (GOME) formaldehyde column measurements, *Journal of Geophysical Research: Atmospheres*, 110, doi:10.1029/2004JD005629, <http://dx.doi.org/10.1029/2004JD005629>, d24301, 2005.
- Singaas, E. L., Lerdau, M., Winter, K., and Sharkey, T. D.: Isoprene increases thermotolerance of isoprene-emitting species, *Plant Physiol.*, pp. 1413–1420, doi:DOI:10.1104/pp.115.4.1413, 1997.
- 745 Stavrakou, T., Müller, J.-F., De Smedt, I., Van Roozendael, M., van der Werf, G. R., Giglio, L., and Guenther, A.: Global emissions of non-methane hydrocarbons deduced from SCIAMACHY formaldehyde columns

- through 2003–2006, *Atmospheric Chemistry and Physics*, 9, 3663–3679, doi:10.5194/acp-9-3663-2009, <https://www.atmos-chem-phys.net/9/3663/2009/>, 2009.
- 750 Stavrakou, T., Müller, J.-F., Bauwens, M., De Smedt, I., Lerot, C., Van Roozendael, M., Clerbaux, C., Boersma, K. F., van der A, R., and Song, Y.: Substantial Underestimation of Post-Harvest Burning Emissions in the North China Plain Revealed by Multi-Species Space Observations, *Nature Scientific Reports*, 6, doi:10.1038/srep32307, 2016.
- Taraborrelli, D., Lawrence, M. G., Butler, T. M., Sander, R., and Lelieveld, J.: Mainz Isoprene Mechanism 2 (MIM2): an isoprene oxidation mechanism for regional and global atmospheric modelling, *Atmospheric Chemistry and Physics*, 9, 2751–2777, doi:10.5194/acp-9-2751-2009, 2009.
- 755 Thomas, W., Hegels, E., Slijkhuis, S., Spurr, R., and Chance, K.: Detection of biomass burning combustion products in Southeast Asia from backscatter data taken by the GOME Spectrometer, *Geophysical Research Letters*, 25, 1317–1320, doi:10.1029/98GL01087, <http://dx.doi.org/10.1029/98GL01087>, 1998.
- 760 Tyagi, S., Tiwari, S., Mishra, A., Hopke, P. K., Attri, S., Srivastava, A., and Bisht, D.: Spatial variability of concentrations of gaseous pollutants across the National Capital Region of Delhi, India, *Atmospheric Pollution Research*, 7, 808–816, doi:10.1016/j.apr.2016.04.008, 2016.
- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, doi:10.5194/essd-9-697-2017, 2017.
- 765 Volkamer, R., Sheehy, P., Molina, L. T., and Molina, M. J.: Oxidative capacity of the Mexico City atmosphere Part 1: A radical source perspective, *Atmospheric Chemistry and Physics*, 10, 6969–6991, doi:10.5194/acp-10-6969-2010, <https://www.atmos-chem-phys.net/10/6969/2010/>, 2010.
- Wang, C., Huang, X.-F., Han, Y., Zhu, B., and He, L.-Y.: Sources and Potential Photochemical Roles of Formaldehyde in an Urban Atmosphere in South China, *Journal of Geophysical Research: Atmospheres*, doi:10.1002/2017JD027266, <http://dx.doi.org/10.1002/2017JD027266>, 2017JD027266, 2017.
- 770 Whalley, L. K., Furneaux, K. L., Goddard, A., Lee, J. D., Mahajan, A., Oetjen, H., Read, K. A., Kaaden, N., Carpenter, L. J., Lewis, A. C., Plane, J. M. C., Saltzman, E. S., Wiedensohler, A., and Heard, D. E.: The chemistry of OH and HO<sub>2</sub> radicals in the boundary layer over the tropical Atlantic Ocean, *Atmospheric Chemistry and Physics*, 10, 1555–1576, doi:10.5194/acp-10-1555-2010, <https://www.atmos-chem-phys.net/10/1555/2010/>, 2010.
- Wolfe, G. M., Kaiser, J., Hanisco, T. F., Keutsch, F. N., de Gouw, J. A., Gilman, J. B., Graus, M., Hatch, C. D., Holloway, J., Horowitz, L. W., Lee, B. H., Lerner, B. M., Lopez-Hilifiker, F., Mao, J., Marvin, M. R., Peischl, J., Pollack, I. B., Roberts, J. M., Ryerson, T. B., Thornton, J. A., Veres, P. R., and Warneke, C.: Formaldehyde production from isoprene oxidation across NO<sub>x</sub> regimes, *Atmospheric Chemistry and Physics*, 16, 2597–2610, doi:10.5194/acp-16-2597-2016, <https://www.atmos-chem-phys.net/16/2597/2016/>, 2016.
- 780 Young, E. L. and Paton-Walsh, C.: Formaldehyde and nitrogen dioxide in smoke plumes from Australia’s Black Saturday fires, *IOP Conference Series: Earth and Environmental Science*, 11, 012 023, <http://stacks.iop.org/1755-1315/11/i=1/a=012023>, 2010.
- 785 Zhu, L., Jacob, D. J., Mickley, L. J., Marais, E. A., Cohan, D. S., Yoshida, Y., Duncan, B. N., Abad, G. G., and Chance, K. V.: Anthropogenic emissions of highly reactive volatile organic compounds in eastern Texas

inferred from oversampling of satellite (OMI) measurements of HCHO columns, *Environmental Research Letters*, 9, 114 004, 2014.

790 Zhu, L., Jacob, D. J., Kim, P. S., Fisher, J. A., Yu, K., Travis, K. R., Mickley, L. J., Yantosca, R. M., Sulprizio,  
M. P., De Smedt, I., González Abad, G., Chance, K., Li, C., Ferrare, R., Fried, A., Hair, J. W., Hanisco, T. F.,  
Richter, D., Jo Scarino, A., Walega, J., Weibring, P., and Wolfe, G. M.: Observing atmospheric formaldehyde  
(HCHO) from space: validation and intercomparison of six retrievals from four  
satellites (OMI, GOME2A, GOME2B, OMPS) with SEAC<sup>4</sup>RS aircraft observations over the southeast  
795 US, *Atmospheric Chemistry and Physics*, 16, 13 477–13 490, doi:10.5194/acp-16-13477-2016, [https://www.  
atmos-chem-phys.net/16/13477/2016/](https://www.atmos-chem-phys.net/16/13477/2016/), 2016.




Figure 1: Geography of India: population density based on the 2001 census (top left); focal points of this study (top right); and annual mean values of the Normalized Difference Vegetation Index (NDVI) for 2014 from the NASA MODIS instrument, averaged on the  $0.25^\circ \times 0.3125^\circ$  GEOS-Chem nested grid. For the map of focal points: the blue outline represent the states of Punjab and Haryana; the red outline denotes Seven sister states; and the yellow outline represents the state of Kerala. The pink and green shaded areas denote the NE and E forest sites, where we infer isoprene emissions from OMI HCHO columns. The grey shaded area denotes the oversampling region used to study Delhi. The stars denote the sites where we study HCHO production using a 0-D photochemical model. The population image is modified with permission from LuminoCity3D.org: data provided by EC JRC & CIESIN, and the design provided by D. A. Smith, Barlett Centre for Advanced Spatial Analysis, University College London.




Figure 2: Seasonal isoprene emission rates ( $\text{atom C}/\text{cm}^2/\text{s}$ ) from the MEGAN inventory (Guenther et al., 2012). Data are described on the GEOS-Chem  $0.25^\circ \times 0.3125^\circ$  grid for A) winter (JF), B) pre-monsoon (MAM), C) monsoon (JJAS), and D) post-monsoon seasons (OND).




Figure 3: Time-dependent cumulative HCHO yield (per-C) produced from isoprene in three contrasting photochemical environments, and from propene, propane, and n-butane in an urban photochemical environments based on Delhi. Calculations are denoted by an alphanumeric code that is explained in Table 1.




Figure 4: Annual mean air mass factors (AMFs) for 2014 calculated using (left) nested  $0.25^\circ \times 0.3125^\circ$  and (right) standard  $2.0^\circ \times 2.5^\circ$  GEOS-Chem grid.




Figure 5: Annual mean clear-sky (left) OMI and (right) GEOS-Chem HCHO vertical columns ( $10^{15}$  molec/cm<sup>2</sup>) over India for 2014. The OMI vertical columns are described on the  $0.25^\circ \times 0.3125^\circ$  GEOS-Chem grid. These vertical columns are transformed from observed slant columns using AMFs informed by the distribution of HCHO described by the (top) nested  $0.25^\circ \times 0.3125^\circ$  and (bottom) standard  $2.0^\circ \times 2.5^\circ$  GEOS-Chem grid. Data exclusion criteria are described in the main text. The model is sampled at the time and location of each observed scene.




Figure 6: Seasonal mean clear-sky (left) OMI and (right) GEOS-Chem HCHO vertical columns ( $10^{15} \text{ molec/cm}^2$ ) for 2014, averaged on a common  $0.25^\circ \times 0.3125^\circ$  grid. Seasonal definitions are described in Section 2.




Figure 7: Timeseries of (top) OMI and GEOS-Chem vertical columns ( $10^{15}$  molec/cm<sup>2</sup>) for 2014, and (bottom) OMI slant columns ( $10^{15}$  molec/cm<sup>2</sup>) from 2008 to 2015 over India. GEOS-FP ground-level temperature (K) is also shown in the top panel. The thick grey line in the bottom panel denotes 2008–2015 HCHO column monthly means.




Figure 8: Annual mean OMI HCHO slant columns ( $10^{15}$  molec/cm<sup>2</sup>) over India for (left) 2014 and (right) 2008–2015, averaged on a common  $0.25^\circ \times 0.3125^\circ$  grid.




Figure 9: Annual mean GEOS-Chem ground-level NO<sub>x</sub> mixing ratios (ppbv) in 2014, averaged on the  $0.25^\circ \times 0.3125^\circ$  grid.




Figure 10: Timeseries of (left) monthly mean OMI and GEOS-Chem vertical HCHO columns ( $10^{15}$  molec/cm<sup>2</sup>), and GEOS-FP ground-level temperature, and (right) PAR ( $W/m^2$ ) and LAI ( $m^2/m^2$ ) over the state of Kerala in 2014.




Figure 11: Timeseries of monthly mean HCHO columns ( $10^{15}$  molec/cm<sup>2</sup>) and the number of MODIS fire detections ( $10^3$  km<sup>2</sup>/day) over (left) the states of Punjab and Haryana and (right) North-eastern India. The green line denotes all HCHO columns over the region, and the red line denotes columns that are most affected by fires as indicated by MODIS firecounts.




Figure 12: Oversampled distribution of OMI vertical HCHO columns ( $10^{15}$  molec/cm<sup>2</sup>) around Delhi. The underlying road network is reproduced with permission from © www.thunderforest.com (map) and from © www.osm.org/copyright (data).




Figure 13: Standardized HCHO slant columns for 2014 (unitless) from (left) the Global Ozone Monitoring Experiment-2 (GOME-2) aboard MetOp and (right) OMI, averaged on a common  $0.25^\circ \times 0.3125^\circ$  grid. GOME-2 and OMI have local equatorial overpass times of 0930 and 1330, respectively.

Table 1: Photochemical and meteorological scenarios used for box model calculations. The emboldened letters E, NE, S denote the forested regions (Figure 1) and D denotes Delhi. Ground-level temperature (T), relative humidity (RH), boundary layer height (BLH) and mixing ratios are GEOS-Chem monthly mean values from each location.

Scenario name	Location	Month of year	T [K]	RH [%]	BLH [m]	Fixed mixing ratio [ppbv]		
						O <sub>3</sub>	CO	NO <sub>2</sub>
E01	<b>E</b> 19.3°N 80.6°E	Jan	295	56	1240	49	232	1.1
E05		May	307	33	1873	63	131	2.4
E08		Aug	299	84	685	39	114	1.3
E11		Nov	304	64	611	49	220	0.8
NE01	<b>NE</b> 24.4°N 93.1°E	Jan	292	65	648	42	277	1.8
NE05		May	302	59	1202	55	154	0.8
NE08		Aug	299	88	666	24	127	1.3
NE11		Nov	296	73	925	43	241	1.4
S01	<b>S</b> 9.8°N 76.6°E	Jan	300	59	1090	48	260	8.9
S05		May	301	78	725	32	146	8.6
S08		Aug	299	84	690	18	122	8.1
S11		Nov	299	77	732	45	274	8.9
D05	<b>D</b> 28.61°N 77.23°E	May	307	18	1975	37	2300	19

Table 2: Results of the photochemical box modelling for forested region scenarios (Table 1).

Scenario	Isoprene lifetime [min]	Time to reach peak HCHO signal [min]	HCHO yield [per-C]
E01	26	79	0.50
E05	14	59	0.55
E08	10	33	0.49
E11	20	83	0.41
NE01	28	89	0.59
NE05	10	33	0.38
NE08	10	33	0.58
NE11	18	57	0.53
S01	64	127	0.63
S05	48	147	0.63
S08	70	189	0.66
S11	52	153	0.62

Table 3: *A priori* and *a posteriori* isoprene emission estimates ( $10^{11}$  atom C/cm<sup>2</sup>/s) over NE and E forest sites (Figure 1), and the model linear regression coefficients that relate model isoprene emissions and HCHO columns.

East Forest Region						
Season	Mean <i>a priori</i> emission [ $10^{11}$ atom C/cm <sup>2</sup> /s]	Slope $S$ [ $10^3$ s]	Intercept $\Omega_b$ [ $10^{15}$ molec/cm <sup>2</sup> ]	$r^2$	Mean <i>a posteriori</i> emission [ $10^{11}$ atom C/cm <sup>2</sup> /s]	% change from <i>a priori</i>
Winter	2.8	4.1	5.3	0.59	3.0	+7
Pre-monsoon	10.2	4.3	7.0	0.67	7.8	-24
Monsoon	7.4	4.3	7.6	0.51	3.3	-55
Post-monsoon	3.6	4.9	5.2	0.47	2.6	-28
Northeast Forest Region						
Winter	3.5	3.9	3.9	0.42	4.4	+26
Pre-monsoon	17.9	2.9	6.2	0.58	12.6	-30
Monsoon	14.6	2.2	8.7	0.27	1.7	-88
Post-monsoon	6.6	4.9	5.1	0.49	2.7	-59