

Interactive comment on “Temporal characteristics of atmospheric ammonia and nitrogen dioxide over China based on emission data, satellite observations and atmospheric transport modeling since 1980” by Lei Liu et al.

M. Van Damme

martin.van.damme@ulb.ac.be

Received and published: 21 April 2017

Martin Van Damme (1), Lieven Clarisse (1), Simon Whitburn (1), Enrico Dammers (2), Pierre-François Coheur (1)

(1) Université libre de Bruxelles (ULB), Atmospheric Spectroscopy, Service de Chimie Quantique et Photophysique CP 160/09, av. F.D. Roosevelt 50, 1050 Bruxelles, Belgium

(2) Cluster Earth and Climate, Department of Earth Sciences, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

[Printer-friendly version](#)

[Discussion paper](#)



This manuscript presents time-series of NH₃ total columns from the IASI satellite instrument. As the authors point out, the retrieved columns for this trace gas are strongly dependent on the meteorological information that is used as input data in the retrieval algorithm (in particular the surface temperature and the temperature profile in the lowest layers of the atmosphere). At present, the publically available IASI-NH₃ product uses the meteorological parameters as provided in EUMETSAT IASI Level 2 data.

The IASI Level 2 data is processed in real time, and there have been several updates to the algorithm since the launch of IASI. We can confirm that in particular, the change from version 5 to version 6 of the EUMETSAT Level 2 data on 30 September 2014 (and not November as stated in the manuscript), causes on average a substantial increase of the retrieved atmospheric NH₃ columns. Hence the conclusions of the authors regarding NH₃ trends should be taken with great care.

[Interactive comment on Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2017-106, 2017.](#)

[Printer-friendly version](#)[Discussion paper](#)