

Seasonality of aerosol optical properties in the Arctic

Lauren Schmeisser^{1,3*}, John Backman², John A. Ogren^{1,3}, Elisabeth Andrews¹, Eija Asmi², Sandra Starkweather^{1,3}, Taneil Uttal³, Markus Fiebig⁴, Sangeeta Sharma⁵, Kostas Eleftheriadis⁶, Stergios Vratolis⁶, Michael Bergin⁷, Peter Tunved⁸, Anne Jefferson¹

¹University of Colorado, Cooperative Institute for Research in Environmental Sciences, Boulder, CO, USA

²Finnish Meteorological Institute, Atmospheric Composition Research, Helsinki, Finland

³National Oceanic and Atmospheric Administration, Earth System Research Laboratory, Boulder, CO, USA

⁴Norwegian Institute for Air Research, Kjeller, Norway

⁵Environment and Climate Change Canada, Science & Technology Branch, Climate Research Division, Toronto, Canada

⁶Institute of Nuclear and Radiological Science & Technology, Energy & Safety, Environmental Radioactivity Laboratory, NCSR Demokritos, Athens, Greece

⁷Duke University, Department of Civil & Environmental Engineering, Durham, NC, USA

⁸Stockholm University, Department of Environmental Science and Analytical Chemistry, Stockholm, Sweden

*Now at University of Washington, Department of Atmospheric Sciences, Seattle, WA, USA

Correspondence to: Lauren Schmeisser (lauren.schmeisser@gmail.com)

Supplemental Materials




Figure S1. Scatter plots comparing Aethalometer and reference absorption data, σ_0 and σ_{ap} , respectively, for each of the six Arctic monitoring stations. Regression coefficients are noted in the top left of each plot. A reference line of $y=3.243x$ is shown in black on each plot.