

Dear Editors and Reviewers,

Thank you very much for your letter and for the reviewers' comments concerning our manuscript entitled "Impacts of emission reduction and meteorological conditions on air quality improvement during the 2014 Youth Olympic Games in Nanjing, China" (doi:10.5194/acp-2017-114). Your comments are all valuable and very helpful for revising and improving our paper, as well as the important guiding significance to our researches. We have investigated the comments carefully and made corrections which we hope to meet with approval. Based on the instructions, we have uploaded the file of the revised manuscript.

Appended to this letter is our point-by-point response to the reviewers' comments, the change list and the marked-up manuscript.

We would like to thank you for allowing us to resubmit a revised copy of the manuscript. We hope that the revised manuscript can be accepted for publication in ACP.

Sincerely,

Qian Huang

Responses to the reviewers' comments:

Reviewer 1

Comment 1: The authors have largely improved the manuscript by addressing my comments and revising the manuscript accordingly, most of the questions have been clarified and additional model simulations were conducted and interpreted, with some new interesting results presented. I recommend acceptance of this manuscript after minor revisions on English and tables/figures.

Response: Thank you very much for your comment. We have carefully checked and modified the tables and figures, like the legend of Fig. 5 and Fig. 6, to make them more standardized. And we have checked the grammar and revise some sentences. Prof. Wang and Doctor Zhuang also helped to check and improve the English.

Change list:

- 1. Line 23-24:** Revise the abstract.
- 2. Line 52-53, 61, 63, 71, 76, 87, 93-94, 103:** Revise Section 1 Introduction.
- 3. Line 119, 121, 127, 131, 148-150, 158:** Revise Section 2 Methodology.
- 4. Line 167:** Revise Table 1.
- 5. Line 171, 173, 188-191:** Revise Section 2.3 Emissions and simulation scenarios.
- 6. Line 194:** Revise the caption of Fig. 2.
- 7. Line 196-197, 204:** Revise Section 2.3.
- 8. Line 208, 210-212, 215-218, 223-225:** Revise Section 3 Results and discussions.
- 9. Line 239-240:** Revise Table 2 and its caption.
- 10. Line 244-245:** Revise Table 3 and its caption.
- 11. Line 247, 260-261, 263:** Revise Section 3 Results and discussions
- 12. Line 271:** Revise the legend of Fig. 5, and make it more clear.
- 13. Line 276:** Revise the legend of Fig. 6, and make it more clear.
- 14. Line 283-285:** Revise Table 4 and its caption.
- 15. Line 289-291:** Revise Table 5 and its caption.
- 16. Line 305, 309, 320-324, 366, 395-397:** Revise Section 3.

- 17. Line 403-405:** Revise the caption of Fig. 12.
- 18. Line 407-411, 418:** Revise Section 3.4.
- 19. Line 436-437:** Revise the caption of Table 6.
- 20. Line 440-441, 443-444, 449, 451-452:** Revise Section 3.4.
- 21. Line 466:** Revise Section 4.
- 22. Line 480-487:** Add Section 5 Data availability and declare about competing interests.
- 23. Line 491:** Revise the acknowledgements.

1 **Impacts of emission reduction and meteorological conditions on air**
2 **quality improvement during the 2014 Youth Olympic Games in**
3 **Nanjing, China**

4

5 Qian Huang ^{1,2}, Tijian Wang ^{1,*}, Pulong Chen ¹, Xiaoxian Huang ³, Jialei Zhu ⁴, and
6 Bingliang Zhuang ¹

7 ¹ School of Atmospheric Sciences, CMA-NJU Joint Laboratory for Climate Prediction
8 Studies, Jiangsu Collaborative Innovation Center for Climate Change, Nanjing
9 University, Nanjing, 210023, China

10 ² Scientific Research Academy of Guangxi Environmental Protection, Nanning,
11 530022, China

12 ³ College of Plant Science & Technology, Huazhong Agricultural University, Wuhan,
13 430070, China

14 ⁴ Department of Climate and Space Sciences and Engineering, University of Michigan,
15 Ann Arbor, 48109, USA

16

17 Correspondence to: Tijian Wang (tjwang@nju.edu.cn)

18

19 **Abstract**

20 As the holding city of the 2nd Youth Olympic Games (YOG), Nanjing is highly
21 industrialized and urbanized facing with several air pollution issues. In order to ensure
22 better air quality during the event, the local government took great efforts to control
23 the ~~pollution~~ emissions from pollutant sources. However, air quality can still be
24 affected by synoptic weather, not only emission. In this paper, the influences of
25 meteorological factors and emission reductions were investigated using observational
26 data and numerical simulations with WRF/CMAQ. During the YOG holding month
27 (Aug., 2014), the observed hourly mean concentrations of SO₂, NO₂, PM₁₀, PM_{2.5}, CO
28 and O₃ were 11.6 μg/m³, 34.0 μg/m³, 57.8 μg/m³, 39.4 μg/m³, 0.9 mg/m³, and 38.8
29 μg/m³, respectively, which were below China National Ambient Air Quality Standard

30 (Level 2). However, model simulation showed that the weather conditions such as
31 weaker winds during the holding time were adverse for better air quality, and could
32 increase SO₂, NO₂, PM₁₀, PM_{2.5} and CO by 17.5%, 16.9%, 18.5%, 18.8%, 7.8% and
33 0.8%, respectively. Taking account of local emission abatement only, the simulated
34 SO₂, NO₂, PM₁₀, PM_{2.5} and CO was decreased by 24.6%, 12.1%, 15.1%, 8.1% and
35 7.2%, respectively. Consequently, stringent emission control measures can reduce the
36 concentrations of air pollutants in short term, and emission reduction is the very
37 important factor for the air quality improvement during the YOG, which has set up a
38 good example in air protection for important social events.

39 **KEY WORDS:** Youth Olympic Games; Emission reduction; Meteorological
40 conditions; WRF/CMAQ; Nanjing

41

42 **1 Introduction**

43 As located in the economically developed Yangtze River Delta (YRD) region of
44 China, Nanjing successfully hosted the second Youth Olympic Games (YOG) during
45 16 - 28 Aug., 2014. Nanjing is a highly urbanized city and its GDP ranked the 12th of
46 all the cities in China by 2013 (National Bureau of Statistics of China, 2014). Due to
47 fast urbanization and industrialization, heavy motor vehicles and construction dust,
48 Nanjing has long been suffered from air pollution (Dong et al., 2013; Chen et al.,
49 2015).

50 In order to realize the promise of “Green YOG”, the local government had taken
51 a series of measures to reduce emissions of air pollutants. The preparatory work
52 started from 1 Jul., 2014. Besides, the local government performed the [work plan](#)
53 [for](#) stringent environmental quality assurance ~~work plan~~ from 1 Aug. (National
54 Bureau of Statistics of China, 2014). The controlled emissions include 5 major
55 categories: industry, power plants, traffic, VOC product-related sources and others.
56 Some local petrochemical, chemical and steel industries were forced to limit or halt
57 the production. Coal-combustion enterprises were required to use high-quality coals
58 with low sulfur content and ash content. And vehicles with heavy pollution called
59 “yellow label buses” were prohibited in Nanjing during 10-28 Aug.. Oil loading and

60 unloading operations were strictly controlled. All construction processes in the city
61 were forced to stop. The ground sSurface with bare soil was covered.

62 It is well known that air quality can be affected by both meteorological factors
63 and pollutant emissions. Many cases verified that both emission abatement ~~efforts~~ and
64 weather conditions can influence the improvement of local air quality. Emission
65 control has been taken in many social events, like Beijing Olympic Games in 2008
66 and Shanghai Expo in 2010. Xing et al. (2011) suggested that emission controls
67 benefit for pollutants reduction, but meteorological effects can be either ways at
68 different locations. Cermak and Knutti (2009), Wang et al. (2009b, 2010) and Xing et
69 al. (2011) reported that typical meteorological conditions accounted more for air
70 improvement during 2008 Beijing Olympics than emission reductions. Zhou et al.
71 (2010) concluded that control measures for transportation ~~control measures~~ resulted in
72 a reduction of 44.5% and 49.0% in daily CO and NO_x emission from motor vehicles
73 during the 2008 Olympics. Cai et al. (2011) and Wang et al. (2009a) also studied the
74 transportation controls on improving air quality during Beijing Olympic Games.
75 Okuda et al. (2011) argued that sources control during Beijing Olympics significantly
76 reduced PM₁₀, NO₂ and SO₂, but did not ~~as~~ effectively reduce PM_{2.5}. Streets et al.
77 (2007) proposed that local sources controlling is inadequate for heavily populated,
78 urbanized, and industrialized city, regional air quality management is in urgent need.
79 Lin et al. (2013) applied monitoring data to analyze the weather impacts on air quality
80 of the World Expo in YRD and concluded that high frequency of marine winds during
81 the Expo had a positive effect on the air quality of coastal cities, but a negative effect
82 on some inland cities in YRD. Satellite data reflected that the tropospheric NO₂
83 column, aerosol optical thickness (AOT), and CO concentration dropped by 8%, 14%
84 and 12%, respectively over Shanghai during the Expo period, compared to the past
85 three years (Hao et al., 2011). Liu et al. (2013) compared the contributions of
86 long-term and short-term emission control via CMAQ simulation and
87 investigatedeompared their effects on air quality in Guangzhou during the Asian
88 Games. Xu et al. (2013) concluded that PM_{2.5} was mainly emitted from anthropogenic
89 sources other than biogenic sources and indicated that cutting down anthropogenic

90 emissions could decrease PM_{2.5} effectively. Dong et al. (2013) found that independent
91 NO_x emission reduction would strengthen O₃ as a side effect in YRD. Chen et al.
92 (2015, 2017) studied the source apportionment of size-fractionated particles in
93 Nanjing, and found that ~~fugitive and~~ construction dust ~~contributes the most in coarse~~
94 ~~particles, and fugitive and construction dust~~ which decreased significantly in YOG.

95 There have been some studies on air quality during the 2nd YOG (Ding et al.,
96 2015; Chen et al., 2017; Zhou et al., 2017), but few work focused on the relative
97 contributions of meteorology and control efforts. This study takes the air quality
98 monitoring data and applies WRF/CMAQ model to estimate the effect of
99 meteorological factors and emission reduction on air quality of Nanjing during the
100 2nd YOG. Data and model descriptions as well as simulation scenarios are described
101 in Section 2. Section 3 examines the characteristics of six major air pollutants (SO₂,
102 NO₂, PM₁₀, PM_{2.5}, CO and O₃) and compares their concentrations during the YOG
103 with those a year ago (Aug., 2013) and the months without emission reduction (Jul.
104 and Sept., 2014). Besides, this section discusses the separate effect of weather
105 conditions and emission abatement qualitatively and quantitatively based on the
106 simulation results. Section 4 summaries the main conclusions, emphasizes the
107 important factor of the air quality promotion during the YOG, and provides some
108 advice for ensuring pleasant future air quality.

109

110 **2 Methodology**

111 2.1 Data description

112 Hourly observed air quality data during Jul. ~~–~~ Sept. 2014 and Aug. 2013 of two
113 representative stations was collected from Nanjing Environmental Monitoring Center
114 (<http://222.190.111.117:8023/>). Both of the two stations are state controlling air
115 monitoring sites. The data quality assurance and quality control procedures for
116 monitoring strictly follow the national standards (State Environmental Protection
117 Administration of China, 2006). Caochangmen (CCM) Station (118.75° E, 32.06° N)
118 locates in Gulou District, the city center of Nanjing. Gulou District is the center of
119 economy, politics, culture and education in Nanjing, ~~where.~~ Here gathers many East

120 China's high-end industrial and corporate headquarters. Besides, over 90% provincial
121 authorities, more than 20 colleges and universities, and more than 120 ~~research-~~
122 institutes situate in Gulou District. It's the most populated area in Nanjing, with lively
123 commercial hub and heavy traffic. Thus, CCM station was chosen to represent the
124 urban status of Nanjing. The other site calls Xianlin (XL) Station (118.92° E, 32.11°
125 N), which locates in Qixia District, the suburb of Nanjing. Compared to Gulou
126 District, Qixia District is much more sparsely populated with few traffic congestion
127 problems. Thus, XL station was chosen to represent the suburban status of Nanjing.

128


129 2.2 Model description

130 The integrated modeling system WRF/CMAQ was employed in this
131 ~~studyresearch~~. Community Multiscale Air Quality (CMAQ) is a third-generation
132 regional air quality model developed by the Environmental Protection Agency of USA
133 (USEPA). It incorporates a set of up-to-date compatible modules and control
134 equations for the atmosphere, and can fully consider complicated physical and
135 chemical processes (Byun and Schere, 2006; Foley et al., 2010). Many applications
136 have proven that CMAQ is a reliable tool in simulating air quality from city scale to
137 mesoscale (Xing et al., 2011; Dong et al., 2013; Liu et al., 2013; Xu et al. 2013; Shu
138 et al., 2016). Community Multiscale Air Quality (CMAQ v4.7.1, Binkowski and
139 Roselle, 2003) model includes the 2005 Carbon Bond gas-phase mechanism (CB05)
140 (Yarwood et al., 2005) and the fourth-generation CMAQ aerosol module (AERO4)
141 (Byun and Schere, 2006). And it was applied to simulate the pollutant distribution
142 over Nanjing in this paper. Weather Research and Forecasting (WRF) is a new
143 generation of mesoscale weather forecast model and assimilation system, developed
144 by the National Center for Atmospheric Research (NCAR). It has been widely applied
145 in China and shows a good performance in all kinds of weather forecasts (Jiang et al.,
146 2008, 2012; Xu et al., 2013; Liao et al., 2014, 2015; Xie et al., 2014, 2016; Li et al.,
147 2016; Shu et al., 2016). WRF v3.2.1 (Skamarock et al., 2008) model was run to
148 provide meteorological fields for CMAQ. Four nested domains were set for both
149 models, with horizontal resolutions of 81km, 27km, 9km, 3km, ~~andwith~~ the innermost

150 domain covering Nanjing (Fig. 1). For all domains, 23 vertical sigma layers from the
 151 surface to the top pressure of 100 hpa was set, with about 10 layers in the planetary
 152 boundary layer. The detail dynamic parameterization in WRF as well as the physical
 153 and chemical schemes of CMAQ applied in this research were the same as those in
 154 Shu et al. (2016)'s work and were proven to have good performance. As for the
 155 innermost domain, Nanjing Environmental Protection Bureau chooses the local 9 state
 156 controlling air monitoring sites (See Fig. 1, Table 1) to represent the whole Nanjing
 157 (NJ) city. In conformity with this, the 9 sites in domain_4 were chosen to represent the
 158 whole Nanjing while analyzing the impacts of weather conditions and emission
 159 reduction.

160

(a) Domain Configuration



161
 162 **Fig. 1.** Modeling domains and state controlling air monitoring sites in Nanjing. ((a) The four nested
 163 modeling domains at 81km (d01: East Asia), 27km (d02: East China), 9km (d03: Yangtze River Delta),
 164 and 3km (d04: Nanjing), (b) Locations of 9 sites in Nanjing).

165

166 **Table 1**

167 The air monitoring sites in Nanjing

Sites	Abbreviations	Location
-------	---------------	----------

Xuanwuhu Station	XWH	32.08° N, 118.80° E
Ruijinlu Station	RJL	32.03° N, 118.82° E
Zhonghuamen Station	ZHM	32.00° N, 118.76° E
Caochangmen Station	CCM	32.06° N, 118.75° E
Shanxilu Station	SXL	32.07° N, 118.77° E
Maigaoqiao Station	MGQ	32.11° N, 118.81° E
Xianlin Station	XL	32.11° N, 118.92° E
Aoti Station	AT	32.01° N, 118.74° E
Pukou Station	PK	32.07° N, 118.64° E

168


169 2.3 Emissions and simulation scenarios

170 In this study, Multi-resolution Emission Inventory for China (MEIC v1.2,
 171 <http://www.meicmodel.org/>) with a resolution of $0.25^\circ \times 0.25^\circ$ was employed to
 172 provide the anthropogenic emissions for species including SO₂, NO_x, CO, NMVOC,
 173 NH₃, CO₂, PM_{2.5}, PM₁₀, BC, and OC, which form 4 sectors as: industry, power plants,
 174 transportation, and residential.

175 For the innermost domain, the local emission inventory before and after emission
 176 control was used with a horizontal resolution of 3km × 3km. The base year of the
 177 local emission is 2012. According to the local emission control program, 5 major
 178 categories: industry, power plants, traffic, VOC product-related sources and others
 179 were in the control list. In Aug. 2014, all coal-combustion enterprises must use
 180 high-quality coals with low sulfur content less than 0.5% and ash content less than
 181 13%. Besides, the local government ordered over 100 local petrochemical, chemical
 182 and steel enterprises to cut or halt their production. Moreover, heavy pollution
 183 vehicles were prohibited in Nanjing during 10-28 Aug. 2014 to reduce traffic
 184 emission. To reduce emissions of volatile organic compounds, loading and unloading
 185 oil operations were prohibited at the docks in Nanjing section of Yangtze River.
 186 What's more, local construction work was halted during Aug. 2014. With these efforts,
 187 the emission would be cut by 25.0% for SO₂, 15.0% for NO_x, 42.8% for PM₁₀, 36.2%
 188 for PM_{2.5}, and 20.0% for CO. The spatial distributions of emission reduction for
 189 different species were showed in Fig. 2. For SO₂, NO_x, PM₁₀ and PM_{2.5}, theythe
 190 emission reduction area centered in the middle of Nanjing city and. ~~And for CO, the~~

191 ~~emission reduction for CO, it~~ centered in several points.

192



193

194 **Fig. 2.** Emission reduction in domain₄ ((a) SO₂, (b) NO_x, (c) PM₁₀, (d) PM_{2.5}, (e) CO (unit: t/month)).

195

196 The simulated period was from 27 Jul.~~27~~ to 1 Sept.~~1~~ (China standard time,
197 CST), but only the holding month (1 Aug. to 31 Aug.) was focused on for discussions.

198 In order to better understand the influence of meteorological factors and emission

199 abatement, three experiments were carried out. Exp. 1 used the weather conditions

200 during Aug. 2014 (CST) and the emission inventory after reduction. Exp. 2 used the

201 same weather conditions as Exp. 1 with the emission inventory before reduction. Exp.

202 3 used the same inventory as Exp. 2 and the weather conditions during Aug. 2013

203 (CST). Besides, Exp. 2 acted as the control experiment. Therefore, Exp. 1 and Exp. 2

204 were performed to investigate the influence of emission reduction on pollutant levelss.

205 Similarly, Exp. 2 and Exp. 3 were conducted to understand the impact of meteorology

206 on air quality.

207

208 **3 Results and discussions**


209 3.1 Observed air quality during the YOG

210 In ~~the most strictly control month~~ Aug. 2014, emission sources including 5 major

211 categories were strictly reduced, and the air quality had great promotion compared to


212 Aug. 2013. Firstly, air quality was good during the Games in accordance with
 213 China's National Ambient Air Quality Standards (NAAQS) (Ministry of
 214 Environmental Protection of the People's Republic of China, 2012) (Fig. 3, Fig. 4).
 215 The hourly mean pollutant concentrations of pollutants at the two sites during Aug.
 216 2014 are 11.6 $\mu\text{g}/\text{m}^3$ for SO_2 , 34.0 $\mu\text{g}/\text{m}^3$ for NO_2 , 57.8 $\mu\text{g}/\text{m}^3$ for PM_{10} , 39.4 $\mu\text{g}/\text{m}^3$
 217 for $\text{PM}_{2.5}$, 0.9 mg/m^3 for CO, and 38.8 $\mu\text{g}/\text{m}^3$ for O_3 . Secondly, as showed in Table 2
 218 and Table 3, the mean concentrations of the six major species dropped by 64.7% for
 219 SO_2 , 29.8% for PM_{10} , 9.8% for $\text{PM}_{2.5}$, 8.9% for CO and 31.7% for O_3 at CCM station,
 220 while 50.0% for SO_2 , 18.6% for NO_2 , 32.8% for PM_{10} , 4.1% for $\text{PM}_{2.5}$, and 31.7% for
 221 O_3 at XL station. Besides, the smaller standard deviation (std) of SO_2 , NO_2 , CO and
 222 O_3 revealed that concentrations of these air pollutants varied more steadily in Aug.
 223 2014. Actually However, the drop of pollutant concentrations could be caused mainly
 224 by meteorologically conditions or emission reduction, whichs. And we will be
 225 discussed this issue based on model simulations in Section 3.2 and Section 3.3.

226



227
 228 **Fig. 3.** Day-to-day variations in SO_2 , NO_2 , PM_{10} , $\text{PM}_{2.5}$, CO and O_3 -8h at CCM station in Aug. 2013
 229 and Aug. 2014 (CST). Observed data in Aug. 2013 and Aug. 2014 are indicated in blue and red,
 230 respectively. NAAQS are indicated in black dotted line.

231



232
 233 **Fig. 4.** Day-to-day variations in SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃-8h at XL station in Aug. 2013 and
 234 Aug. 2014 (CST). Observed data in Aug. 2013 and Aug. 2014 are indicated in blue and red,
 235 respectively. NAAQS are indicated in black dotted line.

236

237 **Table 2**

238 Statistical analysis of hourly data in Aug. 2013 and Aug. 2014 at CCM station (The unit is µg/m³
 239 except CO (mg/m³))

<u>S</u> species	<u>T</u> ime	<u>M</u> max	<u>M</u> min	<u>M</u> mean	<u>M</u> median	<u>S</u> std	Δ
SO ₂	Aug. 2013	169.0	1.0	33.7	27.0	23.7	
	Aug. 2014	72.0	2.0	11.9	10.0	7.8	-64.7%
NO ₂	Aug. 2013	111.0	1.0	35.4	32.0	19.4	
	Aug. 2014	110.0	1.0	37.3	35.0	18.6	5.0%
PM ₁₀	Aug. 2013	213.0	19.0	86.0	84.0	29.5	
	Aug. 2014	198.0	6.0	60.4	54.0	36.6	-29.8%
PM _{2.5}	Aug. 2013	123.0	10.0	45.2	43.5	16.2	
	Aug. 2014	165.0	3.0	40.7	36.0	23.8	-9.8%
CO	Aug. 2013	3.1	0.4	1.0	0.9	0.4	
	Aug. 2014	2.2	0.3	0.9	0.9	0.3	-8.9%
O ₃	Aug. 2013	198.0	1.0	56.9	42.0	46.2	
	Aug. 2014	150.0	9.0	38.9	34.0	22.6	-31.7%

240 Δ : The change percentage of species in Aug. 2014 based on Aug. 2013.

241

242 **Table 3**

243 Statistical analysis of hourly data in Aug. 2013 and Aug. 2014 at XL station (The unit is µg/m³ except
 244 CO (mg/m³))

<u>S</u> pecies	<u>T</u> ime	<u>M</u> max	<u>M</u> min	<u>M</u> mean	<u>M</u> median	<u>S</u> std	Δ
SO ₂	Aug. 2013	139.0	0.0	22.8	19.0	16.1	
	Aug. 2014	71.0	1.0	11.4	8.0	10.4	-50.0%
NO ₂	Aug. 2013	129.0	0.0	37.7	32.0	21.7	


	Aug. 2014	95.0	7.0	30.7	27.0	15.0	-18.6%
PM ₁₀	Aug. 2013	215.0	0.0	82.1	79.0	32.4	
	Aug. 2014	196.0	6.0	55.2	47.0	35.9	-32.8%
PM _{2.5}	Aug. 2013	122.0	0.0	39.7	37.5	18.9	
	Aug. 2014	157.0	3.0	38.0	34.0	24.1	-4.1%
CO	Aug. 2013	3.2	0.0	0.8	0.7	0.4	
	Aug. 2014	2.0	0.3	0.8	0.7	0.3	<0.1%
O ₃	Aug. 2013	193.0	0.0	56.6	44.0	37.5	
	Aug. 2014	148.0	2.0	38.7	32.0	28.3	-31.7%

245 | Δ : The change percentage of species in Aug. 2014 based on Aug. 2013.


246

247 | Analogously, when comparinged the observational data in Aug. 2014 with that in
248 | Jul. and Sept. 2014 (the months before and after the most aggressive abatement), the
249 | concentrations of most species also decreased obviously. As presented in Fig. 5 and
250 | Fig. 6, without abatement, NO₂, PM₁₀, PM_{2.5} and O₃ were likely to exceed NAAQS,
251 | especially for PM_{2.5} and O₃. As shown in Table 4 and Table 5, compared with Jul.
252 | 2014, the concentration of NO₂, PM₁₀, PM_{2.5}, CO and O₃ dropped by 0.7%, 31.8%,
253 | 33.7%, 1.1%, and 52.8%, respectively at CCM station in Aug. 2014, while the
254 | concentration of SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃ decreased by 15.8%, 39.6%,
255 | 34.6%, 7.1%, and 50.7%, respectively at XL station in Aug. 2014. Without emission
256 | control, the concentration of air pollutants rebounded in Sept. 2014. Compared to
257 | Aug., the concentration of SO₂, NO₂, PM₁₀, PM_{2.5} and O₃ increased by 37.4%, 19.8%,
258 | 37.6%, 22.3%, and 47.2%, respectively at CCM station in Sept. 2014 (Table 4), while
259 | the concentration of SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃ increased by 24.6%, 21.8%,
260 | 28.7%, 17.7%, 4.9%, and 39.9%, respectively at XL station in Sept. 2014 (Table 5). In
261 | additionBesides, for most species, the standard deviation was the lowest in Aug.,
262 | which meant that the potential of extreme events was the least in Aug.. Assume that
263 | the weather conditions in Jul., Aug., Sept., 2014 were nearly similar, it can be
264 | estimated that emission sources could be the major impact factor of explaining the
265 | concentration changes during the three months. These results proved that
266 | concentrations of most species decreased and had less potential in extreme events
267 | after aggressive emission abatement. However, the concentration became higher

268 without emission control. Section 3.3 would further discuss the change of pollutant
 269 concentration with and without emission reduction based on model simulation.
 270



271
 272 **Fig. 5.** Day-to-day variations in SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃-8h at CCM station in Jul., Aug.
 273 and Sept. 2014 (CST). Observed data in Jul., Aug. and Sept. 2014 are indicated in green, red and blue,
 274 respectively. NAAQS are indicated in black dotted line.
 275



276
 277 **Fig. 6.** Day-to-day variations in SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃-8h at XL station in Jul., Aug. and
 278 Sept. 2014 (CST). Observed data in Jul., Aug. and Sept. 2014 are indicated in green, red and blue,
 279 respectively. NAAQS are indicated in black dotted line.

280

281 **Table 4**

282 Statistical analysis of hourly data in Jul. - Sept. 2014 at CCM station (The unit is µg/m³ except CO
 283 (mg/m³))

Species	M _{month}	M _{max}	M _{min}	M _{mean}	M _{median}	S _{std}	Δa	Δb
---------	--------------------	------------------	------------------	-------------------	---------------------	------------------	----	----

	Jul. 2014	83.0	1.0	11.3	9.0	9.8		
SO ₂	Aug. 2014	72.0	2.0	11.9	10.0	7.8	5.1%	-37.4%
	Sept. 2014	70.0	4.0	19.0	18.0	9.9		
	Jul.-Sept. 2014	83.0	1.0	14.0	12.0	9.8		
	Jul. 2014	161.0	1.0	37.5	32.0	28.3		
NO ₂	Aug. 2014	110.0	1.0	37.3	35.0	18.6	-0.7%	-19.8%
	Sept. 2014	151.0	8.0	46.5	42.0	24.5		
	Jul.-Sept. 2014	161.0	1.0	40.2	37.0	24.4		
	Jul. 2014	255.0	6.0	88.5	88.0	50.7		
PM ₁₀	Aug. 2014	198.0	6.0	60.4	54.0	36.6	-31.8%	-37.6%
	Sept. 2014	243.0	6.0	96.7	90.0	45.8		
	Jul.-Sept. 2014	255.0	6.0	81.7	76.0	47.4		
	Jul. 2014	171.0	1.0	61.5	58.0	33.9		
PM _{2.5}	Aug. 2014	165.0	3.0	40.7	36.0	23.8	-33.7%	-22.3%
	Sept. 2014	143.0	3.0	52.4	46.0	27.2		
	Jul.-Sept. 2014	171.0	1.0	51.5	45.0	29.9		
	Jul. 2014	2.7	0.2	0.9	0.9	0.3		
CO	Aug. 2014	2.2	0.3	0.9	0.9	0.3	-1.1%	21.1%
	Sept. 2014	2.1	0.1	0.8	0.7	0.4		
	Jul.-Sept. 2014	2.7	0.1	0.9	0.8	0.4		
	Jul. 2014	281.0	4.0	82.4	69.0	57.6		
O ₃	Aug. 2014	150.0	9.0	38.9	34.0	22.6	-52.8%	-47.2%
	Sept. 2014	240.0	6.0	73.6	61.0	49.2		
	Jul.-Sept. 2014	281.0	4.0	64.7	51.0	49.3		

284 | Δa: The change percentage of species in Aug.2014 based on Jul. 2014.

285 | Δb: The change percentage of species in Aug. 2014 based on Sept. 2014.

286

287 | **Table 5**

288 | Statistical analysis of hourly data in Jul. - Sept.2014 at XL station (The unit is μg/m³ except CO

289 | (mg/m³))

<u>S</u> pecies	<u>M</u> month	<u>M</u> max	<u>M</u> min	<u>M</u> mean	<u>M</u> median	<u>S</u> std	Δa	Δb
	Jul. 2014	61.0	1.0	14.5	12.0	10.3		
SO ₂	Aug. 2014	71.0	1.0	11.4	8.0	10.4	-21.2%	-24.6%
	Sept. 2014	75.0	1.0	15.1	14.0	10.3		
	Jul.-Sept. 2014	75.0	1.0	13.7	11.0	10.4		
	Jul. 2014	123.0	9.0	36.4	33.0	18.9		
NO ₂	Aug. 2014	95.0	7.0	30.7	27.0	15.0	-15.8%	-21.8%
	Sept. 2014	127.0	11.0	39.2	36.0	18.7		
	Jul.-Sept. 2014	127.0	7.0	35.4	32.0	18.0		
	Jul. 2014	300.0	4.0	91.3	85.0	48.9		

PM ₁₀	Aug. 2014	196.0	6.0	55.2	47.0	35.9	-39.6%	-28.7%
	Sept. 2014	226.0	9.0	77.3	70.0	40.3		
	Jul.-Sept. 2014	300.0	4.0	74.5	64.0	44.6		
	Jul. 2014	158.0	2.0	58.2	51.0	34.8		
PM _{2.5}	Aug. 2014	157.0	3.0	38.0	34.0	24.1	-34.6%	-17.7%
	Sept. 2014	144.0	3.0	46.2	38.0	29.0		
	Jul.-Sept. 2014	158.0	2.0	47.4	40.5	30.7		
	Jul. 2014	2.0	0.3	0.8	0.8	0.4		
CO	Aug. 2014	2.0	0.3	0.8	0.7	0.3	-7.1%	-4.9%
	Sept. 2014	2.8	0.3	0.8	0.7	0.4		
	Jul.-Sept. 2014	2.8	0.3	0.8	0.7	0.4		
	Jul. 2014	238.0	2.0	78.4	67.0	55.6		
O ₃	Aug. 2014	148.0	2.0	38.7	32.0	28.3	-50.7%	-39.9%
	Sept. 2014	226.0	2.0	64.4	54.0	46.4		
	Jul.-Sept. 2014	238.0	2.0	60.3	48.0	47.7		

290 Δa : [T](#)he change percentage of species in Aug.2014 based on Jul. 2014.

291 Δb : [T](#)he change percentage of species in Aug. 2014 based on Sept. 2014.

292

293 3.2 Simulated impact of meteorological conditions


294 In this paper, the model configurations were the same as those set by Shu et al.
 295 (2016), who has evaluated the model performance of WRF/CMAQ and proved the
 296 model's reliability in simulating air quality in Nanjing.

297 As we know, meteorology is an important impact factor on air quality. Good
 298 diffusion conditions can alleviate air pollution in the short term (Cermak and Knutti,
 299 2009; Wang et al., 2009b). In this premise, if two experiments (Exp. 2 and Exp. 3) use
 300 the same emission inventory but different weather conditions, it can be concluded that
 301 the higher concentrations may result from poor meteorological conditions. According
 302 to model simulation, Exp. 2 exhibited higher pollutant concentrations for all species in
 303 most part of Nanjing as shown in Fig. 7. For SO₂, NO₂, PM₁₀, PM_{2.5}, CO, and O₃,
 304 their concentrations were increased by 17.5%, 16.9%, 18.5%, 18.8%, 7.8% and 0.8%
 305 during Aug. 2014 compared to Aug. 2013. [Additionally Besides](#), the contributions of
 306 meteorological conditions to primary and secondary particulate matters differed (See
 307 Fig. 8). Secondary PM₁₀ (PM_{10s}) was raised by 21.5%, while primary PM₁₀ (PM_{10p})
 308 rose by 12.6% during Aug. 2014 compared to Aug. 2013. And secondary PM_{2.5}


309 (PM_{2.5s}) was increased by 21.5%, while primary PM_{2.5} (PM_{2.5p}) was ~~enhanced~~ by
310 9.5%. Thus, the weather conditions had a slightly greater impact on secondary fine
311 particulate matters. Moreover, for SO₂, NO₂, PM₁₀, PM_{2.5}, CO, PM_{10p}, PM_{10s}, PM_{2.5p},
312 and PM_{2.5s}, there were some small decreasing areas in the northeast Nanjing (Fig. 7
313 and Fig. 8). In domain 4, the simulated predominant wind was northeast wind in Aug.
314 2014, while that was southeast wind in Aug. 2013. So, the diffusion condition of
315 northeast Nanjing might be better in Aug. 2014 and resulted in small decrease in these
316 areas.

317 The overall increasing pollutant levels in Aug. 2014 suggested that the diffusion
318 conditions in Aug. 2014 were worse than those in Aug. 2013. Focus on the weather
319 conditions during the YOG holding period (16-28 Aug., 2014) and the same period in
320 2013, the simulated hourly mean 10-m wind speed in Nanjing was 1.5 m/s larger in
321 16-28 Aug., 2013, ~~and it was 1.5 m/s larger~~ than that of 16-28 Aug., 2014 (Fig. 9).
322 Also, the simulated 2-m temperature was 2.0 K higher in 16-28 Aug., 2013, ~~and it was~~
323 ~~2.0 K larger~~ than that of 16-28 Aug., 2014 (Fig. 9). ~~Also~~Besides, the simulated
324 planetary boundary layer height (PBLH) was higher in Aug. 2013, especially in 16-28
325 Aug., and it was 27.5 m higher than that of 16-28 Aug., 2014 (Fig. 9). Larger wind
326 speed and higher PBLH benefited the diffusion of air pollutants. Warming on the
327 ground surface was conducive to the promotion of convective instability and was also
328 good for the vertical dilution and diffusion of pollutant. Thus, the diffusion conditions
329 in 16-28 Aug. 2013 were better than those in 16-28 Aug. 2014. Rather worse
330 meteorological conditions in 16-28 Aug. 2014 implied that abatement controls might
331 play a more important role in improving air quality in YOG compared with the same
332 period in 2013. What's more, relative humidity, cloud cover and shortwave solar
333 radiation all affect ozone chemical reaction (Katragkou et al., 2011; Pu et al., 2017).
334 The generation and photochemical reaction of surface ozone depends on the
335 availability of solar radiation. During the heat wave period, less relative humidity
336 leads to less cloud cover, which could result in more net downward shortwave solar
337 radiation and more production of O₃ (Pu et al., 2017). For ordinary O₃, its production
338 also corresponded well to the cloud cover. As shown in Fig. 9, more relative humidity


339 resulted in more cloud cover in northern and eastern Nanjing during 16-28 Aug., 2013,
 340 which resulted in less O₃ in Aug. 2013, but more O₃ in Aug. 2014 (Fig. 7).
 341



342
 343 **Fig. 7.** Influence of meteorology on hourly mean concentrations of air pollutants in Aug. 2014
 344 compared with Aug. 2013. (Black thick lines draw the outline of Nanjing. Picture a - f are hourly
 345 average values of impact percentage ($d_{species}(\%) = (Exp. 2 - Exp. 3)/Exp. 2 * 100\%$) of SO₂, NO₂,
 346 PM₁₀, PM_{2.5}, CO, and O₃, respectively.)
 347



348
 349 **Fig. 8.** Influence of meteorology on hourly mean concentrations of primary and secondary particulate
 350 matters in Aug. 2014 compared with Aug. 2013. (Black thick lines draw the outline of Nanjing. Picture
 351 a - d are hourly average values of impact percentage ($d_{species}(\%) = (Exp. 2 - Exp. 3)/Exp. 2 * 100\%$)
 352 of PM_{10p}, PM_{10s}, PM_{2.5p}, and PM_{2.5s}, respectively.)
 353



354

355 **Fig. 9.** Bias of simulated hourly mean meteorological conditions during the YOG. (Bias =
 356 Meteorological Factors in 16-28 Aug., 2013 - Meteorological Factors in 16-28 Aug., 2014. (a) Bias of
 357 Wind at 10m during 16-28 Aug. (unit: m/s), (b) Bias of temperature at 2m during 16-28 Aug. (unit: K),
 358 (c) Bias of planetary boundary layer height during 16-28 Aug. (unit: m), (d) Bias of relative humidity at
 359 2m during 16-28 Aug. (unit: %), (e) Bias of cloud fraction during 16-28 Aug.).


360

361 3.3 Simulated impact of emission reduction

362 As for SO₂, NO₂, PM₁₀, PM_{2.5}, and CO, the distributions of such short-lived
 363 chemical compositions are largely affected by their sources and sinks. As seen in Fig.
 364 10, the simulated spatial distributions of concentration changes were uneven, large
 365 variations were found in the west of Nanjing corresponding to the downwind regions
 366 of heavy reduction districts (See Fig. 2). ~~—Besides, i~~ Impact percentages (dspecies (%)
 367 = (Exp. 1 - Exp. 2)/ Exp. 2*100%) of species were negative except O₃, implying that
 368 emission regulatory efforts were effective on the other species, but counterproductive
 369 to O₃. Statistically, the concentrations of SO₂, NO₂, PM₁₀, PM_{2.5}, and CO in Nanjing
 370 were reduced by 24.6%, 12.1%, 15.1%, 8.1% and 7.2% during Aug. 2014. As for O₃,
 371 the variation was positive (1.3%), especially in the downwind area of NO_x with heavy
 372 reduction, which might due to the less titration of O₃ by NO_x (Liu et al., 2013; Dong
 373 et al., 2013). For primary and secondary particulate matters, the influence of emission
 374 reduction varied dramatically. As shown in Fig. 11, PM_{10p} was dropped by 39.6%,
 375 while PM_{10s} only declined by 2.9%. And PM_{2.5p} was decreased by 26.2%, while
 376 PM_{2.5s} merely cut down by 2.9%. It seems that emission controls had much more

377 impacts on primary pollutants, especially on coarse particulate matters.

378



379


380 **Fig. 10.** Influence of emission reduction on hourly mean concentrations of air pollutants in Aug. 2014.

381 (Black thick lines draw the outline of Nanjing. Picture a - f are hourly average values of impact

382 percentage ($\text{dspecies} (\%) = (\text{Exp. 1} - \text{Exp. 2}) / \text{Exp. 2} * 100\%$) of SO_2 , NO_2 , PM_{10} , $\text{PM}_{2.5}$, CO and O_3 ,

383 respectively.).

384



385

386 **Fig. 11.** Influence of emission reduction on hourly mean concentrations of primary and secondary

387 particulate matters in Aug. 2014. (Black thick lines draw the outline of Nanjing. Picture a - d are hourly

388 average values of impact percentage ($\text{dspecies} (\%) = (\text{Exp. 1} - \text{Exp. 2}) / \text{Exp. 2} * 100\%$) of PM_{10p} , PM_{10s} ,

389 $\text{PM}_{2.5p}$ and $\text{PM}_{2.5s}$, respectively.).

390

391 3.4 Comparison of simulated meteorological factors and emission reduction


392 Fig. 12 displays the simulated effect of meteorological factors and emission

393 reduction in Nanjing on air quality improvement during Aug. 2014. In general,

394 meteorological conditions played a negative role in air quality promotion in most days,

395 only played a positive role in a few days. ~~And the negative effect of weather~~

396 ~~conditions exceeded its positive effect during the whole month~~ (See discussion in
 397 Section 3.2). On the other hand, emission reduction contributed to the decline of SO₂,
 398 NO₂, PM₁₀, PM_{2.5}, CO, PM_{10p}, PM_{10s}, PM_{2.5p}, and PM_{2.5s} all the time, especially for
 399 primary coarse particulate matters. However, reduction of NO_x could cause a slight
 400 rise of O₃.
 401



402
 403 **Fig. 12.** The simulated effect of meteorology and emission reduction on pollutant concentrations in
 404 Nanjing during 1-31 Aug. , 2014, Met. (Exp. 2-Exp. 3) represents the effect of meteorology, while Red.
 405 (Exp. 1-Exp. 2) represents the simulated effect of emission -reduction.

406
 407 ~~Moreover,~~ the opposite effects of meteorology and emission abatement on SO₂,
 408 NO₂, PM₁₀, PM_{2.5}, CO, PM_{10p}, PM_{10s}, PM_{2.5p} and PM_{2.5s} during the whole month were
 409 significantmore apparent as statistically listed in Table 6. CCM station represents the
 410 urban status, XL station represents the suburban status and NJ represents the whole
 411 city. Adverse meteorological conditiony was found to raise the concentrations of the
 412 six pollutants as 17.4% for SO₂, 15.1% for NO₂, 15.6% for PM₁₀, 14.9% for PM_{2.5},
 413 6.4% for CO and 0.9% for O₃ at CCM station, and 14.1% for SO₂, 12.4% for NO₂,
 414 22.4% for PM₁₀, 24.5% for PM_{2.5}, 2.3% for CO, and 1.6% for O₃ at XL station. On the

415 contrary, emission abatement reduced their levels in most cases, especially in the
 416 urban site. It seems that the levels of air pollutants reduced more at CCM station
 417 compared to XL station. Emission abatement independently led to a 24.3% decrease
 418 in SO₂ at CCM station, which was 5.1% higher ~~compared to~~~~than that at~~ XL station.
 419 Moreover, the cutbacks of NO₂, PM₁₀, PM_{2.5} and CO were 11.7%, 13.9%, 7.5% and
 420 7.0%, respectively at CCM station, being larger by 1.0% to 2.0% compared with XL
 421 station. Though O₃ under emission reduction scenarios resulted in a slightly rise
 422 (0.9% to 1.3%) at both sites, the effectiveness of emission abatement was remarkable
 423 generally. As for primary and secondary particulate matters, meteorological factors
 424 also played a negative role during Aug. 2014, and had slightly more effect on
 425 secondary fine particles. Emission controls seemed to cause tremendous impact on
 426 primary particulate matters, especially for primary coarse particles. Emission
 427 abatement led to a 38.3% decrease in PM_{10p} at CCM station, a 33.2% decrease in
 428 PM_{10p} at XL station, and a 39.6% decrease in PM_{10p} for the whole city. For secondary
 429 particulate matters, including sulfate, nitrate, and ammonium salt, emission reduction
 430 played rather minor role in cutting the pollutants, with merely a 2.4% decrease in
 431 PM_{10s} and PM_{2.5s} at CCM station, a 2.9% decrease in PM_{10s} and PM_{2.5s} at XL station.

432

433 **Table 6**

434 Comparison between the simulated effect of meteorology and emission reduction at CCM and XL
 435 station

Species	Met. (CCM)	Red. (CCM)	Met. (XL)	Red. (XL)	Met. (NJ)	Red. (NJ)
SO ₂	17.4%	-24.3%	14.1%	-19.2%	17.5%	-24.6%
NO ₂	15.1%	-11.7%	12.4%	-10.2%	16.9%	-12.1%
PM ₁₀	15.6%	-13.9%	22.4%	-11.9%	18.5%	-15.1%
PM _{2.5}	14.9%	-7.5%	24.5%	-6.3%	18.8%	-8.1%
CO	6.4%	-7.0%	2.3%	-5.5%	7.8%	-7.2%
O ₃	0.9%	1.3%	1.6%	0.9%	0.7%	1.5%
PM _{10p}	13.2%	-38.3%	5.9%	-33.2%	12.6%	-39.6%
PM _{10s}	16.7%	-2.4%	29.4%	-2.9%	21.5%	-2.9%
PM _{2.5p}	8.4%	-25.8%	4.9%	-20.1%	9.5%	-26.2%
PM _{2.5s}	16.7%	-2.4%	29.4%	-2.9%	21.5%	-2.9%

436 Met.: [T](#)he change percentage of species in Exp. 2 based on Exp3, represents the effect of meteorology.

437 Red.: [T](#)he change percentage of species in Exp. 1 based on Exp 2, represents the effect of Nanjing

438 local emission reduction.

439

440 The decrease of SO₂ might ~~result from~~due to the limit and halt of power plants
441 and improvement of coal-combustion. ~~T~~Besides, the prohibition of heavy pollution
442 vehicles could contribute to the drop of NO₂ and CO. Also, limiting the production of
443 industries helped to reduce NO₂ and CO. The cut of particulate matters might ~~be~~due
444 attributable to the stop of construction process and use of low ash content coal for
445 industry. For secondary particulate matters, controlling the emission of SO₂ and NO_x
446 could help to reduce the formation of sulfate and nitrate, but no control on the
447 emission of NH₃ could still result in quite a part of ammonium salt. The response of
448 O₃ to emission control could be due to its non-linearity chemistry (Fu et al., 2012),
449 reducing NO₂ ~~pollution~~ may have side-effect by increasing O₃ because of the titration
450 effect. On the whole, the meteorological factors and emission reduction during the
451 YOG had opposite effects on SO₂, NO₂, PM₁₀, PM_{2.5}, and CO, and emission reduction
452 indeed played a very important role in the air quality improvement.

453

454 **4 Summary and conclusions**

455 The air quality during the 2nd YOG was superior according to the current
456 NAAQS. Both observation and modeling confirmed that stringent emission reductions
457 were effective to ambient air quality promotion during the Youth Olympic Games,
458 especially to SO₂, NO₂, PM₁₀, PM_{2.5} and CO. Compared to Aug. 2013, the observed
459 concentrations in Aug. 2014 were dropped by 64.7% for SO₂, 29.8% for PM₁₀, 9.8%
460 for PM_{2.5}, 8.9% for CO and 31.7% for O₃ at CCM station, while 50.0% for SO₂,
461 18.6% for NO₂, 32.8% for PM₁₀, 4.1% for PM_{2.5}, and 31.7% for O₃ at XL station. The
462 simulated impact percentage of emission reductions were -24.6%, -12.1%, -15.1%,
463 -8.1% and -7.2% for SO₂, NO₂, PM₁₀, PM_{2.5}, and CO, respectively.

464 The meteorological conditions in the holding time of the YOG were inferior to
465 those of the same period in 2013, with lower temperature and weaker winds. Model
466 simulations show that unless—favorable weather conditions caused higher
467 concentrations for all species, including O₃ which was increased due to less cloud

468 cover. Besides, meteorological conditions had slightly more effect on secondary fine
469 particular matters compared to primary fine particular matters. Emission reduction
470 could cut down the levels of SO₂, NO₂, CO and particular matters, especially for
471 primary coarse particles. Thus, emission reduction control is the very important factor
472 for the air quality improvement during the YOG.

473 In general, better air quality during the YOG benefit a lot from emission
474 reduction, which has set up a good example in air protection for important social
475 events. However, the enhanced concentrations of air pollutants after the YOG (in Sept.
476 2014) suggest that short-term emission control can only ease air pollution effectively
477 and temporarily. Long-term control policies are necessary to ensure pleasant future air
478 quality.

479

480 **5 Data availability**

481 [The air quality monitoring records are available at http://222.190.111.117:8023/.](http://222.190.111.117:8023/)
482 [The Multi-resolution Emission Inventory for China \(MEIC\) is available at](http://www.meicmodel.org/)
483 [http://www.meicmodel.org/.](http://www.meicmodel.org/) [The WRF model is available at](http://www2.mmm.ucar.edu/wrf/users/downloads.html)
484 [http://www2.mmm.ucar.edu/wrf/users/downloads.html.](http://www2.mmm.ucar.edu/wrf/users/downloads.html) [The CMAQ model is](https://www.cmascenter.org/cmaq/)
485 [available at https://www.cmascenter.org/cmaq/.](https://www.cmascenter.org/cmaq/)

486

487 [*Competing interests.* The authors declare that they have no conflict of interest.](#)

488

489 **Acknowledgements**

490 This work was supported by the National Natural Science Foundation of China
491 (~~91544230~~, 41575145, 41621005, [91544230](#)), the National Key Research
492 Development Program of China (2016YFC0203303, 2016YFC0208504,
493 2014CB441203), the National Special Fund for the Weather Industry
494 (GYHY201206011) and the National Special Fund for the Environmental Protection
495 Industry (GYHY201409008) . We are grateful to Prof. Yu Zhao from School of
496 Environment of Nanjing University for supply the emission data of Nanjing. The
497 contents of this paper are solely the responsibility of the authors and do not

498 necessarily represent the official views of sponsors.

499

500 **References**

501 Binkowski, F. S. and Roselle, S. J.: Models-3 Community Multiscale Air Quality
502 (CMAQ) model aerosol component 1. Model description, *J. Geophys. Res.*, 108,
503 4183, doi:10.1029/2001JD001409, 2003.

504 Byun, D. and Schere, K. L.: Review of the governing equations, computational
505 algorithms, and other components of the models-3 Community Multiscale Air
506 Quality (CMAQ) modeling system, *Appl. Mech. Rev.*, 59, 51-77,
507 doi:10.1115/1.2128636, 2006.

508 Cai, H. and Xie, S.: Traffic-related air pollution modeling during the 2008 Beijing
509 Olympic Games: the effects of an odd-even day traffic restriction scheme,
510 *Science of the total environment*, 409, 1935-1948, doi:
511 10.1016/j.scitotenv.2011.01.025, 2011.

512 Center for Earth System Science, Tsinghua University: Multi-resolution Emission
513 Inventory for China, <http://www.meicmodel.org/>, 2015.

514 Cermak, J. and Knutti, R.: Beijing Olympics as an aerosol field experiment,
515 *Geophysical Research Letters*, 36, 1-5, doi:10.1029/2009GL038572, 2009.

516 Chen P., Wang T., and Hu X.: Chemical Mass Balance Source Apportionment of
517 Size-Fractionated Particulate Matter in Nanjing, China, *Aerosol & Air Quality*
518 *Research*, 15, 1855-1867. doi:10.4209/aaqr.2015.03.0172, 2015.

519 Chen, P., Wang, T., Lu, X., Yu, Y., Kasoar, M., Xie, M., and Zhuang, B.: Source
520 apportionment of size-fractionated particles during the 2013 Asian Youth Games
521 and the 2014 Youth Olympic Games in Nanjing, China, *Science of the Total*
522 *Environment*. 579, 860-870. <http://dx.doi.org/10.1016/j.scitotenv.2016.11.014>,
523 2017.

524 Ding, J., Van, d. A. R. J., Mijling, B., Levelt, P. F., and Hao, N.: NO_x emission
525 estimates during the 2014 Youth Olympic Games in Nanjing, *Atmos. Chem.*
526 *Phys.*, 15, 9399-9412, doi: 10.5194/acp-15-9399-2015, 2015.

527 Dong, X. Y., Gao, Y., Fu, J. S., Li, J., Huang, K., Zhuang, G. S., and Zhou, Y.: Probe

528 into gaseous pollution and assessment of air quality benefit under sector
529 dependent emission control strategies over megacities in Yangtze River Delta,
530 China, *Atmos. Environ.*, 79, 841-852, doi:10.1016/j.atmosenv.2013.07.041,
531 2013.

532 Foley, K. M., Roselle, S. J., Appel, K. W., Bhawe, P. V., Pleim, J. E., Otte, T. L.,
533 Mathur, R., Sarwar, G., Young, J. O., Gilliam, R. C., Nolte, C. G., Kelly, J. T.,
534 Gilliland, A. B., and Bash, J. O.: Incremental testing of the Community
535 Multiscale Air Quality (CMAQ) modeling system version 4.7, *Geosci. Model*
536 *Dev.*, 3, 205-226, doi:10.5194/gmd-3-205-2010, 2010.

537 Fu, J. S., Dong, X. Y., Gao, Y., Wong, D. C., and Lam, Y. F.: Sensitivity and linearity
538 analysis of ozone in East Asia: the effects of domestic emission and
539 intercontinental transport, *Journal of the Air and Waste Management Association*,
540 62, 1102-1114, doi:10.1080/10962247.2012.699014, 2012.

541 Hao, N., Valks, P., Loyola, D., Cheng, Y. F., and Zimmer, W.: Space-based
542 measurements of air quality during the World Expo 2010 in Shanghai,
543 *Environmental Research Letters*, 6, doi:10.1088/1748-9326/6/4/044004, 2011.

544 Jiang, F., Wang, T. J., Wang, T. T., Xie, M., and Zhao, H.: Numerical modeling of a
545 continuous photochemical pollution episode in Hong Kong using WRF-chem,
546 *Atmos. Environ.*, 42, 8717-8727, doi:10.1016/j.atmosenv.2008.08.034, 2008.

547 Jiang, F., Zhou, P., Liu, Q., Wang, T. J., Zhuang, B. L., and Wang, X. Y.: Modeling
548 tropospheric ozone formation over East China in springtime, *J. Atmos. Chem.*,
549 69, 303-319, doi:10.1007/s10874-012-9244-3, 2012.

550 Katragkou E., Zanis P., Kioutsioukis I., Tegoulas I., Melas D., Krüger, B. C., and
551 Coppola E.: Future climate change impacts on summer surface ozone from
552 regional climate-air quality simulations over Europe, *Journal of Geophysical*
553 *Research Atmospheres*, 116, D22307, doi:10.1029/2011JD015899, 2011.

554 Li, M. M., Song, Y., Mao, Z. C., Liu, M. X., and Huang, X.: Impacts of thermal
555 circulations induced by urbanization on ozone formation in the Pearl River Delta
556 region, China, *Atmos. Environ.*, 127, 382-392,
557 doi:10.1016/j.atmosenv.2015.10.075, 2016.

558 Liao, J. B., Wang, T. J., Wang, X. M., Xie, M., Jiang, Z. Q., Huang, X. X., and Zhu, J.
559 L.: Impacts of different urban canopy schemes in WRF/Chem on regional
560 climate and air quality in Yangtze River Delta, China, *Atmos. Res.*, 145, 226-243,
561 doi:10.1016/j.atmosres.2014.04.005, 2014.

562 Liao, J. B., Wang, T. J., Jiang, Z. Q., Zhuang, B. L., Xie, M., Yin, C. Q., Wang, X. M.,
563 Zhu, J. L., Fu, Y., and Zhang, Y.: WRF/Chem modeling of the impacts of urban
564 expansion on regional climate and air pollutants in Yangtze River Delta, China,
565 *Atmos. Environ.*, 106, 204-214, doi:10.1016/j.atmosenv.2015.01.059, 2015.

566 Lin Y., Huang, K., Zhuang, G., Fu, J. S., Xu, C., Shen, J., and Chen, S.: Air Quality
567 over the Yangtze River Delta during the 2010 Shanghai Expo, *Aerosol & Air
568 Quality Research*, 13, 1655-1666, doi: 10.4209/aaqr.2012.11.0312, 2013.

569 Liu, H., Wang, X. M., Zhang, J. P., He, K. B., Wu, Y., and Xu, J. Y.: Emission controls
570 and changes in air quality in Guangzhou during the Asian Games, *Atmos.
571 Environ.*, 76, 81-93, doi:10.1016/j.atmosenv.2012.08.004, 2013.

572 Ministry of Environmental Protection of the People's Republic of China: The second
573 Summer Youth Olympic Games environmental quality assurance work plan,
574 Beijing, http://www.js.xinhuanet.com/2014-07/19/c_1111695515_2.htm/, 2014.

575 National Bureau of Statistics of China: <http://www.stats.gov.cn/>, 2014.

576 Okuda, T., Matsuura, S., Yamaguchi, D., Umemura, T., Hanada, E., Orihara, H.,
577 Tanaka, S., He, K.B., Ma, Y. L., Cheng, Y., and Liang, L. L.: The impact of the
578 pollution control measures for the 2008 Beijing Olympic Games on the chemical
579 composition of aerosols, *Atmos. Environ.*, 45, 2789-2794,
580 doi:10.1016/j.atmosenv.2011.01.053, 2011.

581 Pu X., Wang T. J., Huang X., Melas D., Zanis P., Papanastasiou D. K., and Poupkou
582 A.: Enhanced surface ozone during the heat wave of 2013 in Yangtze River Delta
583 region, China. *Science of the Total Environment*, 603-604, 807-816,
584 doi:10.1016/j.scitotenv.2017.03.056, 2017.

585 Shu L., Xie M., Wang T. J., Gao D., Chen P. L., Han Y., Li S., Zhuang B. L., Li M. M.:
586 Integrated studies of a regional ozone pollution synthetically affected by subtrop
587 ical high and typhoon system in the Yangtze River Delta region, China. *Atmos.*

588 Chem. Phys., 16, 15801-15819. doi:10.5194/acp-16-15801-2016, 2016.

589 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G.,
590 Huang, X. Y., Wang, W., and Powers, J. G.: A Description of the Advanced
591 Research WRF Version 3. NCAR Tech Notes-475+STR, 2008.

592 State Environmental Protection Administration of China: China National
593 Environmental Protection Standard: Automated Methods for Ambient Air
594 Quality Monitoring, China Environmental Science Press, Beijing, 2006.

595 Streets, D. G., Fu, J. S., Jang, C. J., Hao, J. M., He, K. B., Tang, X. Y., Zhang, Y. H.,
596 Wang, Z. F., Li, Z. P., Zhang, Q., Wang, L. T., Wang, B. Y., and Yu, C.: Air
597 quality during the 2008 Beijing Olympic Games, *Atmos. Environ.*, 41, 480-492,
598 doi:10.1016/j.atmosenv.2006.08.046, 2007.

599 Wang, X., Westerdahl, D., Chen, L. C., Wu, Y., Hao, J. M., Pan, X. C., Guo, X. B.,
600 and Zhang, K. M.: Evaluating the air quality impacts of the 2008 Beijing
601 Olympic Games: On-road emission factors and black carbon profiles, *Atmos.*
602 *Environ.* 43, 4535-4543, doi:10.1016/j.atmosenv.2009.06.054, 2009a.

603 Wang, W. T., Primbs, T., Tao, S., Zhu, T., and Simonich, S. L.: Atmospheric
604 particulate matter pollution during the 2008 Beijing Olympics, *Environmental*
605 *Science & Technology*, 43, 5314-5320, 2009b.

606 Wang, T., Nie, W., Gao, J., Xue, L. K., Gao, X. M., Wang, X. F., Qiu, J., Poon, C. N.,
607 Meinardi, S., Blake, D., Wang, S. L., Ding, A. J., Chai, F. H., Zhang, Q. Z., and
608 Wang, W. X.: Air quality during the 2008 Beijing Olympics: secondary
609 pollutants and regional impact, *Atmos. Chem. Phys.*, 10, 7603-7615,
610 doi:10.5194/acp-10-7603-2010, 2010.

611 Xie, M., Zhu, K. G., Wang, T. J., Yang, H. M., Zhuang, B. L., Li, S., Li, M. G., Zhu, X.
612 S., and Ouyang, Y.: Application of photochemical indicators to evaluate ozone
613 nonlinear chemistry and pollution control countermeasure in China, *Atmos.*
614 *Environ.*, 99, 466-473, doi:10.1016/j.atmosenv.2014.10.013, 2014.

615 Xie, M., Liao, J., Wang, T., Zhu, K., Zhuang, B., Han, Y., Li, M., and Li, S.: Modeling
616 of the anthropogenic heat flux and its effect on regional meteorology and air
617 quality over the Yangtze River Delta region, China, *Atmos. Chem. Phys.*, 16,

618 6071-6089, doi:10.5194/acp-16-6071-2016, 2016.

619 Xing, J., Zhang, Y., Wang, S. X., Liu, X. H., Cheng, S. H., Zhang, Q., Chen, Y. S.,
620 Streets, D. G., Jang, G., Hao, J. M., and Wang, W. X.: Modeling study on the air
621 quality impacts from emission reductions and atypical meteorological conditions
622 during the 2008 Beijing Olympics, *Atmos. Environ.*, 45, 1786-1798,
623 doi:10.1016/j.atmosenv.2011.01.025, 2011.

624 Xu, H. M., Tao, J., Ho, S. S. H., Ho, K. F., Cao, J. J., Li, N., Chow, J. C., Wang, G. H.,
625 Han, Y. M., Zhang, R. J., Watson, J. G., Zhang, J. Q.: Characteristics of fine
626 particulate non-polar organic compounds in Guangzhou during the 16th Asian
627 Games: Effectiveness of air pollution controls, *Atmos. Environ.*, 76, 94-101,
628 doi:10.1016/j.atmosenv.2012.12.037, 2013.

629 Yarwood, G., Rao, S., Yocke, M., and Whitten G.: Updates to the Carbon Bond
630 chemical mechanism: CB05, Final Report to the US EPA, RT-0400675, 2005.

631 Zhou, D., Li, B., Huang, X., Virkkula, A., Wu, H., Zhao, Q., Zhang, J., Liu, Q., Li, L.,
632 Li, C., Chen, F., Yuan, S., Qiao, Y., Shen, G., and Ding, A.: The Impacts of
633 Emission Control and Regional Transport on PM_{2.5} Ions and Carbon Components
634 in Nanjing during the 2014 Nanjing Youth Olympic Games, *Aerosol & Air
635 Quality Research*, 17, 730-740, doi: 10.4209/aaqr.2016.03.0131, 2017.

636 Zhou, Y., Wu, W., Yang, L., Fu, L. X., He, K. B., Wang, S. X., Hao, J. M., Chen, J. C.,
637 Li, C. Y.: The impact of transportation control measures on emission reductions
638 during the 2008 Olympic Games in Beijing, China, *Atmos. Environ.*, 44,
639 285-293, doi:10.1016/j.atmosenv.2009.10.040, 2010.