

1 **Impacts of emission reduction and meteorological conditions on air**
2 **quality improvement during the 2014 Youth Olympic Games in**
3 **Nanjing, China**

4
5 Qian Huang ^{1,2}, Tijian Wang ^{1,*}, Pulong Chen ¹, Xiaoxian Huang ³, Jialei Zhu ⁴, and
6 Bingliang Zhuang ¹

7 ¹ School of Atmospheric Sciences, CMA-NJU Joint Laboratory for Climate Prediction
8 Studies, Jiangsu Collaborative Innovation Center for Climate Change, Nanjing
9 University, Nanjing, 210023, China

10 ² Scientific Research Academy of Guangxi Environmental Protection, Nanning,
11 530022, China

12 ³ College of Plant Science & Technology, Huazhong Agricultural University, Wuhan,
13 430070, China

14 ⁴ Department of Climate and Space Sciences and Engineering, University of Michigan,
15 Ann Arbor, 48109, USA

16
17 Correspondence to: Tijian Wang (tjwang@nju.edu.cn)

18
19 **Abstract**

20 As the holding city of the 2nd Youth Olympic Games (YOG), Nanjing is highly
21 industrialized and urbanized facing with several air pollution issues. In order to ensure
22 better air quality during the event, the local government took great efforts to control
23 the pollution emissions. However, air quality can still be affected by synoptic weather.
24 In this paper, the influences of meteorological factors and emission reductions were
25 investigated using observational data and numerical simulations with WRF/CMAQ.
26 During the YOG holding month (Aug., 2014), the observed hourly mean
27 concentrations of SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃ were 11.6 μg/m³, 34.0 μg/m³,
28 57.8 μg/m³, 39.4 μg/m³, 0.9 mg/m³, and 38.8 μg/m³, respectively, which were below
29 China National Ambient Air Quality Standard (Level 2). However, model simulation

30 showed that the weather conditions such as weaker winds during the holding time
31 were adverse for better air quality, and could increase SO₂, NO₂, PM₁₀, PM_{2.5} and CO
32 by 17.5%, 16.9%, 18.5%, 18.8%, 7.8% and 0.8%, respectively. Taking account of
33 local emission abatement only, the simulated SO₂, NO₂, PM₁₀, PM_{2.5} and CO was
34 decreased by 24.6%, 12.1%, 15.1%, 8.1% and 7.2%, respectively. Consequently,
35 stringent emission control measures can reduce the concentrations of air pollutants in
36 short term, and emission reduction is the very important factor for the air quality
37 improvement during the YOG, which has set up a good example in air protection for
38 important social events.

39 **KEY WORDS:** Youth Olympic Games; Emission reduction; Meteorological
40 conditions; WRF/CMAQ; Nanjing

41

42 **1 Introduction**

43 As located in the economically developed Yangtze River Delta (YRD) region of
44 China, Nanjing successfully hosted the second Youth Olympic Games (YOG) during
45 16 - 28 Aug., 2014. Nanjing is a highly urbanized city and its GDP ranked the 12th of
46 all the cities in China by 2013 (National Bureau of Statistics of China, 2014). Due to
47 fast urbanization and industrialization, heavy motor vehicles and construction dust,
48 Nanjing has long been suffered from air pollution (Dong et al., 2013; Chen et al.,
49 2015).

50 In order to realize the promise of “Green YOG”, the local government had taken
51 a series of measures to reduce emissions of air pollutants. The preparatory work
52 started from 1 Jul., 2014. Besides, the local government performed the stringent
53 environmental quality assurance work plan from 1 Aug. (National Bureau of Statistics
54 of China, 2014). The controlled emissions include 5 major categories: industry, power
55 plants, traffic, VOC product-related sources and others. Some local petrochemical,
56 chemical and steel industries were forced to limit or halt the production.
57 Coal-combustion enterprises were required to use high-quality coals with low sulfur
58 content and ash content. And vehicles with heavy pollution called “yellow label
59 buses” were prohibited in Nanjing during 10-28 Aug.. Oil loading and unloading

60 operations were strictly controlled. All construction processes in the city were forced
61 to stop. Surface with bare soil was covered.

62 It is well known that air quality can be affected by both meteorological factors
63 and pollutant emissions. Many cases verified that both emission abatement efforts and
64 weather conditions can influence the improvement of local air quality. Emission
65 control has been taken in many social events, like Beijing Olympic Games in 2008
66 and Shanghai Expo in 2010. Xing et al. (2011) suggested that emission controls
67 benefit for pollutants reduction, but meteorological effects can be either ways at
68 different locations. Cermak and Knutti (2009), Wang et al. (2009b, 2010) and Xing et
69 al. (2011) reported that typical meteorological conditions accounted more for air
70 improvement during 2008 Beijing Olympics than emission reductions. Zhou et al.
71 (2010) concluded that transportation control measures resulted in a reduction of
72 44.5% and 49.0% in daily CO and NO_x emission from motor vehicles during the 2008
73 Olympics. Cai et al. (2011) and Wang et al. (2009a) also studied the transportation
74 controls on improving air quality during Beijing Olympic Games. Okuda et al. (2011)
75 argued that sources control during Beijing Olympics significantly reduced PM₁₀, NO₂
76 and SO₂, but did not as effectively reduce PM_{2.5}. Streets et al. (2007) proposed that
77 local sources controlling is inadequate for heavily populated, urbanized, and
78 industrialized city, regional air quality management is in urgent need. Lin et al. (2013)
79 applied monitoring data to analyze the weather impacts on air quality of the World
80 Expo in YRD and concluded that high frequency of marine winds during the Expo
81 had a positive effect on the air quality of coastal cities, but a negative effect on some
82 inland cities in YRD. Satellite data reflected that the tropospheric NO₂ column,
83 aerosol optical thickness (AOT), and CO concentration dropped by 8%, 14% and 12%,
84 respectively over Shanghai during the Expo period, compared to the past three years
85 (Hao et al., 2011). Liu et al. (2013) compared the contributions of long-term and
86 short-term emission control via CMAQ simulation and compared their effects on air
87 quality in Guangzhou during the Asian Games. Xu et al. (2013) concluded that PM_{2.5}
88 was mainly emitted from anthropogenic sources other than biogenic sources and
89 indicated that cutting down anthropogenic emissions could decrease PM_{2.5} effectively.

90 Dong et al. (2013) found that independent NO_x emission reduction would strengthen
91 O₃ as a side effect in YRD. Chen et al. (2015, 2017) studied the source apportionment
92 of size-fractionated particles in Nanjing, and found that construction dust contributes
93 the most in coarse particles, and fugitive and construction dust decreased significantly
94 in YOG.

95 There have been some studies on air quality during the 2nd YOG (Ding et al.,
96 2015; Chen et al., 2017; Zhou et al. 2017), but few work focused on the relative
97 contributions of meteorology and control efforts. This study takes the air quality
98 monitoring data and applies WRF/CMAQ model to estimate the effect of
99 meteorological factors and emission reduction on air quality of Nanjing during the
100 2nd YOG. Data and model descriptions as well as simulation scenarios are described
101 in Section 2. Section 3 examines the characteristics of six major air pollutants (SO₂,
102 NO₂, PM₁₀, PM_{2.5}, CO and O₃) and compares their concentrations during the YOG
103 with those a year ago and the months without emission reduction (Jul. and Sept.,
104 2014). Besides, this section discusses the separate effect of weather conditions and
105 emission abatement qualitatively and quantitatively based on the simulation results.
106 Section 4 summaries the main conclusions, emphasizes the important factor of the air
107 quality promotion during the YOG, and provides some advice for ensuring pleasant
108 future air quality.

109

110 **2 Methodology**

111 2.1 Data description

112 Hourly observed air quality data during Jul.- Sept. 2014 and Aug. 2013 of two
113 representative stations was collected from Nanjing Environmental Monitoring Center
114 (<http://222.190.111.117:8023/>). Both of the two stations are state controlling air
115 monitoring sites. The data quality assurance and quality control procedures for
116 monitoring strictly follow the national standards (State Environmental Protection
117 Administration of China, 2006). Caochangmen (CCM) Station (118.75° E, 32.06° N)
118 locates in Gulou District, the city center of Nanjing. Gulou District is the center of
119 economy, politics, culture and education in Nanjing. Here gathers many East China's

120 high-end industrial and corporate headquarters. Besides, over 90% provincial
121 authorities, more than 20 colleges and universities, and more than 120 research
122 institutes situate in Gulou District. It's the most populated area in Nanjing, with lively
123 commercial hub and heavy traffic. Thus, CCM station was chosen to represent the
124 urban status of Nanjing. The other site calls Xianlin (XL) Station (118.92° E, 32.11°
125 N), which locates in Qixia District, the suburb of Nanjing. Compared to Gulou
126 District, Qixia District is much more sparsely populated with few traffic congestion
127 problem. Thus, XL station was chosen to represent the suburban status of Nanjing.


128

129 2.2 Model description

130 The integrated modeling system WRF/CMAQ was employed in this research.
131 Community Multiscale Air Quality (CMAQ) is a third-generation regional air quality
132 model developed by the Environmental Protection Agency of USA (USEPA). It
133 incorporates a set of up-to-date compatible modules and control equations for the
134 atmosphere, and can fully consider complicated physical and chemical processes
135 (Byun and Schere, 2006; Foley et al., 2010). Many applications have proven that
136 CMAQ is a reliable tool in simulating air quality from city scale to mesoscale (Xing
137 et al., 2011; Dong et al., 2013; Liu et al., 2013; Xu et al. 2013; Shu et al., 2016).
138 Community Multiscale Air Quality (CMAQ v4.7.1, Binkowski and Roselle, 2003)
139 model includes the 2005 Carbon Bond gas-phase mechanism (CB05) (Yarwood et al.,
140 2005) and the fourth-generation CMAQ aerosol module (AERO4) (Byun and Schere,
141 2006). And it was applied to simulate the pollutant distribution over Nanjing in this
142 paper. Weather Research and Forecasting (WRF) is a new generation of mesoscale
143 weather forecast model and assimilation system, developed by the National Center for
144 Atmospheric Research (NCAR). It has been widely applied in China and shows a
145 good performance in all kinds of weather forecasts (Jiang et al., 2008, 2012; Xu et
146 al.,2013; Liao et al., 2014, 2015; Xie et al., 2014, 2016; Li et al., 2016; Shu et al.,
147 2016). WRF v3.2.1 (Skamarocket al., 2008) model was run to provide meteorology
148 fields for CMAQ. Four nested domains were set for both models, with horizontal
149 resolutions of 81km, 27km, 9km, 3km, with the innermost domain covering Nanjing

150 (Fig. 1). For all domains, 23 vertical sigma layer from the surface to the top pressure
 151 of 100 hpa was set, with about 10 layers in the planetary boundary layer. The detail
 152 dynamic parameterization in WRF as well as the physical and chemical schemes of
 153 CMAQ applied in this research were the same as those in Shu et al. (2016)'s work and
 154 were proven to have good performance. As for the innermost domain, Nanjing
 155 Environmental Protection Bureau chooses the local 9 state controlling air monitoring
 156 sites (See Fig. 1, Table 1) to represent the whole Nanjing (NJ) city. In conformity with
 157 this, the 9 sites in domain4 were chosen to represent the whole Nanjing while
 158 analyzing the impacts of weather conditions and emission reduction.
 159

(a) Domain Configuration



160
 161 **Fig. 1.** Modeling domains and state controlling air monitoring sites in Nanjing. ((a) The four nested
 162 modeling domains at 81km (d01: East Asia), 27km (d02: East China), 9km (d03: Yangtze River Delta),
 163 and 3km (d04: Nanjing), (b) Locations of 9 sites in Nanjing).

164

165 **Table 1**

166 The air monitoring sites in Nanjing

Sites	Abbreviations	Location
Xuanwuhu Station	XWH	32.08° N, 118.80° E
Ruijinlu Station	RJL	32.03° N, 118.82° E

Zhonghuamen Station	ZHM	32.00° N, 118.76° E
Caochangmen Station	CCM	32.06° N, 118.75° E
Shanxilu Station	SXL	32.07° N, 118.77° E
Maigaoqiao Station	MGQ	32.11° N, 118.81° E
Xianlin Station	XL	32.11° N, 118.92° E
Aoti Station	AT	32.01° N, 118.74° E
Pukou Station	PK	32.07° N, 118.64° E


167

168 2.3 Emissions and simulation scenarios

169 In this study, Multi-resolution Emission Inventory for China (MEIC v1.2,
170 <http://www.meicmodel.org/>) with a resolution of $0.25^{\circ} \times 0.25^{\circ}$ was employed to
171 provide the anthropogenic emissions for species including SO₂, NO_x, CO, NMVOC,
172 NH₃, CO₂, PM_{2.5}, PM₁₀, BC, and OC, from 4 sectors: industry, power plants,
173 transportation, and residential.

174 For the innermost domain, the local emission inventory before and after emission
175 control was used with a horizontal resolution of 3km \times 3km. The base year of the
176 local emission is 2012. According to the local emission control program, 5 major
177 categories: industry, power plants, traffic, VOC product-related sources and others
178 were in the control list. In Aug. 2014, all coal-combustion enterprises must use
179 high-quality coals with low sulfur content less than 0.5% and ash content less than
180 13%. Besides, the local government ordered over 100 local petrochemical, chemical
181 and steel enterprises to cut or halt their production. Moreover, heavy pollution
182 vehicles were prohibited in Nanjing during 10-28 Aug. 2014 to reduce traffic
183 emission. To reduce emissions of volatile organic compounds, loading and unloading
184 oil operations were prohibited at the docks in Nanjing section of Yangtze River.
185 What's more, local construction work was halted during Aug. 2014. With these efforts,
186 the emission would be cut by 25.0% for SO₂, 15.0% for NO_x, 42.8% for PM₁₀, 36.2%
187 for PM_{2.5}, and 20.0% for CO. The spatial distributions of emission reduction were
188 showed in Fig. 2. For SO₂, NO_x, PM₁₀ and PM_{2.5}, the emission reduction area centered
189 in the middle of Nanjing city. And for CO, the emission reduction centered in several
190 points.

191



192
193 **Fig. 2.** Emission reduction in domain4 ((a) SO₂, (b) NO_x, (c) PM₁₀, (d) PM_{2.5}, (e) CO (unit: t/month)).
194

195 The simulated period was from Jul. 27 to Sept. 1 (China standard time, CST), but
196 only the holding month (1 Aug. to 31 Aug.) was focused on for discussions. In order
197 to better understand the influence of meteorological factors and emission abatement,
198 three experiments were carried out. Exp. 1 used the weather conditions during Aug.
199 2014 (CST) and the emission inventory after reduction. Exp. 2 used the same weather
200 conditions as Exp. 1 with the emission inventory before reduction. Exp. 3 used the
201 same inventory as Exp. 2 and the weather conditions during Aug. 2013 (CST).
202 Besides, Exp. 2 acted as the control experiment. Therefore, Exp. 1 and Exp. 2 were
203 performed to investigate the influence of emission reduction on pollutants. Similarly,
204 Exp. 2 and Exp. 3 were conducted to understand the impact of meteorology on air
205 quality.


206
207 **3 Results and discussion**

208 **3.1 Observed air quality during the YOG**

209 In the most strictly control month Aug. 2014, emission sources including 5 major
210 categories were reduced, and the air quality had great promotion compared to Aug.
211 2013. Firstly, it was good during the Games in accordance with China's National
212 Ambient Air Quality Standards (NAAQS) (Ministry of Environmental Protection of

213 the People's Republic of China, 2012) (Fig. 3, Fig. 4). The hourly mean pollutant
 214 concentration of the two sites during Aug. 2014 is $11.6 \mu\text{g}/\text{m}^3$ for SO_2 , $34.0 \mu\text{g}/\text{m}^3$ for
 215 NO_2 , $57.8 \mu\text{g}/\text{m}^3$ for PM_{10} , $39.4 \mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$, $0.9 \text{ mg}/\text{m}^3$ for CO , and $38.8 \mu\text{g}/\text{m}^3$
 216 for O_3 . Secondly, as showed in Table 2 and Table 3, the mean concentration of the six
 217 major species dropped by 64.7% for SO_2 , 29.8% for PM_{10} , 9.8% for $\text{PM}_{2.5}$, 8.9% for
 218 CO and 31.7% for O_3 at CCM station, while 50.0% for SO_2 , 18.6% for NO_2 , 32.8%
 219 for PM_{10} , 4.1% for $\text{PM}_{2.5}$, and 31.7% for O_3 at XL station. Besides, the smaller
 220 standard deviation (std) of SO_2 , NO_2 , CO and O_3 revealed that concentrations of these
 221 air pollutants varied more steadily in Aug. 2014. However, the drop of pollutant
 222 concentration could be caused mainly by meteorology conditions or emission
 223 reductions. And we will discuss this issue based on model simulations in Section 3.2
 224 and Section 3.3.


225



226

227 **Fig. 3.** Day-to-day variations in SO_2 , NO_2 , PM_{10} , $\text{PM}_{2.5}$, CO and O_3 -8h at CCM station in Aug. 2013
 228 and Aug. 2014 (CST). Observed data in Aug. 2013 and Aug. 2014 are indicated in blue and red,
 229 respectively. NAAQS are indicated in black dotted line.

230



231
 232 **Fig. 4.** Day-to-day variations in SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃-8h at XL station in Aug. 2013 and
 233 Aug. 2014 (CST). Observed data in Aug. 2013 and Aug. 2014 are indicated in blue and red,
 234 respectively. NAAQS are indicated in black dotted line.

235

236 **Table 2**

237 Statistical analysis of hourly data in Aug. 2013 and Aug. 2014 at CCM station (The unit is µg/m³
 238 except CO (mg/m³))

species	time	max	min	mean	median	std	Δ
SO ₂	Aug. 2013	169.0	1.0	33.7	27.0	23.7	
	Aug. 2014	72.0	2.0	11.9	10.0	7.8	-64.7%
NO ₂	Aug. 2013	111.0	1.0	35.4	32.0	19.4	
	Aug. 2014	110.0	1.0	37.3	35.0	18.6	5.0%
PM ₁₀	Aug. 2013	213.0	19.0	86.0	84.0	29.5	
	Aug. 2014	198.0	6.0	60.4	54.0	36.6	-29.8%
PM _{2.5}	Aug. 2013	123.0	10.0	45.2	43.5	16.2	
	Aug. 2014	165.0	3.0	40.7	36.0	23.8	-9.8%
CO	Aug. 2013	3.1	0.4	1.0	0.9	0.4	
	Aug. 2014	2.2	0.3	0.9	0.9	0.3	-8.9%
O ₃	Aug. 2013	198.0	1.0	56.9	42.0	46.2	
	Aug. 2014	150.0	9.0	38.9	34.0	22.6	-31.7%

239 Δ : the change percentage of species in Aug. 2014 based on Aug. 2013.

240

241 **Table 3**

242 Statistical analysis of hourly data in Aug. 2013 and Aug. 2014 at XL station (The unit is µg/m³ except
 243 CO (mg/m³))

species	time	max	min	mean	median	std	Δ
SO ₂	Aug. 2013	139.0	0.0	22.8	19.0	16.1	
	Aug. 2014	71.0	1.0	11.4	8.0	10.4	-50.0%
NO ₂	Aug. 2013	129.0	0.0	37.7	32.0	21.7	

	Aug. 2014	95.0	7.0	30.7	27.0	15.0	-18.6%
PM ₁₀	Aug. 2013	215.0	0.0	82.1	79.0	32.4	
	Aug. 2014	196.0	6.0	55.2	47.0	35.9	-32.8%
PM _{2.5}	Aug. 2013	122.0	0.0	39.7	37.5	18.9	
	Aug. 2014	157.0	3.0	38.0	34.0	24.1	-4.1%
CO	Aug. 2013	3.2	0.0	0.8	0.7	0.4	
	Aug. 2014	2.0	0.3	0.8	0.7	0.3	<0.1%
O ₃	Aug. 2013	193.0	0.0	56.6	44.0	37.5	
	Aug. 2014	148.0	2.0	38.7	32.0	28.3	-31.7%


244 Δ : the change percentage of species in Aug. 2014 based on Aug. 2013.

245

246 Analogously, compared the observational data in Aug. 2014 with that in Jul. and
247 Sept. 2014 (the months before and after the most aggressive abatement), the
248 concentrations of most species also decreased obviously. As presented in Fig. 5 and
249 Fig. 6, without abatement, NO₂, PM₁₀, PM_{2.5} and O₃ were likely to exceed NAAQS,
250 especially for PM_{2.5} and O₃. As shown in Table 4 and Table 5, compared with Jul.
251 2014, the concentration of NO₂, PM₁₀, PM_{2.5}, CO and O₃ dropped by 0.7%, 31.8%,
252 33.7%, 1.1%, and 52.8%, respectively at CCM station in Aug. 2014, while the
253 concentration of SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃ decreased by 15.8%, 39.6%,
254 34.6%, 7.1%, and 50.7%, respectively at XL station in Aug. 2014. Without emission
255 control, the concentration of air pollutants rebounded in Sept. 2014. Compared to
256 Aug., the concentration of SO₂, NO₂, PM₁₀, PM_{2.5} and O₃ increased by 37.4%, 19.8%,
257 37.6%, 22.3%, and 47.2%, respectively at CCM station in Sept. 2014 (Table 4), while
258 the concentration of SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃ increased by 24.6%, 21.8%,
259 28.7%, 17.7%, 4.9%, and 39.9%, respectively at XL station in Sept. 2014 (Table 5).
260 Besides, for most species, the standard deviation was the lowest in Aug., which meant
261 that the potential of extreme events was the least in Aug.. Assume that the weather
262 conditions in Jul., Aug., Sept., 2014 were similar, it can be estimated that emission
263 sources could be the major impact factor of explaining the concentration changes
264 during the three months. These results proved that concentrations of most species
265 decreased and had less potential in extreme events after aggressive emission
266 abatement. However, the concentration became higher without emission control.

267 Section 3.3 would further discuss the change of pollutant concentration with and
 268 without emission reduction based on model simulation.


269



270

271 **Fig. 5.** Day-to-day variations in SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃-8h at CCM station in Jul., Aug.
 272 and Sept. 2014 (CST). Observed data in Jul., Aug. and Sept. 2014 are indicated in green, red and blue,
 273 respectively. NAAQS are indicated in black dotted line.

274



275

276 **Fig. 6.** Day-to-day variations in SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃-8h at XL station in Jul., Aug. and
 277 Sept. 2014 (CST). Observed data in Jul., Aug. and Sept. 2014 are indicated in green, red and blue,
 278 respectively. NAAQS are indicated in black dotted line.

279

280 **Table 4**

281 Statistical analysis of hourly data in Jul. - Sept. 2014 at CCM station (The unit is µg/m³ except CO
 282 (mg/m³))

species	month	max	min	mean	median	std	Δa	Δb
SO ₂	Jul. 2014	83.0	1.0	11.3	9.0	9.8		
	Aug. 2014	72.0	2.0	11.9	10.0	7.8	5.1%	-37.4%

	Sept. 2014	70.0	4.0	19.0	18.0	9.9		
	Jul.-Sept. 2014	83.0	1.0	14.0	12.0	9.8		
NO ₂	Jul. 2014	161.0	1.0	37.5	32.0	28.3		
	Aug. 2014	110.0	1.0	37.3	35.0	18.6	-0.7%	-19.8%
	Sept. 2014	151.0	8.0	46.5	42.0	24.5		
	Jul.-Sept. 2014	161.0	1.0	40.2	37.0	24.4		
PM ₁₀	Jul. 2014	255.0	6.0	88.5	88.0	50.7		
	Aug. 2014	198.0	6.0	60.4	54.0	36.6	-31.8%	-37.6%
	Sept. 2014	243.0	6.0	96.7	90.0	45.8		
	Jul.-Sept. 2014	255.0	6.0	81.7	76.0	47.4		
PM _{2.5}	Jul. 2014	171.0	1.0	61.5	58.0	33.9		
	Aug. 2014	165.0	3.0	40.7	36.0	23.8	-33.7%	-22.3%
	Sept. 2014	143.0	3.0	52.4	46.0	27.2		
	Jul.-Sept. 2014	171.0	1.0	51.5	45.0	29.9		
CO	Jul. 2014	2.7	0.2	0.9	0.9	0.3		
	Aug. 2014	2.2	0.3	0.9	0.9	0.3	-1.1%	21.1%
	Sept. 2014	2.1	0.1	0.8	0.7	0.4		
	Jul.-Sept. 2014	2.7	0.1	0.9	0.8	0.4		
O ₃	Jul. 2014	281.0	4.0	82.4	69.0	57.6		
	Aug. 2014	150.0	9.0	38.9	34.0	22.6	-52.8%	-47.2%
	Sept. 2014	240.0	6.0	73.6	61.0	49.2		
	Jul.-Sept. 2014	281.0	4.0	64.7	51.0	49.3		

283 Δa: the change percentage of species in Aug.2014 based on Jul. 2014.

284 Δb: the change percentage of species in Aug. 2014 based on Sept. 2014.

285

286 **Table 5**

287 Statistical analysis of hourly data in Jul. - Sept.2014 at XL station (The unit is μg/m³ except CO

288 (mg/m³))

species	month	max	min	mean	median	std	Δa	Δb
SO ₂	Jul. 2014	61.0	1.0	14.5	12.0	10.3		
	Aug. 2014	71.0	1.0	11.4	8.0	10.4	-21.2%	-24.6%
	Sept. 2014	75.0	1.0	15.1	14.0	10.3		
	Jul.-Sept. 2014	75.0	1.0	13.7	11.0	10.4		
NO ₂	Jul. 2014	123.0	9.0	36.4	33.0	18.9		
	Aug. 2014	95.0	7.0	30.7	27.0	15.0	-15.8%	-21.8%
	Sept. 2014	127.0	11.0	39.2	36.0	18.7		
	Jul.-Sept. 2014	127.0	7.0	35.4	32.0	18.0		
PM ₁₀	Jul. 2014	300.0	4.0	91.3	85.0	48.9		
	Aug. 2014	196.0	6.0	55.2	47.0	35.9	-39.6%	-28.7%
	Sept. 2014	226.0	9.0	77.3	70.0	40.3		

	Jul.-Sept. 2014	300.0	4.0	74.5	64.0	44.6		
	Jul. 2014	158.0	2.0	58.2	51.0	34.8		
PM _{2.5}	Aug. 2014	157.0	3.0	38.0	34.0	24.1	-34.6%	-17.7%
	Sept. 2014	144.0	3.0	46.2	38.0	29.0		
	Jul.-Sept. 2014	158.0	2.0	47.4	40.5	30.7		
	Jul. 2014	2.0	0.3	0.8	0.8	0.4		
CO	Aug. 2014	2.0	0.3	0.8	0.7	0.3	-7.1%	-4.9%
	Sept. 2014	2.8	0.3	0.8	0.7	0.4		
	Jul.-Sept. 2014	2.8	0.3	0.8	0.7	0.4		
	Jul. 2014	238.0	2.0	78.4	67.0	55.6		
O ₃	Aug. 2014	148.0	2.0	38.7	32.0	28.3	-50.7%	-39.9%
	Sept. 2014	226.0	2.0	64.4	54.0	46.4		
	Jul.-Sept. 2014	238.0	2.0	60.3	48.0	47.7		

289 Δ a: the change percentage of species in Aug.2014 based on Jul. 2014.

290 Δ b: the change percentage of species in Aug. 2014 based on Sept. 2014.

291

292 3.2 Simulated impact of meteorological conditions


293 In this paper, the model configurations were the same as those set by Shu et al.
 294 (2016), who has evaluated the model performance of WRF/CMAQ and proved the
 295 model's reliability in simulating air quality in Nanjing.

296 As we know, meteorology is an important impact factor on air quality. Good
 297 diffusion conditions can alleviate air pollution in the short term (Cermak and Knutti,
 298 2009; Wang et al., 2009b). In this premise, if two experiments (Exp. 2 and Exp. 3) use
 299 the same emission inventory but different weather conditions, it can be concluded that
 300 the higher concentrations may result from poor meteorological conditions. According
 301 to model simulation, Exp. 2 exhibited higher pollutant concentrations for all species in
 302 most part of Nanjing as shown in Fig. 7. For SO₂, NO₂, PM₁₀, PM_{2.5}, CO, and O₃,
 303 their concentrations were increased by 17.5%, 16.9%, 18.5%, 18.8%, 7.8% and 0.8%
 304 during Aug. 2014 compared to Aug. 2013. Besides, the contributions of
 305 meteorological conditions to primary and secondary particulate matters differed (See
 306 Fig. 8). Secondary PM₁₀ (PM_{10s}) was raised by 21.5%, while primary PM₁₀ (PM_{10p})
 307 rose by 12.6% during Aug. 2014 compared to Aug. 2013. And secondary PM_{2.5}
 308 (PM_{2.5s}) was increased by 21.5%, while primary PM_{2.5} (PM_{2.5p}) was added by 9.5%.
 309 Thus, the weather conditions had a slightly greater impact on secondary fine

310 particulate matters. Moreover, for SO₂, NO₂, PM₁₀, PM_{2.5}, CO, PM_{10p}, PM_{10s}, PM_{2.5p},
311 and PM_{2.5s}, there were some small decreasing areas in the northeast Nanjing (Fig. 7
312 and Fig. 8). In domain 4, the simulated predominant wind was northeast wind in Aug.
313 2014, while that was southeast wind in Aug. 2013. So, the diffusion condition of
314 northeast Nanjing might be better in Aug. 2014 and resulted in small decrease in these
315 areas.

316 The overall increasing pollutant levels in Aug. 2014 suggested that the diffusion
317 conditions in Aug. 2014 were worse than those in Aug. 2013. Focus on the weather
318 conditions during the YOG holding period (16-28 Aug., 2014) and the same period in
319 2013, the simulated hourly mean 10-m wind speed in Nanjing was larger in 16-28
320 Aug., 2013, and it was 1.5 m/s larger than that of 16-28 Aug., 2014 (Fig. 9). Also, the
321 simulated 2-m temperature was higher in 16-28 Aug., 2013, and it was 2.0 K larger
322 than that of 16-28 Aug., 2014 (Fig. 9). Besides, the simulated planetary boundary
323 layer height (PBLH) was higher in Aug. 2013, especially in 16-28 Aug., and it was
324 27.5 m higher than that of 16-28 Aug., 2014 (Fig. 9). Larger wind speed and higher
325 PBLH benefited the diffusion of air pollutants. Warming on the ground surface was
326 conducive to the promotion of convective instability and was also good for the
327 vertical dilution and diffusion of pollutant. Thus, the diffusion conditions in 16-28
328 Aug. 2013 were better than those in 16-28 Aug. 2014. Rather worse meteorological
329 conditions in 16-28 Aug. 2014 implied that abatement controls might play a more
330 important role in improving air quality in YOG compared with the same period in
331 2013. What's more, relative humidity, cloud cover and shortwave solar radiation all
332 affect ozone chemical reaction (Katrakou et al., 2011; Pu et al., 2017). The
333 generation and photochemical reaction of surface ozone depends on the availability of
334 solar radiation. During the heat wave period, less relative humidity leads to less cloud
335 cover, which could result in more net downward shortwave solar radiation and more
336 production of O₃ (Pu et al., 2017). For ordinary O₃, its production also corresponded
337 well to the cloud cover. As shown in Fig. 9, more relative humidity resulted in more
338 cloud cover in northern and eastern Nanjing during 16-28 Aug., 2013, which resulted
339 in less O₃ in Aug. 2013, but more O₃ in Aug. 2014 (Fig. 7).


340



341

342 **Fig. 7.** Influence of meteorology on hourly mean concentrations of air pollutants in Aug. 2014
343 compared with Aug. 2013. (Black thick lines draw the outline of Nanjing. Picture a - f are hourly
344 average values of impact percentage (dspecies (%))= (Exp. 2 - Exp. 3)/Exp. 2 * 100%) of SO₂, NO₂,
345 PM₁₀, PM_{2.5}, CO, and O₃, respectively.).


346



347

348 **Fig. 8.** Influence of meteorology on hourly mean concentrations of primary and secondary particulate
349 matters in Aug. 2014 compared with Aug. 2013. (Black thick lines draw the outline of Nanjing. Picture
350 a - d are hourly average values of impact percentage (dspecies (%))= (Exp. 2 - Exp. 3)/Exp. 2 * 100%)
351 of PM_{10p}, PM_{10s}, PM_{2.5p}, and PM_{2.5s}, respectively.)

352



353

354 **Fig. 9.** Bias of simulated hourly mean meteorological conditions during the YOG. (Bias =
 355 Meteorological Factors in 16-28 Aug., 2013 - Meteorological Factors in 16-28 Aug., 2014. (a) Bias of
 356 Wind at 10m during 16-28 Aug. (unit: m/s), (b) Bias of temperature at 2m during 16-28 Aug. (unit: K),
 357 (c) Bias of planetary boundary layer height during 16-28 Aug. (unit: m), (d) Bias of relative humidity at
 358 2m during 16-28 Aug. (unit: %), (e) Bias of cloud fraction during 16-28 Aug.).


359

360 3.3 Simulated impact of emission reduction

361 As for SO₂, NO₂, PM₁₀, PM_{2.5}, and CO, the distributions of such short-lived
 362 chemical compositions are largely affected by their sources and sinks. As seen in Fig.
 363 10, the simulated spatial distributions of concentration changes were uneven, large
 364 variations were found in the west of Nanjing corresponding to the downwind regions
 365 of heavy reduction districts (See Fig. 2). Besides, impact percentages ($ds_{\text{species}}(\%) =$
 366 $(\text{Exp. 1} - \text{Exp. 2}) / \text{Exp. 2} * 100\%$) of species were negative except O₃, implying that
 367 emission regulatory efforts were effective on the other species, but counterproductive
 368 to O₃. Statistically, the concentrations of SO₂, NO₂, PM₁₀, PM_{2.5}, and CO in Nanjing
 369 were reduced by 24.6%, 12.1%, 15.1%, 8.1% and 7.2% during Aug. 2014. As for O₃,
 370 the variation was positive (1.3%), especially in the downwind area of NO_x with heavy
 371 reduction, which might due to the less titration of O₃ by NO_x (Liu et al., 2013; Dong
 372 et al., 2013). For primary and secondary particulate matters, the influence of emission
 373 reduction varied dramatically. As shown in Fig. 11, PM_{10p} was dropped by 39.6%,
 374 while PM_{10s} only declined by 2.9%. And PM_{2.5p} was decreased by 26.2%, while
 375 PM_{2.5s} merely cut down by 2.9%. It seems that emission controls had much more

376 impacts on primary pollutants, especially on coarse particulate matters.

377



378


379 **Fig. 10.** Influence of emission reduction on hourly mean concentrations of air pollutants in Aug. 2014.

380 (Black thick lines draw the outline of Nanjing. Picture a - f are hourly average values of impact

381 percentage ($dspecies (\%) = (Exp. 1 - Exp. 2) / Exp. 2 * 100\%$) of SO₂, NO₂, PM₁₀, PM_{2.5}, CO and O₃,

382 respectively.).

383



384

385 **Fig. 11.** Influence of emission reduction on hourly mean concentrations of primary and secondary

386 particulate matters in Aug. 2014. (Black thick lines draw the outline of Nanjing. Picture a - d are hourly

387 average values of impact percentage ($dspecies (\%) = (Exp. 1 - Exp. 2) / Exp. 2 * 100\%$) of PM_{10p}, PM_{10s},

388 PM_{2.5p} and PM_{2.5s}, respectively.).

389

390 3.4 Comparison of simulated meteorological factors and emission reduction


391 Fig. 12 displays the simulated effect of meteorological factors and emission

392 reduction in Nanjing on air quality improvement during Aug. 2014. In general,

393 meteorological conditions played a negative role in air quality promotion in most days,

394 only played a positive role in a few days. And the negative effect of weather

395 conditions exceeded its positive effect during the whole month (See discussion in
 396 Section 3.2). On the other hand, emission reduction contributed to the decline of SO₂,
 397 NO₂, PM₁₀, PM_{2.5}, CO, PM_{10p}, PM_{10s}, PM_{2.5p}, and PM_{2.5s} all the time, especially for
 398 primary coarse particulate matters. However, reduction of NO_x could cause a slight
 399 rise of O₃.
 400



401
 402 **Fig. 12.** The simulated effect of meteorology and reduction on pollutant concentrations in Nanjing
 403 during 1-31 Aug. , 2014, Met. (Exp. 2-Exp. 3) represents the effect of meteorology, while Red. (Exp.
 404 1-Exp. 2) represents the simulated effect of reduction.

405
 406 Moreover, the opposite effects of meteorology and emission abatement on SO₂,
 407 NO₂, PM₁₀, PM_{2.5}, CO, PM_{10p}, PM_{10s}, PM_{2.5p} and PM_{2.5s} during the whole month were
 408 more apparent as statistically listed in Table 6. CCM station represents the urban
 409 status, XL station represents the suburban status and NJ represents the whole city.
 410 Adverse meteorology was found to raise the concentration of the six pollutants as
 411 17.4% for SO₂, 15.1% for NO₂, 15.6% for PM₁₀, 14.9% for PM_{2.5}, 6.4% for CO and
 412 0.9% for O₃ at CCM station, and 14.1% for SO₂, 12.4% for NO₂, 22.4% for PM₁₀,
 413 24.5% for PM_{2.5}, 2.3% for CO, and 1.6% for O₃ at XL station. On the contrary,
 414 emission abatement reduced their levels in most cases, especially in the urban site. It
 415 seems that the levels of air pollutants reduced more at CCM station compared to XL
 416 station. Emission abatement independently led to a 24.3% decrease in SO₂ at CCM
 417 station, which was 5.1% higher than that at XL station. Moreover, the cutbacks of

418 NO₂, PM₁₀, PM_{2.5} and CO were 11.7%, 13.9%, 7.5% and 7.0%, respectively at CCM
 419 station, being larger by 1.0% to 2.0% compared with XL station. Though O₃ under
 420 emission reduction scenarios resulted in a slightly rise (0.9% to 1.3%) at both sites,
 421 the effectiveness of emission abatement was remarkable generally. As for primary and
 422 secondary particulate matters, meteorological factors also played a negative role
 423 during Aug. 2014, and had slightly more effect on secondary fine particles. Emission
 424 controls seemed to cause tremendous impact on primary particulate matters,
 425 especially for primary coarse particles. Emission abatement led to a 38.3% decrease in
 426 PM_{10p} at CCM station, a 33.2% decrease in PM_{10p} at XL station, and a 39.6% decrease
 427 in PM_{10p} for the whole city. For secondary particulate matters, including sulfate,
 428 nitrate, and ammonium salt, emission reduction played rather minor role in cutting the
 429 pollutants, with merely a 2.4% decrease in PM_{10s} and PM_{2.5s} at CCM station, a 2.9%
 430 decrease in PM_{10s} and PM_{2.5s} at XL station.

431

432 **Table 6**

433 Comparison between the simulated effect of meteorology and emission reduction at CCM and XL
 434 station

Species	Met. (CCM)	Red. (CCM)	Met. (XL)	Red. (XL)	Met. (NJ)	Red. (NJ)
SO ₂	17.4%	-24.3%	14.1%	-19.2%	17.5%	-24.6%
NO ₂	15.1%	-11.7%	12.4%	-10.2%	16.9%	-12.1%
PM ₁₀	15.6%	-13.9%	22.4%	-11.9%	18.5%	-15.1%
PM _{2.5}	14.9%	-7.5%	24.5%	-6.3%	18.8%	-8.1%
CO	6.4%	-7.0%	2.3%	-5.5%	7.8%	-7.2%
O ₃	0.9%	1.3%	1.6%	0.9%	0.7%	1.5%
PM _{10p}	13.2%	-38.3%	5.9%	-33.2%	12.6%	-39.6%
PM _{10s}	16.7%	-2.4%	29.4%	-2.9%	21.5%	-2.9%
PM _{2.5p}	8.4%	-25.8%	4.9%	-20.1%	9.5%	-26.2%
PM _{2.5s}	16.7%	-2.4%	29.4%	-2.9%	21.5%	-2.9%

435 Met.: the change percentage of species in Exp. 2 based on Exp3, represents the effect of meteorology.

436 Red.: the change percentage of species in Exp. 1 based on Exp 2, represents the effect of Nanjing local
 437 emission reduction.

438

439 The decrease of SO₂ might due to the limit and halt of power plants and
 440 improvement of coal-combustion. Besides, the prohibition of heavy pollution vehicles

441 could contribute to the drop of NO₂ and CO. Also, limiting the production of
442 industries helped to reduce NO₂ and CO. The cut of particulate matters might due to
443 the stop of construction process and use of low ash content coal for industry. For
444 secondary particulate matters, controlling the emission of SO₂ and NO_x could help to
445 reduce the formation of sulfate and nitrate, but no control on the emission of NH₃
446 could still result in quite a part of ammonium salt. The response of O₃ to emission
447 control could be due to its non-linearity chemistry (Fu et al., 2012), reducing NO₂
448 pollution may have side-effect by increasing O₃ because of the titration effect. On the
449 whole, the meteorological factors and emission reduction during the YOG had
450 opposite effects on SO₂, NO₂, PM₁₀, PM_{2.5}, and CO, and emission reduction played a
451 very important role in the air quality improvement.

452

453 **4 Summary and conclusions**

454 The air quality during the 2nd YOG was superior according to the current
455 NAAQS. Both observation and modeling confirmed that stringent emission reductions
456 were effective to ambient air quality promotion during the Youth Olympic Games,
457 especially to SO₂, NO₂, PM₁₀, PM_{2.5} and CO. Compared to Aug. 2013, the observed
458 concentrations in Aug. 2014 were dropped by 64.7% for SO₂, 29.8% for PM₁₀, 9.8%
459 for PM_{2.5}, 8.9% for CO and 31.7% for O₃ at CCM station, while 50.0% for SO₂,
460 18.6% for NO₂, 32.8% for PM₁₀, 4.1% for PM_{2.5}, and 31.7% for O₃ at XL station. The
461 simulated impact percentage of emission reductions were -24.6%, -12.1%, -15.1%,
462 -8.1% and -7.2% for SO₂, NO₂, PM₁₀, PM_{2.5}, and CO, respectively.

463 The meteorological conditions in the holding time of the YOG were inferior to
464 those of the same period in 2013, with lower temperature and weaker winds. Model
465 simulations show that less favorable weather conditions caused higher concentrations
466 for all species, including O₃ which was increased due to less cloud cover. Besides,
467 meteorological conditions had slightly more effect on secondary fine particular
468 matters compared to primary fine particular matters. Emission reduction could cut
469 down the levels of SO₂, NO₂, CO and particular matters, especially for primary coarse
470 particles. Thus, emission reduction control is the very important factor for the air

471 quality improvement during the YOG.

472 In general, better air quality during the YOG benefit a lot from emission
473 reduction, which has set up a good example in air protection for important social
474 events. However, the enhanced concentrations of air pollutants after the YOG (in Sept.
475 2014) suggest that short-term emission control can only ease air pollution effectively
476 and temporarily. Long-term control policies are necessary to ensure pleasant future air
477 quality.

478

479 **Acknowledgements**

480 This work was supported by the National Natural Science Foundation of China
481 (91544230, 41575145, 41621005), the National Key Research Development Program
482 of China (2016YFC0203303, 2016YFC0208504, 2014CB441203), the National
483 Special Fund for the Weather Industry (GYHY201206011) and the National Special
484 Fund for the Environmental Protection Industry (GYHY201409008) . We are grateful
485 to Prof. Yu Zhao from School of Environment of Nanjing University for supply the
486 emission data of Nanjing. The contents of this paper are solely the responsibility of
487 the authors and do not necessarily represent the official views of sponsors.

488

489 **References**

- 490 Binkowski, F. S. and Roselle, S. J.: Models-3 Community Multiscale Air Quality
491 (CMAQ) model aerosol component 1. Model description, *J. Geophys. Res.*, 108,
492 4183, doi:10.1029/2001JD001409, 2003.
- 493 Byun, D. and Schere, K. L.: Review of the governing equations, computational
494 algorithms, and other components of the models-3 Community Multiscale Air
495 Quality (CMAQ) modeling system, *Appl. Mech. Rev.*, 59, 51-77,
496 doi:10.1115/1.2128636, 2006.
- 497 Cai, H. and Xie, S.: Traffic-related air pollution modeling during the 2008 Beijing
498 Olympic Games: the effects of an odd-even day traffic restriction scheme,
499 *Science of the total environment*, 409, 1935-1948, doi:
500 10.1016/j.scitotenv.2011.01.025, 2011.
- 501 Center for Earth System Science, Tsinghua University: Multi-resolution Emission
502 Inventory for China, <http://www.meicmodel.org/>, 2015.
- 503 Cermak, J. and Knutti, R.: Beijing Olympics as an aerosol field experiment,
504 *Geophysical Research Letters*, 36, 1-5, doi:10.1029/2009GL038572, 2009.
- 505 Chen P., Wang T., and Hu X.: Chemical Mass Balance Source Apportionment of
506 Size-Fractionated Particulate Matter in Nanjing, China, *Aerosol & Air Quality
507 Research*, 15, 1855-1867. doi:10.4209/aaqr.2015.03.0172, 2015.
- 508 Chen, P., Wang, T., Lu, X., Yu, Y., Kasoar, M., Xie, M., and Zhuang, B.: Source
509 apportionment of size-fractionated particles during the 2013 Asian Youth Games
510 and the 2014 Youth Olympic Games in Nanjing, China, *Science of the Total
511 Environment*. 579, 860-870. <http://dx.doi.org/10.1016/j.scitotenv.2016.11.014>,
512 2017.
- 513 Ding, J., Van, d. A. R. J., Mijling, B., Levelt, P. F., and Hao, N.: NO_x emission
514 estimates during the 2014 Youth Olympic Games in Nanjing, *Atmos. Chem.
515 Phys.*, 15, 9399-9412, doi: 10.5194/acp-15-9399-2015, 2015.
- 516 Dong, X. Y., Gao, Y., Fu, J. S., Li, J., Huang, K., Zhuang, G. S., and Zhou, Y.: Probe
517 into gaseous pollution and assessment of air quality benefit under sector
518 dependent emission control strategies over megacities in Yangtze River Delta,

519 China, *Atmos. Environ.*, 79, 841-852, doi:10.1016/j.atmosenv.2013.07.041,
520 2013.

521 Foley, K. M., Roselle, S. J., Appel, K. W., Bhawe, P. V., Pleim, J. E., Otte, T. L.,
522 Mathur, R., Sarwar, G., Young, J. O., Gilliam, R. C., Nolte, C. G., Kelly, J. T.,
523 Gilliland, A. B., and Bash, J. O.: Incremental testing of the Community
524 Multiscale Air Quality (CMAQ) modeling system version 4.7, *Geosci. Model*
525 *Dev.*, 3, 205-226, doi:10.5194/gmd-3-205-2010, 2010.

526 Fu, J. S., Dong, X. Y., Gao, Y., Wong, D. C., and Lam, Y. F.: Sensitivity and linearity
527 analysis of ozone in East Asia: the effects of domestic emission and
528 intercontinental transport, *Journal of the Air and Waste Management Association*,
529 62, 1102-1114, doi:10.1080/10962247.2012.699014, 2012.

530 Hao, N., Valks, P., Loyola, D., Cheng, Y. F., and Zimmer, W.: Space-based
531 measurements of air quality during the World Expo 2010 in Shanghai,
532 *Environmental Research Letters*, 6, doi:10.1088/1748-9326/6/4/044004, 2011.

533 Jiang, F., Wang, T. J., Wang, T. T., Xie, M., and Zhao, H.: Numerical modeling of a
534 continuous photochemical pollution episode in Hong Kong using WRF-chem,
535 *Atmos. Environ.*, 42, 8717-8727, doi:10.1016/j.atmosenv.2008.08.034, 2008.

536 Jiang, F., Zhou, P., Liu, Q., Wang, T. J., Zhuang, B. L., and Wang, X. Y.: Modeling
537 tropospheric ozone formation over East China in springtime, *J. Atmos. Chem.*,
538 69, 303-319, doi:10.1007/s10874-012-9244-3, 2012.

539 Katragkou E., Zanis P., Kioutsioukis I., Tegoulas I., Melas D., Krüger, B. C., and
540 Coppola E.: Future climate change impacts on summer surface ozone from
541 regional climate-air quality simulations over Europe, *Journal of Geophysical*
542 *Research Atmospheres*, 116, D22307, doi:10.1029/2011JD015899, 2011.

543 Li, M. M., Song, Y., Mao, Z. C., Liu, M. X., and Huang, X.: Impacts of thermal
544 circulations induced by urbanization on ozone formation in the Pearl River Delta
545 region, China, *Atmos. Environ.*, 127, 382-392,
546 doi:10.1016/j.atmosenv.2015.10.075, 2016.

547 Liao, J. B., Wang, T. J., Wang, X. M., Xie, M., Jiang, Z. Q., Huang, X. X., and Zhu, J.
548 L.: Impacts of different urban canopy schemes in WRF/Chem on regional

549 climate and air quality in Yangtze River Delta, China, *Atmos. Res.*, 145, 226-243,
550 doi:10.1016/j.atmosres.2014.04.005, 2014.

551 Liao, J. B., Wang, T. J., Jiang, Z. Q., Zhuang, B. L., Xie, M., Yin, C. Q., Wang, X. M.,
552 Zhu, J. L., Fu, Y., and Zhang, Y.: WRF/Chem modeling of the impacts of urban
553 expansion on regional climate and air pollutants in Yangtze River Delta, China,
554 *Atmos. Environ.*, 106, 204-214, doi:10.1016/j.atmosenv.2015.01.059, 2015.

555 Lin Y., Huang, K., Zhuang, G., Fu, J. S., Xu, C., Shen, J., and Chen, S.: Air Quality
556 over the Yangtze River Delta during the 2010 Shanghai Expo, *Aerosol & Air
557 Quality Research*, 13, 1655-1666, doi: 10.4209/aaqr.2012.11.0312, 2013.

558 Liu, H., Wang, X. M., Zhang, J. P., He, K. B., Wu, Y., and Xu, J. Y.: Emission controls
559 and changes in air quality in Guangzhou during the Asian Games, *Atmos.
560 Environ.*, 76, 81-93, doi:10.1016/j.atmosenv.2012.08.004, 2013.

561 Ministry of Environmental Protection of the People's Republic of China: The second
562 Summer Youth Olympic Games environmental quality assurance work plan,
563 Beijing, http://www.js.xinhuanet.com/2014-07/19/c_1111695515_2.htm/, 2014.

564 National Bureau of Statistics of China: <http://www.stats.gov.cn/>, 2014.

565 Okuda, T., Matsuura, S., Yamaguchi, D., Umemura, T., Hanada, E., Orihara, H.,
566 Tanaka, S., He, K.B., Ma, Y. L., Cheng, Y., and Liang, L. L.: The impact of the
567 pollution control measures for the 2008 Beijing Olympic Games on the chemical
568 composition of aerosols, *Atmos. Environ.*, 45, 2789-2794,
569 doi:10.1016/j.atmosenv.2011.01.053, 2011.

570 Pu X., Wang T. J., Huang X., Melas D., Zanis P., Papanastasiou D. K., and Poupkou
571 A.: Enhanced surface ozone during the heat wave of 2013 in Yangtze River Delta
572 region, China. *Science of the Total Environment*, 603-604, 807-816,
573 doi:10.1016/j.scitotenv.2017.03.056, 2017.

574 Shu L., Xie M., Wang T. J., Gao D., Chen P. L., Han Y., Li S., Zhuang B. L., Li M. M.:
575 Integrated studies of a regional ozone pollution synthetically affected by subtrop
576 ical high and typhoon system in the Yangtze River Delta region, China. *Atmos.
577 Chem. Phys.*, 16, 15801-15819. doi:10.5194/acp-16-15801-2016, 2016.

578 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G.,

579 Huang, X. Y., Wang, W., and Powers, J. G.: A Description of the Advanced
580 Research WRF Version 3. NCAR Tech Notes-475+STR, 2008.

581 State Environmental Protection Administration of China: China National
582 Environmental Protection Standard: Automated Methods for Ambient Air
583 Quality Monitoring, China Environmental Science Press, Beijing, 2006.

584 Streets, D. G., Fu, J. S., Jang, C. J., Hao, J. M., He, K. B., Tang, X. Y., Zhang, Y. H.,
585 Wang, Z. F., Li, Z. P., Zhang, Q., Wang, L. T., Wang, B. Y., and Yu, C.: Air
586 quality during the 2008 Beijing Olympic Games, *Atmos. Environ.*, 41, 480-492,
587 doi:10.1016/j.atmosenv.2006.08.046, 2007.

588 Wang, X., Westerdahl, D., Chen, L. C., Wu, Y., Hao, J. M., Pan, X. C., Guo, X. B.,
589 and Zhang, K. M.: Evaluating the air quality impacts of the 2008 Beijing
590 Olympic Games: On-road emission factors and black carbon profiles, *Atmos.*
591 *Environ.* 43, 4535-4543, doi:10.1016/j.atmosenv.2009.06.054, 2009a.

592 Wang, W. T., Primbs, T., Tao, S., Zhu, T., and Simonich, S. L.: Atmospheric
593 particulate matter pollution during the 2008 Beijing Olympics, *Environmental*
594 *Science & Technology*, 43, 5314-5320, 2009b.

595 Wang, T., Nie, W., Gao, J., Xue, L. K., Gao, X. M., Wang, X. F., Qiu, J., Poon, C. N.,
596 Meinardi, S., Blake, D., Wang, S. L., Ding, A. J., Chai, F. H., Zhang, Q. Z., and
597 Wang, W. X.: Air quality during the 2008 Beijing Olympics: secondary
598 pollutants and regional impact, *Atmos. Chem. Phys.*, 10, 7603-7615,
599 doi:10.5194/acp-10-7603-2010, 2010.

600 Xie, M., Zhu, K. G., Wang, T. J., Yang, H. M., Zhuang, B. L., Li, S., Li, M. G., Zhu, X.
601 S., and Ouyang, Y.: Application of photochemical indicators to evaluate ozone
602 nonlinear chemistry and pollution control countermeasure in China, *Atmos.*
603 *Environ.*, 99, 466-473, doi:10.1016/j.atmosenv.2014.10.013, 2014.

604 Xie, M., Liao, J., Wang, T., Zhu, K., Zhuang, B., Han, Y., Li, M., and Li, S.: Modeling
605 of the anthropogenic heat flux and its effect on regional meteorology and air
606 quality over the Yangtze River Delta region, China, *Atmos. Chem. Phys.*, 16,
607 6071-6089, doi:10.5194/acp-16-6071-2016, 2016.

608 Xing, J., Zhang, Y., Wang, S. X., Liu, X. H., Cheng, S. H., Zhang, Q., Chen, Y. S.,

609 Streets, D. G., Jang, G., Hao, J. M., and Wang, W. X.: Modeling study on the air
610 quality impacts from emission reductions and atypical meteorological conditions
611 during the 2008 Beijing Olympics, *Atmos. Environ.*, 45, 1786-1798,
612 doi:10.1016/j.atmosenv.2011.01.025, 2011.

613 Xu, H. M., Tao, J., Ho, S. S. H., Ho, K. F., Cao, J. J., Li, N., Chow, J. C., Wang, G. H.,
614 Han, Y. M., Zhang, R. J., Watson, J. G., Zhang, J. Q.: Characteristics of fine
615 particulate non-polar organic compounds in Guangzhou during the 16th Asian
616 Games: Effectiveness of air pollution controls, *Atmos. Environ.*, 76, 94-101,
617 doi:10.1016/j.atmosenv.2012.12.037, 2013.

618 Yarwood, G., Rao, S., Yocke, M., and Whitten G.: Updates to the Carbon Bond
619 chemical mechanism: CB05, Final Report to the US EPA, RT-0400675, 2005.

620 Zhou, D., Li, B., Huang, X., Virkkula, A., Wu, H., Zhao, Q., Zhang, J., Liu, Q., Li, L.,
621 Li, C., Chen, F., Yuan, S., Qiao, Y., Shen, G., and Ding, A.: The Impacts of
622 Emission Control and Regional Transport on PM_{2.5} Ions and Carbon Components
623 in Nanjing during the 2014 Nanjing Youth Olympic Games, *Aerosol & Air
624 Quality Research*, 17, 730-740, doi: 10.4209/aaqr.2016.03.0131, 2017.

625 Zhou, Y., Wu, W., Yang, L., Fu, L. X., He, K. B., Wang, S. X., Hao, J. M., Chen, J. C.,
626 Li, C. Y.: The impact of transportation control measures on emission reductions
627 during the 2008 Olympic Games in Beijing, China, *Atmos. Environ.*, 44,
628 285-293, doi:10.1016/j.atmosenv.2009.10.040, 2010.