

# Author Response and Manuscript Revision

Sonya Fiddes

April 2018

Dear Prof. Takemura,

Please find our response to the Reviewer's comments and our revised manuscript 'Cloud, precipitation and radiation responses to large perturbations in global dimethyl sulfide'. We are grateful for the Reviewer's careful and considered comments and believe that our manuscript has benefited from the close attention provided.

We apologise for not including this cover letter in the previous replies. This may have lead to some confusion about our replies to the inline comments of Reviewer 2, many of which are general and apply throughout the manuscript. We have now highlighted where and how we have made changes in response to all significant comments in the replies. A version of the new manuscript containing the 'tracked changes' (resulting from the latexdiff tool) is also now included.

In particular, both Reviewers commented on the novelty of this study. Whilst we agree that there have been other similar studies, we find that none of them address the question of the role of DMS in the climate system as completely as we do, or with the broad perspectives we do. Specifically, our study considers the complete range of chemical, aerosol, meteorological and climate responses to perturbed DMS fluxes, and discusses these results in the context of climate observational and modelling uncertainty. This is unique in the literature at the present time. Considering these Reviewer comments, and taking suggestions from the Reviewers, have added extra emphasis to this novelty in the manuscript. Furthermore, we have included greater discussion in the introduction and conclusion about the limitations of previous work and continued that discussion throughout the manuscript. Whilst our study does not resolve all of the limitations of the field, we believe, as Reviewer 1 notes, that we address 'current scientific questions, which are not yet fully answered' and as Reviewer 2 notes, these questions are 'timely and relevant'. We have also now provided suggestions as to what directions future studies could take to address these ongoing questions, as suggested by Reviewer 1. We believe that our manuscript, in highlighting these limitations and placing them in a broader perspective, can encourage and provide a starting point for the scientific community to improve our knowledge of this complex and non-linear system.

We hope that you find our responses and revisions satisfactory and that our manuscript is now acceptable for publication in *Atmospheric Chemistry and Physics*.

Kind regards, Sonya Fiddes

# Response to Reviewer 1

We would like to thank Reviewer 1 for their comments on our manuscript. We agree that while similar simulations have been conducted, our focus has been on providing a complete end-to-end evaluation of the role of DMS in the climate system, from changes in the chemistry through to effects on clouds, precipitation and radiation. Furthermore, we have highlighted the problems remaining when trying to model and understand this complex system. We have made these discussion points clearer in both the introduction and conclusion, which has strengthened the arguments presented in this work. Furthermore, we have clarified our methods, which hopefully leaves no remaining ambiguity. We have addressed the specific comments below, and we hope you find our revisions satisfactory.

## General Comments:

*In the abstract not all results are summarized, like the comparison of the control simulation with observations, the second experiment with increased DMS emissions, and the estimate of temperature response to changes in DMS TOA radiative effect. The final sentence of the abstract is too general, the authors should try to briefly explain what kind of model study are further needed and why.*

We have revised the abstract to include the model evaluation, results of both experiments and the temperature sensitivity to DMS flux. We have also made the abstract more specific.

*The explanation given at the end of section 2.1.1 about nudging is not very clear. I understand that doing nudged simulation is better to compare the control model simulation with observation. On the other hand the disadvantage is to not have the full impact on meteorology when comparing the control simulation with the two sensitivity experiments. As far as I understand the model simulation includes both direct and indirect effect of aerosols on radiation, so it is not very clear the sentence at page 4, lines 15-18. Maybe also the paragraph at page 14, lines 3-5 should be included in section 2.1.1.*

We have revised our explanation and justification of using nudging in our simulations in the methods section as suggested (page 4, lines 15-21). The reviewer is correct that the model can simulate direct and indirect aerosol effects. Our use of the word 'direct' in this section may have been misleading and we have rectified this. We have also revised the text on the use of nudging in the discussion section (page 15, line 16).

*The quality of the figures is not always satisfying. In particular the figures which include multiple maps are too small and it is difficult to visualize the fields. Also sometime the colors does not help the data visualization. In particular I would recommend to improve Figures 2,3,4,5,6,8 and 10.*

We have adjusted the colour scales by removing the darkest colours to increase visibility of the figures. We have also removed white space where we can and enlarged the figures. We hope these changes make our figures easier to interpret. We will also ensure the figures are readable after the typesetting process.

*The section 5 is too long and somehow difficult to read. I would recommend to split in two or to shorten it by removing some of the details which are repeated from the previous sections. I would try to explain better the last part of the discussion, providing more details on what kind of experiments are needed to better understand the role of DMS future, considering the ocean acidification, as mentioned in the last paragraph.*

We have significantly shortened this section, cutting out repetitive paragraphs summarising results previously discussed as suggested by the reviewer. We strengthened our discussion involving not only the questions we are addressing but also pointing towards what steps need to be taken for future studies. We are hesitant to include more information about the impacts of ocean acidification as it is beyond the scope of this study (and admittedly, not our area of expertise). However, we do wish to highlight potential implications of ocean acidification on climate that may not have previously been considered.

## Minor Comments:

*Page 3, line 1: Six et al. (2013)*

Typo has been amended

*Page 3, Line 21: the control simulation is not explained, only the two DMS perturbation C2 simulations are described*

We have included a description of the control simulation: 'Three simulations are performed to explore the chemical, aerosol and meteorological implications of large DMS<sub>w</sub> perturbations. In the first experiment, a control simulation is compared to a simulation in which all DMS<sub>w</sub> is removed from the system to determine its current contribution to the climate. In the second experiment, the control simulation is compared to a simulation in which DMS<sub>w</sub> is significantly increased, and the results are compared to that of the work by Grandey and Wang(2015)'. More information about the control simulation is provided in Section 2.1 and 2.3.

*Page 4, Line 1: In this study, . . .*

We are unsure what the reviewer meant here. We do not feel starting the paragraph with 'in this study' is appropriate.

*Page 4, Line 10: did you forget CO2 from the list of GHG gases?*

CO2 has now been explicitly mentioned, it is prescribed to the model as global mean concentration: 'Long-lived greenhouse gas concentrations (e.g.CO<sub>2</sub>,CH<sub>4</sub>, and N<sub>2</sub>O) are prescribed from Coupled Model Intercomparison Project Phase 5 (CMIP5) and RCP6.0 recommendations'

*Page 6, Line 1: 50-440 hPa*

Typo has been amended

*Page 6, Line 17-19: WHY these three regions were chosen? A short motivation should be added. I would include the boundaries of these regions in one of the figures.*

We have included motivation as to why these regions have been chosen and included the boundaries in Figure 2: 'Three regions of interest are defined for their relevance to the broader Australian community (for which ACCESS is purposed) or are of particular interest in the DMS-climate system'.

*Page 6, Line 18: Pacific*

Typo has been amended

*Page 6, Line 29: for the first time the CTL name is used, should be introduced before*

The Ctl name has been introduced a few paragraphs above.

*Page 6, Line31: "without the need for an . . .", I would remove or rephrase as an ensemble experiment of free-running climate simulations is needed to better quantify the impact of DMS forcing on the temperature.*

We have rephrased this sentence as suggested 'In ACCESS-UKCA, an ensemble experiment would be required to provide equivalent temperature difference estimates ...'

*Page 7, Line 9: fraction larger then 0.5 instead of  $\delta$*

Typo has been amended

*Page 8, Line 18: outgoing TOA, LW or SW?*

We have clarified this (SW)

*Page 8, Line 30: (Fig 5c)? positive bias if is the difference between the model and observations, not clear from figure caption.*

We have revised the figure caption for clarity: 'The mean (2000-2009) annual total precipitation of a) the satellite climatology from TRMM (Huffman et al., 2007)), b) the ACCESS-UKCA climatology and c) the difference between ACCESS-UKCA and the TRMM product (model - observations). Units are in year<sup>-1</sup>,

*Page 8, Line 30: over regions to the north and south if the equator, .. but only in the tropics is over 2000 mm/yr*

We have revised this sentence for clarity: 'Precipitation in ACCESS-UKCA has large positive biases in regions that receive the most annual rainfall and align with the Intertropical and South Pacific Convergence Zones (ITCZ/SPCZ). These regions overestimate precipitation by over 2000mm year<sup>-1</sup> ,

*Page 9, Line 4: the figure with % differences is not shown, but the values of the largest % differences could be inserted in the text.*

We have included the % differences in the text as suggested: 'the largest differences occur in the eastern basins of the South Pacific (493% over the SEP region) and South Atlantic Oceans (275% from 0 to 25S,330 to 10E).'

*Page 9, Line 17: The largest absolute differences are in the tropics and mid-latitudes over the Oceans.*  
Sentence amended as suggested: 'The largest absolute differences are in the tropics and mid-latitudes over the oceans'

*Page 10, Line 5: Fig6 g-h is not correct*  
Typo has been amended

*Page 10, Line 21: the largest absolute differences are in clean terrestrial regions. C3 Which regions? Not easy to visualize in the figures. Too small.*  
We have removed reference to these differences due to the difficulty of visualisation.

*Page 11, Line 11: the results presented here suggest a lower CCN . . .*  
Sentence amended as suggested

*Page 11, Line 18: would title the section "Clouds and precipitation response"*  
Title amended as suggested

*Page 11, Line 21-23: this explanation about nudging should be explained also before when describing the experiments.*  
We have revised our explanation of nudging in the Methods section (see previous comment).

*Page 11, Line 32: (see section 3) Page 12, Line 4: remove ) after fig 10a-b*  
Typo has been amended

*Page 13, Section 4.3: Is it possible to put the uncertainties of these estimates of temperature changes per Tg of S emitted?*  
We have performed a moving block bootstrap to determine the uncertainty of the mean changes in radiation, via the 10th and 90th percentile confidence intervals. This can then be translated into a change in temperature by the FAIR model and applied to the flux sensitivity calculations. Further description of how this was performed is now included in the Methods and Tables 3 and 4 have been updated to include these uncertainties.

*Page 14, Line 33: put ref of thomas and mahajan in parenthesis*  
Typo has been amended

*Page 16, Line 3: can you better explain the role of coral-reef derived DMS? Why it is important?*  
We have included a sentence describing why coral reef derived DMS is important (the fact that it is unaccounted for in current global modelling): 'For example, recent studies have indicated that coral reefs produce significant amounts of DMS, and are an unaccounted for source of sulfur (Hopkins et al., 2016; Swan et al., 2017; Jones et al., 2017)'. Note that this section has been significantly revised and re-structured as suggested in the general comments.

*Page 16, Line 8: our results imply that a 25% decrease in . . . would result in an increase of 0.1C. Is it possible to put the uncertainty on this estimate?*  
We have now included confidence intervals for our temperature/flux sensitivity calculations, and the estimates of potential temperature change with decreasing DMS flux.

*Caption of figure 3: third column is not absolute differences, as negative numbers are shown.*  
Caption has been amended

*Caption Figure 5: c) not clear if difference model-obs or contrary while reading the description of the figure in the manuscript.*  
We have revised the caption for clarity (see previous comment)



*Caption Figure 7: blue lines show the SO (Southern Ocean) mean, red the australian (aus). The short name aus is used only here and not in the manuscript.*

We have only shortened Australia to Aus. for clarity in this figure.

## Response to Reviewer 2

We would like to thank Reviewer 2 for their extensive comments.

We acknowledge that there are a number of recent studies examining the role of DMS in the climate system. Our study provides a thorough end-to-end analysis of all aspects of the chemical, aerosol and meteorological impacts of large DMS perturbations in the current climate. To our knowledge, previous studies have not considered the entire system, and typically only considered short time periods (1yr) or future projections. We have made these points clear in the Introduction (page 4, line 1-13), and more specifically noted this on page 4, line 10-13: 'Many of the studies noted above focused on one or two aspects of the DMS-climate system, commonly reporting on the flux<sub>DMS</sub> and its radiative and temperature effects. In this study we evaluate the whole system, examining chemical, aerosol and meteorological changes, including cloud and precipitation effects'

We would also like to thank Reviewer 2 for bringing to our attention the two Tesdal et al. 2016 papers, in addition to the other papers listed in the point by point comments. They have provided a useful reference for discussion around DMS climatology and flux uncertainties. We have included the references suggested by the Reviewer where appropriate, as listed in the point by point comments.

We have taken the Reviewer's advice and have strengthened not only our problem statement in the Introduction (in particular on Page 4 line 18-20: 'We aim to discuss these sensitivities not only within the specific context of the DMS-climate system, as mentioned above, over a 10 year time period, but also in the broader context of the current uncertainties in the DMS-climate system and climate modelling') but also the points mentioned above in the Introduction, which help to better identify gaps in the literature and frame our study.

We agree that just using a different model does not provide sufficient novelty to a study and it was not our intention for this to be a main aim of the paper. By clarifying our problem statement above, we have highlighted the true aim and novelty of this study. ACCESS-UKCA is an established and respected global climate-chemistry model, and for this reason we have not provided any comment comparing ACCESS-UKCA to other global chemistry-climate models. Nevertheless, we have included some more information in the Methods about why ACCESS-UKCA, but more particularly GLOMAP-mode, is a desirable tool (including references suggested by this Reviewer, Page 5, line 31-33: 'A full description of the scheme can be found in Mann et al. (2010) with improvements detailed in Mann et al. (2012). Bellouin et al. (2013) compare GLOMAP-mode with an older generation aerosol scheme, finding significant differences in aerosol response to perturbations between the two schemes'), and further elucidated the novel aspects of the study in the manuscript.

We also agree that the experiments performed here are extreme cases. However, there are significant insights into the DMS-climate system gained from these experiments that will serve to inform more targeted, realistic scenario experiments. Specifically, our large scale perturbations allow quantification of the overall contribution of DMS to the climate system (Section 4.1, page 11). More subtly, this study has identified non-linearities in the DMS-climate system, highlights areas of particular sensitivity, provided evidence of important cloud-aerosol processes and has placed these results in a broad context of our current understanding of the DMS-climate system and the associated uncertainties. We have enhanced these results throughout the Conclusion (in particular the paragraphs, page 16 line 3-20).

### Point by point comments:

*The introduction provides relevant background regarding the role of aerosols in the global radiation budget. It would benefit from more background information regarding the uncertainty in DMS concentrations in*

the ocean. See for example: Belviso et al. (2004), Tesdal et al. (2016a).

We have included further background information regarding the uncertainties surrounding the DMS climatology and flux as suggested by the Reviewer (see paragraph page 2, line 29 - page 3, line 3).

(page 2 line 9): Additional support for the use of DMS fields in climate model are given in Belviso et al. (2004) and Tesdal et al. (2016a), and it is recommended to include these references. Hopkins et al. (2016) is not an appropriate reference for the uncertainty around observed DMS concentrations. Royer et al. (2015) is suggested.

We have revised this section, removed parts and included more appropriate references. It now reads as 'Global DMS concentrations and fluxes remain poorly constrained by observations (Tesdal et al., 2016; Royer et al., 2015), ...'

(page 2 line 10) Charlson et al (1987) is not an appropriate reference for the role of DMS in climate systems being subject to debate, as this is the paper that introduced the hypothesis. The debate came later. A more appropriate article that also serves as a review of the CLAW hypothesis is Ayers and Cainey (2007).

We have included the recommended citation as the Reviewer has suggested: '... its role in the climate system is subject to debate (Ayers and Cainey, 2007; Quinn and Bates, 2011)'.

(page 2 line 16) Add "s" to "contribute"

Typo amended

(page 2 line 23) Add "in certain regions" following "...local DMS concentrations"

Sentence amended as suggested

(page 2 line 29) Add "in order" between "... a one year simulation" and "to quantify its importance. . ."

Sentence amended as suggested

(page 2 line 32) Replace "for example" with "e.g."

Sentence amended as suggested

(page 2 line 34) Is ocean acidification meant to be an example of "anthropogenic climate change"? Please clarify.

This sentence has been clarified to reflect that ocean acidification is caused by anthropogenic emission of CO<sub>2</sub>. '... found DMS emissions were reduced by 17% by the end of the century, primarily due to decreasing ocean pH (caused by anthropogenic CO<sub>2</sub> emissions)'.

(page 3 line 3 and throughout manuscript) Both Celsius and Kelvin temperature units are used throughout. Please use one or the other (preferably Celsius) for consistency.

We have made all temperature units consistent throughout the text

(page 3 line 10) Replace "low-mid level" with "low- and mid-level"

Sentence amended as suggested

(page 3 line 26) Add "analyzes" after "Section 4"

Sentence amended as suggested

(page 4 line 23) Replace "of" with "detailed in"

Sentence amended as suggested

(page 4 line 28) "A full description of the scheme can be found in Mann et al. . ." Here it would be appropriate to mention the study by Bellouin et al. (2013), who compared GLOMAP-mode and CLASSIC and determine that GLOMAP-mode is more sophisticated and accurate.

We have made reference to the Bellouin et al. (2013) paper, as well as the Mann et al (2012) paper. We have retained the Mann et al. (2010) citation as this is the model description paper. This now reads as: 'A full description of the scheme can be found in Mann et al. (2010) with improvements detailed in Mann et al. (2012). Bellouin et al. (2013) compare GLOMAP-mode with an older generation aerosol

scheme, finding significant differences in aerosol response to perturbations between the two schemes'

(page 5 line 1) *"Significant sampling biases...Northern Hemisphere (Lana et al 2011)" This sentence is not entirely clear. Is the point that the data is biased towards springsummer and towards Northern Hemisphere?*

Yes, we have clarified this sentence to reflect this: 'Significant sampling biases exist within the Lana et al. (2011) data set, with approximately half of observations collected in late spring through summer, and more than two-thirds of the data collected in the Northern Hemisphere.'

(page 5 line 2) *Replace "spring-summer" with "spring through summer"*  
Sentence amended as suggested

(page 5 line 5) *"The Liss and Merlivat (1986) parameterization. . ." There should be justification for why this parameterization is used. Several different parameterizations of the piston velocity in terms of wind speed have been used in modelling studies (e.g., Liss and Merlivat, 1986; Wanninkhof, 1992; Nightingale et al., 2000), leading to substantially different flux fields for a given concentration field (Tesdal et al., 2016a).*

We have made reference to the alternative flux parameterisations as well as the modelling studies that detail their uncertainties in the Introduction (page 2, line 29 - page 3, line 3). Of the parameterisations available within ACCESS-UKCA, we chose the Liss Merlivat (1986) method because it agrees more closely with the latest generation of DMS flux parameterisations, e.g. Vlahos Monahan (2009), Bell et al. (2017), which were specifically derived for DMS, unlike the above parameterisations. We have updated the text to reflect this: 'The most common flux parameterisations exhibit considerable ranges in flux<sub>DMS</sub>, from 15 – 35 Tg year<sup>-1</sup> of sulfur in Elliott (2009) to 9-34 Tg year<sup>-1</sup> of sulfur found in Tesdal et al. (2016), who recommend a range of 18-24 Tg year<sup>-1</sup> of sulfur as a reasonable estimate. Vlahos and Monahan (2009) and Bell et al. (2017) show that current parameterisations overestimate the flux<sub>DMS</sub> at high wind speeds and suggest that annual global flux<sub>DMS</sub> is likely to be in the lower range of current estimates. Of the flux<sub>DMS</sub> parameterisations available in ACCESS-UKCA, the Liss and Merlivat (1986) method yields a low to moderate flux comparable to those calculated in Vlahos and Monahan (2009) and Bell et al. (2017), and is used in this study.'

(page 5 line 16-17) *Following Eq. 4, Replace "Where" with "Here," . Insert "is" before "determined following the method of Saltzman.."*  
Sentence amended as suggested

(page 5 line 20) *"seawater" should not be*  
hyphenated Typo amended

(Section 2.2 Model Evaluation) *The evaluation should include comparison to observation of atmospheric concentration of DMS and other sulfur species, for example as described in Tesdal et al. (2016b).*

We appreciate that an evaluation of atmospheric DMS and other sulfur species is desirable, as done in the Tesdal et al. (2016b) study. However, we have limited this evaluation to meteorological data sets that are globally available from satellite products. We are confident in the use of this model for DMS studies due to GLOMAP-mode being extensively evaluated against observations in many previous studies (Mann et al. 2010, 2012, Woodhouse et al 2010, 2013, 2015).

(page 6 line 1) *add "between" between "medium" and "440-680 hPa" and between "low" and "680-1000 hPa"*  
Sentence amended as suggested

(page 6 line 17) *Add period at end of sentence that begins "The Australian region...."* (page 6 line 24) *Replace "that" with "the one"*  
Sentence amended as suggested

(page 6 line 30) *"By providing this radiative effect. . ." The text appears to imply that FAIR is a tool that provides estimates of climate response, given a simple value of radiative effect. Thus, by feeding estimates of radiative effects to FAIR one can analyze the effect on temperature and other climate variables. The text would benefit from more elaboration regarding the relationship between the radiative effect*

*and the FAIR climate component.*

We have included some more information on how the FAIR emulator links radiative forcing changes to temperature and we have included some additional references for readers to pursue. This section now reads as: 'FAIR's climate component is a simple impulse response model which emulates the behaviour of more complex Earth System Models, given a certain radiative forcing (in this case due to DMS). FAIR has been designed to determine temperature responses to radiative forcing of similar magnitudes to the DMS radiative effect (Millar et al., 2017). FAIR's temperature response is calculated as the sum of two components, approximately representing the response of the upper mixed layer and deep ocean to a change in radiative forcing (Millar et al., 2017). Due to its simplicity, FAIR cannot capture the non-linearities and feedbacks in the climate system, and hence the temperature response calculated must be taken as an estimate only. Furthermore, in this work we consider only a single mid-range estimate of climate sensitivity'

*(page 7 line 12) Add comma after "e.g."*

Sentence amended as suggested

*(page 8 line 12) "...instead reflecting more sunlight thus enhancing the albedo..." This clause is confusing. What is it that is reflecting more sunlight?*

We have restructured this sentence to be clearer: 'Over the Antarctic ice sheets, both TOA outgoing and surface incoming SW radiation are overestimated, due to an underestimation of low clouds which allows the high albedo to reflect too much incoming SW radiation back out to space.'

*(page 8 line 20) Replace "...low clouds allowing.." with "...low clouds, which allows. . ."*

Sentence amended as suggested

*(page 8 line 24) "This estimation is slightly greater. . ." Greater than what?*

We have amended this sentence to say 'greater than that found for CMIP5 GCMs'

*(page 9 line 20) Replace "representing" with "which is due to" or "which represents" to avoid awkward verb-gerund construction ("...representing lofting...").*

Sentence amended as suggested

*(page 9 line 23) Use appropriate subscript formatting for H<sub>2</sub>SO<sub>4</sub>.*

Typo amended

*(page 9 line 25) Replace "...in new particle formation, forming secondary sulfate aerosol. . ." with "...in formation of secondary sulfate aerosols,..."*

Sentence amended as suggested

*(page 9 line 26) Correct formatting of H<sub>2</sub>SO<sub>4</sub>.*

Typo amended, this has been checked throughout the text

*(page 10 line 5 and throughout manuscript) Replace "Whilist" " with "While"*

This has been amended throughout the manuscript as suggested

*(page 10 line 26) Replace "cloud condensation nucleii" with "CCN"*

Sentence amended as suggested

*(page 10 line 26-33) "Fig. 8d-e show the number. . ." The paragraph does not clearly describe the figures; sentence structure is awkward. Consider revision as follows: Fig. 8d-e show the number concentration of CCN with dry diameters greater than 70 nm (CCN<sub>70</sub>) for the Ctl and the differences resulting from Exp.1. The largest absolute differences are in the tropics, which, similarly to the N3, have the highest concentration. Relatively, there is a global decrease of 5%, whilst decreases of 7% were found over the Australian region, decreases of 8% over the SO and decreases of 20% over the SEP. Differences in cloud droplet number (CDN) are shown in Fig. 8g-h. The relative differences in CDN (Exp.1-Ctl) show a similar spatial pattern to that of the CCN. Global mean CDN decreases by 5%. A decrease of 5% is also found for the Australian region, whereas the SO shows an 8% decrease, and the SEP shows an 18% decrease. In both the CCN<sub>70</sub> and CDN, the marine Southern Hemisphere mid-latitudes have the largest*

decreases of 14% (averaged between 5-35 $\ddot{E}$ ŠS) despite the SO having some of the larger decreases in SO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub>.

The paragraph has been amended as suggested by the Reviewer.

(page 10 line 30) "Exp 1- Ctl" Is this explaining the ratio for relative difference? It is not given elsewhere when talking about relative difference.

The 'Exp 1- Ctl' notation simply implies Experiment 1 minus the Control. We have removed it to avoid confusion.

(page 11 line 1) Replace "tropics mid-latitudes" with "mid-latitude tropics".

Sentence has been amended

(page 11 line 13) "Similar CCN sensitivities are reported in the Woodhouse et al. (2010) study." State actual values reported in the reference.

Actual values have been provided 'Similar CCN sensitivities are reported in the Woodhouse et al. (2010) study (63cm<sup>3</sup>/mg m<sup>2</sup> day<sup>1</sup> global average).'

(page 11 line 24) "...radiation scheme via aerosols and some (see Section 2.1.1)." Missing word/phrase following "some".

Missing words was 'gases'

(page 11 line 28) "...slightly removed from the coastline." Not clear. From the Southern Hemisphere's coastline? Recommend revising the whole first sentence, ending with a period after "(Fig. 9a)" and then beginning a new sentence. This would help to clarify what difference is seen in the SH generally and would parallel the structure used for the rest of the paragraph.

We have revised this sentence and removed the confusing reference to the coastline: 'The largest differences occur in eastern basins of the Southern Hemisphere's oceans. '

(page 11 line 32) "...see Section 3 in comparison to other areas. . ." Confusing. The author is comparing all of Section 3 to the areas of low cloud formation?

A typo has been amended in the sentence (a missing bracket after Section 3) which resolves the clarity of this section.

(page 11 line 35) "...biases exacerbate this. . ." Clarify what "this" refers to.

We have clarified what 'this' refers to (the differing responses of areas of low cloud formation): '... whether the long standing model biases, especially those around the formation of supercooled liquid water, have contributed to the differing responses requires further investigation'

(page 12 line 4) Complete parentheses for "shown in Fig. 10a-b"

Typo amended

(page 12 line 8) Capitalize "Southern Hemisphere"

Typo amended

(page 12 line 13) Add "the" before "Ctl"

Sentence amended as suggested

(page 12 line 25) Delete the comma after "magnitude"

Sentence amended as suggested

(page 12 line 28) Revise text to "...the SEP: increases of 42%, 172%, and 89% respectively."

Sentence amended as suggested

(page 12 line 29) Insert "occurs" after "...a decrease of 37%"

Sentence amended as suggested

(page 12 line 33) Insert "respectively" after "...increases by 6%, 4% and 5%"

Sentence amended as suggested

(page 12 line 34) Insert "the" before "...SEP of 14%"  
Sentence amended as suggested

(page 13 line 2) Replace "showing" with "which show"  
Sentence amended as suggested

(page 13 line 2) Suggest "Incoming surface SW radiation" rather than "Surface incoming SW radiation. . ."  
Sentence amended as suggested

(page 13 line 6) Replace "analyzed under" with "which analyzed"  
Sentence amended as suggested

(page 13 line 8) Replace "south east" with "southeast"; Replace "these results presented" with "the results presented here"  
Sentence amended as suggested

(page 13 line 10) Insert "however" between "...the surface," and "the precipitation. . ."  
Sentence amended as suggested

(page 13 line 18) Delete "Although"; insert "and" between "...warming" and "we. . ."  
Sentence amended as suggested

(page 13 line 21) Delete comma after "...study"  
Typo amended

(page 13 line 26) Replace "on" with "in"  
Typo amended

(page 13 line 30) Delete duplicate "the"; insert "that" between "...this study is" and "the model. . ."; replace "underestimation of" with "underestimates"  
Sentence amended as suggested

(page 14 line 1) Replace "rather a multitude of theories" with "multiple theories have been proposed"  
Sentence amended as suggested

(page 14 line 7) "representation" should be plural  
Typo amended as suggested

(page 14 line 13) Change "Figure 1" to "Fig. 1" for consistency with rest of manuscript  
Sentence amended as suggested

(page 14 line 14) Delete comma after SO<sub>2</sub>  
Sentence amended as suggested

(page 14 line 32) Break run-on sentence: Period after "importance". Next sentence: "Instead, the studies highlighted SO (Thomas et al., 2010; Mahajan et al., 2015)." Highlighted the SO in terms of what parameter(s)?

These studies have focused on cloud feedbacks, which we have clarified in the text: '... instead, these studies focused on cloud feedbacks in the SO'

(page 14 line 35) Revise sentence as: "...slightly lower than the estimation of 2.03 Wm<sup>2</sup> by Thomas. . ."  
Sentence amended as suggested

(page 15 line 1) Replace "are" with "were"  
Sentence amended as suggested

*(page 15 line 5) Revise reference format: (Albrecht, 1989)*

Sentence amended as suggested

*(page 15 line 16) Revise placement of subordinate clause in the sentence beginning "The cause of the discrepancy. . ." as: "Without further information, it is difficult to speculate on the cause of the discrepancy. . ."*

Sentence amended as suggested

*(page 15 line 18) "Suggest" should be plural; delete comma after constraints*

Sentence amended as suggested

*(page 15 line 22) Replace "Whilst" with "Though"*

Sentence amended as suggested

*(page 15 line 23) Replace the colon after "For example" with a comma; replace "an as of yet" with currently; replace "for" with "as a"*

Sentence amended as suggested

*(page 15 line 31) Replace the semicolon after "Six et al (2013)" with "and"*

Sentence amended as suggested

*(page 16 line 3) "coral-reef-derived" (2 hyphens), or "DMS derived from coral reefs"*

Sentence amended as suggested

*(page 16 line 5) Replace semicolon after "Hopkins et al. (2011)" with a comma*

Sentence amended as suggested

*(page 16 line 8) Insert "those" between "than" and "found"*

Sentence amended as suggested

*(page 16 line 12) Make example given in this sentence parenthetical: "...increases (e.g., via solar radiation management) may have a short term cooling effects, however, without. . ."*

Sentence amended as suggested

*(page 16 line 13) Delete the "a" between "...may have" and "short term..."; insert comma after "...however"*

Sentence amended as suggested

*(page 16 line 14) Delete "on" between "...impact" and "marine life. . ."*

Sentence amended as suggested

# Cloud, precipitation and radiation responses to large perturbations in global dimethyl sulfide

Sonya L. Fiddes<sup>1,2,3</sup>, Matthew T. Woodhouse<sup>3</sup>, Zebedee Nicholls<sup>1</sup>, Todd P. Lane<sup>4</sup>, and Robyn Schofield<sup>2</sup>

<sup>1</sup>Australian-German Climate and Energy College, University of Melbourne, Parkville, 3010 Australia

<sup>2</sup>ARC Centre of Excellence for Climate System Science, School of Earth Sciences, University of Melbourne, Parkville, 3010, Australia

<sup>3</sup>Climate Science Centre, Oceans and Atmosphere, Commonwealth Scientific and Industrial Research Organisation, Aspendale, 3195, Australia

<sup>4</sup>ARC Centre of Excellence for Climate Extremes, School of Earth Sciences, University of Melbourne, Parkville, 3010, Australia

*Correspondence to:* S. Fiddes

(sonya.fiddes@climate-energy-college.org)

**Abstract.** Natural aerosol emission represents one of the largest uncertainties in our understanding of the radiation budget. Sulfur emitted by marine organisms, as dimethyl sulfide (DMS), constitutes one-fifth of the global sulfur budget and yet the distribution, fluxes and fate of DMS remain poorly constrained. This study evaluates the Australian Community Climate and Earth System Simulator (ACCESS) - United Kingdom Chemistry and Aerosol (UKCA) model in terms of cloud fraction, radiation and precipitation, and then quantifies the role of DMS in the chemistry-climate system. We find that ACCESS-UKCA has similar cloud and radiation biases to other global climate models. By removing all, or alternatively significantly enhancing marine DMS, we find a top of atmosphere radiative effect of 1.7 and -1.4  $\text{Wm}^{-2}$  respectively. The largest responses to these DMS perturbations (removal/enhancement) are in stratiform cloud decks in the Southern Hemisphere's eastern ocean basins. These regions show significant differences in low-cloud (-9/+6%), surface incoming shortwave radiation (+7/-5  $\text{Wm}^{-2}$ ) and large-scale rainfall (+15/-10%). We demonstrate a precipitation suppression effect of DMS-derived aerosol in stratiform cloud deck regions due to DMS, coupled with an increase in low cloud fraction. The difference in low cloud fraction is an example of the aerosol lifetime effect. Globally, we find a sensitivity of temperature to annual DMS flux of 0.027 and 0.019 K per  $\text{Tg year}^{-1}$  of sulfur, respectively. Other areas of low cloud formation, such as the Southern Ocean and stratiform cloud decks in the Northern Hemisphere, have a relatively weak response to DMS perturbations. We highlight the need for greater understanding of the DMS-climate cycle within the context of uncertainties and biases of climate models as well as those of DMS-climate observations.

## 1 Introduction

Current understanding of the global climate is underpinned by the concept of the radiation budget; the balance of energy entering and leaving the Earth's atmosphere. Aerosols play an important role in this budget, having direct (McCormick and Ludwig, 1967) and indirect effects via cloud processes (Twomey, 1974; Albrecht, 1989). Aerosols produce a net cooling effect



at the surface, with the total aerosol effective radiative forcing estimated as  $-0.9 \text{ W m}^{-2}$  by the most recent Intergovernmental Panel on Climate Change (IPCC) report, ~~substantially offsetting~~. The aerosol radiative forcing substantially offsets the effect of well mixed greenhouse gases effective radiative forcing of  $2.8 \text{ W m}^{-2}$  (Myhre et al., 2013b). However, large uncertainty in aerosol radiative forcing remains ( $\pm 0.5 \text{ W m}^{-2}$  in the 2013 IPCC report), and is in fact the largest source of uncertainty to the overall radiation budget for the current climate (Myhre et al., 2013b; Carslaw et al., 2013). Uncertainties due to aerosols affect not only the radiation budget, but also chemical and meteorological parameters such as ozone concentration and photolysis (Kushta et al., 2014), cloud formation, albedo, temperature and precipitation (Seinfeld et al., 2016; Rotstayn et al., 2015; Rosenfeld et al., 2014).

Natural aerosol sources account for the largest portion of this uncertainty, explaining up to 45% of the variance of aerosol forcing, compared to 34% from anthropogenic aerosol emissions (Carslaw et al., 2013). ~~Second to volcanic activity~~, DMS produced by marine organisms makes up approximately 19% of global sulfur emissions, producing a DMS flux ( $\text{flux}_{\text{DMS}}$ ) of  $17.6 \text{ Tg year}^{-1}$  (Sheng et al., 2015), though estimates range from ~~15-32-9-35~~  $\text{Tg year}^{-1}$  (~~Elliott, 2009; Woodhouse et al., 2010~~) ~~of sulfur~~ (Belviso et al., 2004a; Elliott, 2009; Woodhouse et al., 2010; Tesdal et al., 2016). Global DMS concentrations and fluxes remain poorly constrained by observations ~~and in climate models~~ (Hopkins et al., 2016) (Tesdal et al., 2016), and its role in the climate system is subject to debate (~~Charlson et al., 1987; Quinn and Bates, 2011~~) (Ayers and Cainey, 2007; Quinn and Bates, 2011).

Charlson, Lovelock, Andreae and Warren (CLAW) proposed a hypothesis by which marine organisms, primarily phytoplankton, regulate their environment via the increased production of dimethyl sulfonium propionate (DMSP) when stressed, for example due to warm sea surface temperatures (SSTs) (Charlson et al., 1987). DMSP is degraded via bacterial processes to DMS in the ocean (Yoch, 2002), some of which is vented into the atmosphere (Charlson et al., 1987). Once in the atmosphere DMS has a lifetime of 1-2 days (Kloster et al., 2007; Korhonen et al., 2008), and oxidises to form sulfuric acid and ultimately contributes to the aerosol burden. This additional source of aerosol can directly or indirectly influence the radiation budget and potentially cool local SSTs (although this has not been proven in the literature), hence reducing phytoplankton stress. The DMS-climate system is summarised in Fig. 1.

Current understanding of the DMS-climate system implies that no bio-regulatory feedback exists as proposed by the CLAW hypothesis (Quinn and Bates, 2011; Woodhouse et al., 2013). However, observations show that seasonal cloud condensation nuclei (CCN) variability cannot be explained without a contribution from DMS (Korhonen et al., 2008; Vallina et al., 2006), implying DMS is important for the longer term climate. Complicating this problem is our ~~poor~~ limited understanding of the global distribution of DMS, ultimately relying on the collection of observations collated and interpolated by Lana et al. (2011), which may ~~considerably misrepresent~~ not capture local DMS concentrations in certain regions such as over coral reefs (Hopkins et al., 2016) and at the poles (Mungall et al., 2016; Kim et al., 2017).

~~The relationship of DMS with~~ A number of studies have parameterised global oceanic DMS (or DMSP) concentrations, using primary productivity, insolation, SSTs and other fields as predictors (Anderson et al., 2001; Simó and Dachs, 2002; Belviso et al., 2004b; Gali et al., 2015). However, numerous issues arise when trying to parameterise oceanic DMS, including the lack of observations as mentioned, but also that DMS production is species dependent and predictors are not uniformly

relevant across marine biota. Halloran et al. (2010) find that two older parameterisations of DMS concentration perform little better than the Kettle and Andreae (2000) climatology. Further uncertainty in  $\text{flux}_{\text{DMS}}$  arises from the parameterisation of the sea-air flux mechanism. Several parameterisations of  $\text{flux}_{\text{DMS}}$  exist (e.g. Liss and Merlivat, 1986; Wanninkhof, 1992; Nightingale et al., 2000), resulting in a large range of annual global  $\text{flux}_{\text{DMS}}$  estimates, for example, 15-35 Tg year<sup>-1</sup> of sulfur (Elliott, 2009) or 9-35 Tg year<sup>-1</sup> of sulfur (Tesdal et al., 2016). Both these uncertainties (in climatology and flux) can have significant consequences for our understanding of climate.

The importance of DMS in large-scale climate has been highlighted by numerous global modelling studies. Mahajan et al. (2015) (using the Lana et al. (2011) DMS climatology) and Thomas et al. (2010) (using the Kettle et al. (1999) DMS climatology) found DMS to have a radiative effect of -1.79 W m<sup>-2</sup> and -2.03 W m<sup>-2</sup> at the top of the atmosphere (TOA) respectively. Thomas et al. (2011) doubled ~~DMS~~, surface water DMS concentrations, ( $\text{DMS}_w$ ), finding a TOA radiative effect of -3.42 W m<sup>-2</sup>. These studies ~~removed DMS from~~ perturbed DMS in the climate system ~~in a one-year simulation to quantify its importance on climate and noted~~ in order to quantify the effect on climate, and noted that the largest changes in radiation and cloud microphysics occurred in the Southern Ocean and southern Pacific and Indian Oceans.

Other modelling studies have explored the impact of anthropogenic climate change on marine DMS production, often with opposing conclusions, making it unclear whether marine DMS production would increase or decrease with warming temperatures (for example Bopp et al., 2004; Gabriele et al., 2004; Kloster et al., 2007; Cameron-Smith et al., 2011). ~~More recent laboratory experiments have confirmed that under anthropogenic climate change, and in particular ocean acidification, marine organisms produce significantly less DMS (Hopkins et al., 2011). A global modelling study by Six et al. (2013) (e.g. Bopp et al., 2004) found DMS emissions were reduced by 17% by the end of the century, primarily due to decreasing ocean pH. Six et al. also explored the resultant feedback of these changes in DMS on the climate, as outlined by Fig. 1. The reduced DMS flux (caused by anthropogenic CO<sub>2</sub> emissions). The reduced  $\text{flux}_{\text{DMS}}$  was found to cause an additional 0.23-0.48 K of warming by the end of the century (Six et al., 2013). Reduced DMS flux- $\text{flux}_{\text{DMS}}$  due to ocean acidification is also found modelled by Schwinger et al. (2017), who found under the Representative Concentration Pathway (RCP) 8.5 to the year 2200, DMS production decreases by 48% assuming a strong sensitivity of DMS production to changes in pH. The authors Schwinger et al. (2017) calculated a DMS temperature sensitivity of -0.041 K per Tg year<sup>-1</sup> of sulfur.~~

~~Conversely,~~ Laboratory experiments have found that under ocean acidification, marine organisms produce significantly less DMS (Hopkins et al., 2011). However, more recently Hopkins et al. (2018) report that polar planktonic communities show resilience to ocean acidification. The Six et al. (2013) study found an increased  $\text{flux}_{\text{DMS}}$  in the polar regions, while the Schwinger et al. (2017) study did not.

Grandey and Wang (2015) attempted to determine if a significant artificial increase of marine DMS production (due to, for example, ocean fertilisation) in the oceanic ecosystem could offset future warming trends. ~~Under such a scenario of drastically increased atmospheric~~ Under a scenario where  $\text{flux}_{\text{DMS}}$  is increased to 46.1 Tg year<sup>-1</sup> of sulfur, Grandey and Wang found global temperature increases due to anthropogenic climate change under RCP 4.5 were partially offset, primarily due to ~~low-mid-level~~ low- and mid-level cloud feedbacks resulting in a ~~change to the radiative effect~~ radiative flux perturbation of -2.0 W m<sup>-2</sup>. Regional changes in precipitation (both increases and decreases) were also noted, up to as much as 30%.

The direct aerosol effect can be approximately linearly related to aerosol concentration (Myhre et al., 2013a). By contrast aerosol-cloud processes, or the secondary aerosol indirect effects, have large uncertainties, with implications for the radiation budget (Myhre et al., 2013b). Global climate models are currently unable to capture many key physical and chemical processes and interactions in the aerosol-cloud system (Rosenfeld et al., 2014). These shortcomings add further uncertainty to quantification of the DMS-climate system. Quantification of model performance is essential in providing context and perspective to any modelling experiment.

Many DMS-climate modelling studies consider DMS under future scenarios (Bopp et al., 2004; Gabric et al., 2004; Kloster et al., 2007; Cameron-Smith et al., 2011; Six et al., 2013; Grandey and Wang, 2015; Schwinger et al., 2017). However, it is clear that our understanding of DMS in the current climate is not yet fully established, considering both modelling and observational uncertainties. Studies exploring DMS changes under current climate conditions (Thomas et al., 2010; Woodhouse et al., 2010; Mahajan et al., 2015) have completed short simulations (approximately one year), which are too short to be indicative of a true climatological response. Furthermore, uncertainties related to DMS emission and fate in the atmosphere are not the only barriers to the DMS-climate question. Climate model uncertainties and biases must also be considered, which have not previously been provided.

The interactions between DMS-derived sulfur, its oxidation products and the atmosphere can be highly non-linear, vary regionally and have far-reaching impacts on multiple processes in the climate system (Thomas et al., 2011). ~~These interactions merit further study and may have important implications in addition to the more commonly reported effects of DMS on global temperature and radiation, for example its influence on cloud and precipitation.~~ Many of the studies noted above focused on one or two aspects of the DMS-climate system, commonly reporting on the  $\text{flux}_{\text{DMS}}$  and its radiative and temperature effects. In this study we evaluate the whole system, examining chemical, aerosol and meteorological changes, including cloud and precipitation effects.

This study has two aims, the first of which is to assess the suitability of the ACCESS (Australian Community Climate and Earth System Simulator) -UKCA (United Kingdom Chemistry and Aerosol) model for examining the role of DMS in the Earth's climate in terms of low, medium and high cloud cover, outgoing TOA shortwave (SW) and longwave (LW) radiation, incoming surface SW radiation, and precipitation. Secondly, ACCESS-UKCA is used to test the large-scale sensitivity of the present day climate to prescribed changes in ~~surface water DMS concentrations,  $\text{DMS}_{\text{w}}$ . Two simulations are performed to explore the chemical, aerosol and meteorological implications of perturbations. In the first simulation, all is removed from the system to determine its current contribution to the climate. In the second experiment, is significantly increased, and the results are compared to that of the work by Grandey and Wang (2015).~~

Three simulations are performed to explore the chemical, aerosol and meteorological implications of large  $\text{DMS}_{\text{w}}$  perturbations. In the first experiment, the control simulation is compared to a simulation in which all  $\text{DMS}_{\text{w}}$  is removed from the system to determine its current contribution to the climate. In the second experiment,  $\text{DMS}_{\text{w}}$  is significantly increased, and the results are compared to that of the work by Grandey and Wang (2015).

This paper is organised as follows: Section 2 outlines the methodology used in this study, Section 3 evaluates how well ACCESS-UKCA performs with respect to certain satellite products, Section 4 analyses the sensitivity of the ACCESS-UKCA climate to large perturbations in DMS ~~is analysed~~ and Section 5 provides some discussion and concluding remarks.

## 2 Methods

### 2.1 Model description and set-up

#### 2.1.1 ACCESS-UKCA

The ACCESS-UKCA coupled climate-chemistry model is a platform from which the influences of DMS on the large-scale climate can be evaluated. The physical atmosphere in the ACCESS model is the United Kingdom Met Office's Unified Model (UM). In this case, UM version 8.4 is used, in conjunction with the UKCA chemistry model (Abraham et al., 2012), which includes the GLObal Model of Aerosol Processes (GLOMAP)-mode aerosol scheme described in Section 2.1.2.

Horizontal grid resolution is  $1.25^\circ$  latitude x  $1.85^\circ$  longitude, with 85 vertical levels, where the model top is located at 85 km. Anthropogenic emissions are prescribed pre-2000 ~~by~~ from the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP) (Lamarque et al., 2010), and post-2000 ~~by~~ from the Representative Concentration Pathway 6.0 (van Vuuren et al., 2011). Biomass burning emissions are from the GFED4s database (van der Werf et al., 2017). Emissions of other species required by ACCESS-UKCA, and their original sources, including biogenic emissions, chemical precursors and primary aerosol, are described in detail in Woodhouse et al. (2015). DMS emissions are calculated within UKCA, and are described in Section 2.1.3. Long-lived greenhouse gas concentrations (e.g. CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) are prescribed from Coupled Model Intercomparison Project Phase 5 (CMIP5) and RCP6.0 recommendations. Monthly mean SST and sea ice coverage are prescribed as per the Atmospheric Model Intercomparison Project (AMIP) (Taylor et al., 2015). UKCA is coupled to the ACCESS radiation scheme via O<sub>3</sub>, CH<sub>4</sub>, N<sub>2</sub>O and aerosol (direct scattering and absorption). Aerosols further influence the large-scale cloud and precipitation schemes via the cloud droplet ~~number~~ number (CDN) concentration, allowing changes in the the chemical/aerosol fields to affect the meteorology.

For this study, ACCESS-UKCA is run for the years 2000-2009, with a one year spin-up, 1999. ~~The simulations are nudged to ERA-Interim (Dee et al., 2011), using the horizontal wind component and potential temperature, at 6 hourly intervals in the free troposphere. Whilst performing a nudged simulation limits the ability of the model to respond to the changes in sulfate aerosol, a free running simulation would not allow us to separate direct aerosol responses from the subsequent process feedbacks that may occur. In this theoretical study, we are interested in the direct impacts of sulfate aerosol on the climate, including the atmospheric composition as well as its effects on radiation, cloud and precipitation.~~ The simulations are nudged to ERA-Interim (Dee et al., 2011), using the horizontal wind component and potential temperature, at 6 hourly intervals in the free troposphere. The use of nudging does not allow aerosol and cloud responses to perturbed DMS to affect synoptic scale meteorology, hence the results here represent instantaneous responses in the climate system. Nudging was deemed desirable for this study to limit computational expense, allowing single runs of 10 years. Due to nudging, responses in the simulation may

be dampened, but can be attributed directly to the DMS perturbations. The complicating effect of internal variability within the modelled system is also avoided in nudged simulations.

### 2.1.2 GLOMAP

The GLOMAP-mode aerosol scheme uses two-moment pseudo-modal aerosol dynamics to simulate aerosol size distributions (Mann et al., 2010, 2012). GLOMAP-mode simulates particle compositions with sulfate, sea-salt, elemental and organic carbon in internally-mixed modes (Mann et al., 2010). Dust is treated outside of GLOMAP-mode according to the scheme of detailed in Woodward (2001). Processes simulated within GLOMAP-mode include primary emissions, new particle formation, particle growth by coagulation, condensation and cloud processing and removal by dry deposition, and in-cloud and below-cloud scavenging (Mann et al., 2010). New particle formation occurs via two mechanisms in ACCESS-UKCA; free tropospheric binary homogeneous nucleation (Kulmala et al., 1998) and organic-mediated boundary layer nucleation (Metzger et al., 2010). The aerosol size distribution is represented in four soluble modes (corresponding to nucleation, Aitken, accumulation and coarse size modes) and one insoluble mode (Aitken). A full description of the scheme can be found in Mann et al. (2010) with improvements detailed in Mann et al. (2012). Bellouin et al. (2013) compare GLOMAP-mode with an older generation aerosol scheme, finding significant differences in aerosol response to perturbations between the two schemes.

### 2.1.3 DMS<sub>w</sub> climatology and flux parameterization

The number of DMS<sub>w</sub> observations have increased dramatically over the last three decades (Kettle et al., 1999; Lana et al., 2011), although significant gaps, both spatially and temporally, remain. Lana et al. (2011) use observations to derive a gridded DMS<sub>w</sub> climatology, [which is](#) used in this study and shown in Fig. 2a. The Lana et al. (2011) climatology shows that high latitude regions have the highest DMS<sub>w</sub> concentrations. Significant sampling biases exist ~~with approximately half of observations collected in late spring – summer, and more than two thirds of the data in the Northern Hemisphere (Lana et al., 2011).~~ within the Lana et al. (2011) data set, with approximately half of observations collected in late spring through summer, and more than two-thirds of the data collected in the Northern Hemisphere.

The flux<sub>DMS</sub> from the ocean to the atmosphere remains poorly parameterised, has large variability in space and time and cannot easily be measured. This subsequently causes large uncertainties in the flux<sub>DMS</sub> parameterisation, ~~a summary of which can be found in Elliott (2009). The Liss and Merlivat (1986) parameterisation is used here. In this method, is given by.~~ The most common flux parameterisations exhibit considerable ranges in flux<sub>DMS</sub>, from 15-35 Tg year<sup>-1</sup> of sulfur in Elliott (2009) to 9-34 Tg year<sup>-1</sup> of sulfur found in Tesdal et al. (2016), who recommend a range of 18-24 Tg year<sup>-1</sup> of sulfur as a reasonable estimate. Vlahos and Monahan (2009) and Bell et al. (2017) show that current parameterisations overestimate the flux<sub>DMS</sub> at high wind speeds and suggest that annual global flux<sub>DMS</sub> is likely to be in the lower range of current estimates. Of the flux<sub>DMS</sub> parameterisations available in ACCESS-UKCA, the Liss and Merlivat (1986) method yields a low to moderate flux comparable to those calculated in Vlahos and Monahan (2009) and Bell et al. (2017), and is used in this study.

The Liss and Merlivat (1986)  $\text{DMS}_{\text{flux}}$  parameterisation is described in Eq. 1 where  $k$ , the piston velocity, is parameterised under three wind induced sea surface regimes: smooth (Eq. 2) and rough (Eq. 3) gas transfer, and a wave breaking/bubble-bursting regime (Eq. 4).

$$\text{DMS}_{\text{flux}} = k(\text{DMS}_{\text{w}} - \frac{\text{DMS}_{\text{a}}}{\alpha}) = k(\text{DMS}_{\text{w}}\alpha - \text{DMS}_{\text{a}}) \quad (1)$$

5 For  $w_{10} < 3.6 \text{ ms}^{-1}$ :

$$k = 0.17w_{10}\left(\frac{SC_{\text{DMS}}}{600}\right)^{\frac{2}{3}} \quad (2)$$

For  $3.6 \text{ ms}^{-1} < w_{10} < 13 \text{ ms}^{-1}$ :

$$k = 2.85(w_{10} - 3.6)\left(\frac{SC_{\text{DMS}}}{600}\right)^{\frac{1}{2}} + 0.612\left(\frac{SC_{\text{DMS}}}{600}\right)^{\frac{2}{3}} \quad (3)$$

For  $w_{10} > 13 \text{ ms}^{-1}$ :

$$10 \quad k = 5.9(w_{10} - 13)\left(\frac{SC_{\text{DMS}}}{600}\right)^{\frac{1}{2}} + 26.79(w_{10} - 3.6)\left(\frac{SC_{\text{DMS}}}{600}\right)^{\frac{1}{2}} + 0.612\left(\frac{SC_{\text{DMS}}}{600}\right)^{\frac{2}{3}} \quad (4)$$

~~Where~~ Here  $\text{DMS}_{\text{w}}$  solubility  $\alpha = 11.4$  at  $26^\circ\text{C}$ ,  $w_{10} = 10 \text{ m}$  wind speed ( $\text{ms}^{-1}$ ) and the Schmidt Number of DMS  $SC_{\text{DMS}}$ , a measure of viscosity/diffusion and a function of sea surface temperature, is determined following the method of Saltzman et al. (1993). The denominator in this function is the Schmidt Number of  $\text{CO}_2$  in fresh water at  $20^\circ\text{C}$ ,  $SC_{\text{CO}_2} = 600$ , which is used to normalise the numerator ( $SC_{\text{DMS}}$ ). We assume that the concentration of  $\text{DMS}_{\text{a}}$  is negligible, as it is orders of magnitude smaller than that of ~~sea-water~~ seawater.

## 2.2 Model evaluation

In order to provide a climatological evaluation of ACCESS-UKCA, and to put the sensitivity testing of DMS into a real-world context, a comparison to observational data sets is presented. Global means at the surface are calculated over the 2000-2009 period, except for the cloud climatologies which were only available from 2006-2009.

The following global data sets were compared to the model output: low, medium and high cloud fractions from the GCM-Oriented Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation Cloud Product (CALIPSO-GOCCP) (Chepfer et al., 2010), radiation fluxes from the Clouds and the Earth's Radiant Energy System (CERES) -Energy Balanced and Filled (EBAF) -TOA Edition 4.0 (Loeb et al., 2009) and CERES-EBAF-Surface Edition 4.0 (Kato et al., 2013) and precipitation from the Tropical Rainfall Measuring Mission (TRMM) (Huffman et al., 2007). Cloud fraction is defined ~~in-line-with~~ according to the CALIPSO-GOCCP: high between ~~50-44hPa, medium~~ 50-440 hPa, medium between 440-680 hPa and low ~~680-1000hPa~~ between 680-1000 hPa. Direct comparison of cloud fractions between model output and satellites cannot take into account satellite measurement biases, which can be resolved using a cloud satellite simulator such as the Cloud Feedback Model Intercomparison Program (CFMIP) Observation Simulator Package (COSP). COSP was not available in the version of ACCESS-UKCA used here, limiting the comparison. Nevertheless, a useful comparison is still possible.

### 2.3 DMS sensitivity testing

To explore the sensitivity of the global climate to large perturbations in  $\text{DMS}_w$  concentrations, two experimental simulations were performed and compared to the control run (Ctl). As described above, the Ctl simulation used the Lana et al. (2011)  $\text{DMS}_w$  climatology, which is shown in Fig. 2a.

- 5 In Experiment 1 (Exp.1),  $\text{DMS}_w$  was set to zero, leaving a flux of  $0.72 \text{ Tg year}^{-1}$  of sulfur derived from terrestrial sources (for example Jardine et al., 2015). From this we can attribute what role ocean-derived DMS plays in shaping our current climate and enhance our understanding of how the physical processes underpinning the DMS-climate system operate. This may further aid our understanding of how natural aerosols interact with the global radiation budget.

- 10 In Experiment 2 (Exp.2), the  $\text{DMS}_w$  field was set to each latitude's (at the model resolution of  $1.25^\circ$ ) monthly zonal maximum value, following a similar method to Grandey and Wang (2015), and shown in Fig. 2b. This simulation allows further exploration of the physical processes by which DMS can influence the climate, when the perturbations are exaggerated.

- ~~Three regions of interest are defined. The~~ Three regions of interest are defined for their relevance to the broader Australian community (for which ACCESS is purposed) or are of particular interest in the DMS-climate system. They are the Australian region:  $45^\circ\text{S}$ - $10^\circ\text{S}$ ,  $110^\circ\text{E}$ - $160^\circ\text{E}$ , the Southern Ocean (SO): ocean grid points south of  $40^\circ\text{S}$  and the South Eastern ~~Pacific~~ Pacific (SEP) that represents an area of significant stratiform cloud decks:  $240^\circ\text{E}$ - $270^\circ\text{E}$ ,  $25^\circ\text{S}$ - $0^\circ$ .

### 2.4 Global energy budget

Due to the nudging of the model to ERA-Interim and the prescribed SSTs, a direct estimate of how global temperatures might respond to DMS perturbations is not possible. For this reason, a simple energy balance model has been used to estimate the effects of the DMS perturbations on global mean temperatures: ~~the~~, a useful metric for comparison to some previous studies.

- 20 We have used the climate component of the Finite Amplitude Impulse Response (FAIR) model. This model is based on ~~that~~ the one first proposed by Boucher and Reddy (2008) and subsequently used in the most recent IPCC Assessment Report 5 for equivalent emission metric calculations (Myhre et al., 2013b). FAIR's climate component is a simple impulse response model which emulates the behaviour of more complex Earth System Models. ~~It is ideally suited to estimating global mean temperature changes resulting from TOA radiative imbalances-~~, given a certain radiative forcing (in this case due to DMS). FAIR has been  
25 designed to determine temperature responses to radiative forcing of similar magnitudes to the DMS radiative effect (Millar et al., 2017). FAIR's temperature response is calculated as the sum of two components, approximately representing the response of the upper mixed layer and deep ocean to a change in radiative forcing (Millar et al., 2017). Due to its simplicity, FAIR cannot capture the non-linearities and feedbacks in the climate system, and hence the temperature response calculated must be taken as an estimate only. Furthermore, in this work we consider only a single mid-range estimate of climate sensitivity.

- 30 For each experimental run, the radiative effect ( $R_{DMS}$ ) due to increasing or decreasing  $\text{DMS}_w$  is defined as the difference between the TOA energy balance ( $Q^*$ ) of an experimental run from the Ctl, which can be taken directly from ACCESS-UKCA. By providing this radiative effect to FAIR's climate component, we can estimate the difference in temperature expected across the ten year period under zero  $\text{DMS}_w$  or enhanced  $\text{DMS}_w$  conditions, ~~without the need for an ensemble of free-running climate~~



~~simulations.~~ In ACCESS-UKCA, an ensemble experiment would be required to provide equivalent temperature difference estimates, which would be computationally expensive.

To provide a measure of model uncertainty of the change in  $Q^*$ , we have used a moving block bootstrap (Wilks, 2011). By selecting, with replacement, blocks of size 2 (as determined by the time series auto-correlation) to create 1000 alternate time series, we are able to provide the 10th and 90th percentile confidence intervals of mean change in  $Q^*$ . This is subsequently translated into an uncertainty range for the change in temperature and flux sensitivity estimates.

### 3 Model evaluation

This section compares selected ACCESS-UKCA fields to satellite-derived observations. In order to give context to this evaluation, the ACCESS-UKCA output is also compared to that of the CMIP5 (~~Coupled-Model-Intercomparison-Project-Phase-5~~) ~~global-climate-models~~ (GCMs).

#### 3.1 Cloud fraction

The cloud fraction comparison is performed for the years 2006-2009, aligning with the availability of CALIPSO-GOCCP data. ACCESS-UKCA simulates too little low cloud fraction (Fig. 3a-c) over the the majority of the globe (mean bias of -0.16), which is consistent with findings for the CMIP5 GCMs (Cesana and Chepfer, 2012; Nam et al., 2012; Klein et al., 2013). Areas of large stratiform cloud decks in eastern ocean basins are significantly underestimated, by a fraction ~~of~~  $\rightarrow$  larger than 0.5, consistent with other CMIP5 and CFMIP Phase 1 and 2 findings (Bony and Dufresne, 2005; Cesana and Chepfer, 2012; Klein et al., 2013). These low level marine clouds have an important impact on the global radiation budget (Leon et al., 2009) and have been identified as the primary source of uncertainty in tropical cloud-climate feedbacks (e.g., the effects of the cloud albedo) in GCMs (Bony and Dufresne, 2005). These biases have been attributed to poor vertical distribution of clouds in the models, difficulty capturing overlapping cloud layers, the misrepresentation of cloud structures, deficiencies with the statistical parameterisation of clouds and likely problems in the cloud microphysics (Nam et al., 2012). Low clouds over the polar regions and some areas of northern Asia and America are slightly overestimated. The ACCESS-UKCA low-cloud biases over the Arctic are within the range of biases found for the CMIP5 GCMs studied in Cesana and Chepfer (2012). It should be noted that satellite observations are subject to biases in detecting low clouds, particularly over the Southern Ocean.

ACCESS-UKCA reproduces medium cloud fraction (Fig. 3d-f) reasonably well, within  $\pm 0.1$  in most regions (global mean bias of -0.01). The largest discrepancies are overestimated medium cloud fraction over the Southern Ocean and Antarctica, where the simulated medium cloud fraction is at its highest globally. The Antarctic bias is of opposite sign to the CMIP5 models compared in Cesana and Chepfer (2012). Bodas-Salcedo et al. (2014) note that issues within GCMs around distinguishing between clouds with tops at actual mid level and low level clouds contribute to such biases. The biases in high cloud fractions (Fig. 3g-i) show similar spatial patterns to that of the low cloud fraction, where an underestimate occurs over most of the tropics and mid-latitude. The global mean bias is 0.05. The largest negative biases, of up to 0.3 occurs over the Maritime Continent.



Moderate overestimation is noted over the polar regions. These biases are within the range of those found for the CMIP5 models studied in Cesana and Chepfer (2012).

Interestingly, Nam et al. (2012) noted that due to underestimated low clouds in the tropics, the CMIP5 models over compensated by producing optically thick and too bright low clouds and more high clouds, impacting the radiation budget. Here, an underestimation of low clouds is also found, although there is no evidence of an over-compensation of high clouds. Predominantly a small underestimation of high cloud fraction is found in this simulation at tropical to mid latitudes (Fig. 3i).

### 3.2 Radiation

The remaining analyses consider means over the period of 2000-2009. The comparison of the observed and simulated TOA outgoing LW radiation is shown in Fig. 4a-c. The observed global mean of  $239.7 \text{ W m}^{-2}$  is closely matched by the simulated  $241.0 \text{ W m}^{-2}$ . Compared to the CMIP5 ensemble, which tends to underestimate TOA outgoing LW radiation,  $238.6 \text{ W m}^{-2}$  from Stephens et al. (2012) and  $238.9 \text{ W m}^{-2}$  from Wang and Su (2013), TOA outgoing LW radiation in ACCESS-UKCA is slightly overestimated. The regions with the largest biases (both positive and negative) occur in regions of deep convection (Fig. 4c), and align well spatially with the biases in high cloud fractions shown in Fig. 3c. Underestimation by  $-3 \text{ W m}^{-2}$  of TOA outgoing LW radiation occurs over the polar oceans, which may partly be explained by an overestimation of cloud fraction at all levels, and especially the mid-level clouds (Fig. 3f) in this region.

Spatial biases in the TOA outgoing SW radiation (Fig. 4d-f) are of greater magnitude than that of the LW radiation. In most regions the sign of the outgoing SW radiation bias is opposite to that of the LW radiation. The same processes as described above that block LW radiation from escaping the atmosphere prevent SW radiation reaching the surface, ~~instead~~ hence reflecting more sunlight ~~thus~~ and enhancing the albedo. Globally, ACCESS-UKCA performs reasonably well, simulating the global mean TOA outgoing SW radiation of  $101.8 \text{ W m}^{-2}$  compared to the observed  $99.6 \text{ W m}^{-2}$ , consistent with the multi-model mean of GCM ensembles from previous studies (Stephens et al., 2012; Wang and Su, 2013). In Fig. 4f, an abrupt change in sign of TOA outgoing SW radiation at  $60^\circ\text{S}$  is found, which is also present in the CFMIP comparisons (Bodas-Salcedo et al., 2014). In the Southern Ocean, wrongly assigned mid-level cloud types have been found to be a leading cause of the model underestimation of TOA outgoing SW radiation (Bodas-Salcedo et al., 2014). In addition, poor representation of the physical processes surrounding super-cooled liquid water in the Southern ocean has been found to account for 27-38% of the total reflected solar radiation (Bodas-Salcedo et al., 2016). Over the Antarctic ice sheets, both TOA outgoing and surface incoming SW radiation are overestimated, due to an underestimation of low clouds ~~allowing~~, which allows the high albedo to reflect too much incoming SW radiation back out to space.

Globally, ACCESS-UKCA overestimates incoming surface SW radiation (Fig. 4g-i), with  $202.4 \text{ W m}^{-2}$  compared to observations of  $198.3 \text{ W m}^{-2}$ . This overestimation is slightly greater ~~, though within the uncertainty of~~ than that found for CMIP5 GCMs of  $2 \pm 6 \text{ W m}^{-2}$  (Stephens et al., 2012), though within their uncertainty. Nevertheless, large regional biases of over  $\pm 30 \text{ W m}^{-2}$  exist. The most notable features, apart from those discussed above, are too much incoming SW radiation over the continents and the tropical regions, which can be attributed in part to the underestimated cloud cover. The northern Pacific

and Atlantic Oceans, the Arctic Ocean and parts of the Southern Ocean all receive too little incoming SW radiation, consistent with overestimated cloud cover.

### 3.3 Precipitation

Precipitation in ACCESS-UKCA has large positive biases (~~Fig. 5d-f). Over regions to the north and south of the equator, regions that receive the most annual rainfall, ACCESS-UKCA performs with the least skill, overestimating precipitation by over 2000.~~ in regions that receive the most annual rainfall and align with the Intertropical and South Pacific Convergence Zones (ITCZ/SPCZ). These regions overestimate precipitation by over 2000 mm year<sup>-1</sup>. Poor performance of GCMs in this region are not unusual however (Stephens et al., 2010), with the current CMIP5 GCM ensemble overestimating precipitation in a similar region by more than 1000 mm year<sup>-1</sup> (Flato et al., 2013). Stephens et al. (2010) found that models in these regions produce light rain too frequently, indicating that convective processes are poorly simulated. Two of Australia's CMIP5 GCMs, ACCESS 1.0 and 1.3, both overestimate precipitation in this region by similar amounts to that of the ACCESS-UKCA model (Bi et al., 2013). If biases of precipitation are considered as a percentage (not shown), ~~the largest differences (positive) occur in the eastern basins of the South Pacific and South Atlantic Oceans.~~

the largest differences occur in the eastern basins of the South Pacific (493% over the SEP region) and South Atlantic Oceans (275% from 0 – 25°S, 330 – 10°E).

## 4 DMS perturbations

This section aims to quantify the role of DMS in the large-scale climate system. Two experimental simulations have been performed, described in Section 2.3 and Table 1, which involve the removal of all DMS<sub>w</sub> (Exp.1) and setting the DMS<sub>w</sub> to the zonal maximum (Exp.2).

### 4.1 Exp.1: Zero DMS<sub>w</sub>

#### 4.1.1 Chemistry response

The 2000-2009 annual mean ocean flux<sub>DMS</sub> from ACCESS-UKCA is 17.41 Tg year<sup>-1</sup> of sulfur, resulting in an atmospheric DMS (DMS<sub>a</sub>) annual mean surface concentration of 81.9 ppt. Taking all marine DMS out of the model (but retaining the terrestrial source of 0.72 Tg year<sup>-1</sup> of sulfur) results in a 94% reduction in DMS<sub>a</sub> at the surface; throughout the troposphere, it results in a 98% reduction of DMS<sub>a</sub>.

The impact of this reduced flux<sub>DMS</sub> on atmospheric sulfur can be seen in Fig. 6a-b, d-e. Globally, there is a net decrease of 15% of SO<sub>2</sub> at the surface. The largest absolute differences are in the tropics and mid-latitudes over the oceans. Large relative decreases in SO<sub>2</sub> occur in the SO and SEP, 84% and 94% respectively. Fig. 7a shows the vertical profile of SO<sub>2</sub> for the Australian region (ref), the SO (blue) and the SEP (green). The large peak in concentration at approximately 500 m occurring in the Australian profile is attributable to industrial and energy-related emissions of SO<sub>2</sub>, ~~representing~~ which is due to lofting

by chimneys and smokestacks. The  $\text{SO}_2$  in Exp.1 is consistently lower than that of the Ctl throughout the troposphere, though for the regional means, the difference begins to decrease closer to the tropopause.

Surface  $\text{H}_2\text{SO}_4$  (Fig. 6d-e) shows significant loss in predominantly clean marine areas; the SO has a 79% decrease and the SEP an 84% decrease, compared to a 49% global mean decrease. Interestingly, heavily polluted regions, especially busy shipping lanes, undergo an increase in  $\text{H}_2\text{SO}_4$ .  $\text{H}_2\text{SO}_4$  is a precursor gas, which can participate ~~in new particle formation forming~~ in the formation of secondary sulfate aerosol, or it can condense onto pre-existing particles. The increased  $\text{H}_2\text{SO}_4$  concentration in heavily polluted regions results from a decreased condensational sink (not shown). Similar non-linearities have been described in Thomas et al. (2011).

The vertical profiles of  $\text{H}_2\text{SO}_4$  in Fig. 7b, show that the largest differences between the Exp.1 and the Ctl occur in the free troposphere (between 1-10 km) for all regions. In all three regions (each considered a clean atmospheric environment), net decreases of  $\text{H}_2\text{SO}_4$  occur.

#### 4.1.2 Aerosol response

The majority of gaseous  $\text{H}_2\text{SO}_4$  is taken up by aerosol formation (99.99%) as opposed to being removed by dry deposition (0.01%) (Mann et al., 2010). The peak in nucleation mode number density in the free troposphere in ~~Fig. 7e~~ Fig. 7c coincides with the peak concentration of  $\text{H}_2\text{SO}_4$ . Surface global nucleation mode number concentration decreases by 9% between Exp.1 and the Ctl (see Fig. ~~6g-h~~ 7c). While in absolute terms, clean terrestrial regions have the largest decreases, the Australian region only has a relative decrease of 18% in nucleation mode particles. Over the oceans, although few nucleation mode particles exist, there are large relative differences of both signs.

In absolute terms, the differences in the aerosol number concentration are greatest in the smaller aerosol modes, particularly the nucleation mode described above. Fig. 7d-f show the number concentrations for the Aitken mode, accumulation mode, and coarse mode (global maps not shown). The Aitken mode (Fig. 7d) shows some differences between the two simulations, with profiles reflecting reduced new particle formation in the free troposphere and reduced condensation-growth of  $\text{H}_2\text{SO}_4$  onto pre-existing particles in the boundary layer. The largest differences are seen over the Australian region. Similar boundary layer differences are also present in the accumulation mode, with the differences between the Exp.1 and Ctl consistent below 1 km (Fig. 7e). Little difference is seen in the coarse mode throughout the troposphere (Fig. 7f), which in marine regions is dominated by sea salt.

As aerosols grow towards the larger end of the Aitken mode, they become relevant to cloud processes. Fig. 8a shows the Ctl's  $\text{N}_3$  (condensation) number concentration ( $\text{N}_3$  signifies particles with a dry diameter greater than 3 nm). The difference in surface  $\text{N}_3$  number concentration between the Ctl and Exp.1 shows the largest relative decreases occur in clean, coastal regions, predominantly in the Southern Hemisphere, as well as some regions of the SO. In heavily polluted terrestrial regions a small increase in the  $\text{N}_3$  number concentration occurs. A decrease of 8% is found globally. For the Australian region (representative of a clean, terrestrial region), a decrease of 17% is found. ~~The largest absolute differences are in clean terrestrial regions.~~

Over the SO a relative decrease of 39% occurs at the surface. The SO and the SEP have far fewer aerosols in all modes except the coarse mode (see Fig. 7c-f), where sea salt dominates. This decrease in number concentration in small aerosol

modes represents a large portion of the aerosol loading in the region. The increase in nucleation mode particles is reflected in the  $N_3$  for the SEP region, via a more moderate decrease of 20%.

Fig. 8d-e show the number concentration of ~~cloud condensation nuclei~~ CCN with dry diameters greater than 70 nm ( $CCN_{70}$ ) for the Ctl and the differences resulting from Exp.1. The largest absolute differences are in the tropics, which, similarly to the  $N_3$ , have the highest concentration. Relatively, there is a global decrease of 5%, ~~whilst over the Australian region a 7% decrease, the SO an 8% decrease and the SEP a 20% decrease~~ while decreases of 7% were found over the Australian region, decreases of 8% over the SO and decreases of 20% over the SEP. Differences in cloud droplet number (CDN) are shown in Fig. 8g-h. The relative differences in CDN (~~Exp.1-Ctl~~) show a similar spatial pattern to that of the CCN. Global mean CDN decreases by 5%, ~~whilst a~~. A decrease of 5% is also found for the Australian region, ~~8% for the SO and 18% for the SEP~~, whereas the SO shows an 8% decrease, and the SEP shows an 18% decrease. In both the  $CCN_{70}$  and CDN, the marine Southern Hemisphere ~~mid-latitudes~~ mid-latitudes have the largest decreases of 14% (averaged between 5-35°S) despite the SO having some of the larger decreases in  $SO_2$  and  $H_2SO_4$ .

The larger differences in concentration of both CCN and CDN in the oceanic Southern Hemisphere ~~tropics-mid~~ tropics and mid latitudes, compared to the SO, warrant further investigation of how sulfate aerosols are interacting with their background environments, for example cloud processes and pre-existing aerosols. The SO has large concentrations of sea salt particles, which like more polluted regions of the Northern Hemisphere, may provide a buffering effect to reduced DMS-derived aerosols. Additionally, in areas of persistent low cloud formation, in-cloud aqueous sulfate oxidation is the dominant reaction (over gaseous nucleation), which allows almost instantaneous condensational growth of existing aerosols, and is temperature dependent. We speculate that poor representation of low clouds in the SO may be having further impacts on atmospheric composition modelling than currently realised. A cloud resolving modelling study may be better suited to gain understanding of the complex system described here.

Following the method of Woodhouse et al. (2010, 2013), global and hemispheric sensitivities of CCN to  $flux_{DMS}$  have been calculated (Table 2). The results presented here ~~a suggest~~ suggest a lower CCN sensitivity to  $flux_{DMS}$  compared to the Woodhouse et al. (2013) study where absolute sensitivities of 94 and 63  $cm^{-3}/mg\ m^{-2}\ day^{-1}$  of sulfur were found globally for June and December respectively. Similar CCN sensitivities are reported in the Woodhouse et al. (2010) study (63  $cm^{-3}/mg\ m^{-2}\ day^{-1}$  global average). The lower sensitivities in our study are likely the result of the large (near 100%) changes in  $flux_{DMS}$  (the denominator). Relative CCN sensitivities calculated here compare well with the Woodhouse et al. (2010) and ~~Woodhouse et al. (2015)~~ Woodhouse et al. (2013) studies. For example Woodhouse et al. (2010) finds mean hemispheric relative CCN sensitivities of 0.02 for the Northern Hemisphere and 0.07 for the Southern Hemisphere. These results highlight the greater relative importance of DMS in the Southern Hemisphere.

#### 4.1.3 Meteorological response

### 4.1.3 Cloud, radiation and precipitation response

Meteorological responses to the DMS perturbations must be considered carefully. As detailed in the methods section, the ACCESS-UKCA simulations are nudged to ERA-Interim potential temperature and horizontal winds, preserving synoptic scale meteorology and limiting any feedbacks. ~~Whilst~~ While performing a non-nudged simulation would allow the meteorology to respond to changes in the chemistry and aerosol more freely, it would make comparison of the aerosol and meteorological responses more difficult. Within ACCESS-UKCA, GLOMAP-mode is directly coupled to the large-scale cloud and precipitation schemes via the CDN (Abraham et al., 2012), as well as the radiation scheme via aerosols and some gases (see Section 2.1.1). Convective rainfall and cloud formation are not directly coupled to the aerosol scheme, but can be indirectly influenced via changes in radiation (which can act on temperature, moisture, etc).

Differences in low cloud fraction occur predominantly in areas with large stratiform cloud decks (Fig. 9a), ~~in the eastern basins of the Southern Hemisphere's oceans and slightly removed from the coastline~~. The largest differences occur in eastern basins of the Southern Hemisphere's oceans. The SEP region shows an annual mean decrease in low cloud fraction of 9%. In the Northern Hemisphere (including the north eastern Pacific where significant stratiform cloud decks are found) and the SO (where persistent low cloud formation occurs) only small differences are evident, which may in part be due to the modest differences in CCN and CDN concentrations discussed in Section 4.1.2. Stratiform cloud deck fractions are consistently underestimated by ACCESS-UKCA and other GCMs (see Section 3) in comparison to other areas of significant low cloud formation such as the SO. The mechanism behind the different responses (between the SO and cloud deck regions), and whether the long standing model biases ~~exacerbate this~~, especially those around the formation of supercooled liquid water, have contributed to the differing responses requires further investigation.

The decrease in low cloud fraction and aerosol number concentration discussed above lead to an increase in surface incoming SW radiation (Fig. 9c). This increase in surface SW radiation is highest in the regions of stratiform cloud deck formation. In the SEP region there is a mean increase of  $7 \text{ W m}^{-2}$ .

Decreases of total liquid water ( $Q_{cl}$ ) at 1700 m height shown in ~~Fig. 10a-b~~ Fig. 10a-b are found in the stratiform cloud deck regions. Little difference in  $Q_{cl}$  occurs at the surface. The decrease in  $Q_{cl}$  is coincident with increases in large-scale precipitation in the stratiform cloud decks, regions with very little precipitation (Fig. 10d-e). In the SEP region large-scale rainfall increases by  $17 \text{ mm year}^{-1}$  (15%) over the Ctl mean of  $111 \text{ mm year}^{-1}$ .

In the ~~southern hemisphere~~ Southern Hemisphere stratiform cloud decks, and in particular the SEP region, the model demonstrates a distinct cloud lifetime effect in response to removing DMS in Exp.1. Decreased CDN concentration and the associated increase in cloud droplet size and increased liquid water lead to increased autoconversion and large scale rainfall. The overall impact is to reduce low cloud fraction.

Figure 10g-h shows the differences in convective rainfall. ~~Whilst~~ While the convective rainfall scheme is not coupled directly to GLOMAP-mode, there are differences between the simulations. Convective rainfall decreases in Exp.1 compared to the Ctl along the Intertropical Convergence Zone (ITCZ) (a mean difference of  $11 \text{ mm year}^{-1}$  between  $20^\circ\text{S}$ - $20^\circ\text{N}$ ). This difference

represents a small fraction (less than 1%) of the total convective rainfall. Relatively, (not shown) the largest differences (a 5% decrease in the SEP) are found once again in eastern basins of Southern Hemisphere stratiform cloud decks.

Seifert and Beheng (2006) note that even when convection schemes are coupled to an aerosol scheme, the effects of CCN on convection, and the resultant precipitation and associated maximum updrafts, differs significantly depending on the cell type and size, making these effects difficult to quantify. Large differences in convective rainfall would not be expected in these results, due to the meteorological nudging used in the experiments.

## 4.2 Exp.2: Zonally increased DMS<sub>w</sub>

This section considers the response to zonally enhanced DMS<sub>w</sub>, resulting in a flux<sub>DMS</sub> of 37.05 Tg year<sup>-1</sup> of sulfur (relative to ~~17.4~~ 17.41 Tg year<sup>-1</sup> of sulfur in the Ctl simulation). For comparison the Grandey and Wang (2015) study used a zonally enhanced flux<sub>DMS</sub> of 46.1 Tg year<sup>-1</sup> of sulfur (up from 18.2 Tg year<sup>-1</sup> of sulfur), under global warming scenarios. Many of the differences resulting from zonally enhancing DMS<sub>w</sub> show similar spatial patterns, with similar magnitude ~~-, but reversed sign,~~ but reversed sign compared to Exp.1.

Globally, the differences in SO<sub>2</sub> (Fig. 6c) are of comparable magnitude to Exp.1. Increased SO<sub>2</sub> concentrations occur over the Australian region, the SO and the SEP: increases of 42%, 172%, 89% respectively. There is a net decrease in H<sub>2</sub>SO<sub>4</sub> of 14% in Australia, and a larger decrease over the tropical oceans. Over the SO there is an increase of 9%, ~~whilst~~ while in the SEP a decrease of 37% occurs. Similar non-linearities are discussed in terms of doubled DMS in the Thomas et al. (2011) study. These differences in SO<sub>2</sub> and H<sub>2</sub>SO<sub>4</sub> are also clear in the vertical profiles shown in Fig. 7a-b.

Differences in the aerosol modes (see Fig. 7c-f) are of similar magnitude but opposite sign to those noted in Section 4.1.2. Global mean N<sub>3</sub>, CCN<sub>70</sub> and CDN increases by 6%, 4% and 5% respectively (Fig. 8c,f,i). Larger differences are seen over the SO of 27%, 15% and 13% and the SEP of 14%, 19% and 17% for N<sub>3</sub>, CCN<sub>70</sub> and CDN respectively.

Globally, there is little difference in low cloud fraction or  $Q_{cl}$ , though increases are noted in regions of large stratiform cloud decks (Fig. 9d), ~~showing~~ which show similar spatial patterns to that of Exp.1. ~~Surface-incoming~~ Incoming surface SW radiation has a global mean decrease of -1.75 W m<sup>-2</sup>. This decrease is comparable to the Grandey and Wang (2015) finding of -2.2 W m<sup>-2</sup> (noting the larger DMS perturbation by Grandey and Wang (2015)). Lastly, decreases in large-scale precipitation are found, again in regions of stratiform cloud decks (Fig. 10f), ~~whilst~~ while general increases in convective precipitation over the tropical oceans occur (Fig. 10i). The Grandey and Wang (2015) study, ~~analysed-under~~ which analysed a warming climate, also found large relative decreases in precipitation rate predominantly in eastern ocean basins. We find, under the current climate, the largest relative increase in total precipitation (not shown) in the ~~south-east~~ southeast basins of the Pacific and Atlantic ocean, however these results presented here are much noisier than the Grandey and Wang (2015) results. ~~Whilst~~ Grandey and Wang (2015) find that artificial enhancement of DMS may offset global warming, which is supported by this study as implied by the decreases in incoming SW radiation at the surface, however the precipitation responses warrant further study.

### 4.3 Temperature response

The global 2000-2009 mean of the TOA radiation budget ( $Q^*$ ), and its main components are provided in Table 3, along with the relevant confidence intervals derived from the bootstrapping technique. Due to the nudging used in the simulations, we do not expect the TOA  $Q^*$  to be balanced (i.e.  $Q^* = 0$ ). The differences in  $Q^*$  seen in Exp.1 and Exp.2,  $1.69 \text{ W m}^{-2}$  (1.13 to 2.43, 10th and 90th confidence intervals) and  $-1.48 \text{ W m}^{-2}$  (-2.33 to -0.98, 10th and 90th confidence intervals) respectively, show a substantial radiative effect of DMS on the energy budget. The  $Q^*$  response found for Exp.1 is consistent with the Mahajan et al. (2015) findings of  $1.79 \text{ W m}^{-2}$ . Using the FAIR model's climate component, the 2000-2009 mean temperature response is calculated to be  $0.45 \text{ K}$  (0.30 to 0.64, 10th and 90th percentile range) for Exp.1 and  $-0.38 \text{ K}$  (-0.26 to -0.61, 10th and 90th percentile range) for Exp.2.

Although other studies generally consider DMS changes under global warming, and we can make comparison comparisons via the sensitivity of the estimated global temperature response to changes in the  $\text{flux}_{\text{DMS}}$  (see Table 4). In this study, we find a response of  $0.027 \text{ K per Tg year}^{-1}$  of sulfur in Exp.1, and  $0.019 \text{ K per Tg year}^{-1}$  of sulfur in Exp.2. These results are of similar magnitude to the Grandey and Wang (2015) study, and in the range of the lowest impact scenario of Six et al. (2013). The other scenarios in the Six et al. (2013) study suggest much higher temperature sensitivities to changes in, as does the Schwinger et al. (2017) study. These results are of similar magnitude to the Grandey and Wang (2015) study ( $0.029 \text{ K per Tg year}^{-1}$  of sulfur) and in the range of the lowest impact scenario of Six et al. (2013) ( $0.03 - 0.060 \text{ K per Tg year}^{-1}$  of sulfur). The other scenarios in the Six et al. (2013) study ( $0.046 - 0.11 \text{ K per Tg year}^{-1}$  of sulfur) suggest much higher temperature sensitivities to changes in  $\text{flux}_{\text{DMS}}$ , as does the Schwinger et al. (2017) study ( $0.041 \text{ K per Tg year}^{-1}$  of sulfur).

## 5 Discussion and Conclusions

We have used a global climate model, ACCESS-UKCA, which includes a detailed microphysical aerosol module. The ACCESS-UKCA chemistry-climate model, which includes a detailed microphysical aerosol module, has been evaluated against satellite observations of cloud fraction, radiation and precipitation and subsequently used to conduct sensitivity experiments to determine the role of DMS on in several aspects of the climate system.

Evaluation using satellite-derived observations shows ACCESS-UKCA is comparable to other GCMs in the assessed climatological fields. Low, medium and high cloud fractions, compared to the CALIPSO-GOCCP satellite climatology show similar biases to that of the CMIP5 ensemble (Bony and Dufresne, 2005; Cesana and Chepfer, 2012; Nam et al., 2012; Klein et al., 2013), as do the radiation fields (Stephens et al., 2012; Wang and Su, 2013). Of particular interest to this study is the model underestimation of large stratiform cloud decks located in the eastern mid-latitude basins of the Earth's oceans. These regions of extensive low cloud produce little rainfall (that is overestimated by the model) and are often regions of high primary productivity. These biases have not been attributed to a single cause (rather a multitude of theories, as discussed in Section 3.1), indicating a gap in understanding of atmospheric processes in these regions (Nam et al., 2012).



The use of nudging does not allow aerosol and cloud responses to perturbed DMS to affect synoptic scale meteorology, hence the results here represent instantaneous responses in the climate system. Nudging was deemed desirable for this study to limit computational expense, allowing single runs of 10 years, rather than ensemble runs of greater duration. Other considerations

Important considerations when using climate models include the inherent uncertainties associated with all climate simulations, including e.g. emissions uncertainties (both natural and anthropogenic), and parameterizations and physical representation parameterisations and physical representations of atmospheric processes. Nevertheless, where clear shortcomings have been found in comparison to the satellite-derived observations, the ACCESS-UKCA model has been found to perform with comparable skill to current CMIP5 GCMs. Additionally, it is important to note biases in the satellite products themselves, for example in cloud fraction retrievals as noted in Mace and Zhang (2014) or Protat et al. (2014).

Removing marine from the climate system leads to significant responses in chemistry and aerosol concentrations. The differences are largest in and , with differences becoming smaller through the latter stages of the DMS-aerosol-cloud system (Figure 1). Areas of Of particular interest, our evaluation of ACCESS-UKCA shows an underestimation of large stratiform cloud decks are particularly sensitive to DMS emissions. On average globally, marine DMS contributes up to 16% of, and 48% of. The Australian region, considered a clean, terrestrial environment, undergoes greater decreases in these fields than the global mean, whilst the SO and the SEP clean marine environments experience even larger decreases. Marine DMS removal results in an 8% decrease of located in the number density and eastern mid-latitude basins of the Earth's oceans. These regions of extensive low cloud produce little rainfall (that is overestimated by the model) and are often regions of high primary productivity. These biases have not been attributed to a 5% decrease of CCN and CDN globally. In less pristine environments, close to coastlines and in the Northern Hemisphere, a weak response is found in , CCN and CDN. Similarly, a weak response is also found in the SO where sea salt makes a large contribution to aerosol. single cause (multiple theories have been proposed, as discussed in Section 3.1), indicating a gap in understanding of atmospheric processes in these regions (Nam et al., 2012).

Changes Globally, removing or enhancing DMS<sub>w</sub> from the climate system leads to significant responses in chemistry and aerosol concentrations. While changes in meteorological parameters (low cloud fraction, large-scale precipitation, moisture, radiation) are largest in regions where large the Southern Hemisphere stratiform cloud decks occur, generally in the Southern Hemisphere. In the SEP region we find a decrease of 9% in low cloud formation and a 15% increase in large scale rainfall when DMS is removed. These stratiform cloud decks are regions where considerable model bias exists in low cloud fraction and precipitation. Global, global mean differences were small, implying a higher sensitivity of these stratiform regions to changes in DMS. We find that DMS in these stratiform regions plays an important role in cloud processes and precipitation suppression (as discussed in Thomas et al. (2011) or with regards to anthropogenic pollution in Ackerman et al. (2004)).

Furthermore, we have demonstrated that marine DMS is responsible for increasing low cloud fraction in stratiform cloud deck regions, a demonstration of the second aerosol indirect (or lifetime) effect (Albrecht, 1989). These results indicate that a greater understanding of natural aerosols and their interaction with cloud processes (both via observations and modelling studies) in these regions may improve model representation, as it is these regions that show considerable model bias in comparison to observations.



In other regions of significant low cloud formation (SO, Northern Hemisphere cloud decks), aerosol sources such as sea salt and anthropogenic aerosols may ~~buffer~~ buffer the regions from changes in DMS-derived aerosols. Additionally, in the SO, representation of cloud processes in global climate models is poor, especially with respect to supercooled liquid water (Bodas-Salcedo et al., 2016). It is likely that these biases are misrepresenting the DMS-climate interactions in these regions.

5 ~~Significant differences in the global radiation budget result from removing marine and the subsequent decrease in sulfate aerosol and low cloud fractions. As a result of the decreases in low cloud fraction in the SEP region, there is an increase of 7 in incoming surface SW radiation. We find removal of leads to a global increase in the TOA radiation of 1.69, resulting in an increase in global temperature of approximately 0.45 K.~~ By nudging these simulations, the model response to the DMS perturbations is limited to fast (aerosol and cloud) changes. We suggest that free running ensemble experiments are performed  
10 to gain insight into the aerosol-cloud-climate processes occurring in regions of significant DMS influence. Such experiments should focus on improving microphysical aspects of aerosol-cloud interaction in these regions (and how it differs among regions) or improving the representation of aerosols, in particular natural aerosols.

Previous studies examining the role of DMS in the climate system have not identified stratiform cloud decks as regions of particular importance, ~~rather highlighting the SO Thomas et al. (2010); Mahajan et al. (2015).~~ Instead, these studies focused on  
15 cloud feedbacks in [the SO](#) (Thomas et al., 2010; Mahajan et al., 2015). Mahajan et al. (2015) estimated the [global](#) TOA radiative effect of DMS to be  $1.79 \text{ Wm}^{-2}$ , which is consistent with our results ( $1.69 \text{ Wm}^{-2}$ ), but slightly lower than the estimation of  $2.03 \text{ Wm}^{-2}$  estimated by Thomas et al. (2010) (who used the previous ~~Kettle et al. (1999)~~ [Kettle and Andreae \(2000\)](#) DMS<sub>w</sub> climatology).

In ~~addition to impacts on radiation budget and temperature, as previously reported in the literature, we are able to quantify the role of DMS in suppressing large-scale precipitation in stratiform cloud deck regions. For example in the SEP region we find an increase in large-scale rainfall of 15% when is removed. Furthermore we have demonstrated that marine is directly responsible for increasing low cloud fraction in stratiform cloud deck regions, a demonstration of the second aerosol indirect (or lifetime) effect (Albrecht (1989)).~~

The zonally enhanced simulation responds in almost the opposite manner, with the same regions of the globe showing  
25 differences of similar magnitude but opposite sign. The Grandey and Wang (2015) study, with a of 44.1 of sulfur estimated a TOA radiative effect of up to -1.28 in low-medium cloud regions for the year 2000, and -1.49 by 2080. Our study, with a of 37.0 of sulfur, finds a global mean radiative effect of -1.48 (2000-2009 mean). Areas of stratiform cloud decks show differences of up to -5 of incoming SW radiation. In this study, the estimated global mean temperature change with increased is -0.38 C (2000-2009 mean). The changes in the aerosol and chemistry noted here imply that significant enhancements of DMS may  
30 have greater implications for some regions than previously identified in the Grandey and Wang (2015) study.

The ~~estimated temperature~~ we find [the estimated temperature](#) responses per unit change in DMS-derived sulfur flux ( $0.027$  or  $0.019 \text{ K per Tg year}^{-1}$  of sulfur) are lower than those reported in the Six et al. (2013) ~~and Schwinger et al. (2017) studies by approximately a factor of two~~ ( $0.046$ - $0.096 \text{ K per Tg year}^{-1}$  of sulfur, medium impact scenario) ~~and Schwinger et al. (2017)~~ ( $0.041 \text{ K per Tg year}^{-1}$  of sulfur) [studies](#). The temperature response sensitivities calculated here are comparable to those  
35 given in Grandey and Wang (2015) ( $0.029 \text{ K per Tg year}^{-1}$  of sulfur). Without further information, it is difficult to speculate

on the cause of the discrepancy between the results presented here and those in Six et al. (2013) and Schwinger et al. (2017). ~~The cause of the discrepancy between the results presented here and those in Six et al. (2013) and Schwinger et al. (2017) is difficult to speculate upon without further information.~~ However, the discrepancy between these results suggests the need for better observational constraints, and highlights the complexity of the DMS-aerosol-cloud system.

5 Natural aerosols account for a significant source of uncertainty in climate modelling and radiation budgets (Carslaw et al., 2013). Our study uses the Lana et al. (2011) DMS<sub>w</sub> climatology with the Liss and Merlivat (1986) flux parameterization. ~~Whilst~~ Though this dataset and method are commonplace for DMS-climate studies, both are limited by sparse observations and ~~biases~~ uncertainties (Tesdal et al., 2016). For example:–, recent studies have indicated that coral reefs ~~are an as of yet unaccounted for source of marine DMS (Hopkins et al., 2016; Swan et al., 2017; Jones et al., 2017)~~ produce significant  
10 amounts of DMS, and are an unaccounted for source of sulfur (Hopkins et al., 2016; Swan et al., 2017; Jones et al., 2017). Furthermore, larger concentrations and/or fluxes of DMS than what we currently consider have also been found at the poles, especially around sea ice and polynyas (Nomura et al., 2011; Jarnikova and Tortell, 2016; Mungall et al., 2016; Kim et al., 2017).

Observational deficiencies become particularly relevant when considering the stratiform cloud deck regions. In the Lana et al.  
15 (2011) dataset, the SEP region contains only two ship campaigns collecting measurements in January and February. The cloud deck in the Southern Hemisphere eastern basin of the Indian Ocean has no DMS<sub>w</sub> observations. The higher susceptibility of cloud and precipitation to changes in DMS in these regions suggest that they should be a priority for future atmospheric composition ~~observation campaigns. Providing further motivation for study in these areas, Six et al. (2013); Schwinger et al. (2017) predict an increase in in these stratiform cloud regions (despite a global mean decrease) due to climate change by the end of the 21st century.~~ field campaigns.

~~Global perturbations of in this study indicate that significant changes in meteorology are restricted to key marine boundary layer cloud regions despite global responses in chemistry and aerosol fields. This suggests that whilst inclusion of updated concentrations may have an impact on regional chemistry and aerosol fields, the impact on regional climate may be small, and in terms of global climate, insignificant. This hypothesis will be tested in future studies. For example, a follow-up to this study will examine the role of coral reef-derived DMS in regional and global climate.–~~

To place the conclusions of this study into a broader perspective, we must consider the DMS-climate system within the context of anthropogenic climate change. ~~Hopkins et al. (2011); Six et al. (2013) despite the uncertainties discussed above and in Section 1.~~ As discussed in the introduction and above, a better understanding of current global DMS is essential before future scenarios can be considered with certainty. Nevertheless, Hopkins et al. (2011), Six et al. (2013) and Schwinger et al. (2017)  
30 have ~~demonstrated that~~ suggested that global production of DMS by marine phytoplankton is vulnerable to ocean acidification, amongst other oceanic changes expected with global warming, for example impacts on nutrient availability, salinity, mixed layer depths, and light penetration (Kloster et al., 2007). ~~Whilst~~ While both the Six et al. (2013) and Schwinger et al. (2017) temperature responses are much larger than those found here, our results imply a 25% decrease in flux<sub>DMS</sub> would result in an increase of ~~0.4~~ 0.12 ~~°C~~ K (0.07 to 0.16, 10th and 90th confidence intervals) globally. Considering the current Paris Agreement  
35 target of limiting global warming to 1.5-2.0 K, the sensitivity of ocean-derived sulfate aerosol to warming temperatures and

ocean acidification becomes important. Strategies to mitigate anthropogenic climate change must consider not only the effect of increased CO<sub>2</sub> on temperatures, but also on ocean pH. Mitigating only temperature increases, (e.g. via solar radiation management, ~~may have a~~) may have short term cooling effects, however, without removing CO<sub>2</sub> from the atmosphere, ocean acidification will continue to impact ~~on~~ marine life, and as demonstrated here, the climate.

- 5 *Author contributions.* SF completed the ACCESS-UKCA simulations, analysis and the initial draft of this manuscript. MTW developed the initial model setup and provided advice as to the specific setup requirements of this study. MTW helped guide the analysis and contributed significantly to the revisions of this manuscript. ZN provided the FAIR analysis in this manuscript and contributed to the revisions of this manuscript. RS and TL provided advice and guidance on the direction of this study and contributed to the revisions of this manuscript.

*Competing interests.* The authors declare that they have no conflict of interest.

- 10 *Acknowledgements.* SLF would like to thank P. J. Rayner and his research group for their helpful discussions. SLF, RS, ~~TPL~~ are supported by the Australian Research Council (ARC) Centre of Excellence for Climate System Science (CE110001028). ~~ZN~~ TPL is supported by the Australian Research Council (ARC) Centre of Excellence for Climate Extremes (CE170100023). SLF, RS, were supported by the ARC Discovery Project: Great Barrier Reef as a significant source of climatically relevant aerosol particles (DP150101649). MTW is supported by the Earth System and Climate Change Hub of the Australian Government's National Environmental Science Programme (NESP). This
- 15 research was undertaken with the assistance of resources and services from the National Computational Infrastructure (Project q90), which is supported by the Australian Government. SF and ZN are supported by the Australian Government Research Training Program Scholarship.

## References

- Abraham, N. L., Archibald, A. T., Bellouin, N., Boucher, O., Braesicke, P., Bushell, A., Carslaw, K., Collins, B., Dalvi, M., Emmerson, K., Folberth, G., Haywood, J., Johnson, C., Kipling, Z., Macintyre, H., Mann, G., Telford, P., Merikanto, J., Morgenstern, O., Connor, F. O., Ord, C., Osprey, S., Pringle, K., Pyle, J., Rae, J., Reddington, C., Savage, N., Spracklen, D., Stier, P., and West, R.: Unified Model Documentation Paper No. 84: United Kingdom Chemistry and Aerosol (UKCA) Technical Description MetUM Version 8.4, Tech. rep., UK Met Office, 2012.
- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing., *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03137.1>, 2004.
- Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness., *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Anderson, T. R., Spall, S. A., Yool, A., Cipollini, P., Challenor, P. G., and Fasham, M. J.: Global fields of sea surface dimethylsulfide predicted from chlorophyll, nutrients and light, *J. Mar. Syst.*, 30, 1–20, [https://doi.org/10.1016/S0924-7963\(01\)00028-8](https://doi.org/10.1016/S0924-7963(01)00028-8), 2001.
- Ayers, G. P. and Cainey, J. M.: The CLAW hypothesis: A review of the major developments, *Environ. Chem.*, 4, 366–374, <https://doi.org/10.1071/EN07080>, 2007.
- Bell, T. G., Landwehr, S., Miller, S. D., De Bruyn, W. J., Callaghan, A. H., Scanlon, B., Ward, B., Yang, M., and Saltzman, E. S.: Estimation of bubble-mediated air-sea gas exchange from concurrent DMS and CO<sub>2</sub> transfer velocities at intermediate-high wind speeds, *Atmos. Chem. Phys.*, 17, 9019–9033, <https://doi.org/10.5194/acp-17-9019-2017>, 2017.
- Bellouin, N., Mann, G. W., Woodhouse, M. T., Johnson, C., Carslaw, K. S., and Dalvi, M.: Impact of the modal aerosol scheme GLOMAP-mode on aerosol forcing in the hadley centre global environmental model, *Atmos. Chem. Phys.*, 13, 3027–3044, <https://doi.org/10.5194/acp-13-3027-2013>, 2013.
- Belviso, S., Bopp, L., Moulin, C., Orr, J. C., Anderson, T. R., Aumont, O., Chu, S., Elliott, S., Maltrud, M. E., and Simó, R.: Comparison of global climatological maps of sea surface dimethyl sulfide, *Global Biogeochem. Cycles*, 18, <https://doi.org/10.1029/2003GB002193>, 2004a.
- Belviso, S., Moulin, C., Bopp, L., and Stefels, J.: Assessment of a global climatology of oceanic dimethylsulfide (DMS) concentration based on SeaWiFS imagery (1998-2001), *Can. J. Fish. Aquat. Sci.*, 61, 804–816, <https://doi.org/10.1139/F04-001>, 2004b.
- Bi, D., Dix, M., Marsland, S. J., O’Farrell, S., Rashid, H. A., Uotila, P., Hirst, A. C., Kowalczyk, E., Golebiewski, M., Sullivan, A., Yan, H., Hannah, N., Franklin, C., Sun, Z., Vohralik, P., Watterson, I., Zhou, X., Fiedler, R., Collier, M., Ma, Y., Noonan, J., Stevens, L., Uhe, P., Zhu, H., Griffies, S. M., Hill, R., Harris, C., and Puri, K.: The ACCESS coupled model: description, control climate and evaluation, *Aust. Met. Oceanogr. J.*, 63, 41–64, 2013.
- Bodas-Salcedo, A., Williams, K. D., Ringer, M. A., Beau, I., Cole, J. N. S., Dufresne, J. L., Koshiro, T., Stevens, B., Wang, Z., and Yokohata, T.: Origins of the solar radiation biases over the Southern Ocean in CFMIP2 models, *J. Climate*, 27, 41–56, <https://doi.org/10.1175/JCLI-D-13-00169.1>, 2014.
- Bodas-Salcedo, A., Hill, P. G., Furtado, K., Williams, K. D., Field, P. R., Manners, J. C., Hyder, P., and Kato, S.: Large contribution of supercooled liquid clouds to the solar radiation budget of the Southern Ocean, *J. Climate*, 29, 4213–4228, <https://doi.org/10.1175/JCLI-D-15-0564.1>, 2016.
- Bony, S. and Dufresne, J. L.: Marine boundary layer clouds at the heart of tropical cloud feedback uncertainties in climate models, *Geophys. Res. Lett.*, 32, 1–4, <https://doi.org/10.1029/2005GL023851>, 2005.

- Bopp, L., Boucher, O., Aumont, O., Belviso, S., Dufresne, J.-L., Pham, M., and Monfray, P.: Will marine dimethylsulfide emissions amplify or alleviate global warming? A model study, *Can. J. Fish. Aquat. Sci.*, 61, 826–835, <https://doi.org/10.1139/f04-045>, 2004.
- Boucher, O. and Reddy, M. S.: Climate trade-off between black carbon and carbon dioxide emissions, *Energ. Policy*, 36, 193–200, <https://doi.org/10.1016/j.enpol.2007.08.039>, 2008.
- 5 Cameron-Smith, P., Elliott, S., Maltrud, M., Erickson, D., and Wingenter, O.: Changes in dimethyl sulfide oceanic distribution due to climate change, *Geophys. Res. Lett.*, 38, 1–5, <https://doi.org/10.1029/2011GL047069>, 2011.
- Carlsaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. a., and Pierce, J. R.: Large contribution of natural aerosols to uncertainty in indirect forcing., *Nature*, 503, 67–71, <https://doi.org/10.1038/nature12674>, <http://www.ncbi.nlm.nih.gov/pubmed/24201280>, 2013.
- 10 Cesana, G. and Chepfer, H.: How well do climate models simulate cloud vertical structure? A comparison between CALIPSO-GOCCP satellite observations and CMIP5 models, *Geophys. Res. Lett.*, 39, 1–6, <https://doi.org/10.1029/2012GL053153>, 2012.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655–661, <https://doi.org/10.1038/326655a0>, 1987.
- Chepfer, H., Bony, S., Winker, D., Cesana, G., Dufresne, J. L., Minnis, P., Stubenrauch, C. J., and Zeng, S.: The GCM-oriented CALIPSO cloud product (CALIPSO-GOCCP), *J. Geophys. Res-Atmos*, 115, 1–13, <https://doi.org/10.1029/2009JD012251>, 2010.
- 15 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 20 Elliott, S.: Dependence of DMS global sea-air flux distribution on transfer velocity and concentration field type, *J. Geophys. Res-Biogeophys.*, 114, 1–18, <https://doi.org/10.1029/2008JG000710>, 2009.
- Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M.: Evaluation of Climate Models, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 741–866, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324>, 2013.
- Gabric, A. J., Simó, R., Cropp, R. A., Hirst, A. C., and Dachs, J.: Modeling estimates of the global emission of dimethylsulfide under enhanced greenhouse conditions, *Global Biogeochem. Cy.*, 18, <https://doi.org/10.1029/2003GB002183>, 2004.
- 30 Gali, M., Devred, E., Levasseur, M., Royer, S. J., and Babin, M.: A remote sensing algorithm for planktonic dimethylsulfoniopropionate (DMSP) and an analysis of global patterns, *Remote Sens. Environ.*, 171, 171–184, <https://doi.org/10.1016/j.rse.2015.10.012>, [http://dx.doi.org/10.1016/j.rse.2015.10.012https://www.researchgate.net/profile/Sarah-Jeanne/\\_/publication/283054397/\\_/A/\\_remote/\\_sensing/\\_algorithm/\\_for/\\_planktonic/\\_dimethylsulfoniopropionate/\\_/DMSP/\\_/and/\\_an/\\_analysis/\\_of/\\_global/\\_patterns/links/5637d44908ae7f7](http://dx.doi.org/10.1016/j.rse.2015.10.012https://www.researchgate.net/profile/Sarah-Jeanne/_/publication/283054397/_/A/_remote/_sensing/_algorithm/_for/_planktonic/_dimethylsulfoniopropionate/_/DMSP/_/and/_an/_analysis/_of/_global/_patterns/links/5637d44908ae7f7), 2015.
- 35 Grandey, B. S. and Wang, C.: Enhanced marine sulphur emissions offset global warming and impact rainfall., *Sci. Rep-UK*, 5, 13 055, <https://doi.org/10.1038/srep13055>, <http://www.nature.com/srep/2015/150821/srep13055/full/srep13055.html>, 2015.

- Halloran, P. R., Bell, T. G., and Totterdell, I. J.: Can we trust empirical marine DMS parameterisations within projections of future climate?, *Biogeosciences*, 7, 1645–1656, <https://doi.org/10.5194/bg-7-1645-2010>, 2010.
- Hopkins, F. E., Nightingale, P., and Liss, P.: Ocean acidification, in: *Ocean acidification*, edited by Gattuso, J.-P. and Hansson, L., p. 326, Oxford University Press, 2011.
- 5 Hopkins, F. E., Bell, T. G., Yang, M., Suggett, D. J., and Steinke, M.: Air exposure of coral is a significant source of dimethylsulfide (DMS) to the atmosphere, *Sci. Rep-UK*, 6, 36 031, <https://doi.org/10.1038/srep36031>, 2016.
- Hopkins, F. E., Nightingale, P. D., Stephens, J. A., Moore, C. M., Cripps, G. L., and Archer, S. D.: Dimethylsulfide ( DMS ) production in polar oceans may be resilient to ocean acidification ., *Biogeosciences Discuss.*, pp. 1–42, <https://doi.org/https://doi.org/10.5194/bg-2018-55>, 2018.
- 10 Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., Hong, Y., Bowman, K. P., and Stocker, E. F.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales, *J. Hydrometeorol.*, 8, 38–55, <https://doi.org/10.1175/JHM560.1>, 2007.
- Jardine, K., Yañez-Serrano, A. M., Williams, J., Kunert, N., Jardine, A., Taylor, T., Abrell, L., Artaxo, P., Guenther, A., Hewitt, C. N., House, E., Florentino, A. P., Manzi, A., Higuchi, N., Kesselmeier, J., Behrendt, T., Veres, P. R., Derstroff, B., Fuentes, J. D., Martin, S. T., and
- 15 Andreae, M. O.: Dimethyl sulfide in the Amazon rain forest, *Global Biogeochem. Cy.*, 29, 19–32, <https://doi.org/10.1002/2014GB004969>, <http://doi.wiley.com/10.1002/2014GB004969>, 2015.
- Jarnikova, T. and Tortell, P. D.: Towards a revised climatology of summertime dimethylsulfide concentrations and sea – air fluxes in the Southern Ocean, *Environ. Chem.*, 13, 364–378, <https://doi.org/10.1071/EN14272>, 2016.
- Jones, G., Curran, M., Swan, H., and Deschaseaux, E.: Dimethylsulfide and Coral Bleaching: Links to Solar Radiation, Low Level
- 20 Cloud and the Regulation of Seawater Temperatures and Climate in the Great Barrier Reef, *Am. J. Climate Change*, 06, 328–359, <https://doi.org/10.4236/ajcc.2017.62017>, <http://www.scirp.org/journal/PaperDownload.aspx?DOI=10.4236/ajcc.2017.62017>, 2017.
- Kato, S., Loeb, N. G., Rose, F. G., Doelling, D. R., Rutan, D. A., Caldwell, T. E., Yu, L., and Weller, R. A.: Surface irradiances consistent with CERES-derived top-of-atmosphere shortwave and longwave irradiances, *J. Climate*, 26, 2719–2740, <https://doi.org/10.1175/JCLI-D-12-00436.1>, 2013.
- 25 Kettle, A. J. and Andreae, M.: Flux of dimethylsulfide from the oceans : A comparison of updated data sets and flux models, *J. Geophys. Res.*, 105, 26,793–26,808, <https://doi.org/10.1029/2000JD900252>, 2000.
- Kettle, A. J., Amouroux, D., Andreae, T. W., Bates, T. S., Berresheim, H., Bingemer, H., Boniforti, R., Helas, G., Leck, C., Maspero, M., Matrai, P., McTaggart, A. R., Mihalopoulos, N., Nguyen, B. C., Novo, A., Putaud, J. P., Rapsomanikis, S., Roberts, G., Schebeske, G., Sharma, S., Simó, R., Staubes, R., Turner, S., and Uher, G.: A global data base of sea surface dimethyl sulfide (DMS) measurements and
- 30 a simple model to predict sea surface DMS as a function of latitude, longitude, and month, *Global Biogeochem. Cy.*, 13, 399–444, 1999.
- Kim, I., Hahm, D., Park, K., Lee, Y., Choi, J.-O., Zhang, M., Chen, L., Kim, H.-C., and Lee, S.: Characteristics of the horizontal and vertical distributions of dimethyl sulfide throughout the Amundsen Sea Polynya, *Sci. Total Environ.*, 584, 154–163, <https://doi.org/10.1016/j.scitotenv.2017.01.165>, <http://dx.doi.org/10.1016/j.scitotenv.2017.01.165>, 2017.
- Klein, S. A., Zhang, Y., Zelinka, M. D., Pincus, R., Boyle, J., and Gleckler, P. J.: Are climate model simulations of clouds improving? An
- 35 evaluation using the ISCCP simulator, *J. Geophys. Res-Atmos*, 118, 1329–1342, <https://doi.org/10.1002/jgrd.50141>, 2013.
- Kloster, S., Six, K. D., Feichter, J., Maier-Reimer, E., Roeckner, E., Wetzol, P., Stier, P., and Esch, M.: Response of dimethylsulfide (DMS) in the ocean and atmosphere to global warming, *J. Geophys. Res. Biogeosciences*, 112, 1–13, <https://doi.org/10.1029/2006JG000224>, 2007.


- Korhonen, H., Carslaw, K. S., Spracklen, D. V., Mann, G. W., and Woodhouse, M. T.: Influence of oceanic dimethyl sulfide emissions on cloud condensation nuclei concentrations and seasonality over the remote Southern Hemisphere oceans: A global model study, *J. Geophys. Res.-Atmos*, 113, 1–16, <https://doi.org/10.1029/2007JD009718>, 2008.
- Kulmala, M., Laaksonen, A., and Pirjola, L.: Parameterizations for sulfuric acid/water nucleation rates, *J. Geophys. Res.*, 103, 8301, <https://doi.org/10.1029/97JD03718>, 1998.
- Kushta, J., Kallos, G., Astitha, M., Solomos, S., Spyrou, C., Mitsakou, C., and Lelieveld, J.: Impact of natural aerosols on atmospheric radiation and consequent feedbacks with the meteorological and photochemical state of the atmosphere, *J. Geophys. Res.-Atmos*, 119, 1463–1491, <https://doi.org/10.1002/2013JD020714>, 2014.
- Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and Van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, <https://doi.org/10.5194/acp-10-7017-2010>, 2010.
- Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochem. Cy.*, 25, 1–17, <https://doi.org/10.1029/2010GB003850>, 2011.
- Leon, D. C., Wang, Z., and Liu, D.: Climatology of drizzle in marine boundary layer clouds based on 1 year of data from CloudSat and Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO), *J. Geophys. Res.-Atmos*, 114, <https://doi.org/10.1029/2008JD009835>, 2009.
- Liss, P. S. and Merlivat, L.: Air-Sea Gas Exchange Rates: Introduction and Synthesis, in: *The Role of Air-Sea Exchange in Geochemical Cycling*, edited by Buat-Ménard, P., pp. 113–127, Springer Netherlands, Dordrecht, [https://doi.org/10.1007/978-94-009-4738-2\\_5](https://doi.org/10.1007/978-94-009-4738-2_5), 1986.
- Loeb, N. G., Wielicki, B. A., Doelling, D. R., Smith, G. L., Keyes, D. F., Kato, S., Manalo-Smith, N., and Wong, T.: Toward optimal closure of the Earth's top-of-atmosphere radiation budget, *J. Climate*, 22, 748–766, <https://doi.org/10.1175/2008JCLI2637.1>, 2009.
- Mace, G. G. and Zhang, Q.: The CloudSat radar-lidar geometrical profile product (RL-GeoProf): Updates, improvements, and selected results, *J. Geophys. Res.-Atmos*, 119, 9441–9462, <https://doi.org/10.1002/2013JD021374>, 2014.
- Mahajan, A. S., Fadnavis, S., Thomas, M. a., Pozzoli, L., Gupta, S., Royer, S.-j., Saiz-Lopez, A., and Simó, R.: Quantifying the impacts of an updated global dimethyl sulfide climatology on cloud microphysics and aerosol radiative forcing, *J. Geophys. Res. Atmos.*, 120, 2524–2536, <https://doi.org/10.1002/2014JD022687>, <http://doi.wiley.com/10.1002/2014JD022687>, 2015.
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, *Geosci. Model Dev*, 3, 519–551, <https://doi.org/10.5194/gmd-3-519-2010>, 2010.
- Mann, G. W., Carslaw, K. S., Ridley, D. A., Spracklen, D. V., Pringle, K. J., Merikanto, J., Korhonen, H., Schwarz, J. P., Lee, L. A., Manktelow, P. T., Woodhouse, M. T., Schmidt, A., Breider, T. J., Emmerson, K. M., Reddington, C. L., Chipperfield, M. P., and Pickering, S. J.: Intercomparison of modal and sectional aerosol microphysics representations within the same 3-D global chemical transport model, *Atmos. Chem. Phys.*, 12, 4449–4476, <https://doi.org/10.5194/acp-12-4449-2012>, 2012.
- McCormick, R. A. and Ludwig, J. H.: Climate Modification by Atmospheric Aerosols, *Science*, 156, 1358–1359, <https://doi.org/10.1126/science.156.3780.1358>, <http://www.sciencemag.org/cgi/doi/10.1126/science.156.3780.1358>, 1967.

- Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S. H., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the role of organics in aerosol particle formation under atmospheric conditions., *P. Natl. Acad. Sci. USA*, 107, 6646–51, <https://doi.org/10.1073/pnas.0911330107>, 2010.
- 5 Millar, J. R., Nicholls, Z. R., Friedlingstein, P., and Allen, M. R.: A modified impulse-response representation of the global near-surface air temperature and atmospheric concentration response to carbon dioxide emissions, *Atmos. Chem. Phys.*, 17, 7213–7228, <https://doi.org/10.5194/acp-17-7213-2017>, 2017.
- Mungall, E. L., Croft, B., Lizotte, M., Thomas, J. L., Murphy, J. G., Lévasseur, M., Martin, R. V., Wentzell, J. J. B., Liggio, J., and Abbatt, J. P. D.: Dimethyl sulfide in the summertime Arctic atmosphere: measurements and source sensitivity simulations, *Atmos. Chem. Phys.*, 16, 6665–6680, <https://doi.org/10.5194/acp-16-6665-2016>, <http://www.atmos-chem-phys.net/16/6665/2016/>, 2016.
- 10 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J. F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., Van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J. H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys.*, 13, 1853–1877, <https://doi.org/10.5194/acp-13-1853-2013>, 2013a.
- 15 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 659–740, <https://doi.org/10.1017/CBO9781107415324.018>, 2013b.
- Nam, C., Bony, S., Dufresne, J. L., and Chepfer, H.: The too few, too bright tropical low-cloud problem in CMIP5 models, *Geophys. Res. Lett.*, 39, 1–7, <https://doi.org/10.1029/2012GL053421>, 2012.
- Nightingale, D., Malin, G., Law, C. S., Watson, J., Liss, S., and Liddicoat, I.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cy.*, 14, 373–387, <https://doi.org/10.1029/1999GB900091>, 2000.
- Nomura, D., Kasamatsu, N., Tateyama, K., Kudoh, S., and Fukuchi, M.: DMSP and DMS in coastal fast ice and under-ice water of Lützow-Holm Bay, eastern Antarctica, *Cont. Shelf Res.*, 31, 1377–1383, <https://doi.org/10.1016/j.csr.2011.05.017>, 2011.
- 25 Protat, A., Young, S. A., McFarlane, S. A., L’Ecuyer, T., Mace, G. G., Comstock, J. M., Long, C. N., Berry, E., and Delanoë, J.: Reconciling ground-based and space-based estimates of the frequency of occurrence and radiative effect of clouds around Darwin, Australia, *J. Appl. Meteorol. Clim.*, 53, 456–478, <https://doi.org/10.1175/JAMC-D-13-072.1>, 2014.
- Quinn, P. K. and Bates, T. S.: The case against climate regulation via oceanic phytoplankton sulphur emissions, *Nature*, 480, 51–56, <https://doi.org/10.1038/nature10580>, 2011.
- 30 Rosenfeld, D., Sherwood, S., Wood, R., and Donner, L.: Climate Effects of Aerosol-Cloud Interactions, *Science*, 343, 379–380, <https://doi.org/10.1126/science.1247490>, <http://www.sciencemag.org/cgi/doi/10.1126/science.1247490>, 2014.
- Rotstain, L. D., Collier, M. A., and Luo, J.-J.: Effects of declining aerosols on projections of zonally averaged tropical precipitation, *Environ. Res. Lett.*, 10, 44018, <https://doi.org/10.1088/1748-9326/10/4/044018>, <http://stacks.iop.org/1748-9326/10/i=4/a=044018>, 2015.
- Saltzman, E. S., King, D. B., Holmen, K., and Leck, C.: Experimental-determination of the Diffusion-coefficient of Dimethylsulfide In Water, *Journal of Geophysical Research-Oceans*, 98, 16481–16486, 1993.
- 35 Schwinger, J., Tjiputra, J., Goris, N., Six, K., Kirkevåg, A., Seland, Ø., Heinze, C., and Ilyina, T.: Amplification of global warming through pH-dependence of DMS-production simulated with a fully coupled Earth system model, *Biogeosciences Discuss.*, 20, 1–26, <https://doi.org/10.5194/bg-2017-33>, 2017.




- Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 2: Maritime vs. continental deep convective storms, *Meteorol. Atmos. Phys.*, 92, 67–82, <https://doi.org/10.1007/s00703-005-0113-3>, 2006.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosolcloud interactions in the climate system, *P. Natl. Acad. Sci. USA*, 113, 5781–5790, <https://doi.org/10.1073/pnas.1514043113>, <http://www.pnas.org/lookup/doi/10.1073/pnas.1514043113>, 2016.
- Sheng, J. X., Weisenstein, D. K., Luo, B. P., Rozanov, E., Stenke, A., Anet, J., Bingemer, H., and Peter, T.: Global atmospheric sulfur budget under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *J. Geophys. Res-Atmos*, 120, 256–276, <https://doi.org/10.1002/2014JD021985>, 2015.
- Simó, R. and Dachs, J.: Global ocean emission of dimethylsulfide predicted from biogeophysical data, *Global Biogeochem. Cycles*, 16, 26–1–26–10, <https://doi.org/10.1029/2001GB001829>, <http://doi.wiley.com/10.1029/2001GB001829>, 2002.
- Six, K. D., Kloster, S., Ilyina, T., Archer, S. D., Zhang, K., and Maier-Reimer, E.: Global warming amplified by reduced sulphur fluxes as a result of ocean acidification, *Nat. Clim. Change*, 3, 975–978, <https://doi.org/10.1038/nclimate1981>, <http://www.nature.com/nclimate/journal/v3/n11/full/nclimate1981.html>{% }5Cn<http://www.nature.com/nclimate/journal/v3/n11/pdf/nclimate1981.pdf>, 2013.
- Stephens, G. L., L’Ecuyer, T., Forbes, R., Gettlemen, A., Golaz, J. C., Bodas-Salcedo, A., Suzuki, K., Gabriel, P., and Haynes, J.: Dreary state of precipitation in global models, *J. Geophys. Res-Atmos*, 115, <https://doi.org/10.1029/2010JD014532>, 2010.
- Stephens, G. L., Li, J., Wild, M., Clayson, C. A., Loeb, N. G., Kato, S., L’Ecuyer, T., Stackhouse, P. W., Lebsock, M., and Andrews, T.: An update on Earth’s energy balance in light of the latest global observations, *Nat. Geosci.*, 5, 691–696, <https://doi.org/10.1038/ngeo1580>, <http://www.nature.com/doi/10.1038/ngeo1580>, 2012.
- Swan, H. B., Jones, G. B., Deschaseaux, E. S. M., and Eyre, B. D.: Coral reef origins of atmospheric dimethylsulfide at Heron Island, southern Great Barrier Reef, Australia, *Biogeosciences Discuss.*, 14, 229–239, <https://doi.org/10.5194/bg-2016-387>, <http://www.biogeosciences-discuss.net/bg-2016-387/>, 2017.
- Taylor, K. E., Williamson, D., and Zwiers, F. W.: Program for Climate Model Diagnosis and Intercomparison AMIP Sea Surface Temperature and Sea Ice Concentration Boundary Conditions, <http://www-pcmdi.llnl.gov/projects/amip/AMIP2EXPDSN/BCS/index.php>, 2015.
- Tesdal, J. E., Christian, J. R., Monahan, A. H., and Von Salzen, K.: Evaluation of diverse approaches for estimating sea-surface DMS concentration and air-sea exchange at global scale, *Environ. Chem.*, 13, 390–412, <https://doi.org/10.1071/EN14255>, 2016.
- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Rast, S., Devasthale, A., Kloster, S., Feichter, J., and Lenton, T. M.: Quantification of DMS aerosol-cloud-climate interactions using the ECHAM5-HAMMOZ model in a current climate scenario, *Atmos. Chem. Phys.*, 10, 7425–7438, <https://doi.org/10.5194/acp-10-7425-2010>, 2010.
- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Devasthale, A., Kloster, S., Rast, S., Feichter, J., and Lenton, T. M.: Rate of non-linearity in DMS aerosol-cloud-climate interactions, *Atmos. Chem. Phys.*, 11, 11 175–11 183, <https://doi.org/10.5194/acp-11-11175-2011>, 2011.
- Twomey, S.: Pollution and the Planetary Albedo, *Atmos. Environ.*, 8, 1251–1256, <http://linkinghub.elsevier.com/retrieve/pii/S1352231007009843>, 1974.
- Vallina, S. M., Simó, R., and Gassó, S.: What controls CCN seasonality in the Southern Ocean? A statistical analysis based on satellite-derived chlorophyll and CCN and model-estimated OH radical and rainfall, *Global Biogeochem. Cy.*, 20, 1–13, <https://doi.org/10.1029/2005GB002597>, 2006.


- van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997-2015, *Earth Syst. Sci. Data Discuss.*, pp. 1–43, <https://doi.org/10.5194/essd-2016-62>, <http://www.earth-syst-sci-data-discuss.net/essd-2016-62/>, 2017.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J. F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: An overview, *Climatic Change*, 109, 5–31, <https://doi.org/10.1007/s10584-011-0148-z>, 2011.
- Vlahos, P. and Monahan, E. C.: A generalized model for the air-sea transfer of dimethyl sulfide at high wind speeds, *Geophys. Res. Lett.*, 36, L21 605, <https://doi.org/10.1029/2009GL040695>, <http://doi.wiley.com/10.1029/2009GL040695>, 2009.
- Wang, H. and Su, W.: Evaluating and understanding top of the atmosphere cloud radiative effects in Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5) Coupled Model Intercomparison Project Phase 5 (CMIP5) models using satellite observations, *J. Geophys. Res-Atmos*, 118, 683–699, <https://doi.org/10.1029/2012JD018619>, 2013.
- Wanninkhof, R.: Relationship Between Wind Speed and Gas Exchange, *J. Geophys. Res.*, 97, 7373–7382, <https://doi.org/10.1029/92JC00188>, 1992.
- Wilks, D. S.: Statistical Methods in the Atmospheric Sciences, vol. 100, Elsevier, 3rd edn., <https://doi.org/10.1016/B978-0-12-385022-5.00026-9>, <http://www.sciencedirect.com/science/article/pii/B9780123850225000269>, 2011.
- Woodhouse, M. T., Carslaw, K. S., Mann, G. W., Vallina, S. M., Vogt, M., Halloran, P. R., and Boucher, O.: Low sensitivity of cloud condensation nuclei to changes in the sea-air flux of dimethyl-sulphide, *Atmos. Chem. Phys.*, 10, 7545–7559, <https://doi.org/10.5194/acp-10-7545-2010>, 2010.
- Woodhouse, M. T., Mann, G. W., Carslaw, K. S., and Boucher, O.: Sensitivity of cloud condensation nuclei to regional changes in dimethyl-sulphide emissions, *Atmos. Chem. Phys.*, 13, 2723–2733, <https://doi.org/10.5194/acp-13-2723-2013>, 2013.
- Woodhouse, M. T., Luhar, A. K., Stevens, L., Galbally, I., Thatcher, M., Uhe, P., Noonan, J., and Molloy, S.: Australian reactive-gas emissions in a global chemistry-climate model and initial results, *Air Qual. Clim. Change*, 49, 31–38, 2015.
- Woodward, S.: Modeling the atmospheric life cycle and radiative impact of mineral dust in the Hadley Centre climate model, *J. Geophys. Res-Atmos*, 106, 18 155–18 166, <https://doi.org/10.1029/2000JD900795>, 2001.
- Yoch, D. C.: Dimethylsulfoniopropionate : Its Sources , Role in the Marine Food Web , and Biological Degradation to Dimethylsulfide, *Appl. Environ. Microb.*, 68, 5804–5815, <https://doi.org/10.1128/AEM.68.12.5804-5815.2002>, 2002.




**Figure 1.** Schematic diagram showing the ocean-atmosphere sulfur life cycle and climate-relevant processes. Acronyms are defined as follows: sea surface temperatures (SSTs), methane sulfonic acid (MSA), dimethyl sulfoniopropionate (DMSP), dimethyl sulfoxide (DMSO), dimethyl sulfide (DMS), cloud condensation nuclei (CCN), cloud droplet number (CDN)




**Figure 2.** The annual mean concentrations of a) the Lana et al. (2011) DMS<sub>w</sub> climatology, b) the DMS<sub>w</sub> field of the second experimental run: zonal maximum DMS<sub>w</sub>. Additionally, three regions of interest are shown in a) by red boxes, the Australian region:  $45^{\circ}S$ - $10^{\circ}S$ ,  $110^{\circ}E$ - $160^{\circ}E$ , the South



**Figure 3.** The 2006-2009 annual mean of the ACCESS-UKCA CTL-Ctl (first column) compared to the CALIPSO-GOCCP (Chepfer et al., 2010) climatology (second column), with the absolute-relative differences between the two shown in the third column (model - observations). The top row shows the low cloud fraction, the middle row shows the middle cloud fraction and the bottom row the high cloud fraction




**Figure 4.** As for Fig. 3 but for annual means for 2000-2009, where the top row shows the TOA outgoing LW radiation and the second row: TOA outgoing SW radiation and the third row: surface incoming SW radiation. The observations are from the CERES-EBAF - TOA and Surface Ed. 4.0 (Loeb et al., 2009; Kato et al., 2013); all units are in  $\text{W m}^{-2}$




**Figure 5.** The mean (2000-2009) annual total precipitation of a) the satellite climatology from TRMM (Huffman et al., 2007)), b) the ACCESS-UKCA climatology and c) the difference between ACCESS-UKCA and the TRMM and ACCESS-UKCA climatologies product (model - observations). Units are in mm year<sup>-1</sup>






**Figure 6.** Mean (2000-2009) values for the Ctl (first column), the difference between Exp.1 (zero DMS<sub>w</sub>) and the Ctl (second column) and the difference between Exp.2 (zonally enhanced DMS<sub>w</sub>) and the Ctl (third column). The first row shows the volume mixing ratio of SO<sub>2</sub> in ppb; the second row the volume mixing ratio of H<sub>2</sub>SO<sub>4</sub> in ppt







**Figure 7.** The vertical profiles of a)  $\text{SO}_2$ ; b)  $\text{H}_2\text{SO}_4$ ; c) nucleation mode number density; d) Aitken mode number density; e) accumulation mode number density; f) coarse mode number density; g)  $\text{N}_3$  nuclei number; h) cloud condensation nuclei number; and i) cloud droplet number. The solid lines represent the Ctl; dashed lines shows Exp.1; and dotted lines show Exp.2. Blue lines show the SO (SO) mean, red the Australian (Aus) region mean and green the South Eastern Pacific (SEP). All units are  $\text{cm}^{-3}$ , apart from a): ppb and b): ppt  $\times 10^{-3}$



**Figure 8.** As for Figure 6, but where the first row shows the number concentration of  $N_3$  (condensation nuclei); the second the number of cloud condensation nuclei greater than 70 nm dry diameter; and the third the cloud droplet number concentration. All units are in  $\text{cm}^{-3}$



**Figure 9.** Comparisons in of the low cloud fraction (as a percentage) (a-b) and incoming SW radiation at surface ( $\text{W m}^{-2}$ ) c-d) over the 2000-2009 period for Exp.1 (first column) and Exp.2 (second column) minus the Ctl. The absolute values for the Ctl of these fields can be seen in Fig. 3 and 4



**Figure 10.** Comparisons of the (a-c) total liquid water at 1700m height ( $Q_{cl}$ ), (d-f) large-scale rainfall, (g-i) convective rainfall over the 2000-2009 period. The Ctl absolute values are shown in the first column, and respectively Exp.1 and Exp.2 minus the Ctl in the second and third columns. Units are in  $\text{kg kg}^{-1}$  and  $\text{mm year}^{-1}$

**Table 1.** Summary of the three global simulations presented in this study, the DMS<sub>w</sub> climatology used and the annual mean (2000-2009) total global flux<sub>DMS</sub>

Simulation	DMS Climatology	Flux <sub>DMS</sub> (Tg year <sup>-1</sup> of sulfur)
Ctl	Lana et al. (2011)	17.41
Exp.1	Zero marine DMS	0.72
Exp.2	Zonal maximum DMS from Lana et al. (2011)	37.05

**Table 2.** Global and hemispheric means of the CCN sensitivity to the  $\text{flux}_{\text{DMS}}$  (as defined by Woodhouse et al. (2010, 2013)) in both absolute ( $\text{cm}^{-3}/\text{mg m}^2 \text{ day}^{-1}$ ) and relative terms, for Exp.1 and Exp.2.

Region	Exp.1 Absolute	Exp.2 Absolute	Exp.1 Relative	Exp.2 Relative
Global	16.9	12.4	0.048	0.036
SH	15.8	11.2	0.090	0.063
NH	18.6	14.5	0.029	0.023

**Table 3.** Summary of the global mean (2000-2009) radiation fields: absolute Ctl values for the TOA shortwave (SW) and longwave (LW) outgoing and Q\*; and the differences in these quantities resulting from Exp.1 and Exp.2 (from the Ctl) as well as the FAIR temperature response. The ranges shown for Exp.1-Ctl and Exp.2-Ctl indicate the 10th and 90th percentile confidence intervals

Simulation	TOA SW ↑ (W m <sup>-2</sup> )	TOA LW ↑ (W m <sup>-2</sup> )	Q* (W m <sup>-2</sup> )	FAIR response (K)
Ctl absolute values	101.79	241.04	-1.35	NA
Exp.1 - Ctl	-1.82 (-2.62 to -1.23)	0.13 (0.04 to 0.20)	1.69 (1.13 to 2.43)	0.45 (0.30 to 0.64)
Exp.2 - Ctl	1.57 (1.08 to 2.48)	-0.12 (-0.20 to -0.04)	-1.45 (-2.33 to -0.98)	-0.38 (-0.61 to -0.26)

**Table 4.** The estimated temperature response to perturbations in the flux<sub>DMS</sub> (K per Tg year<sup>-1</sup> of sulfur) for the current study's experiments (Exp.1 and Exp.2) and those found in the literature. The ranges shown for Exp.1 and Exp.2 indicate the 10th and 90th percentile confidence intervals

Experiment	K per Tg year <sup>-1</sup> of sulfur
Exp.1	0.027 <u>(0.018 to 0.038)</u>
Exp.2	0.019 <u>(0.013 to 0.031)</u>
Schwinger et al. (2017)	0.041
Six et al. (2013) - low pH impact scenario	0.03 - 0.060
Six et al. (2013) - medium pH impact scenario	0.046 - 0.096
Six et al. (2013) - high pH impact scenario	0.051 - 0.11
Grandey and Wang (2015)	0.029