

## Responses to reviewers:

We thank the two anonymous reviewers for their helpful comments and suggestions. We have revised the manuscript following the comments as described below. Reviewer comments are shown in blue. Our responses are shown in black. The revised texts are shown in italics.

### Reviewer #1

(1) Treatment of uncertainty: aside from several brief mentions of NMB or similar metrics, there is little discussion of model uncertainties. For example, uncertainties in emission inventories surely propagate into derived O<sub>3</sub> partitioning. In particular, it would be helpful to estimate the confidence in the values shown in Fig. 9, which is the key result of the paper.

Thanks for the comment. We added discussions on uncertainties of (1) emission inventory, (2) chemical mechanism, and (3) biogenic VOC measurements

(1) Emission inventory

Anthropogenic emissions were obtained from the Multi-resolution Emission Inventory for China (Li et al., 2017). Emission estimates from bottom-up inventories are uncertain due to lack of complete knowledge of human activities and emission from different sources. The overall uncertainties of the bottom-up SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, VOC, and PM<sub>2.5</sub> emission estimates are ±31%, ±37%, ±153%, ±78% and ±133%, respectively, with 95% confidence intervals (CI) (Zhang et al., 2009; Lei et al., 2011; Zhao et al., 2011; Lu et al., 2011; Kurokawa et al., 2013). We updated Table S1 and added text in the Section 2.3 of the revised manuscript to describe this issue.

*The emission estimates and uncertainties of VOCs, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, and PM<sub>2.5</sub> in the domain during the simulation period are summarized in Table S1.*

(2) Chemical mechanism

The uncertainty of chemical mechanism is stated in the comment (4).

(3) Biogenic VOCs

The uncertainties of biogenic VOC observation and simulation are stated in the comment (7). In addition, we added discussion about sub-month variability of the simulated biogenic VOCs in Section 3.2 in the revised manuscript.

*We also analyzed the temporal variation of simulated biogenic VOC during the whole simulation period and found the sub-month variability was relatively small (the standard deviation < 25%).*

(2) NO<sub>x</sub> measurements: the “NO<sub>x</sub>” measurement uses a hot molybdenum converter,

which is known to convert a lot more than NO<sub>2</sub> (probably most of NO<sub>y</sub>). This point may or may not impact the model measurement comparison shown in Fig. 3 depending on the NO<sub>y</sub> partitioning, but it should be acknowledged and addressed.

5 Thank you for pointing this out. The observed NO<sub>x</sub> (NO+NO<sub>2</sub>) concentration in the center of Xi'an was measured by a gas-phase chemiluminescence detection analyzer. The principle of this method is converting ambient NO<sub>2</sub> to NO on the hot surface of a molybdenum oxide (MoO) catalyst followed by a chemiluminescence detection of NO. The conversion of the thermal catalyst is not specific to NO<sub>2</sub>, but also other nitrogen-containing components (NO<sub>z</sub>: NO<sub>z</sub>=NO<sub>y</sub>-NO<sub>x</sub>, including PANs, HNO<sub>3</sub>, HO<sub>2</sub>NO<sub>2</sub>, HONO, RO<sub>2</sub>NO<sub>2</sub>, and organic nitrates), resulting in a higher NO<sub>2</sub> concentration. 10 However, despite its drawbacks, it remains a widely used method in air-quality monitoring networks and research, owing in part to its low cost, sensitive detection, and ease of operation.

15 Xu et al. (2013) evaluated this potential uncertainty of the MoO converter at four different sites in China, and found that the MoO converter worked well at the urban site (overestimation less than 10%), but overestimated NO<sub>2</sub> more than 30% at suburban and background sites. In the Guanzhong basin, we used the simulated NO<sub>y</sub> to evaluate the uncertainty of the NO<sub>x</sub> measurements, due to the lack of referenced NO<sub>y</sub> or NO<sub>x</sub> observation. The calculated results indicated that the NO<sub>z</sub> accounted for 11% of the total NO<sub>y</sub> at the urban Xi'an during the no-raining period of Fig. 3. We noted the uncertainty in the NO<sub>x</sub> measurements, but thought this would not crucially impact the model-measurement comparison. 20

We modified the text in Section 2.1.2 and 3.3 in the revised manuscript to describe this issue.

25 *Section 2.1.2*

*NO<sub>x</sub> was measured by a gas-phase chemiluminescence detection analyzer EC9841, coupled with a hot molybdenum converter.*

*Section 3.3*

30 *It is worth noting that the observed NO<sub>x</sub> were detected by a chemiluminescence analyzer coupled with a hot molybdenum converter (MoO), and this method was recognized to cause higher NO<sub>2</sub> concentration due to the positive interference of other nitrogen-containing components (NO<sub>z</sub>, e.g. PAN, HNO<sub>3</sub> and HONO). Xu et al. (2013) found that the uncertainty caused by the MoO converter was much smaller at urban sites (less than 10%) than that at suburban and background sites (more than 30%). In the GZ basin, to evaluate the uncertainty, we estimated the ratio of NO<sub>z</sub>/(NO<sub>x</sub>+NO<sub>z</sub>) from the model. The calculated results indicated that the NO<sub>z</sub> accounted for 11% of the total NO<sub>x</sub>+NO<sub>z</sub> at urban Xi'an during the no-raining period. We noted the uncertainty in our NO<sub>x</sub> measurements, but thought this would not crucially impact the model-measurement comparison.* 35

40

(3) Sect. 2.2.1: It should be made clear here that only 6 samples total were collected.

Thanks for the suggestion. We modified the text in Section 2.1.1 in the revised manuscript as follows.

*We selected six field sites in the Qinling Mountains (Fig. 1b, the triangles) and*

collected one ambient air sample at each site on 6<sup>th</sup> – 7<sup>th</sup> August 2011 under sunny weather conditions (details are presented in Table 1).

(4) Page 7, Line 5: This mechanism seems outdated given the recent leaps in understanding of isoprene chemistry. Of particular concern is treatment of alkyl nitrates (which can be temporary or permanent NO<sub>x</sub> sinks, depending on the mechanism) and the general assumed fate of Isoprene RO<sub>2</sub> radicals. The authors should consider either 1) justifying why this does not impact their results, or 2) assessing potential uncertainties arising from use of an outdated mechanism.

Thank you for pointing this out. Isoprene is the most important NMVOC (Guenther et al., 2006), affecting tropospheric O<sub>3</sub>, OH and aerosols in complex ways. Being highly reactive, isoprene has a lifetime of only ~3h at OH concentration of  $1 \times 10^6 \text{ mol cm}^{-3}$ . In the daytime, isoprene reacts with OH radical to form hydroperoxy radicals (ISOPO<sub>2</sub>). Subsequently, in the presence of NO<sub>x</sub>, ISOPO<sub>2</sub> reacts with NO leading to the production of hydroxynitrates (ISOPN) by a minor branch, which sequester NO<sub>x</sub> and therefore regulate O<sub>3</sub> formation locally. With the presence of a double bond, ISOPNs are highly reactive with respect to OH and O<sub>3</sub>. The products from ISOPN oxidation might either release NO<sub>x</sub> or generate secondary organic nitrates. Some of the secondary organic nitrates products are found to be considerably longer-lived than ISOPN and can serve as temporary reservoirs for NO<sub>x</sub> (Paulot et al., 2009a, b; 2012). Besides daytime oxidation, nighttime oxidation of isoprene by NO<sub>3</sub> contributes significantly to the budget of organic nitrates (Horowitz et al., 2007; von Kuhlmann et al., 2004; Xie et al., 2013).

Many laboratory, field observations and simulation studies (e.g. von Kuhlmann et al., 2004; Perring et al., 2009; Paulot et al., 2009a) agreed that tropospheric O<sub>3</sub> production was highly sensitive to the ISOPN yield (4.4% to 15%) and NO<sub>x</sub> recycling efficiency (3% to 50%). However, large uncertainties still remain in estimating these parameters. Horowitz et al. (2007) found a 4% ISOPN yield best captured the alkyl and multifunctional nitrates measured by aircraft, and unreasonably high ISOPN yield (18%) would let ISOPN to be a terminal sink for NO<sub>x</sub> (Hudman et al., 2009).

Some recent modeling studies evaluated the sensitivity of tropospheric O<sub>3</sub> to isoprene oxidation chemistry. Xie et al. (2013) incorporated recent advances in isoprene oxidation chemistry in CMAQ (including a more explicit representation of isoprene nitrate formation from OH/NO and NO<sub>3</sub> pathways as well as modification to the isoprene chemistry under low-NO<sub>x</sub> conditions) and found the model can capture observations by assuming the ISOPN yield of 6% to 12%. O<sub>3</sub> increased by 2 ppbv in the eastern U.S as a result of these updates. Also focusing on the eastern U.S., Mao et al. (2013) implemented an updated isoprene chemistry mechanism in GEOS-Chem and found O<sub>3</sub> increased by 3-5 ppbv.

We added the discussion above in Section 2.2 to state the uncertainty.

*We noted some advances in isoprene nitrate chemistry in recent year. A number of laboratory, field observation and simulation studies (e.g. Paulot et al., 2009a, b, 2012; Horowitz et al., 2007; Hudman et al., 2009) highlighted the importance of the yield of isoprene nitrate (4-15%), the NO<sub>x</sub> recycling efficiency (3-50%) and the representation of secondary nitrate photochemistry for simulation of tropical O<sub>3</sub>. They all agreed there were still large uncertainties in isoprene nitrate chemistry. Some recent*

5 modeling studies evaluated the sensitivity of tropospheric O<sub>3</sub> to isoprene oxidation chemistry. Xie et al. (2013) incorporated recent advances in isoprene oxidation chemistry in CMAQ and found the model can capture observations by assuming the isoprene nitrate yield of 6% to 12%. Simulated O<sub>3</sub> increased by 2 ppbv in the eastern U.S. as a result of these updates. Also focusing on the eastern U.S., Mao et al. (2013) implemented an updated isoprene chemistry mechanism in GEOS-Chem and found O<sub>3</sub> increased by 3-5 ppbv.

10 (5) Page 9, Line 5: “Brute force” does not, to the reviewer’s knowledge, refer a specific method of source characterization. Please refine or define.

Thank you for the comment. We defined the “Brute force” method in Section 2.4 in the revised manuscript as follows.

15 *The approach referred to as the “brute-force” method (sensitivity analysis used to measure the model output response to emission changes) is traditionally used in air quality model to identify source contributions from specific non-reactive species in a linear process....*

20 (6) Page 10, Line 22: Does the model use assimilated meteorology? If so and the model is being nudged with observations, this agreement may not be especially remarkable.

We appreciate the reviewer's concern. We did not include assimilated meteorology in this study.

25 (7) Sect. 3.2: There are only 6 observations here, and looking at Table 1 the model seems to have little skill in capturing the variability of those. It would be worthwhile to point this out and justify why that’s not a big deal for the present analysis.

30 We agree that our model didn’t well capture the variability of the observed BVOCs. We think it is mainly because that, in our 3×3 km grid, different terpene emitters are not homogeneously distributed in the grid and the point measurements are influenced by the microenvironment and meteorology (Zare et al., 2012; Kota et al., 2015).

35 However, in this study, our goal is estimating the biogenic effect on O<sub>3</sub> formation in an urban area 50 km away from the foothills. We pay more attention to the magnitude of BVOC concentration at the regional scale, instead of capturing the hourly- and microenvironment-scale variability in either the observation or the simulation. Therefore, we think it is reasonable to compare the averaged simulations of 6 grids with the measurements.

We modified the text in Section 3.2 of the revised manuscript to state the uncertainty and to describe the above rationale.

40 *In general, different terpene emitters are not homogeneously distributed in a kilometer-scale grid and the point measurements are influenced by the microenvironment and meteorology (Zare et al., 2012; Kota et al., 2015). However, in this study, our goal is to estimate the biogenic effects on urban O<sub>3</sub> 50 km away from the*

foothills, which requests more concern on the regional scale VOC level, rather than the microenvironment-scale variability in either the observation or the simulation. Thus, we compared the average of VOC measurements with model simulations to validate whether the calculated results were reasonable. The isoprene mean concentration simulated in the six grids (corresponding to the time of observations) was 1.4 ppb, which is close to the observed average value of 1.3 ppb at the six sampling sites. Monoterpenes performed quite similarly, simulated 0.22 ppb comparing with observed 0.21 ppb. We also analyzed the temporal variation of simulated biogenic VOC during the whole simulation period and found the sub-month variability was relatively small (the standard deviation < 25%). The evaluation indicates that biogenic VOCs simulations reasonably agreed with the observations in the Qinling Mountains, on average, which provides a basis for us to further evaluate biogenic effects on O<sub>3</sub>.

(8) Page 11, Line 30: “agree well” seems optimistic. Looking at Fig. 4, 5 of the 11 non-rain days show model measurement disagreement by more than a factor of 2, and it is not evident that the model captures observed day-to-day variability.

Thank you for pointing this out. We modified text in Section 3.3 in the revised manuscript to make our statements more precise.

The model predicted PM<sub>2.5</sub> concentration to be 94.6±28.2 μg m<sup>-3</sup>, slightly lower (NMB=-12%) than measured 107 μg m<sup>-3</sup> averaged for the no-raining period, but didn't perform well in capturing temporal correlation (r=0.17). The simulated PM<sub>2.5</sub> showed the similar compositions to the observation (Fig. S1b and S1c)....

(9) Page 13, Line 7: Is this JNO<sub>2</sub> calculation based on model output? Please clarify.

Yes, the J(NO<sub>2</sub>) is calculated by model output “photor\_no2”. We added the text in Section 4.1 in the revised manuscript as follows.

Figure 6 shows the changes of J(NO<sub>2</sub>) (calculated by model track output photor\_no2) with the participation of PM averaged for urban Xi'an.

(10) Page 13, Line 15: presumably, aerosol lifetime is also longer than NO<sub>x</sub>.

Thank you for the suggestion. We added this in Section 4.1 in the revised manuscript.

The spatial distribution of high-values of PM<sub>2.5</sub> was similar to that of NO<sub>x</sub>, but covered a wider area mostly in the downwind region of urban Xi'an, which is expected due to longer lifetime of aerosols compared with NO<sub>x</sub> and the time required for secondary aerosol formation, thus further dispersion.

(11) Page 15, Line 18: Calling this the “original state” is confusing. If it is the anthropogenic simulation, just refer to it as such.

Thank you for the comment. We modified the text in Section 4.3 in the revised manuscript to make our statements clearer.

*To make it more specific, we started the discussion from the result of ANTH simulation without the biogenic sources (Table 2). Figure 10b shows the O<sub>3</sub> production regime in the ANTH simulation.*

5 (12) Page 16, Line 10: It is not clear that the discussion of the “actual” contributions adds much to the message of the paper. Indeed, it is a bit confusing because “pure” and “actual” have similar connotations, and because it convolutes several of the separated contributions.

10 Thank you for the comment. We deleted the discussion of the actual contributions to avoid ambiguity.

15 (13) Sect. 4.3: The lack of a “synergistic” contribution to PM is noteworthy and may deserve a few more sentences of discussion, especially given its importance for air quality. In other regions (e.g. the SE US), there seem to be relatively strong links between anthro/bio emissions and PM.

Thank you for the suggestion. We added the text in Section 4.3 in the revised manuscript to discuss this issue.

20 *It is worth noting that the biogenic contribution to PM<sub>2.5</sub> is not obvious (less than 3%) in GZ basin, which might be different from some other regions (e.g. Fu et al., 2012; Li et al., 2013). The main reasons are that 1) organic matter, the most important biogenic PM<sub>2.5</sub> component, only accounted for 14-16% of PM<sub>2.5</sub> in GZ basin in August; 2) Undeniably, uncertainties still exist in organic matter simulations in the model.*

25 (14) Table 2: Not sure this is necessary for the story. Could be moved to supplement. Your call.

Thanks for the suggestion. We moved this table to the supplement of the revised manuscript.

(15) Page 12, Line 14: discuss

30 Thanks. Corrected


(16) Page 16, Line 20: delete “and discussion”

Thanks. Corrected

35 (17) Fig. 12: suggest modifying colorbar to better separate NO<sub>x</sub> vs VOC controlled regions (e.g. blue-whitered gradient) and mentioning the limits for each regime in the caption.

Thank you for the suggestion. We modified the colorbar of Fig. 10 in the revised manuscript.





**Figure 10.** The monthly mean ratio of  $\text{H}_2\text{O}_2/\text{HNO}_3$  during the daily  $\text{O}_3$  peak time (14:00-18:00 local time) in August 2011 in the (a) Base simulation and (b) the simulation without biogenic sources.

5

## Reviewer #2

(1) My major concern with the work comes from the model uncertainty analysis, particularly in the simulation of VOCs and in emission inputs. In my opinion, six off-line VOC measurement points in space and time are not sufficient to validate the model accuracy for the following ozone source apportionment analysis. In addition, an agreement in Aug. 6 & 7 would not necessary indicate an agreement in the rest of month August. The current state of Section 3.2 is insufficient. More material is needed to show and quantify the model uncertainty in VOC estimation for the area under study. The uncertainty in emission input is needed as it is one of the main sources of model uncertainties.

Thanks for the comment. We added discussions about uncertainties of emission inventory and biogenic VOC simulations.

### (1) Biogenic VOC

We agree that only six off-line measurement points are not sufficient to describe the spatial and temporal variability of BVOCs. However, in this study, our goal is estimating the biogenic effect on  $\text{O}_3$  formation in urban Xi'an. We pay more attention to the magnitude of BVOC concentration at the regional scale, instead of capturing the hourly- and microenvironment-scale variability in either the observation or the simulation.

To verify the representativeness of BVOC observation (in Aug. 6 & 7), we analyzed the magnitude and temporal variability of simulated BVOC in the Qinling Mountains for the whole simulation period. The results indicated that the mean BVOC (isoprene + monoterperens) concentration is 1.73 ppb, close to the observed value (1.51 ppb) during Aug 6 & 7, and the sub-month variability (characterized by the standard deviation of daily mean simulation (averaged for 08:00-16:00 local time)) were <25%, indicating that the sub-month variability was relatively small.

We modified the text in the Section 3.2 of the revised manuscript to state the

uncertainty.

*In general, different terpene emitters are not homogeneously distributed in a kilometer-scale grid and the point measurements are influenced by the microenvironment and meteorology (Zare et al., 2012; Kota et al., 2015). However, in this study, our goal is to estimate the biogenic effects on urban O<sub>3</sub> 50 km away from the foothills, which requests more concern on the regional scale VOC level, rather than the microenvironment-scale variability in either the observation or the simulation. Thus, we compared the average of VOC measurements with model simulations to validate whether the calculated results were reasonable. The isoprene mean concentration simulated in the six grids (corresponding to the time of observations) was 1.4 ppb, which is close to the observed average value of 1.3 ppb at the six sampling sites. Monoterpenes performed quite similarly, simulated 0.22 ppb comparing with observed 0.21 ppb. We also analyzed the temporal variation of simulated biogenic VOC during the whole simulation period and found the sub-month variability was relatively small (the standard deviation < 25%). The evaluation indicates that biogenic VOCs simulations reasonably agreed with the observations in the Qinling Mountains, on average, which provides a basis for us to further evaluate biogenic effects on O<sub>3</sub>.*

## (2) Emission inventory

Anthropogenic emissions were obtained from the Multi-resolution Emission Inventory for China (Li et al., 2017). Emission estimates from bottom-up inventories are uncertain due to lack of complete knowledge of human activities and emission from different sources. The overall uncertainties of the bottom-up SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, VOC, and PM<sub>2.5</sub> emission estimates are ±31%, ±37%, ±153%, ±78% and ±133%, respectively, with 95% confidence intervals (CI) (Zhang et al., 2009; Lei et al., 2011; Zhao et al., 2011; Lu et al., 2011; Kurokawa et al., 2013). We updated Table S1 and added text in Section 2.3 of the revised manuscript to describe this issue.

*The emission estimates and uncertainties of VOCs, SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub>, and PM<sub>2.5</sub> in the domain during the simulation period are summarized in Table S1.*

(2) Another minor comment is with the application of brute-force comparison method in assessing the contributions of different emission sources to ozone concentration. The brute-force method has inherent disadvantages when applied to secondary species such as ozone and PM due to the non-linearity in responses. A critical question I would expect the authors to discuss in the manuscript is about the difference between actual and pure contribution from anthropogenic and biogenic.

The sensitivity analysis (brute-force method) used to measure the model output response to emission changes, is traditionally used in air quality model to identify source contributions from non-reactive species in a linear process. However, as the reviewer mentioned, it is not proper to secondary species such as O<sub>3</sub> and PM, because secondary species are generated nonlinearly, and interactions between different sources cannot be ignored. The difference between the actual and pure contribution from anthropogenic or biogenic source is the synergistic effect between the two sources. In this work, we combined the brute-force method and Factor Separation Technique (FST) (section 2.4) to discuss the importance of synergistic effect on O<sub>3</sub> formation. We



modified the text in the Section 2.4 in the revised manuscript to make our statement clearer.

5 *O<sub>3</sub> is formed by complicated nonlinear reactions of anthropogenic and biogenic precursors (NO<sub>x</sub> and VOCs) in the presence of sunlight. The approach referred to as the “brute-force” method (sensitivity analysis used to measure the model output response to emission changes) is traditionally used in air quality model to identify source contributions from specific non-reactive species in a linear process, but it cannot straightforward apply to secondary species due to the non-linearity in responses. In practice, the actual impact of one factor in a nonlinear process in the presence of others can be separated into 1) pure impact from the factor and 2) interactional impacts from all those factors. In this study, we adopted the factor separation approach (FSA) (Stein and Alpert, 1993) to decompose the pure contribution of a factor from its interaction with other factors.*

15 (3) In Fig. 7, is the black line based on theoretical calculation or model simulation? It seems to me that the smooth black line is based on the theoretical calculation for clear sky condition. If it was true, then I doubt if the decrease (red line) includes both the cloud and aerosol effects.

20 The black line is based on model simulated output “photor\_no2”. We modified the text in Section 4.1 in the revised manuscript to make this issue clear.

*Figure 6 shows the changes of J(NO<sub>2</sub>) (calculated by model track output photor\_no2) under the participation of PM averaged for urban Xi’an.*


25 (4) In Section 3.2, is the simulated isoprene/monoterpene mean concentration the mean over the one-month simulation periods or Aug. 6-7th or the mean of six data points corresponding to the time and location of the six measurements?

30 The simulated isoprene and monoterpenes concentrations in Section 3.2 were the mean of six data points corresponding to the time and location of the six measurements. We modified the text in Section 3.2 in the revised manuscript to make the statement clearer.

*The isoprene mean concentration simulated in the six grids (corresponding to the time of observations) was 1.4 ppb, which is close to the observed average value of 1.3 ppb at the six sampling sites.*

35 (5) Why modeled wind speed is discontinuous in Fig. 3c?

We appreciate the reviewer's concern. Figure 3c is presented as a “feather picture”, in which arrow direction indicates wind direction and length of arrow indicates wind speed. We give a line graph as follow to show wind speed more clearly.



## Reference

- 5 Fu, T.M., Cao, J.J., Zhang, X.Y., Lee, S.C., Zhang, Q., Han, Y.M., Qu, W.J., Han, Z., Zhang, R., Wang, Y.X., Chen, D., and Henze, D. K.: Carbonaceous aerosols in China: top-down constraints on primary sources and estimation of secondary contribution, *Atmos. Chem. Phys.*, 12, 2725-2746, doi:10.5194/acp-12-2725-2012, 2012.
- 10 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, doi:10.5194/acp-6-3181-2006, 2006.
- 15 Horowitz, L. W., Fiore, A.M., Milly, G.P., Cohen, R.C., Perring, A., Wooldridge, P.J., Hess, P.G., Emmons, L.K., and Lamarque J.F.: Observational constraints on the chemistry of isoprene nitrates over the eastern United States, *J. Geophys. Res.*, 112, D12S08, doi:10.1029/2006JD007747, 2007.
- 20 Hudman, R.C., Murray, L.T., Jacob, D.J., Turquety, S., Wu, S., Millet, D.B., Avery, M., Goldstein, A.H., and Holloway, J.: North American influence on tropospheric ozone and the effects of recent emission reductions: Constraints from ICARTT observations, *J. Geophys. Res.*, 114, D07302, doi:10.1029/2008JD010126, 2009.
- Kota, S. H., Schade, G., Estes, M., Boyer, D., and Ying, Q.: Evaluation of MEGAN predicted biogenic isoprene emissions at urban locations in Southeast Texas, *Atmospheric Environment*, 110, 54-64, 10.1016/j.atmosenv.2015.03.027, 2015.
- 25 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058, doi:10.5194/acp-13-11019-2013, 2013.
- 30 Lei, Y., Zhang, Q., He, K. B., and Streets, D. G.: Primary anthropogenic aerosol emission trends for China, 1990-2005, *Atmos. Chem. Phys.*, 11, 931-954, doi:10.5194/acp-11-931-2011, 2011.

- Li, M., Zhang, Q., Kurokawa, J.I., Woo, J.H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D.G., Carmichael, G.R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963, doi:10.5194/acp-17-935-2017, 2017.
- Li, N., Fu, T.-M., Cao, J., Lee, S., Huang, X.-F., He, L.-Y., Ho, K.-F., Fu, J. S., and Lam, Y.-F.: Sources of secondary organic aerosols in the Pearl River Delta region in fall: Contributions from the aqueous reactive uptake of dicarbonyls, *Atmos. Environ.*, 76, 200-207, doi:10.1016/j.atmosenv.2012.12.005, 2013.
- Lu, Z., Zhang, Q., and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996-2010, *Atmos. Chem. Phys.*, 11, 9839-9864, doi:10.5194/acp-11-9839-2011, 2011.
- Mao, J., Paulot, F., Jacob, D. J., Cohen, R.C., Crouse, J.D., Wennberg, P.O., Keller, C.A., Hudman, R.C., Barkley, M.P., and Horowitz, L.W.: Ozone and organic nitrates over the eastern United States: Sensitivity to isoprene chemistry, *J. Geophys. Res.*, 118, 11256-11268, doi:10.1002/jgrd.50817, 2013.
- Paulot, F., Crouse, J.D., Kjaergaard, H.G., Kroll, J.H., Seinfeld, J.H., and Wennberg, P.O.: Isoprene photooxidation: New insights into the production of acids and organic nitrates, *Atmos. Chem. Phys.*, 9(4), 1479–1501, 2009a.
- Paulot, F., Crouse, J.D., Kjaergaard, H.G., Kurten, A., St Clair, J.M., Seinfeld, J.H., and Wennberg, P.O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325(5941), 730–733, 2009b.
- Paulot, F., Henze, D.K., and Wennberg, P.O.: Impact of the isoprene photochemical cascade on tropical ozone, *Atmos. Chem. Phys.*, 12(3), 1307–1325, 2012.
- Perring, A.E., Wisthaler, A., Graus, M., Wooldridge, P.J., Lockwood, A.L., Mielke, L.H., Shepson, P.B., Hansel, A., and Cohen, R.C.: A product study of the isoprene + NO<sub>3</sub> reaction, *Atmos. Chem. Phys.*, 9(14), 4945–4956, 2009.
- von Kuhlmann, R., Lawrence, M.G., Poschl, U., and Crutzen, P.J.: Sensitivities in global scale modeling of isoprene, *Atmos. Chem. Phys.*, 4, 1–17, 2004.
- Xie, Y., Paulot, F., Carter, W.P.L., Nolte, C.G., Luecken, D.J., Hutzell, W.T., Wennberg, P.O., Cohen, R.C., and Pinder R.W.: Understanding the impact of recent advances in isoprene photooxidation on simulations of regional air quality, *Atmos. Chem. Phys.*, 13(16), 8439–8455, 2013.
- Xu, Z., Wang, T., Xue, L. K., Louie, P. K. K., Luk, C. W. Y., Gao, J., Wang, S. L., Chai, F. H., and Wang, W. X.: Evaluating the uncertainties of thermal catalytic conversion in measuring atmospheric nitrogen dioxide at four differently polluted sites in China, *Atmos. Environ.*, 76, 221-226, doi: 10.1016/j.atmosenv.2012.09.043, 2013.
- Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131-5153, doi:10.5194/acp-9-5131-2009, 2009.
- Zare, A., Christensen, J. H., Irannejad, P., and Brandt, J.: Evaluation of two isoprene

emission models for use in a long-range air pollution model, *Atmos Chem Phys*, 12, 7399-7412, 10.5194/acp-12-7399-2012, 2012.

- 5 Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B., and Hao, J.: Quantifying the uncertainties of a bottom-up emission inventory of anthropogenic atmospheric pollutants in China, *Atmos. Chem. Phys.*, 11, 2295-2308, doi:10.5194/acp-11-2295-2011, 2011.

**Impacts of Biogenic and Anthropogenic Emissions on Summertime  
Ozone Formation in the Guanzhong Basin, China**

5

Nan Li <sup>1,2</sup>, Qingyang He <sup>2,3</sup>, Jim Greenberg <sup>4</sup>, Alex Guenther <sup>5</sup>, Junji Cao <sup>2,6</sup>, Jun Wang <sup>7</sup>, Hong Liao <sup>1</sup>, **Qiyuan Wang** <sup>2</sup>, Qiang Zhang <sup>8</sup>

<sup>1</sup>Jiangsu Key Laboratory of Atmospheric Environment Monitoring and Pollution Control, Jiangsu Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing, 210044, China

<sup>2</sup>Key Lab of Aerosol Chemistry & Physics, SKLLQG, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, 710061, China

15 <sup>3</sup>Nanjing Star-jelly Environmental Consultants Co., Ltd, Nanjing, 210013, China

<sup>4</sup>National Center for Atmospheric Research, CO, USA

<sup>5</sup>Department of Earth System Science, University of California, Irvine, Irvine CA 92697-3100, USA

<sup>6</sup>Institute of Global Environmental Change, Xi'an Jiaotong University, Xi'an, 710049, China

<sup>7</sup>Center of Global and Regional Environmental Research, Department of Chemical and Biochemical Engineering, University of Iowa, Iowa City, Iowa, USA

20 <sup>8</sup>Department of Earth System Science, Tsinghua University, Beijing, 100084, China

*Correspondence to: Nan Li, [linan@nuist.edu.cn](mailto:linan@nuist.edu.cn)*

## Abstract

This study is the first attempt to understand the synergistic impact of anthropogenic and biogenic emissions on summertime ozone (O<sub>3</sub>) formation in the Guanzhong (GZ) basin where Xi'an, the oldest and the most populous city (with a population of 9 million) in the northwest China, is located. Month-long (August 2011) WRF-Chem simulations with different sensitivity experiments were conducted and compared with near-surface measurements. Biogenic volatile organic compounds (VOCs) concentrations were characterized from 6 surface sites among the Qinling Mountains, and urban air composition was measured in the Xi'an city at a tower 100 m above the surface. The WRF-Chem control experiment reasonably reproduced the magnitudes and variations of observed O<sub>3</sub>, VOCs, NO<sub>x</sub>, PM<sub>2.5</sub> and meteorological parameters, with normalized mean biases for each parameter within ±21%. Subsequent analysis employed the factor separation approach (FSA) to quantitatively disentangle the pure and synergistic impacts of anthropogenic and/or biogenic sources on summertime O<sub>3</sub> formation. The impact of anthropogenic sources alone was found to be dominant for O<sub>3</sub> formation. Although anthropogenic particles reduced NO<sub>2</sub> photolysis by up to 60%, the anthropogenic sources contributed 19.1 ppb O<sub>3</sub> formation on average for urban Xi'an. The abundant biogenic VOCs from the nearby forests promoted O<sub>3</sub> formation in urban areas by interacting with the anthropogenic NO<sub>x</sub>. The calculated synergistic contribution (from both biogenic and anthropogenic sources) was up to 14.4 ppb in urban Xi'an, peaking in the afternoon. Our study reveals that the synergistic impact of individual source contributions to O<sub>3</sub> formation should be considered in the formation of air pollution control strategies, especially for big cities in the vicinity of forests.

**Keywords:** ozone, emission, biogenic volatile organic compounds, WRF-Chem



## 1. Introduction

Elevated ozone (O<sub>3</sub>) levels in China has drawn increasing attention in recent years (e.g. Xue et al., 2014; Hu et al., 2016; Wang T. et al., 2016; Wang M. et al., 2016). O<sub>3</sub>, a secondary pollutant, is mainly formed by complex photochemical reactions of nitrogen oxides (NO<sub>x</sub> = NO + NO<sub>2</sub>) and volatile organic compounds (VOCs). High O<sub>3</sub> concentration at ground level is harmful to human health and ecosystems (WHO, 2005; Feng et al., 2015; Brauer et al., 2016). O<sub>3</sub> in the troposphere is an important greenhouse gas that has the third-highest radiative forcing after CO<sub>2</sub> and CH<sub>4</sub> (Stevenson et al., 2013; IPCC, 2013). In addition, O<sub>3</sub> is the primary source of the hydroxyl radical (OH), which has a major influence on the oxidizing capacity of the atmosphere and thus impacts the oxidation chemistry of secondary pollutants (e.g. sulfate and secondary organic aerosol) (Ehhalt et al., 2000; Rohrer et al., 2006). In recent years, the surface O<sub>3</sub> level has been increasing in most Chinese cities. For instance, in the highly urbanized areas in China, maximum 8h O<sub>3</sub> concentration increased by 19% (16.9 ppb) from 2015 to 2017 in the Beijing-Tianji-Hebei (BTH) region, the growth trend was 14% (11.3 ppb) for the Pearl River Delta (PRD) region and 12% (10.5 ppb) for the Yangtze River Delta (YRD) region (<http://datacenter.mep.gov.cn>). In the future, the pollution trend is likely to worsen due to potential changes of climate and emissions (Wang et al., 2013; Liu et al., 2013; Zhu et al., 2016, 2017).

On the global scale, VOC emissions from natural vegetation is estimated to be one order greater than that from anthropogenic activities, in spite of large uncertainties in different studies (Guenther et al., 2006; Wu et al., 2007, 2008; Jiang et al., 2013; Zhu et al., 2017). In addition, biogenic VOCs (e.g. isoprene) are highly reactive, reacting more efficiently with OH than most anthropogenic VOC species (Carlo et al., 2004). Previous studies have demonstrated the significant impacts of biogenic VOCs on surface O<sub>3</sub> formation under strong solar radiation, high temperature and NO<sub>x</sub> level (e.g. Fiore et al., 2005, 2011; Wang et al., 2008; Curci et al., 2009; Geng et al., 2011; Strong et al., 2013; Squire et al., 2014; Lee et al., 2014; Zhang et al., 2017). In China, biogenic VOC emissions are estimated to be 17-44 TgC yr<sup>-1</sup> (Guenther et al., 1995; Lin et al., 2008; Fu et al., 2012, 2104; Li et al., 2014), and are concentrated in the warm summer season. Current studies report that biogenic VOCs contribute to surface O<sub>3</sub> concentrations in China (Geng et al., 2011; Qu et al., 2013, 2014) and play an important role in intercontinental transport of O<sub>3</sub> (Zhu et al., 2017).

The impacts of biogenic VOCs on O<sub>3</sub> formation may vary in different regions and different seasons (Im et al., 2011; Strong et al., 2013; Wagner et al., 2014; Lee et al., 2014). Qu et al. (2013) employed the RAQM model to examine the influence of biogenic emissions on daily maximum surface O<sub>3</sub> concentration in China. Their calculations showed that in general the impact from biogenic sources on O<sub>3</sub> were more obvious in South China than in North China, but the O<sub>3</sub> increments in different regions didn't follow the same seasonality. Geng et al. (2011) used the WRF-Chem model to evaluate the effect of biogenic emissions on O<sub>3</sub> production in Shanghai in summer, and suggested that the carbonyls produced by the continuous oxidation of isoprene have important

impacts on O<sub>3</sub> level in the city. In addition, some studies suggest that biogenic emissions may increase due to global warming and land-use change, and the impact on O<sub>3</sub> formation could be more significant in the future (Lin et al., 2008; Fiore et al., 2011; Liu et al., 2013; Fu et al., 2015). Fiore et al. (2011) pointed out that the potential increases in biogenic isoprene in North American (NA) could offset the regional and intercontinental surface O<sub>3</sub> decreases produced by controls on NA anthropogenic emissions during warm seasons.

The Guanzhong (GZ) basin is the most developed region in northwestern China. In the past few years, air pollution has grown up to be a severe issue in the GZ basin (Wang et al., 2012; Xue et al., 2017), due to its specific basin topography and abundant anthropogenic emissions (Li et al., 2017). According to the data from national environmental monitoring stations in the GZ basin, 43% of days in 2013-2017 have AQI > 100 (e.g., unhealthy air quality category), and in summer O<sub>3</sub> was regarded as the primary pollutant in 70% of polluted cases. In this study, we employed a regional chemical model WRF-Chem to simulate O<sub>3</sub> concentration in the GZ basin for summer 2011. Our aim is the source apportionment of urban O<sub>3</sub> formation in this city surrounded by forests, specifically quantifying the individual and synergistic contributions of anthropogenic and biogenic sources. The paper is organized as follows. We first describe the sampling campaign, the chemical model and the emission data used for driving the model (Section 2). We then evaluate the model performance by comparing the observed urban air quality and biogenic VOCs with the simulated results (Section 3). Finally, we analyze the sensitivity of summertime O<sub>3</sub> formation to biogenic and anthropogenic sources (Section 4).

## 2. Methodology

### 2.1 Sampling sites and descriptions

The study was conducted in Xi'an, one of the oldest cities in the world. Xi'an is the most populous city in the northwest of China with a population of greater than 9 million. It is located in the heartland of the GZ basin between the Qinling Mountains and the Loess Plateau (Fig. 1). The topographical features result in air pollutants often being trapped in the valley with limited dispersion. The city borders the northern foot of the Qinling Mountains, only around 50 kilometers from downtown to the foothills. The Qinling Mountains are an east-west mountain range of 1600 kilometers in length and 300 kilometers in width and are regarded as a natural boundary between Northern and Southern China. Climate, and even culture are significantly different from the north to the south. The Xi'an city has high temperatures and strong solar intensity in summer, making it an ideal location to assess the importance of biogenic contributions to urban air quality.

#### 2.1.1 Biogenic VOC measurements in the Qinling Mountains

We selected six field sites in the Qinling Mountains (Fig. 1b, the triangles) and collected one

ambient air sample at each site on 6<sup>th</sup> – 7<sup>th</sup> August 2011 under sunny weather conditions (details are presented in Table 1). Sampling was conducted between 9:30 am to 16:30 pm local time to target expected daily maximum isoprene concentrations. At each site, ambient air samples were pulled in parallel onto three cartridges filled with Tenax GR and Carbograph 5TD solid adsorbents using a mass flow controlled pump for 30 min. Samples were shipped to the lab at NCAR (Boulder CO, USA) for chemical analysis. Cartridges were desorbed using an Ultra<sup>TM</sup> TD auto sampler with a Unity thermal desorption system (MARKES International Series 2, Llantrisant, UK) interfaced with a temperature programmed Agilent 7890A series Gas Chromatograph with a 5975C Electron Impact Mass Spectrometer and a Flame Ionization Detector (GC-MS/FID, Agilent Technologies, Santa Clara, CA, USA). We used nitrogen as a carrier gas at the flow rate of 3 mL min<sup>-1</sup>. Isoprene and monoterpene identifications were based on the comparison of retention time of authentic standards and mass spectra in the National Institute of Standards and Technologies (NIST) databases. Quantifications were calculated using FID calibrated with a NIST traceable standard.

### 2.1.2 Air quality monitoring in urban Xi'an

We set up an urban air quality monitoring site (Fig. 1b) at the roof (107 m above ground) of the main building (108.984°E, 34.245°N) on the campus of Xi'an Jiaotong University to minimize the ground level influences of local emissions. The campus is in the southeastern part of the downtown surrounded by residential areas. We obtained reliable observations of the concentrations of O<sub>3</sub>, NO<sub>x</sub> and PM<sub>2.5</sub> during 15<sup>th</sup> – 30<sup>th</sup> August 2011. Gases were measured by Ecotech analyzers (Ecotech Pty Ltd, Australia). O<sub>3</sub> were measured by an UV photometric analyzer EC9801. NO<sub>x</sub> was measured by a gas-phase chemiluminescence detection analyzer EC9841, coupled with a hot molybdenum converter. We collected 24h PM<sub>2.5</sub> filter samples by a mini-volume sampler (Airmetrics, USA) at a flow rate of 5 L min<sup>-1</sup> using both 47 mm quartz-fiber (Whatman, Mid-dlesex, UK) and teflon-membrane (Gelman, Ann Arbor, MI) filters. We calculated the PM<sub>2.5</sub> mass concentrations gravimetrically by weighing the teflon-membrane filters pre and post-collection for at least 4 times using an electronic microbalance (MC5, Sartorius, Göttingen, Germany) with ±1 µg sensitivity under controlled conditions. EC and OC concentrations were analyzed based on a 0.5 cm<sup>2</sup> punch from the quartz-fiber filter following the IMPROVE\_A (Interagency Monitoring of Protected Visual Environments) thermal/optical reflectance (TOR) protocol (Chow et al., 2007) using a DRI model 2001 Carbon Analyzer. The concentrations of ions were quantified from a 10.8 cm<sup>2</sup> of the teflon-membrane filter by a Dionex DX-600 ion chromatography (Dionex Inc., Sunnyvale, CA, USA) (Zhang et al., 2014).

## 2.2 The WRF-Chem model

We employed the WRF-Chem model to study biogenic VOC emissions from the Qinling Mountains and their contributions to regional O<sub>3</sub> formation in urban Xi'an. WRF-Chem is a 3-D

online-coupled meteorology and chemistry model consisting of the components of emission, transport, chemical transformation, photolysis and radiation (Tie et al., 2003, Li et al., 2011), dry and wet deposition (Wesely, 1989), and aerosol interactions (replaced with CMAQ aerosol module, Binkowski and Roselle, 2003; Li et al., 2010). WRF is a non-hydrostatic mesoscale dynamical system with various options for physical parameterizations (Skamarock et al., 2008). The chemical modules were implemented into the WRF framework obeying the same schemes for the simultaneous simulations (Grell et al., 2005).

We adopted RADM2 (Regional Acid Deposition Model) as the gas phase chemical mechanism to predict O<sub>3</sub> formation. RADM2 is an aggregated species type using the reactivity based weighting scheme to adjust for lumping (Stockwell et al., 1990). The mechanism implemented in our WRF-Chem model covers 158 reactions among 36 species, containing the complete reaction paths for isoprene, monoterpenes and the relevant inorganic reactions. As an explicit species, isoprene chemistry is based on an updated CB4 gas-phase mechanism (Carter and Atkinson., 1996). We noted some advances in isoprene nitrate chemistry in recent year. A number of laboratory, filed observation and simulation studies (e.g. Paulot et al., 2009a, b, 2012; Horowitz et al., 2007; Hudman et al., 2009) highlighted the importance of the yield of isoprene nitrate (4-15%), the NO<sub>x</sub> recycling efficiency (3-50%) and the representation of secondary nitrate photochemistry for simulation of tropical O<sub>3</sub>. They all agreed there were still large uncertainties in isoprene nitrate chemistry. Some recent modeling studies evaluated the sensitivity of tropospheric O<sub>3</sub> to isoprene oxidation chemistry. Xie et al. (2013) incorporated recent advances in isoprene oxidation chemistry in CMAQ and found the model could capture observations by assuming the isoprene nitrate yield of 6% to 12%. Simulated O<sub>3</sub> increased by 2 ppbv in the eastern U.S. as a result of these updates. Also focusing on the eastern U.S., Mao et al. (2013) implemented an updated isoprene chemistry mechanism in GEOS-Chem and found O<sub>3</sub> increased by 3-5 ppbv.

The simulated domain (Fig. 1) is 600×600 km<sup>2</sup> centered on urban Xi'an with 3 km horizontal grid spacing. We set up 28 vertical layers from the surface up to 50 hPa with 7 layers below 1 km to assure a high near-ground vertical resolution. The National Centers for Environmental Prediction (NCEP) FNL Operational Global Analysis data provided the initial and boundary fields of meteorology. Initial and boundary conditions of chemistry were derived by a global chemical transport model (Model for Ozone and Related chemical Tracers, MOZART) (Emmons et al., 2010). We considered the first 7 days as spin-up period, and the study focused on 6<sup>th</sup> – 7<sup>th</sup> and 15<sup>th</sup> – 30<sup>th</sup> August 2011 because of the available field observation datasets (as described in Section 2.1).

### 2.3 Biogenic and anthropogenic emissions

Biogenic emissions were quantified by the widely used model MEGAN (Model of Emissions of Gas and Aerosols from Nature) (Guenther et al., 2006). MEGAN coupled into the WRF-Chem

model, referred to here as WRF-MEGAN2, provides on-line estimates of the net landscape-averaged biogenic emissions from terrestrial ecosystems into the above-canopy atmosphere. The on-line estimated emissions of isoprene, individual monoterpenes and other biogenic VOCs serve as the inputs for the further chemistry simulation. To drive MEGAN, we need the following inputs:  
5 emission factors (EFs), leaf area index (LAI), plant functional types (PFTs), as well as meteorology conditions. The meteorology was obtained from WRF simulations and the LAI and PFT data were extracted from MODIS (Tian et al., 2004). We adopted the canopy-scale emission factors of dominant species from Guenther et al. (2006).

Estimated the whole year 2011 by WRF-MEGAN2, the isoprene and monoterpene emissions  
10 in the Qinling Mountains were mostly concentrated in summer (71% and 58%, respectively). During our simulation period, the isoprene emission from the domain is  $157 \text{ Gg mon}^{-1}$ , accounting for ~80% of total biogenic VOC emissions (Table S1). The rest are monoterpenes and other biogenic VOCs (e.g. acetone and MBO (2-methyl-3-buten-2-ol)). Figures 2a and 2b show the spatial distributions of biogenic isoprene and monoterpenes emission fluxes during the simulation period, indicating the  
15 high emission zone of isoprene in the Qinling Mountains lying to the south of the Xi'an city.

The anthropogenic emissions were obtained from the Multi-resolution Emission Inventory for China (MEIC, Li et al., 2017) for the year of 2010, which was downscaled to a resolution of 3 km using locations of point sources and various spatial proxies (Geng et al., 2017). The upgraded highly resolved emission data were based on a collection of statistics and newly developed emission factors.  
20 The emission inventory used in our model includes all major anthropogenic sources, but excluded open biomass burning which occupies a low proportion in the GZ basin during our simulating period (estimated by Fire Inventory from NCAR, <https://www2.acom.ucar.edu/modeling/finn-fire-inventory-ncar>). The anthropogenic emission sources are composed of power, industry, residential, transportation, and agriculture. **The emission estimates and uncertainties of VOCs,  $\text{SO}_2$ ,  $\text{NO}_x$ ,  $\text{NH}_3$ , and  $\text{PM}_{2.5}$  in the domain during the simulation period are summarized in Table S1.** The estimated anthropogenic VOCs emissions are 72.2 Gg, contributing up to ~30% of total VOC emissions. Figures 2c and 2d present the spatial distributions of anthropogenic VOC and  $\text{NO}_x$  emissions in the simulation period. The highest emission intensity of anthropogenic VOCs and  $\text{NO}_x$  are in Xi'an city and the GZ basin due to the frequent vehicle and industrial activities of this area.

30

## 2.4 Factor Separation Technique

$\text{O}_3$  is formed by complicated nonlinear reactions of anthropogenic and biogenic precursors ( $\text{NO}_x$  and VOCs) in the presence of sunlight. **The approach referred to as the “brute-force” method (sensitivity analysis used to measure the model output response to emission changes) is traditionally  
35 used in air quality model to identify source contributions from specific non-reactive species in a linear process, but it cannot straightforward apply to secondary species due to the non-linearity in**

responses. In practice, the actual impact of one factor in a nonlinear process in the presence of others can be separated into 1) pure impact from the factor and 2) interactional impacts from all those factors. In this study, we adopted the factor separation approach (FSA) (Stein and Alpert, 1993) to decompose the pure contribution of a factor from its interaction with other factors.

5 We considered anthropogenic and biogenic sources as two interactional factors to influence the O<sub>3</sub> formation.  $f_{anth-bio}$ ,  $f_{anth}$ ,  $f_{bio}$  and  $f_0$  are the simulation results including both anthropogenic and biogenic sources, anthropogenic source only, biogenic source only, and neither, respectively. Pure contributions of anthropogenic and biogenic sources are expressed as Eq. (1) and (2), respectively:

$$f'_{anth} = f_{anth} - f_0 \quad (1)$$

10  $f'_{bio} = f_{bio} - f_0 \quad (2)$

The calculated result including both anthropogenic and biogenic sources should include both pure contributions of the two factors, the synergistic impact, and the impact of background transport (Eq. 3):

$$f_{anth-bio} = f'_{anth} + f'_{bio} + f'_{anth-bio} + f_0 \quad (3)$$

15 Thus, the synergistic effect between anthropogenic and biogenic sources is represented as:

$$\begin{aligned} f'_{anth-bio} &= f_{anth-bio} - f'_{anth} - f'_{bio} - f_0 \\ &= f_{anth-bio} - (f_{anth} - f_0) - (f_{bio} - f_0) - f_0 \\ &= f_{anth-bio} - f_{anth} - f_{bio} + f_0 \end{aligned} \quad (4)$$

20 Based on the FSA, we conducted four simulations, namely BASE, ANTH, BIO and NEITHER, to explore the pure and synergistic impacts of anthropogenic and/or biogenic sources on O<sub>3</sub> production in the GZ basin. Detailed simulation settings and the various contribution definitions are summarized in Table 2.

### 3. Observation data and model validation

#### 25 3.1 Meteorology

The specific topographical features of Xi'an make the meteorological conditions crucial for the accumulation and dispersion of urban pollutants. To validate our model performance in wind, temperature and relative humidity, we compared the hourly meteorological data (<http://www.meteomanz.com>) observed at the Jinghe site (108.58°E, 34.26°N, in the west of Xi'an) with model simulations. Figure 3c illustrates the observed and simulated near-surface wind speed and directions during 15<sup>th</sup> – 30<sup>th</sup> August 2011. The WRF-Chem model successfully captured the prevailing wind direction from north and northeast, consistent with the in-situ observations. It should be noted that in our simulation period the prevailing wind blew from south, which enhanced the transport of biogenic emissions from the Qinling Mountains to urban Xi'an. In addition, a continuous rainfall event during 18<sup>th</sup>-22<sup>nd</sup> August (green shadow in Fig. 3) was characterized by



lower temperature and near-saturated humidity.

We conducted the statistical verification of meteorological variables in Table 3, including the  $r$  (correlation coefficient), NMB (normalized mean bias) and RMSE (root mean square errors). Modeled meteorological variables were in good agreement with observations (Fig. 3a-c) with the  
5 NMB less than  $\pm 6\%$ .

### 3.2 Biogenic VOC concentrations in the Qinling Mountains

Samples from the Qinling Mountains show that the dominant VOC species was isoprene, and  $\alpha$ -pinene was the main constituent of monoterpenes (Table 1). The ratio of isoprene to monoterpenes  
10 varies considerably. In general, different terpene emitters are not homogeneously distributed in a kilometer-scale grid and the point measurements are influenced by the microenvironment and meteorology (Zare et al., 2012; Kota et al., 2015). However, in this study, our goal is to estimate the biogenic effects on urban  $O_3$  50 km away from the foothills, which requests more concern on the regional scale VOC level, rather than the microenvironment-scale variability in either the  
15 observation or the simulation. Thus, we compared the average of VOC measurements with model simulations to validate whether the calculated results were reasonable. The isoprene mean concentration simulated in the six grids (corresponding to the time of observations) was 1.4 ppb, which is close to the observed average value of 1.3 ppb at the six sampling sites. Monoterpenes performed quite similarly, simulated 0.22 ppb comparing with observed 0.21 ppb. We also analyzed  
20 the temporal variation of simulated biogenic VOC during the whole simulation period and found the sub-month variability was relatively small (the standard deviation  $< 25\%$ ). The evaluation indicates that biogenic VOCs simulations reasonably agreed with the observations in the Qinling Mountains, on average, which provides a basis for us to further evaluate biogenic effects on  $O_3$ .

### 25 3.3 Gaseous and particulate pollutants in urban Xi'an

The sampling campaign was organized in summer. Based on the gaseous and particulate pollutant observations, the daily mean  $PM_{2.5}$  concentration was  $90.0 \pm 53.5 \mu g m^{-3}$ , with 57% of days exceeding the WHO Interim target-1 (IT-1)  $75 \mu g m^{-3}$ . The daily mean  $NO_x$  concentrations were 25.8-63.2 ppb, with 40% of days exceeding the  
30 guideline 48.7 ppb ( $\approx 100 \mu g m^{-3}$ , GB 3095-2012). The maximum 8h  $O_3$  concentration was 3.5-95.6 ppb, with most of the values around the national first grade standard of 46.6 ppb ( $\approx 100 \mu g m^{-3}$ , GB 3095-2012). Summer in Xi'an is monitored as the least polluted season of the year, and the case we picked is regarded as a typical situation in summer Xi'an.

Figure 3 compares the simulated hourly O<sub>3</sub> and NO<sub>x</sub> concentrations with in-situ observations. During the rainy episode (the green shadow in Fig. 3), our model overestimated NO<sub>x</sub> concentration and underestimated O<sub>3</sub> concentration. The deviation can be explained by the failure to simulate the precipitation in the WRF model resulting in underestimates in wet deposition. So, we focused our analysis on the period excluding the rainy period. During the no-raining days, our model well reproduced the diurnal variations and magnitudes of O<sub>3</sub> and NO<sub>x</sub> concentrations. The calculated O<sub>3</sub> averaged for no-raining period was 38.7 ppb, ~20% higher than the observed value of 31.5 ppb. Our simulated O<sub>3</sub> also reproduced the temporal variation of the observed O<sub>3</sub> (r = 0.72). For NO<sub>x</sub>, the simulated hourly NO<sub>x</sub> averaged for the no-raining period was 46.6 ppb, close to the observed 47.0 ppb (NMB=-1%). **It is worth noting that the observed NO<sub>x</sub> were detected by a chemiluminescence analyzer coupled with a hot molybdenum converter (MoO), and this method was recognized to cause higher NO<sub>2</sub> concentration due to the positive interference of other nitrogen-containing components (NO<sub>z</sub>, e.g. PAN, HNO<sub>3</sub> and HONO). Xu et al. (2013) found that the uncertainty caused by the MoO converter was much smaller at urban sites (less than 10%) than that at suburban and background sites (more than 30%). In the GZ basin, to evaluate the uncertainty, we estimated the ratio of NO<sub>z</sub>/(NO<sub>x</sub>+NO<sub>z</sub>) from the model. The calculated results indicated that the NO<sub>z</sub> accounted for 11% of the total NO<sub>x</sub>+NO<sub>z</sub> at urban Xi'an during the no-raining period. We noted the uncertainty in our NO<sub>x</sub> measurements, but considered this would not crucially impact the model-measurement comparison.**

We analyzed PM<sub>2.5</sub> concentration and composition (sulfate, nitrate, ammonium, EC, organic matter) with the filter-base measurements. **The model predicted PM<sub>2.5</sub> concentration to be 94.6±28.2 μg m<sup>-3</sup>, slightly lower (NMB=-12%) than measured 107 μg m<sup>-3</sup> averaged for the no-raining period, but didn't perform well in capturing temporal correlation (r=0.17). The simulated PM<sub>2.5</sub> showed the similar compositions to the observation (Fig. S1b and S1c).** Sulfate is the dominant constituent of both simulated (32%) and observed (37%) PM<sub>2.5</sub>. High sulfate concentration was mainly attributed to the high SO<sub>2</sub> emission in the GZ basin as well as the humid weather conditions (Wang et al., 2014). The secondary constituent of observed PM<sub>2.5</sub> is organic matter, which accounted for 16% of the total observed PM<sub>2.5</sub>, close to the simulated result (14%). Secondary organic matter contributed half to total simulated organic matter, mainly due to the abundant precursor (i.e. VOCs) emissions and the high atmospheric oxidation capacity in summer.

#### 4. Impacts of biogenic and anthropogenic sources on O<sub>3</sub> formation

In this section, we analyze the results from the four simulations (BASE, ANTH, BIO and NEITHER, Table 2) to characterize the fate of O<sub>3</sub> and its precursors in the GZ basin and to quantify the pure and synergistic impacts of anthropogenic and/or biogenic sources on summertime O<sub>3</sub> formation.

#### 5 4.1 Base simulation of O<sub>3</sub>

Firstly, we discuss the spatial and temporal characteristics of the simulated O<sub>3</sub> and the precursors (VOCs and NO<sub>x</sub>) in the GZ basin in the BASE simulation.

Figure 4a shows spatial distribution of the simulated VOCs during the no-raining period, overlaid with the simulated wind vectors. The highest concentration (more than 50 ppb) was in urban Xi'an and its downwind region (the southwest of urban Xi'an), due to anthropogenic activities. In addition, another high-value area (~30 ppb) was found in the Qinling Mountains, which was probably due to biogenic sources. To better understand the composition of VOCs, we analyzed some typical individual VOC species. Figure S2a shows the spatial distribution of xylenes, representing anthropogenic VOCs, and Fig. S3a and S4a show isoprene and monoterpenes, representing biogenic VOCs. The anthropogenic xylenes were mainly distributed in the GZ basin, while the high biogenic isoprene and monoterpene concentrations were found over the Qinling Mountains. These results explain the spatial feature of total VOCs and the dominant sources. Detailed discussion of source apportionment is given in Section 4.2

The spatial distribution of NO<sub>x</sub> was slightly different (Fig. 5a). The highest concentrations of NO<sub>x</sub> were in the GZ basin (average of 11.1 ppb), especially in urban Xi'an (averaged of 30.1 ppb), while among the Qinling Mountains, NO<sub>x</sub> was low and dominated by biogenic sources.

PM, even though not directly involved in the formation pathways of O<sub>3</sub>, influences the chemical equilibrium indirectly. In the daytime, NO<sub>2</sub> photolysis frequency ( $J(\text{NO}_2)$ ) is determined by the solar radiation influenced by PM via scattering and absorption. **Figure 6 shows the changes of  $J(\text{NO}_2)$  (calculated by model track output photor\_no2) with the participation of PM averaged for urban Xi'an.**  $J(\text{NO}_2)$  was reduced by 40-60%, most significantly in morning and evening rush hours. In the night time, PM<sub>2.5</sub> can remove N<sub>2</sub>O<sub>5</sub> from the NO<sub>x</sub> cycle via heterogeneous reactions, as one of the major NO<sub>x</sub> sinks in the atmosphere (Xue et al., 2014). Figure S5a shows the spatial feature of PM<sub>2.5</sub>. The densest area was urban Xi'an (averaged for 102 μg m<sup>-3</sup>) followed by the western part of the GZ basin. **The spatial distribution of high-values of PM<sub>2.5</sub> was similar to that of NO<sub>x</sub>, but covered a wider area mostly in the downwind region of urban Xi'an, which is expected due to longer lifetime of aerosols compared with NO<sub>x</sub> and the time required for secondary aerosol formation, thus further dispersion.**

The typical diurnal variation of O<sub>3</sub> (Fig. 7b) demonstrates there are higher concentrations in the afternoon and lower at night. For better understanding of O<sub>3</sub> concentration characteristics and

source/sink mechanisms, we discussed two different time scales: 1) O<sub>3</sub> peak time (14:00-18:00) (Fig. 8) and 2) O<sub>3</sub> 24-hour average (Fig. 9). During the peak time, simulated near-surface O<sub>3</sub> was high in the GZ basin, with averaged concentration of 75 ppb. In the downwind region of high NO<sub>x</sub> and VOCs in the west of urban Xi'an, the concentration reached up to 110 ppb. We employed the ratio of H<sub>2</sub>O<sub>2</sub>/HNO<sub>3</sub> to investigate the chemistry regime of O<sub>3</sub> formation (Sillman 1995; Wang et al., 2017). If the ratio is greater than 0.5, the O<sub>3</sub> production regime is considered NO<sub>x</sub>-controlled, otherwise VOC-controlled if the ratio less than 0.3. The range between 0.3 and 0.5 is defined as the transition regime from NO<sub>x</sub>- to VOC-controlled, indicating the competition of both NO<sub>x</sub> and VOCs in O<sub>3</sub> production. Figure 10a shows the spatial distribution of the simulated H<sub>2</sub>O<sub>2</sub>/HNO<sub>3</sub> ratio during the O<sub>3</sub> peak time. The west and southeast of the GZ basin were right in the transition regime with a complicated O<sub>3</sub> production mechanism sensitive to both NO<sub>x</sub> and VOCs. Most of the rest of the simulation region was VOC-controlled, excluding the Yuncheng and Hejing cities in the neighboring Shanxi provinces.

On the 24h average scale, the spatial distribution of O<sub>3</sub> presented a different picture (Fig. 9). The original high-value area during the peak time in the GZ city cluster shifted to low-value region due to the consumption of O<sub>3</sub> by abundant NO<sub>x</sub> emissions. At night time, the titration effect of freshly emitted NO dominates, and the O<sub>3</sub> concentration tends to drop to a lower level. The high value of 24h averaged O<sub>3</sub> converged in the south and northwest outside of the GZ Basin. Those areas have elevated O<sub>3</sub> due to high daytime production, similar to the nearby zone of peak O<sub>3</sub>, but also have lower emissions of NO resulting in lower loss of O<sub>3</sub>.

#### 4.2 Pure impact of biogenic or anthropogenic sources

Using the FSA method, we evaluated the pure contribution of anthropogenic or biogenic sources to the summertime O<sub>3</sub> formation in the GZ basin. In the scenario of pure contribution of anthropogenic emissions, the VOC concentrations were mostly distributed over the GZ city cluster (8.0 ppb), especially in urban Xi'an (26.4 ppb) (Fig. 4c). In the scenario of pure contribution of biogenic emissions, the VOCs were widely dispersed over the Qinling Mountains (Fig. 4d), with a calculated 9.9 ppb for the GZ basin and 12.4 ppb for urban Xi'an. NO<sub>x</sub> concentration has the similar pattern as VOCs in the scenario of pure contribution of anthropogenic emissions, with averaged concentrations of 11.0 ppb for the GZ basin and 30.3 ppb for urban Xi'an (Fig. 5c). However, in the GZ basin and urban Xi'an, biogenic sources contributed less than 0.2 ppb to NO<sub>x</sub> concentration (Fig. 5d). In the scenario of pure contribution of anthropogenic emissions, PM<sub>2.5</sub> spread over a wider area (Fig. S5c), due to the time required for secondary aerosol formation. In the scenario of pure contribution of biogenic emissions, PM<sub>2.5</sub> was mostly distributed among the Qinling Mountains (Fig. S5d), but the concentration was lower by one order of magnitude (Table 4, Fig. S6).

In the scenario of pure contribution of anthropogenic emissions, daily peak O<sub>3</sub> accumulated in

the downwind region in the center and western GZ due to high VOCs and NO<sub>x</sub> concentrations. Daily peak O<sub>3</sub> concentrations reached 22.6 ppb for GZ and 19.1 ppb for urban Xi'an (Fig. 8c). Lower daily peak O<sub>3</sub> concentration was found outside of GZ basin where less anthropogenic VOCs and NO<sub>x</sub> was emitted. In contrast, daily peak O<sub>3</sub> was negligible (less than 3 ppb) in the scenario of pure contribution of biogenic emissions (Fig. 8d) due to the low NO<sub>x</sub> emissions. However, the distribution of 24h averaged O<sub>3</sub> was different from daily peak O<sub>3</sub>. 24h averaged O<sub>3</sub> concentration in the scenario of pure contribution of anthropogenic emissions was more diluted in the GZ city cluster than for surrounding areas (Fig. 9c). Due to the abundant NO emission and its titration effect on O<sub>3</sub>, the pure effect of anthropogenic sources was negative, calculated to be -2.2 ppb in urban Xi'an.

10

### 4.3 Synergistic impact of the interaction between biogenic and anthropogenic sources

The synergistic impact on O<sub>3</sub> formation includes the interactions between anthropogenic and biogenic sources. In other word, it reflects the potential production trend of either "O<sub>3</sub>-promoted" or "O<sub>3</sub>-suppressed" under the natural coexistence of all emission sources. In the cases of NO<sub>x</sub>, VOCs and PM<sub>2.5</sub>, the synergistic impacts contributed less than ±3% of total concentrations (Fig. 4b, 5b, S5b, S6, and Table 4). However, the synergistic impact on O<sub>3</sub> played a remarkable role showing positive impacts for both daily peak (Fig. 8b) and 24h averaged O<sub>3</sub> (Fig. 9b). It means that the mixed state of anthropogenic and biogenic sources potentially enhanced the O<sub>3</sub> production more than each single source. To make it more specific, we started the discussion from the result of ANTH simulation without the biogenic sources (Table 2). Figure 10b shows the O<sub>3</sub> production regime in the ANTH simulation. VOC-controlled O<sub>3</sub> production regime covered the west of urban Xi'an and the southeast of the GZ basin. In the rest of the GZ basin and the neighboring Shanxi provinces, the O<sub>3</sub> production was in the transition regime, controlled by both NO<sub>x</sub> and VOC. NO<sub>x</sub>-controlled O<sub>3</sub> production regime dominated the rest of the region. After we included biogenic VOC emissions in the simulation, the O<sub>3</sub> concentration was significantly enhanced in the VOC-controlled regions, and partly enhanced in the mix-controlled region. However, in the VOC-controlled region, the synergistic impact contributed little.

The synergistic impact is of great importance, approximately the same magnitude as the impact from pure contributions of anthropogenic sources. The synergistic impact contributed daily peak O<sub>3</sub> concentrations of 10.5 ppb for the GZ basin and 14.3 ppb for urban Xi'an, while the pure anthropogenic impact contributed 22.6 ppb for GZ basin and 19.1 ppb for urban Xi'an. However, the extent was ~50% smaller on the 24h averaged scale, but still increased O<sub>3</sub> concentration by 5.8 ppb for the GZ basin and 6.8 ppb for urban Xi'an. Figure 7 shows the diurnal variation of the observed and simulated O<sub>3</sub> concentration at Xi'an Jiaotong University, as well as the tested contributing components. Transport dominated O<sub>3</sub>, constantly contributing 30-40 ppb as background. The impact of pure anthropogenic sources was positive on O<sub>3</sub> production during 13:00-

35

19:00 but negative during the rest of the time, and the impact of pure biogenic sources was negligible. Synergistic impact of both anthropogenic and biogenic sources resulted in a positive contribution during 10:00-21:00, comparable to the impact of pure anthropogenic sources.

5 It is worth noting that the biogenic contribution to  $PM_{2.5}$  is not obvious (less than 3%) in GZ basin, which might be different from some other regions (e.g. Fu et al., 2012; Li et al., 2013). The main reasons are that 1) organic matter, the most important biogenic  $PM_{2.5}$  component, only accounted for 14-16% of  $PM_{2.5}$  in GZ basin in August; 2) Undeniably, uncertainties still exist in organic matter simulations in the model.

## 10 5. Conclusions

The GZ basin is a representative region in the northwest of China, suffering serious air pollution in recent years. Geographically, the GZ basin borders the northern foot of the Qinling Mountains. For this reason, in addition to the anthropogenic emissions from metropolitan areas, biogenic emissions are of great importance in the region, especially in warm season with active photochemistry. In this study, we used the WRF-Chem model to simulate  $O_3$  in the GZ basin and compared the results to near-surface measurements, with the aim of quantifying the pure and synergistic impacts of anthropogenic and/or biogenic sources on summertime  $O_3$  formation. The simulation was driven by the best currently available inventory of anthropogenic emissions and online calculated biogenic emissions. Near-surface measurements were captured from 6 surface sites among the Qinling Mountains for biogenic VOCs and one 100-m-high site in the Xi'an city for air quality ( $NO_x$ , VOCs,  $O_3$  and  $PM_{2.5}$ ).

Our model successfully reproduced the observed air quality and meteorological parameters. The biogenic VOCs simulation showed a reasonable agreement. Our model also well-reproduced the magnitudes and variations of  $O_3$ ,  $NO_x$  and  $PM_{2.5}$  concentrations excluding rainy days, with normalized mean bias less than  $\pm 21\%$ .

We further conducted three scenario simulations to explore the pure and synergistic impacts of anthropogenic and/or biogenic sources on  $O_3$  and the precursors, by using the factor separation approach (FSA). The results concluded that, for the precursors, pure impact of anthropogenic source contributed 99% of  $NO_x$ , 80% of  $PM_{2.5}$ , and 33% of VOCs in the GZ basin, and pure impact of biogenic source contributed 40% of VOCs but only 1-5% of  $PM_{2.5}$  and  $NO_x$ . Meanwhile, synergistic impacts from the combination of anthropogenic and biogenic sources did not bring significant changes on  $NO_x$ , VOCs and  $PM_{2.5}$  (less than  $\pm 4\%$ ). In the case of daily peak  $O_3$ , the pure impact of anthropogenic source remained the dominant contributor (19.1 ppb for urban Xi'an), even after anthropogenic particles reduced the  $NO_2$  photolysis by up to 60%. The abundant biogenic VOCs from the nearby forests promoted the  $O_3$  formation by interaction with anthropogenic  $NO_x$ , contributing 14.4 ppb to  $O_3$  in urban Xi'an. This synergistic impact presented a positive contribution



to O<sub>3</sub> production throughout the day and the positive effect was much more prominent during 12:00-19:00.

O<sub>3</sub> pollution in China has been raising increasing concerns in recent years. Some scientists hold the view that excessive concentration of PM<sub>2.5</sub> suppressed the formation of O<sub>3</sub> in the past, hiding the problem temporally. However, with the effective control of PM<sub>2.5</sub>, O<sub>3</sub> pollution is manifested. The phenomenon can also be demonstrated by the government control action during G20 summit (The Group of Twenty Finance Ministers and Central Bank Governors) in Hangzhou in 2016. The concentration of PM<sub>2.5</sub> was depressed sharply under the strict emission control, but O<sub>3</sub> concentration was even higher than usual. Better understanding of O<sub>3</sub> pollution sources/sinks and formation mechanisms in high PM<sub>2.5</sub> exposed area in China will benefit and guide the implementation of PM<sub>2.5</sub>/O<sub>3</sub> cooperative control. Our results suggest that, in big cities geographically close to forest, O<sub>3</sub> pollution can be enhanced by the synergistic impact from the combination of biogenic and anthropogenic sources. The synergistic contribution of each single source to O<sub>3</sub> formation cannot be neglected when making pollution control strategies.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (41705128), Opening Project of State Key Laboratory of Loess and Quaternary Geology (SKLLQG1709), and Opening Project of Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention (LAP<sup>3</sup>) (FDLAP17003).

## Reference

- 25 Binkowski, F.S., and Roselle, S.J.: Models-3 community multiscale air quality (CMAQ) model aerosol component-1. Model description, *J. Geophys. Res.*, 108, 4183, doi:10.1029/2001jd001409, 2003.
- Bossioli, E., Tombrou, M., Karali, A., Dandou, A., Paronis, D., and Sofiev, M.: Ozone production from the interaction of wildfire and biogenic emissions: a case study in Russia during spring 2006, *Atmos. Chem. Phys.*, 12, 7931-7953, doi:10.5194/acp-12-7931-2012, 2012.
- 30 Brauer, M., Freedman, G., Frostad, J., van Donkelaar, A., Martin, R.V., Dentener, F., van Dingenen, R., Estep, K., Amini, H., Apte, J.S., Balakrishnan, K., Barregard, L., Broday, D., Feigin, V., Ghosh, S., Hopke, P.K., Knibbs, L.D., Kokubo, Y., Liu, Y., Ma, S., Morawska, L., Texcalac Sangrador, J. L., Shaddick, G., Anderson, H. R., Vos, T., Forouzanfar, M. H., Burnett, R.T., and Cohen, A.: Ambient Air Pollution Exposure Estimation for the Global Burden of Disease 2013, *Environ. Sci. Technol.*, 50, 79-88, doi:10.1021/acs.est.5b03709, 2016.
- 35 Carlo, P.D., Brune, W.H., Martinez, M., Harder, H., Leshner, R., Ren, X., Thornberry, T., Carroll, M.

- A., Young, V., and Shepson, P. B.: Missing OH Reactivity in a Forest: Evidence for Unknown Reactive Biogenic VOCs, *Science*, 304, 722-725, doi: 10.1126/science.1094392, 2004.
- Carter W.P.L. and Atkinson R.: Development and evaluation of a detailed mechanism for the atmospheric reactions of isoprene and NO<sub>x</sub>. *Int. J. Chem. Kinet.*, 28, 497-530, 1996.
- 5 Chow, J.C., Watson, J.G., Chen, L.W.A., Chang, M.C.O., Robinson, N.F., Trimble, D., and Kohl, S.: The IMPROVE-A temperature protocol for thermal/optical carbon analysis: maintaining consistency with a long-term database, *J. Air. Waster. Manage.*, 57, 1014-1023, doi:10.3155/1047-3289.57.9.1014, 2007.
- 10 Curci, G., Beekmann, M., Vautard, R., Smiatek, G., Steinbrecher, R., Theloke, J., and Friedrich, R.: Modelling study of the impact of isoprene and terpene biogenic emissions on European ozone levels, *Atmos. Environ.*, 43, 1444-1455, doi:10.1016/j.atmosenv.2008.02.070, 2009.
- Ehhalt, D.H., and Rohrer, F.: Dependence of the OH concentration on solar UV, *J. Geophys. Res.*, 105, 3565-3571, 2000.
- 15 Emmons, L.K., Walters, S., Hess, P.G., Lamarque, J.F., Pfister, G.G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S.L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geosci. Model Dev.*, 3, 43-67, doi:10.5194/gmd-3-43-2010, 2010.
- 20 Feng, T., Bei, N., Huang, R.J., Cao, J., Zhang, Q., Zhou, W., Tie, X., Liu, S., Zhang, T., Su, X., Lei, W., Molina, L. T., and Li, G.: Summertime ozone formation in Xi'an and surrounding areas, China, *Atmos. Chem. Phys.*, 16, 4323-4342, doi:10.5194/acp-16-4323-2016, 2016.
- Feng, Z., Hu, E., Wang, X., Jiang, L., and Liu, X.: Ground-level O<sub>3</sub> pollution and its impacts on food crops in China: A review, *Environ. Pollut.*, 199, 42-48, doi:10.1016/j.envpol.2015.01.016, 2015.
- 25 Fiore, A.M., Horowitz, L.W., Purves, D.W., Levy, H., Evans, M.J., Wang, Y.X., Li, Q.B., and Yantosca, R.M.: Evaluating the contribution of changes in isoprene emissions to surface ozone trends over the eastern United States, *J. Geophys. Res.*, 110, doi:10.1029/2004jd005485, 2005.
- Fiore, A.M., Levy, H. II, and Jaffe, D.A.: North American isoprene influence on intercontinental ozone pollution, *Atmos. Chem. Phys.*, 11, 1697-1710, doi:10.5194/acp-11-1697-2011, 2011.
- 30 Fu, T.M., Cao, J.J., Zhang, X.Y., Lee, S.C., Zhang, Q., Han, Y.M., Qu, W.J., Han, Z., Zhang, R., Wang, Y.X., Chen, D., and Henze, D. K.: Carbonaceous aerosols in China: top-down constraints on primary sources and estimation of secondary contribution, *Atmos. Chem. Phys.*, 12, 2725-2746, doi:10.5194/acp-12-2725-2012, 2012.
- 35 Fu, T.-M., Zheng, Y., Paulot, F., Mao, J., and Yantosca, R.M.: Positive but variable sensitivity of August surface ozone to large-scale warming in the southeast United States, *Nat. Clim. Change*, 5, 454-458, doi:10.1038/nclimate2567, 2015.
- Fu, Y., and Liao, H.: Simulation of the interannual variations of biogenic emissions of volatile organic compounds in China: impacts on tropospheric ozone and secondary organic aerosol, *Atmos. Environ.*, 59, 170-185, doi:10.1016/j.atmosenv.2012.05.053, 2012.
- 40 Fu, Y., and Liao, H.: Impacts of land use and land cover changes on biogenic emissions of volatile organic compounds in China from the late 1980s to the mid-2000s: implications for tropospheric ozone and secondary organic aerosol, *Tellus B*, 66, doi:10.3402/tellusb.v66.24987, 2014.
- 45 Geng, F., Tie, X., Guenther, A., Li, G., Cao, J., and Harley, P.: Effect of isoprene emissions from major forests on ozone formation in the city of Shanghai, China, *Atmos. Chem. Phys.*, 11, doi:10449-10459, 10.5194/acp-11-10449-2011, 2011.
- Geng, G., Zhang, Q., Martin, R. V., Lin, J., Huo, H., Zheng, B., Wang, S., and He, K.: Impact of

spatial proxies on the representation of bottom-up emission inventories: A satellite-based analysis, *Atmos. Chem. Phys.*, 17, 4131-4145, doi:10.5194/acp-17-4131-2017, 2017.

- 5 Grell, G.A., Peckham, S.E., Schmitz, R., McKeen, S.A., Frost, G., Skamarock, W.C., and Eder, B.: Fully coupled "online" chemistry within the WRF model, *Atmos. Environ.*, 39, 6957-6975, doi:10.1016/j.atmosenv.2005.04.027, 2005.
- Guenther, A., Hewitt, C.N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W.A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: a global-model of natural volatile organic-compound emissions, *J. Geophys. Res.*, 100, 8873-8892, doi:10.1029/94jd02950, 1995.
- 10 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, doi:10.5194/acp-6-3181-2006, 2006.
- Horowitz, L. W., Fiore, A.M., Milly, G.P., Cohen, R.C., Perring, A., Wooldridge, P.J., Hess, P.G., Emmons, L.K., and Lamarque J.F.: Observational constraints on the chemistry of isoprene nitrates over the eastern United States, *J. Geophys. Res.*, 112, D12S08, doi:10.1029/2006JD007747, 2007.
- 15 Hu, J., Chen, J., Ying, Q., and Zhang, H.: One-Year Simulation of Ozone and Particulate Matter in China Using WRF/CMAQ Modeling System, *Atmos. Chem. Phys.*, 16, 10333-10350, doi:10.5194/acp-16-10333-2016, 2016.
- 20 Hudman, R.C., Murray, L.T., Jacob, D.J., Turquety, S., Wu, S., Millet, D.B., Avery, M., Goldstein, A.H., and Holloway, J.: North American influence on tropospheric ozone and the effects of recent emission reductions: Constraints from ICARTT observations, *J. Geophys. Res.*, 114, D07302, doi:10.1029/2008JD010126, 2009.
- Im, U., Poupkou, A., Incecik, S., Markakis, K., Kindap, T., Unal, A., Melas, D., Yenigun, O., Topcu, S., Odman, M.T., Tayanc, M., and Guler, M.: The impact of anthropogenic and biogenic emissions on surface ozone concentrations in Istanbul, *Sci. Total Environ.*, 409, 1255-1265, doi:10.1016/j.scitotenv.2010.12.026, 2011.
- 25 IPCC: Working group I contribution to the IPCC Fifth Assessment Report Climate Change 2013: the Physical Science Basis, Cambridge University Press, Cambridge, New York, 2013
- 30 Jiang, H., Liao, H., Pye, H.O.T., Wu, S., Mickley, L.J., Seinfeld, J.H., and Zhang, X.Y.: Projected effect of 2000-2050 changes in climate and emissions on aerosol levels in China and associated transboundary transport, *Atmos. Chem. Phys.*, 13, 7937-7960, doi:10.5194/acp-13-7937-2013, 2013.
- Kota, S. H., Schade, G., Estes, M., Boyer, D., and Ying, Q.: Evaluation of MEGAN predicted biogenic isoprene emissions at urban locations in Southeast Texas, *Atmospheric Environment*, 110, 54-64, doi:10.1016/j.atmosenv.2015.03.027, 2015.
- 35 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., Kawashima, K., and Akimoto, H.: Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission inventory in ASia (REAS) version 2, *Atmos. Chem. Phys.*, 13, 11019-11058, doi:10.5194/acp-13-11019-2013, 2013.
- 40 Lee, K.-Y., Kwak, K.-H., Ryu, Y.-H., Lee, S.-H., and Baik, J.-J.: Impacts of biogenic isoprene emission on ozone air quality in the Seoul metropolitan area, *Atmos. Environ.*, 96, 209-219, doi:10.1016/j.atmosenv.2014.07.036, 2014.
- Lei, Y., Zhang, Q., He, K. B., and Streets, D. G.: Primary anthropogenic aerosol emission trends for China, 1990-2005, *Atmos. Chem. Phys.*, 11, 931-954, doi:10.5194/acp-11-931-2011, 2011.
- 45 Li, G., Lei, W., Zavala, M., and Volkamer, R.: Impacts of HONO sources on the photochemistry in Mexico City during the MCMA-2006/MILAGO Campaign, *Atmos. Chem. Phys.*, 10, 14, 6551-6567, doi:10.5194/acp-10-6551-2010, 2010.


- Li, G., Bei, N., Tie, X., and Molina, L. T.: Aerosol effects on the photochemistry in Mexico City during MCMA-2006/MILAGRO campaign, *Atmos. Chem. Phys.*, 11, 5169-5182, doi:10.5194/acp-11-5169-2011, 2011.
- 5 Li, L.Y., and Xie, S. D.: Historical variations of biogenic volatile organic compound emission inventories in China, 1981-2003, *Atmos. Environ.*, 95, 185-196, doi:10.1016/j.atmosenv.2014.06.033, 2014.
- 10 Li, M., Zhang, Q., Kurokawa, J.I., Woo, J. H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D.G., Carmichael, G.R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935-963, doi:10.5194/acp-17-935-2017, 2017.
- 15 Li, N., Fu, T.-M., Cao, J., Lee, S., Huang, X.-F., He, L.-Y., Ho, K.-F., Fu, J. S., and Lam, Y.-F.: Sources of secondary organic aerosols in the Pearl River Delta region in fall: Contributions from the aqueous reactive uptake of dicarbonyls, *Atmos. Environ.*, 76, 200-207, doi:10.1016/j.atmosenv.2012.12.005, 2013.
- Lin, J.-T., Patten, K.O., Hayhoe, K., Liang, X.-Z., and Wuebbles, D.J.: Effects of future climate and biogenic emissions changes on surface ozone over the United States and China, *J. Appl. Meteorol. Clim.*, 47, 1888-1909, doi:10.1175/2007jamc1681.1, 2008.
- 20 Liu, Q., Lam, K.S., Jiang, F., Wang, T.J., Xie, M., Zhuang, B.L., and Jiang, X.Y.: A numerical study of the impact of climate and emission changes on surface ozone over South China in autumn time in 2000-2050, *Atmos. Environ.*, 76, 227-237, doi:10.1016/j.atmosenv.2013.01.030, 2013.
- 25 Mao, J., Paulot, F., Jacob, D. J., Cohen, R.C., Crounse, J.D., Wennberg, P.O., Keller, C.A., Hudman, R.C., Barkley, M.P., and Horowitz, L.W.: Ozone and organic nitrates over the eastern United States: Sensitivity to isoprene chemistry, *J. Geophys. Res.*, 118, 11256-11268, doi:10.1002/jgrd.50817, 2013.
- Paulot, F., Crounse, J.D., Kjaergaard, H.G., Kroll, J.H., Seinfeld, J.H., and Wennberg, P.O.: Isoprene photooxidation: New insights into the production of acids and organic nitrates, *Atmos. Chem. Phys.*, 9(4), 1479–1501, doi: 10.5194/acp-9-1479-2009, 2009a.
- 30 Paulot, F., Crounse, J.D., Kjaergaard, H.G., Kurten, A., St Clair, J.M., Seinfeld, J.H., and Wennberg, P.O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325(5941), 730–733, doi: 10.1126/science.1172910, 2009b.
- Paulot, F., Henze, D.K., and Wennberg, P.O.: Impact of the isoprene photochemical cascade on tropical ozone, *Atmos. Chem. Phys.*, 12(3), 1307–1325, doi: 10.5194/acp-12-1307-2012, 2012.
- 35 Qu, Y., An, J., and Li, J.: Synergistic impacts of anthropogenic and biogenic emissions on summer surface O<sub>3</sub> in East Asia, *J. Environ. Sci.*, 25, 520-530, doi:10.1016/s1001-0742(12)60069-2, 2013.
- Qu, Y., An, J., Li, J., Chen, Y., Li, Y., Liu, X., and Hu, M.: Effects of NO<sub>x</sub> and VOCs from five emission sources on summer surface O<sub>3</sub> over the Beijing-Tianjin-Hebei region, *Adv. Atmos. Sci.*, 31, 787-800, doi:10.1007/s00376-013-3132-x, 2014.
- 40 Rohrer, F., and Berresheim, H.: Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet radiation, *Nature*, 442, 184, doi:10.1038/nature04924, 2006.
- Sillman, S.: The use of NO<sub>y</sub>, H<sub>2</sub>O<sub>2</sub>, and HNO<sub>3</sub> as indicators for ozone-NO<sub>x</sub>-hydrocarbon sensitivity in urban locations, *J. Geophys. Res.*, 100, 14175-14188, doi:10.1029/94jd02953, 1995.
- 45 Skamarock, W. C., Klemp, J.B., Dudhia, J., Gill, D.O., Barker, D.M., Duda, M.G., Huang, X.Y., Wang, W., and Powers, J.G.: A Description of the Advanced Research WRF Version 3, Technical Report, National Center for Atmospheric Research, TN-475,+STR, 2008.
- Squire, O.J., Archibald, A.T., Abraham, N.L., Beerling, D.J., Hewitt, C.N., Lathiere, J., Pike, R.C.,

- Telford, P.J., and Pyle, J.A.: Influence of future climate and cropland expansion on isoprene emissions and tropospheric ozone, *Atmos. Chem. Phys.*, 14, 1011-1024, doi:10.5194/acp-14-1011-2014, 2014.
- 5 Stein, U., and Alpert, P.: Factor separation in numerical simulations, *J. Atmos. Sci.*, 50, 2107-2115, doi:10.1175/1520-0469(1993)050<2107:fsins>2.0.co;2, 1993.
- 10 Stevenson, D.S., Young, P.J., Naik, V., Lamarque, J.F., Shindell, D.T., Voulgarakis, A., Skeie, R.B., Dalsoren, S.B., Myhre, G., Bernsten, T.K., Folberth, G.A., Rumbold, S.T., Collins, W.J., MacKenzie, I.A., Doherty, R.M., Zeng, G., van Noije, T.P.C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D.A., Strode, S.A., Horowitz, L., Lee, Y.H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K.W., Wild, O., and Archibald, A.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmos. Chem. Phys.*, 13, 3063-3085, doi:10.5194/acp-13-3063-2013, 2013.
- 15 Stockwell, W.R., Middleton, P., Chang, J.S., and Tang, X.: The second generation regional acid deposition model chemical mechanism for regional air quality modeling, *J. Geophys. Res.*, 95, D10, 16343-16367, doi: 10.1029/1990JD00461, 1990.
- 20 Strong, J., Whyatt, J.D., Metcalfe, S.E., Derwent, R.G., and Hewitt, C.N.: Investigating the impacts of anthropogenic and biogenic VOC emissions and elevated temperatures during the 2003 ozone episode in the UK, *Atmos. Environ.*, 74, 393-401, doi:10.1016/j.atmosenv.2013.04.006, 2013.
- Tian, Y., Dickinson, R.E., Zhou, L., Myneni, R.B., Friedl, M., Schaaf, C.B., Carroll, M., Gao, F.: Land boundary conditions from MODIS data and consequences for the albedo of a climate model. *Geophys. Res. Lett.*, 31, L05504, doi: 10.1029/2003GL019104, 2004.
- 25 Tie, X.X., Madronich, S., Walters, S., Zhang, R.Y., Rasch, P., and Collins, W.: Effect of clouds on photolysis and oxidants in the troposphere, *J. Geophys. Res.*, 108, doi:10.1029/2003jd003659, 2003.
- Wagner, P., and Kuttler, W.: Biogenic and anthropogenic isoprene in the near-surface urban atmosphere-A case study in Essen, Germany, *Sci. Total Environ.*, 475, 104-115, doi:10.1016/j.scitotenv.2013.12.026, 2014.
- 30 Wang, D., Hu, J., Xu, Y., Lv, D., Xie, X., Kleeman, M., Xing, J., Zhang, H., and Ying, Q.: Source contributions to primary and secondary inorganic particulate matter during a severe wintertime PM<sub>2.5</sub> pollution episode in Xi'an, China, *Atmos. Environ.*, 97, 182-194, doi:10.1016/j.atmosenv.2014.08.020, 2014.
- 35 Wang, M., Sampson, P.D., Hu, J., Kleeman, M., Keller, J. P., Olives, C., Szpiro, A.A., Vedal, S., and Kaufman, J.D.: Combining Land-Use Regression and Chemical Transport Modeling in a Spatiotemporal Geostatistical Model for Ozone and PM<sub>2.5</sub>, *Environ. Sci. Technol.*, 50, 5111-5118, doi:10.1021/acs.est.5b06001, 2016.
- 40 Wang, Q., Han, Z., Wang, T., and Zhang, R.: Impacts of biogenic emissions of VOC and NO<sub>x</sub> on tropospheric ozone during summertime in eastern China, *Sci. Total Environ.*, 395, 41-49, doi:10.1016/j.scitotenv.2008.01.059, 2008.
- Wang, T., Xue, L., Brimblecombe, P., Yun, F.L., Li, L., and Zhang, L.: Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects, *Sci. Total Environ.*, 575, 1582-1596, doi:10.1016/j.scitotenv.2016.10.081, 2017.
- 45 Wang, X., Shen, Z., Cao, J., Zhang, L., Liu, L., Li, J., Liu, S., and Sun, Y.: Characteristics of surface ozone at an urban site of Xi'an in Northwest China, *J. Environ. Monitor.*, 14, 116-126, doi:10.1039/c1em10541h, 2012.
- Wang, Y., Shen, L., Wu, S., Mickley, L., He, J., and Hao, J.: Sensitivity of surface ozone over China to 2000-2050 global changes of climate and emissions, *Atmos. Environ.*, 75, 374-382, doi:10.1016/j.atmosenv.2013.04.045, 2013.


- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical-models, *Atmos. Environ.*, 23, 1293-1304, doi:10.1016/0004-6981(89)90153-4, 1989.
- World Health Organization (WHO): Air Quality Guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide. Global update, 2005.
- 5 Wu, S., Mickley, L.J., Jacob, D.J., Logan, J.A., Yantosca, R.M., and Rind, D.: Why are there large differences between models in global budgets of tropospheric ozone?, *J. Geophys. Res.*, 112, doi:10.1029/2006jd007801, 2007.
- 10 Wu, S., Mickley, L.J., Leibensperger, E.M., Jacob, D.J., Rind, D., and Streets, D.G.: Effects of 2000-2050 global change on ozone air quality in the United States, *J. Geophys. Res.*, 113, doi:10.1029/2007jd008917, 2008.
- 15 Xie, Y., Paulot, F., Carter, W.P.L., Nolte, C.G., Luecken, D.J., Hutzell, W.T., Wennberg, P.O., Cohen, R.C., and Pinder R.W.: Understanding the impact of recent advances in isoprene photooxidation on simulations of regional air quality, *Atmos. Chem. Phys.*, 13(16), 8439–8455, doi: 10.5194/acp-13-8439-2013, 2013.
- Xu, Z., Wang, T., Xue, L. K., Louie, P. K. K., Luk, C. W. Y., Gao, J., Wang, S. L., Chai, F. H., and Wang, W. X.: Evaluating the uncertainties of thermal catalytic conversion in measuring atmospheric nitrogen dioxide at four differently polluted sites in China, *Atmos. Environ.*, 76, 221-226, doi: 10.1016/j.atmosenv.2012.09.043, 2013.
- 20 Xue, L.K., Wang, T., Gao, J., Ding, A.J., Zhou, X.H., Blake, D.R., Wang, X.F., Saunders, S.M., Fan, S.J., Zuo, H.C., Zhang, Q.Z., and Wang, W.X.: Ground-level ozone in four Chinese cities: precursors, regional transport and heterogeneous processes, *Atmos. Chem. Phys.*, 14, 13175-13188, doi:10.5194/acp-14-13175-2014, 2014.
- 25 Xue, Y., Ho, S., Huang, Y., Li, B., Wang, L., Dai, W., Cao, J., and Lee, S.: Source apportionment of VOCs and their impacts on surface ozone in an industry city of Baoji, Northwestern China, *Sci. Rep.*, 7, 9979, doi:10.1038/s41598-017-10631-4, 2017.
- Zare, A., Christensen, J. H., Irannejad, P., and Brandt, J.: Evaluation of two isoprene emission models for use in a long-range air pollution model, *Atmos Chem Phys*, 12, 7399-7412, 10.5194/acp-12-7399-2012, 2012.
- 30 Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, *Atmos. Chem. Phys.*, 9, 5131-5153, doi:10.5194/acp-9-5131-2009, 2009.
- 35 Zhang, R., Cohan, A., Biazar, A.P., and Cohan, D.S.: Source apportionment of biogenic contributions to ozone formation over the United States, *Atmos. Environ.*, 164, 8-19, doi:10.1016/j.atmosenv.2017.05.044, 2017.
- Zhang, T., Cao, J.J., Chow, J.C., Shen, Z.X., Ho, K.F., Ho, S.S.H., Liu, S.X., Han, Y.M., Watson, J.G., and Wang, G.H.: Characterization and seasonal variations of levoglucosan in fine particulate matter in Xi'an, China, *J. Air Waste Manage.*, 64, 1317-1327, doi:10.1080/10962247.2014.944959, 2014.
- 40 Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B., and Hao, J.: Quantifying the uncertainties of a bottom-up emission inventory of anthropogenic atmospheric pollutants in China, *Atmos. Chem. Phys.*, 11, 2295-2308, doi:10.5194/acp-11-2295-2011, 2011.
- 45 Zhu, J., and Liao, H.: Future ozone air quality and radiative forcing over China owing to future changes in emissions under the Representative Concentration Pathways (RCPs), *J. Geophys. Res.*, 121, 1978-2001, doi:10.1002/2015JD023926, 2016.
- Zhu, J., Liao, H., Mao, Y., Yang, Y., and Jiang, H.: Interannual variation, decadal trend, and future change in ozone outflow from East Asia, *Atmos. Chem. Phys.*, 17, 1-37, doi:10.5194/acp-17-




3729-2017, 2017.



**Figure 1.** The simulation domain and the locations of six biogenic VOCs sites (triangles), one meteorological site (square) and one air quality site (snowflake). Underlain are land types from MODIS. The area of red line indicates the urban area of the Xi'an city. The area of blue line indicates the GZ basin.




**Figure 2.** Monthly mean emissions of (a) biogenic isoprene, (b) biogenic monoterpenes, (c) anthropogenic VOCs and (d) anthropogenic  $\text{NO}_x$  in the GZ basin and surrounding areas in August 2011.




**Figure 3.** Observed (black) and simulated (red) temporal patterns of temperature (a), relative humidity (b) and wind (c) at the Jinghe site and NO<sub>x</sub> (d) and O<sub>3</sub> (e) concentrations at Xi'an Jiaotong University during the period from 15<sup>th</sup> to 30<sup>th</sup> August 2011. The green shadow (18<sup>th</sup> -22<sup>nd</sup> August) indicates rainy days.







**Figure 4.** Spatial distributions of monthly mean concentrations of VOCs in August 2011. (a) is the result from the BASE simulation, overlaid with simulated wind vectors. (b)-(f) are simulated VOCs concentrations contributed from synergistic anthropogenic and biogenic, pure anthropogenic, pure biogenic, actual anthropogenic and actual biogenic sources, respectively.



**Figure 5.** Spatial distributions of monthly mean concentrations of  $\text{NO}_x$  in August 2011. (a) is the result from the BASE simulation, overlaid with simulated wind vectors. (b)-(f) are simulated  $\text{NO}_x$  concentrations contributed from synergistic anthropogenic and biogenic, pure anthropogenic, pure biogenic, actual anthropogenic and actual biogenic sources, respectively.




**Figure 6.** Diurnal variations of  $J(\text{NO}_2)$  (black) and the changes in  $J(\text{NO}_2)$  (red) averaged in urban Xi'an due to PM effects in August 2011.




**Figure 7.** Temporal patterns of the simulated O<sub>3</sub> concentrations and the tested contributing components during the period from 15<sup>th</sup> to 30<sup>th</sup> August 2011, excluding the rainy days (18<sup>th</sup> -22<sup>nd</sup> August). The orange shadow (14:00-18:00) indicates daily O<sub>3</sub> peak time.






**Figure 8.** Spatial distributions of monthly mean concentrations of daily peak O<sub>3</sub> in August 2011. (a) is the result from the BASE simulation, overlaid with simulated wind vectors. (b)-(f) are simulated daily peak O<sub>3</sub> concentrations contributed from synergistic anthropogenic and biogenic, pure anthropogenic, pure biogenic, actual anthropogenic and actual biogenic sources, respectively.



**Figure 9.** Spatial distributions of monthly mean concentrations of 24h averaged O<sub>3</sub> in August 2011. (a) is the result from the BASE simulation, overlaid with simulated wind vectors. (b)-(f) are simulated 24h averaged O<sub>3</sub> concentrations contributed from synergistic anthropogenic and biogenic, pure anthropogenic, pure biogenic, actual anthropogenic and actual biogenic sources, respectively.

5



**Figure 10.** The monthly mean ratio of  $H_2O_2/HNO_3$  during the daily  $O_3$  peak time (14:00-18:00 local time) in August 2011 in the (a) Base simulation and (b) the simulation without biogenic sources.

**Table 1.** Ambient biogenic VOCs observations in the Qinling Mountains during 6<sup>th</sup>– 7<sup>th</sup> August 2011

Site <sup>a</sup>	Date	Start Time <sup>b</sup>	Location	Isoprene (ppb)		Monoterpenes (ppb)		Dominant monoterpenes <sup>c</sup>
				Observation	Simulation	Observation	Simulation	
NBL	2011/8/6	10:20	33.78°E, 108.88°N	3.8	4.5	0.42	0.29	$\alpha$ -pinene
ZA	2011/8/6	12:44	33.40°E, 109.05°N	0.1	0.9	0.16	0.22	$\alpha$ -pinene, limonene, menthone
XY	2011/8/6	16:14	32.87°E, 109.40°N	1.0	0.3	0.24	0.13	$\alpha$ -pinene
AK	2011/8/7	09:45	32.71°E, 109.01°N	0.8	0.4	0.10	0.08	$\alpha$ -pinene
TJB	2011/8/7	13:00	32.99°E, 107.78°N	0.5	1.1	0.27	0.26	$\alpha$ -pinene
YS	2011/8/7	14:50	33.27°E, 107.73°N	1.6	0.9	0.04	0.32	$\alpha$ -pinene
Average				1.3	1.4	0.21	0.22	

<sup>a</sup> Site names: NBL (Niubeiliang), ZA (Zhenan), XY (Xunyang), AK (Ankang), TJB (Tangjiaba), YS (Youshui)

<sup>b</sup> The sampling duration is 30 minutes.

<sup>c</sup> Other monoterpenes detected (0.1% to 9%) include tricyclene,  $\alpha$ -thujene, camphene, sabinene, myrcene,  $\alpha$ -phellandrene,  $\Delta$ -carene, o-cymene,  $\beta$ -ocimene, cineole, isopulegol, isomenthone.

**Table 2.** Summary of different simulation settings and definitions of the various contributions from anthropogenic and/or biogenic sources.

Simulation	Simulation results	Anthropogenic emission	Biogenic emission
BASE	$f_{\text{anth-bio}}$	✓	✓
ANTH	$f_{\text{anth}}$	✓	✗
BIO	$f_{\text{bio}}$	✗	✓
NEITHER	$f_0$	✗	✗
Contribution			
	$f_{\text{anth-bio}} - f_{\text{bio}}$	Actual contribution of anthropogenic emissions	
	$f_{\text{anth-bio}} - f_{\text{anth}}$	Actual contribution of biogenic emissions	
	$f'_0 = f_0$	The contribution of background transport	
	$f'_{\text{anth}} = f_{\text{anth}} - f_0$	Pure contribution of anthropogenic emissions	
	$f'_{\text{bio}} = f_{\text{bio}} - f_0$	Pure contribution of biogenic emissions	
	$f'_{\text{anth-bio}} = f_{\text{anth-bio}} - (f_{\text{anth}} + f_{\text{bio}}) + f_0$	Synergistic contribution of anthropogenic and biogenic emissions	

**Table 3.** Statistics of meteorological and air quality variables over the GZ basin in August 2011<sup>a</sup>

	Mean		r <sup>d</sup>	NMB <sup>d</sup>	RMSE <sup>d</sup>
	Observation	Simulation			
<b>Meteorology<sup>b</sup></b>					
Wind speed (m s <sup>-1</sup> )	2.6	2.5	-	-6%	1.8
Temperature (°C)	25.1	24.2	0.86	4%	2.5
Relative humidity	73.6%	74.2%	0.72	1%	12%
<b>Air quality<sup>c</sup></b>					
NO <sub>x</sub> (ppb)	47.0	46.6	0.36	-1%	18.1
O <sub>3</sub> (ppb)	31.5	38.7	0.72	21%	8.1
PM <sub>2.5</sub> (µg m <sup>-3</sup> )	107	94.6	-	-12%	49.3

<sup>a</sup> Averaged for the period from 15<sup>th</sup> to 30<sup>th</sup> August 2011, excluding the rainy days.

<sup>b</sup> Meteorological data were obtained from the hourly surface measurements at Jinghe station (108.58°E, 34.26°N)

<sup>c</sup> Air quality data were measured at the roof (107 m above ground) of the main building (108.98°E, 34.25°N) on the campus of Xi'an Jiaotong University

<sup>d</sup> r: correlation coefficient; NMB: normalized mean bias; RMSE: root mean square errors.

**Table 4.** The various contribution components of the simulated O<sub>3</sub> (and the precursors) and PM<sub>2.5</sub> in August 2011

	NO <sub>x</sub> [ppb]	VOCs [ppb]	O <sub>3</sub> [ppb]		PM <sub>2.5</sub> [μg m <sup>-3</sup> ]
			daily peak <sup>a</sup>	24h	
<i>The GZ basin</i>					
Base	11.1	24.5	74.1	44.4	65.1
Pure ANTH	11.0	8.0	22.6	5.0	52.0
Pure BIO	0.1	9.9	2.0	1.1	3.3
Actual ANTH	10.6	7.7	33.0	10.8	53.5
Actual BIO	-0.3	9.6	12.5	7.0	4.9
Syn ANTH-BIO	-0.4	-0.3	10.5	5.8	1.5
<i>Urban Xi'an</i>					
Base	30.1	44.8	74.7	38.7	102
Pure ANTH	30.3	26.4	19.1	-2.2	88.7
Pure BIO	0.15	12.4	2.6	1.4	3.4
Actual ANTH	29.6	26.2	33.4	4.6	91.1
Actual BIO	-0.6	12.2	16.8	8.2	5.8
Syn ANTH-BIO	-0.7	-0.2	14.3	6.8	2.4

<sup>a</sup> Daily O<sub>3</sub> peak time is from 14:00 to 18:00 local time