

Supplement of:

Source apportionment of organic aerosol from two-year highly time-resolved measurements by an aerosol chemical speciation monitor in Beijing, China

Yele Sun^{1,2,3}, Weiqi Xu^{1,2}, Qi Zhang⁴, Qi Jiang^{1*}, Francesco Canonaco⁵, André S. H. Prévôt⁵, Pingqing Fu^{1,2,3}, Jie Li¹, John Jayne⁶, Douglas R. Worsnop⁶, and Zifa Wang^{1,2,3}

¹State Key Laboratory of Atmospheric Boundary Layer Physics and Atmospheric Chemistry, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

²University of Chinese Academy of Sciences, Beijing 100049, China

³Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

⁴Department of Environmental Toxicology, University of California, 1 Shields Ave., Davis, CA 95616, USA

⁵Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen PSI 5232, Switzerland

⁶Aerodyne Research, Inc., Billerica, MA 01821, USA

*Now at National Meteorological Centre, Beijing 100081, China

Correspondence to: Yele Sun (sunyele@mail.iap.ac.cn)




Figure S1: Mass spectral profiles of HOA and OOA from two-factor solution of PMF analysis of ACSM OA mass spectra in (a) S11, summer 2011, (a2) S12, summer 2012, (b1) F11, fall 2011, (b2) F12, fall 2012, (c1) W11, winter 2011, (c2) W12, winter 2012, (d1) Sp12, spring 2012, and (d2) Sp13, spring 2013.




Figure S2: Comparisons of monthly-average mass concentrations and mass fractions of POA and SOA between three different analysis, including seasonal PMF-ACSM and ME2-ACSM, and ME2-ACSM for the entire dataset. In addition, the results of SOA by subtracting LO-OOA and POA by adding LO-OOA are also shown.




Figure S3: Scatter plots of monthly average POA and SOA concentrations from ME2-ACSM and PMF-ACSM.




Figure S4: Comparisons of POA and SOA from ME2-ACSM in this study with those reported previously from PMF-ACSM in July – August 2011 (Sun et al., 2012), November 2011 – January 2012 (Sun et al., 2013), September 2012 (Jiang et al., 2013), January 2013 (Sun et al., 2014), and February – March 2013 (Jiang et al., 2015).




Figure S5: Correlation coefficients of five OA factors with other species, including Org, SO₄, NO₃, NH₄, Chl, black carbon (BC), NO, NO₂, NO_y, O₃, SO₂, CO, PM_{2.5}, light extinction, FOA, COA, BBOA, LO-OOA and MO-OOA in (a1) S11, summer 2011, (a2) S12, summer 2012, (b1) F11, fall 2011, (b2) F12, fall 2012, (c1) W11, winter 2011, (c2) W12, winter 2012, (d1) Sp12, spring 2012, and (d2) Sp13, spring 2013.




Figure S6: Average diurnal OA composition during four seasons.




Figure S7: RH/ T dependence of (a) wind speed and (b) FOA/COA for the entire study. The data are grouped into grids with increments of RH and T being 5% and 3 °C, respectively. Grid cells with the number of data points fewer than 10 are excluded.




Figure S8: Wind rose plots color coded by wind speed (m s^{-1}) during four seasons. The wind speed and wind direction were measured at 240 m, and the scales are set to the same during all seasons.

References:

- Jiang, Q., Sun, Y., Wang, Z., and Yin, Y.: Real-time online measurements of the inorganic and organic composition of haze fine particles with an Aerosol Chemical Speciation Monitor (ACSM) (in Chinese), *Chin. Sci. Bull. (Chin Ver)*, 58, 3818-3828, doi: 10.1360/972013-501, 2013.
- Jiang, Q., Sun, Y. L., Wang, Z., and Yin, Y.: Aerosol composition and sources during the Chinese Spring Festival: fireworks, secondary aerosol, and holiday effects, *Atmos. Chem. Phys.*, 15, 6023-6034, 10.5194/acp-15-6023-2015, 2015.
- Sun, Y. L., Wang, Z., Dong, H., Yang, T., Li, J., Pan, X., Chen, P., and Jayne, J. T.: Characterization of summer organic and inorganic aerosols in Beijing, China with an Aerosol Chemical Speciation Monitor, *Atmos. Environ.*, 51, 250-259, 10.1016/j.atmosenv.2012.01.013, 2012.
- Sun, Y. L., Wang, Z. F., Fu, P. Q., Yang, T., Jiang, Q., Dong, H. B., Li, J., and Jia, J. J.: Aerosol composition, sources and processes during wintertime in Beijing, China, *Atmos. Chem. Phys.*, 13, 4577-4592, 10.5194/acp-13-4577-2013, 2013.
- Sun, Y. L., Jiang, Q., Wang, Z., Fu, P., Li, J., Yang, T., and Yin, Y.: Investigation of the sources and evolution processes of severe haze pollution in Beijing in January 2013, *J. Geophys. Res.*, 119, 4380-4398, 10.1002/2014JD021641, 2014.