

Dear Co-editor,

Thanks for this review, which offers us the opportunity to improve and clarify our manuscript. We have corrected the latest version of the manuscript of **12th June 2018** and added some clarifications (Line numbers in this document refer to the previously reviewed version of the manuscript submitted on 12th June 2018). All corrections are commented below and included in the new version of the manuscript.

This review is from the Co-Editor.

This paper describes measurements of NH₃ and NO fluxes in Benin, West Africa and in June and July, 2016. The article has received one round of reviews. My primary concern is that the new version has not adequately addressed concerns around the NH₃ flux measurements. Data of soil fluxes in West Africa are rare and I hope more information can be provided on the measurements so the paper can be published. I have provided other comments as well.

1/The application of experimental results from Vaaitinen et al. (2013) is not yet convincing. The value for PFA of 13.9×10^{12} molecules/cm² was determined under constant conditions that do not represent the field sites: 10% sea-level atmospheric pressure, a constant temperature of 295K, constant humidity, and ambient NH₃ of ~8 ppm (not 8 ppb). Vaaitinen et al. (2013) themselves say the reported uptake values are minimums, rather than absolute quantities. Diurnal and day-to-day changes in these conditions could drive significant variability in air-chamber exchange, despite the low standard deviations reported in Vaaitinen et al. (2013). If the chambers were frequently washed, then maybe uptake is the dominant direction of the air-chamber exchange; however, a key conclusion is that soils are largely an NH₃ sink. For this reason, additional work is need to verify that the reported fluxes are between the air and soil rather than the air and chamber.

Authors' response :

The chambers were not exactly “washed”, but rather wiped with dry clean paper cloth. This clarification was added to line 173.

“The Teflon chamber was cleaned with a dry clean paper cloth at the beginning of each day of measurement, and during the day when the deposition of sand could potentially interfere with the measurements”

We propose to modify line 183 with the following text:

“Moreover, experimental tests with and without the Teflon chamber attached to the analyzer were made in ambient air to verify that deposition on the walls of the Teflon chamber is negligible. The results of this experiment are reported in Appendix A.”

We propose to add the following text in Appendix A, line 576:

“We ran a laboratory experiment to verify that deposition on the walls of the Teflon chamber is negligible.

Ambient air concentrations were measured by the analyzer, inside the room where the analyzer and the chamber were placed. Measurements of NH₃ concentrations were made in ambient air with and without the Teflon chamber attached to the analyzer. The Teflon chamber was placed on a Teflon frame, and they were sealed together with Teflon tape. Measurements of NH₃ concentrations with the Teflon chamber attached to the analyzer were followed by measurements without the chamber 30 to 60 minutes later. The two sets of measurements were made under similar conditions of temperature and humidity. Average values of NH₃ concentrations were calculated for 10 to 30

minutes before and after connecting the chamber. Average NH₃ concentrations during this time interval varied between 8 and 36 ppb, with a variation between 1.5 to 13% around the mean. The lowest NH₃ concentrations correspond to air samples previously passed through charcoal and desiccant cartridges (NO and NO₂ zero air). Measured NH₃ concentrations are reported in Table A1, along with temperature, humidity and the ratio between average concentration with and without the Teflon chamber attached to the analyzer..

This test was made at different times of the day on different days: air humidity varied between 46 and 54%, temperature varied between 25 to 29°C, while pressure varied between 1006 and 1008 hPa (not reported).

Results show negligible variation between concentrations of air reaching the analyzer via the chamber or going directly to the analyzer.

Temperature (°C)	Humidity (%)	[NH ₃] TC	[NH ₃] D	Ratio [NH ₃]TC / [NH ₃]D
25	54	26.4±0.4	25.9±0.5	1.02
27	49	9.0±0.8	8.8±1.2	1.12
29	46	35.2±2.4	36.0±1.4	0.98
28	46	25.6±0.7	24.0±1.0	1.07
27	45	26.2±0.9	27.4±0.9	0.96
28	46	23.4±0.3	24.1±0.4	0.97
26	50	19.2±0.6	18.4±0.8	1.04

Table A1: Measurements of NH₃ concentrations (ppb) through the chamber (TC) or directly (D) to the analyzer. “

 2/Referee 1 makes a valid point in comment 2 (below):

“The authors use the measured NO and NH₃ fluxes for a stepwise linear multiple regression analysis, upscaling to country-wide soil fluxes, and comparison with soil emission estimates from the GEOSChem model. These analyses give valuable information on the importance of soil NO and NH₃ exchange and our current knowledge about them. However, while the authors state that a process understanding of the NO and NH₃ fluxes is not within the scope of the presented study, in my opinion it is important to understand the underlying processes of the measured fluxes. For example, the estimated emissions from soil characteristics only poorly agree with the measured fluxes in some parts, which indicates that a more detailed process understanding is necessary.”

However, the authors have made no substantive changes to the manuscript in response. Or, if they have, they have not indicated this in the formal response. Please clarify.

Authors’ response :

As mentioned in the previous authors’ response of 12th June 2018, the authors agree that a better understanding of the underlying processes of NO and NH₃ exchanges is important. As explained in lines 420-425 our experiment set-up was chosen to give an estimate of soil fluxes at a large ecosystem scale, rather than reproducing the relationships between soil fluxes and meteorological variables, like soil temperature and soil moisture. Our experiment does not show the details of microbial and physical processes driving soil fluxes at a single point because measurements are

done at different locations every day, but aims to estimate the spatial variability of fluxes at the ecosystem scale.

We have re-written the two paragraphs starting from line 415 to clarify this:

« Due to this non linear character of the NO fluxes, no direct correlation was found between NO fluxes and environmental variables such as soil moisture or soil temperature taken individually. Moreover, soil temperature and soil moisture were not measured on the same soil parcel where the soil fluxes were measured, and the location of the soil flux measurements was not kept constant even for the same land cover type on the same measurement day. This measurement protocol was designed to give an estimate of soil fluxes at a larger ecosystem scale, rather than reproducing the relationships between soil fluxes and meteorological variables, like soil temperature and soil moisture.

A multiple linear regression analysis was performed between daily mean NO fluxes and the following variables: wind speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, upward longwave radiation and downward shortwave radiation. This regression gives $R^2=0.49$ (p-value=0.004), indicating a weak but existing relationship between those variables and NO soil emissions, while the regression was weak between NO fluxes and each individual variable. This correlation shows the influence of these environmental variables considered collectively on NO fluxes, highlighting the underlying mechanisms responsible for NO release to the atmosphere. Our experiment does not show the details of microbial and physical processes driving soil fluxes at a single point, because measurements are done at different locations every day, but aims to estimate the spatial variability of fluxes at the ecosystem scale.”

Other studies have been made to investigate the processes behind soil fluxes, see for example lines 45-48 in the Introduction.

Moreover, the comparison of measured and simulated NO soil emissions made in this study is done against GEOS-Chem which includes a process-based model for NO (Hudman et al., 2012). A better process understanding could generally be useful to improve soil emissions models, including the already advanced process-based model by Hudman et al., 2012 used in this study, but other issues should be considered, as the model uncertainty in reproducing weather and soil conditions, as well as the inevitable simplification of the actual highly variable land and vegetation cover introduced by the use of grid cell land cover and vegetation type in the model simulation.

We have modified lines 485-489 to better explain this concept :

« Simulated NO emissions are often higher than those recorded over the grassland areas, however, simulated NO emissions are often within the error bars of measurements (Fig. 6). The model uses land cover and vegetation types to simulate the highly variable land and vegetation cover of the observation site, for this reason we do not expect the model to reproduce the site-to-site variability of the measured soil fluxes, but to at least reproduce their average magnitude and behaviour. It appears that when the model is able to reproduce the length and the intensity of the rain events, NO emissions are especially well simulated, e.g. the model is able to reproduce the longest rain period (from 20 to 26 July 2016) and the decrease of emissions at the end of the measurement campaign. »

3/Rearding Referee 1's comment:

"The assumption that the concentration in the chamber is equal to the concentration leaving the chamber to the analyzer is questionable. Due to the low flow rate required for the practical use of the closed-dynamic chamber technique, the residence time within the chamber is substantial (17–18 min). As no active mixing (e.g. with fan) is used, the chamber geometry in relation to the positioning of the ambient air inlet and sample outlet is of importance."

Please modify the text so the reader does not need to read another paper to understand the fundamentals of the experiment. Consider this comment broadly and add a brief overview of all relevant information from Delon et al. (2017). You may refer readers to Delon et al. (2017) for additional detail, but provide the basics here.

Authors' response:

We have added the details of the chamber geometry at line 170:

"The external volume of the chamber was 40 cm × 20 cm × 20 cm. The useful volume was 18 × 38 × 18 cm³, due to the thickness of the Teflon walls".

The structure of the chamber and the positions of the inlet was already included in the latest manuscript, at line 170.

4/For the new text: "This is different from Delon et al., 2017, but fluxes were anyway superior to this value." Change the word superior to "greater than."

Authors' response

We thank the co-editor for this correction. That sentence was only used in the response to the referee and it is not included in the manuscript.

5/Regarding Referee 1's comment (a concern also raised by Referee 2):

"This assumption seems brave if it was not tested with a set of test experiments. Although the microbial activity is reduced due to the dry conditions, there is a chance that NH₃ volatilizes with the drying of the soil sample material."

I see you have added a reference to this phenomenon, but discussion for how this relates qualitatively and quantitatively to your results is still needed.

Authors' response:

The paragraph starting from line 247 was re-written to fully discuss this issue.

"Soil samples were collected with a cylinder of known volume (290 cm³) during the measurement campaign to analyze the biogeochemical characteristics of the site. Soil samples (0-5 cm) were taken for each land cover type where NO and NH₃ fluxes were measured. Fifteen samples were collected at the four different land cover types, three to four times during the campaign.

Samples were dried in ambient conditions (mean day-time temperature is approximately 26 °C, Kalthoff et al., 2017), and stored in the dark. After drying, the weight of the samples was measured to determine the bulk density (d_a = dry soil mass / total volume), which was found to be 1.24 ± 0.14 g cm⁻³. Assuming a density of soil particles (d_r) of 2.6 g cm⁻³, the Water Filled Pore Space (WFPS) is calculated with the following equation:

$$\text{WFPS} = \text{SM}/(1-d_a/d_r) \quad (5)$$

where SM is the soil moisture in %.

Soil samples were analyzed for the determination of texture, ammonium concentrations $[\text{NH}_4^+]$, C/N ratio, total C, total N, and pH at the GALYS Laboratoire (<http://www.galys-laboratoire.fr>, NF EN ISO/CEI 17025: 2005). The analyses were performed two months after sampling. We assume that the ammonium content in litter or soils is not modified by volatilization or chemical transformation during transport and storage, because of the very low soil moisture level in samples. Indeed, when collected, WFPS of the samples ranged between 6 and 14% (mean=8.5±3.5), and soil temperature between 35 and 38 °C (data obtained from the databases described in Brooks et al., 2017). Bai et al. (2013, and references therein) have found that significant changes in nitrification and net mineralization (influencing the ammonium content) may occur when soil temperature is raised up to 35°C (the optimum for nitrification), for optimal soil moisture conditions (WFPS =20%, Oswald et al., 2013). In the present study, soil temperatures when sampling were equal or above the optimum, and WFPS was below the optimum, reducing the nitrification efficiency and the change in ammonium content. Past studies have demonstrated that microbial activity and mineralization processes are inhibited in low soil moisture conditions, even when soil temperature is high (Bai et al., 2013, and references therein). Several authors have published results of ammonium concentrations measured in soils dried in ambient air. For example, Dick et al., (2006) collected top soil after the wet season in two sites in Senegal. The authors state that their soils were considered dry when collected and were air-dried in the mid-day sun immediately after collection. The protocol used in our study is identical. Other studies (Bai et al., 2010, Cassity-Duffrey et al., 2015, Vanlauwe et al., 2002) also published ammonium measurements made on air-dried soils from seasonally dry climates with comparable textures to the soil in Savé.”

6/New text: “Daily means of NO concentration vary from 1.28 to 5.40 ppb for all sites.” In Fig. 2 (and 4), NO concentrations appear to be equal 7–9 ppb on multiple days.

Authors’ response:

An error was found in the legend of the figures 2 to 5. The symbols for NO and NH₃ concentrations were inverted. This has been corrected. The NO concentration range is correct in the text.

7/As far as I can see, Figs. 2 and 4 are identical. Is this an error?

Authors’ response:

Yes that is an error. The authors apologize for this error and the wrong figure has been replaced by the correct one for the bare soil site (Figure 2). All legends have been corrected.

8/New text: “Average NH₃ concentration is 6.28 ± 3.90 ppb for bare soils.” Over bare soil, it appears the NH₃ concentration never exceeds ~3 ppb.

Authors’ response:

As mentioned above the legends for NH₃ and NO concentrations were inverted, and figure 2 for bare soils was not the right figure. This was corrected. The concentration ranges given in the text are correct.

9/Regarding Referee 2's comment:

"While I am sure location and resources had much to do with this, air-drying may result in large changes to ammonium concentrations. Additionally, significant changes in the amounts of ammonium can take place over prolonged storage at room temperature, even if soils are dried. It seems that the authors are aware of this issue and attempted to justify their method by citing a meta-analysis of warming experiments on N-cycle activity. (Bai et al. 2013). However, this meta-analysis found that warming and moisture reduction had no significant effect on mineralization (Bai et al, 2013: Table 1), indicating even in dried samples, pools of inorganic-N may change over time. To remedy this, the authors could have compared their ammonium concentrations to similar studies from this region; however, this was not included in the results/discussion."

Please expand the new discussion, which at just two sentences (line 255 and line 360), is insufficient.

Authors' response:

Please see authors' response number 5 above, in this document. The paragraph starting from line 247 was re-written to fully discuss this issue.

Some new references were added as asked by the reviewers, however measurements of ammonium content from the same region of this study are apparently not available in the literature.

The discussion on inorganic pool changes with temperature and moisture was included in the newly rewritten paragraph:

10/Other comments:

All plots are of poor visual quality, can the authors remake the plots with a program capable of higher resolution graphics and that allow the authors to improve the readability of the plots?

Authors' response:

Figures are provided as pdf for better resolution.

11/Abstract: I don't follow the logic of the first sentence.

Authors' response:

The first sentence was modified to:

"Biogenic fluxes from soil at a local and regional scale are crucial to study air pollution and climate. Here we present field measurements of soil fluxes of nitric oxide (NO) and ammonia (NH₃) observed over four different land cover types, i.e. bare soil, grassland, maize field and forest, at an inland rural site in Benin, West Africa, during the DACCIWA field campaign in June and July 2016. At the regional scale, urbanization and a massive growth in population in West Africa has been causing a strong increase in anthropogenic emissions. Anthropogenic pollutants are transported inland and northward from the mega cities located on the coast, where the reaction with biogenic emissions may lead to enhanced ozone production outside urban areas, as well as

secondary organic aerosol formation, with detrimental effects on humans, animals, natural vegetation and crops.”

12/Line 20: change aerosols to aerosol.

Authors' response:

Correction done

13/Abstract should be a single paragraph.

Authors' response:

The abstract was rewritten as a single paragraph.

Small errors have been corrected:

Line 156: “reference NO₂ air” becomes “reference NO air”.

Line 183: “chemical” becomes “photochemical”

Line 188: “increase” becomes “variation”

Line 350: “compared to Le Roux et al. (1995) and Lata et al. (2004). » becomes « compared to these studies »

Measurements of nitric oxide and ammonia soil fluxes from a wet savanna ecosystem site in West Africa during the DACCIWA field campaign.

5 Federica Pacifico¹, Claire Delon¹, Corinne Jambert¹, Pierre Durand¹, Eleanor Morris², Mat J. Evans²,
Fabienne Lohou¹, Solène Derrien¹, Venance H. E. Donnou³, Arnaud V. Houeto³, Irene Reinares
Martínez¹, Pierre-Etienne Brilouet¹

¹Laboratoire d'Aérodologie, University of Toulouse, CNRS, UPS, Toulouse, 31400, France

²Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, YO10 5DD, UK

10 ³Laboratoire de Physique du Rayonnement, Université d'Abomey-Calavi, Cotonou, 01 BP 526, Benin

15 *Correspondence to:* Claire Delon (claire.delon@aero.obs-mip.fr)

Abstract.

Biogenic fluxes from soil at a local and regional scale are crucial to study air pollution and climate. Here we present field measurements of soil fluxes of nitric oxide (NO) and ammonia (NH₃) observed over four different land cover types, i.e. bare soil, grassland, maize field and forest, at an inland rural site in Benin, West Africa, during the DACCIWA field campaign in June and July 2016. At the regional scale, urbanization and a massive growth in population in West Africa has been causing a strong increase in anthropogenic emissions. Anthropogenic pollutants are transported inland and northward from the mega cities located on the coast, where the reaction with biogenic emissions may lead to enhanced ozone production outside urban areas, as well as secondary organic aerosol formation, with detrimental effects on humans, animals, natural vegetation and crops. We observe NO fluxes up to 48.05 ngN m⁻² s⁻¹. NO fluxes averaged over all land cover types are 4.79 ± 5.59 ngN m⁻² s⁻¹, maximum soil emissions of NO are recorded over bare soil. NH₃ is dominated by deposition for all land cover types. NH₃ fluxes range between -6.59 and 4.96 ngN m⁻² s⁻¹. NH₃ fluxes averaged over all land cover types are -0.91 ± 1.27 ngN m⁻² s⁻¹ and maximum NH₃ deposition is measured over bare soil. The observations show high spatial variability even for the same soil type, same day and same meteorological conditions. We compare point daily average measurements of NO emissions recorded during the field campaign with those simulated by GEOS-Chem (Goddard Earth Observing System Chemistry Model) for the same site and find good agreement. In an attempt to quantify NO emissions at the regional and national scale, we also provide a tentative estimate of total NO emissions for the entire country of Benin for the month of July using two distinct methods: upscaling point measurements and using the GEOS-Chem model. The two methods give similar results:

20
25
30

1.17 ± 0.6 GgN/month and 1.44 GgN/month, respectively. Total NH₃ deposition estimated by upscaling point measurements for the month of July is 0.21 GgN/month.

35 It is important to correctly simulate biogenic fluxes from soil in atmospheric chemistry models at a local and regional scale to study air pollution and climate in an area of the world, West Africa, that has been subject to a strong increase in anthropogenic emissions due to a massive growth in population and urbanization. Anthropogenic pollutants are transported inland and northward from the mega-cities located on the coast, where the reaction with biogenic emissions may lead to enhanced ozone production outside urban areas, as well as secondary organic aerosols formation, with detrimental effects on
40 humans, animals, natural vegetation and crops.

Here we present field measurements of soil fluxes of nitric oxide (NO) and ammonia (NH₃) observed over four different land cover types, i.e. bare soil, grassland, maize field and forest, at an inland rural site in Benin, West Africa, during the DACCIWA field campaign in June and July 2016.

45 We observe NO fluxes up to 48.05 ngN m⁻² s⁻¹. NO fluxes averaged over all land cover types are 4.79 ± 5.59 ngN m⁻² s⁻¹; maximum soil emissions of NO are recorded over bare soil. NH₃ is dominated by deposition for all land cover types. NH₃ fluxes range between -6.59 and 4.96 ngN m⁻² s⁻¹. NH₃ fluxes averaged over all land cover types are -0.91 ± 1.27 ngN m⁻² s⁻¹ and maximum NH₃ deposition is measured over bare soil. The observations show high spatial variability even for the same
50 soil type, same day and same meteorological conditions.

We compare point daily average measurements of NO emissions recorded during the field campaign with those simulated by GEOS Chem (Goddard Earth Observing System Chemistry Model) for the same site and find good agreement. In an attempt to quantify NO emissions at the regional and national scale, we also provide a tentative estimate of total NO emissions for
55 the entire country of Benin for the month of July using two distinct methods: upscaling point measurements and using the GEOS Chem model. The two methods give similar results: 1.17 ± 0.6 GgN/month and 1.44 GgN/month, respectively. Total NH₃ deposition estimated by upscaling point measurements for the month of July is 0.21 GgN/month.

1 Introduction

Biogenic soil fluxes of nitric oxide (NO) and ammonia (NH₃) play an important role on tropospheric chemistry. Nitric Oxide emitted by soil influences the concentration of nitrogen oxides (NO_x) in the atmosphere, consequently modifying the rates of ozone (O₃) production, where O₃ is a pollutant, harmful to humans and plants, and also a greenhouse gas (Steinkamp et al., 2009). The production and consumption of NO in soil is regulated by microbial activity, mainly nitrification/denitrification processes, and chemical reactions (Pilegaard et al., 2013). Measurements using soil chambers in the field and laboratory experiments show that nitrification/denitrification, and consequently NO emissions, vary greatly with climate and soil conditions, in particular they are strongly correlated with nitrogen (N) availability, temperature and soil moisture, making soil NO emissions dependent on regional temperature and precipitation patterns, and fertilizer management practices (e.g., Bouwman et al., 2002; Meixner and Yang, 2006; Hudman et al., 2010).

Soil NO emissions are about 20% of total NO sources to the atmosphere (IPCC, 2007) and almost of the same order of magnitude of fossil fuel NO emissions. Soil emission of biogenic NO plays a prominent role in the regional atmospheric chemistry of non-urbanized areas, where anthropogenic emissions are negligible (Pilegaard, 2013). The main inputs of N compounds onto semi arid uncultivated soils, like the savanna ecosystem, are biological nitrogen fixation, atmospheric wet and dry deposition and lightning. NO fluxes are considered as one way only, even if NO deposition exists in very specific conditions (Grote et al., 2009).

Soil N losses towards the atmosphere also involve NH₃. The largest source of NH₃ emissions is agriculture, via the application of synthetic fertilizer. When released into the atmosphere, NH₃ increases the level of air pollution. In the atmosphere NH₃ has a relatively short life time of less than five days and high deposition rates, it is converted into ammonium (NH₄⁺) aerosols, which has a life time of the order of fifteen days, can travel long distances and it is relevant for air quality and climate (Fuzzi et al., 2015). The exchange of soil NH₃ is bi-directional as it also includes deposition. NH₃ returned to the surface by deposition can potentially cause eutrophication, reducing biodiversity and water quality (Sutton et al., 2009a).

The net flux of NH₃ is the combination of different exchange pathways between plant (cuticle and stomata), soil, leaf litter and atmosphere. The overall NH₃ flux for a given surface may switch from net emission to net deposition at sub-hourly, diurnal and seasonal scales. Moreover, NH₃ can be rapidly deposited onto cuticles due to its high solubility (e.g. Sutton et al., 2009b; Massad et al., 2010; Loubet et al., 2012).

The direction and magnitude of NH_3 exchanges depend on the difference in NH_3 concentration between the canopy and the atmosphere, and on a large range of environmental factors, in particular air humidity, which influence surface wetness, and soil moisture conditions, but also vegetation cover and soil characteristics. The relationships between NO and NH_3 soil fluxes have been identified through the ammonium content in the soil (McCalley and Sparks, 2008). Ammonia is mainly emitted by agricultural activities, and also by the decomposition of litter and volatilization of animal excreta (Sutton et al., 2009b; Massad et al., 2010).

Soil fluxes in West Africa have only been measured in a limited number of studies due to the challenging experimental conditions (remote sites, no power supply, very hot temperatures), and mainly with manual chamber techniques rather than more complex micrometeorological techniques (Serça et al., 1998, Le Roux et al. 1995 for NO , Delon et al., 2017 for NO and NH_3). However, tropical savanna has been recognized as one of the ecosystems characterized by the largest NO emissions (Davidson and Kinglerlee, 1997, Hudman et al., 2012).

Anthropogenic emissions of pollutants from mega cities located on the Guinean coast in South West Africa have been increasing, and are likely to keep increasing in the next decades, due to a strong anthropogenic pressure, land use change and urbanization. When transported northward on the African continent, polluted air masses meet biogenic emissions from rural areas which contributes to increased O_3 and secondary organic aerosol production, in high temperature and solar radiation conditions, highly favorable to enhance photochemistry (Knippertz et al., 2015a, 2015b).

The objectives of this study are to quantify soil fluxes of NO and NH_3 for the different land cover types typical of rural West Africa, suggest a tentative strategy to scale point measurements in the field to ecosystem and larger regional scale, and provide data for inventories and model evaluation to improve air quality and climate modelling.

In this paper we present the soil fluxes of NO and NH_3 measured in a rural site near the city of Savè, Benin, West Africa, during the DACCIWA (Dynamics-Aerosol-Chemistry-Cloud Interactions in West Africa) field campaign which lasted from 14th June to 30th July 2016 (wet season). The DACCIWA campaign was lead to investigate the possible role of local air pollution on climate change in West Africa, focusing on atmospheric composition, air pollution and cloud-aerosol interactions over several sites in the region (Knippertz et al., 2015a, 2015b, 2017). The Savè site is part of the savanna ecosystem, where grassland is intercut with crops and degraded forest. Biogenic soil fluxes measurements were taken using the manual chamber technique, which is robust and of reduced costs (Delon et al., 2017). Along with these observations we also present measurements of soil characteristics and meteorological variables from the same site. We include the comparison of measured NO soil emissions with those simulated by the Hudman et al. (2012) process-based model for NO soil emission implemented into GEOS-Chem.

2 Material and method

2.1 Site description

125 The Savè site for ground-based observations is located in a hinterland area of Benin, 6 km south west from the city of Savè (8°02'03" N, 2°29'11" E, 166 m a.s.l.). The Savè ground-based observation site is located within the Gobè site managed by the Institut National des Recherches Agricoles du Bénin (INRAB).

130 The site is characterized by a wet savanna ecosystem. The climate of the region is Sudano-Guinean, with a rainy season from March to October and a dry season from November to February (Michels et al., 2000). The average annual rainfall is about 1100 mm (Savè weather station, data averaged from 1969 to 2004, Michels et al., 2000 and Säïdou et al., 2004) and the average yearly temperature is about 27.5 °C with little variation from year to year (data averaged from 1984 to 2004, Säïdou et al., 2004). Average minimum temperature, based on 1969-1990 data, is 21.5 °C and mean maximum temperature is 35.5 °C.

135 The tree coverage in the Savè region is low with most of the land occupied by subsistence agriculture and grassland (CILSS, 2016). Four land cover types are identified at the observation site: bare soil, grassland, maize field and degraded forest. Bare soil is defined as a patch of land of minimum five by five meters wide, without vegetation growing or hanging over the plot. Ground photographs of the four land cover types are shown in Fig.1.

140 The most abundant tree species next to the grassland site and in the forest are: *Anacardium occidentale*, *Daniellia oliveri*, and *Pterocarpus erinaceus*; while the most abundant tree species next to the maize field are: *Mangifera indica*, *Cocos nucifera*, *Carica papaya L.*, *Tectona grandis*, and *Azadirachta indica*. The herbaceous vegetation is dominated by *Cleome* sp., *Crotalaria* sp., *Mucuna* sp., *Imperata cylindrica* and *Rhynchelytrum repens* next to the grassland site and in the forest, and *Commelina benghalensis*, *Euphorbia* sp., *Boerhavia diffusa*, *Phyllanthus amarus*, *Digitaria horizontalis* by maize field.

145 In the maize field, the main species, *Zea Mays*, is intercropped with *Sesamum indicum* and, to a lesser extent, with other species: *Dioscorea* sp., *Manihot esculenta*, *Arachis hypogaea*, *Vigna unguiculata*, *Gossypium* sp., *Sorghum* sp. and *Solanum lycopersicum*. The maize field was not treated with mineral fertilizer. The only livestock consist of a few dozens of domestic fowls belonging to small subsistence-oriented family farms, mainly grazing in the maize field.

150 At the Savè site, the soil is sandy, with 87% of sand and 4.1% of clay (the rest being silt) for the 0-5 cm horizon. Surface pH ranges from 6.32 to 8.46, depending on the place where the measurement is done. Mean meteorological and average soil characteristics for the observation site are reported in Table 1, dominant vegetation species and soil composition for each land cover type are given in Table 2 and 3, respectively. Sunrise and sunset UTC time at the beginning and at the end of the campaign are: 05:33 and 18:08 on 14th June, and 05:42 and 18:11 on 30th July.

155 2.2 Sampling sites

The samples were taken at the four land cover types (bare soil, grassland, maize field and forest), one location per day. Two to three samplings spots were chosen each day for each location, collecting eight to twenty-five flux measurements for both NO and NH₃ soil fluxes each day. Each location was sampled during daytime, approximately from 7 a.m. to 6 p.m., alternating measurements at the four different land cover types from one day to the other, over the entire campaign. Bare soil and the maize field were sampled for both NO and NH₃ soil fluxes on eight different, generally non-consecutive, days, grassland on ten days, and the forest site on four different days.

2.3 Chamber flux measurements

The technique used to measure NO and NH₃ soil fluxes makes use of a ThermoScientific 17i (ThermoFischer Scientific, MA, USA) to measure the concentration of NO and NH₃ through chemiluminescence, and the closed dynamic chamber technique to calculate fluxes. The details of this technique are fully described in Delon et al. (2017).

The remoteness of the study site limited installation of permanent structures and we were unable to automate our chamber measurements, thus all measurements were made manually. The instrument was powered by a generator (>100 m away) and carried around on a wheeled-table to reach the locations of the four soil types where the NO and NH₃ soil fluxes were measured. The analyzer was connected via a Teflon tube to the Teflon chamber which was put on the ground to detect the fluxes. The external sides of the chamber were covered with sand or soil to isolate it during the measurement. The soil under the chamber was left unperturbed. Adjustments were taken in order to make sure the analyzer did not reach temperatures that would invalidate the measurements.

The calibration of the NO sensor of the 17i analyzer was made before and after the campaign, with a reference NO₂ air mixture, i.e. NO in N₂ diluted with zero air. Two post-campaign calibrations were made: a first one to validate the efficiency of the NO₂ converter using a reference dilution of NO₂ in zero air, and a second one to validate the efficiency of the NH₃ converter with a NH₃/N₂ mixture diluted in pure air (Alphagaz 1, Airliquide). The zero air for NO, NO₂ calibration was obtained by filtering ambient air, previously passed on charcoal and desiccant cartridges. The dilution for all the calibration experiments was made with the 146i module (ThermoFischer Scientific, MA, USA) and the dilution module, equipped with certified mass flow meters, on board of the ATR-42 research aircraft during an inter-calibration with other NO_x instrumentation of the DACCIWA campaign (i.e. the instrumentation on the Savè measurement site tower and the instrumentation on the ATR-42 aircraft; Brito et al., 2017, Derrien et al., 2016). Reference NO, and NO₂ were ISO 6141:2015 certified at 8.73 and 8.58 ppm for NO, before and after the campaign, respectively, and 9.28 ppm for NO₂, both with 5% precision. Reference NH₃ mixture was certified at 14.78 ppm with 2% precision for NH₃. Multipoint (at least 4

points) calibrations between 50 to 250 ppb were done to ensure the linearity of the response, obtaining regression coefficients over 0.9993 for both NO and NO₂. The global precision of the analyzer is ±0.4 ppb.

190 The external volume of the chamber was 40 cm×20 cm×20 cm. The useful volume was 18×38×18 cm³, due to the thickness
of the Teflon walls. The air inlet is located on one side of the chamber, where a small vent of 4 mm in diameter provided the
pressure equilibrium between the inside and outside of the chamber. The air outlet on the other side is connected to the
analyzer with a 4 m Teflon tube. The chamber is continuously swept with an air flow of 0.7 L min⁻¹ insured by the instrument
pump, and the air flow is controlled inside the analyzer by a flow meter. The air residence time in the chamber is
195 approximately 20 min (Volume/flow), and the chamber is maintained in place for 10 min. The Teflon chamber was cleaned
(with a dry clean paper cloth) at the beginning of each day of measurement, and during the day when the deposition of sand
could potentially interfere with the measurements.

The opaque walls minimize photochemical reactions inside the chamber, which are therefore considered as negligible. The
chamber is placed on the soil for 10 min. After 10 min, the chamber is turned over to let the analyzer be swept by ambient air
200 for 5 min, then the chamber is placed again on the soil to begin a new cycle.

The calculation of the fluxes are based on the closed dynamic chamber technique, with the following assumptions: the
concentration in the chamber is equal to the concentration leaving the chamber to the analyzer, no deposition occurs onto the
Teflon walls chamber as Vaittinen et al. (2013, and references therein) have demonstrated that the adsorption of ammonia on
205 Teflon is negligible; Moreover, experimental tests with and without the Teflon chamber attached to the analyzer were made
in ambient air to verify that deposition on the walls of the Teflon chamber is negligible. The results of this experiment are
reported in Appendix A. Photochemical reactions in the gas phase inside the chamber are limited, thanks to the opaque walls.
All the details of the calculation and the chamber design are given in Delon et al. (2017). In brief:

$$210 \quad F_x = \frac{V}{A_0} \frac{\delta C_x}{\delta t} \quad (1)$$

Where F_x is the flux (NO or NH₃) in nmol m⁻² s⁻¹, δC_x is the concentration increase-variation in the chamber in nmol m⁻³
during the temporal interval δt. A₀=0.0684 m² is the surface of the ground covered by the chamber, V=0.0123 m³ is the
volume of the chamber. This equation is similar to the one in Davidson et al. (1990). The flux is then converted to ngN m⁻² s⁻¹.
215

The linear regression is calculated over a 100 to 300 s time interval after the installation of the chamber on soil for both NO
and NH₃. Based on the methodology developed in Delon et al. (2017), the dilution effect due to mixing of outside air in the

chamber is calculated for each flux separately and is in average 6.7(±1.6) % for NO and 7.7(±1.7) % for NH₃. Considering the precision of the analyzer (±0.4 ppbv), the detection limit is 0.4 ngN m⁻² s⁻¹ for NO and NH₃ fluxes

The chemical reactions inside the chamber can determine NO consumption, and consequently an underestimation of the NO fluxes calculated with our method. This underestimation is taken into account and calculated following the method by Pape et al. (2009) with the relation $k \cdot [\text{NO}] \cdot [\text{O}_3]$. In this relation k is the temperature-dependent reaction rate constant (Pape et al., 2009, Atkinson et al., 2004), $[\text{NO}]$ is measured by the ThermoScientific 17i at soil level just before positioning the chamber for the measurement of soil fluxes, and $[\text{O}_3]$ at soil level is derived by measurements of NO and NO₂ at soil level made with the ThermoScientific 17i and measurements of NO, NO₂ and O₃ taken on an 8 m high tower. On the 8 m high tower, NO and NO₂ were measured with a Model 42C TraceLevel NO-NO₂-NO_x by Thermo-environmental Instruments Inc., calibrated with the same method as the ThermoScientific 17i and with 0.05 ppb (2-sigma) detection limit. Ozone was measured on the tower with a Model 49i Ozone Analyzer by Thermo-environmental Instruments Inc. with 1 ppb detection limit. The Model 49i Ozone Analyzer was calibrated by comparison with a Thermo Scientific Model 49PS reference instrument. The reference instrument is sent twice a year to the French Laboratoire national d'Essais (LNE) for comparison with a National Institute of Standards and technology (NIST). All data on the tower were sampled at 10 seconds. $[\text{O}_3]$ at soil level was then calculated considering the diurnal steady state of the reactions described in equation (2) and (3), using equation (4):



$$[\text{O}_3]_{sl} = \frac{[\text{NO}]_l [\text{O}_3]_l [\text{NO}_2]_l}{[\text{NO}_2]_l [\text{NO}]_l} \quad [\text{O}_3]_{sl} = \frac{[\text{NO}]_{tl} [\text{O}_3]_{tl} [\text{NO}_2]_{sl}}{[\text{NO}_2]_{tl} [\text{NO}]_{sl}} \quad (4)$$

240 Where $[\]_{sl}$ is the concentration at the soil level and $[\]_{tl}$ is the concentration measured on the tower. In conclusion, we correct NO fluxes for the underestimation of NO fluxes due to chemical reactions inside the chamber with values ranging between 0 and 63% (8% on average for the whole campaign).

As studied by Kristensen et al. (2010a) and Kristensen et al. (2010b), O₃ deposition can decrease O₃ concentration close to soil surface further. However, considering that O₃ concentrations calculated near the soil are already very low (1 ppb at soil level compared to 24 ppb at 8 m, averaged for the entire measurement campaign), O₃ deposition has been considered of secondary importance in this calculation and has not been included. If O₃ deposition were to be included it would possibly decrease the correction of NO fluxes and consequently slightly decrease NO emissions in a negligible proportion compared to the correction already applied for the chemical reactions inside the chamber.

250 The measurements have not been corrected from a possible interaction with particulate matter.

2.4 Data quality check

A quality check method based on the following criteria is used to select observed fluxes (Delon et al., 2017):

- The coefficient of determination for linear regression R^2 has to be higher than 0.4 (considered as a significant correlation) for NH_3 fluxes, and higher than 0.8 for NO fluxes.

255 - A flux error was estimated by calculating the dispersion of points around the linear regression's slope. According to this method, the dispersion for NO flux calculation is comprised between 5 and 12%, and the dispersion for NH_3 flux calculation is comprised between 15 and 20%

- The concentration difference between the last and the first NH_3 measurement point has to be more than 0.4 ppb (sensitivity of the analyzer). R^2 was generally lower than 0.4 for concentration differences below 0.4 ppb.

260 Finally, 351/488 (72%) NH_3 flux measurements and 459/488 (94%) NO flux measurements are considered valid.

2.5 Meteorological station

Continuous in situ observations of meteorological variables, including air and soil temperature and moisture, rainfall, wind speed, wind direction, radiation and energy balance components were taken at the Savè site as part of the DACCIWA campaign. Data are provided as 1 min averages, apart from energy fluxes which are given as 30 min averages (Derrien et al., 265 2016, Kohler et al., 2016, Handwerker et al., 2016, Wieser et al., 2016). An overview of the complete set of instrumentation and measurements is given by Brooks et al. (2017), while a summary of the available ground-based meteorological observations is given by Kalthoff et al. (2017). In this study we present soil moisture measured in two distinct locations of the Savè site by the Karlsruhe Institute of Technology (KIT) instrumentation at 5cm depth on grassland, and average soil moisture, between 0 and 30 cm, measured by the Université Paul Sabatier (UPS) instrumentation in the maize field. Details 270 of the instrumentation is given by Brooks et al. (2017). We include soil moisture measured with both systems as the inter-comparison of the two methods is out of the scope of this study.

2.6 Soil characteristics (texture, pH, N content)

Soil samples were collected with a cylinder of known volume (290 cm^3) during the measurement campaign to analyze the biogeochemical characteristics of the site. Soil samples (0-5 cm) were taken for each land cover type where NO and NH_3 275 fluxes were measured. Fifteen S_s samples were collected at the four different land cover types, three to four times during the campaign.

Samples were dried in ambient conditions (mean day-time temperature is approximately $26 \text{ }^\circ\text{C}$, Kalthoff et al., 2017), and stored in the dark. After drying, the weight of the samples was measured to determine the bulk density ($d_b = \text{dry soil mass} / \text{total volume}$), which was found to be $1.24 \pm 0.14 \text{ g cm}^{-3}$. Assuming a density of soil particles (d_p) of 2.6 g cm^{-3} , the Water Filled Pore Space (WFPS) is calculated with the following equation 5:

$$WFPS = SM / (1 - d_a / d_r) \quad (5)$$

where SM is soil moisture in %.

285 Soil samples were analyzed for the determination of texture, ammonium concentrations $[NH_4^+]$, C/N ratio, total C, total N, and pH at the GALYS Laboratoire (<http://www.galys-laboratoire.fr>, NF EN ISO/CEI 17025: 2005). The analyses were performed two months after sampling. We assume that the ammonium content in litter or soils is not modified by volatilization or chemical transformation during transport and storage, because of the very low soil moisture level in samples. Indeed, when collected, WFPS of the samples ranged between 6 and 14% (mean=8.5±3.5%), and soil temperature
290 between 35 and 38 °C (data obtained from the databases described in Brooks et al., 2017). Bai et al. (2013, and references therein) have found that significant changes in nitrification and net mineralization (influencing the ammonium content) may occur when soil temperature raises until 35°C (the optimum for nitrification), for optimal soil moisture conditions (WFPS =20%, Oswald et al., 2013. In the present study, soil temperatures when sampling were equal or above the optimum, and WFPS was below the optimum, reducing the nitrification efficiency and the change in ammonium content. Past studies have
295 demonstrated that microbial activity and mineralization processes are inhibited in low soil moisture conditions, even when soil temperature is high (Bai et al., 2013, and references therein). Several authors have published results of ammonium concentrations measured in soils dried in ambient air. For example, Dick et al., (2006) collected top soil after the wet season in two sites in Senegal. The authors state that their soils were considered dry when collected and were air-dried in the mid-day sun immediately after collection. The protocol used in our study is identical. Some authors have also published results of ammonium concentrations measured in soils that were dried in ambient air. Other studies (Bai et al., 2010, Dick et al., 2006, Cassity-Duffrey et al., 2015, Vanlauwe et al., 2002) also published ammonium measurements made on air dried soils from seasonally dry climates with comparable textures to the soil in Savé. Soil texture is determined following norm NF X 31e107. Clay (<2 µm), fine silt (2 to 20 µm), coarse silt (20 to 50 µm) and total sand (50 to 2000 µm) are determined without decarbonation. Organic carbon and total carbon are determined following norm NF ISO 10694. The whole carbon of the sample is transformed into CO₂. Then CO₂ is measured by thermal conductivity. NF ISO 13878 is used for Total N. Mineral nitrogen is determined following an internal method MT-AZM adapted from norm NF ISO 14256-2. This method uses a
300 potassium chloride solution and is COFRAC certified. The sample is heated at 1000°C with O₂. Products of combustion or decomposition are reduced in N₂. N₂ is then measured by thermal conductivity (catharometer). pH is determined according to norm NF ISO 10390, with soil samples stirred with water (ratio 1/5).

310 2.7 Soil ammonia emission potential Γ_g and compensation point χ_g

Measurements of soil pH and ammonium concentrations $[NH_4^+]$ are used to quantify the soil emission potentials for the different land cover types at the measurement site. The soil emission potential Γ_g is the ratio of $[NH_4^+]$ to $[H^+]$ concentrations in the water solution of the soil (mol L⁻¹). A large Γ_g indicates that the soil has a high propensity to emit NH₃, considering that the potential emission of NH₃ depends on the availability of ammonium in the soil and on pH.

The soil compensation point (χ_g) has been calculated from the emission potential Γ_g , as a function of soil surface temperature (T_g in K) according to Wentworth et al. (2014):

$$\chi_g \text{ (ppb)} = 13\,587 \cdot \Gamma_g \cdot e^{-(10\,396\text{K}/T_g)} \times 10^9, \quad (56)$$

The soil compensation point indicates the equilibrium between gaseous NH_3 in the soil pore space and $[\text{NH}_4^+]$ in the soil solution, i.e. the concentration of NH_3 for which the NH_3 flux switches from emission to deposition (or viceversa).

2.8 Stepwise multiple regression analysis

A stepwise linear multiple regression analysis was performed between daily averaged gas fluxes of NO and NH_3 and relevant available daily averaged variables such as wind speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, outgoing longwave radiation and incoming shortwave radiation. Soil parameters such as mineral nitrogen, total N and organic C, soil texture and pH could not be used for this regression analysis since their relative measurements did not have the same temporal resolution as the other parameters. The R software (<http://www.R-project.org>) was used to provide the results of this linear regression analysis.

330 2.9 GEOS-Chem

GEOS-Chem is a global three-dimensional model of tropospheric chemistry driven by meteorological input from the NASA Goddard Earth Observing System (www.geos-chem.org, Bey et al., 2001). In this study we use GEOS-Chem Version 10-01 which includes the process-based parameterization of soil NO emission by Hudman et al. (2012). This parameterization represents available nitrogen (N) in soils using biome specific emission factors, online wet and dry deposition of N, and fertilizer and manure N derived from a spatially explicit dataset, distributed using seasonality derived from data obtained by the Moderate Resolution Imaging Spectrometer (MODIS). Emissions are a smooth function of soil moisture and temperature consistent with point measurements and ecosystem scale experiments. This parameterization also included pulsing following soil wetting by rain or irrigation, represented as a function dependent on dry spell length. The parameterization by Hudman et al. (2012) was successfully evaluated for pulsing events in central Sahel (0–30° W, 12–18° N).

340

Boundary conditions for our experiment are generated from a global GEOS-Chem simulation at 4° x 5° horizontal resolution. The regional GEOS-Chem model for West Africa runs at a horizontal resolution of 0.25° x 0.3125° (latitudes 6°S–16°N, longitudes 18.125°W–26.875°E) and a vertical resolution of 47 levels (up to 0.01hPa). Meteorology is driven by the NASA GMAO (Global Modeling and Assimilation Office) GEOS-FP (Forward Processing) assimilated meteorological data. The global model is spun-up from 1st May 2015 to 1st May 2016. The global simulation is then run from 1st May 2016 to 1st August 2016, outputting boundary condition files for West Africa. The regional West Africa simulation is then run from

1st May 2016 to 1st August 2016 using the 4° x 5° boundary conditions from the global simulation. All simulations use the GEOS-FP meteorology which has a three-hour time resolution. We used the same MODIS/Koppen land cover map as in Hudman et al. (2012; <http://glcf.umd.edu/data/lc>) which includes 24 land cover types. In this simulation we use EDGAR v4.2 (EC-JRC/PBL, 2011) for anthropogenic emissions, GFED4 (Giglio et al., 2013) for biomass burning emissions and MEGAN v2.1 (Guenther et al., 2012) for biogenic emissions of volatile organic compounds. The same emission inventories are used for both the boundary conditions and the West Africa simulation.

3 Results and discussion

3.1 Meteorological data

355 Mean air temperature averaged over the whole campaign was 25.4 ± 2.6 °C, mean wind speed was 1.3 ± 0.6 m s⁻¹, mean relative air humidity is 86.3 ± 10.5 %, mean soil temperature was 25.2 ± 3.4 °C, mean KIT soil moisture at 5 cm was 7.1 ± 3.6 %, while mean UPS soil moisture averaged between 0 and 30 cm was 4.5 ± 2.8 %. Total KIT precipitation was 198 mm for the whole campaign, and total UPS precipitation was 215 mm.

360 Median diurnal cycles of air temperature, specific humidity and precipitation are reported in Kalthoff et al., (2017). Knippertz et al. (2017) distinguish four different phases of the monsoon season during the DACCIWA campaign (14th June to 30th July 2016) over the DACCIWA focus region (5–10° N, 8° W - 8° E), which covers a wide area of West Africa (see Fig. 1, Knippertz et al. (2017)). The division into phases is mainly based on the north–south precipitation difference between the coastal zone (0–7.5° N) and the Sudanian–Sahelian zone (7.5–15° N), both averaged across the longitude range 8° W–8° E. Savè (8.03° N) is located very close to the border between the two zones, with a rainfall pattern that seems to follow more closely that of the coastal zone rather than that of the northern inland Sudanian–Sahelian zone. These four phases are: the pre-onset phase characterized by a rainfall maximum near the coast (before 21st June, phase 1); the post-onset phase during which the rainfall maximum occurred inland (22nd June - 20th July, phase 2); the wet westerly regime when the rainfall maximum shifted back to the coast (21st - 26th July, phase 3); and the recovery of the monsoon with a shift of the rainfall maximum inland (27th July until the end of the campaign, phase 4). A specific period within phase 2 is indicated “vortex”, during which an unusual development occurred (09th - 16th July): in the north, a cyclonic feature slowly propagated from eastern Mali to Cape Verde and in the south, an anticyclonic vortex tracked in the west-northwesterly direction along the Guinean coast (see Knippertz et al. 2017 for a more detailed description). At the Savè site the most intense rainfall events happened the day before the first soil fluxes observation, on 15th June 2016, and towards the end of the measurement campaign between 20th and 23rd July 2016. Other minor rainfall events are recorded on 19th and 27th June, 8th, 12th, 13th, 24th and 26th July. Daily rainfall measurements are reported in Figs. 2 to 5.

3.2 Soil texture, soil organic carbon, total nitrogen, pH and ammonium content

Bare soil recorded the lower amount of total sand ($83.75 \pm 1.82\%$) and the higher amount of clay ($5.13 \pm 0.63\%$), fine ($5.13 \pm 0.96\%$) and coarse silt ($5.98 \pm 0.51\%$). Grassland recorded the higher amount of total sand ($89.20 \pm 0.71\%$) and the lower amount of clay ($3.15 \pm 0.50\%$) and fine silt ($2.93 \pm 0.32\%$), while intermediate values were found for the maize field and forest (Table 3). These values determine the classification of sandy soil for all measurements sites.

Soil organic carbon (C) and total nitrogen (N) are respectively $12.2 \pm 5.7 \text{ g kg}^{-1}$ and $0.95 \pm 0.51 \text{ g kg}^{-1}$, averaged for all land cover types over the entire campaign. Table 4 gives soil characteristics for each land cover types, including individual values of C/N ratio, soil organic C and total N for the entire field campaign. The highest average soil organic C was measured for bare soil ($17.3 \pm 5.9 \text{ g kg}^{-1}$) and the lowest soil organic C was measured for grassland ($6.2 \pm 1.3 \text{ g kg}^{-1}$), while the maize field and forest site accounted for $14.1 \pm 2.9 \text{ g kg}^{-1}$ and $11.4 \pm 4.5 \text{ g kg}^{-1}$ soil organic C, respectively. The highest average total N was measured for bare soil ($1.44 \pm 0.51 \text{ g kg}^{-1}$) and the lowest total N was measured for grassland ($0.44 \pm 0.04 \text{ g kg}^{-1}$), while the maize field and forest site accounted similar amounts of total N, $0.99 \pm 0.19 \text{ g kg}^{-1}$ and $0.94 \pm 0.48 \text{ g kg}^{-1}$, respectively. Values of C/N, soil organic C and total N recorded for grassland at the Savè site compare closely to those reported by Delon et al. (2017, table 2) for the semi-arid site of Dahra ($15^{\circ}24' \text{ N } 15^{\circ}25' \text{ W}$), Senegal. Our values of C/N and total N for grassland are also close to those reported by Le Roux et al. (1995, table 1) and Lata et al. (2004) for the wet savanna ecosystem of Lamto ($6^{\circ}13' \text{ N}, 5^{\circ}20' \text{ W}$), Ivory Coast, although we observe lower values of soil organic C compared to ~~Le Roux et al. (1995) and Lata et al. (2004)~~ these studies. Values of C/N and soil organic C recorded for the maize field at the Savè site are slightly higher than those recorded by Barthes et al. (2004) in a maize field at Agonkanmey ($6^{\circ}24' \text{ N}, 2^{\circ}20' \text{ E}$), near Cotonou in southern Benin.

All the sites listed in the comparison in the previous paragraph are sandy, as the Savè site. The Dahra site (Delon et al., 2017) also shows similar pH than our site (Table 5), while lower pH (acidic or near-neutral) was recorded at the sites of Lamto (Le Roux et al., 1995, Lata et al., 2004) and Agonkanmey (Barthes et al., 2004). Table 5 provides individual values of pH, $[\text{NH}_4^+]$, Γ_g and χ_g for the entire field campaign. The highest average pH was observed for bare soil (8.23) and the lowest for the forest site (7.07), while measured average pH was 7.27 for grassland and 7.70 for the maize field. The $[\text{NH}_4^+]$ content averaged for all land cover types over the entire campaign is $5.33 \pm 4 \text{ mg kg}^{-1}$. The highest average $[\text{NH}_4^+]$ was recorded for the maize field ($7.9 \pm 6 \text{ mg kg}^{-1}$) and the lowest for grassland ($2.0 \pm 0.3 \text{ mg kg}^{-1}$). Average $[\text{NH}_4^+]$ is $6.2 \pm 5 \text{ mg kg}^{-1}$ and $7.0 \pm 2.2 \text{ mg kg}^{-1}$ for forest and bare soil, respectively. Dick et al. (2006) have found NH_4^+ concentrations between 2 and 8 mgN.kg^{-1} in Senegalese soils, which is very close from our results. Vanlauwe et al. (2002) have found values between 0.8 and 1.4 mgN.kg^{-1} in West African moist savanna soils (in Togo and Nigeria).

Higher soil organic C and N over bare soil could be due to the fact that these bare soil patches experienced recent burning
410 (Santín and Doerr, 2016). The higher $[\text{NH}_4^+]$ over the maize field can be caused by chicken excreta, as chickens were
roaming over the maize field (Paillat et al., 2005, Tiquia and Tam, 2000).

3.3 Soil emission potential Γ_g and compensation point χ_g

The mean soil emission potentials for the Savè site is $43\,714 \pm 58\,077$, with values ranging from 380 to 159 343. The highest
values of soil emission potentials are observed for bare soil ($113\,672 \pm 67\,788$), followed by maize field ($33\,880 \pm 20\,680$),
415 forest ($11\,982 \pm 11\,061$) and grassland ($4\,929 \pm 4\,409$). The ammonia compensation point ranges between 5 to 2 215 ppb,
with soil temperatures between 25 and 29 °C. The highest values of χ_g are observed for bare soil ($1\,607 \pm 993$), followed by
maize field (473 ± 317), forest (175 ± 167) and grassland (58 ± 47). Our values of soil emission potential for bare soil and
maize (no fertilization) are comparable with those presented in Massad et al. (2010, table 4), although those data come from
measurements taken on different ecosystems. Both Γ_g and χ_g values recorded at the Savè site exceed those recorded by Delon
420 et al. (2017) over a grazed semi-arid Sahelian ecosystem in Senegal.

3.4 NO fluxes

NO fluxes from soil measured during the field campaign range between 0 and $48.05 \text{ ngN m}^{-2} \text{ s}^{-1}$. NO fluxes averaged over all
land cover types are $4.79 \pm 5.59 \text{ ngN m}^{-2} \text{ s}^{-1}$, while average NO fluxes for each land cover type are: $8.05 \pm 3.49 \text{ ngN m}^{-2} \text{ s}^{-1}$
for bare soil, $3.73 \pm 1.76 \text{ ngN m}^{-2} \text{ s}^{-1}$ for the maize field, $2.87 \pm 1.49 \text{ ngN m}^{-2} \text{ s}^{-1}$ for forest and $2.82 \pm 3.46 \text{ ngN m}^{-2} \text{ s}^{-1}$ for
425 grassland. Soil emissions of NO from the different land cover types provide similar values, NO emissions from bare soil are
higher on average, but have a larger standard deviation.

Other measurements of biogenic NO soil emissions from the West African wet savanna can be found in Delon et al. (2012,
table 7). We find that our measured NO soil emissions averaged over all land cover types are higher than those measured
430 from other wet savanna sites. Our measurements are in better agreement with emissions from dry savanna grasslands (Delon
et al., 2012), and with measurements from a semi-arid savanna, with over 80% sandy soil, in South Africa (Parsons et al.,
1996, Scholes et al., 1997). However, these studies measured NO emissions during different seasons and soil moisture
conditions compared to our study. For example, Parsons et al. (1996) recorded NO emissions up to $20 \text{ ngN m}^{-2} \text{ s}^{-1}$ over an
open savanna during the period going from the end of the dry season to the beginning of the wet season. Nitric oxide
435 emissions of the same magnitude as in our study were also recorded over a grazed semi-arid Sahelian ecosystem in Senegal
during the month of July by Delon et al. (2017): $5.7 \pm 3.1 \text{ ngN m}^{-2} \text{ s}^{-1}$ in July 2012 and $5.1 \pm 2.1 \text{ ngN m}^{-2} \text{ s}^{-1}$ in July 2013.

Daily means of NO concentrations are measured close to the soil (0.1m, half height of the chamber) and reported in fig. 2 to
5. Daily means of NO concentration vary from 1.28 to 5.40 ppb for all sites. The average concentration during the whole
440 campaign on all sites is 2.70 ± 1.03 ppb. Average NO concentration is 2.97 ± 1.49 ppb on bare soil, 2.57 ± 0.96 ppb on

grassland, 2.55 ± 0.83 on maize, and 2.76 ± 0.65 ppb on forest soil. The concentrations are quasi equivalent for all sites. As these concentrations are low, they do not lead to NO deposition on soil and the NO flux stays positive. In fact, NO deposition has been measured in other studies only in the case of high NO concentrations (>60 ppb, Laville et al., 2011).

445 Figures 2 to 5 show daily averaged NO and NH₃ fluxes (± 1 standard deviation) for each land cover type, along with precipitation and soil moisture. The spatial variability of NO fluxes is high, especially for bare soil, forest and the maize field where underground roots, not visible at the surface, are heterogeneously distributed. These roots are likely to influence the ammonium content of the soil, and the subsequent NO flux measurement. Standard deviation is generally smaller for
450 grassland (except for two days, July 9th and 13th), where the vegetation (and the root distribution) is more homogeneous. The variation of soil moisture is consistent with the presence of rain events, showing a sharper increase of soil moisture at 5 cm, especially after rainfall following dry periods.

NO emissions from bare soil and grassland show an increase, sharper for grassland, one to two days after the rain event on 8th July. The longer rain event between 20th and 24th July does not seem to produce an increase in NO emissions (data
455 available only for maize field and forest). This might be linked with the non-linear relationship between NO biogenic soil emissions and soil water content (Oswald et al., 2013). In fact, a light precipitation event (5-15 mm) occurring on dry soils can result in a large flux of NO (Meixner & Yang, 2006, Hartley & Sclesinger, 2000). However, when soil moisture stays at an equivalent level, after several rain events, pulse emissions do not occur (Millet et al., 2004). Due to this non linear character of the NO fluxes, no direct correlation was found between NO fluxes and environmental variables such as soil moisture or soil temperature taken individually. Moreover, soil temperature and soil moisture were not measured on the same soil parcel where the soil fluxes were measured and the location of the soil flux measurements was not kept constant even for the same land cover type on the same measurement day. This measurement protocol was designed to give an estimate of soil fluxes at a large ecosystem scale, rather than reproducing the relationships between soil fluxes and meteorological variables, like soil temperature and soil moisture.

465 A multiple linear regression analysis was performed between daily mean NO fluxes and the following variables: wind speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, upward longwave radiation and downward shortwave radiation. This regression gives $R^2=0.49$ (p-value=0.004), indicating a weak but existing relationship between those variables and NO soil emissions. while the regression was weak between NO fluxes and each individual variable. This correlation shows the influence of these environmental variables considered collectively on NO fluxes, highlighting the underlying mechanisms responsible for NO release to the atmosphere. However, the aim of our experiment was not to reproduce the known relationships between soil fluxes and meteorological variables, like soil temperature and soil moisture, but to give an estimate of soil fluxes at the larger ecosystem scale. For this reason the location of the soil flux measurements was not kept constant even for the same land cover type on the same measurement day. Moreover, soil temperature and soil moisture were

475 ~~not measured on the same soil parcel where the soil fluxes were measured. Therefore,~~ Our experiment does not show the
480 details of microbial and physical processes driving soil fluxes at a single point, ~~because measurements are done at different~~
locations every day. -but aims to estimate the spatial variability of fluxes at the ecosystem scale.

The NO flux estimated in this study does not consider the impact of vegetation on the net ecosystem flux, as we focus on soil
480 fluxes only. However, the net emission to the atmosphere should take into account the oxidation of NO to NO₂ and the
eventual re-deposition of NO₂ on the vegetation, i.e. what is called Canopy Reduction Factor and is assumed to be a linear
function of the Leaf Area Index (e.g. Yienger and Levy, 1995, and Ganzeveld et al., 2002)).

3.5 NH₃ fluxes

NH₃ fluxes measured during the field campaign range between -6.59 and 4.96 ngN m⁻² s⁻¹. Ammonia fluxes averaged over
485 all land cover types are -0.91 ± 1.27 ngN m⁻² s⁻¹, showing a predominance of NH₃ deposition over emission, which is
verified for every land cover type, with an average value of: -1.33 ± 0.86 ngN m⁻² s⁻¹ for bare soil, -0.75 ± 0.31 ngN m⁻² s⁻¹
for the maize field, -0.48 ± 0.55 ngN m⁻² s⁻¹ for grassland, and -0.30 ± 0.38 ngN m⁻² s⁻¹ for forest. Low positive ammonia
fluxes, indicating average NH₃ emission, are only recorded during three days, between 6th and 8th July, after the longest dry
period of the measurement campaign (Figures 2 to 5).

490 To our knowledge, NH₃ soil fluxes from west African wet savanna are not available in the scientific literature. In Delon et al.
(2017) NH₃ soil fluxes measured in Dahra (15°24' N 15°25' W), Senegal, on a dry savanna ecosystem, show low fluxes with
a predominance of NH₃ emission: 1.3 ± 1.1 ngN m⁻² s⁻¹, -0.1 ± 1.1 ngN m⁻² s⁻¹ and 0.7 ± 0.5 ngN m⁻² s⁻¹ over three different
measurements campaigns. However, Sutton et al. (2007) shows how pre-cut grassland is characterized by NH₃ deposition, as
495 in our study, in contrast to post-cut grassland, which is marked by NH₃ emission. It is interesting to notice that the literature
provides up to about 700 ngN m⁻² s⁻¹ NH₃ emission for fertilized Zea Mays fields (Walker et al., 2013) while in our study site
NH₃ deposition was recorded for the maize field, which is not treated with mineral fertilizer.

As for NO concentrations, NH₃ concentrations are reported in fig. 2 to 5. Daily means of NH₃ concentration vary from 0 to
500 12.46 ppb for all sites, and the average concentration is 4.42 ± 3.23 ppb during the whole campaign. Average NH₃
concentration is 6.28 ± 3.90 ppb for bare soils, 3.28 ± 1.79 ppb for grassland, 4.36 ± 3.99 for the maize field, and 3.68 ± 2.13
440 ppb for forest. The largest deposition fluxes are found on bare soils, where the largest concentrations are measured.

A multiple linear regression analysis was performed between daily mean NH₃ fluxes and the following variables: wind
505 speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, outgoing longwave radiation and incoming shortwave
radiation). This regression gives a weak but existing relationship, with $R^2=0.37$ (p-value=0.03). This correlation highlights

the link between NH_3 fluxes and relevant environmental parameters. However, the same considerations explained in Sect. 3.4 for NO emissions are also valid for the correlation between NH_3 fluxes and meteorological variables.

510 According to the current parameterization of soil ammonia emission potential (Sect. 2.7), high values of pH and $[\text{NH}_4^+]$ in
the liquid phase will determine high values of Γ_g indicating that the soil has a high propensity to emit NH_3 . However, despite
the high values of Γ_g recorded, our measurement site remains a net sink for NH_3 . The reasons for this can be manifold. One
explanation could be that soil particles on our site may have a high adsorption capacity limiting the amount of soil gaseous
 NH_3 concentrations (Neftel et al., 1998) and the largest part of the estimated ammonium content in the soil may not be in the
515 liquid phase, but adsorbed by solid soil particles. In these conditions ammonium will not be available for gas exchange to
open porosity and the atmosphere (Flechard et al., 2013). Another explanation could be given by the presence of a water film
at the soil surface (linked to high air humidity at the site), which will increase the net deposition process. David et al. (2009)
conclude from their measurements that the bare soil can be a significant source of NH_3 only for a limited period and only
when the cut vegetation is removed, but not if the soil surface remains covered by grass. Measurements in Ferrara et al.
520 (2014) show other occurrences of high soil ammonia emission potential and NH_3 deposition.

Our measurements were conducted without vegetation inside the chambers, but vegetation was present in the fields. It is
important to mention that the role of vegetation on NH_3 bidirectional fluxes is essential, especially during the wet season
(time of the experiment), when deposition on the vegetation through stomata and cuticles dominate the exchange (during rain
525 events, the cuticular resistance becomes small and cuticular deposition dominates), due to an increase of the deposition
velocity of NH_3 (consecutive to the humidity response of the surface) and a decrease of the canopy compensation point,
sensitive to the surface temperature and the surface wetness (Wichink-Kruit et al., 2007).

3.6 Comparison of observed and modelled NO soil emissions

We have compared observed daily averaged (8 a.m. to 6 p.m.) soil NO emissions with those modelled by GEOS-Chem for
530 the entire period of the campaign over the model grid box including the measurement site. The model grid box is positioned
at latitude $8.0^\circ \text{ N} - 8.25^\circ \text{ N}$ and longitude $2.19^\circ \text{ E} - 2.5^\circ \text{ E}$. The area of this grid box is 958 km^2 . The land cover type within
this grid box is classified as “Savannah (Warm)” but the surrounding area also consists of “Woody Savannah”, while the
observations were taken over the four land cover types representative of the region: bare soil, grassland, maize field and
forest.

535 The model is able to reproduce mean air temperature ($25.3 \pm 0.7 \text{ }^\circ\text{C}$) and the main rain events. Soil emissions of NO are well
simulated in magnitude. Simulated NO emissions are often higher than those recorded over the grassland areas, however,
simulated NO emissions are often within the error bars of measurements (Fig. 6). [The model uses land cover and vegetation](#)

540 types to simulate the highly variable land and vegetation cover of the observation site, for this reason we do not expect the model to reproduce the site-to-site variability of the measured soil fluxes, but to at least reproduce their average magnitude and behaviour. It appears that when the model is able to reproduce the length and the intensity of the rain events, NO emissions are especially well simulated, e.g. the model is able to reproduce the longest rain period (from 20th to 26th July 2016) and the decrease of emissions at the end of the measurement campaign.

3.7 Estimate of total NO soil emissions and NH₃ deposition for Benin

545 In order to give a tentative estimate of NO and NH₃ soil fluxes for Benin we have used the land use/land cover map of Benin provided by the US Geographical Survey Atlas: Landscapes of West Africa – A Window on a Changing World (CILSS, 2016, Fig. 7a). The method of mapping land use/land cover used in this atlas was based on Landsat imagery and expert visual interpretation. In particular, these maps provide an accurate indication of cropland distribution using visual interpretation. According to CILSS (2016) Benin's present-day (2013) land cover is mainly savanna, almost 60%, followed by agricultural
550 land, 31%, while forest is only a small fraction under 1% (the rest of the surface is mainly gallery forest and, on a smaller extent, settlements). In the Atlas (CILSS, 2016) bare soils are defined as those surfaces that are bare even in the green/rainy season. For Benin, the amount of bare soil estimated by CILSS (2016) is very small, not big enough to appear on the Atlas' maps. We have multiplied average NO emissions measured at the Savè site for each land cover type by an estimate of the land cover area of each class given by the Atlas. We have made some approximations: as the land use/land cover maps do not
555 distinguish between shrub savanna, tree savanna, and wooded savanna, we have considered NO soil emissions from Savè's grassland savanna to be representative of the general savanna category in CILSS (2016). Moreover, the Atlas has a crop category that does not distinguish the type of crop and we only have observations of NO soil emissions from an intercropped maize field. We have taken NO soil emissions from the maize field as representative of NO emissions of Benin's agricultural land, but other cultures are present in other parts of the country, e.g. oil palm plantations, where possibly stronger
560 fertilization could determine higher NO soil emissions. This tentative calculation gives that Benin's NO soil emissions for the month of July (wet season) is 1.17 ± 0.6 GgN/month, i.e. 0.09% of the average global monthly NO soil emissions as given by Davidson et al., (1997).

We have also calculated Benin's total monthly NO soil emissions with GEOS-Chem adding together the NO soil emissions
565 from the grid boxes where 50% or greater of the box lies within Benin. Benin's total monthly NO soil emissions calculated with GEOS-Chem for the month of July are 1.44 GgN/month and agree with the tentative calculation given above (1.54 ± 0.8 GgN/month). However, the land cover types covering Benin in GEOS-Chem differ from those in the US Geographical Survey Atlas (CILSS, 2016). In GEOS-Chem Benin is cover by 60.9% savanna, 31.4% woody savannah, 4.5% grassland, 1.3% mixed forest and 0.6% urban and built-up lands. Benin's total monthly NO soil emissions calculated with GEOS-Chem
570 for the months of May and June are higher, 3.51 GgN/month and 2.59 GgN/month, respectively, given those months are at the beginning of the wet season and are characterized by more predominant pulse emissions. Using the same method

described above we have upscaled point measurements of NH_3 fluxes with relevant land cover surfaces from CILSS (2016) and obtained that total NH_3 dry deposition for the month of July is $0.21 \pm 0.11 \text{ GgN/month}$ ($0.22 \text{ kgN ha}^{-1} \text{ yr}^{-1}$). This value is about ten times smaller than the estimation of NH_3 dry deposition given in Adon et al. (2013) for the wet savanna site of Djougou (Benin, 9.7°N , 1.7°E) for the month of July, which is around $2.5 \text{ kgN ha}^{-1} \text{ yr}^{-1}$.

4. Conclusion

We provide soil flux measurements along with soil characteristics for a land cover type, savanna, that is considered to have large NO emissions (Davidson and Kinglerlee, 1997), and for an area of the world, West Africa, with little observations. The aim of this study is to contribute to our knowledge in biogenic soil nitrogen exchanges, provide data for inventories and model evaluation to improve air quality and climate modelling.

In situ measurements were made in a wet savanna site in central Benin from mid-June to the end of July 2016. Complementary to these exchange fluxes, soil N and C content, as well as soil pH, soil moisture, soil temperature and meteorological data were measured. Soil fluxes of NO and NH_3 were measured over four different land cover types in order to give a tentative estimate of regional soil fluxes.

Given the set up of the experiment, the known relationships between soil fluxes, soil temperature and soil moisture were not reproduced. Rather than looking at the microbial and physical processes behind soil fluxes, we are able to provide observations that are representative of a bigger surface area and that represent the spatial variability of fluxes. However, we observe that while shorter rain events determine an increase in NO soil emissions, the longer rain event at the end of the campaign (20th to 24th July 2016) is accompanied by a decrease in NO soil emissions, in agreement with the fact that the relationship between NO soil emissions and soil moisture is not univocal. Soil emissions of NO increase until an optimum value of soil moisture is reached and then decrease (Oswald et al., 2013).

NH_3 emissions measured in this study probably underestimate total NH_3 emissions for the entire country, as possibly higher localized NH_3 emissions are present in the south of the country where industrial scale agriculture would probably deploy mineral N fertilization.

Soil NO emissions simulated by GEOS-Chem are in good agreement with the local observations taken at the site of Savè, providing a good baseline for simulating local atmospheric chemistry. Moreover, GEOS-Chem is also in good agreement with the tentative total monthly NO soil emission estimate for Benin for the month of July made with local observation in Savè and US Geographical Survey Atlas (CILSS, 2016). All these elements contribute to improve our confidence in the results of modelling studies of local and regional air quality and climate over this region.

605 Agriculture is the first form of economic activity in Benin, occupying a majority of the active population. The most obvious recent change in land cover is the major expansion of agricultural land across most regions of Benin. Agricultural areas (including plantations and irrigated agriculture) progressed from 9.2 to 27.1 % of the total country area between 1975 and 2013, improving food security. Oil palm trees are the main crop, and oil palms farmland already covered most of the southern Terre de Barre plateau of Benin by 1975, and increased by about 28 percent over the following 38-year period. A century or more ago, Benin was covered by dense, biologically diverse forest. Since then, Benin has lost nearly all of that forest cover, by 2013, 58 % of the 1975 forest cover had been lost, leaving only 0.2 percent of the country covered with dense forest. Savanna area has also decreased by 23 percent since 1975, but it still remains the dominant land cover type in Benin and covers more than half of the country (CILSS, 2016).

615 More measurements of NO and NH₃ exchanges between soil-vegetation-atmosphere in areas of Benin (or West Africa) interested by land-use change could improve our estimate of the impact of biogenic soil emissions on air quality and climate, as biogenic soil fluxes influence for example the amount of aerosol and tropospheric O₃, a greenhouse gas and pollutant, in the atmosphere. Management practices of agriculture affect biogenic soil emissions. Moreover, loosing savanna to oil palm plantations or other crop would have different impacts on air quality, carbon budget and climate than the conversion of forest into crop or oil palm plantation. Furthermore, oil palm plantations are generally closer to the coast and likely to be more influenced by anthropogenic emissions from industry and coastal cities (Knippertz et al., 2015a, 2015b). Oil palm trees are also a strong isoprene emitters. Isoprene emissions influence ozone concentration and the oxidizing capacity of the atmosphere, and it is a source of secondary organic aerosol, thus affecting local air quality and global climate. Large-scale land use change in the tropics – specifically the conversion of tropical rain forest to oil palm plantations in Malaysia – were shown to cause changes in atmospheric composition and chemistry (Hewitt et al., 2009), indicating that the management of the emissions of reactive nitrogen species is essential to prevent damaging levels of ground-level ozone in those regions.

630

635

Appendix A

We ran a laboratory experiment to verify that deposition on the walls of the Teflon chamber is negligible.

Ambient air concentrations were measured by the analyzer, inside the room where the analyzer and the chamber were placed.

Measurements of NH₃ concentrations were made in ambient air with and without the Teflon chamber attached to the analyzer. The Teflon chamber was placed on a Teflon frame, and they were sealed together with Teflon tape. Measurements

of NH₃ concentrations with the Teflon chamber attached to the analyzer were followed by measurements without the chamber 30 to 60 minutes later. The two sets of measurements were made under similar conditions of temperature and humidity. Average values of NH₃ concentrations were calculated for 10 to 30 minutes before and after connecting the chamber. Average NH₃ concentrations during this time interval varied between 8 and 36 ppb, with a variation between 1.5 to 13% around the mean. The lowest NH₃ concentrations correspond to air samples previously passed through charcoal and desiccant cartridges (NO and NO₂ zero air). Measured NH₃ concentrations are reported in Table A1, along with temperature, humidity and the ratio between average concentration with and without the Teflon chamber attached to the analyzer.

This test was made at different times of the day on different days: air humidity varied between 46 and 54%, temperature varied between 25 to 29°C, while pressure varied between 1006 and 1008 hPa (not reported).

Results show negligible variation between concentrations of air reaching the analyzer via the chamber or going directly to the analyzer;

Temperature (°C)	Humidity (%)	[NH ₃] TC	[NH ₃] D	Ratio [NH ₃]TC / [NH ₃]D
25	54	26.4±0.4	25.9±0.5	1.02
27	49	9.0±0.8	8.8±1.2	1.12
29	46	35.2±2.4	36.0±1.4	0.98
28	46	25.6±0.7	24.0±1.0	1.07
27	45	26.2±0.9	27.4±0.9	0.96
28	46	23.4±0.3	24.1±0.4	0.97
26	50	19.2±0.6	18.4±0.8	1.04

Table A1 Measurements of NH₃ concentrations (ppb) through the chamber (TC) or directly (D) to the analyzer.

Acknowledgements

The DACCIWA project has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 603502. We also want thank the staffs from NCAS (National Centre for Atmospheric Science), KIT (Karlsruhe Institute of Technology) and UPS (Université Paul Sabatier, Toulouse III) for helping to install the equipment as well as the people from INRAB in Savè for allowing the equipment on their ground.

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of Ox, HOx, NOx and SOx species, *Atmos. Chem. Phys.*, 4, 1461–1738, 2004, <http://www.atmos-chem-phys.net/4/1461/2004/>.
- Bai, E., Li, S., Xu, W., Li, W., Dai, W., Jiang, P.: A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics, *New Phytol.* 199, 441–451. <http://dx.doi.org/10.1111/nph.12252>, 2013.
- Bai Junhong, Haifeng Gao, Wei Deng, Zhifeng Yang, Baoshan Cui, Rong Xiao, Nitrification potential of marsh soils from two natural saline–alkaline wetlands, *Biol Fertil Soils*, 46:525–529, 2010.
- Barthès, B., Azontonde, A., Blanchart, E., Girardin, C., Villenave, C., Lesaint, S., Oliver, R. and Feller, C.: Effect of a legume cover crop (*Mucuna pruriens* var. *utilis*) on soil carbon in an Ultisol under maize cultivation in southern Benin. *Soil Use and Manage.*, 20, 231–239, 2004.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J. and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106 (D19), 23,073 – 23,095, doi:10.1029/2001JD000807, 2001.
- Bouwman, A.F., Boumans, L.J.M., Batjes, N.H.: Emissions of N₂O and NO from fertilized fields: summary of available measurement data. *Glob. Biogeochem. Cycles* 16 (4), 1058. <http://dx.doi.org/10.1029/2001GB001811>, 2002.
- Brito, J., Freney, E., Dominutti, P., Borbon, A., Haslett, S. L., Colomb, A., Dupuy, R., Denjean, C., Burnet, F., Bourriane, T., Deroubaix, A., Sellegri, K., Coe, H., Flamant, C., Knippertz, P. and Schwarzenboeck, A., Assessing the role of anthropogenic and biogenic sources on PM₁ over Southern West Africa using aircraft measurements, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-717>, 2017.
- Brooks, B., Bessardon, G., Smith, V., Groves, J., Sharpe, S., Kalthoff, N., Adler, B., Handwerker, J., Kohler, M., Kunka, N., Tan, N., Wieser, A., Lohou, F., Bezombes, Y., Bret, G., Delon, C., Derrien, S., Dione, C., Durand, P., Etienne, P., Gabella, O., Jambert, C., Leclercq, J., Lothon, M., Medina, P., Pedruzo, X., Reinales, I., Jegede, G., Ayoola, M., Sunmonu, L., Ajao, A., Abiye, O., Ajileye, O., Aryee, J., Fosu-Amankwah, K., Cayle-Aethelhard, F., Atiah, W. A., Amekudzi, L., Danuor, S.: A new high-quality dataset of the southern West African atmospheric boundary layer diurnal cycle during the Monsoon season, *Sci. Data*, 2017, in preparation.

- Cassity-Duffey Kate, Miguel Cabrera, John Rema, Ammonia Volatilization from Broiler Litter: Effect of Soil Water Content and Humidity, *Soil Sci. Soc. Am. J.* 79:543–550, 2014.
- CILSS (2016). *Landscapes of West Africa – A Window on a Changing World*. U.S. Geological Survey, EROS, 47914 252nd St, Garretson, SD 57030, UNITED STATES, 2016.
- David, M., Loubet, B., Cellier, P., Mattsson, M., Schjoerring, J.K., Nemitz, E., Roche, R., Riedo, M., Sutton, M.A.: Ammonia sources and sinks in an intensively managed grassland canopy. *Biogeosciences* 6, 1903-1915, 2009.
- Davidson, E. A. and Kinglerlee, W.: A global inventory of nitric oxide emissions from soils, *Nutr Cycl Agroecosys*, 48: 37–50, 1997.
- 700 Delon, C., Galy-Lacaux, C., Boone, A., Lioussé, C., Serça, D., Adon, M., Diop, B., Akpo, A., Lavenu, F., Mougín, E., and Timouk, F.: Atmospheric nitrogen budget in Sahelian dry savannas, *Atmos. Chem. Phys.*, 10, 2691–2708, doi:10.5194/acp-10-2691-2010, 2010.
- Delon, C., Galy-Lacaux, C., Adon, M., Lioussé, C., Serça, D., Diop, B., and Akpo, A.: Nitrogen compounds emission and deposition in West African ecosystems: comparison between wet and dry savanna, *Biogeosciences*, 9, 385–402, doi:10.5194/bg-9-385-2012, 2012.
- 705 Delon, C., Galy-Lacaux, C., Serça, D., Loubet, B., Camara, N., Gardrat, E., Saneh, I., Fensholt, R., Tagesson, T., Le Dantec, V., Sambou, B., Diop, C., Mougín, E.: Soil and vegetation-atmosphere exchange of NO, NH₃, and N₂O from field measurements in a semi arid grazed ecosystem in Senegal, *Atmos. Environ.*, 156, 36–51, 2017.
- Derrien, S., Bezombes, Y., Bret, G., Gabella, O., Jarnot, C., Medina, P., Pique, E., Delon, C., Dione, C., Campistron, B., Durand, P., Jambert, C., Lohou, F., Lothon, M., Pacifico, F., Meyerfeld, Y.: DACCIWA field campaign, Savè super-site, UPS instrumentation; SEDOO OMP. <https://doi.org/10.6096/DACCIWA.1618>, 2016.
- Dick Jan, Ute Skiba, Robert Munro and Douglas Deans, Effect of N-fixing and non N-fixing trees and crops on NO and N₂O emissions from Senegalese soils, *Journal of Biogeography (J. Biogeogr.)*, 33, 416–423, 2006.
- EC-JRC / PBL: Emission Database for Global Atmospheric Research (EDGAR), release version 4.2. European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL), <http://edgar.jrc.ec.europa.eu>, 2011.
- 715 Ferrara, R. M., Loubet, B., Decuq, C., Palumbo, A. D., Di Tommasi, P., Magliulo, V., Masson, S., Personne, E., Cellier, P., Rana, G.: Ammonia volatilisation following urea fertilisation in an irrigated sorghum crop in Italy, *Agr. Forest Meteorol.* 195–196, 179–191, 2014.
- Flechard, C.R., Massad, R.-S., Loubet, B., Personne, E., Simpson, D., Bash, J.O., Cooter, E.J., Nemitz, E., Sutton, M.A.: Advances in understanding, models and parameterizations of biosphere-atmosphere ammonia exchange, *Biogeosciences*, 10, 5183-5225, 2013.
- 720

- Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs. *Atmos. Chem. Phys.* 15, 8217-8299. <http://dx.doi.org/10.5194/acp-15-8217-2015>, 2015.
- 725 Ganzeveld L.N., J. Lelieveld, F. J. Dentener, M. C. Krol, A. J. Bouwman, and G.-J. Roelofs, Global soil-biogenic NO_x emissions and the role of canopy processes, *JOURNAL OF GEOPHYSICAL RESEARCH*, VOL. 107, NO. D16, 10.1029/2001JD001289, 2002.
- Giglio, L., Randerson, J. T. and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res. Biogeosci.*, 118, 317–328, doi:10.1002/jgrg.20042, 2013.
- 730 Grote, R., Lehmann, E., Brümmner, C., Brüggemann, N., Szarzynski, J., Kunstmann, H.: Modelling and observation of biosphere-atmosphere interactions in natural savannah in Burkina Faso, West Africa. *Phys. Chem. Earth* 34, 251-260, 2009.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, doi:10.5194/gmd-5-1471-2012, 2012.
- 735 Jegede, O. O., Ayoola, M. A., Sunmonu, L. A., Ajao A. I., and Babić K.: DACCIWA field campaign, Ile-Ife super-site, Surface measurements; SEDOO OMP.
- Kalthoff, N., Lohou, F., Brooks, B., Jegede, G., Adler, B., Babić, K., Dione, C., Ajao, A., Amekudzi, L. K., Aryee, J. N. A., Ayoola, M., Bessardon, G., Danour, S. K., Handwerker, J., Kohler, M., Lothon, M., Pedruzo-Bagazgoitia, X., Smith, V., Sunmonu, L., Wieser, A., Fink, A. H.1, and Knippertz, P.: An overview of the diurnal cycle of the atmospheric boundary layer during the West African monsoon season: Results from the 2016 observational campaign, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-631>, 2017.
- 740 Kohler, M., Kalthoff, N., Seringer, J., and Kraut, S.: DACCIWA field campaign, Savè super-site, Surface measurements; SEDOO OMP. <https://doi.org/10.6096/DACCIWA.1690>, 2016.
- Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Liousse, C., Mari, C., Allan, R. P., Brooks, B., Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCIWA project: Dynamics-aerosol-chemistry-cloud interactions in West Africa, *B. Am. Meteorol. Soc.*, 96, 1451–1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, 2015a.
- 750 Knippertz, P., Evans, M. J., Field, P. R., Fink, A. H., Liousse, C., and Marsham, J. H.: The possible role of local air pollution in climate change in West Africa, *Nature Climate Change*, 5, 815–822, <https://doi.org/10.1038/nclimate2727>, 2015b.

- Knippertz, P., Fink, A. H., Affo-Dogo, A., Bahaga, T., Brosse, F., Deroubaix, A., Evans, M., Gaetani, M., Guebsi, R., Latifou, I., Lavaysse, C., Maranan, M., Mari, M., Marsham, J. H., Meynadier, R., Morris, E., Rosenberg, P. D., Schlueter, A., and Tocquer, F.: A meteorological and chemical overview of the DACCIWA field campaign in West Africa in June–July 2016, *Atmos. Chem. Phys. Discuss.*, doi.org/10.5194/acp-2017-345, 2017.
- Handwerker, J., Scheer, S., and Gamer, T.: DACCIWA field campaign, Savè super-site, Cloud and precipitation; SEDOO OMP. <https://doi.org/10.6096/DACCIWA.1686>, 2016.
- Hewitt, C. N., MacKenzie, A. R., Di Carlo, P., Di Marco, C. F., Dorsey, J. R., Evans, M., Fowler, D., Gallagher, M. W., Hopkins, J. R., Jones, C. E., Langford, B., Lee, J. D., Lewis, A. C., Lim, S. F., McQuaid, J., Misztal, P., Moller, S. J., Monks, P. S., Nemitz, E., Oram, D. E., Owen, S. M., Phillips, G. J., Pugh, T. A. M., Pyle, J. A., Reeves, C. E., Ryder, J., Siong, J., Skiba, U., and Stewart, D. J.: Nitrogen Management Is Essential to Prevent Tropical Oil Palm Plantations from Causing Ground-Level Ozone Pollution, *P. Natl. Acad. Sci.*, 106, 18447–18451, 2009.
- Hudman, R.C., Russell, A.R., Valin, L.C., Cohen, R.C.: Interannual variability in soil nitric oxide emissions over the United States as viewed from space. *Atmos. Chem. Phys.* 10, 9943e9952. <http://dx.doi.org/10.5194/acp-10-9943-2010>, 2010.
- 765 Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C., and Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions: implementation and space based-constraints, *Atmos. Chem. Phys.*, 12, 7779-7795, doi:10.5194/acp-12-7779-2012, 2012.
- Kristensen, L., Lenschow, D. H., Gurarie, D., Jensen, N. O.: A simple model for the reactive species in the convective atmospheric boundary layer, *Bound-Lay Meteorol.*, 134, 195-221, doi:10.1007/s10546-009-9443-x, 2010a.
- 770 Kristensen, L., Lenschow, D. H., Gurarie, D., Jensen, N. O.: Erratum to: A Simple Model for the Vertical Transport of Reactive Species in the Convective Atmospheric Boundary Layer, *Bound-Lay Meteorol.*, 135, 181–183, DOI 10.1007/s10546-010-9473-4, 2010b.
- Lata, J.-C., Degrange, V. Raynaud, X., Maron, P.-A., Lensis, R. and Abbadie, L.: Grass populations control nitrification in savanna soils, *Funct. Ecol.*, 18, 605-611, 2004.
- 775 Le Roux, X., Abbadie, L., Lensi, R., Serça, D.: Emission of nitrogen monoxide from African tropical ecosystems: control of emission by soil characteristics in humid and dry savannas of West Africa. *J. Geophys. Res.* 100, 23133-23142, 1995.
- Loubet, B., Decuq, C., Personne, E., Massad, R.S., Flechard, C., Fanucci, O., Mascher, N., Gueudet, J.-C., Masson, S., Durand, B., Genermont, S., Fauvel, Y., Cellier, P.: Investigating the stomatal, cuticular and soil ammonia fluxes over a growing tritical crop under high acidic loads. *Biogeosciences* 9, 1537-1552, 2012.
- 780 Massad, R.-S., Nemitz, E., Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere. *Atmos. Chem. Phys.* 10, 10359-10386. <http://dx.doi.org/10.5194/acp-10-10359-2010>, 2010.

- McCalley, C. K. and Sparks, J. P.: Controls over nitric oxide and ammonia emissions from Mojave Desert soils. *Oecologia* 156, 871-881, 2008.
- Meixner, F.X., Yang, W.X., In: DO doroco, P., Porporato, A. (Eds.), *Biogenic Emissions of Nitric Oxide and Ditrour Oxide from Arid and Semi-arid Land, Dryland Ecohydrology*. Kluwer Academic Publishers B.V, Dordrecht, The Netherlands, pp. 23-46, 2006.
- Michiels, B., Babatounde, S., Dahouda, M., Chabi, S. L. W. and Buldgen, A.: Botanical composition and nutritive value of forage consumed by sheep during the rainy season in a Sudano-guinean savanna (central Benin), *Trop Grasslands*, 34, 43-47, 2000.
- 790 Neftel, A., Blatter, A., Gut, A., Högger, D., Meixner, F., Ammann, C., Nathaus, F. J.: NH₃ soil and soil surface gas measurements in a triticale wheat field, *Atmos. Environ.*, 32, 3, 499-505, 1998.
- Oswald, R., Behrendt, T., Ermel, M., Wu, D., Su, H., Cheng, Y., et al.: HONO Emissions from Soil Bacteria as a Major Source of Atmospheric Reactive Nitrogen. *Science*, 341(6151), 1233-1235, 2013.
- Paillat, J, -M, , Robin, P., Hassouna, M., Leterme, P.: Predicting ammonia and carbon dioxide emissions from carbon and nitrogen biodegradability during animal waste composting, *Atmos. Environ.* 39, 6833–6842, 2005.
- 795 Pape, L., Ammann, C., Nyfeler-Brunner, A., Spirig, C., Hens, K., Meixner, F.X.: An automated dynamic chamber system for surface exchange measurement of non-reactive and reactive trace gases of grassland ecosystems. *Biogeosciences*, 6, 405e429. <http://dx.doi.org/10.5194/bg-6-405-2009>, 2009.
- Parsons, D. A. B., Scholes, M. C. and Levine, J. S.: Biogenic NO emissions from savanna soils as a function of fire regime, soil type, soil nitrogen and water status, *J Geophys Res* 101: 23683–23688, 1996.
- 800 Pilegaard, K.: Processes regulating nitric oxide emissions from soils, *Phil. Trans. R. Soc. B* 2013 (368), 20130126, 2013.
- Säïdou, A., Kuyper, T. W., Kossou, D. K., Tossou, R., and Richards, P.: Sustainable soil fertility management in Benin: learning from farmers, *NJAS-WAGEN J LIFE SC*, 52-3/4, 349-369, 2004.
- Santín C., and Doerr, S. H.: Fire effects on soils: the human dimension. *Phil. Trans. R. Soc. B* 371: 20150171. <http://dx.doi.org/10.1098/rstb.2015.0171>, 2016.
- 805 Scholes, M. C., Scholes, R. J., Parsons, D., Martin, R., and Winstead, E.: NO and N₂O emissions from savanna soils following the first rains. *Nut. Cycling Agroecosys*. 48: 115–122, 1997.
- Serça, D., Delmas, R., Le Roux, X., Parsons, D. A. B., Scholes, M. C., Abbadie, L., Lensi, R., Ronce, O., Labroue, L.: Comparison of nitrogen monoxide emissions from several African tropical ecosystems and influence of season and fire. *Glob. Biogeochem. Cy.* 12 (4), 637-651, 1998.
- 810

- Steinkamp, J., Ganzeveld, L.N., Wilcke, W., Lawrence, M.G.: Influence of modelled soil biogenic NO emissions on related trace gases and the atmospheric oxidizing efficiency. *Atmos. Chem. Phys.* 9, 2663e2677. [http:// dx.doi.org/10.5194/acp-9-2663-2009](http://dx.doi.org/10.5194/acp-9-2663-2009), 2009.
- 815 Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Butterbach Bahl, K., Cellier, C., de Vries, W., Cotrufo, F., Skiba, U., Di Marco, C., Jones, S., Laville, P., Soussana, J. F., Loubet, B., Twigg, M., Famulari, D., Whitehead, J., Gallagher, M. W., Neftel, A., Flechard, C., Herrmann, B., Calanca, P.L., Schjoerring, J. K., Daemmgen, U., Horvath, L., Tang, Y. S., Emmett, B. A., Tietema, A., Peñuelas, J., Kesik, M., Brüggemann, N., Pilegaard, K., Vesala, T., Campbell, C. L., Olesen, J. E., Dragosits, U., Theobald, M. R., Levy, P., Mobbs, D. C., Milne, R., Viovy, N., Vuichard, N., Smith, J. U., Smith, P. E., Bergamaschi, P., Fowler, D., and Reis, S.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, 820 *Environ. Pollut.*, 150, 125–139, 2007.
- Sutton, M., Reis, S., and Baker, S. M. H.: Atmospheric Ammonia, detecting emission changes and environmental impacts. Results of an expert workshop under the convention of long range transboundary air pollution, Springer Edition, New York, ISBN 978-1-4020-9120-9, 2009a.
- 825 Sutton, M. A., Nemitz, E., Milford, C., Campbell, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M., Dorsey, J. R., Gallagher, M. W., Horvath, L., Weidinger, T., Meszaros, R., Dammgén, U., Neftel, A., Herrmann, B., Lehman, B. E., Flechard, C., Burkhardt: Dynamics of ammonia exchange with cut grassland: synthesis of results and conclusions of the GRAMINAE integrated experiment, *Biogeosciences* 6, 2907-2934, 2009b.
- Tiquia, S. M. and Tam, N. F. Y.: Fate of nitrogen during composting of chicken litter, *Environ. Pollut.*, 110, 535-541, 2000.
- 830 Vaittinen O., M. Metsälä, S. Persijn, M. Vainio, L. Halonen, Adsorption of ammonia on treated stainless steel and polymer surfaces, *Appl. Phys. B*, DOI 10.1007/s00340-013-5590-3, 2013.
- Vanlauwe B., J. Diels, O. Lyasse, K. Aihou, E.N.O. Iwuafor, N. Sanginga, R. Merckx & J. Deckers, Fertility status of soils of the derived savanna and northern guinea savanna and response to major plant nutrients, as influenced by soil type and land use management, *Nutrient Cycling in Agroecosystems* 62: 139–150, 2002.
- 835 Walker, J. T., Jones, M. R., Bash, J. O., Myles, L., Meyers, T., Schwede, D., Herrick, J., Nemitz, E., and W. Robarge: Processes of ammonia air-surface exchange in a fertilized *Zea mays* canopy, *Biogeosciences*, 10, 981–998, doi:10.5194/bg-10-981-2013, 2013.
- Wentworth, G. R., Murphy, J. G., Gregoire, P. K., Cheyne, C. A. L., Tevlin, A. G. and Hems, R.: Soil–atmosphere exchange of ammonia in a non-fertilized grassland: measured emission potentials and inferred fluxes, *Biogeosciences*, 11, 5675-5686, 840 2014.

Wichink Kruit, R. J., van Pul, W. A. J., Otjes, R. P., Hofschreuder, P., Jacobs, A. F. G., and Holtslag, A. A. M, Ammonia fluxes and derived canopy compensation points over non-fertilized agricultural grassland in the Netherlands using the new gradient ammonia – high accuracy – monitor (GRAHAM), *Atmos. Environ.*, 41, 1275–1287, 2007.

845 Wieser, A., Adler, B., and Deny, B.: DACCIWA field campaign, Savè super-site, Thermodynamic data sets; SEDOO OMP. <https://doi.org/10.6096/DACCIWA.1659>, 2016.

Yienger, J. J., and H. Levy II, Global inventory of soil-biogenic NO_x emissions, *J. Geophys. Res.*, 100, 11,447– 11,464, 1995.

Savè ground-based observation site

Location	8°02'03" N, 2°29'11" E
Elevation	166 m a.s.l.
Mean annual precipitation	1100 mm
Mean annual temperature	27.5 °C
Soil type	sandy
Sand percentage	87%
Clay percentage	4.1%

850

Table 1

Main characteristics of the Savè site.

Soil type		Plant family	Plant species	Common name/s	
Next to grassland and forest	Dominant tree species	Anacardiaceae	<i>Anacardium occidentale</i>	cashew tree	
		Fabaceae	<i>Daniellia oliveri</i> <i>Pterocarpus erinaceus</i>	African copaiba balsam tree barwood, muninga, vène, mukwa	
Next to maize field	Dominant ground species	Cleomaceae	<i>Cleome</i> sp.	spider flowers, spider plants	
		Fabaceae	<i>Crotalaria</i> sp. <i>Mucuna</i> sp.	rattlepod or rattlebox velvet bean	
	Dominant tree species	Poaceae	<i>Imperata cylindrica</i> <i>Rhynchelytrum repens</i>	cogon grass, cotton wool grass, kura-kura rose natal grass	
		Anacardiaceae	<i>Mangifera indica</i>	mango	
	Crops	Dominant ground species	Areaceae	<i>Cocos nucifera</i>	coconut tree
			Caricaceae	<i>Carica papaya</i> L.	papaya
Lamiaceae			<i>Tectona grandis</i>	teak	
Meliaceae			<i>Azadirachta indica</i>	neem, nimtree, Indian lilac	
Commelinaceae			<i>Commelina benghalensis</i>	benghal dayflower, tropical spiderwort	
Euphorbiaceae			<i>Euphorbia</i> sp.	spurge	
Crops	Crops	Nyctaginaceae	<i>Boerhavia diffusa</i>	punarnava, red spiderling	
		Phyllanthaceae	<i>Phyllanthus amarus</i>	gale of the wind, stonebreaker	
		Poaceae	<i>Digitaria horizontalis</i>	Jamaican crabgrass	
		Dioscoreaceae	<i>Dioscorea</i> sp.	yam	
		Euphobiaceae	<i>Manihot esculenta</i>	cassava	
		Fabaceae	<i>Arachis hypogaea</i> <i>Vigna unguiculata</i>	peanut cowpea	
		Malvaceae	<i>Gossypium</i> sp.	cooton	
		Pedaliaceae	<i>Sesamum indicum</i>	sesame	
		Poaceae	<i>Zea mays</i> <i>Sorghum</i> sp.	maize sorghum	
		Solanaceae	<i>Solanum lycopersicum</i>	tomato	

Table 2

List of plant species at the Savè site. The list of common names is not considered to be exhaustive.

860

	Bare Soil	Grassland	Maize Field	Forest
Clay (<2µm) (%)	5.13±0.63	3.15±0.50	4.40±0.35	3.70±1.25
Fine Silt (2 to 20µm) (%)	5.13±0.96	2.93±0.32	4.13±1.00	3.40±1.21
Coarse Silt (20 to 50µm) (%)	5.98±0.51	4.78±0.66	4.37±0.38	4.67±1.05
Total Sand (50 to 2000 µm) (%)	83.75±1.82	89.20±0.71	87.13±0.99	88.20±3.50

Table 3

865 List of soil characteristics for each land cover type at the Savè site, including standard deviation.

Soil Type	Date	C/N ratio	Organic C (g kg ⁻¹)	Total N (g kg ⁻¹)
Bare soil	06/07/2016	14.90	24.28	1.63
	09/07/2016	12.20	10.47	0.86
	19/07/2016	12.30	15.05	1.22
	28/07/2016	9.50	19.30	2.04
Grassland	07/07/2016	15.20	5.78	0.38
	09/07/2016	16.00	7.36	0.46
	19/07/2016	14.90	7.16	0.48
	28/07/2016	10.90	4.56	0.42
Maize	09/07/2016	14.80	17.33	1.17
	19/07/2016	16.40	13.08	0.8
	28/07/2016	11.80	11.83	1.00
Forest	06/07/2016	14.80	7.98	0.54
	19/07/2016	11.90	9.62	0.81
	28/07/2016	11.30	16.56	1.47

870

Table 4

List of soil characteristics for each land cover type at the Savè site for each soil sampling day: Carbon-to-Nitrogen ratio (C/N), organic carbon (g kg⁻¹) and total nitrogen (g kg⁻¹). The accuracy for the C/N ratio is 14%. The measurement accuracy
875 for organic carbon and total nitrogen is 14 and 13%, respectively.



Soil Type	Date	pH	[NH ₄ ⁺] (mg kg ⁻¹)	Γ _g [NH ₄ ⁺]/[H ⁺]	X _g (ppb)
Bare soil	06/07/2016	8.38	6.82	136 334	1891
	09/07/2016	7.73	2.90	12 978	134
	19/07/2016	8.46	6.63	159 343	2188
	28/07/2016	8.34	8.01	146 033	2215
Grassland	07/07/2016	7.15	1.96	2 307	29
	09/07/2016	7.91	1.55	10 499	108
	19/07/2016	7.52	2.28	6 291	86
	28/07/2016	6.51	2.30	620	9
Maize	09/07/2016	7.46	4.40	10 575	109
	19/07/2016	7.61	14.74	50 040	687
	28/07/2016	8.04	4.49	41 027	622
Forest	06/07/2016	6.32	2.18	380	5
	19/07/2016	7.51	4.88	13 159	181
	28/07/2016	7.37	11.47	22 407	340

880

Table 5

List of soil pH, ammonium concentrations [NH₄⁺] (mg kg⁻¹), soil emission potential Γ_g and soil compensation point χ_g (ppb) for each land cover type at the Savè site for each soil sampling day. The measurement accuracy for pH is 0.15 when pH ≤ 7 and 0.20 when pH > 7. The accuracy for ammonium concentrations [NH₄⁺], soil emission potential Γ_g and soil compensation point χ_g is 25%.


885



910

915


Fig. 1(a) Location of the Savè site in West Africa, (b) one of the bare soil sampling sites, (c) the grassland sampling site, (d) the maize field sampling site and (e) the forest sampling site at the Savè site.



920

Fig. 2 Upper panel: Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, **daily mean NO fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ measured at the bare soil site**; Middle panel: **daily mean NO and daily mean NH₃ fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ measured at the bare soil site**; Lower panel: daily mean NO and NH₃ concentrations in ppb **measured at the bare soil site**. Vertical bars show the standard deviation from individual fluxes and concentrations


925




930

Fig. 3 Upper panel: Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, daily mean NO fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ measured at the grassland site; Middle panel: daily mean ~~NO~~ and NH_3 fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ measured at the grassland site; Lower panel: daily mean NO and NH_3 concentrations in ppb measured at the grassland site. Vertical bars show the standard deviation from individual fluxes and concentrations.

935




940 Fig. 4 Upper panel: Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, daily mean NO fluxes in ngN m⁻² s⁻¹ measured at the maize field site; Middle panel: daily mean ~~NO~~ and NH₃ fluxes in ngN m⁻² s⁻¹ measured at the maize field site; Lower panel: daily mean NO and NH₃ concentrations in ppb measured at the maize field site. Vertical bars show the standard deviation from individual fluxes and concentrations.



945

Fig. 5 Upper panel: Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, daily mean NO fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ measured at the forest site; Middle panel: daily mean ~~NO and~~ NH_3 fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ ~~measured at the forest site~~; Lower panel: daily mean NO and NH_3 concentrations in ppb ~~measured at the forest site~~. Vertical bars show the standard deviation from individual fluxes and concentrations.

950



955 Fig. 6 Nitric oxide emissions in $\text{ngN m}^{-2} \text{s}^{-1}$ measured over each land cover type (orange dot for bare soil, green for grassland, yellow for the maize field and brown for forest) and simulated with GEOS-Chem, along with rainfall measured and modelled with GEOS-Chem. Soil NO emissions are daily average between 8 a.m. and 6 p.m..




Fig. 7 (a) Land cover map of Benin for 2013 from the US Geographical Survey Atlas: Landscapes of West Africa – A Window on a Changing World (CILSS, 2016) and (b) land cover map of Benin used in the GEOS-Chem simulation.