



Measurements of nitric oxide and ammonia soil fluxes from a wet savanna ecosystem site in West Africa during the DACCIWA field campaign.

Federica Pacifico¹, Claire Delon¹, Corinne Jambert¹, Pierre Durand¹, Eleanor Morris², Mat J. Evans²,
5 Fabienne Lohou¹, Solène Derrien¹, Venance H. E. Donnou³, Arnaud V. Houeto³, Irene Reinares Martínez¹, Pierre-Etienne Brilouet¹

¹Laboratoire d'Aérodologie, University of Toulouse, CNRS, UPS, Toulouse, 31400, France

²Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, YO10 5DD, UK

10 ³ Laboratoire de Physique du Rayonnement, Université d'Abomey-Calavi, Cotonou, 01 BP 526, Benin

15 *Correspondence to:* Claire Delon (claire.delon@aero.obs-mip.fr)

Abstract. It is important to correctly simulate biogenic fluxes from soil in atmospheric chemistry models at a local and regional scale to study air pollution and climate in an area of the world, West Africa, that has been subject to a strong increase in anthropogenic emissions due to a massive growth in population and urbanization. Anthropogenic pollutants are transported inland and northward from the mega cities located on the coast, where the reaction with biogenic emissions may
20 lead to enhanced ozone production outside urban areas, as well as secondary organic aerosols formation, with detrimental effects on humans, animals, natural vegetation and crops.

Here we present field measurements of soil fluxes of nitric oxide (NO) and ammonia (NH₃) observed over four different land cover types, i.e. bare soil, grassland, maize field and forest, at an inland rural site in Benin, West Africa, during the
25 DACCIWA field campaign in June and July 2016.

We observe NO fluxes up to 48.05 ngN m⁻² s⁻¹. NO fluxes averaged over all land cover types are 4.79 ± 5.59 ngN m⁻² s⁻¹, maximum soil emissions of NO are recorded over bare soil. NH₃ is dominated by deposition for all land cover types. NH₃ fluxes range between -6.59 and 4.96 ngN m⁻² s⁻¹. NH₃ fluxes averaged over all land cover types are -0.91 ± 1.27 ngN m⁻² s⁻¹
30 and maximum NH₃ deposition is measured over bare soil. The observations show high spatial variability even for the same soil type, same day and same meteorological conditions.



35 We compare point daily average measurements of NO emissions recorded during the field campaign with those simulated by GEOS-Chem (Goddard Earth Observing System Chemistry Model) for the same site and find good agreement. In an attempt to quantify NO emissions at the regional and national scale, we also provide a tentative estimate of total NO emissions for the entire country of Benin for the month of July using two distinct methods: upscaling point measurements and using the GEOS-Chem model. The two methods give similar results: 1.17 ± 0.6 GgN/month and 1.44 GgN/month, respectively. Total NH₃ deposition estimated by upscaling point measurements for the month of July is 0.21 GgN/month.



1 Introduction

40 Biogenic soil fluxes of nitric oxide (NO) and ammonia (NH₃) play an important role on tropospheric chemistry. Nitric Oxide emitted by soil influences the concentration of nitrogen oxides (NO_x) in the atmosphere, consequently modifying the rates of ozone (O₃) production, where O₃ is a pollutant, harmful to humans and plants, and also a greenhouse gas (Steinkamp et al., 2009). The production and consumption of NO in soil is regulated by microbial activity, mainly nitrification/denitrification processes, and chemical reactions (Pilegaard et al., 2013). Measurements using soil chambers in the field and laboratory
45 experiments show that nitrification/denitrification, and consequently NO emissions, vary greatly with climate and soil conditions, in particular they are strongly correlated with nitrogen (N) availability, temperature and soil moisture, making soil NO emissions dependent on regional temperature and precipitation patterns, and fertilizer management practices (e.g., Bouwman et al., 2002; Meixner and Yang, 2006; Hudman et al., 2010).

50 Soil NO emissions are about 20% of total NO sources to the atmosphere (Oswald et al., 2013) and almost of the same order of magnitude of fossil fuel NO emissions. Soil emission of biogenic NO plays a prominent role in the regional atmospheric chemistry of non-urbanized areas, where anthropogenic emissions are negligible (Pilegaard, 2013). The main inputs of N compounds onto semi arid uncultivated soils, like the savanna ecosystem, are biological nitrogen fixation, atmospheric wet and dry deposition and lightning. NO fluxes are considered as one way only, even if NO deposition exists in very specific
55 conditions (Grote et al., 2009).

Soil N losses towards the atmosphere also involve NH₃. The largest source of NH₃ emissions is agriculture, via the application of synthetic fertilizer. When released into the atmosphere, NH₃ increases the level of air pollution. In the atmosphere NH₃ has a relatively short life time of less than five days and high deposition rates, it is converted into
60 ammonium (NH₄⁺) aerosols, which has a life time of the order of fifteen days, can travel long distances and it is relevant for air quality and climate (Fuzzi et al., 2015). The exchange of soil NH₃ is bi-directional as it also includes deposition. NH₃ returned to the surface by deposition can potentially cause eutrophication, reducing biodiversity and water quality (Sutton et al., 2009a).

65 The net flux of NH₃ is the combination of different exchange pathways between plant (cuticle and stomata), soil, leaf litter and atmosphere. The overall NH₃ flux for a given surface may switch from net emission to net deposition at sub-hourly, diurnal and seasonal scales. Moreover, NH₃ can be rapidly deposited onto cuticles due to its high solubility (e.g. Sutton et al., 2009b; Massad et al., 2010; Loubet et al., 2012).



70 The direction and magnitude of NH_3 exchanges depend on the difference in NH_3 concentration between the canopy and the
atmosphere, and on a large range of environmental factors, in particular air humidity, which influence surface wetness, and
soil moisture conditions, but also vegetation cover and soil characteristics. The relationships between NO and NH_3 soil
fluxes have been identified through the ammonium content in the soil (McCalley and Sparks, 2008). Ammonia is mainly
emitted by agricultural activities, and also by the decomposition of litter and volatilization of animal excreta (Sutton et al.,
75 2009b; Massad et al., 2010).

Soil fluxes in West Africa have only been measured in a limited number of studies due to the challenging experimental
conditions (remote sites, no power supply, very hot temperatures), and mainly with manual chamber techniques rather than
more complex micrometeorological techniques (Serça et al., 1998, Le Roux et al. 1995 for NO, Delon et al., 2017 for NO
80 and NH_3). However, tropical savanna has been recognized as one of the ecosystems characterized by the largest NO
emissions (Davidson and Kinglerlee, 1997, Hudman et al., 2012).

Anthropogenic emissions of pollutants from mega cities located on the Guinean coast in South West Africa have been
increasing, and are likely to keep increasing in the next decades, due to a strong anthropogenic pressure, land use change and
85 urbanization. When transported northward on the African continent, polluted air masses meet biogenic emissions from rural
areas which contributes to increased O_3 and secondary organic aerosol production, in high temperature and solar radiation
conditions, highly favorable to enhance photochemistry (Knippertz et al., 2015a, 2015b).

The objectives of this study are to quantify soil fluxes of NO and NH_3 for the different land cover types typical of rural West
90 Africa, suggest a tentative strategy to scale point measurements in the field to ecosystem and larger regional scale, and
provide data for inventories and model evaluation to improve air quality and climate modelling.

In this paper we present the soil fluxes of NO and NH_3 measured in a rural site near the city of Savè, Benin, West Africa,
during the DACCIWA (Dynamics-Aerosol-Chemistry-Cloud Interactions in West Africa) field campaign which lasted from
95 14th June to 30th July 2016 (wet season). The DACCIWA campaign was lead to investigate the possible role of local air
pollution on climate change in West Africa, focusing on atmospheric composition, air pollution and cloud-aerosol
interactions over several sites in the region (Knippertz et al., 2015a, 2015b, 2017). The Savè site is part of the savanna
ecosystem, where grassland is intercut with crops and degraded forest. Biogenic soil fluxes measurements were taken using
the manual chamber technique, which is robust and of reduced costs (Delon et al., 2017). Along with these observations we
100 also present measurements of soil characteristics and meteorological variables from the same site. We include the
comparison of measured NO soil emissions with those simulated by the Hudman et al. (2012) process-based model for NO
soil emission implemented into GEOS-Chem.



2 Material and method

2.1 Site description

105 The Savè site for ground-based observations is located in a hinterland area of Benin, 6 km south west from the city of Savè (8°02'03" N, 2°29'11" E, 166 m a.s.l.). The Savè ground-based observation site is located within the Gobè site managed by the Institut National des Recherches Agricoles du Bénin (INRAB).

The site is characterized by a wet savanna ecosystem. The climate of the region is Sudano-guinean, with a rainy season from March to October and a dry season from November to February (Michels et al., 2000). The average annual rainfall is about 1100 mm (Savè weather station, data averaged from 1969 to 2004, Michels et al., 2000 and Sâidou et al., 2004) and the average yearly temperature is about 27.5 °C with little variation from year to year (data averaged from 1984 to 2004, Sâidou et al., 2004). Average minimum temperature, based on 1969-1990 data, is 21.5 °C and mean maximum temperature is 35.5 °C.

115 The tree coverage in the Savè region is low with most of the land occupied by subsistence agriculture and grassland (CILSS, 2016). Four land cover types are identified at the observation site: bare soil, grassland, maize field and degraded forest. Bare soil is defined as a patch of land of minimum five by five meters wide, without vegetation growing or hanging over the plot. Ground photographs of the four land cover types are shown in Fig.1.

120 The most abundant tree species next to the grassland site and in the forest are: *Anacardium occidentale*, *Daniellia oliveri*, and *Pterocarpus erinaceus*; while the most abundant tree species next the maize field are: *Mangifera indica*, *Cocos nucifera*, *Carica papaya L.*, *Tectona grandis*, and *Azadirachta indica*. The herbaceous vegetation is dominated by *Cleome* sp., *Crotalaria* sp., *Mucuna* sp., *Imperata cylindrica* and *Rhynchelytrum repens* next to grassland site and in the forest, and 125 *Commelina benghalensis*, *Euphorbia* sp., *Boerhavia diffusa*, *Phyllanthus amarus*, *Digitaria horizontalis* by maize field. In the maize field, the main species, *Zea Mays*, is intercropped with *Sesamum indicum* and, to a lesser extent, with other species: *Dioscorea* sp., *Manihot esculenta*, *Arachis hypogaea*, *Vigna unguiculata*, *Gossypium* sp., *Sorghum* sp. and *Solanum lycopersicum*. The maize field was not treated with mineral fertilizer. The only livestock consist of a few dozens of domestic fowls belonging to small subsistence-oriented family farms, mainly grazing in the maize field.

130 At the Savè site, the soil is sandy, with 87% of sand and 4.1% of clay (the rest being silt) for the 0-5 cm horizon. Surface pH ranges from 6.32 to 8.46, depending on the place where the measurement is done. Mean meteorological and average soil characteristics for the observation site are reported in Table 1, dominant vegetation species and soil composition for each land cover type are given in Table 2 and 3, respectively. Sunrise and sunset UTC time at the beginning and at the end of the 135 campaign are: 05:33 and 18:08 on 14th June, and 05:42 and 18:11 on 30th July.



2.2 Sampling sites

The samples were taken at the four land cover types (bare soil, grassland, maize field and forest), one location per day. Two to three samplings spots were chosen each day for each location, collecting eight to twenty-five flux measurements for both NO and NH₃ soil fluxes each day. Each location was sampled during daytime, approximately from 7 a.m. to 6 p.m., alternating measurements at the four different land cover types from one day to the other, over the entire campaign. Bare soil and the maize field were sampled for both NO and NH₃ soil fluxes on eight different, generally non-consecutive, days, grassland on ten days, and the forest site on four different days.

2.3 Chamber flux measurements

The technique used to measure NO and NH₃ soil fluxes makes use of a ThermoScientific 17i (ThermoFischer Scientific, MA, USA) to measure the concentration of NO and NH₃ through chemiluminescence, and the closed dynamic chamber technique to calculate fluxes. The details of this technique are fully described in Delon et al. (2017).

The remoteness of the study site limited installation of permanent structures and we were unable to automate our chamber measurements, thus all measurements were made manually. The instrument was powered by a generator (>100 m away) and carried around on a wheeled-table to reach the locations of the four soil types where the NO and NH₃ soil fluxes were measured. The analyzer was connected via a Teflon tube to the Teflon chamber which was put on the ground to detect the fluxes. The external sides of the chamber were covered with sand or soil to isolate it during the measurement. The soil under the chamber was left unperturbed. Adjustments were taken in order to make sure the analyzer did not reach temperatures that would invalidate the measurements.

The calibration of the NO sensor of the 17i analyzer was made before and after the campaign, with a reference NO₂ air mixture, i.e. NO in N₂ diluted with zero air. Two post-campaign calibrations were made: a first one to validate the efficiency of the NO₂ converter using a reference dilution of NO₂ in zero air, and a second one to validate the efficiency of the NH₃ converter with a NH₃/N₂ mixture. The zero air was obtained by filtering ambient air, previously passed on charcoal and desiccant cartridges. The dilution for all the calibration experiments was made with the 146i module (ThermoFischer Scientific, MA, USA) and the dilution module, equipped with certified mass flow meters, on board of the ATR-42 research aircraft during an inter-calibration with other NO_x instrumentation of the DACCIWA campaign (i.e. the instrumentation on the Savè measurement site tower and the instrumentation on the ATR-42 aircraft; Brito et al., 2017, Derrien et al., 2016). Reference NO, and NO₂ were ISO 6141:2015 certified at 8.73 and 8.58 ppm for NO, before and after the campaign, respectively, and 9.28 ppm for NO₂, both with 5% precision. Reference NH₃ mixture was certified at 14.78 ppm with 2% precision for NH₃. Multipoint (at least 4 points) calibrations between 50 to 250 ppb were done to ensure the linearity of the



response, obtaining regression coefficients over 0.9993 for both NO and NO₂. The global precision of the analyzer is ±0.4 ppb.

- 170 The air inlet is located on one side of the chamber, where a small vent of 4 mm in diameter provided the pressure equilibrium between the inside and outside of the chamber. The air outlet on the other side is connected to the analyzer with a 4 m Teflon tube. The chamber is continuously swept with an air flow of 0.7 L min⁻¹ insured by the instrument pump, and the air flow is controlled inside the analyzer by a flow meter.
- 175 The opaque walls minimize photochemical reactions inside the chamber, which are therefore considered as negligible. The chamber is placed on the soil for 10 min. After 10 min, the chamber is turned over to let the analyzer be swept by ambient air for 5 min, then the chamber is placed again on the soil to begin a new cycle.

- The calculation of the fluxes are based on the closed dynamic chamber technique, with the following assumptions: the
- 180 concentration in the chamber is equal to the concentration leaving the chamber to the analyzer, no deposition occurs onto the Teflon walls chamber; chemical reactions in the gas phase inside the chamber are limited, thanks to the opaque walls. All the details of the calculation are given in Delon et al. (2017). In brief:

$$F_x = \frac{V}{A_0} \frac{\delta C_x}{\delta t} \quad (1)$$

185

Where F_x is the flux (NO or NH₃) in nmol m⁻² s⁻¹, δC_x is the concentration increase in the chamber in nmol m⁻³ during the temporal interval δt . $A_0=0.0684$ m² is the surface of the ground covered by the chamber, $V=0.0123$ m³ is the volume of the chamber. This equation is similar to the one in Davidson et al. (1990). The flux is then converted to ngN m⁻² s⁻¹.

- 190 The linear regression is calculated over a 120 to 360 s time interval, about 120s after the installation of the chamber on soil for NO, and 180-300s after the installation of the chamber on soil for NH₃. Based on the methodology developed in Delon et al. (2017), the dilution effect due to mixing of outside air in the chamber is 6.6% for NO and 7.6% for NH₃. Considering the precision of the analyzer (±0.4 ppbv), the detection limit is 0.2 ngN m⁻² s⁻¹ for NO and 0.3 ngN m⁻² s⁻¹ for NH₃ fluxes.
- 195 The chemical reactions inside the chamber can determine NO consumption, and consequently an underestimation of the NO fluxes calculated with our method. This underestimation is taken into account and calculated following the method by Pape et al. (2009) with the relation $k \cdot [\text{NO}] \cdot [\text{O}_3]$. In this relation k is the temperature-dependent reaction rate constant (Pape et al., 2009, Atkinson et al., 2004), [NO] is measured by the ThermoScientific 17i at soil level just before positioning the chamber



for the measurement of soil fluxes, and $[O_3]$ at soil level is derived by measurements of NO and NO_2 at soil level made with
200 the ThermoScientific 17i and measurements of NO, NO_2 and O_3 taken on an 8 m high tower. On the 8 m high tower, NO and
 NO_2 were measured with a Model 42C TraceLevel NO- NO_2 -NO $_x$ by Thermo-environmental Instruments Inc., calibrated
with the same method as the ThermoScientific 17i and with 0.05 ppb detection limit. Ozone was measured on the tower with
a Model 49i Ozone Analyzer by Thermo-environmental Instruments Inc. with 1 ppb detection limit. The Model 49i Ozone
Analyzer was calibrated by comparison with a Thermo Scientific Model 49PS reference instrument. The reference
205 instrument is sent twice a year to the French Laboratoire national d'Essais (LNE) for comparison with a National Institute of
Standards and technology (NIST). All data on the tower were sampled at 10 seconds. $[O_3]$ at soil level was then calculated
considering the diurnal steady state of the reactions described in equation (2) and (3), using equation (4):



$$[O_3]_{sl} = \frac{[NO]_{tl} [O_3]_{tl} [NO_2]_{sl}}{[NO_2]_{tl} [NO]_{sl}} \quad (4)$$

Where $[]_{sl}$ is the concentration at the soil level and $[]_{tl}$ is the concentration measured on the tower. In conclusion, we correct
NO fluxes for the underestimation of NO fluxes due to chemical reactions inside the chamber with values ranging between 0
215 and 63% (8% on average for the whole campaign).

As studied by Kristensen et al. (2010a) and Kristensen et al. (2010b), O_3 deposition can decrease O_3 concentration close to
soil surface further. However, considering that O_3 concentrations calculated near the soil are already very low (1 ppb at soil
level compared to 24 ppb at 8 m, averaged for the entire measurement campaign), O_3 deposition has been considered of
220 secondary importance in this calculation and has not been included. If O_3 deposition were to be included it would possibly
decrease the correction of NO fluxes and consequently slightly decrease NO emissions in a negligible proportion compared
to the correction already applied for the chemical reactions inside the chamber.

2.4 Data quality check

A quality check method based on the following criteria is used to select observed fluxes (Delon et al., 2017):

225 - The coefficient of determination for linear regression R^2 has to be higher than 0.4 (considered as a significant correlation)
for NH_3 fluxes, and higher than 0.8 for NO fluxes.

- The concentration difference between the last and the first NH_3 measurement point has to be more than 0.4 ppb (sensitivity
of the analyzer). R^2 was generally lower than 0.4 for concentration differences below 0.4 ppb.

Finally, 351/488 (72%) NH_3 flux measurements and 459/488 (94%) NO flux measurements are considered valid.



230 2.5 Meteorological station

Continuous in situ observations of meteorological variables, including air and soil temperature and moisture, rainfall, wind speed, wind direction, radiation and energy balance components were taken at the Savè site as part of the DACCIWA campaign. Data are provided as 1 min averages, apart from energy fluxes which are given as 30 min averages (Derrien et al., 2016, Kohler et al., 2016, Handwerker et al., 2016, Wieser et al., 2016). An overview of the complete set of instrumentation and measurements is given by Brooks et al. (2017), while a summary of the available ground-based meteorological observations is given by Kalthoff et al. (2017). In this study we present soil moisture measured in two distinct locations of the Savè site by the Karlsruhe Institute of Technology (KIT) instrumentation at 5cm depth on grassland, and average soil moisture, between 0 and 30 cm, measured by the Université Paul Sabatier (UPS) instrumentation in the maize field. Details of the instrumentation is given by Brooks et al. (2017). We include soil moisture measured with both systems as the inter-comparison of the two methods is out of the scope of this study.

2.6 Soil characteristics (texture, pH, N content)

Soil samples were collected during the measurement campaign to analyze the biogeochemical characteristics of the site. Soil samples (0-5 cm) were taken for each land cover type where NO and NH₃ fluxes were measured. Three to four samples were collected for each land cover type on different days during the campaign.

Samples were dried in ambient conditions (mean day-time temperature is approximately 26 °C, Kalthoff et al., 2017), and stored in the dark for the determination of texture, ammonium concentrations [NH₄⁺], C/N ratio, total C, total N, and pH at the GALYS Laboratoire (<http://www.galys-laboratoire.fr>, NF EN ISO/CEI 17025: 2005). The analyses were performed two months after sampling. We assume that the ammonium content in litter or soils is not modified by volatilization or chemical transformation during transport and storage, because of the very low soil moisture level in samples. Past studies have demonstrated that microbial activity and mineralization processes are inhibited in low soil moisture conditions, even when soil temperature is high (Bai et al., 2013, and references therein). Soil texture is determined following norm NF X 31e107. Clay (<2 µm), fine silt (2 to 20 µm), coarse silt (20 to 50 µm) and total sand (50 to 2000 µm) are determined without decarbonation. Organic carbon and total nitrogen are determined following norm NF ISO 10694. The whole carbon of the sample is transformed into CO₂. Then CO₂ is measured by thermal conductivity. Mineral and organic nitrogen are determined following norm NF ISO 13878. The sample is heated at 1000°C with O₂. Products of combustion or decomposition are reduced in N₂. N₂ is then measured by thermal conductivity (catharometer). pH is determined according to norm NF ISO 10390, with soil samples stirred with water (ratio 1/5).



2.7 Soil ammonia emission potential Γ_g and compensation point χ_g

- 260 Measurements of soil pH and ammonium concentrations $[\text{NH}_4^+]$ are used to quantify the soil emission potentials for the different land cover types at the measurement site. The soil emission potential Γ_g is the ratio of $[\text{NH}_4^+]$ to $[\text{H}^+]$ concentrations in the water solution of the soil (mol L^{-1}). A large Γ_g indicates that the soil has a high propensity to emit NH_3 , considering that the potential emission of NH_3 depends on the availability of ammonium in the soil and on pH.
- 265 The soil compensation point (χ_g) has been calculated from the emission potential Γ_g , as a function of soil surface temperature (T_g in K) according to Wentworth et al. (2014):

$$\chi_g \text{ (ppb)} = 13\,587 \cdot \Gamma_g \cdot e^{-(10\,396\text{K}/T_g)} \times 10^9, \quad (5)$$

- 270 The soil compensation point indicates the equilibrium between gaseous NH_3 in the soil pore space and $[\text{NH}_4^+]$ in the soil solution, i.e. the concentration of NH_3 for which the NH_3 flux switches from emission to deposition (or viceversa).

2.8 Stepwise multiple regression analysis

- A stepwise linear multiple regression analysis was performed between daily averaged gas fluxes of NO and NH_3 and relevant available daily averaged variables such as wind speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, outgoing longwave radiation and incoming shortwave radiation. Soil parameters such as mineral nitrogen, total N and organic C, soil texture and pH could not be used for this regression analysis since their relative measurements did not have the same temporal resolution as the other parameters. The R software (<http://www.R-project.org>) was used to provide the results of this linear regression analysis.
- 275

2.9 GEOS-Chem

- 280 GEOS-Chem is a global three-dimensional model of tropospheric chemistry driven by meteorological input from the NASA Goddard Earth Observing System (www.geos-chem.org, Bey et al., 2001). In this study we use GEOS-Chem Version 10-01 which includes the process-based parameterization of soil NO emission by Hudman et al. (2012). This parameterization represents available nitrogen (N) in soils using biome specific emission factors, online wet and dry deposition of N, and fertilizer and manure N derived from a spatially explicit dataset, distributed using seasonality derived from data obtained by the Moderate Resolution Imaging Spectrometer (MODIS). Emissions are a smooth function of soil moisture and temperature consistent with point measurements and ecosystem scale experiments. This parameterization also included pulsing following soil wetting by rain or irrigation, represented as a function dependent on dry spell length. The parameterization by Hudman et al. (2012) was successfully evaluated for pulsing events in central Sahel ($0\text{--}30^\circ \text{W}$, $12\text{--}18^\circ \text{N}$).
- 285



290 Boundary conditions for our experiment are generated from a global GEOS-Chem simulation at $4^\circ \times 5^\circ$ horizontal resolution. The regional GEOS-Chem model for West Africa runs at a horizontal resolution of $0.25^\circ \times 0.3125^\circ$ (latitudes 6°S – 16°N , longitudes 18.125°W – 26.875°E) and a vertical resolution of 47 levels (up to 0.01hPa). Meteorology is driven by the NASA GMAO (Global Modeling and Assimilation Office) GEOS-FP (Forward Processing) assimilated meteorological data. The global model is spun-up from 1st May 2015 to 1st May 2016. The global simulation is then run from 1st May 2016 to
295 1st August 2016, outputting boundary condition files for West Africa. The regional West Africa simulation is then run from 1st May 2016 to 1st August 2016 using the $4^\circ \times 5^\circ$ boundary conditions from the global simulation. All simulations use the GEOS-FP meteorology which has a three-hour time resolution. We used the same MODIS/Koppen land cover map as in Hudman et al. (2012; <http://glcf.umd.edu/data/lc>) which includes 24 land cover types. In this simulation we use EDGAR v4.2 (EC-JRC/PBL, 2011) for anthropogenic emissions, GFED4 (Giglio et al., 2013) for biomass burning emissions and
300 MEGAN v2.1 (Guenther et al., 2012) for biogenic emissions of volatile organic compounds. The same emission inventories are used for both the boundary conditions and the West Africa simulation.

3 Results and discussion

3.1 Meteorological data

Mean air temperature averaged over the whole campaign is 25.4 ± 2.6 °C, mean wind speed is 1.3 ± 0.6 m s⁻¹, mean relative
305 air humidity is 86.3 ± 10.5 %, mean soil temperature is 25.2 ± 3.4 °C, mean KIT soil moisture at 5 cm is 7.1 ± 3.6 %, while mean UPS soil moisture averaged between 0 and 30 cm is 4.5 ± 2.8 %.

Median diurnal cycles of air temperature, specific humidity and precipitation are reported in Kalthoff et al., (2017). Knippertz et al. (2017) distinguish four different phases of the monsoon season during the DACCIWA campaign (14th June
310 to 30th July 2016) over the DACCIWA focus region (5 – 10° N, 8° W - 8° E), which covers a wide area of West Africa (see Fig. 1, Knippertz et al. (2017)). The division into phases is mainly based on the north–south precipitation difference between the coastal zone (0 – 7.5° N) and the Sudanian–Sahelian zone (7.5 – 15° N), both averaged across the longitude range 8° W– 8° E. Savè (8.03° N) is located very close to the border between the two zones, with a rainfall pattern that seems to follow more closely that of the coastal zone rather than that of the northern inland Sudanian–Sahelian zone. These four phases are: the
315 pre-onset phase characterized by a rainfall maximum near the coast (before 21st June, phase 1); the post-onset phase during which the rainfall maximum occurred inland (22nd June - 20th July, phase 2); the wet westerly regime when the rainfall maximum shifted back to the coast (21st – 26th July, phase 3); and the recovery of the monsoon with a shift of the rainfall maximum inland (27th July until the end of the campaign, phase 4). A specific period within phase 2 is indicated “vortex”, during which an unusual development occurred (09th – 16th July): in the north, a cyclonic feature slowly propagated from
320 eastern Mali to Cape Verde and in the south, an anticyclonic vortex tracked in the west-northwesterly direction along the Guinean coast (see Knippertz et al. 2017 for a more detailed description). At the Savè site the most intense rainfall events



happened the day before the first soil fluxes observation, on 15th June 2016, and towards the end of the measurement campaign between 20th and 23rd July 2016. Other minor rainfall events are recorded on 19th and 27th June, 8th, 12th, 13th, 24th and 26th July. Daily rainfall measurements are reported in Figs. 2 to 5.

325 3.2 Soil texture, soil organic carbon, total nitrogen, pH and ammonium content

Bare soil recorded the lower amount of total sand (83.75 ± 1.82 %) and the higher amount of clay (5.13 ± 0.63 %), fine (5.13 ± 0.96 %) and coarse silt (5.98 ± 0.51 %). Grassland recorded the higher amount of total sand (89.20 ± 0.71 %) and the lower amount of clay (3.15 ± 0.50 %) and fine silt (2.93 ± 0.32 %), while intermediate values were found for the maize field and forest (Table 3). These values determine the classification of sandy soil for all measurements sites.

330

Soil organic carbon (C) and total nitrogen (N) are respectively 12.2 ± 5.7 g kg⁻¹ and 0.95 ± 0.51 g kg⁻¹, averaged for all land cover types over the entire campaign. Table 4 gives soil characteristics for each land cover types, including individual values of C/N ratio, soil organic C and total N for the entire field campaign. The highest average soil organic C was measured for bare soil (17.3 ± 5.9 g kg⁻¹) and the lowest soil organic C was measured for grassland (6.2 ± 1.3 g kg⁻¹), while the maize field and forest site accounted for 14.1 ± 2.9 g kg⁻¹ and 11.4 ± 4.5 g kg⁻¹ soil organic C, respectively. The highest average total N was measured for bare soil (1.44 ± 0.51 g kg⁻¹) and the lowest total N was measured for grassland (0.44 ± 0.04 g kg⁻¹), while the maize field and forest site accounted similar amounts of total N, 0.99 ± 0.19 g kg⁻¹ and 0.94 ± 0.48 g kg⁻¹, respectively. Values of C/N, soil organic C and total N recorded for grassland at the Savè site compare closely to those reported by Delon et al. (2017, table 2) for the semi-arid site of Dahra (15°24' N 15°25' W), Senegal. Our values of C/N and total N for grassland are also close to those reported by Le Roux et al. (1995, table 1) and Lata et al. (2004) for the wet savanna ecosystem of Lamto (6°13' N, 5°20' W), Ivory Coast, although we observe lower values of soil organic C compared to Le Roux et al. (1995) and Lata et al. (2004). Values of C/N and soil organic C recorded for the maize field at the Savè site are slightly higher than those recorded by Barthes et al. (2004) in a maize field at Agonkanmey (6°24' N, 2°20' E), near Cotonou in southern Benin.

345

All the sites listed in the comparison in the previous paragraph are sandy, as the Savè site. The Dahra site (Delon et al., 2017) also shows similar pH than our site (Table 5), while lower pH (acidic or near-neutral) was recorded at the sites of Lamto (Le Roux et al., 1995, Lata et al., 2004) and Agonkanmey (Barthes et al., 2004). Table 5 provides individual values of pH, [NH₄⁺], Γ_g and χ_g for the entire field campaign. The highest average pH was observed for bare soil (8.23) and the lowest for the forest site (7.07), while measured average pH was 7.27 for grassland and 7.70 for the maize field. The [NH₄⁺] content averaged for all land cover types over the entire campaign is 5.33 ± 4 mg kg⁻¹. The highest average [NH₄⁺] was recorded for the maize field (7.9 ± 6 mg kg⁻¹) and the lowest for grassland (2.0 ± 0.3 mg kg⁻¹). Average [NH₄⁺] is 6.2 ± 5 mg kg⁻¹ and 7.0 ± 2.2 mg kg⁻¹ for forest and bare soil, respectively.

350



355 Higher soil organic C and N over bare soil could be due to the fact that these bare soil patches experienced recent burning (Santín and Doerr, 2016). The higher $[\text{NH}_4^+]$ over the maize field can be caused by chicken excreta, as chickens were roaming over the maize field (Paillat et al., 2005, Tiquia and Tam, 2000).

3.3 Soil emission potential Γ_g and compensation point χ_g

The mean soil emission potentials for the Savè site is $43\,714 \pm 58\,077$, with values ranging from 380 to 159 343. The highest values of soil emission potentials are observed for bare soil ($113\,672 \pm 67\,788$), followed by maize field ($33\,880 \pm 20\,680$), forest ($11\,982 \pm 11\,061$) and grassland ($4\,929 \pm 4\,409$). The ammonia compensation point ranges between 5 to 2 215 ppb, with soil temperatures between 25 and 29 °C. The highest values of χ_g are observed for bare soil ($1\,607 \pm 993$), followed by maize field (473 ± 317), forest (175 ± 167) and grassland (58 ± 47). Our values of soil emission potential for bare soil and maize (no fertilization) are comparable with those presented in Massad et al. (2010, table 4), although those data come from measurements taken on different ecosystems. Both Γ_g and χ_g values recorded at the Savè site exceed those recorded by Delon et al. (2017) over a grazed semi-arid Sahelian ecosystem in Senegal.

3.4 NO fluxes

NO fluxes from soil measured during the field campaign range between 0 and $48.05 \text{ ngN m}^{-2} \text{ s}^{-1}$. NO fluxes averaged over all land cover types are $4.79 \pm 5.59 \text{ ngN m}^{-2} \text{ s}^{-1}$, while average NO fluxes for each land cover type are: $8.05 \pm 3.49 \text{ ngN m}^{-2} \text{ s}^{-1}$ for bare soil, $3.73 \pm 1.76 \text{ ngN m}^{-2} \text{ s}^{-1}$ for the maize field, $2.87 \pm 1.49 \text{ ngN m}^{-2} \text{ s}^{-1}$ for forest and $2.82 \pm 3.46 \text{ ngN m}^{-2} \text{ s}^{-1}$ for grassland. Soil emissions of NO from the different land cover types provide similar values, NO emissions from bare soil are higher on average, but have a larger standard deviation.

Other measurements of biogenic NO soil emissions from the West African wet savanna can be found in Delon et al. (2012, table 7). We find that our measured NO soil emissions averaged over all land cover types are higher than those measured from other wet savanna sites. Our measurements are in better agreement with emissions from dry savanna grasslands (Delon et al., 2012), and with measurements from a semi-arid savanna, with over 80% sandy soil, in South Africa (Parsons et al., 1996, Scholes et al., 1997). However, these studies measured NO emissions during different seasons and soil moisture conditions compared to our study. For example, Parsons et al. (1996) recorded NO emissions up to $20 \text{ ngN m}^{-2} \text{ s}^{-1}$ over an open savanna during the period going from the end of the dry season to the beginning of the wet season. Nitric oxide emissions of the same magnitude as in our study were also recorded over a grazed semi-arid Sahelian ecosystem in Senegal during the month of July by Delon et al. (2017): $5.7 \pm 3.1 \text{ ngN m}^{-2} \text{ s}^{-1}$ in July 2012 and $5.1 \pm 2.1 \text{ ngN m}^{-2} \text{ s}^{-1}$ in July 2013.

Figures 2 to 5 show daily averaged NO and NH_3 fluxes (± 1 standard deviation) for each land cover type, along with precipitation and soil moisture. The spatial variability of NO fluxes is high, especially for bare soil and the maize field where underground roots, not visible at the surface, are likely to influence the ammonium content of the soil, and the subsequent



NO flux measurement. Standard deviation is generally smaller for grassland. The variation of soil moisture is consistent with the presence of rain events, showing a sharper increase of soil moisture at 5 cm, especially after rainfall following dry periods.

390

NO emissions from bare soil and grassland show an increase, sharper for grassland, one to two days after the rain event on 8th July. The longer rain event between 20th and 24th July does not seem to produce an increase in NO emissions (data available only for maize field and forest). This might be linked with the non-linear relationship between NO biogenic soil emissions and soil water content (Oswald et al., 2013). In fact, a light precipitation event (5-15 mm) occurring on dry soils
395 can result in a large flux of NO (Meixner & Yang, 2006, Hartley & Sclesinger, 2000). However, when soil moisture stays at an equivalent level, after several rain events, pulse emissions do not occur (Millet et al., 2004).

A multiple linear regression analysis was performed between daily mean NO fluxes and the following variables: wind speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, upward longwave radiation and downward shortwave
400 radiation. This regression gives $R^2=0.49$ (p-value=0.004), indicating a weak but existing relationship between those variables and NO soil emissions. However, the aim of our experiment was not to reproduce the known relationships between soil fluxes and meteorological variables, like soil temperature and soil moisture, but to give an estimate of soil fluxes at the larger ecosystem scale. For this reason the location of the soil flux measurements was not kept constant even for the same land cover type on the same measurement day. Moreover, soil temperature and soil moisture were not measured on the same soil
405 parcel where the soil fluxes were measured. Therefore, our experiment does not show the details of microbial and physical processes driving soil fluxes at a single point, but aims to estimate the spatial variability of fluxes at the ecosystem scale.

3.5 NH₃ fluxes

NH₃ fluxes measured during the field campaign range between -6.59 and 4.96 ngN m⁻² s⁻¹. Ammonia fluxes averaged over all land cover types are -0.91 ± 1.27 ngN m⁻² s⁻¹, showing a predominance of NH₃ deposition over emission, which is verified
410 for every land cover type, with an average value of: -1.33 ± 0.86 ngN m⁻² s⁻¹ for bare soil, -0.75 ± 0.31 ngN m⁻² s⁻¹ for the maize field, -0.48 ± 0.55 ngN m⁻² s⁻¹ for grassland, and -0.30 ± 0.38 ngN m⁻² s⁻¹ for forest. Low positive ammonia fluxes, indicating average NH₃ emission, are only recorded during three days, between 6th and 8th July, after the longest dry period of the measurement campaign (Figures 2 to 5).

415 To our knowledge, NH₃ soil fluxes from west African wet savanna are not available in the scientific literature. In Delon et al. (2017) NH₃ soil fluxes measured in Dahra (15°24' N 15°25' W), Senegal, on a dry savanna ecosystem, show low fluxes with a predominance of NH₃ emission: 1.3 ± 1.1 ngN m⁻² s⁻¹, -0.1 ± 1.1 ngN m⁻² s⁻¹ and 0.7 ± 0.5 ngN m⁻² s⁻¹ over three different measurements campaigns. However, Sutton et al. (2007) shows how pre-cut grassland is characterized by NH₃ deposition, as in our study, in contrast to post-cut grassland, which is marked by NH₃ emission. It is interesting to notice that the literature



420 provides up to about $700 \text{ ngN m}^{-2} \text{ s}^{-1}$ NH_3 emission for fertilized Zea Mays fields (Walker et al., 2013) while in our study site NH_3 deposition was recorded for the maize field, which is not treated with mineral fertilizer.

A multiple linear regression analysis was performed between daily mean NH_3 fluxes and the following variables: wind speed, soil temperature at 5 cm, soil moisture at 5 cm, soil heat flux, outgoing longwave radiation and incoming shortwave
425 radiation). This regression gives a weak but existing relationship, with $R^2=0.37$ (p-value=0.03). This correlation highlights the link between NH_3 fluxes and relevant environmental parameters. However, the same considerations explained in Sect. 3.4 for NO emissions are also valid for the correlation between NH_3 fluxes and meteorological variables.

According to the current parameterization of soil ammonia emission potential (Sect. 2.7), high values of pH and $[\text{NH}_4^+]$ in
430 the liquid phase will determine high values of Γ_g indicating that the soil has a high propensity to emit NH_3 . However, despite the high values of Γ_g recorded, our measurement site remains a net sink for NH_3 . The reasons for this can be manifold. One explanation could be that soil particles on our site may have a high adsorption capacity limiting the amount of soil gaseous NH_3 concentrations (Neftel et al., 1998) and the largest part of the estimated ammonium content in the soil may not be in the liquid phase, but adsorbed by solid soil particles. In these conditions ammonium will not be available for gas exchange to
435 open porosity and the atmosphere (Flechard et al., 2013). Another explanation could be given by the presence of a water film at the soil and vegetation surface (linked to high air humidity at the site), which will increase the net deposition process. David et al. (2009) conclude from their measurements that the bare soil can be a significant source of NH_3 only for a limited period and only when the cut vegetation is removed, but not if the soil surface remains covered by grass. Measurements in Ferrara et al. (2014) show other occurrences of high soil ammonia emission potential and NH_3 deposition.

440 **3.6 Comparison of observed and modelled NO soil emissions**

We have compared observed daily averaged (8 a.m. to 6 p.m.) soil NO emissions with those modelled by GEOS-Chem for the entire period of the campaign over the model grid box including the measurement site. The model grid box is positioned at latitude $8.0^\circ \text{ N} - 8.25^\circ \text{ N}$ and longitude $2.19^\circ \text{ E} - 2.5^\circ \text{ E}$. The area of this grid box is 958 km^2 . The land cover type within this grid box is classified as “Savannah (Warm)” but the surrounding area also consists of “Woody Savannah”, while the
445 observations were taken over the four land cover types representative of the region: bare soil, grassland, maize field and forest.

The model is able to reproduce mean air temperature ($25.3 \pm 0.7 \text{ }^\circ\text{C}$) and the main rain events. Soil emissions of NO are well simulated in magnitude. Simulated NO emissions are often higher than those recorded over the grassland areas, however,
450 simulated NO emissions are often within the error bars of measurements (Fig. 6). It appears that when the model is able to reproduce the length and the intensity of the rain events, NO emissions are especially well simulated, e.g. the model is able



to reproduce the longest rain period (from 20th to 26th July 2016) and the decrease of emissions at the end of the measurement campaign.

3.7 Estimate of total NO soil emissions and NH₃ deposition for Benin

455 In order to give a tentative estimate of NO and NH₃ soil fluxes for Benin we have used the land use/land cover map of Benin provided by the US Geographical Survey Atlas: Landscapes of West Africa – A Window on a Changing World (CILSS, 2016, Fig. 7a). The method of mapping land use/land cover used in this atlas was based on Landsat imagery and expert visual interpretation. In particular, these maps provide an accurate indication of cropland distribution using visual interpretation. According to CILSS (2016) Benin's present-day (2013) land cover is mainly savanna, almost 60%, followed by agricultural
460 land, 31%, while forest is only a small fraction under 1% (the rest of the surface is mainly gallery forest and, on a smaller extent, settlements). In the Atlas (CILSS, 2016) bare soils are defined as those surfaces that are bare even in the green/rainy season. For Benin, the amount of bare soil estimated by CILSS (2016) is very small, not big enough to appear on the Atlas' maps. We have multiplied average NO emissions measured at the Savè site for each land cover type by an estimate of the land cover area of each class given by the Atlas. We have made some approximations: as the land use/land cover maps do not
465 distinguish between shrub savanna, tree savanna, and wooded savanna, we have considered NO soil emissions from Savè's grassland savanna to be representative of the general savanna category in CILSS (2016). Moreover, the Atlas has a crop category that does not distinguish the type of crop and we only have observations of NO soil emissions from an intercropped maize field. We have taken NO soil emissions from the maize field as representative of NO emissions of Benin's agricultural land, but other cultures are present in other parts of the country, e.g. oil palm plantations, where possibly stronger
470 fertilization could determine higher NO soil emissions. This tentative calculation gives that Benin's NO soil emissions for the month of July (wet season) is 1.17 ± 0.6 GgN/month, i.e. 0.09% of the average global monthly NO soil emissions as given by Davidson et al., (1997).

We have also calculated Benin's total monthly NO soil emissions with GEOS-Chem adding together the NO soil emissions
475 from the grid boxes where 50% or greater of the box lies within Benin. Benin's total monthly NO soil emissions calculated with GEOS-Chem for the month of July are 1.44 GgN/month and agree with the tentative calculation given above (1.54 ± 0.8 GgN/month). However, the land cover types covering Benin in GEOS-Chem differ from those in the US Geographical Survey Atlas (CILSS, 2016). In GEOS-Chem Benin is cover by 60.9% savanna, 31.4% woody savannah, 4.5% grassland, 1.3% mixed forest and 0.6% urban and built-up lands. Benin's total monthly NO soil emissions calculated with GEOS-Chem
480 for the months of May and June are higher, 3.51 GgN/month and 2.59 GgN/month, respectively, given those months are at the beginning of the wet season and are characterized by more predominant pulse emissions. Using the same method described above we have upscaled point measurements of NH₃ fluxes with relevant land cover surfaces from CILSS (2016) and obtained that total NH₃ dry deposition for the month of July is 0.21 ± 0.11 GgN/month ($0.22 \text{ kgN ha}^{-1} \text{ yr}^{-1}$). This value is



about ten times smaller than the estimation of NH_3 dry deposition given in Adon et al. (2013) for the wet savanna site of
485 Djougou (Benin, 9.7°N , 1.7°E) for the month of July, which is around $2.5 \text{ kgN ha}^{-1} \text{ yr}^{-1}$.

4. Conclusion

We provide soil flux measurements along with soil characteristics for a land cover type, savanna, that is considered to have
large NO emissions (Davidson and Kinglerlee, 1997), and for an area of the world, West Africa, with little observations. The
aim of this study is to contribute to our knowledge in biogenic soil nitrogen exchanges, provide data for inventories and
490 model evaluation to improve air quality and climate modelling.

In situ measurements were made in a wet savanna site in central Benin from mid-June to the end of July 2016.
Complementary to these exchange fluxes, soil N and C content, as well as soil pH, soil moisture, soil temperature and
meteorological data were measured. Soil fluxes of NO and NH_3 were measured over four different land cover types in order
495 to give a tentative estimate of regional soil fluxes.

Given the set up of the experiment, the known relationships between soil fluxes, soil temperature and soil moisture were not
reproduced. Rather than looking at the microbial and physical processes behind soil fluxes, which is not the scope of this
study, we are able to provide observations that are representative of a bigger surface area and that represent the spatial
500 variability of fluxes. However, we observe that while shorter rain events determine an increase in NO soil emissions, the
longer rain event at the end of the campaign (20th to 24th July 2016) is accompanied by a decrease in NO soil emissions, in
agreement with the fact that the relationship between NO soil emissions and soil moisture is not univocal. Soil emissions of
NO increase until an optimum value of soil moisture is reached and then decrease (Oswald et al., 2013).

505 NH_3 emissions measured in this study probably underestimate total NH_3 emissions for the entire country, as possibly higher
localized NH_3 emissions are present in the south of the country where industrial scale agriculture would probably deploy
mineral N fertilization.

Soil NO emissions simulated by GEOS-Chem are in good agreement with the local observations taken at the site of Savè,
510 providing a good baseline for simulating local atmospheric chemistry. Moreover, GEOS-Chem is also in good agreement
with the tentative total monthly NO soil emission estimate for Benin for the month of July made with local observation in
Savè and US Geographical Survey Atlas (CILSS, 2016). All these elements contribute to improve our confidence in the
results of modelling studies of local and regional air quality and climate over this region.

515 Agriculture is the first form of economic activity in Benin, occupying a majority of the active population. The most obvious
recent change in land cover is the major expansion of agricultural land across most regions of Benin. Agricultural areas



(including plantations and irrigated agriculture) progressed from 9.2 to 27.1 % of the total country area between 1975 and 2013, improving food security. Oil palm trees are the main crop, and oil palms farmland already covered most of the southern Terre de Barre plateau of Benin by 1975, and increased by about 28 percent over the following 38-year period. A century or more ago, Benin was covered by dense, biologically diverse forest. Since then, Benin has lost nearly all of that forest cover, by 2013, 58 % of the 1975 forest cover had been lost, leaving only 0.2 percent of the country covered with dense forest. Savanna area has also decreased by 23 percent since 1975, but it still remains the dominant land cover type in Benin and covers more than half of the country (CILSS, 2016).

More measurements of NO and NH₃ exchanges between soil-vegetation-atmosphere in areas of Benin (or West Africa) interested by land-use change could improve our estimate of the impact of biogenic soil emissions on air quality and climate, as biogenic soil fluxes influence for example the amount of aerosol and tropospheric O₃, a greenhouse gas and pollutant, in the atmosphere. Management practices of agriculture affect biogenic soil emissions. Moreover, loosing savanna to oil palm plantations or other crop would have different impacts on air quality, carbon budget and climate than the conversion of forest into crop or oil palm plantation. Furthermore, oil palm plantations are generally closer to the coast and likely to be more influenced by anthropogenic emissions from industry and coastal cities (Knippertz et al., 2015a, 2015b). Oil palm trees are also a strong isoprene emitters. Isoprene emissions influence ozone concentration and the oxidizing capacity of the atmosphere, and it is a source of secondary organic aerosol, thus affecting local air quality and global climate. Large-scale land use change in the tropics – specifically the conversion of tropical rain forest to oil palm plantations in Malaysia – were shown to cause changes in atmospheric composition and chemistry (Hewitt et al., 2009), indicating that the management of the emissions of reactive nitrogen species is essential to prevent damaging levels of ground-level ozone in those regions.



Acknowledgements

The DACCIWA project has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 603502. We also want thank the staffs from NCAS (National Centre for Atmospheric Science), KIT (Karlsruhe Institute of Technology) and UPS (Université Paul Sabatier, Toulouse III) for helping to install the equipment as well as the people from INRAB in Savè for allowing the equipment on their ground.

References

- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of Ox, HOx, NOx and SOx species, *Atmos. Chem. Phys.*, 4, 1461–1738, 2004, <http://www.atmos-chem-phys.net/4/1461/2004/>.
- Bai, E., Li, S., Xu, W., Li, W., Dai, W., Jiang, P.: A meta-analysis of experimental warming effects on terrestrial nitrogen pools and dynamics, *New Phytol.* 199, 441-451. <http://dx.doi.org/10.1111/nph.12252>, 2013.
- Barthès, B., Azontonde, A., Blanchart, E., Girardin, C., Villenave, C., Lesaint, S., Oliver, R. and Feller, C.: Effect of a legume cover crop (*Mucuna pruriens* var. *utilis*) on soil carbon in an Ultisol under maize cultivation in southern Benin. *Soil Use and Manage.*, 20, 231-239, 2004.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J. and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106 (D19), 23,073 – 23,095, doi:10.1029/2001JD000807, 2001.
- Bouwman, A.F., Boumans, L.J.M., Batjes, N.H.: Emissions of N₂O and NO from fertilized fields: summary of available measurement data. *Glob. Biogeochem. Cycles* 16 (4), 1058. <http://dx.doi.org/10.1029/2001GB001811>, 2002.
- Brito, J., Freney, E., Dominutti, P., Borbon, A., Haslett, S. L., Colomb, A., Dupuy, R., Denjean, C., Burnet, F., Bourriane, T., Deroubaix, A., Sellegri, K., Coe, H., Flamant, C., Knippertz, P. and Schwarzenboeck, A., Assessing the role of anthropogenic and biogenic sources on PM₁ over Southern West Africa using aircraft measurements, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-717>, 2017.
- Brooks, B., Bessardon, G., Smith, V., Groves, J., Sharpe, S., Kalthoff, N., Adler, B., Handwerker, J., Kohler, M., Kunka, N., Tan, N., Wieser, A., Lohou, F., Bezombes, Y., Bret, G., Delon, C., Derrien, S., Dione, C., Durand, P., Etienne, P., Gabella, O., Jambert, C., Leclercq, J., Lathon, M., Medina, P., Pedruzo, X., Reinales, I., Jegede, G., Ayoola, M., Sunmonu, L., Ajao, A., Abiye, O., Ajileye, O., Aryee, J., Fosu-Amankwah, K., Cayle-Aethelhard, F., Atiah, W. A., Amekudzi, L., Danuor, S.: A new high-quality dataset of the southern West African atmospheric boundary layer diurnal cycle during the Monsoon season, *Sci. Data*, 2017, in preparation.



- CILSS (2016). Landscapes of West Africa – A Window on a Changing World. U.S. Geological Survey, EROS, 47914 252nd St, Garretson, SD 57030, UNITED STATES, 2016.
- David, M., Loubet, B., Cellier, P., Mattsson, M., Schjoerring, J.K., Nemitz, E., Roche, R., Riedo, M., Sutton, M.A.:
570 Ammonia sources and sinks in an intensively managed grassland canopy. *Biogeosciences* 6, 1903-1915, 2009.
- Davidson, E. A. and Kinglerlee, W.: A global inventory of nitric oxide emissions from soils, *Nutr Cycl Agroecosys*, 48: 37–50, 1997.
- Delon, C., Galy-Lacaux, C., Boone, A., Lioussé, C., Serça, D., Adon, M., Diop, B., Akpo, A., Lavenu, F., Mougín, E., and Timouk, F.: Atmospheric nitrogen budget in Sahelian dry savannas, *Atmos. Chem. Phys.*, 10, 2691–2708, doi:10.5194/acp-
575 10-2691-2010, 2010.
- Delon, C., Galy-Lacaux, C., Adon, M., Lioussé, C., Serça, D., Diop, B., and Akpo, A.: Nitrogen compounds emission and deposition in West African ecosystems: comparison between wet and dry savanna, *Biogeosciences*, 9, 385–402, doi:10.5194/bg-9-385-2012, 2012.
- Delon, C., Galy-Lacaux, C., Serça, D., Loubet, B., Camara, N., Gardrat, E., Saneh, I., Fensholt, R., Tagesson, T., Le Dantec,
580 V., Sambou, B., Diop, C., Mougín, E.: Soil and vegetation-atmosphere exchange of NO, NH₃, and N₂O from field measurements in a semi arid grazed ecosystem in Senegal, *Atmos. Environ.*, 156, 36–51, 2017.
- Derrien, S., Bezombes, Y., Bret, G., Gabella, O., Jarnot, C., Medina, P., Pique, E., Delon, C., Dione, C., Campistron, B., Durand, P., Jambert, C., Lohou, F., Lothon, M., Pacifico, F., Meyerfeld, Y.: DACCIWA field campaign, Savè super-site, UPS instrumentation; SEDOO OMP. <https://doi.org/10.6096/DACCIWA.1618>, 2016.
- 585 EC-JRC / PBL: Emission Database for Global Atmospheric Research (EDGAR), release version 4.2. European Commission, Joint Research Centre (JRC)/Netherlands Environmental Assessment Agency (PBL), <http://edgar.jrc.ec.europa.eu>, 2011.
- Ferrara, R. M., Loubet, B., Decuq, C., Palumbo, A. D., Di Tommasi, P., Magliulo, V., Masson, S., Personne, E., Cellier, P., Rana, G.: Ammonia volatilisation following urea fertilisation in an irrigated sorghum crop in Italy, *Agr. Forest Meteorol.* 195–196, 179–191, 2014.
- 590 Flechard, C.R., Massad, R.-S., Loubet, B., Personne, E., Simpson, D., Bash, J.O., Cooter, E.J., Nemitz, E., Sutton, M.A.: Advances in understanding, models and parameterizations of biosphere-atmosphere ammonia exchange, *Biogeosciences*, 10, 5183-5225, 2013.
- Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V.,
595 Vignati, E., Wild, M., Williams, M., Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs. *Atmos. Chem. Phys.* 15, 8217-8299. <http://dx.doi.org/10.5194/acp-15-8217-2015>, 2015.



- Giglio, L., Randerson, J. T. and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res. Biogeosci.*, 118, 317–328, doi:10.1002/jgrg.20042, 2013.
- 600 Grote, R., Lehmann, E., Brümmner, C., Brüggemann, N., Szarzynski, J., Kunstmann, H.: Modelling and observation of biosphere-atmosphere interactions in natural savannah in Burkina Faso, West Africa. *Phys. Chem. Earth* 34, 251-260, 2009.
- Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, doi:10.5194/gmd-5-1471-2012, 2012.
- 605 Jegede, O. O., Ayoola, M. A., Sunmonu, L. A., Ajao A. I., and Babić K.: DACCIWA field campaign, Ile-Ife super-site, Surface measurements; SEDOO OMP.
- Kalthoff, N., Lohou, F., Brooks, B., Jegede, G., Adler, B., Babić, K., Dione, C., Ajao, A., Amekudzi, L. K., Aryee, J. N. A., Ayoola, M., Bessardon, G., Danour, S. K., Handwerker, J., Kohler, M., Lothon, M., Pedruzo-Bagazgoitia, X., Smith, V., Sunmonu, L., Wieser, A., Fink, A. H.1, and Knippertz, P.: An overview of the diurnal cycle of the atmospheric boundary
610 layer during the West African monsoon season: Results from the 2016 observational campaign, *Atmos. Chem. Phys. Discuss.*, <https://doi.org/10.5194/acp-2017-631>, 2017.
- Kohler, M., Kalthoff, N., Seringer, J., and Kraut, S.: DACCIWA field campaign, Savè super-site, Surface measurements; SEDOO OMP. <https://doi.org/10.6096/DACCIWA.1690>, 2016.
- Knippertz, P., Coe, H., Chiu, J. C., Evans, M. J., Fink, A. H., Kalthoff, N., Liousse, C., Mari, C., Allan, R. P., Brooks, B.,
615 Danour, S., Flamant, C., Jegede, O. O., Lohou, F., and Marsham, J. H.: The DACCIWA project: Dynamics-aerosol-chemistry-cloud interactions in West Africa, *B. Am. Meteorol. Soc.*, 96, 1451–1460, <https://doi.org/10.1175/BAMS-D-14-00108.1>, 2015a.
- Knippertz, P., Evans, M. J., Field, P. R., Fink, A. H., Liousse, C., and Marsham, J. H.: The possible role of local air pollution in climate change in West Africa, *Nature Climate Change*, 5, 815–822, <https://doi.org/10.1038/nclimate2727>, 2015b.
- 620 Knippertz, P., Fink, A. H., Affo-Dogo, A., Bahaga, T., Brosse, F., Deroubaix, A., Evans, M., Gaetani, M., Guebsi, R., Latifou, I., Lavaysse, C., Maranan, M., Mari, M., Marsham, J. H., Meynadier, R., Morris, E., Rosenberg, P. D., Schlueter, A., and Tocquer, F.: A meteorological and chemical overview of the DACCIWA field campaign in West Africa in June–July 2016, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2017-345, 2017.
- Handwerker, J., Scheer, S., and Gamer, T.: DACCIWA field campaign, Savè super-site, Cloud and precipitation; SEDOO
625 OMP. <https://doi.org/10.6096/DACCIWA.1686>, 2016.



- Hewitt, C. N., MacKenzie, A. R., Di Carlo, P., Di Marco, C. F., Dorsey, J. R., Evans, M., Fowler, D., Gallagher, M. W., Hopkins, J. R., Jones, C. E., Langford, B., Lee, J. D., Lewis, A. C., Lim, S. F., McQuaid, J., Misztal, P., Moller, S. J., Monks, P. S., Nemitz, E., Oram, D. E., Owen, S. M., Phillips, G. J., Pugh, T. A. M., Pyle, J. A., Reeves, C. E., Ryder, J., Siong, J., Skiba, U., and Stewart, D. J.: Nitrogen Management Is Essential to Prevent Tropical Oil Palm Plantations from Causing
630 Ground-Level Ozone Pollution, *P. Natl. Acad. Sci.*, 106, 18447–18451, 2009.
- Hudman, R.C., Russell, A.R., Valin, L.C., Cohen, R.C.: Interannual variability in soil nitric oxide emissions over the United States as viewed from space. *Atmos. Chem. Phys.* 10, 9943e9952. <http://dx.doi.org/10.5194/acp-10-9943-2010>, 2010.
- Hudman, R. C., Moore, N. E., Mebust, A. K., Martin, R. V., Russell, A. R., Valin, L. C., and Cohen, R. C.: Steps towards a mechanistic model of global soil nitric oxide emissions: implementation and space based-constraints, *Atmos. Chem. Phys.*,
635 12, 7779-7795, doi:10.5194/acp-12-7779-2012, 2012.
- Kristensen, L., Lenschow, D. H., Gurarie, D., Jensen, N. O.: A simple model for the reactive species in the convective atmospheric boundary layer, *Bound-Lay Meteorol.*, 134, 195-221, doi:10.1007/s10546-009-9443-x, 2010a.
- Kristensen, L., Lenschow, D. H., Gurarie, D., Jensen, N. O.: Erratum to: A Simple Model for the Vertical Transport of Reactive Species in the Convective Atmospheric Boundary Layer, *Bound-Lay Meteorol.*, 135, 181–183, DOI
640 10.1007/s10546-010-9473-4, 2010b.
- Lata, J.-C., Degrange, V. Raynaud, X., Maron, P.-A., Lensis, R. and Abbadie, L.: Grass populations control nitrification in savanna soils, *Funct. Ecol.*, 18, 605-611, 2004.
- Le Roux, X., Abbadie, L., Lensi, R., Serça, D.: Emission of nitrogen monoxide from African tropical ecosystems: control of emission by soil characteristics in humid and dry savannas of West Africa. *J. Geophys. Res.* 100, 23133-23142, 1995.
- Loubet, B., Decuq, C., Personne, E., Massad, R.S., Flechard, C., Fanucci, O., Mascher, N., Gueudet, J.-C., Masson, S., Durand, B., Genermont, S., Fauvel, Y., Cellier, P.: Investigating the stomatal, cuticular and soil ammonia fluxes over a growing tritical crop under high acidic loads. *Biogeosciences* 9, 1537-1552, 2012.
- Massad, R.-S., Nemitz, E., Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere. *Atmos. Chem. Phys.* 10, 10359-10386. <http://dx.doi.org/10.5194/acp-10-10359-2010>, 2010.
- 650 McCalley, C. K. and Sparks, J. P.: Controls over nitric oxide and ammonia emissions from Mojave Desert soils. *Oecologia* 156, 871-881, 2008.
- Meixner, F.X., Yang, W.X., In: DO doroco, P., Porporato, A. (Eds.), *Biogenic Emissions of Nitric Oxide and Ditrour Oxide from Arid and Semi-arid Land, Dryland Ecohydrology*. Kluwer Academic Publishers B.V, Dordrecht, The Netherlands, pp. 23-46, 2006.



- 655 Michiels, B., Babatounde, S., Dahouda, M., Chabi, S. L. W. and Buldgen, A.: Botanical composition and nutritive value of forage consumed by sheep during the rainy season in a Sudano-guinean savanna (central Benin), *Trop Grasslands*, 34, 43-47, 2000.
- Neftel, A., Blatter, A., Gut, A., Högger, D., Meixner, F., Ammann, C., Nathaus, F. J.: NH_3 soil and soil surface gas measurements in a triticale wheat field, *Atmos. Environ.*, 32, 3, 499-505, 1998.
- 660 Oswald, R., Behrendt, T., Ermel, M., Wu, D., Su, H., Cheng, Y., et al.: HONO Emissions from Soil Bacteria as a Major Source of Atmospheric Reactive Nitrogen. *Science*, 341(6151), 1233-1235, 2013.
- Paillat, J., -M., Robin, P., Hassouna, M., Leterme, P.: Predicting ammonia and carbon dioxide emissions from carbon and nitrogen biodegradability during animal waste composting, *Atmos. Environ.* 39, 6833–6842, 2005.
- Pape, L., Ammann, C., Nyfeler-Brunner, A., Spirig, C., Hens, K., Meixner, F.X.: An automated dynamic chamber system for
665 surface exchange measurement of non-reactive and reactive trace gases of grassland ecosystems. *Biogeosciences*, 6, 405e429. <http://dx.doi.org/10.5194/bg-6-405-2009>, 2009.
- Parsons, D. A. B., Scholes, M. C. and Levine, J. S.: Biogenic NO emissions from savanna soils as a function of fire regime, soil type, soil nitrogen and water status, *J Geophys Res* 101: 23683–23688, 1996.
- Pilegaard, K.: Processes regulating nitric oxide emissions from soils, *Phil. Trans. R. Soc. B* 2013 (368), 20130126, 2013.
- 670 Säïdou, A., Kuyper, T. W., Kossou, D. K., Tossou, R., and Richards, P.: Sustainable soil fertility management in Benin: learning from farmers, *NJAS-WAGEN J LIFE SC*, 52-3/4, 349-369, 2004.
- Santín C., and Doerr, S. H.: Fire effects on soils: the human dimension. *Phil. Trans. R. Soc. B* 371: 20150171. <http://dx.doi.org/10.1098/rstb.2015.0171>, 2016.
- Scholes, M. C., Scholes, R. J., Parsons, D., Martin, R., and Winstead, E.: NO and N_2O emissions from savanna soils
675 following the first rains. *Nut. Cycling Agroecosys.* 48: 115–122, 1997.
- Serça, D., Delmas, R., Le Roux, X., Parsons, D. A. B., Scholes, M. C., Abbadie, L., Lensi, R., Ronce, O., Labroue, L.: Comparison of nitrogen monoxide emissions from several African tropical ecosystems and influence of season and fire. *Glob. Biogeochem. Cy.* 12 (4), 637-651, 1998.
- Steinkamp, J., Ganzeveld, L.N., Wilcke, W., Lawrence, M.G.: Influence of modelled soil biogenic NO emissions on related
680 trace gases and the atmospheric oxidizing efficiency. *Atmos. Chem. Phys.* 9, 2663e2677. [http:// dx.doi.org/10.5194/acp-9-2663-2009](http://dx.doi.org/10.5194/acp-9-2663-2009), 2009.
- Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Butterbach Bahl, K., Cellier, C., de Vries, W., Cotrufo, F., Skiba, U., Di Marco, C., Jones, S., Laville, P., Soussana, J. F., Loubet, B., Twigg, M., Famulari, D., Whitehead, J., Gallagher, M. W., Neftel, A., Flechard, C., Herrmann, B., Calanca, P.L., Schjoerring, J. K., Daemmgen, U., Horvath, L., Tang, Y. S., Emmett, B.



- 685 A., Tietema, A., Peñuelas, J., Kesik, M., Brüggemann, N., Pilegaard, K., Vesala, T., Campbell, C. L., Olesen, J. E., Dragosits, U., Theobald, M. R., Levy, P., Mobbs, D. C., Milne, R., Viovy, N., Vuichard, N., Smith, J. U., Smith, P. E., Bergamaschi, P., Fowler, D., and Reis, S.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, *Environ. Pollut.*, 150, 125–139, 2007.
- Sutton, M., Reis, S., and Baker, S. M. H.: Atmospheric Ammonia, detecting emission changes and environmental impacts. Results of an expert workshop under the convention of long range transboundary air pollution, Springer Edition, New York, ISBN 978-1-4020-9120-9, 2009a.
- 690 Sutton, M. A., Nemitz, E., Milford, C., Campbell, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M., Dorsey, J. R., Gallagher, M. W., Horvath, L., Weidinger, T., Meszaros, R., Dammgén, U., Neftel, A., Herrmann, B., Lehman, B. E., Flechard, C., Burkhardt: Dynamics of ammonia exchange with cut grassland: synthesis of results and conclusions of the GRAMINAE integrated experiment, *Biogeosciences* 6, 2907-2934, 2009b.
- Tiquia, S. M. and Tam, N. F. Y.: Fate of nitrogen during composting of chicken litter, *Environ. Pollut.*, 110, 535-541, 2000.
- Walker, J. T., Jones, M. R., Bash, J. O., Myles, L., Meyers, T., Schwede, D., Herrick, J., Nemitz, E., and W. Robarge: Processes of ammonia air-surface exchange in a fertilized *Zea mays* canopy, *Biogeosciences*, 10, 981–998, doi:10.5194/bg-700 10-981-2013, 2013.
- Wentworth, G. R., Murphy, J. G., Gregoire, P. K., Cheyne, C. A. L., Tevlin, A. G. and Hems, R.: Soil–atmosphere exchange of ammonia in a non-fertilized grassland: measured emission potentials and inferred fluxes, *Biogeosciences*, 11, 5675-5686, 2014.
- Wieser, A., Adler, B., and Deny, B.: DACCIWA field campaign, Savè super-site, Thermodynamic data sets; SEDOO OMP. 705 <https://doi.org/10.6096/DACCIWA.1659>, 2016.



	Savè ground-based observation site
Location	8°02'03" N, 2°29'11" E
Elevation	166 m a.s.l.
Mean annual precipitation	1100 mm
Mean annual temperature	27.5 °C
Soil type	sandy
Sand percentage	87%
Clay percentage	4.1%

Table 1

710 Main characteristics of the Savè site.



Soil type		Plant family	Plant species	Common name/s
Next to grassland and forest	Dominant tree species	Anacardiaceae	<i>Anacardium occidentale</i>	cashew tree
		Fabaceae	<i>Daniellia oliveri</i>	African copaiba balsam tree
			<i>Pterocarpus erinaceus</i>	barwood, muninga, vène, mukwa
	Dominant ground species	Cleomaceae	<i>Cleome</i> sp.	spider flowers, spider plants
		Fabaceae	<i>Crotalaria</i> sp.	rattlepod or rattlebox
			<i>Mucuna</i> sp.	velvet bean
		Poaceae	<i>Imperata cylindrica</i>	cogon grass, cotton wool grass, kura-kura
			<i>Rhynchelytrum repens</i>	rose natal grass
Next to maize field	Dominant tree species	Anacardiaceae	<i>Mangifera indica</i>	mango
		Arecaceae	<i>Cocos nucifera</i>	coconut tree
		Caricaceae	<i>Carica papaya</i> L.	papaya
		Lamiaceae	<i>Tectona grandis</i>	teak
		Meliaceae	<i>Azadirachta indica</i>	neem, nimtree, Indian lilac
	Dominant ground species	Commelinaceae	<i>Commelina benghalensis</i>	benghal dayflower, tropical spiderwort
		Euphorbiaceae	<i>Euphorbia</i> sp.	spurge
		Nyctaginaceae	<i>Boerhavia diffusa</i>	punarnava, red spiderling
		Phyllanthaceae	<i>Phyllanthus amarus</i>	gale of the wind, stonebreaker
		Poaceae	<i>Digitaria horizontalis</i>	Jamaican crabgrass
Crops		Dioscoreaceae	<i>Dioscorea</i> sp.	yam
		Euphorbiaceae	<i>Manihot esculenta</i>	cassava
		Fabaceae	<i>Arachis hypogaea</i>	peanut
			<i>Vigna unguiculata</i>	cowpea
		Malvaceae	<i>Gossypium</i> sp.	cooton
		Pedaliaceae	<i>Sesamum indicum</i>	sesame
		Poaceae	<i>Zea mays</i>	maize
	<i>Sorghum</i> sp.		sorghum	
	Solanaceae	<i>Solanum lycopersicum</i>	tomato	



715

Table 2

List of plant species at the Savè site. The list of common names is not considered to be exhaustive.



720

	Bare Soil	Grassland	Maize Field	Forest
Clay (<2µm) (%)	5.13±0.63	3.15±0.50	4.40±0.35	3.70±1.25
Fine Silt (2 to 20µm) (%)	5.13±0.96	2.93±0.32	4.13±1.00	3.40±1.21
Coarse Silt (20 to 50µm) (%)	5.98±0.51	4.78±0.66	4.37±0.38	4.67±1.05
Total Sand (50 to 2000 µm) (%)	83.75±1.82	89.20±0.71	87.13±0.99	88.20±3.50

Table 3

List of soil characteristics for each land cover type at the Savè site, including standard deviation.

725



Soil Type	Date	C/N ratio	Organic C (g kg ⁻¹)	Total N (g kg ⁻¹)
Bare soil	06/07/2016	14.90	24.28	1.63
	09/07/2016	12.20	10.47	0.86
	19/07/2016	12.30	15.05	1.22
	28/07/2016	9.50	19.30	2.04
Grassland	07/07/2016	15.20	5.78	0.38
	09/07/2016	16.00	7.36	0.46
	19/07/2016	14.90	7.16	0.48
	28/07/2016	10.90	4.56	0.42
Maize	09/07/2016	14.80	17.33	1.17
	19/07/2016	16.40	13.08	0.8
	28/07/2016	11.80	11.83	1.00
Forest	06/07/2016	14.80	7.98	0.54
	19/07/2016	11.90	9.62	0.81
	28/07/2016	11.30	16.56	1.47

730 Table 4

List of soil characteristics for each land cover type at the Savè site for each soil sampling day: Carbon-to-Nitrogen ratio (C/N), organic carbon (g kg⁻¹) and total nitrogen (g kg⁻¹). The accuracy for the C/N ratio is 14%. The measurement accuracy for organic carbon and total nitrogen is 14 and 13%, respectively.

735




Soil Type	Date	pH	[NH ₄ ⁺] (mg kg ⁻¹)	Γ _g [NH ₄ ⁺]/[H ⁺]	X _g (ppb)
Bare soil	06/07/2016	8.38	6.82	136 334	1891
	09/07/2016	7.73	2.90	12 978	134
	19/07/2016	8.46	6.63	159 343	2188
	28/07/2016	8.34	8.01	146 033	2215
Grassland	07/07/2016	7.15	1.96	2 307	29
	09/07/2016	7.91	1.55	10 499	108
	19/07/2016	7.52	2.28	6 291	86
	28/07/2016	6.51	2.30	620	9
Maize	09/07/2016	7.46	4.40	10 575	109
	19/07/2016	7.61	14.74	50 040	687
	28/07/2016	8.04	4.49	41 027	622
Forest	06/07/2016	6.32	2.18	380	5
	19/07/2016	7.51	4.88	13 159	181
	28/07/2016	7.37	11.47	22 407	340

Table 5

740 List of soil pH, ammonium concentrations [NH₄⁺] (mg kg⁻¹), soil emission potential Γ_g and soil compensation point χ_g (ppb) for each land cover type at the Savè site for each soil sampling day. The measurement accuracy for pH is 0.15 when pH ≤ 7 and 0.20 when pH > 7. The accuracy for ammonium concentrations [NH₄⁺], soil emission potential Γ_g and soil compensation point χ_g is 14%.




745



750

755




760

765


770

775

Fig. 1(a) Location of the Savè site in West Africa, (b) one of the bare soil sampling sites, (c) the grassland sampling site, (d) the maize field sampling site and (e) the forest sampling site at the Savè site.



780 Fig. 2 Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, daily mean NO and NH₃ fluxes in $\text{ngN m}^{-2} \text{s}^{-1}$ measured at the bare soil site. Vertical bars show the standard deviation from individual fluxes.



785

Fig. 3 Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, daily mean NO and NH₃ fluxes in ngN m⁻² s⁻¹ measured at the grassland site. Vertical bars show the standard deviation from individual fluxes.




Fig. 4 Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) instrumentation, daily mean NO and NH₃ fluxes in ngN m⁻² s⁻¹ measured at the maize field site. Vertical bars show the standard deviation from individual fluxes.

795




Fig. 5 Daily total precipitation (mm), daily mean soil moisture at 5 cm (%) measured by the Karlsruhe Institute of Technology (KIT), daily mean soil moisture averaged between 0 and 30 cm measured by the Université Paul Sabatier (UPS) 800 instrumentation, daily mean NO and NH₃ fluxes in ngN m⁻² s⁻¹ measured at the forest site. Vertical bars show the standard deviation from individual fluxes.




Fig. 6 Nitric oxide emissions measured over each land cover type (orange dot for bare soil, green for grassland, yellow for the maize field and brown for forest) and simulated with GEOS-Chem, along with rainfall measured and modelled with GEOS-Chem. Soil NO emissions are daily average between 8 a.m. and 6 p.m..




Fig. 7 (a) Land cover map of Benin for 2013 from the US Geographical Survey Atlas: Landscapes of West Africa – A Window on a Changing World (CILSS, 2016) and (b) land cover map of Benin used in the GEOS-Chem simulation.