

Answers to reviewer comments

We would like to start by thanking the reviewer for the many constructive comments that help us improve the manuscript. The reviewer's questions and remarks have been answered in the text below, where reviewer comments are marked in black and our responses in this [blue](#) color.

Reviewer 1

This contribution to our understanding of stratospheric aerosol optical depth (AOD) focuses on the CALIOP backscatter data set after it is cleared of clouds. A significant fraction of the paper is devoted to a procedure to correct CALIOP measurements beneath volcanic clouds when the laser is attenuated by the volcanic cloud. The procedure is relatively straight forward, although I found the discussion of how the threshold is set to instigate the procedure, and the relevance of Figure 3 to that procedure, confusing. I don't have specific suggestions to help here, but perhaps the authors could have a close read of this part to see if it may be improved.

Then the procedure is applied and the CALIOP data set presented over the 14 year period. The results are quite interesting and I appreciate the separation of the stratosphere into three layers associated with their relevance to cross tropopause transport. This separation led to some nice generalizations concerning the volcanic eruptions which occurred over the 14 year period and how these eruptions influenced AOD and hence the cooling associated with stratospheric aerosol.

The paper is interesting, especially once the data are discussed, and should be published. There are a few things that need to be taken care of before that happens.

Aside from the detailed comments below the authors should add labels a), b), . . . to all the figure panels. The authors are very good at explaining what each of the a), b), . . . panels are in the captions, and refer to Figure Xa in the text. But none of the panels in the figures have labels. [We apologize for overlooking this, and have now included the indexes to all subfigures.](#)

Overall the writing is quite good, but there are a few awkward places. I have tried to flag these. One general comment the phrase "in order" is never necessary and can be deleted with no change in the sentence meaning. [Thank you for pointing that out. We have included the suggested changes, and make further language corrections adjusting to comment #4 by the second reviewer.](#)

1.9-11. Why switch between CALIOP and CALIPSO in the abstract when neither are defined. CALIOP is the instrument and should suffice in both places. [We agree with the reviewer. We will use CALIOP throughout the manuscript, and add the definition in the abstract.](#)

1.19-20. Awkward sentence, beginning with "of which". Try. Trends in the abundance of aerosol particles are an important component of the climate system, although their influence on climate is still highly uncertain. . . [We have used the reviewer's suggestion.](#)

2.10-14. Change forming to contributing. If OCS is known to form the Junge layer (which it isn't) then why the discussion about so₂? [The reviewer is right, and we have changed accordingly.](#)

2.21 LMS? – We forgot to define the lowermost stratosphere. Thank you for pointing that out.

5.20 residing . . . We changed accordingly.

6.9 becomes . . . We changed accordingly.

7.29-31. This is confusing. Why would you expect aerosol scattering to compare with extinction? There is at least a factor of 50 between them. In Fig. 3 it is much more. The AS is noted in a range of $1e-8$ to $1e-7$, while the extinctions range from 0.02-0.1. The units are not of importance for the comparison. It is simply done to highlight that the Level 2 extinction data do not follow that of the volcanic aerosol. We understand that it is not obvious for the reader, and therefore added a sentence to clarify this (8.26).

Fig. 3. There are no labels a), b), . . . Why is there a big 0 between the panels at the bottom? We have updated the figure to include the requested labels and there is no 0 between the panels in the new version.

8.1 Where is Fig. 2b? There are no labels on Figure 2 either. Why strangely? We apologize for missing this. Subfigure indexing is added in the new version (June, 2018).

11.2 first 8 months . . . We changed this.

11.3. Are these the averages, maxima, value after 8 months, or? They are the average, and we have now added that info in the text (12.3).

11.28-29, “A feature appearing in the southern tropics in April has been identified as smoke from bush fires in February 2009 in Victoria, Australia (Vernier et al., 2011).” Is this the feature near 20 km, if so it is worth mentioning the altitude as this may be surprising for some. Best to be clear. It is the feature around 20 km altitude. We agree with the reviewer and have clarified this in the manuscript (12.28).

Figure 6. Perhaps the vertical solid lines represent the tropical eruptions? The figure caption is confusing. Please add short labels to the eruptions, which are listed in Table 1, so the reader can easily see the effects from Sarychev, Calbuco, and Kasatochi, without having to search them out. Similar labels would be appreciated in Figures, 5, 7-9. The indexing was incorrect. Thank you for pointing that out. We change this, and add indexing in the figures.

13.18-19. What PV levels are considered the LMS. This is needed to understand the statement, “the aerosol signal is strong also in the lowest LMS layer of Fig. 6a”. What is the lowest LMS layer? All of the PV-levels in the figure pertain to the LMS. The lowest LMS layer is 1.5-2 PVU. We have clarified this in the manuscript (14.15).

13.25. The discussion previous to this conclusion has been descriptive of Figure 6, but it’s not clear why the last half of this sentence holds. The volcanoes affected all the PV layers investigated more or less equally. It would be nice to see a PV level that wasn’t affected, then maybe there would be a justification for the choice. The value of 1.5 PVU is the lowest dynamic tropopause commonly used. We find volcanism to impact all the way down to this lowest (stratospheric) PV-level (and in the case

of Kasatochi there is an impact also below this PV-level). Therefore, we choose the 1.5 PVU to represent the lower altitude limit where volcanism impacts the stratospheric aerosol. It has been clarified in the manuscript (14.29).

14.6-7. This conclusion about the source of the springtime increase in aerosol scattering is much too strong based on the evidence given. What do “These observations” refer to, to the work here or that by Martinsson? Perhaps they “suggest”, but “indicate”, no. Much more work would be required to come to such a definite conclusion. We agree with the reviewer and have changed to “suggest”.

Figure 8. Why don't the sum of the three lines add up to the total? There is usually at least a difference of 0.002 in AOD between the sum of the three layers and the total. We have controlled this and do not find any discrepancy, neither in the codes out-print, nor in the figure in the manuscript. We find that the three lines (representing the three layers) in each subfigure adds up to the total (red lines). We would be grateful if the reviewer could point out specific coordinates as an example of the stated discrepancy.

14.26-29. This is a bit surprising. Can some more detail be provided? Using Figure 6 the spring time maxima in 2007 and 2008 indicate SR in the lowest layers equivalent to the SR after Nabro, which shows a large AOD. Is it the fact that the impact of the dust layers, or whatever is causing this, so narrow that the impact on AOD is small?

The seasonal increase in AOD from dust (and other tropospheric sources), are estimated to constitute *less than 5%* of that of the AOD increase from Nabro. The dust signals are rapidly decreasing in strength in the first 2 km above the tropopause (1.5-5.5 PVU, the ExTL). Thus, they constitute a small fraction of the total stratospheric AOD, since they are contained within such a small fraction of the stratospheric air-mass. The picture is the opposite for the Nabro aerosol. The sulfate concentrations are increasing from the tropopause into the stratosphere, and this effect is larger for tropical eruptions than for the local eruptions in midlatitudes. In midlatitudes the subsidence of air (Nabro's aerosol) from the 380 K isentrope through the LMS to the tropopause takes several months, during which the stratospheric air gradually mixes with tropospheric air which decreases the sulfate concentrations and induces a gradient of sulfate from the tropopause throughout the LMS. The sulfur gradient and the weak Nabro aerosol signals around the tropopause are both evident in Figure 5a. We have estimated that *the Nabro aerosol in the ExTL constituted less than 6%* of the total stratospheric AOD during the period of 1-8 months after its eruption.

This may be counterintuitive from the appearance of figure 6. We therefore added sentences to clarify this (15.14-15.16).

15.3. The small peak near the equator in the upper layer can hardly be considered to be “high AOD”. It is only slightly higher than in the mid latitudes. The reviewer is right. We have reformulated our statement.

15.26-31. It would be helpful to indicate here that Figure 7 is being discussed. This is the figure I used to follow the discussion. Thanks for pointing that out. We have added a reference to Figure 7.

16.4. “A small sudden increase in the AOD is observed in the southern extratropics approximately one and a half years after the Kelut eruption.” At what level? Which figure is being referred to? [This is indeed confusing. We have added a more descriptive text.](#)

16.8-9. “The aerosol in the upper layer is eventually transported out to the next lower layer at midlatitudes, i.e the one located between the 380-470 K isentropes.” It would be nice to know what figure is being used to make this claim. Figure 7 for example does not support this statement.

[The AOD in Figure 7b indicates this. For example, the AOD in the northern midlatitudes of the layer 380-470 K was higher in year 2007 and beginning of 2008 compared to the background \(year 2013\). The increased AOD in midlatitudes most likely comes from subsidence from the overlying layer. We will make changes to the text to clarify this.](#)

19.2. “. . . forcing, and is added . . .” [We will make the suggested change.](#)

Reviewer 2

Volcanic eruptions represent one of the largest source of natural variability of our climate system. Thus, the construction of long-term database of stratospheric aerosol optical depth is highly important to constrain global climate models. In this study, Friberg et al. (2018) used the CALIPSO space-born lidar to derive the time evolution of stratospheric AOD between 2005 and 2016. They proposed two new techniques to correct the effect of particle attenuation on retrieved optical parameters (backscatter and extinction) and remove Polar Stratospheric Clouds. After selecting a definition of the tropopause based on Potential Vorticity, they show time series of stratospheric AOD and discuss the influence of several volcanic eruptions. Overall, the paper is interesting, and the technique developed to correct the effect of particle attenuation is relatively well explained. But there are several points which would need serious considerations before its publication in ACP.

1. Purpose of this work. The purpose of this study is not clear. Do the authors intend to create a new aerosol dataset from the CALIPSO data? If yes, do they have a plan for archiving the data and make them publicly available? If this is the case, is there any established collaboration with the CALIPSO team to work on this dataset or it is an independent endeavor?

The purpose of this work is to investigate the stratospheric aerosol in a period when several volcanic eruptions affected the stratosphere. Compared to previous studies we have put emphasis on understanding the aerosol load in relation to important transport features of the stratosphere. We have also included a larger part of the stratosphere compared with previous studies, approaching a complete coverage of the stratospheric aerosol of the period. We highly appreciate the work of the CALIPSO team, and would be interested in collaboration just as we have been in the past (Andersson et al., 2015).

2. Lack of validation. This study does not make use of additional datasets to compare/ validate the retrieved AOD. Several satellite datasets (e.g. OSIRIS, OMPS) are available and could provide a source of validation data but are not used. What about In situ data from the CARIBIC program such as aerosol size distribution in the Lowermost stratosphere? Why not to use those data to infer lidar ratio using Mie Calculations?

CALIOP data are validated already in for example publications by Vernier, and used in the Science paper of Solomon et al. (2011). Our methods to retrieve the AOD are not significantly different from Vernier's and that used in our previous CALIOP publication in Nature Communications 2015 (Andersson et al., 2015). We have also validated that we retrieve the same scattering and AOD values (at 20-30 km) as in Vernier et al. (2011). We will compare our AODs with those in Rieger et al. (2015), Vernier et al. (2011), and Thomason et al. (2018), and add a more thorough description about the lidar ratio in the manuscript.

Our intension is to discuss the volcanic impact from the global observations of CALIOP, and to include the entire stratosphere. Furthermore, a paper on comparison between CALIOP and CARIBIC is in preparation by one of the co-authors.

Our paper extends the study by Andersson et al. (2015) highlighting the significance of the LMS, which has been neglected until very recently, and the overlying stratosphere has to our knowledge not been separated in individual layers before.

3. *Retrieving AOD from CALIPSO. The authors propose an approach to correct the particle attenuation effect which is especially important after a significant volcanic eruption. A major issue with the proposed technique is its dependency to the type of volcanic eruptions. The major assumption of the correction technique is to assume that the Upper Troposphere is clear of volcanic aerosol, but this is not always the case as shown after the Kasatochi eruption. Any techniques applied for this purpose should be independent from volcanoes and therefore could only be achieved by the iterative approach developed in Hostetler et al. (2006). The overall impact on the corrected AOD is relatively small (impact between 4-7 %).*

It is true that it depends on the type of eruption and cannot be used directly for all types of eruptions. Kasatochi is a special case. In-situ observations by CARIBIC shows that volcanic elevations of the stratospheric aerosol load generally have small impact on the UT aerosol concentrations. Thus, the proposed method can be used directly for eruptions which injections mainly reached above the UT, and by our method we risk doing a small underestimation of the true volcanic elevations.

We believe that the method described in Hostetler et al. (2006)/Young et al. (2005) is good for correcting for attenuation in detected volcanic layers, but that it is evident that the method cannot detect the volcanic aerosol once it has mixed with the background. Hence, the two methods have different weaknesses.

The computed changes are relatively small in perspective to the total stratospheric AOD for the period studied here, but could become much more important for stronger eruptions. The attenuation depends on the overlying aerosol column, and is thus particularly important for the representation of the lowest layer of the stratosphere, i.e. the LMS.

The authors never discussed thoughtfully the other sources of uncertainties that could have bigger influences (e.g. calibration of the lidar, lidar ratio conversion factor). We agree that these uncertainties are important and that the attenuation corrections made in the manuscript are small in comparison to e.g. that in the lidar ratio, and we decided to include a description in the methods section according to the suggestion of the reviewer. Our corrections make a significant difference for the retrieved aerosol scattering, especially in the LMS.

For example, they rapidly mentioned that the lidar ratio values of 50sr used to convert backscatter into extinction agree with Prata et al. (2017). This is not correct, Prata et al. (2017) found a mean lidar ratio of 69 sr for the Cordon plume, 66 for Kasatochi and 63 for Sarychev. This would increase the volcanic AOD during volcanically influenced periods by 30-40 %. The lidar ratio assumption is therefore one of the main source of uncertainty for AOD retrieved from CALIPSO but poorly discussed here.

We agree with the reviewer that the motivation for using the factor 50 sr was discussed too shortly.

The assumption of value on the lidar ratio is indeed a large uncertainty for the retrievals by CALIOP. The figures by Prata et al. (2017) are higher than those used in our manuscript, but according to the uncertainties reported in Prata et al. (2017) the factor 50 sr is not significantly different from the values reported in their paper, but we are on the lower side of their estimations. They reported

means and standard deviations for the lidar ratios of volcanic layers to be 69 ± 13 sr (Puyehue-Cordón Caulle, Jun 2011), 66 ± 19 sr (Kasatochi, Aug 2008) and 63 ± 14 sr (Sarychev).

There are differences in methodology in our manuscript and that of the Prata et al. (2017) paper. They found relatively fresh volcanic aerosol plumes in the stratosphere and computed the AOD from that, while we use zonal means, i.e. mixtures of layers of volcanic aerosol and the stratospheric background aerosol. Also, the values reported in Prata et al. (2017) are based on fresh aerosol. Aging of the aerosol and mixing with background changes the particle size distribution. Thus, the lidar ratio is not necessarily equal for the fresh volcanic aerosol, and aerosol that has been aged and mixed with the background. Most studies that dealt with stratospheric aerosol show lower lidar ratios than those reported in Prata et al., (2017). In a most recent study Thomason et al. (2018) find the CALIOP lidar ratio to have a value of 53 sr. We therefore believe that using the values reported in Jäger and Deshler (2003) is well-suited for our study.

We added the estimation on the (low) bias of CALIOP that have been computed from comparison to an airborne lidar (Kar et al., 2018). The bias has been significantly reduced in the latest version of the CALIOP data (V4), which is the one we used in the present study. We have included a more thorough discussion on the lidar ratio. We point out (in the manuscript) that we used a conservative value of the lidar ratio, along with the comparison to other AOD data-sets (GISS, CALIOP/GOMOS and OSIRIS from Rieger et al.(2015), the mean of OSIRIS, GOMOS, CALIOP and OMPS from Khaykin et al. (2017) and the GloSSAC (global space-based stratospheric aerosol climatology) based on CALIOP/OSIRIS from Thomason et al. (2018)).

4. English language. This is an overall issue which could be difficult to address without a native English speaker person. However, the level of English in the paper is relatively poor and would need to be improved. I recommend the co-authors of the paper to take part of this effort to improve the English.

We are aware that we as non-native speakers are in a difficult position language-wise. Reviewer #1 wrote that “...Overall the *writing is quite good, but there are a few awkward places...*”, and also provided a thorough list of suggested changes. We have corrected many language-mistakes by taking into account the many points made by Reviewer #1, and gone through the text with a critical eye. We believe that these changes, as well as all other changes, have improved the manuscript.

References

Andersson, S. M., Martinsson, B. G., Vernier, J. P., Friberg, J., Brenninkmeijer, C. A. M., Hermann, M., Van Velthoven, P. F. J. and Zahn, A.: Significant radiative impact of volcanic aerosol in the lowermost stratosphere, *Nat. Commun.*, 6(May), 1–8, doi:10.1038/ncomms8692, 2015.

Hostetler, C. a, Liu, Z., Reagan, J., Vaughan, M., Winker, D., Osborn, M., Hunt, W. H., Powell, K. a and Trepte, C.: CALIOP Algorithm Theoretical Basis Document - Calibration and Level 1 Data Products, , (PC-SCI-201 Release 1.0) [online] Available from: <http://www->

calipso.larc.nasa.gov/resources/pdfs/PC-SCI-201v1.0.pdf, 2006.

Jäger, H. and Deshler, T.: Erratum: Lidar backscatter to extinction, mass and area conversions for stratospheric aerosols based on midlatitude balloonborne size distribution measurements (Geophysical Research Letters (2002) 29:19 (1929) DOI:10.1029/2002GL015609), Geophys. Res. Lett., 30(7), 1–4, doi:10.1029/2003GL017189, 2003.

Kar, J., Vaughan, M. A., Lee, K. P., Tackett, J. L., Avery, M. A., Garnier, A., Getzewich, B. J., Hunt, W. H., Josset, D., Liu, Z., Lucker, P. L., Magill, B., Omar, A. H., Pelon, J., Rogers, R. R., Toth, T. D., Treppe, C. R., Vernier, J. P., Winker, D. M. and Young, S. A.: CALIPSO lidar calibration at 532 nm: Version 4 nighttime algorithm, Atmos. Meas. Tech., 11(3), 1459–1479, doi:10.5194/amt-11-1459-2018, 2018.

Khaykin, S. M., Godin-Beekmann, S., Keckhut, P., Hauchecorne, A., Jumelet, J., Vernier, J. P., Bourassa, A., Degenstein, D. A., Rieger, L. A., Bingen, C., Vanhellemont, F., Robert, C., DeLand, M. and Bhartia, P. K.: Variability and evolution of the midlatitude stratospheric aerosol budget from 22 years of ground-based lidar and satellite observations, Atmos. Chem. Phys., 17(3), 1829–1845, doi:10.5194/acp-17-1829-2017, 2017.

Prata, A. T., Young, S. A., Siems, S. T. and Manton, M. J.: Lidar ratios of stratospheric volcanic ash and sulfate aerosols retrieved from CALIOP measurements, Atmos. Chem. Phys., 17(13), 8599–8618, doi:10.5194/acp-17-8599-2017, 2017.

Rieger, L. A., Bourassa, A. E. and Degenstein, D. A.: Merging the OSIRIS and SAGE II stratospheric aerosol records, J. Geophys. Res., 120(17), 8890–8904, doi:10.1002/2015JD023133, 2015.

Solomon, S., Daniel, J. S., Neely, R. R., Vernier, J.-P., Dutton, E. G. and Thomason, L. W.: The Persistently Variable “Background” Stratospheric Aerosol Layer and Global Climate Change, Science (80-.), 333(6044), 866–870, doi:10.1126/science.1206027, 2011.

Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J. P., Manney, G., Luo, B., Arfeuille, F. and Peter, T.: A global space-based stratospheric aerosol climatology: 1979–2016, Earth Syst. Sci. Data, 10(1), 469–492, doi:10.5194/essd-10-469-2018, 2018.

Vernier, J. P., Thomason, L. W., Pommereau, J. P., Bourassa, A., Pelon, J., Garnier, A., Hauchecorne, A., Blanot, L., Treppe, C., Degenstein, D. and Vargas, F.: Major influence of tropical volcanic eruptions on the stratospheric aerosol layer during the last decade, Geophys. Res. Lett., 38(12), 1–8, doi:10.1029/2011GL047563, 2011.

Young, S. A., Winker, D. M., Noel, V., Vaughan, M. A., Hu, Y. and Kuehn, R. E.: Cloud-Aerosol Lidar Infrared Pathfinder Satellite Observations CALIOP Algorithm Theoretical Basis Document Document No: PC-SCI-201., 2005.

Changes made in manuscript

Language corrections and other minor changes, such as indexing of subfigures, were made that are included in the response to the reviewers’ comments. We have included the language changes suggested by reviewer 1, and made further language corrections adjusting to comment #4 by the second reviewer. At one point the indexing was incorrect, which we also changed.

We added the new Table 2 which summarizes our AOD comparisons to AODs in other studies.

The page- and line-numbers refer to the document with track-changes.

1.7-19. (page 1, lines 7-19). The abstract was updated to include e.g. findings in a new section of the manuscript.

2.12 and 2.15. OCS contributes to the Junge layer. Previously we stated that it was formed by OCS.

5.1-5.5. We added the estimation on the (low) bias of CALIOP that have been computed from comparison to an airborne lidar. The bias has been significantly reduced in the latest version of the CALIOP data (V4), which is the one we used in the present study.

5.8-6.6. We have made substantial changes in the methods section (section 3) including a survey on the values of the lidar ratio. We have also explained why we choose to use a value of 50 sr.

8.25-26. We have clarified the discussion on Fig 3, where the aerosol scattering (AS) is compared to the particle extinction provided in the CALIOP level 2 data.

11.27. We pointed out that it is the mean stratospheric AOD.

12.23. We clarified that we mean the feature at 20 km altitude.

14.16. Clarified that the lowest LMS layer is the 1.5-2 PVU.

14.23. Clarified why we used the 1.5 PVU as the dynamical tropopause.

15.6. Regarding the spring maximum in the ExTL. We used the suggestion of the reviewer and now writes that the “observations suggest that the spring/summer peaks in the lowest part of the LMS are caused by upwelling dust”.

15.7-9. We explain why the dust impact on the stratosphere is small.

16.4. Changed from “The upper layer has high AOD in the tropics that decreases to the poles” to “The upper layer has its highest AOD in the tropics and it decreases to the poles.”

16.27. We added a reference to figure 7.

17.5-7. We added a more descriptive text to clarify that the eruption of Calbuco may have impacted the upper stratospheric, the layer shown in figure 7a.

17.11-12. We have clarified why the increased AOD in northern midlatitudes most likely comes from subsidence from the overlying layer.

20.1-28. We wrote a new section on comparison to stratospheric AODs from other long-term studies, and included a new table (Table 2) summarizing the comparison. There is generally a good agreement between our AODs and those in other studies.

21.17-20. We included the findings from Table 2 in the conclusions section.

Volcanic impact on the climate – the stratospheric aerosol load in the period 2006 – 2015

Johan Friberg¹, Bengt G. Martinsson¹, Sandra M. Andersson^{1,2}, Oscar S. Sandvik¹

¹Division of Nuclear Physics, Lund University, Lund, 22100, Sweden

² Now at the Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

5 *Correspondence to:* Johan Friberg (johan.friberg@nuclear.lu.se)

Abstract. We present a study on the stratospheric aerosol load during 2006-2015, discuss the influence from volcanism and other sources, and reconstruct an [aerosol optical depth \(AOD\)](#) data-set in a resolution of 1° latitudinally and 8 days timewise. ~~Our attempt~~ [The purpose](#) is to include the “entire” stratosphere, from the tropopause to the almost particle free altitudes of the mid-stratosphere. A dynamic tropopause of 1.5 PVU was ~~used~~ [found to be the best suited one](#), [since it enclosed](#) almost all of the volcanic signals in the CALIOP data-set. ~~New methods were developed to handle bias in the data.~~ The data were successfully cleaned from polar stratospheric clouds using a temperature threshold of 195 K. Furthermore, a method was developed to correct data when the [CALIPSO-CALIOP](#) laser beam was strongly attenuated by volcanic aerosol, preventing a negative bias in the AOD data-set. Tropospheric influence, likely from upwelling dust, was found in the extratropical transition layer in spring. Eruptions of both extratropical and tropical volcanoes that injected aerosol into the stratosphere impacted the stratospheric aerosol load for up to a year if their clouds reached lower than 20 km altitudes, ~~whereas~~ [deeper reaching tropical injections rose in the tropical pipe and impacted it for several years.](#) [Our AODs mostly compare well to other long-term studies of the stratospheric AOD.](#) Over the years 2006-2015, volcanic eruptions increased the stratospheric AOD on average by ~40%. In absolute numbers the stratospheric AOD and radiative forcing amounted to 0.008 and -0.2 Wm⁻², respectively.

20 **1 Introduction**

Aerosol particles have a large impact on the Earth’s climate. [Trends in the abundance of aerosol particles are an important component of the climate system, although their influence on climate is still highly uncertain](#) ~~Trends in the abundance of aerosol particles remain as an important component in the climate system of which influence on climate is still highly uncertain~~ (IPCC, 2014). For example, volcanic eruptions inject particles and the sulfate forming precursor gas sulfur dioxide (SO₂) into the stratosphere, where [submicron](#) particles may remain for up to several years. There, the aerosol particles scatter and absorb solar radiation hindering it from heating the surface of the Earth. The eruption of Mount Pinatubo in 1991 resulted in a global cooling in excess of 0.5°C in the following years (McCormick et al., 1995).

In recent years much attention ~~(Fyfe et al., 2016; Rajaratnam et al., 2015; Trenberth, 2015; Xie et al., 2016; Yan et al., 2016)~~ has been drawn to a discrepancy between the Earth system models’ (CMIP5) projections of temperatures to that of an observed slow-down in the warming trend in the beginning of this century (Fyfe et al., 2013, 2016; Rajaratnam et al., 2015;

Trenberth, 2015; Xie et al., 2016; Yan et al., 2016), often called the hiatus. Part of the discrepancy was caused by bias in the temperature data (Karl et al., 2015). Likely reasons behind the remaining discrepancy are fluctuations in oceanic heat sequestration, solar blocking by volcanic aerosol in the stratosphere and solar forcing not captured by the state of the art models (Andersson et al., 2015; Medhaug et al., 2017; Meehl and Teng, 2014; Myhre et al., 2013; Santer et al., 2014; Solomon et al., 2011).

In the past decade several volcanic eruptions perturbed the stratospheric aerosol load, causing periods of increased stratospheric aerosol optical depth (AOD), that has been observed via remote sensing from satellites and ground (Khaykin et al., 2017; Sakai et al., 2016; Vernier et al., 2011; Zuev et al., 2017), and in-situ observations (Martinsson et al., 2017).

10

In the tropics, upwelling tropospheric air enters the stratosphere, bringing aerosol particles and precursor gases including SO₂ and carbonyl sulfide (OCS), ~~that form contributing to~~ the background stratospheric aerosol. It is still debated which of these components that dominates as source (Brühl et al., 2012; Sheng et al., 2015). While the water soluble SO₂ may be scavenged by cloud processes in convective transport, a large fraction of the OCS ~~is gets~~ transported to the stratosphere where it ~~is~~ oxidized ~~ds~~ by UV-radiation at ~25 km altitude ~~forming and contributes to~~ the Junge layer (Crutzen, 1976). A ~~most~~ recent study by Rollins et al. (2017) indicates that SO₂ concentrations in the tropical UT are too low for general tropical SO₂ upwelling to be a significant source of the stratospheric background aerosol.

Sulfate is the dominating constituent in the ~~stratospheric~~ background as well as in the volcanic aerosol (Deshler, 2008). In addition, organics (Friberg et al., 2014; Murphy et al., 2014), meteoritic matter, soil dust, and volcanic ash can be found in stratospheric aerosol particles. The ash component decreases ~~rapidly readily~~ after a volcanic eruption as the large ash particles ~~are get~~ gravitationally separated from the sulfate forming volcanic SO₂ layers and subside out of the stratosphere. ~~It thus~~ ~~Therefore it~~ generally constitutes only a small fraction of the stratospheric aerosol.

Sandwiched between the tropopause and the overlying 380 K isentrope, the ~~lowermost stratosphere (LMS)~~ constitutes the lowest ~~lying portion part~~ of the stratosphere. It is connected with the troposphere through isentropic surfaces crossing the tropopause, where significant bidirectional transport of air takes place (Holton et al., 1995). Local air-exchange occurs in a region extending a ~~couple offew~~ kilometers above the tropopause, termed the extra-tropical transition layer (ExTL; Hoor et al., 2002). ~~The ExTL#~~ is characterized by steep gradients in chemical species that have different concentration levels in the troposphere and stratosphere (Gettelman et al., 2011).

The Brewer-Dobson circulation transports air meridionally from the tropical stratosphere, in an overturning circulation (Gettelman et al., 1997). Most of the air is transported in a low altitude, or shallow, branch that spans an isentropic range of

380 K to ~450-470 K (Fueglistaler et al., 2009; Lin and Fu, 2013). A portion of the air ascends in the so-called tropical pipe. This high-altitude, or deep, branch extends to the stratopause (~50 km altitude) and transports the air slowly, resulting in residence times of several years (Bönisch et al., 2009). Air is then subsiding through the midlatitude and polar stratosphere, bringing aerosol particles down to the LMS from higher stratospheric altitudes that eventually end up in the troposphere, where it is ~~rapidly~~readily scavenged by cloud processes.

The CALIOP (Cloud-Aerosol Lidar with Orthogonal Polarization) instrument ~~onboard the CALIPSO satellite is based on the use of~~ a lidar to retrieve observations of the backscattering from aerosol particles and clouds at high vertical resolution (Winker et al., 2007). Vernier et al. (2009) developed a method of filtering out clouds, to study the backscattering from aerosol particles above 15 km altitude. The stratospheric AOD have been computed from the lidar backscattering from particles by assumptions on particle size distribution and composition (Jäger and Deshler, 2002, 2003). Solomon et al. (2011) estimated the radiative forcing induced by the injection of volcanic aerosol above 15 km altitude during year 2000 - 2010 to be on average ~-0.1 Wm⁻². Their approach excluded the lower-lying LMS, which contains approximately 40% of the stratospheric mass. Including the LMS, Ridley et al. (2014) and Andersson et al. (2015) revealed that the ~~lowermost stratosphere~~ (LMS) contains a significant fraction of the stratospheric aerosol, which influence on the AOD had been neglected in previous studies. Hence, for estimation of the full climate impact of volcanism one needs to consider the whole stratospheric column.

In this paper we present the volcanic influence on the “entire” stratospheric AOD (from the tropopause to 35 km altitude) over time and space, starting with studies on the transport of volcanic aerosol within the stratosphere. New techniques of handling the CALIOP data will be presented and discussed: one being the removal of signals from polar stratospheric clouds (PSC) and the other a means of correcting data in periods when the lidar signal ~~becomes~~gets attenuated by dense aerosol layers. Finally, the regional and global AODs are presented for the entire stratosphere in relation to transport patterns. As a means to represent the residence time of the aerosol from eruptions at several latitudes, reaching various altitudes, the AOD is presented for three stratospheric layers: the LMS, the potential temperature range of 380 to 470 K, and for altitudes above the 470 K isentrope.

2 The CALIOP data

This study of the stratospheric aerosol in the time period 2006 – 2015 is based on measurements with the CALIOP (Cloud-Aerosol Lidar with Orthogonal Polarization) instrument aboard the satellite CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation) in a joint mission of NASA and the French space agency, CNES. The almost nadir-viewing (3°) CALIOP is a two-wavelength lidar utilizing three receiver channels (Winker et al., 2007, 2009). One channel measures

backscatter intensity at 1064 nm and two channels measure orthogonally polarized backscatter intensity at 532 nm. The measurements result in high resolution vertical profiles of aerosols and clouds: in the altitude ranges 8.2 – 20.2, 20.2 – 30.1 and 30.1 – 40 km, ~~with respective the~~ vertical resolutions ~~of are~~ 60, 180 and 300 m, ~~respectively~~. During one day CALIPSO performs approximately 15 orbits between 82°S and 82°N with a repeat cycle of 16 days (Winker et al., 2010).

5

This work is based on the Level 1B (version 4-10) night-time products of the 532 nm perpendicular and parallel polarized channels. ~~Together, the~~The total from these two channels provides the backscatter intensity. Their ratio, ~~called the~~ polarization ratio, is used to infer the shape of the scattering objects. In the present study each swath measured by CALIOP was averaged over 1° in latitude and averaged or interpolated to 180 m in altitude. ~~The p~~Polarization ratios ~~was were~~ used to identify pixels containing clouds. From that a mask was formed for removal of pixels containing signals from clouds, using a 5% threshold in the polarization ratio (Vernier et al., 2009). The mask was expanded to remove weak signals from the edges of clouds, and data below thick clouds were excluded (Andersson et al., 2015); compare Figure 1b with Figure 1a for an example.

10

15

The measured backscattering intensity is the sum of that by aerosol particles and air molecules. ~~In order to~~To separate these two components, the molecular scattering of the air was modeled based on ozone number density, atmospheric temperature and pressure from the Global Modeling and Assimilation Office (GMAO; <http://gmao.gsfc.nasa.gov/>). In this study we use the ratio of the total scattering (β) to the modeled molecular scattering (β_m), the so-called scattering ratio (SR), and the difference between them, the aerosol scattering (AS):

20

$$SR = \frac{\beta}{\beta_m} \tag{1}$$

$$AS = \beta - \beta_m = (SR - 1) \cdot \beta_m \tag{2}$$

25 The SR is an optical equivalent to mixing ratio and the AS to aerosol concentration.

The measured backscattering intensity is ~~really~~an ‘attenuated backscattering’ (β'), due to light extinction of the laser pulses caused by molecules and particles. The attenuation that the laser pulses experience while passing through the atmosphere is estimated using the so-called two-way transmission parameter (T^2), which is the product of the two-way transmissions by the two components (molecules and particles). The molecular part is estimated from modelling data of the molecular background (Hostetler et al., 2006; Winker et al., 2009), while the ~~influence of the other~~attenuation from particles is generally ~~is~~considered negligible (e.g. Khaykin et al., 2017; Vernier et al., 2009). We will discuss this in section 5, along with a method we developed to correct for the increased attenuation occurring during periods when volcanism increased the

30

stratospheric aerosol load. The improved calibration in the version 4 (V4) data has reduced the bias of CALIOP, and by comparison to airborne lidar (the HSRL: High Spectral Resolution Lidar) Kar et al. (2018) estimate the bias to $1.6 \pm 2.4\%$. Up to 60% of the mean bias (1.6%) may have come from undetected clouds and aerosols, suggesting it to be even lower. We use the latest version of the CALIOP V4 data (V4-10), and therefore expect potential bias in β' to have low impact on our data analysis.

3. Computing the stratospheric AOD

~~The AOD as a function of latitude was obtained by integrating the AS in the vertical direction, and multiplying with the so-called lidar ratio, i.e. the extinction to backscattering ratio. In order to obtain the AOD from the CALIOP lidar measurements at 532 nm wavelength the aerosol particle size distribution is needed. The lidar ratio has been estimated in many studies, usually reported to span a range of values from ~40-70 sr for stratospheric aerosol (Jäger et al., 1995; Kremser et al., 2016; Prata et al., 2017 and references therein). In a thorough study Jäger and Deshler (2003) reported a value of 50 sr for background stratospheric aerosol concentration and size distribution occurring in the end of the 1990s. This value is commonly used for CALIOP (e.g. Hostetler et al., 2006a; Khaykin et al., 2017; Solomon et al., 2011) and with that value Khaykin et al. (2017) found good agreement between CALIOP, other satellite instruments and ground based lidar. In a recent long-term study, Thomason et al. (2018) found a CALIOP lidar ratio of 53 sr, by comparing to data from the OSIRIS instrument. Prata et al. (2017) found that lidar ratios of volcanic aerosol after the eruptions of Kasatochi (2008), Sarychev (2009) and Puyehue-Cordón Caulle (2011), where ~20-30% higher but not statistically differing from the values reported by Jäger and Deshler (2003) and Thomason et al. (2018). They used data from detected layers of relatively fresh volcanic aerosol, whereas we use zonal means where the volcanic and background aerosol are averaged together.~~

The aerosol particle size distribution is needed to obtain the AOD from the CALIOP lidar measurements. It is well established that large volcanic eruptions like the Mt Pinatubo in 1991 cause a shift towards larger particle sizes in the stratosphere (Deshler, 2008), resulting in a variable relation between lidar backscatter and extinction (Jäger and Deshler, 2003). Measurements in the LMS show that volcanic eruptions of Grimsvötn and Nabro in 2011 only had minor influence on the particle size distribution (Martinsson et al., 2014), and they were similar to size distributions obtained in periods of small volcanic impact (Jäger and Deshler, 2002). We will therefore use the more conservative, lidar ratio of 50 sr, corresponding to the background stratospheric aerosol (Jäger and Deshler, 2003), which is also close to the value of 53 sr retrieved by Thomason et al. (2018). Therefore, the background size distribution reported by the latter authors was used in this work. ~~The AOD as a function of latitude was obtained by integrating the AS in the vertical direction, and multiplying with the so-called lidar ratio, i.e. the extinction to backscattering ratio.~~

(2018)

We use a lidar ratio of 50 sr provided by Jäger and Deshler (2003). That value corresponds to the background stratospheric aerosol concentration and size distribution occurring in the end of the 1990s. In a recent study Prata et al. (2017) find lidar ratios after volcanic eruptions, of Kasatochi (2008), Sarychev (2009) and Puyehue Cordón Caulle (2011), to be somewhat higher but not significantly differing from the factor 50 sr, reported by Jäger and Deshler (2003).

Existing stratospheric AOD data-sets excludes the LMS. They are based on measurements where the AOD was integrated with the lower altitude limit being either the 380 K isentrope (Bourassa et al., 2012; Sato et al., 1993) or 15 km altitude (Solomon et al., 2011; Vernier et al., 2011). A first AOD estimation based on high resolution measurements that included the LMS were presented by Andersson et al. (2015). As the lower altitude limit of the AOD integration they used the tropopause supplied with the CALIOP data-data-set, a thermalstatic tropopause (Pan and Munchak, 2011) based on GEOS5 (Goddard Earth Observing System Model Version 5) data. The thermal tropopause according to the WMO definition on average resides more than 1 km above the dynamic tropopause of 1.5 PVU in the extra-tropics (Wilcox et al., 2012).

In the present study we use a dynamic tropopause as it captures the chemistry better than the thermal one does. This is reflected in that part of the volcanic aerosol in the present study residingresided below the thermal tropopause. The location of the dynamic tropopause was computed from potential vorticity (PV) values obtained from ERA-Interim reanalysis data provided by the ECMWF (European Center for Medium Ranged Weather Forecasts), and averaged to the same resolution as was the CALIOP data (1° latitudinally, 180 m vertically). Using a lower tropopause will of course increase the stratospheric AOD and the duration of effects on stratospheric AOD from volcanic eruptions. It will thereby provide a better estimate of the total stratospheric AOD. However, lowering the tropopause leads to more influence from the ExTL (Hoor et al., 2004), thus increasing the influence from tropospheric aerosol on the estimate of the stratospheric AOD. The location of the dynamic tropopause will be discussed more in section 6.1.2.

4 Handling PSCs

The Occasional presence of PSCs sometimes results in strong back-scattering signals in the winter polar stratosphere that can cause bias to the stratospheric AOD. This is especially problematic in the Southern Hemisphere where the PSCs occur at latitudes from 60°S. This is illustrated in Figure 1a as the high SR extending up to 25 km altitude in the Antarctic region. For investigation of the volcanic influence on the stratospheric aerosol, the influence from PSCs needs to be negligible, requiring a means of excluding data affected by these clouds. Andersson et al. (2015) solved this problem by manually omitting periods with PSCs. In the present study, we present a general approach to exclude the PSCs from the aerosol data.

Since PSC formation requires temperatures below 195 K, that temperature was used as a minimum threshold for a PSC mask applied to the Polar Regions, as means of minimizing the bias from PSCs. Furthermore, data below the clouds were removed to avoid bias from the strongly attenuated signals below the optically dense PSCs. This automated approach resulted in an almost complete removal of PSC signals, as illustrated by comparing Figure 1b and c, while keeping almost all of the data. Weak PSC signals remain, but its influence becomes negligible when averaged globally or hemispherically since the Antarctic region (60-90°S) constitutes only 6% of the Earth's surface area.

5 Correcting for errors caused by particle extinction

5.1 Attenuation of the laser

Passing through the atmosphere, the laser beam is attenuated by scattering and absorption from aerosol particles, molecules and O₃. Hence, the CALIOP instrument retrieves an 'attenuated backscattering' from laser pulses. Computing the true backscattering requires information on the extinction through the overlying atmosphere. The attenuation is accounted for by the two-way transmission (T^2), where an expression for the corrected backscattering becomes

$$\beta = \frac{\beta'}{T^2} \quad (3)$$

where β' is the attenuated backscattering retrieved by CALIOP. The two-way transmission depends on two attenuating components, i.e. that of molecules (T_m^2) and particles (T_p^2) (Young et al., 2005), and can be expressed as their product:

$$T^2 = T_m^2 \cdot T_p^2 \quad (4)$$

where the absorption from ozone molecules here is included in T_m^2 . By combining equations 1, 3 and 4, an expression for the SR corrected from both molecular extinction and particles becomes:

$$SR_{mpc} = \frac{\beta'}{T_m^2 \cdot T_p^2 \cdot \beta_m} \quad (5)$$

Attenuation from molecules is computed based on modelling, but that by particles are generally not considered in studies based on CALIOP data. For example, Vernier et al. (2009) and Khaykin et al. (2017) discuss that the attenuation from particles is less than 1% at 15 km altitude in absence of strong volcanic eruptions, arguing that corrections are unnecessary during these time periods. Assuming $T_p^2 = 1$, the expression for the SR corrected from molecular extinction becomes:

$$SR_{mC} = \frac{\beta'}{\tau_m^2 \cdot \beta_m} \quad (6)$$

from which the molecular extinction corrected AS (AS_{mC}) can be computed using eq.2.

5 Volcanic clouds may however result in non-negligible attenuation of the lidar signals. By neglecting the attenuation caused by particles, part of the aerosol signal will be accounted as signal coming from molecules, because the latter signal is also attenuated by the aerosol. Unaccounted for, the attenuation by volcanic aerosol particles can result in underestimation of the full effect of volcanic impact of the stratospheric aerosol load and the corresponding AOD. The attenuation is naturally increasing as the laser beams (and scattered light) passes through the atmosphere causing the largest errors to occur for
10 signals retrieved from the lower altitude side of volcanic clouds, and below them. In the present work this corresponds to the strongest attenuation being at the tropopause and in the UT.

In the present study, we first corrected the data by computing the two-way transmission caused by the attenuation from molecules (eq.6). This correction is sufficient in the absence of volcanic clouds, but light extinction caused by aerosol
15 particles in dense volcanic clouds resulted in further attenuation of the CALIOP laser. This is evident in [Figure 2](#) (black dashed line) where apparent decreases in SR arise in the UT after the volcanic eruptions of Sarychev, Nabro and Calbuco. The most dramatic decrease is observed for the SR data after the eruption of Calbuco in 2015, where strong attenuation resulted in unphysically low SR, i.e. below 1.

20 **5.2. Estimating the two-way transmission from particle extinction**

The two-way transmission from particles can be retrieved by a complicated technique where strongly attenuating features in the [CALIPSO/CALIOP](#) data are first identified and then an iterative process is used as means of estimating particle extinctions in attenuating layers (Young et al., 2005). The layer detection requires the use of a threshold value for identification of attenuating features. The resulting particle extinction parameter in the CALIOP Level 2 data is compared
25 with the AS in [Figure- 3](#). [The patterns of the Level 2 particle extinction \(Fig. 3a and c\) and that of the AS \(Fig. 3b and d\) do not match.](#) The algorithm obviously fails to detect most of the relatively fresh volcanic clouds (~~compare Fig 3a with 3b~~), and when the volcanic aerosol is mixed with the background aerosol 5 months after the eruption (Fig. 3c and d) the Level 2 procedure does not detect any attenuating aerosol. In [Figure- 2b](#) it is clear that the SR in the UT still is strangely lowered 5 months after the Calbuco eruption. We therefore developed a means of correcting for the attenuation caused by volcanic
30 aerosol particles, where the particle related two-way transmission (T_p^2) is calculated and applied ~~to~~ the SR_{mC} data. This procedure follows below.

The apparent decrease in the UT SR (Fig. 2) following the largest eruptions of the time period studied indicates attenuation of the laser signal induced by particles. By assuming the UT conditions to remain approximately unchanged after volcanic eruptions the degree of attenuation in a given volcanically perturbed time and place can be estimated by comparison with the signals in the years without strong volcanic influence. However, a volcanically perturbed stratosphere results in a small elevation of the UT aerosol concentration. That effect can be assumed small for most of the eruptions of this study, but the lower cloud of Kasatochi had a direct impact on the UT aerosol load (Andersson et al., 2015). In section 5.2.1 we present a method to handle that eruption separately. By assuming the UT conditions to remain approximately unchanged after all but the Kasatochi eruption, we can estimate the two-way transmission from particles at the altitude of the tropopause

$$\langle SR_{ut} \rangle_v = \langle SR_{ut} \rangle_{bg} \Rightarrow \frac{\langle SR_{mc,ut} \rangle_v}{\langle T_{p,ut}^2 \rangle_v} = \frac{\langle SR_{mc,ut} \rangle_{bg}}{\langle T_{p,ut}^2 \rangle_{bg}} \Rightarrow \frac{\langle T_{p,tp}^2 \rangle_v}{\langle T_{p,tp}^2 \rangle_{bg}} = \frac{\langle SR_{mc,ut} \rangle_v}{\langle SR_{mc,ut} \rangle_{bg}}$$

where $T_{p,tp}^2$ is the two-way transmission at the tropopause, and the indexes v and bg represent volcanic and background conditions. Furthermore, we assume that $\langle T_{p,tp}^2 \rangle_{bg} = 1$ since the particle extinction is small in background conditions, and we end up with

$$\langle T_{p,tp}^2 \rangle_v = \frac{\langle SR_{mc,ut} \rangle_v}{\langle SR_{mc,ut} \rangle_{bg}} \quad (7)$$

~~It is obvious that~~ T^2 values can theoretically ~~can~~ be in the range of 0 to 1. In this work $\langle T_{p,tp}^2 \rangle_v$ are distributed from 0.95 to 1, with the majority of values being >0.97 . Thus, the decreases in SR under volcanic clouds are mostly on the orders of 2-5%, which may appear to be negligibly low. For example, Figure 2a shows typical decreases in SR to be on the order of 0.025 in absolute terms, from for example $\langle SR_{mc,ut} \rangle_{bg}$ of 1.050 to $\langle SR_{mc,ut} \rangle_v$ of 1.025. Using eq.7 these values correspond to a $\langle T_{p,tp}^2 \rangle_v$ of 0.976. However, transferred to the AS and AOD these deviations become relevant. Using eq. 2, and assuming βm equal during volcanic periods and background, we end up with

$$\frac{\langle AS_{mc,ut} \rangle_v}{\langle AS_{mc,ut} \rangle_{bg}} = \frac{\langle SR_{mc,ut} \rangle_v - 1}{\langle SR_{mc,ut} \rangle_{bg} - 1} = \frac{1.025 - 1}{1.050 - 1} = 0.50$$

the $\langle AS_{mc,ut} \rangle_v$ and $\langle AS_{mc,ut} \rangle_{bg}$ being the ~~respective~~ AS in the UT during volcanic and background conditions. Thus, if the attenuation by particles is unaccounted for, the AOD ~~of~~ the UT gets underestimated by as much as 50% in the current example, since we retrieve the AOD linearly from the AS. A small effect on the SR is transformed into a large effect on the AS and AOD because before the correction half of the aerosol signal was accounted as signal from molecules in the example

above. This simple example illustrates that apparent tiny decreases in the T^2 may result in large underestimations in the computed AOD of low altitude layers, i.e. the LMS and especially the ExTL and UT. Once computed, the $\langle T_{p,tp}^2 \rangle_v$ are combined with the column integrated AS_{mc} data to compute two-way transmission matrices for the whole stratospheric column (T_p^2).

5

To compute T_p^2 values for the whole stratospheric column, the integrated column of stratospheric AS_{mc} values are used as a means of normalizing the T_p^2 values from the ‘top of the atmosphere’, here 35 km altitude, down to the tropopause. In this procedure the uppermost altitude bin gets a T_p^2 value of 1, altitude bins below get lower T_p^2 values, down to the tropopause where the $T_{p,tp}^2$ is computed from eq.7. We start by using the fraction that gets extinct by particles

10

$$Ext_p = 1 - T_p \quad (8)$$

and formulate an expression for estimating the fraction extinct in a single altitude bin (a)

$$15 \quad Ext_{p,a} = \frac{AS_{mc,a}}{\sum_{TP}^{TOA} AS_{mc}(z)} \cdot Ext_{p,tp} \quad (9)$$

where $Ext_{p,tp}$ is the extinct fraction at the tropopause ($1 - T_{p,tp}$), $AS_{mc,a}$ is the molecular corrected AS in altitude bin a , and the denominator holds the integrated backscattering for the total stratospheric column where z is the altitude bin. The transmission through the column can be computed by multiplying the transmission through each layer. Such approach may result in instabilities when the AS_{mc} values become close to zero or negative, as were the case after the eruption of Calbuco, i.e. the $SR_{mc} < 1$ in [Figure 2b](#) corresponds to negative AS_{mc} . By instead assuming linearity we can compute the extinction at each altitude simply by summing the extinction in each overlying layer. Combined with eq.8, this procedure yields the following expression for the altitude dependent two-way transmission from particles:

$$25 \quad T_p^2(z) = \left\{ 1 - \frac{\sum_{z+1}^{TOA} AS_{mc}(z)}{\sum_{TP}^{TOA} AS_{mc}(z)} (1 - T_{p,tp}) \right\}^2 \quad (10)$$

where z is the altitude bin, and TOA is the ‘top of the atmosphere’ (35 km altitude). This linear method produces T_p^2 values that in the current study deviate by $\ll 1\%$ from those computed by the multiplicative method, and the resulting AOD values deviate by $< 1\%$. However, the deviations grow with the size of the particle extinction values and may become more relevant for eruptions of the size of the 1991 Pinatubo eruption.

30

5.2.1 The Kasatochi eruption

Since Kasatochi's lower cloud increased the UT aerosol load the $\langle T_{p,tp}^2 \rangle_v$ could not be retrieved directly using eq.7 for that eruption. Instead we made use of the [similar extratropical eruption of Sarychev-eruption](#). For Sarychev we found the extinct fraction $(1 - \langle T_{p,tp} \rangle_{Sar})$ to be related to the amount of aerosol in the column, i.e. the column integrated AS ($\sum_{TP}^{TOA} AS_{Sar}$).

5 The $\langle T_{p,tp}^2 \rangle_v$ after Kasatochi was estimated indirectly from that relation:

$$\frac{\sum_{TP}^{TOA} AS_{Kas}}{1 - \langle T_{p,tp} \rangle_{Kas}} = \frac{\sum_{TP}^{TOA} AS_{Sar}}{1 - \langle T_{p,tp} \rangle_{Sar}} \quad (11)$$

where the indexes indicate the respective volcano. Combined with eq.8 and rearranging yields the expression for estimating

10 the UT particle related two-way transmission after the Kasatochi eruption

$$\langle T_{p,tp}^2 \rangle_{Kas} = \left\{ 1 - \left(\frac{\sum_{TP}^{TOA} AS_{Kas}}{\sum_{TP}^{TOA} AS_{Sar}} \right) \cdot \langle T_{p,tp} \rangle_{Sar} \right\}^2 \quad (12)$$

5.3. Correcting the data

Once the T_p^2 is known, the scattering ratios can be corrected in regard to the attenuation by particles by combining equations

15 5 and 6

$$SR_{mpc} = \frac{\beta'}{T_m^2 \cdot T_p^2 \cdot \beta_m} = \frac{SR_{mc}}{T_p^2} \quad (13)$$

and the corresponding aerosol scattering (AS_{mpc}) can be computed from insertion into eq.2. By this, our method introduces a

20 simple means of correcting the data to account for particle attenuation.

To study the sensitivity of the method, an iterative process was performed where the AS_{mpc} data were used to compute new T_p^2 values. We found only negligible changes in the SR_{mpc} and AS_{mpc} after these iterations. Thus, the developed procedure was found to be a robust method for correcting stratospheric CALIOP data for the attenuation caused by particle extinction.

25

Correcting the CALIOP data following volcanic eruptions, according to the equations above, significantly changes the SR as shown in Figure 2 (dashed vs. full lines). The global [mean](#) stratospheric AOD in the [8-first 8](#) months after eruptions increases [d](#) by [the percentages](#) [4%](#) (Kasatochi), [6%](#) (Sarychev), [6%](#) (Nabro), and Calbuco [7%](#) (Fig. 2b). These numbers are the result of relatively small volcanic elevations in the stratospheric aerosol load. The influence of particle extinction

naturally grows with the strength in elevation of the aerosol concentrations. Thus, eruptions of the size of the 1991 eruption of Mt Pinatubo, which was more than a magnitude larger than the eruptions of this study, would result in a strong need of correcting the data for the particle extinction.

6 Results

5 Large quantities of background sulfuric acid aerosol are produced deep into the stratosphere from OCS that is transported from the tropical troposphere in the Brewer-Dobson circulation (Crutzen, 1976) together with particles, SO₂ and other particle precursors from natural and anthropogenic sources. To this rather stable stratospheric background, special events inject large amounts of particles and precursor gases into the stratosphere, causing large variability in the stratospheric aerosol load. The most common cause of this variability is explosive volcanic eruptions (Robock, 2000), but ~~also~~ large fires
10 can occasionally influence the stratospheric aerosol (Fromm et al., 2010). For the time period studied here, mid-2006 to 2015, several volcanic eruptions and two large fires had such a potential. Table 1 lists these events starting more than a year before mid-2006 ~~in order~~ to account for the long residence time of volcanic aerosol going deep into the stratosphere, e.g. the aerosol load following the Mt Pinatubo eruption in 1991 declined over several years (McCormick et al., 1995).

6.1 The volcanic impact on the SR and AS

15 The stratospheric aerosol load varied substantially during the studied period as indicated in Figure 4, where the SR is illustrated in relation to latitude and altitude averaged for the months January, April, July and October over the years 2006 - 2015. Already in the first month, July 2006, a feature is seen in the tropics at 19 km altitude. This was caused by the eruption of the Soufriere Hills volcano in May 20, 2006 (Vernier et al., 2009), approximately one month before the start of measurements from CALIPSO. In October the same year a second feature appears to the south and at slightly lower altitude,
20 as a result of the eruption of Rabaul in October 7, 2006. The next major feature appears in October 2008. Aerosol from the eruption of the extratropical volcano Kasatochi in August 7, 2008 formed two layers at different altitudes. The influence from this eruption remained in January 2009, and in ~~the frame for~~ April more volcanic aerosol was added close to the tropopause by the eruptions of Redoubt in March and April 2009. A feature appearing above 20 km altitude in the southern tropics in April has been identified as smoke from bush fires in February 2009 in Victoria, Australia (Vernier et al., 2011). In
25 July 2009 the Northern Hemispheric stratosphere was perturbed again, this time by the extratropical volcano Sarychev (June 12, 2009), with the SR signal still clearly visible ~~until in~~ January 2010. In January 2011 aerosol from the eruption of the tropical volcano Merapi (November 5, 2010) had affected mainly the Southern Hemispheric stratosphere. ~~The SR of three volcanic eruptions can be observed~~ influenced the stratosphere in July 2011. The Southern Hemispheric Puyehue-Cordón Caulle (June 5, 2011) and the Northern Hemispheric Grimsvötn (May 21, 2011) affected the respective LMS, and aerosol
30 from the tropical Nabro (June 12, 2011) went to the north and affected the stratosphere during approximately one year. After

a couple of years with lower activity the tropical volcano Kelut erupted (February 13, 2014), reaching above 20 km altitude. The majority of the SR signal was confined to the tropics, but over a year, part of the aerosol was transported to the extratropics while the remaining tropical aerosol rose to higher altitudes. Finally, the Southern Hemispheric extratropical volcano Calbuco (April 23, 2015), induced the highest SR that occurred in the Southern Hemisphere during 2006-2015, which The effect of this eruption still remains at the end of the period.

Another interesting feature in Figure 4 is the variability at altitudes above 30 km, which is connected with the Quasi-Biennial Oscillation (QBO) (Vernier et al., 2011). The stratospheric temperature varies with the QBO, in the 30 – 50 hPa layer by almost 10°C, with the highest temperatures in the westerly shear (Baldwin et al., 2001). Modeling indicates that the QBO-associated temperature variability induces QBO-related altitude dependence in sulfuric acid evaporation from the aerosol (Hommel et al., 2015). Hence, at these altitudes less amount of aerosol is present during the westerly shear.

The SR deals with properties relative to the air mass, but from climatic point of view the AS, i.e. the absolute measure of the scattering by aerosols (eq.2), is of importance. Figure 5 shows the AS as a function of time and altitude in six latitude bands, each constituting 16% of the earth's surface area. The white lines mark the mean upper and lower limits of the LMS, i.e. the 380 K isentrope and the tropopause.

Elevated AS are clearly observed both after the stronger and weaker volcanic eruptions, and a striking feature in Figure 5 is the transport patterns where volcanic clouds in the tropic (extra-tropics) ascend (descend). Several eruptions reached above the 380 K, but it is also shown that a large portion of the aerosol was located in the LMS. Part of that aerosol came from the direct injection of volcanic clouds in the extratropics. Volcanic injections to higher altitudes eventually reach the LMS and due to the latitudinal transport and subsidence in the midlatitudes within the Brewer-Dobson circulation and subsiding in the midlatitudes, also the aerosol from tropical eruptions eventually ended up in the LMS. Furthermore, a seasonal variation is revealed in the strong signal from particles in the extra-tropical troposphere with the maximum AS occurring in spring/summer. A small part of this tropospheric source extends into the ExTL. Its possible influence on the stratospheric aerosol load will be discussed in section 6.3.

6.2 Using the dynamic tropopause

The dynamic tropopause was chosen to represent the lower boundary of the LMS as it is expected to best enclose the stratospheric air carrying volcanic aerosol. The PV of the dynamic tropopause varies between different studies (Gettelman et al., 2011), and is generally considered to lie somewhere in the range of 1.5 – 3.5 PVU (Hoerling et al., 1991; Hoinka, 1997; Kunz et al., 2011). The PV fields become vertical close to the equator. Therefore the 380 K isentrope is used as a limit of the dynamic tropopause's maximum possible altitude. Thus, the altitude of the dynamic tropopause becomes the lowest of the

380 K isentrope and that of a chosen PV-surface. The lowest commonly used dynamic tropopause, the 1.5 PVU surface, is on average located approximately 1.3 km below the thermal tropopause. Integration from this dynamic tropopause increases the total stratospheric AOD compared to when using the thermal tropopause. Here the aim is to use as low PV level as possible for best possible enclosure of the stratospheric volcanic aerosol. However, before deciding the tropopause level we need to consider the impact from tropospheric aerosol in the ExTL.

In [Figures 1 to 5](#), we assumed the 1.5 PVU level to be a good representation of the tropopause. Figure 6 illustrates a simple means of investigating the best suited PV value of a dynamic tropopause where the SR in approximately 0.25 km thick layers of the atmosphere above the 1.5 PVU level (LMS) are plotted for the midlatitudes of each hemisphere over the entire time frame of our study.

It is evident from ~~the~~ [Figure 6](#) that the strong volcanic eruptions of Sarychev and Calbuco induced strong gradients in the SR throughout the ExTL, and that volcanic influence is present down to the lowest PV-range, i.e. the 1.5-2 PVU. This is further corroborated by in situ particulate sulfur measurements (Martinsson et al., 2017). Unlike these eruptions, Kasatochi injected large amounts of volcanic aerosol in the lower parts of the LMS and the UT. As a result, the aerosol signal is strong also in the lowest LMS layer of [Figure: 6a \(1.5-2 PVU\)](#).

The local tropospheric sources' contribution to the stratospheric aerosol load are small in comparison to the signals of strong volcanic eruptions ~~such as those~~ from Kasatochi, Sarychev or Calbuco, making them negligible in the perspective of the stratosphere's total AOD. They do however have significant relative influence on the ExTL in the absence of strong volcanic eruptions.

[The value of 1.5 PVU is the lowest dynamic tropopause commonly used, and b](#)Based on the investigation above, a PV-value of 1.5 PVU was chosen to represent the dynamic tropopause in the following analyses, as it best captures the “full” volcanic impact on the stratosphere and on the climate.

6.3 Volcanic and tropospheric impact in the ExTL

Many interesting features can be seen in comparisons of the atmospheric slices in the [lower part of the LMS \(Ffig. 6\)](#) and of the two hemispheres. The largest peaks in the SR occurred after major volcanic eruptions in the extratropics (Kasatochi, Sarychev and Calbuco), which affected [their](#)s respective hemisphere. In the aftermath, the SR is highest deep into the LMS (at the highest PVs) and decreases down to the tropopause. This gradient is caused by gradual mixing of stratospheric air, carrying volcanic aerosol, with cleaner tropospheric air.

In the absence of major volcanic eruptions, the highest SR is found in spring close to the tropopause. Thus, the SR gradient is reversed compared to the periods dominated by volcanism, and it is more evident in the Northern than in the Southern Hemisphere. Figure 5 shows that the gradient is connected to a spring-/summertime increase in AS in the troposphere. Hence, we conclude that tropospheric local sources have significant influence in the ExTL during spring. These sources are obviously less important in the Southern Hemisphere. Air-craft measurements in the ExTL revealed that upwelling dust peaks in spring (Martinsson et al., 2005). These observations ~~indicate-suggest~~ that the spring/summer peaks in the lowest part of the LMS are caused by upwelling dust. Its impact on the total stratosphere is small, and is rapidly decreasing in strength in the first 2 km above the tropopause (1.5-5.5 PVU, the ExTL). Fluctuations caused by local tropospheric sources therefore have only minor influence on the stratospheric aerosol.

10 7 Discussions

The stratospheric AOD is obtained by converting aerosol scattering to extinction based on the particle size distribution and chemical composition (Jäger and Deshler, 2002, 2003), and integrating in the vertical direction. The patterns of volcanism are evident in Figure 7 where we divided the stratosphere into three layers for which we calculated AOD. This sub-division was based on transport patterns: the upper layer extends down to the 470 K isentrope, and represents the region where the latitudinal transport is weak (Fueglistaler et al., 2009; Lin and Fu, 2013), the LMS constitute the lowest layer, and in between is the mid-layer spanning isentropes of 380 – 470 K, where the shallow Brewer-Dobson branch is strong. This categorization of stratospheric layers will be used in the following discussions.

The temporal trends of the global and hemispheric mean AODs are compared in Figure 8 for the three layers and for that of the “entire” stratosphere. Even though the LMS is mostly ~~is~~-confined to the extratropics, and that altitudes above 470 K constitute a small portion of the stratospheric mass, it is evident that aerosol in these layers make up a significant portion of the global stratospheric AOD. In times of low volcanic impact the three layers contribute in approximately equal amounts to the total stratospheric AOD. Furthermore, Figures 7 and 8 show that the global stratospheric AOD reached its lowest values in the studied decade around the year 2013. This is in agreement with in-situ observations that find the stratospheric aerosol load to be at background conditions in 2013 (Martinsson et al., 2017).

~~The influence from spring time dust in the ExTL (section 6.3; Fig 6) on the total stratospheric aerosol AOD is very small (Fig 8). Thus, fluctuations caused by local tropospheric sources (upwelling dust) have only minor influence on the AOD estimation, and, in that respect, validates once again that the 1.5 PVU level is a well suited tropopause for studying the influence from volcanism.~~

We further explore the AOD distribution by investigating the strong latitudinal patterns shown in Figure 7. This is illustrated in Figure 9, where the AOD was averaged over different time periods, ~~starting with the entire decade~~ In Figure 9a, ~~There~~ we find the AOD to be distributed very differently in the respective layers. The upper layer has its highest AOD in the tropics and it that decreases to the poles. The LMS naturally shows highest AODs in the extratropics. The largest elevations in the LMS AOD relative to background conditions are observed in the Northern Hemisphere due to ~~the stronger~~ volcanic influence ~~being stronger there~~. The mid-layer is shown to have more evenly distributed AOD, with slightly higher AOD in the extra-tropics, and highest in the Northern Hemisphere. In ~~F~~figure 9b – d we separate the layers, to compare the influence from three types of volcanic eruptions to that of the decadal mean and that of the background conditions (year 2013).

7.1 The different eruption types

10 It is evident that the impact on the stratospheric AOD varied between eruptions and that only few eruptions reached into the uppermost layer. The eruptions that had significant impact on the stratospheric AOD were grouped into three categories depending on the observations in Figure 7. These types are as follows:

15 Trop I - Tropical eruptions with deep-reaching volcanic clouds, which were to a large degree incorporated into the deep Brewer-Dobson branch and ascended in the tropical pipe.

Trop II - Tropical eruptions with clouds that were confined in the mid-layer, transported in the shallow Brewer-Dobson branch, and not significantly transported in the deep branch.

20 Extrop - Volcanic clouds from extratropical eruptions. Some of these were confined within the LMS, and others partly penetrated into the mid-layer where the aerosol was incorporated into the shallow Brewer-Dobson branch and spread to the tropics.

25 The impact of the volcanic eruptions will be discussed based on these categories in the following sections, along with the two wildfires.

7.1.1 Deep reaching tropical volcanic eruptions

30 Two volcanoes significantly affected the upper layer (Fig. 7a), i.e Soufriere Hills (May 2006) and Kelut (February 2014). In addition, part of the elevations in AOD from summer 2006 may have come from the January 2005 Manam eruption (Vernier et al., 2009). The high reaching part of the Soufriere Hills cloud slowly ascended in the tropics. A strong elevation of the AOD is visible for more than one and a half years after the eruption after which the AOD decreases slowly over the following years until reaching its lowest values in 2013, ~~before the Kelut eruption~~. The Kelut volcano erupted ~~exploded~~ after

[this](#) period of stratospheric background levels, making it easy to track the spread of the volcanic aerosol. A large fraction of the volcanic cloud slowly ascended with the deep Brewer-Dobson branch in the tropical pipe, whereas the lower part of it was spread latitudinally in the mid-layer, mostly to the southern extratropics (Figures 4 and 9). The high-altitude part of the cloud was confined within the tropics. A small sudden increase in the AOD is observed in the southern extratropics (Fig. 7a) approximately one and a half years after the Kelut eruption. ~~That~~ ~~The~~ elevation [is shown as in Figure 7a and](#) was likely connected with [aerosol injected by](#) the midlatitude eruption of Calbuco (April 2015). [The aerosol load of upper layer show](#) ~~Minor-minor~~ fluctuations ~~which can be in the upper layer's aerosol load may have been~~ caused by evaporation/condensation of sulfuric acid (Vernier et al., 2011) in the upper part of the layer in connection with temperature variability induced by the quasi-biennial oscillation (Hommel et al., 2015). The aerosol in the upper layer is eventually transported out to the next lower layer at midlatitudes, i.e the one ~~confined located~~ between the 380-470 K isentropes. [We observe this in the 380-470K layer in northern midlatitudes \(Fig. 7b\), where the AOD was higher in the year 2007 and beginning of 2008 compared to the background \(year 2013\).](#)

7.1.2 Tropical eruptions below the 470 K isentrope

The volcanic clouds of the tropical eruptions from Rabaul (October 2006), Merapi (November 2010) and Nabro (June 2011) all penetrated the tropopause but did not reach altitudes above the 470 K isentrope (~20 km). While the two former eruptions impacted both hemispheres, the Nabro eruption mostly influenced the Northern Hemisphere (Figures 7 and 9). The small increase shown after Nabro in the Southern Hemisphere LMS (Fig. 9) was caused by the extratropical eruption of Puyehue-Cordón Caulle (June 2011). The aerosol from the Trop II eruptions was incorporated into the shallow Brewer-Dobson branch and spread to the midlatitudes within weeks. There the volcanic aerosol subsided and increased the AOD of the LMS, while decreasing in the mid-layer (Figures 7 and 8). The AOD in the mid-layer decreased over ~9 months in the case of the strong Nabro eruption, and the subsidence through the LMS resulted in several months prolonging of the volcanic impact of the eruption (Figures 7 and 8).

7.1.3 Extra-tropical volcanic clouds

Several extra-tropical eruptions impacted the Northern Hemispheric stratosphere, and some influenced the southern one. In the Northern Hemisphere, Kasatochi (August 2008) and Sarychev (June 2009) induced the strongest elevations in the AOD. Part of the aerosol from these eruptions reached the mid-layer, where it was spread to the tropical stratosphere within the shallow Brewer-Dobson branch. Figure 8 shows that some patterns of the global and Northern Hemispheric mean AOD in the mid-layer after Sarychev are similar to that of the Nabro eruption. The latitudinal distributions in the mid-layer shows similarities (Fig. 9c), but being a tropical eruption Nabro naturally impacted more in the tropics.

The Kasatochi eruption formed two clouds. Kasatochi's dense lower cloud was confined to the LMS and UT. Its stratospheric part was transported to the tropopause within approximately three months. The remaining weaker signal comes from the upper cloud that descended to the LMS (Fig. 8). The upper cloud reached the 380 – 470 K layer, and shows some similarities with that from the Sarychev and Nabro eruptions, having a rapid latitudinal transport in the lower BD branch. For all three eruptions, the elevation of the AOD remained until the following spring until the strong subsidence transported the aerosol down to the troposphere.

The Calbuco eruption (April 2015) was by far the largest one in the Southern Hemisphere. The volcanic cloud reached above the LMS, and induced a rapid strong elevation of the AOD of the mid-layer. The subsiding aerosol increased the AOD of the LMS so that it peaked a few months later than in the mid-layer. Calbuco reached higher altitudes than Kasatochi and Sarychev did, explaining the slower transport down to the LMS.

There were also a number of minor influences from extratropical volcanic eruptions that are shown as small increases in Figures 7 and 8, e.g. the eruptions of Redoubt (March 2009), Grimsvötn (May 2011) and Puyehue-Cordón Caulle (June 2011). Their clouds contained lower amounts of aerosol and did not penetrate as deep into the stratosphere, as the extratropical eruptions discussed above. Hence, the aerosol was rapidly transported out of the stratosphere, similarly to the lower cloud following the Kasatochi eruption.

7.1.4 The forest fires

The two forest fires affecting the stratosphere in this 10 year period had significantly lower impact on the stratospheric AOD than the volcanic eruptions. However, the fire in February 2009 reached altitudes of more than 20 km (Figures 4 and 7a). A concurrent small increase in the AOD is shown in the southern tropics upper layer (Fig. 7a), that mixed with the volcanic aerosol from the Soufriere Hills eruption. This fire therefore could have made a small, but long-term impact on the stratospheric AOD. The fire in December 2006 mainly affected the southern LMS, as evidenced in Figure 6c, with duration of a few months.

7.2 Patterns of volcanic aerosol in the stratosphere

Summarizing the findings of volcanic perturbations and transport within the stratosphere we make the following observations:

1. Aerosol from extratropical eruptions injected into the LMS remains there until transported out to the troposphere, e.g. the lower Kasatochi (2008) cloud. The AOD isgets elevated during-for less than a year.

2. Volcanic injections close to the extratropical tropopause only briefly impacts the stratospheric aerosol load, for example Grimsvötn (May 2011).
3. Aerosol from extratropical eruptions reaching the mid-layer gets dispersed hemispherically, and will be found at higher altitude in the tropics than their extratropical injection altitude, but no clear case of interhemispheric exchange. Examples are the upper Kasatochi cloud, the upper part of the Sarychev (2009) cloud and Calbuco (2015).
4. Volcanic clouds from tropical eruptions reaching above the 470 K isentrope (approximately 20 km altitude) tend to move upwards without a strong poleward transport, e.g. Soufriere Hills (2006) and Kelut (2014). Hence, the aerosol remains in the stratosphere for several years.
5. Volcanic clouds injected close to the tropical tropopause tend primarily to be transported poleward and within 1 – 2 months reach midlatitudes, e.g. Rabaul (2006), Merapi (2010) and Nabro (2011). The Nabro eruption shows that the AOD ~~was elevated for gets increased during up to~~ approximately one year. For the other two eruptions the signals are too weak for such estimations.
6. The last, clear indications of influence from extratropical as well as tropical volcanic eruptions appear in the LMS at mid- and high latitudes, as manifested by the three largest eruptions in this study (according to Table 1), i.e. Kasatochi, Sarychev and Nabro.

These observations agree well with the large-scale circulation pattern in the stratosphere. The general circulation of the stratosphere (Brewer-Dobson) is directed upwards in the tropics and downwards in the extratropics. As a consequence volcanic clouds injected into the LMS, which is not isentropically connected with other parts of the stratosphere, are transported downwards to the troposphere (point 1), and clouds injected close to the extratropical tropopause are rapidly removed due to that transport (2). Volcanic clouds reaching above approximately 20 km altitude in the tropics (4) get incorporated in the upper Brewer-Dobson branch and become relatively isolated from the extratropics for years (Fueglistaler et al., 2009), whereas tropical eruptions reaching lower altitudes are rapidly mixed meridionally (5) via the shallow Brewer-Dobson branch. This also explains the observation of reverse transport of extratropical volcanic clouds where aerosol reaching above the upper boundary of the LMS (3) is transported in the shallow Brewer-Dobson branch. Finally, the observation that the volcanic clouds of both tropical and extratropical eruptions leave the stratosphere via the LMS (6) is connected with the direction of the Brewer-Dobson circulation. Obviously, volcanic aerosol that reached the stratosphere ends up in the LMS sooner or later, no matter the injection latitude or altitude of the stratospheric injection.

7.3 Comparison to other AOD data-sets

Our AODs were compared with AODs reported in other long-term studies. The comparison was undertaken by forming annual means for various parts of the stratosphere, and is summarized in Table 2. In the comparison we have computed AODs for the same stratospheric regions in terms of latitude and altitude intervals as used in the referenced data, with the exception that our tropopause differs from that of Thomason et al (2018) by being in the order of 1 km lower altitude at midlatitudes. Our AODs are presented along with values on how much higher or lower the AODs are in the other studies. Khaykin et al. (2017) computed AODs between 17-30 km altitude at wavelength of 532 nm (same as that of CALIOP) by combining satellite and ground based instruments. Their values are very similar to ours. We also compared our data to AODs at wavelength of 525 nm (Rieger et al., 2015; Thomason et al., 2018), which are expected to be 2% higher than the AODs at 532 nm (Jäger and Deshler, 2002, 2003; Khaykin et al., 2017). Thomason et al., (2018) combined data from space based platforms to obtain the AOD of the “entire” stratosphere, and their study extends until 2009. We compare the annual means of 2007, 2008, and 2009 which were produced by OSIRIS and CALIOP in their data-set. Their southern midlatitude AODs are higher, and tropical AODs are slightly higher, than in our study. In the northern midlatitudes after the eruptions of Kasatochi (2008) and Sarychev (2009) the AODs from Thomason et al. (2018) are lower than AODs obtained in the present study. The relative importance of LMS aerosol in the Northern Hemisphere was elevated during 2008 and 2009, suggesting differences in the LMS aerosol quantification between their data-sets and ours. In part this can be explained by differences in tropopause definition, they using a monthly mean thermal tropopause as their lower limit, whereas we use the local 1.5 PVU dynamic tropopause situated at lower altitude. The OSIRIS data in Thomason et al. (2018) comes from the data-set presented in Rieger et al. (2015). The AODs from the merged data-set of SAGE II and OSIRIS in Rieger et al. (2015) are higher than ours. Their AOD data were converted from retrievals at a wavelength of 750 nm with commonly used assumptions on the aerosol properties. The CALIOP/GOMOS data, in the same paper show higher AODs than ours in 2007 and 2008, whereas the data agree well with ours from 2009-2013. Finally, the stratospheric AODs in the GISS climate model denoted OSIRIS/Sato in Table 2, also obtained from Rieger et al. (2015), are lower than those presented here.

Some obvious differences between the data-sets, such as differences in measurement techniques can explain part of the difference. Detailed knowledge on stratospheric aerosol composition, size distribution and optical properties is needed to improve the accuracy of stratospheric AOD retrievals from all the satellite-based instruments in this comparison. This problem requires further attention, but is outside the scope of this paper.

7.4.3 Climate relevance

Comparison of the mean AOD for 2006 – 2015 to that in the background (the year 2013) reveals that the volcanic eruptions increased the stratospheric AOD by ~40% (Table 32). The eruptions of the extratropical Sarychev and tropical Nabro volcanoes both increased the AOD with ~70% over the course of a year. The higher-reaching Kelut eruption had a smaller

initial influence, i.e. ~20% in the first year after eruption. It is expected to impact the stratospheric AOD over several years due to the slow transport in the deep Brewer-Dobson branch. Hence, averaged over longer time-spans the Kelut eruption likely caused an impact on the stratospheric AOD of similar size as the more sulfur-rich eruptions of Sarychev and Nabro.

5 The resulting radiative forcing was estimated as in Hansen et al. (2005) and Solomon et al. (2011), using a conversion factor of -25 for AOD to radiative forcing, and is added to Figure 8 as a secondary y-axis. The global radiative forcing from stratospheric aerosol ranged from approximately -0.15 in the background to -0.35 Wm^{-2} after the strongest volcanic eruptions of the period (Fig. 8). Most of the elevation in radiative forcing appeared in the Northern Hemisphere, and the lowest occurring during the year 2013. On average the global stratospheric radiative forcing amounted to -0.2 Wm^{-2} during a time-
10 period of a decade (Table 32).

8 Conclusions

We present a study on the stratospheric aerosol optical depth (AOD) and radiative forcing over a period of almost a decade (mid 2006 - 2015), covering periods of varying volcanic impact as well as stratospheric background conditions, with a resolution of 1° latitudinally and 8 days timewise. This required development of new methods in order to prevent influence
15 from polar stratospheric clouds, and to correct data when the lidar was attenuated by volcanic aerosol. The latter correction increased the AOD by 4-7% in the first year after the volcanic eruptions of Kasatochi (2008), Sarychev (2009), Nabro (2011) and Calbuco (2015). We find that our stratospheric AODs generally compare well with values reported in other long-term studies. This comparison was undertaken for various parts of the stratosphere. Looking at the reported total stratospheric AODs the results reported here usually are the highest, because we, in contrast to most of the other data-sets, include the
20 LMS.

Strong volcanic impact was found in the extra-tropical tropopause layer (ExTL) down to potential vorticities (PV) of 1.5 PVU, i.e. more than 1 km below the thermal tropopause. We therefore used the 1.5 PVU level as the tropopause in our analysis to include the full impact of volcanism. In spring/summer upwelling dust clearly elevated the aerosol load, up to
25 PV-levels of 5-6 PVU, i.e. ~2 km above the dynamic tropopause, but it had insignificant influence on the total AOD of the lowermost stratosphere (LMS).

The stratospheric AOD was studied by dividing the stratosphere into three layers that incidentally carry approximately the same global AOD during conditions close to the stratospheric background; the LMS, the altitude range between the 380 and
30 470 K isentropes, and altitudes above the 470 K isentrope (~20 – 35 km). Several eruptions were found to influence the two lower layers, both extratropical and tropical ones.

Only the high-reaching aerosol from the tropical volcanoes Soufriere Hills (May 2006) and Kelut (February 2014) clearly impacted the upper layer (>470 K). Their volcanic clouds were first observed at altitudes of ~20 km, after which the clouds rose to higher altitudes incorporated in the deep Brewer-Dobson branch, impacting on the stratospheric AOD over several years. After the Soufriere Hills eruption the aerosol gradually decreased over the following years, which also has been observed for the far stronger Pinatubo eruption (1991). A similar decay was observed after the Kelut eruption, up till the end of the period studied here.

Volcanic clouds reaching into the mid-layer (380 - 470K) were not found to rise, but spread latitudinally in the shallow Brewer-Dobson branch, before being transported down to the LMS and eventually out of the stratosphere in the extratropics. For example, aerosol from the eruptions of the extratropical volcanoes Sarychev (June 2009) and Calbuco (April 2015) spread to the tropics within weeks, whereas aerosol from the eruption of the tropical volcano Nabro (June 2011) spread in the opposite direction followed by subsiding to the LMS. Such transport was limited within a Hemisphere and impacted the stratosphere for up to a year.

The stratospheric AOD was elevated the most in the extratropics, due to the combined effect of latitudinal transport and the larger stratospheric column. The majority of that elevation came from aerosol located in the LMS. Subsidence through the LMS causes the AOD to remain elevated in the LMS for several months after that the overlying stratosphere has returned to its background aerosol levels.

We have included the LMS in an estimation of the AOD of the “entire” stratosphere over the period 2006 – 2015. The stratospheric background AOD and the volcanic impact were found to be ~50% higher in midlatitudes than in the tropics. Volcanism was found to have elevated the average global stratospheric AOD by ~40%. The stratospheric aerosol had a cooling effect of the Earth, which in terms of radiative forcing, is estimated to -0.2 Wm^{-2} .

Acknowledgements

Financial support from the Swedish National Space Board (contract 130/15) and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (contract 942-2015-995) is gratefully acknowledged. Aerosol products from the CALIOP sensor were produced by NASA Langley Research Center.

References

- Andersson, S. M., Martinsson, B. G., Vernier, J. P., Friberg, J., Brenninkmeijer, C. A. M., Hermann, M., Van Velthoven, P. F. J. and Zahn, A.: Significant radiative impact of volcanic aerosol in the lowermost stratosphere, *Nat. Commun.*, 6(May), 1–8, doi:10.1038/ncomms8692, 2015.
- 5 Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnersley, J. S., Marquardt, C., Sato, K. and Takahashi, M.: The quasi-biennial oscillation, *Rev. Geophys.*, 39(2), 179–229, doi:10.1029/1999RG000073, 2001.
- Bourassa, A. E., Robock, A., Randel, W. J., Deshler, T., Rieger, L. A., Lloyd, N. D., Llewellyn, E. J. (Ted) and Degenstein, D. A.: Large Volcanic Aerosol Load in the Stratosphere Linked to Asian Monsoon Transport, *Science* (80-.), 337(6090), 78
- 10 LP-81 [online] Available from: <http://science.sciencemag.org/content/337/6090/78.abstract>, 2012.
- Brühl, C., Lelieveld, J., Crutzen, P. J. and Tost, H.: The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate, *Atmos. Chem. Phys.*, 12(3), 1239–1253, doi:10.5194/acp-12-1239-2012, 2012.
- Bönisch, H., Engel, A., Curtius, J. and Hoor, P.: Quantifying transport into the lowermost stratosphere using simultaneous in-situ measurements of SF₆ and CO₂, *Atmos. Chem. Phys.*, 5905–5919, 2009.
- 15 Carn, S. A. and Prata, F. J.: Satellite-based constraints on explosive SO₂ release from Soufrière Hills Volcano, Montserrat, *Geophys. Res. Lett.*, 37(18), 1–5, doi:10.1029/2010GL044971, 2010.
- Carn, S. A., Paluster, J. S., Lara, L., Ewert, J. W., Watt, S., Prata, A. J., Thomas, R. J. and Villarosa, G.: The Unexpected Awakening of Chaitén Volcano, Chile, *Eos* (Washington, DC), 90(24), 205–206, doi:10.1029/2009EO240001, 2009.
- Clarisse, L., Hurtmans, D., Clerbaux, C., Hadji-Lazaro, J., Ngadi, Y. and Coheur, P. F.: Retrieval of sulphur dioxide from the
- 20 infrared atmospheric sounding interferometer (IASI), *Atmos. Meas. Tech.*, 5(3), 581–594, doi:10.5194/amt-5-581-2012, 2012.
- Crutzen, P. J.: The possible importance of COS for the sulphate layer of the stratosphere, *Geophys. Res. Lett.*, 3(2), 73–76, 1976.
- Deshler, T.: A review of global stratospheric aerosol: Measurements, importance, life cycle, and local stratospheric aerosol,
- 25 *Atmos. Res.*, 90(2–4), 223–232, doi:10.1016/j.atmosres.2008.03.016, 2008.
- Friberg, J., Martinsson, B. G., Andersson, S. M., Brenninkmeijer, C. A. M., Hermann, M., Van Velthoven, P. F. J. and Zahn, A.: Sources of increase in lowermost stratospheric sulphurous and carbonaceous aerosol background concentrations during 1999–2008 derived from CARIBIC flights, *Tellus, Ser. B Chem. Phys. Meteorol.*, 66(1), doi:10.3402/tellusb.v66.23428, 2014.
- 30 Fromm, M., Lindsey, D. T., Servranckx, R., Yue, G., Trickl, T., Sica, R., Doucet, P. and Godin-Beekmann, S.: The untold story of pyrocumulonimbus, *Bull. Am. Meteorol. Soc.*, 91(9), 1193–1209, doi:10.1175/2010BAMS3004.1, 2010.
- Fueglistaler, S., Dessler, a E., Dunkerton, T. J., Folkins, I., Fu, Q. and Ote, P. W.: Tropical tropopause layer, *Rev. Geophys.* 47, RG1004, doi(2008), 101029/, doi:10.1029/2008RG000267.1.INTRODUCTION, 2009.

- Fyfe, J. C., Gillett, N. P. and Zwiers, F. W.: Overestimated global warming over the past 20 years, *Nat. Clim. Chang.*, 3(9), 767–769, doi:10.1038/nclimate1972, 2013.
- Fyfe, J. C., Meehl, G. A., England, M. H., Mann, M. E., Santer, B. D., Flato, G. M., Hawkins, E., Gillett, N. P., Xie, S. P., Kosaka, Y. and Swart, N. C.: Making sense of the early-2000s warming slowdown, *Nat. Clim. Chang.*, 6(3), 224–228, doi:10.1038/nclimate2938, 2016.
- 5 Gettelman, A., Holton, J. R. and Rosenlof, K. H.: Mass fluxes of O₃, CH₄, N₂O and CF₂Cl₂ in the lower stratosphere calculated from observational data, *J. Geophys. Res.*, 102(D15), 19149, doi:10.1029/97JD01014, 1997.
- Gettelman, a., Pan, L. L., Randel, W. J., Hoor, P., Birner, T. and Hegglin, M. I.: the Extratropical Upper Troposphere and Lower Stratosphere, *Rev. Geophys.*, 49(3), 1–31, doi:10.1029/2011RG000355.1.INTRODUCTION, 2011.
- 10 Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G., Aleinov, I., Bauer, M., Bauer, S., Bell, N., Cairns, B., Canuto, V., Chandler, M., Cheng, Y., Del Genio, A., Faluvegi, G., Fleming, E., Friend, A., Hall, T., Jackman, C., Kelley, M., Kiang, N., Koch, D., Lean, J., Lerner, J., Lo, K., Menon, S., Miller, R., Minnis, P., Novakov, T., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D., Romanou, A., Shindell, D., Stone, P., Sun, S., Tausnev, N., Thresher, D., Wielicki, B., Wong, T., Yao, M. and Zhang, S.: Efficacy of climate forcings, *J. Geophys. Res. D Atmos.*, 110(18), 1–45, doi:10.1029/2005JD005776, 2005.
- 15 Haywood, J. M., Jones, A., Clarisse, L., Bourassa, A., Barnes, J., Telford, P., Bellouin, N., Boucher, O., Agnew, P., Clerbaux, C., Coheur, P., Degenstein, D. and Braesicke, P.: Observations of the eruption of the Sarychev volcano and simulations using the HadGEM2 climate model, *J. Geophys. Res. Atmos.*, 115(21), 1–18, doi:10.1029/2010JD014447, 2010.
- Hoerling, M. P., Schaack, T. K. and Lenzen, A. J.: Global Objective Tropopause Analysis, *Mon. Weather Rev.*, 119(8), 20 1816–1831, doi:10.1175/1520-0493(1991)119<1816:GOTA>2.0.CO;2, 1991.
- Hoinka, K. P.: The tropopause: Discovery, definition and demarcation, *Meteorol. Zeitschrift*, 6(6), 281–303, 1997.
- Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B. and Pfister, L.: Stratosphere-troposphere exchange, *Rev. Geophys.*, 33(4), 403–439, doi:10.1029/95RG02097, 1995.
- Hommel, R., Timmreck, C., Giorgetta, M. A. and Graf, H. F.: Quasi-biennial oscillation of the tropical stratospheric aerosol 25 layer, *Atmos. Chem. Phys.*, 15(10), 5557–5584, doi:10.5194/acp-15-5557-2015, 2015.
- Hoor, P., Fischer, H., Lange, L., Lelieveld, J. and Brunner, D.: Seasonal variations of a mixing layer in the lowermost stratosphere as identified by the CO-O₃ correlation from in situ measurements, *J. Geophys. Res. Atmos.*, 107(D5), ACL 1-1-ACL 1-11, doi:10.1029/2000JD000289, 2002.
- Hoor, P., Gurk, C., Brunner, D., Hegglin, M. I., Wernli, H. and Fischer, H.: Seasonality and extent of extratropical TST 30 derived from in-situ CO measurements during SPURT, *Atmos. Chem. Phys. Discuss.*, 4(2), 1691–1726, doi:10.5194/acpd-4-1691-2004, 2004.
- Hostetler, C. a, Liu, Z., Reagan, J., Vaughan, M., Winker, D., Osborn, M., Hunt, W. H., Powell, K. a and Trepte, C.: CALIOP Algorithm Theoretical Basis Document - Calibration and Level 1 Data Products, , (PC-SCI-201 Release 1.0)

- [online] Available from: <http://www-calipso.larc.nasa.gov/resources/pdfs/PC-SCI-201v1.0.pdf>, 2006.
- IPCC Working Group 1: IPCC Fifth Assessment Report (AR5) - The physical science basis., 2014.
- Jäger, H. and Deshler, T.: Lidar backscatter to extinction, mass and area conversions for stratospheric aerosols based on midlatitude balloonborne size distribution measurements, *Geophys. Res. Lett.*, 29(19), 35-1-35-4, doi:10.1029/2002GL015609, 2002.
- Jäger, H. and Deshler, T.: Erratum: Lidar backscatter to extinction, mass and area conversions for stratospheric aerosols based on midlatitude balloonborne size distribution measurements (*Geophysical Research Letters* (2002) 29:19 (1929) DOI:10.1029/2002GL015609), *Geophys. Res. Lett.*, 30(7), 1–4, doi:10.1029/2003GL017189, 2003.
- Jäger, H., Deshler, T. and Hofmann, D. J.: Midlatitude lidar backscatter conversions based on balloonborne aerosol measurements, *Geophys. Res. Lett.*, 22(13), 1729–1732, doi:10.1029/95GL01521, 1995.
- Kar, J., Vaughan, M. A., Lee, K. P., Tackett, J. L., Avery, M. A., Garnier, A., Getzewich, B. J., Hunt, W. H., Josset, D., Liu, Z., Lucker, P. L., Magill, B., Omar, A. H., Pelon, J., Rogers, R. R., Toth, T. D., Trepte, C. R., Vernier, J. P., Winker, D. M. and Young, S. A.: CALIPSO lidar calibration at 532 nm: Version 4 nighttime algorithm, *Atmos. Meas. Tech.*, 11(3), 1459–1479, doi:10.5194/amt-11-1459-2018, 2018.
- Karl, T. R., Arguez, A., Huang, B., Lawrimore, J. H., McMahon, J. R., Menne, M. J., Peterson, T. C., Vose, R. S. and Zhang, H.-M.: Possible artifacts of data biases in the recent global surface warming hiatus, *Science* (80-.), 348(6242), 1469–1472, doi:10.1126/science.aaa5632, 2015.
- Khaykin, S. M., Godin-Beekmann, S., Keckhut, P., Hauchecorne, A., Jumelet, J., Vernier, J. P., Bourassa, A., Degenstein, D. A., Rieger, L. A., Bingen, C., Vanhellemont, F., Robert, C., DeLand, M. and Bhartia, P. K.: Variability and evolution of the midlatitude stratospheric aerosol budget from 22 years of ground-based lidar and satellite observations, *Atmos. Chem. Phys.*, 17(3), 1829–1845, doi:10.5194/acp-17-1829-2017, 2017.
- Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M., Stenke, A., Schwarz, J. P., Weigel, R., Fueglistaler, S., Prata, F. J., Vernier, J. P., Schlager, H., Barnes, J. E., Antuña-Marrero, J. C., Fairlie, D., Palm, M., Mahieu, E., Notholt, J., Rex, M., Bingen, C., Vanhellemont, F., Bourassa, A., Plane, J. M. C., Klocke, D., Carn, S. A., Clarisse, L., Trickl, T., Neely, R., James, A. D., Rieger, L., Wilson, J. C. and Meland, B.: Stratospheric aerosol—Observations, processes, and impact on climate, *Rev. Geophys.*, 54(2), 278–335, doi:10.1002/2015RG000511, 2016.
- Kunz, A., Konopka, P., Müller, R. and Pan, L. L.: Dynamical tropopause based on isentropic potential vorticity gradients, *J. Geophys. Res. Atmos.*, 116(1), doi:10.1029/2010JD014343, 2011.
- Li, C., Krotkov, N. A., Carn, S., Zhang, Y., Spurr, R. J. D. and Joiner, J.: New-generation NASA Aura Ozone Monitoring Instrument (OMI) volcanic SO₂ dataset: Algorithm description, initial results, and continuation with the Suomi-NPP Ozone Mapping and Profiler Suite (OMPS), *Atmos. Meas. Tech.*, 10(2), 445–458, doi:10.5194/amt-10-445-2017, 2017.
- Lin, P. and Fu, Q.: Changes in various branches of the Brewer-Dobson circulation from an ensemble of chemistry climate models, *J. Geophys. Res. Atmos.*, 118(1), 73–84, doi:10.1029/2012JD018813, 2013.

- Lopez, T., Carn, S., Werner, C., Fee, D., Kelly, P., Doukas, M., Pfeffer, M., Webley, P., Cahill, C. and Schneider, D.: Evaluation of Redoubt Volcano's sulfur dioxide emissions by the Ozone Monitoring Instrument, *J. Volcanol. Geotherm. Res.*, 259, 290–307, doi:10.1016/j.jvolgeores.2012.03.002, 2013.
- Martinsson, B. G., Nguyen, H. N., Brenninkmeijer, C. A. M., Zahn, A., Heintzenberg, J., Hermann, M. and van Velthoven, P. F. J.: Characteristics and origin of lowermost stratospheric aerosol at northern midlatitudes under volcanically quiescent conditions based on CARIBIC observations, *J. Geophys. Res. D Atmos.*, 110(12), 1–12, doi:10.1029/2004JD005644, 2005.
- Martinsson, B. G., Friberg, J., Andersson, S. M., Weigelt, A., Hermann, M., Assmann, D., Voigtländer, J., Brenninkmeijer, C. A. M., Van Velthoven, P. J. F. and Zahn, A.: Comparison between CARIBIC aerosol samples analysed by accelerator-based methods and optical particle counter measurements, *Atmos. Meas. Tech.*, 7(8), 2581–2596, doi:10.5194/amt-7-2581-2014, 2014.
- Martinsson, B. G., Friberg, J., Sandvik, O. S., Hermann, M., Van Velthoven, P. F. J. and Zahn, A.: Particulate sulfur in the upper troposphere and lowermost stratosphere - Sources and climate forcing, *Atmos. Chem. Phys.*, 17(18), 10937–10953, doi:10.5194/acp-17-10937-2017, 2017.
- McCormick, M. P., Thomason, L. W. and Trepte, C. R.: Atmospheric effects of the Mt Pinatubo eruption, *Nature*, 373(6513), 399–404, doi:10.1038/373399a0, 1995.
- Medhaug, I., Stolpe, M. B., Fischer, E. M. and Knutti, R.: Reconciling controversies about the “global warming hiatus,” *Nature*, 545(7652), 41–47, doi:10.1038/nature22315, 2017.
- Meehl, G. A. and Teng, H.: CMIP5 multi-model hindcasts for the mid-1970s shift and early 2000s hiatus and predictions for 2016-2035, *Geophys. Res. Lett.*, 41(5), 1711–1716, doi:10.1002/2014GL059256, 2014.
- Murphy, D. M., Froyd, K. D., Schwarz, J. P. and Wilson, J. C.: Observations of the chemical composition of stratospheric aerosol particles, *Q. J. R. Meteorol. Soc.*, 140(681), 1269–1278, doi:10.1002/qj.2213, 2014.
- Myhre, G., Shindell, D., Bréon, F., Collins, W., Fuglestedt, J., Huang, J., Koch, D., Lamarque, J., Lee, D., Mendoza, B. and others: Anthropogenic and natural radiative forcing, , 658–740, doi:0.1017/CBO9781107415324.018, 2013.
- Pan, L. L. and Munchak, L. A.: Relationship of cloud top to the tropopause and jet structure from CALIPSO data, *J. Geophys. Res. Atmos.*, 116(12), doi:10.1029/2010JD015462, 2011.
- Pardini, F., Burton, M., Arzilli, F., La Spina, G. and Polacci, M.: Satellite-derived SO₂ flux time-series and magmatic processes during the 2015 Calbuco eruptions, *Solid Earth Discuss.*, (July), 1–25, doi:10.5194/se-2017-64, 2017.
- Prata, A. J. and Bernardo, C.: Retrieval of volcanic SO₂ column abundance from Atmospheric Infrared Sounder data, *J. Geophys. Res. Atmos.*, 112(20), 1–17, doi:10.1029/2006JD007955, 2007.
- Prata, A. T., Young, S. A., Siems, S. T. and Manton, M. J.: Lidar ratios of stratospheric volcanic ash and sulfate aerosols retrieved from CALIOP measurements, *Atmos. Chem. Phys.*, 17(13), 8599–8618, doi:10.5194/acp-17-8599-2017, 2017.
- Rajaratnam, B., Romano, J., Tsiang, M. and Diffenbaugh, N. S.: Debunking the climate hiatus, *Clim. Change*, 133(2), 129–140, doi:10.1007/s10584-015-1495-y, 2015.

- Ridley, D. A., Solomon, S., Barnes, J. E., Burlakov, V. D., Deshler, T., Dolgii, S. I., Herber, A. B., Nagai, T., Neely, R. R., Nevzorov, A. V., Ritter, C., Sakai, T., Santer, B. D., Sato, M., Schmidt, A., Uchino, O. and Vernier, J. P.: Total volcanic stratospheric aerosol optical depths and implications for global climate change, *Geophys. Res. Lett.*, 41(22), 7763–7769, doi:10.1002/2014GL061541, 2014.
- 5 Rieger, L. A., Bourassa, A. E. and Degenstein, D. A.: Merging the OSIRIS and SAGE II stratospheric aerosol records, *J. Geophys. Res.*, 120(17), 8890–8904, doi:10.1002/2015JD023133, 2015.
- Robock, A.: Volcanic eruptions and climate, *Rev. Geophys.*, 38(2), 191–219, doi:10.1029/1998RG000054, 2000.
- Rollins, A. W., Thornberry, T. D., Watts, L. A., Yu, P., Rosenlof, K. H., Mills, M., Baumann, E., Giorgetta, F. R., Bui, T. V., Höpfner, M., Walker, K. A., Boone, C., Bernath, P. F., Colarco, P. R., Newman, P. A., Fahey, D. W. and Gao, R. S.: The
10 role of sulfur dioxide in stratospheric aerosol formation evaluated by using in situ measurements in the tropical lower stratosphere, *Geophys. Res. Lett.*, 44(9), 4280–4286, doi:10.1002/2017GL072754, 2017.
- Sakai, T., Uchino, O., Nagai, T., Liley, B., Morino, I. and Fujimoto, T.: Long-term variation of stratospheric aerosols observed with lidars over Tsukuba, Japan, from 1982 and Lauder, New Zealand, from 1992 to 2015, *J. Geophys. Res.*, 121(17), 10283–10293, doi:10.1002/2016JD025132, 2016.
- 15 Santer, B. D., Bonfils, C., Painter, J. F., Zelinka, M. D., Mears, C., Solomon, S., Schmidt, G. A., Fyfe, J. C., Cole, J. N. S., Nazarenko, L., Taylor, K. E. and Wentz, F. J.: Volcanic contribution to decadal changes in tropospheric temperature, *Nat. Geosci.*, 7(3), 185–189, doi:10.1038/ngeo2098, 2014.
- Sato, M., Hansen, J. E., McCormick, M. P. and Pollack, J. B.: Stratospheric aerosol optical depths, 1850–1990, *J. Geophys. Res.*, 98(D12), 22987, doi:10.1029/93JD02553, 1993.
- 20 Sheng, J. X., Weisenstein, D. K., Luo, B. P., Rozanov, E., Stenke, A., Anet, J., Bingemer, H. and Peter, T.: Global atmospheric sulfur budget under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *J. Geophys. Res.*, 120(1), 256–276, doi:10.1002/2014JD021985, 2015.
- Solomon, S., Daniel, J. S., Neely, R. R., Vernier, J.-P., Dutton, E. G. and Thomason, L. W.: The Persistently Variable “Background” Stratospheric Aerosol Layer and Global Climate Change, *Science* (80-.), 333(6044), 866–870,
25 doi:10.1126/science.1206027, 2011.
- Surono, Jousset, P., Pallister, J., Boichu, M., Buongiorno, M. F., Budisantoso, A., Costa, F., Andreastuti, S., Prata, F., Schneider, D., Clarisse, L., Humaida, H., Sumarti, S., Bignami, C., Griswold, J., Carn, S., Oppenheimer, C. and Lavigne, F.: The 2010 explosive eruption of Java’s Merapi volcano-A “100-year” event, *J. Volcanol. Geotherm. Res.*, 241–242, 121–135, doi:10.1016/j.jvolgeores.2012.06.018, 2012.
- 30 Thomas, H. E., Watson, I. M., Carn, S. A., Prata, A. J. and Realmuto, V. J.: A comparison of AIRS, MODIS and OMI sulphur dioxide retrievals in volcanic clouds, *Geomatics, Nat. Hazards Risk*, 2(3), 217–232, doi:10.1080/19475705.2011.564212, 2011.
- Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J. P., Manney, G., Luo, B., Arfeuille, F. and

- Peter, T.: A global space-based stratospheric aerosol climatology: 1979-2016, *Earth Syst. Sci. Data*, 10(1), 469–492, doi:10.5194/essd-10-469-2018, 2018.
- Trenberth, K. E.: Has there been a hiatus?, *Science* (80-.), 349(6249), 691–692, doi:10.1126/science.aac9225, 2015.
- Vernier, J. P., Pommereau, J. P., Garnier, A., Pelon, J., Larsen, N., Nielsen, J., Christensen, T., Cairo, F., Thomason, L. W.,
5 Leblanc, T. and McDermid, I. S.: Tropical Stratospheric aerosol layer from CALIPSO Lidar observations, *J. Geophys. Res. Atmos.*, 114(24), 1–12, doi:10.1029/2009JD011946, 2009.
- Vernier, J. P., Thomason, L. W., Pommereau, J. P., Bourassa, A., Pelon, J., Garnier, A., Hauchecorne, A., Blanot, L., Trepte, C., Degenstein, D. and Vargas, F.: Major influence of tropical volcanic eruptions on the stratospheric aerosol layer during the last decade, *Geophys. Res. Lett.*, 38(12), 1–8, doi:10.1029/2011GL047563, 2011.
- 10 Wilcox, L. J., Hoskins, B. J. and Shine, K. P.: A global blended tropopause based on ERA data. Part I: Climatology, *Q. J. R. Meteorol. Soc.*, 138(664), 561–575, doi:10.1002/qj.951, 2012.
- Winker, D. M., Hunt, W. H. and McGill, M. J.: Initial performance assessment of CALIOP, *Geophys. Res. Lett.*, 34(19), 1–5, doi:10.1029/2007GL030135, 2007.
- Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H. and Young, S. A.: Overview of the
15 CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Ocean. Technol.*, 26(11), 2310–2323, doi:10.1175/2009JTECHA1281.1, 2009.
- Winker, D. M., Pelon, J., Coakley, J. A., Ackerman, S. A., Charlson, R. J., Colarco, P. R., Flamant, P., Fu, Q., Hoff, R. M., Kittaka, C., Kubar, T. L., Le Treut, H., McCormick, M. P., Mégie, G., Poole, L., Powell, K., Trepte, K., Vaughan, M. A. and Wielicki, B. A.: The Calipso Mission: A Global 3D View of Aerosols and Clouds, *Bull. Am. Meteorol. Soc.*, 91(9), 1211–
20 1229, doi:10.1175/2010BAMS3009.1, 2010.
- Xie, S. P., Kosaka, Y. and Okumura, Y. M.: Distinct energy budgets for anthropogenic and natural changes during global warming hiatus, *Nat. Geosci.*, 9(1), 29–33, doi:10.1038/ngeo2581, 2016.
- Yan, X. H., Boyer, T., Trenberth, K., Karl, T. R., Xie, S. P., Nieves, V., Tung, K. K. and Roemmich, D.: The global warming hiatus: Slowdown or redistribution?, *Earth’s Futur.*, 4(11), 472–482, doi:10.1002/2016EF000417, 2016.
- 25 Young, S. A., Winker, D. M., Noel, V., VaughanOmar, A., Hu, Y. and Kuehn, R. E.: CALIOP algorithm theoretical basis document, Part 5: Extinction Retrieval and Particle Property Algorithms, *Theor. Basis Doc.*, June, PC-SCI-203 Part 5, 2005.
- Zuev, V. V., Burlakov, V. D., Nevzorov, A. V., Pravdin, V. L., Savelieva, E. S. and Gerasimov, V. V.: 30-year lidar observations of the stratospheric aerosol layer state over Tomsk (Western Siberia, Russia), *Atmos. Chem. Phys.*, 17(4), 3067–3081, doi:10.5194/acp-17-3067-2017, 2017.

Tables

Table 1. Volcanic eruptions and wild fires in the 21st century that affect (or had the potential to affect) the aerosol loading of the stratosphere

Volcano		Date	Lat.	Long.	VEI [†]	SO ₂ (Tg)
Ulrawun	U	29 Sep. 2000	5° S	151° E	4	*
Sheveluch	S	22 May 2001	57° N	161° E	4	*
Ruang	R	25 Sep. 2002	2° N	125° E	4	0.03 ^a
Reventador	R	3 Nov. 2002	0° S	78° W	4	0.07 ^a
Anatahan	A	10 May 2003	16° N	146° E	3	0.03 ^a
Manam	M	27 Jan. 2005	4° S	145° E	4	0.09 ^a
Sierra Negra	S	22 Oct. 2005	1° S	91° W	3	*
Soufrière Hills	S	20 May 2006	17° N	62° W	3	0.2 ^b
Rabaul	Rb	7 Oct. 2006	4° S	152° E	4	0.2 ^a
Jebel at Tair	J	30 Sep. 2007	16° N	42° E	3	0.08 ^c
Great Divides Fire	Gd	1 Dec. 2006	37° S	144° E	-	-
Chaitén	C	2 May 2008	43° S	73° W	4	0.01 ^d
Okmok	O	12 Jul. 2008	53° N	168° W	4	0.1 ^c
Kasatochi	Ka	7 Aug. 2008	52° N	176° W	4	1.7 ^c
Fire in Victoria	Vi	7 Feb. 2009	37° S	145° E	-	-
Redoubt	R	23 Mar. 2009	60° N	153° W	3	0.01 ^e
Sarychev	Sa	12 Jun. 2009	48° N	153° E	4	1.2 ^f
Eyjafjallajökull	E	14 Apr. 2010	64° N	20° W	4	*
Merapi	Me	5 Nov. 2010	8° S	110° E	4	0.4 ^g
Grimsvötn	Gr	21 May 2011	64° N	17° W	4	0.4 ^h
Puyehue-Cordón Caulle	Pu	6 Jun. 2011	41° S	72° W	5	0.3 ^h
Nabro	Na	12 Jun. 2011	13° N	42° E	4	1.5 ^h
Kelut	Ke	13 Feb. 2014	8° S	112° E	4	0.2 ⁱ
Calbuco	Ca	23 Apr. 2015	41° S	73° W	4	0.3 ^j

[†] Volcanic Explosivity Index (from Global Volcanism Program (<http://www.volcano.si.edu/>))

5 * Not available

^a Prata and Bernardo (2007)

^b Carn and Prata (2010)

^c Thomas et al. (2011)

^d Carn et al. (2009)

10 ^e Lopez et al. (2013)

^f Haywood et al. (2010)

^g Surono et al. (2012)

^h Clarisse et al. (2012)

ⁱ Li et al. (2017)

^j Pardini et al.(2017)

|

Table 2. Comparison of the annual mean AODs from our study and other studies, for the years 2007-2013 and various latitude intervals.

Instruments	Lat intervals	Altitude ^d	AOD in present study (other groups deviations from our AODs)							
			2007	2008	2009	2010	2011	2012	2013	All years
CALIOP/ OSIRIS ^a	35-45°N	TP-35 km ^c	0.0100 (-4.8%)	0.0128 (-7.2%)	0.0181 (-7.8%)					0.0136 (-4.4%)
	5°S-5°N	TP-35 km ^c	0.0052 (15%)	0.0044 (20%)	0.0052 (18%)					0.0049 (17%)
	35-45°S	TP-35 km ^c	0.0093 (29%)	0.0082 (38%)	0.0088 (32%)					0.0088 (32%)
OSIRIS ^b	50°S-50°N	1 km above 380K-35 km	0.0046 (32%)	0.0041 (19%)	0.0048 (17%)	0.0039 (21%)	0.0050 (25%)	0.0039 (27%)	0.0031 (22%)	0.0042 (23%)
CALIOP/ ^b GOMOS	50°S-50°N	15-35 km	0.0051 (30%)	0.0047 (23%)	0.0056 (0.3%)	0.0045 (10%)	0.0060 (0.1%)	0.0043 (12%)	0.0036 (12%)	0.0048 (12%)
OSIRIS/Sato ^b	50°S-50°N	15-35 km	0.0051 (-7.7%)	0.0047 (-19%)	0.0056 (-16%)	0.0045 (-16%)	0.0060 (-12%)	0.0043 (-8.2%)		0.0048 (-13%)
GOMOS/ OSIRIS/ CALIOP ^e	40-50°N	17-30 km	0.0037 (0.0%)	0.0041 (-6.8%)	0.0045 (-4.8%)	0.0035 (-8.4%)	0.0040 (-6.6%)	0.0034 (1.4%)	0.0029 (-10%)	0.0037 (-5.0%)

^aThomason et al. (2018)

^bRieger et al. (2015), higher lidar ratio than 50 sr for CALIOP

^cTP refers to the tropopause in the comparison with Thomason et al. (2018)


^dThomason et al. (2018) used a climatic tropopause, where we used the the dynamic tropopause at 1.5 PVU.

^eKhaykin et al. (2017), mean of satellite retrievals, lidar ratio of 50 sr for CALIOP


Table 32. Mean Global mean stratospheric AOD and radiative forcing for years 2006-2015, the year 2013 and the first year after the volcanic eruptions in Figure 9.

Period	AOD	RF (Wm^{-2})
Years 2006 - 2015	0.0082	-0.21
The year 2013	0.0059	-0.15
Sarychev +1year	0.0102	-0.26
Nabro +1year	0.0099	-0.25
Kelut +1year	0.0073	-0.18

Figures




5 | **Figure 1. Illustration of the removal of cloud signals from the CALIOP data. (a) The original data, (b) and (c) removal of signals from ice clouds and polar stratospheric clouds, respectively. Data averaged over August 2009 were used in this example. The strong aerosol signal in the Northern Hemisphere comes from aerosol connected to the Sarychev eruption. The white lines mark the average altitudes of the tropopause and the 380 K isentrope.**



10 | **Figure 2. Examples illustrating the effects from corrections made to compensate for the laser attenuation caused by aerosol particles. The scattering ratios (SR) were averaged over the latitude intervals a) 40-50°N and b) 40-50°S for the UT (black; 250 -**

1000 m below the tropopause) and the PV-layer of 6-7.5 PVU (magenta). [Volcanic eruptions are indicated by their abbreviations.](#) Both layers are on average approximately 750 m thick). Corrected (Uncorrected) data are marked as full (dashed) lines.



5 Figure 3. Comparisons of the particle extinction parameter from the level 2 data v4-10 (a and c), and the aerosol scattering (b and d) taken after the Calbuco eruption. The upper figures were taken three weeks after (on May 13, 2015), and the lower almost four months after (on September 17, 2015) the eruption. Data in b and d were based on the attenuated backscattering at 532 nm in the level 1 v4-10 data, and modeling of the molecular scattering. The white lines mark the [thermalstatic](#) tropopause.




Figure 4. Monthly mean scattering ratio as a function of latitude and altitude (7 - 35 km) for years 2006–2015 and in the months January, April, July and October for years 2006 - 2015.





Figure 5. Aerosol scattering against time and altitude in six latitude bands each covering 16% of the earth's surface area. [The white lines mark the mean altitudes of the tropopause and the 380 K isentrope.](#)



5

Figure 6. Temporal trends of the scattering ratio in altitude layers in the lower part of the LMS (PV intervals of 0.5 PVU), for the latitude bands 40-50°N (a) and 40-50°S (b). Vertical soliddashed lines marks extratropical volcanic eruptions, and dashed indicates extratropical eruptions that impacted the respective Hemisphere.





Figure 7. The latitudinal distribution over ten years for the latitude weighted stratospheric AOD: a) above the 470 K isentrope, b) between the 380 and 470 K isentropes, c) the LMS, and d) that of the entire stratosphere. The color bar for a-c spans a range of 1/3 of that in d. Abbreviations and Triangles (circles) marks time and latitude of relevant volcanic eruptions (forest fires).



5 | **Figure 8. Temporal trends of the stratospheric AOD and radiative forcing for a) the entire globe, b) the Northern Hemisphere, and c) the Southern Hemisphere. The lines mark the AOD of the LMS (magenta), altitudes between the 380 and 470 K isentropes (cyan), altitudes above the 470 K isentrope (black), and the entire stratosphere (thick red). Significant volcanic eruptions are indicated by their abbreviations. Note the reversed scale for the radiative forcing on the right y-axis. Data were extrapolated to the Polar Regions and latitude weighted before averaging.**




Figure 9. The latitudinal distribution of the latitude weighted AOD. a) The AOD of the different stratospheric layers and of the total stratospheric AOD (thick red) for the entire period (2006-2015). b-d) shows the respective layers and e) the AOD of the entire stratospheric column, for the time periods noted in the legend. Lines represent AODs in the altitude intervals 470 K to 35 km (black), 380 – 470 K (cyan), the LMS (magenta), and the entire stratosphere (thick red). 2013 (grey filling) is considered as background conditions, and the line styles mark the different periods as noted in the legend.

5