

Disentangling the rates of carbonyl sulphide (COS) production and consumption and their dependency with soil properties across biomes and land use types

5 Aurore Kaisermann¹, Jérôme Ogée¹, Joana Sauze¹, Steve Wohl¹, Sam P. Jones¹, Ana Gutierrez¹, Lisa Wingate¹

¹ INRA/Bordeaux Science Agro, UMR 1391 ISPA, Villenave d'Ornon, 33140, France

Correspondence to: Aurore Kaisermann (aurore.kaisermann@inra.fr)

Abstract. Soils both emit and consume the trace gas carbonyl sulphide (COS) leading to a soil-air COS exchange rate that is the net result of two opposing fluxes. Partitioning these two gross fluxes and understanding their drivers are necessary to estimate the contribution of soils to the current and future atmospheric budget of COS.

Previous efforts to disentangle the gross COS fluxes from soils have used flux measurements on air-dried soils as a proxy for the COS emission rates of moist soils. However, this method implicitly assumes that COS uptake becomes negligible and COS emission remains steady while soils are drying. We tested this assumption by estimating simultaneously the soil COS sources and sinks and their temperature sensitivity (Q_{10}) from soil-air COS flux measurements on fresh soils at different COS concentrations and two soil temperatures. Measurements were performed on 27 European soils from different biomes and land use types in order to obtain a large range of physical-chemical properties and identify the drivers of COS consumption and production rates.

We found that COS production rates from moist and air-dried soils were not significantly different for a given soil and that the COS production rates had Q_{10} values (3.96 ± 3.94) that were larger and more variable than the Q_{10} for COS consumption (1.17 ± 0.27). COS production generally contributed less to the net flux at lower temperatures but this contribution of COS production increased rapidly at higher temperatures, lower soil moisture contents and lower COS concentrations. Consequently, measurements at higher COS concentrations (*viz.* 1000 ppt) always increased the robustness of COS consumption estimates. Across the range of biomes and land use types, COS production rates co-varied with total soil nitrogen concentrations ($r = 0.52$, $P < 0.05$) and mean annual precipitation ($r = 0.53$, $P < 0.05$) whilst the gross COS uptake rate and the first-order COS hydrolysis rate constant co-varied significantly with the microbial biomass N content of the soils ($r = -0.74$ and 0.64 , $P < 0.05$, respectively). Collectively our findings suggest a strong interaction between soil nitrogen and water cycling on COS production and uptake, providing new insights on how to upscale the contribution of soils to the global atmospheric COS budget.

30

1 Introduction

Carbonyl sulphide (COS) is a powerful greenhouse gas whose atmospheric concentration has varied considerably during the Earth's history (Ueno et al., 2009). In the present day stratosphere, COS photolysis contributes to the formation of aerosol particles that also cool the planet, consequently offsetting the global warming potential of COS (Brühl et al., 2012). Currently, the concentration of COS in the troposphere is around 500 ppt (or pmol mol^{-1}) almost 1 million times less than current atmospheric CO_2 concentrations. However, the relative seasonal amplitude of COS is about 5 times larger than that of CO_2 and has led to COS being proposed as a powerful tracer of gross primary production (GPP) over land (e.g. Montzka et al., 2007; Sandoval-Soto et al., 2005; Whelan et al., 2017). This is because the removal of COS from the atmosphere during summer months in the Northern hemisphere is driven by the activity of plants over land that remove COS through an enzymatic reaction with carbonic anhydrase (CA) (Sandoval-Soto et al., 2005; Stimler et al., 2012), an ubiquitous enzyme particularly abundant in leaf mesophyll cells (Fabre et al., 2007).

Soils are also teeming with a diverse range of organisms such as bacteria, fungi and algae that also contain CAs (Elleuche and Pöggeler, 2010; Moroney et al., 2001; Smith and Ferry, 2000) and thus have the potential to remove COS from the atmosphere. For this reason, oxic soils are generally considered net sinks for atmospheric COS (Van Diest and Kesselmeier, 2008; Kesselmeier et al., 1999; Li et al., 2005; Whelan et al., 2016b) albeit with a much weaker sink strength than vegetation (Berry et al., 2013; Campbell et al., 2017; Kettle et al., 2002; Launois et al., 2015). Some soils have also been found to be strong producers of COS, notably anoxic soils (Fried et al., 1993; Hines and Morrison, 1992; Whelan et al., 2013) in addition to some oxic soils with and without plant litter (Bunk et al., 2017; Kitz et al., 2017; Maseyk et al., 2014; Melillo et al., 1993; Sun et al., 2016; Whelan et al., 2016a; Whelan and Rhew, 2015).

Despite the importance of soil COS fluxes for improving the global COS mass budget, the mechanisms underlying COS production and consumption by soils remain unclear. This is partly because it is difficult to disentangle the functional response of these two opposing fluxes in isolation and characterise how changes in the environment or soil properties impact the net COS flux. Recently physical and enzymatic models describing the consumption of COS by soils have been advanced (Ogée et al., 2016; Sun et al., 2015). However, equivalent mechanistic understanding of COS production is still lacking.

One approach for estimating COS emission rates from soils is to measure the net COS flux rate of air-dried soil samples (Whelan and Rhew, 2016; Whelan et al., 2016b). This assumes that the COS consumption by dry soils is negligible as hydrolysis by CA requires the presence of water to proceed (Van Diest and Kesselmeier, 2008; Ogée et al., 2016; Sun et al., 2015; Whelan and Rhew, 2016). Thus with a further assumption that COS emission rates do not vary while soils are drying, COS emission rates can be retrieved from the net COS flux measured on fresh soils (Whelan et al., 2016b). However, it is still not clear whether the COS production by soils is related to biological activity and potentially varying with soil moisture. If this was the case, the described method for estimating COS production rates on dry soils could create strong biases in the partitioning of the net COS flux under wet or moist field conditions.

An alternative approach, presented by Conrad (1994), facilitates the simultaneous estimation of COS production and consumption on fresh soils, thereby providing a test of whether COS production changes with soil water content or not. Because this alternative approach requires the measurement of net COS fluxes at different atmospheric COS concentrations, it cannot be easily implemented in the field without large artefacts (Castro and Galloway, 1991; Mello and Hines, 1994), but it is well adapted to measurements on soil microcosms (Lehmann and Conrad, 1996; Conrad & Meuser, 2000; Whelan et al., 2016b). So far very few studies have implemented this approach, thus the partitioning of COS fluxes at ambient concentrations still remains poorly explored (Whelan et al., 2017).

In this study, we used the approach of Conrad (1994) to estimate COS production and consumption rates simultaneously from moist soils near ambient COS concentrations. We combined this approach with a recently developed theoretical framework (Ogée et al., 2016) to retrieve COS production and first-order consumption rates on a range of soils from different biomes and land use types located in Europe and Israel. We then evaluated the effects of varying COS concentration, soil moisture and temperature on the COS production and consumption rates and assessed the potential role of soil properties such as pH, texture, soil carbon and nitrogen, microbial biomass, etc., as drivers of COS gross fluxes.

2 Materials and methods

2.1 Theory

Assuming that the soil-air COS exchange rate in soil microcosms is governed by only three processes, namely diffusion through the soil column, production and uptake *via* hydrolysis, Ogée *et al.* (2016) proposed a steady-state, analytical model of the COS efflux at the soil surface (F , $\text{pmol m}^{-2} \text{s}^{-1}$) as follows:

$$F = \left(-\sqrt{kB\theta D} C_a + \sqrt{\frac{D}{kB\theta}} \rho_b P \right) \tanh \left(z_{\max} \sqrt{\frac{kB\theta}{D}} \right), \quad (1)$$

where D ($\text{m}^2 \text{s}^{-1}$) is the COS diffusivity through the soil matrix, k (s^{-1}) is the first-order COS hydrolysis rate constant and P ($\text{pmol kg}^{-1} \text{s}^{-1}$) is the COS production rate. Other symbols are B ($\text{m}^3 \text{m}^{-3}$), the COS solubility in soil water, θ ($\text{m}^3 \text{m}^{-3}$), the soil volumetric water content, C_a (mol m^{-3}), the *molar* concentration of COS in air at the soil surface, ρ_b (kg m^{-3}), the soil bulk density and z_{\max} (m), the maximum soil depth. This equation assumes a finite soil depth (z_{\max}) and uniform soil properties (θ , ρ_b ...) and is therefore only suited for soil microcosm studies (Ogée et al., 2016).

Noting that $\tanh(ax)/x = a$ if $x \rightarrow 0$, when soil moisture tends to zero ($\theta \rightarrow 0$), Eq. 1 simplifies to $F_{\text{dry}} = \rho_b P_{\text{dry}z_{\max}}$ where F_{dry} and P_{dry} represent the net COS flux F and the COS production rate P of an air-dry soil, respectively. Thus, assuming that P does not vary with soil moisture ($P = P_{\text{dry}}$), the COS production rate can be estimated from measurements of the soil-to-air COS flux performed on air-dry soil samples. This method, hereafter called Method 1, has been proposed recently for partitioning soil COS sources and sinks (Whelan et al., 2016b). By knowing the value of P , the COS hydrolysis rate constant k can then be estimated using Eq. 1, from measurements of the soil-to-air COS flux performed on moist soils. In practice a

numerical iterative method must be implemented to find the value of k that minimises the discrepancy between the observed and modelled fluxes (Ogée et al., 2016; Sauze et al., 2017a).

Note that if the steady-state net soil COS flux F is measured at different COS concentrations C_a , and provided that P is known, it is possible to derive a different k value for each (F, C_a) pair of measurements. In theory each pair of measurements should provide the same hydrolysis rate constant, unless COS hydrolysis does not follow first-order kinetics and k is not a true rate constant and varies with C_a . In fact this might well be the case as k was originally hypothesised to be a catalysed rate constant (Kesselmeier et al., 1999; Lehmann and Conrad, 1996) that follows Michaelis-Menten kinetics (Ogée et al., 2016; Protoschill-Krebs and Kesselmeier, 1992). In this case the hydrolysis rate would saturate to a maximal rate V_{\max} at very high COS concentrations and would respond linearly to C_a with a slope $k = V_{\max}/K_m$ only at low concentrations, i.e., when $C_a \ll K_m$ where K_m (mol m⁻³) is the so-called Michaelis-Menten coefficient and corresponds to the COS concentration at which k equals $0.5 V_{\max} / k_m$.

Carbonic anhydrases (CA) are a family of enzymes ubiquitous in soil micro-organisms (Wingate et al., 2009) that are known to catalyse COS hydrolysis (Protoschill-Krebs et al., 1992; 1996). However, studies reporting their kinetic parameters are very scarce but tend to show relatively high values of K_m , around 40-60 μM at 20°C (Protoschill-Krebs et al., 1992; Haritos et al., 2005; Ogawa et al., 2013; Ogée et al., 2016), i.e., more than 2 million times the atmospheric COS concentration (500 ppt or 20 pM). With such high K_m values, and assuming that COS consumption by soils is only caused by CA-driven hydrolysis, the rate constant k should be well approximated by V_{\max} / k_m and thus F should respond linearly to C_a (see Eq. 1), without any sign of saturation. A near-linear response of F to C_a has indeed been demonstrated on all soils tested so far, even at COS concentrations 100 times higher than ambient levels (Lehmann & Conrad, 1996). Based on this observation, Eq. 1 can be re-written:

$$F = F_0 - V_{d0}C_a, \quad (2)$$

where F_0 is the gross flux of COS production, i.e., the flux F when $C_a = 0$, and V_{d0} (m s⁻¹) is the COS deposition velocity onto the soil surface that would occur in the absence of COS production (i.e. if $P = 0$). The gross flux of COS uptake is calculated by subtracting F_0 from the net COS flux F .

From Eq. 2 we can see that, by performing steady-state COS flux measurements at different COS concentrations, it is possible to estimate simultaneously V_{d0} and F_0 from the slope and the intercept of a linear regression between the two variables, leading to the determination of P and k :

$$F_0 = \sqrt{\frac{D}{kB\theta}} \rho_b P \tanh\left(z_{\max} \sqrt{\frac{kB\theta}{D}}\right), \quad (3a)$$

$$V_{d0} = \sqrt{kB\theta D} \tanh\left(z_{\max} \sqrt{\frac{kB\theta}{D}}\right), \quad (3b)$$

In practice, this method, hereafter called Method 2, is performed in three steps. First a 3-point linear regression is performed between F and C_a data to estimate F_0 and V_{d0} for each microcosm using the `lm` function in R. Then the value of k that satisfies Eq. 3b is obtained using an iterative numerical method (`fzero` function in the `Pracma` Package; Brochers, 2017). Finally this k value is introduced into Eq. 3a to estimate P from F_0 . Values for B are estimated from soil temperature using Wilhelm et al. (1977) and the COS effective diffusivity D is estimated using the empirical formulation of Moldrup et al. (2003) for repacked soils (see also Ogée et al., 2016).

2.2 Soil sampling and preparation

Soils from 27 locations were collected along a latitudinal gradient in Europe and Israel during the summer of 2016. These locations were selected to cover a range of biomes and land use as well as soil physico-chemical properties (see Tables 1 and S1). The first ten centimetres of soil were collected at three locations at each site and sealed in plastic bags and sent to INRA Bordeaux after collection with no special requirements imposed for the transportation of the soil samples. Upon reception, the different soils were sieved using a 4mm mesh, homogenised and stored at 4°C. The sieving was performed to ensure a representative sample of only soil, to avoid introducing any additional and (uncontrolled) plant litter effects that could potentially introduce variability between sample replicates and complicate the interpretation of the net and gross COS fluxes. We justify this experimental choice as our overall goal was to derive and validate a model of soil COS fluxes regulated by commonly quantified soil physical, chemical and biological characteristics. Our main hypothesis is that the drivers identified in our study would still be applicable in undisturbed soils. Sieved soils were then separated into two batches. One batch was air-dried by spreading soil in a tray and regularly mixing every 2-3 days for 1-2 weeks before being measured to estimate the air-dried COS production rate (P_{dry}) hereafter referred to as “dry”. The other batch was stored at 4°C for several weeks until it was measured to estimate the COS production and consumption rates of fresh soils (P and k) hereafter referred to as “moist”. For both methods, measurements were performed in triplicates, using soil microcosms consisting of custom-made glass jars (0.825 dm³ volume, 8.85 cm internal diameter). During a 2-week incubation period in a climate-controlled chamber, the microcosms remained open to the air circulating in the climate chamber (MD1400, Snijders, Tillburg, NL) that had CO₂ and COS concentrations controlled at around 400 ppm and 500 ppt, respectively, i.e., close to ambient levels. The same CO₂ concentration was also used to purge the air inlet during the gas exchange measurements, so that the microcosms were never exposed to CO₂ levels much higher than those of ambient air or those used during the gas-exchange measurements (see below). For Method 1, 350-400 g of air-dried soil was placed in each microcosm and acclimated for 2-3 days in the dark prior to gas exchange measurements. For Method 2, 200-300 g of equivalent dry soil were used and maintained at 30% of maximum water holding capacity (WHC) for 2 weeks in the dark until gas exchange measurements were performed. Prior to this incubation period, the WHC of each soil was estimated using the method of Haney and Haney (2010) and when fresh soils were too wet (> 30% WHC), they were air-dried until they reached 10-20% WHC, then re-humidified to 30% WHC at the start of the 2-week incubation period. Care was taken to avoid measuring the so-called “Birch effect” (Jarvis et al., 2007) during gas exchange measurements. Because the largest water addition was performed just before the incubation period (especially on

soils that were shipped from very dry places) all gas exchange measurements were delayed for 2 weeks to ensure fluxes had stabilised during the incubation period and that they were outside the decay curve of the Birch effect. Throughout these 2 weeks, microcosms were kept unsealed in the dark and the moisture contents were monitored gravimetrically every two days whereupon, extra but small, water additions were made (about 2-5 g of distilled water for an average water content of 58.7 ± 16 g) and no later than 24h before the start of the gas exchange measurements.

2.3 Gas-exchange measurements

Glass soil microcosms were equipped with customised screw-tight glass lids cut to the exact size of the microcosm area and finely ground to provide a glass-on-glass seal that was held in place by screwing the threaded metal lid onto the jar over the glass panel. The lids were equipped with two stainless steel Swagelok® (Swagelok, Solon, OH, USA) fittings to connect to the 1/8" Teflon inlet and outlet lines of the measurement system. Sealing was ensured using PTFE washers that were previously tested and shown not to emit COS. A stainless steel temperature probe (3-wire PT100, 15 cm length, 3 mm diameter, reference RS 362 9935) continuously recorded the average soil temperature in each microcosm. Dry synthetic air was then adjusted to the desired CO₂ (399 ± 6 ppm) and COS mixing ratios and supplied to the microcosms using the same system as described in Gimeno et al. (2017). All pots were weighed before and after gas exchange measurements to calculate water loss. On average, the water loss was 0.2 ± 0.005 g per hour or 4 ± 1% of the initial water amount (i.e. a reduction of the volumetric water content of less than 0.01 cm³ cm⁻³). Additional gas exchange measurements were also performed after the end of the sequence and we could verify that the COS and CO₂ fluxes of a given microcosm (at a given temperature and COS concentration) were not significantly different between the first and second sequences (16h apart; see below). This was a clear indication that the small water loss during the duration of the gas exchange measurements did not impact significantly the COS and CO₂ fluxes. The inlet and outlet airstreams of each microcosm were analysed sequentially using a mid-infrared quantum cascade laser spectrometer (QCLS, Aerodyne Research Inc Billerica, MA, USA), coupled upstream to a Nafion dryer (MD-070-24-S-2, Perma Pure LLC, Lakewood, NJ, USA) to remove matrix effects caused by water vapour (Kooijmans et al., 2016). To account for instrument drift, an auto-background was implemented regularly (typically every 38 minutes) for 120 s using a dry N₂ bottle. A 2-point calibration scheme was also implemented using the same dry N₂ bottle (measured every 14 minutes) and an Aculife-treated cylinder (Air Liquide USA, Houston, TX, USA) filled with compressed air and 524.8 pmol(COS) mol⁻¹ calibrated to the NOAA-Scripps Institution of Oceanography provisional scale. This second cylinder was measured every 14 to 56 minutes depending on the sequence used.

Using a custom-made multiplexed system (Sauze et al., 2017a), six jars with six different soils and one empty jar (blank) were measured sequentially over *ca.* 18h to investigate simultaneously 6 different soils under identical conditions (Supplement Fig. S2). Over this period the measuring sequence consisted of 8 steps that measured the COS fluxes from all the microcosms at 2 different temperatures (18°C and 23°C) and 3 different COS concentration levels (around 100, 500 and 1000 ppt), with an acclimation time of *ca.* 2 hours following a change in temperature and 40 minutes following a change in COS concentration (see Supplement Fig. S2). While only 14 mins were usually required to stabilise the COS mixing ratio on the chamber lines

after a step change in the COS mixing ratio of the inlet line, two hours seemed the minimum time required to stabilize the soil temperature to a new temperature.

For each temperature and COS concentration level, three inlet/outlet pairs were measured on each microcosm, every 44 minutes to partially take into account the possible variability caused by the small water loss. Each line was measured for 120 s and only the last 15 s were retained to compute the mean COS concentration, accounting for the residence time of air in the tubing and gas analyser. The median standard deviation during these last 15s was 12.4 ppt for COS and 0.09 ppm of CO₂. From each inlet/outlet pair the net soil-to-air COS flux was computed as follows:

$$F = \frac{\phi}{S} (c_a - c_{in}), \quad (4)$$

10

where F is the net COS flux ($\text{pmol m}^{-2} \text{s}^{-1}$), ϕ is the flow rate of dry air through the chamber (mol s^{-1}), S (0.00615 m^2) is the soil surface area, c_{in} (pmol mol^{-1}) is the COS mixing ratio on the inlet and c_a (pmol mol^{-1}) is the COS mixing ratio on the outlet. The air flow rate ϕ was set at 0.250 nlpm, i.e., $186 \mu\text{mol s}^{-1}$. The COS flux for the blank chamber was never significantly different from zero.

15 The molar COS concentration (C_a) was estimated from the molar ratio (c_a) and soil temperature measurements using the ideal gas law and an air pressure of 106000 Pa. The slight over pressure in the glass jars (of about 5 kPa) had been estimated previously during a preliminary experiment using a pressure transducer (BME280; Bosch GmbH, Gerlingen, Germany).

2.4 Estimation of soil COS production and hydrolysis rates

The COS production rate was first estimated on air-dried soils at 18°C and under atmospheric concentration (*ca.* 500 ppt) levels of COS. This “dry” production rate (P_{dry} , $\text{pmol kg}^{-1} \text{s}^{-1}$) was deduced from the COS flux (Eq. 4) according to:

$$P_{dry} = \frac{SF_{dry}}{M_{dry}}, \quad (5)$$

where M_{dry} (kg) is the mass of dry soil in the microcosm. The COS production and hydrolysis rates on fresh soils (P_{moist} and k_{moist} , respectively) were estimated using COS flux measurements performed at the three COS concentrations and Eqs. 2 and 3 described above. The linear relationship between F and C_a observed over a wide range of COS concentrations was confirmed using our set-up over the range of COS mixing ratios used in our experiments, i.e., 0-1200 ppt (see Supplement Fig. S3). These results justified the use of only three COS levels (referred to as “low”, “med” and “high” hereafter) to perform the linear regression and calculate P_{moist} and k_{moist} in subsequent analyses. The COS mixing ratio in the inlet airstream of each microcosm was thus set to 1111 ± 29 ppt (“high”), 557 ± 10 ppt (“med”) or 124 ± 8 ppt (“low”), while the CO₂ mixing ratio was always maintained around 399 ± 6 ppm.

In order to evaluate whether the method used to estimate the COS production rate influenced the calculation of the COS hydrolysis rate of moist soils, we also used P_{dry} to re-calculate the hydrolysis rate of moist soils as in previous studies. To do so, we inserted P_{dry} into Eq. 1 and solved for the hydrolysis rate that satisfied the equation for a given level of COS concentration (referred to as $k_{recal,low}$, $k_{recal,med}$ and $k_{recal,high}$ hereafter).

- 5 The COS production and hydrolysis rates for the wet soils (P_{moist} and k_{moist}) were measured at two temperatures (18°C and 23°C) to estimate their temperature sensitivity (Q_{10}) in this temperature range:

$$Q_{10}(k) = \left(\frac{k_{wet(23^{\circ}C)}}{k_{wet(18^{\circ}C)}} \right)^2, \quad (6a)$$

10
$$Q_{10}(P) = \left(\frac{P_{wet(23^{\circ}C)}}{P_{wet(18^{\circ}C)}} \right)^2, \quad (6b)$$

2.5 Soil physico-chemical properties

At the end of each gas exchange measurement, the soils were analysed for a range of physico-chemical properties. Soil texture and total C, N and CaCO₃ contents were measured using standard procedures at the INRA soil analyses platform (<http://www.lille.inra.fr/las>). Soil pH and redox potential were measured using a 1:5 soil-water ratio using a Fisher Scientific
 15 Accumet™ series XL250 dual Channel pH/mV with pH epoxy gel electrode, gelled redox platinum electrode and temperature probe. pH and redox electrodes were calibrated with buffer solutions pH 4, 7 and 10 and with ORP solution 470mV from Hanna Instruments, respectively. Bulk density was estimated from the weight and volume of each soil microcosm. Soil water content was estimated gravimetrically as the weight difference between moist and oven-dried soil extracts. The concentration of phosphate ions was measured as in Van Veldhoven and Mannaerts, (1987). Microbial biomass for carbon (C) and nitrogen
 20 (N) were estimated as the difference of dissolved C and N contents between fumigated (24h of chloroform fumigation) and non-fumigated soil extracts consisting of 10g of soil mixed with 40ml of 0.5 M of K₂SO₄ and shaken for 30min.

2.6 Statistical analyses

All data processing and graphs were made using the R software (Version 3.3.3, R core Team, 2015) packages dplyr, lubridate, data.table and ggplot2. These were used to test the effects of biome and land use on the gross COS production (P_{moist}) and first-
 25 order hydrolysis rate (k_{moist}) constants, as well as assessing the effects of atmospheric COS concentration on $k_{recal,med}$ and k_{moist} . The temperature response of P_{moist} and k_{moist} was compared using ANOVA and Tukey's HSD tests. To investigate the correlation between soil properties and all COS fluxes (gross COS production and uptake, hydrolysis rate at 18°C) spearman coefficient correlations were calculated and tests for significance were performed with the corrplot package (Wei and Simko, 2017).

3 Results

All moist soils were net COS sinks at 18°C, with net fluxes ranging in magnitude from -7.66 to -0.78 pmol m⁻² s⁻¹ (Fig. 1). In comparison, the blank was not significantly different from zero with mean net COS flux values of -0.11 ± 0.24 pmol m⁻² s⁻¹. Using the theoretical framework presented above we partitioned the net COS fluxes measured on moist soils to assess COS production and uptake rates and to compare moist (P_{moist}) with dry (P_{dry}) soil COS production rates. As illustrated in Fig. 2a the COS production rates measured on moist soils (P_{moist}) were not significantly different from those measured on dry soils (P_{dry}). Overall the relationship between P_{moist} and P_{dry} was highly significant ($P < 0.001$) and followed a linear regression slope of 0.98 with an intercept of 0.02 pmol kg⁻¹ s⁻¹ (Fig. 2a). Dispersion of data around the linear regression ($r^2 = 0.59$) indicated that some soils were occasionally underestimated by one method compared to the other. Upon investigation we found that some soils having much higher P_{moist} than P_{dry} may not have air-dried soils completely dry. This hypothesis is partially supported by the persistence of a small net CO₂ flux (Table S1) for these soils, indicating that some uptake of COS could still be possible contributing to some of the dispersion in the data.

Our study also indicated that the rates of COS production from moist soils measured at 18°C (P_{moist}) were significantly higher in temperate regions compared to those measured in boreal and Mediterranean regions (ANOVA $P = 0.0009$, Tukey's HSD tests: Temperate-Mediterranean $P=0.0009$, Temperate-Boreal $P=0.03$, Mediterranean-Boreal $P=0.4$; Figs. 3 and S4). The highest COS production rates were measured on soils coming from temperate grassland sites (Figs. 3 and S4). Further analysis indicated that the eight temperate soils exhibiting the highest COS production rates also contained high C and N contents (Fig. 2a). The total C and N contents of the different soils were positively correlated with high microbial C and N biomass as well as redox potential, whilst negatively correlated with bulk density (Figs. 4 and S5 and Table S1). In addition, the total N content was further correlated with site mean annual precipitation (MAP) (Fig. 4). No significant effect was detected between P_{moist} and latitude, longitude or land use cover. However, COS production rates were significantly and positively correlated with soil N content ($r = 0.52$), MAP ($r=0.53$) and soil redox potential ($r = 0.50$) and negatively correlated with pH ($r = -0.41$) (Fig. 4). However, a principle component analysis indicated that the production rate was most closely associated with soil N content (Fig. S5).

The partitioned gross COS uptake rates ($V_{40}C_a$) measured at 30% WHC were always much larger in absolute values (between -7.66 and -1.34 pmol m² s⁻¹) than the COS production rates P_{moist} (less than 0.81 pmol m² s⁻¹) and thus dominated the net COS flux F (Fig. 3). The first-order COS hydrolysis rate constant k_{moist} was estimated to vary between 0.05 to 0.47 s⁻¹ and the relationship between the two estimates, k_{moist} and $k_{\text{reca, mid}}$, were strongly ($r^2 = 0.96$) and linearly related, exhibiting a slope and intercept of 0.94 and 0.02 s⁻¹, respectively (Fig. 2b). Thus demonstrating that the use of P_{moist} or P_{dry} had little influence on the retrieval of the first-order COS hydrolysis rate constants (Fig. 2b). However, the relative difference between the COS hydrolysis rates k_{moist} and those re-calculated using P_{dry} and Eq. 1 were significantly different when measured at different COS

concentrations (Fig. 5; $P = 0.002$). Indeed, the COS hydrolysis rate that satisfied the equation for low COS concentrations, $k_{\text{reca},\text{low}}$ (estimated using F and C_a data from the low COS concentration measurements), was significantly ($P = 0.0011$) lower than that estimated at high COS concentration, $k_{\text{reca},\text{high}}$ (estimated using F and C_a data from the high COS concentration measurements), while $k_{\text{reca},\text{mid}}$ (estimated using the medium COS concentration measurements) was intermediate and not significantly different from either $k_{\text{reca},\text{high}}$ ($P = 0.52$) or $k_{\text{reca},\text{low}}$ ($P = 0.056$). Furthermore, $k_{\text{reca},\text{low}}$ values exhibited a larger spread in the deviation from k_{moist} than both $k_{\text{reca},\text{mid}}$ and $k_{\text{reca},\text{high}}$ (Fig. 5). This demonstrated that the estimation of the COS hydrolysis rate using Eq. 1 becomes more sensitive to the value of P prescribed when flux measurements are performed at low COS concentrations as there is a decrease in the fraction of COS uptake with respect to COS production. In contrast to the results found for the COS production rates, the first-order COS hydrolysis rate constants k_{moist} were not related to land use or biome. On the other hand, k_{moist} values were positively and significantly correlated with microbial N ($r = 0.64$) and C ($r = 0.45$) biomass contents (Fig. 4).

The temperature sensitivity (Q_{10}) of P_{moist} had a mean and standard deviation of 4.36 ± 4.45 . This was significantly higher ($P < 0.0001$) than the Q_{10} of the hydrolysis rate that had a mean and standard deviation of 1.26 ± 0.29 (Fig. 6). The variability in Q_{10} values across the 27 soils was also much larger for COS production rates than for COS hydrolysis rates. The temperature sensitivity of P_{moist} did not correlate with any of the measured soil properties (Fig. 4). However, the Q_{10} values of the COS hydrolysis rate constants were significantly and negatively correlated with soil total C content ($r = -0.46$) and positively correlated to bulk density ($r = 0.32$) (Fig. 4).

4 Discussion

4.1 COS production rates measured on dry soils are a reasonable proxy for those occurring in moist soils

Net COS fluxes measured from oxic soils commonly exhibit a unimodal response to water-filled pore space (WFPS) (Kesselmeier, Teusch & Kuhn, 1996; Van Diest & Kesselmeier, 2007; Whelan et al., 2016). Recently a theoretical framework was advanced describing how WFPS influences the diffusion of COS in the soil matrix and how this partially regulates the rate of COS hydrolysis by the enzyme CA in addition to temperature and COS concentration (Ogée et al., 2016). Guided by this theoretical framework we estimated that by maintaining moisture levels in our soils at ~30% water holding capacity we would be conducting our experiments very close to the optimum WFPS (between 15 to 37%) for gross COS uptake. However, it was not clear whether COS production should also respond to variable soil water content. As it is much easier and convenient to store dry soil and measure in the lab we felt it was important to test this assumption as this could open up the possibility of greater spatial characterisation of soil COS production rates if the dry soil was demonstrated to be a robust proxy for COS production rates at optimum WFPS conditions. In general, the experimental results supported the use of dry soil COS production rates to confidently infer COS production rates in moist soils at optimal moisture conditions (Fig 2a). In addition, a recent study by Bunk et al. (2017) measuring the net COS flux of a tropical soil over a range of WHC (between 3 and 90%)

also showed that the COS production rate (estimated as the net COS flux measured after a fungicide treatment) was not responsive to soil moisture variations providing support for our experimental results. However, in contrast Bunk et al. (2017) also reported that a temperate agricultural soil did exhibit a soil moisture response that they attributed to variations in the COS production flux rate, in contradiction with the current theory presented in Eq. 1. Indeed when the hydrolysis rate constant tends to zero ($k \rightarrow 0$), Eq. 1 simplifies to $F = \rho_b P / z_{\max}$ so that the net COS flux F should become independent of soil moisture, provided P does not respond to soil moisture. Thus to reconcile with theory the results from Bunk et al. (2017), we would need to invoke a partial and/or non-uniform inhibition of k by the fungicide application (used to infer the production rate effect). A non-uniform reduction of soil moisture upon drying could also create a soil moisture response without the need to evoke a dependence of P on soil water availability. Interestingly, if we extrapolate the results of Bunk et al. (2017) at 0% WHC, the (fully-dry) net COS flux would correspond reasonably well to the fungicide-inhibited flux measured at 30% WHC, and would completely corroborate the results presented in the current study (Fig. 2a).

A further result of our study showed that when using Eq. 1 assuming P_{dry} as a proxy for P to estimate the COS hydrolysis rate constant k , the uncertainty on k increased at lower atmospheric COS concentrations (Fig. 5). This is because, as the atmospheric COS concentration decreases, the contribution of the sink term to the overall net COS flux becomes progressively smaller. In addition and more importantly, this increased uncertainty was biased towards smaller k values (Fig. 5), with the median k value significantly lower (by around 20%) at *ca.* 100 ppt compared to that obtained at *ca.* 800 ppt. Even at *ca.* 800 ppt, the recalculated k ($k_{\text{recal,high}}$) was still on average smaller than k_{moist} (Fig. 2b). In addition, it is worth noting that the bias between k_{recal} and C_a cannot be explained by Michaelis-Menten kinetics. Indeed, according to such an enzymatic model, k should remain constant for $C_a \ll K_m$ and should decrease, not increase, at higher C_a concentrations. Thus, overall, our results suggest that studies using P_{dry} (Whelan et al., 2016) can provide a good proxy of P in moist conditions but this may lead to a small but probably insignificant underestimation of the COS uptake rate constant if measurements are made over the same range of COS concentrations tested in our study.

4.2 Soils generally act as COS sinks at cool temperatures but become COS sources rapidly upon warming

Across a range of biome and land use types we found that the relative contribution of COS production to the net soil COS flux was generally smaller than the COS uptake rate at 18°C, but increased at higher temperatures (23°C) and lower atmospheric COS concentrations. At COS concentrations close to those found in the atmosphere (~500 ppt), net COS fluxes were always negative in our soils when measured at 18°C, indicating that the compensation point (*i.e.* the COS concentration at which the net flux is zero) was always below the atmospheric COS concentration. Even at 100 ppt, only five of the soils had positive net COS fluxes indicating that the COS compensation point was generally lower than 100 ppt for the majority of the soils. This is consistent with previous studies on oxic soils (Kesselmeier et al., 1999; Liu et al., 2010) but contradictory to the results of Lehman and Conrad (1996) who found much higher compensation points. This apparent contradiction might be explained by the much higher and wider range of COS concentrations (60-410 ppb) explored by Lehmann & Conrad (1996) where different COS consumption processes might take place (e.g. physio-sorption; Conrad and Meuser, 2000). There are further studies

conducted in sub-tropical monsoon humid climates that have also reported COS compensation points above 100 ppt (Geng and Mu, 2004; Yi and Wang, 2011), but still below atmospheric concentrations (i.e. around 300 ppt, respectively). These higher compensation points might be explained by the warmer temperatures expected in this type of climate that should favour COS production over consumption (Fig. 6), shifting the compensation point to higher COS concentration values and even causing some soils to become net COS emitters. Because the temperature sensitivity of the production rate is always larger than that of the hydrolysis constant, a potential shift in the optimum temperature of COS uptake would not be enough to offset the larger production rates at the higher temperatures. This relatively greater temperature sensitivity of COS production rates found in our experiment are also consistent with a number of previous studies reporting the temperature sensitivities (Q_{10}) of production-dominated net COS soil fluxes in the range of 1.7 to 3.3 (Maseyk et al., 2014; Saito et al., 2002; Saito et al., 2002; Whelan and Rhew, 2015, 2016). Altogether, our results show that soil COS production (and its contribution to the net COS flux) varies across different biome and temperature regimes and must be accounted for when performing atmospheric COS budgets.

4.3 Soil COS production rates also increase with soil N content and mean annual precipitation

Currently, COS emissions by oxic soils are considered to be abiotic in origin (Kitz et al., 2017; Whelan and Rhew, 2015). Dramatic COS production rates have been observed across US and Chinese agricultural soils (Billesbach et al., 2014; Liu et al., 2010; Maseyk et al., 2014; Whelan and Rhew, 2015). However, the exact mechanisms underlying COS production are still under debate (Whelan et al., 2016). A number of hypotheses including the thermal degradation of soil organic matter or desorption of COS from soil surfaces have been proposed and are partially supported by the persistence of COS emissions after autoclaving (Kato et al., 2008; Whelan & Rhew, 2015; Whelan et al., 2016). Another abiotic process that could lead to COS production is the chemical reaction that occurs in flue gas from molecules present during combustion such as $\text{CH}_4 + \text{SO}_2 \Leftrightarrow \text{COS} + \text{H}_2\text{O} + \text{H}_2$ (Rhodes et al., 2000). Both sulphur dioxide (SO_2) and methane (CH_4) can be produced in soils, however CH_4 is generally produced in anaerobic zones of submerged soils and tends not to accumulate at the soil surface (Le Mer and Roger, 2001). Thus it is not clear whether this reaction would be possible in aerobic, dry soils and maintain the same rate when moist. The thermal decomposition of CH_3SCO radicals (Barnes et al., 1994) and the oxidation of thioformaldehyde and DMS (Barnes et al., 1996) present other reactions that could lead to the production of COS. However, these two reactions are unlikely to explain our results on dark-incubated soils, as both reactions require the photolysis or photoproduction of certain compounds for this reaction to proceed.

On the other hand there is growing evidence that biotic processes may also contribute to COS emission rates (Whelan et al., 2017). In particular a number of studies provide direct evidence for the production of COS during the hydrolysis of thiocyanates when catalysed by thiocyanate hydrolase, an enzyme found in a range of bacteria (Katayama et al., 1992; Kim and Katayama, 2000; Ogawa et al., 2013) and a few fungi (Masaki et al., 2016). If COS production rates were even partially driven by such biotic processes, this contribution might be sensitive to soil water content and expected to decrease at very low soil water content as microbial activity tends to slow down and microbes enter either a stationary growth phase and/or a dormant state

(Roszak and Colwell, 1987). However, we did not observe any significant reduction in COS production rates after air drying of the soils (Fig. 2a). One potential explanation for this could be that some microorganisms can persist for prolonged periods of time in drought conditions, utilising energy reserves at a very slow rate (Raubuch et al., 2002) but nonetheless remain metabolically active (Manina and McKinney, 2013). For example, Zoppini and Marxsen (2010) demonstrated that some extracellular activities in river sediments were not reduced even after one year of drying. This can arise as air-dried soils can still contain some residual water in soil micropores that maintain enzymatic activity. The amount of liquid water required for maintaining such biological activity, including thiocyanate hydrolase activity, could be extremely small and still result in a detectable amount of COS emitted. In addition, Maire et al. (2013) showed that endoenzymes released from dead organisms were stabilised in soils and could still lead to extracellular oxidative metabolism. This could also partly explain the continuation of COS production even at very low water content in our soils. In this context, even sterilised (autoclaved) soils might still produce COS as microbial death can release nutrients and intracellular metabolites into the soil environment, including endoenzymes capable of resisting the autoclaving process. Although our results cannot rule out any of the above mechanistic hypotheses, our results and previous studies indicate overall that the COS emission rates of air-dried and autoclaved soils are related to past biotic activity and in particular the soil nitrogen status. Our study showed that the magnitude of COS emitted from soils was positively correlated to total N concentration (Figs. 2a and 4) over a range of soil N concentrations between 0.38 and 10.2 g kg⁻¹ (Table S2). Although this is the first study to demonstrate a significant relationship between soil N concentration and gross COS production rates, previous studies have measured shifts in the magnitude of net COS and CS₂ fluxes upon fertilisation with nitrate in both deciduous and evergreen coniferous forests (Melillo & Steudler, 1989). In addition a number of studies on agricultural soils (typically fertilised) in the US and China have observed large temperature-sensitive emissions of COS (Billesbach et al., 2014; Liu et al., 2010; Maseyk et al., 2014; Whelan & Rhew, 2015). Currently, the mechanism(s) for the observed increase in COS production rates with increasing N content in soils is clearly not understood. However it is known that S-containing amino acids such as methionine, cystine and cysteine are all potential precursors of COS and CS₂ formation (Bremner & Steele, 1978; Minami & Fukushi, 1981a; Minami & Fukushi, 1981b) and could be degraded abiotically for example in the presence of ozone (Sharma & Graham, 2010) or light (Whelan & Rhew, 2015). However, all measurements in our study were conducted in the dark, at least eliminating this latter mechanism as a potential driver of our results. Thus ecosystems exposed to higher nitrogen inputs either naturally or by enhanced fertilisation may be creating plant and soil organic matter that contains relatively more N- and S-containing precursors such as amino acids and proteins that then become available substrates in soils for temperature-sensitive abiotic degradation. Further studies investigating the link between soil N inputs and soil COS fluxes would now be useful to assess whether total soil N and soil microbial N biomass as traits could be helpful integrated predictors of how soil COS production and uptake rates, respectively vary across large spatial scales irrespective of whether the underlying mechanism of COS production is abiotic or biotic in nature. Meanwhile parallel studies clarifying the mechanistic processes underlying the production will aid the development of models attempting to describe dynamically the instantaneous exchange between soils and the atmosphere and their link to climate, vegetation type and management regime.

4.4 The soil COS uptake rate constant increases with soil microbial content and has a small temperature sensitivity

Direct evidence for the role of carbonic anhydrase (CA) in the uptake of COS has been established in past lab experiments with plant extracts (Protoschill-Krebs et al., 1996) and indirectly on soils treated with CA inhibitors (Kesselmeier et al., 1999). Based on the theoretical framework that exists for the catalysis of CO₂ uptake by CA in soils (Wingate et al. 2010; Sauze et al., 2017b), Ogée et al. (2016) developed an analogous framework to describe the uptake of COS by CA in soils (Eq. 1) and was able to reproduce the observed response of the net and gross COS uptake rate with water-filled pore space and its optimum. Our study unequivocally suggests that the response of the net COS uptake to soil water content is dominated by changes in the gross COS uptake, not the COS production rate, in agreement with Eq. 1.

An important parameter in this modelling framework is the temperature sensitivity (Q_{10}) of the CA-catalysed COS hydrolysis rate k . In the present study a mean value of 1.23 ± 0.29 was estimated for the Q_{10} of k_{moist} over the entire range of 27 soils and exhibited much lower variability than the temperature sensitivity response of gross COS production (Fig. 6). Although the range of Q_{10} for the hydrolysis rate was linearly and negatively related to soil C content (Fig. 4), this parameter appeared fairly conservative amongst the different soils, and its mean value was also consistent with a range of published Q_{10} values (1.22 to 1.9) for plant CA extracts (Burnell and Hatch, 1988; Boyd et al., 2015, Ogée et al., 2016), reinforcing the idea that the uptake of COS by the soils studied is driven by CA activity.

The large scale variability in the COS hydrolysis rate (at a given temperature and 30% WHC) was mostly related to variations in microbial C and N biomass (Fig. 4), and the majority of the smaller COS hydrolysis rates were indeed found in soils with the lowest microbial biomass. This result is consistent with the model of Ogée et al. (2016) that proposes soil CA activity to vary proportionally to the total volume of all the microbes, present in a soil provided that their CA requirements are similar. Our study, in addition to two further field studies (Saito et al., 2002; Yi et al., 2007), provide support for such a hypothesis, although differences in pH (Ogée et al. 2016; Sauze et al., 2017b) and microbial community structure (Sauze et al. 2017b) may complicate the relationship between the COS uptake rate constant and microbial biomass.

5 Conclusions

Uncertainties in the contribution of oxic soils to the atmospheric mass balance are large, with estimates for the global soil sink strength varying from between 70 and 510 GgS y⁻¹ (Berry et al., 2013; Campbell et al., 2017; Launois et al., 2015; Montzka et al., 2007; Suntharalingam et al., 2008). Although developments in the mechanistic understanding and modelling of soil-atmosphere COS have been made recently (Ogée et al., 2016; Sun et al., 2016) it still remains a challenge to extend the observations of a limited set of experimentally different datasets to robust descriptions of soil-atmosphere COS exchange in land surface models. This lack of coherently collected data across multiple biomes and land use types currently hinders advances in modelling the variability in atmospheric COS concentrations at the large scale. Our study goes some way towards addressing this gap by providing a comprehensive dataset of partitioned COS fluxes across Europe and Israel alongside the prominent soil characteristics that are commonly measured and mapped, providing potential transfer functions that can

translate soil physical and chemical properties into globally gridded maps of COS production and uptake rates by soils. In particular we showed that COS hydrolysis rates were linked to microbial C biomass whilst COS production rates were linked to soil N content and MAP. In addition both of these gross COS fluxes exhibited distinctly different temperature and moisture sensitivities. These different soil proprieties should now be explored more deeply to determine their added-value in the prediction of soil COS fluxes and their ability to reconcile the contribution of soil COS fluxes to the atmospheric COS budget.

Author contribution

AK, JO, LW, JS designed the experiments and AK, JS, SW, SJ and AG carried them out. AK prepared the manuscript with contributions from JO and LW.

Competing interests

10 The authors declare that they have no conflict of interest.

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP7/2007-2013) (grant agreement No. 338264, awarded to L.W), the French Agence Nationale de la Recherche (ANR) (Grant Agreement No. ANR-13-BS06- 0005-01) and the Institut National de la Recherche Agronomique (INRA) departments EFPA and EA (PhD studentship for JS). We are also grateful to all the scientific teams across Europe and Israel that provided their valuable time to collect soils from their local experimental sites for this study especially Jorge Curiel-Yuste, Alexandria Correia, Jean-Marc Ourcival, Jukka Pumpanen, Huizhong Zhang, Carmen Emmel, Nina Buchmann, Sabina Keller, Irene Lehner, Anders Lindroth, Andreas Ibrom, Jens Schaarup Sorensen, Dan Yakir, Fulin Yang, Michal Heliasz, Susanne Burri, Penelope Serrano Ortiz, Maria Rosario Moya Jimenez, Jose Luis Vicente, Holger Tulp, Per Marklund, John Marshall, Nils Henriksson, Raquel Lobo de Vale, Lukas Siebicke, Bernard Longdoz, Pascal Courtois, Katja Klumpp.

References

- Barnes, I., Becker, K. H. and Patroescu, I.: The tropospheric oxidation of dimethyl sulfide: A new source of carbonyl sulfide, *Geophys. Res. Lett.*, 21(22), 2389–2392, doi:10.1029/94GL02499, 1994.
- 25 Barnes, I., Becker, K. H. and Patroescu, I.: FTIR product study of the OH initiated oxidation of dimethyl sulphide: Observation of carbonyl sulphide and dimethyl sulphoxide, *Atmospheric Environment*, 30(10), 1805–1814, doi:10.1016/1352-2310(95)00389-4, 1996.

- Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., Seibt, U., Stimler, K., Yakir, D. and Zhu, Z.: A coupled model of the global cycles of carbonyl sulfide and CO₂: A possible new window on the carbon cycle, *Journal of Geophysical Research: Biogeosciences*, 118(2), 842–852, doi:10.1002/jgrg.20068, 2013.
- 5
- Billesbach, D. P., Berry, J. A., Seibt, U., Maseyk, K., Torn, M. S., Fischer, M. L., Abu-Naser, M. and Campbell, J. E.: Growing season eddy covariance measurements of carbonyl sulfide and CO₂ fluxes: COS and CO₂ relationships in Southern Great Plains winter wheat, *Agricultural and Forest Meteorology*, 184, 48–55, doi:10.1016/j.agrformet.2013.06.007, 2014.
- 10
- Boyd, R. A., Gandin, A. and Cousins, A. B.: Temperature Responses of C₄ Photosynthesis: Biochemical Analysis of Rubisco, Phosphoenolpyruvate Carboxylase, and Carbonic Anhydrase in *Setaria viridis*, *Plant Physiology*, 169(3), 1850–1861, doi:10.1104/pp.15.00586, 2015.
- Bremner, J. M. and Steele, C. G.: Role of Microorganisms in the Atmospheric Sulfur Cycle, in *Advances in Microbial Ecology*, pp. 155–201, Springer, Boston, MA., 1978.
- 15
- Borchers, H.W. (2017). *pracma: Practical Numerical Math Functions*. R package version 2.0.7.
<https://CRAN.R-project.org/package=pracma>
- 20
- Bunk, R., Behrendt, T., Yi, Z., Andreae, M. O. and Kesselmeier, J.: Exchange of carbonyl sulfide (OCS) between soils and atmosphere under various CO₂ concentrations, *J. Geophys. Res. Biogeosci.*, 122(6), 2016JG003678, doi:10.1002/2016JG003678, 2017.
- Burnell, J. N. and Hatch, M. D.: Low Bundle Sheath Carbonic Anhydrase Is Apparently Essential for Effective C₄ Pathway Operation, *Plant Physiology*, 86(4), 1252–1256, doi:10.1104/pp.86.4.1252, 1988.
- 25
- Brühl, C., Lelieveld, J., Crutzen, P.J. & Tost, H.: The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate, *Atmospheric, Chemistry and Physics*, 12, 1239-1253, doi:10.5194/acp-12-1239-2012, 2012.
- 30
- Campbell, J. E., Berry, J. A., Seibt, U., Smith, S. J., Montzka, S. A., Launois, T., Belviso, S., Bopp, L. and Laine, M.: Large historical growth in global terrestrial gross primary production, *Nature*, 544(7648), 84, doi:10.1038/nature22030, 2017.
- Castro, M. S. and Galloway, J. N.: A comparison of sulfur-free and ambient air enclosure techniques for measuring the exchange of reduced sulfur gases between soils and the atmosphere, *J. Geophys. Res.*, 96(D8), 15427–15437, doi:10.1029/91JD01399, 1991.
- 35

- Conrad, R.: Compensation concentration as critical variable for regulating the flux of trace gases between soil and atmosphere, *Biogeochemistry*, 27(3), 155–170, doi:10.1007/BF00000582, 1994.
- Conrad, R. and Meuser : Soils contain more than one activity consuming carbonyl sulfide, *Atmospheric Environment*, 34(21),
5 3635-3639, doi.org/10.1016/S1352-2310(00)00136-9, 2000.
- Elleuche, S. and Pöggeler, S.: Carbonic anhydrases in fungi, *Microbiology*, 156(1), 23–29, doi:10.1099/mic.0.032581-0, 2010.
- Fabre, N., Reiter, I. M., Becuwe-Linka, N., Genty, B. and Rumeau, D.: Characterization and expression analysis of genes
10 encoding α and β carbonic anhydrases in Arabidopsis, *Plant, Cell & Environment*, 30(5), 617–629, doi:10.1111/j.1365-3040.2007.01651.x, 2007.
- Fried, A., Klinger, L. F. and Erickson, D. J.: Atmospheric carbonyl sulfide exchange in bog microcosms, *Geophys. Res. Lett.*,
15 20(2), 129–132, doi:10.1029/93GL00062, 1993.
- Geng, C. and Mu, Y.: Carbonyl sulfide and dimethyl sulfide exchange between lawn and the atmosphere, *J. Geophys. Res.*,
109(D12), D12302, doi:10.1029/2003JD004492, 2004.
- Gimeno, T. E., Ogée, J., Royles, J., Gibon, Y., West, J. B., Burlett, R., Jones, S. P., Sauze, J., Wohl, S., Benard, C., Genty, B.
20 and Wingate, L.: Bryophyte gas-exchange dynamics along varying hydration status reveal a significant carbonyl sulphide (COS) sink in the dark and COS source in the light, *New Phytol*, 215(3), 965–976, doi:10.1111/nph.14584, 2017.
- Haney, R. L. and Haney, E. B.: Simple and Rapid Laboratory Method for Rewetting Dry Soil for Incubations, *Communications
in Soil Science and Plant Analysis*, 41(12), 1493–1501, doi:10.1080/00103624.2010.482171, 2010.
- 25 Haritos, V. S. and Dojchinov, G.: Carbonic anhydrase metabolism is a key factor in the toxicity of CO₂ and COS but not CS₂ toward the flour beetle *Tribolium castaneum* [Coleoptera: Tenebrionidae], *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 140(1), 139–147, doi:10.1016/j.cca.2005.01.012, 2005.
- 30 Hines, M. E. and Morrison, M. C.: Emissions of biogenic sulfur gases from Alaskan tundra, *J. Geophys. Res.*, 97(D15), 16703–16707, doi:10.1029/90JD02576, 1992.
- Jarvis, P. G., Rey, A., Petsikos, C., Wingate, L., Rayment, M.B., Pereira, J.S., Banza, J., David, J., Miglietta, F., Borgetti, M.
and Valentini, R. (2007) Drying and wetting of Mediterranean soils stimulates decomposition and carbon dioxide emission:
35 the “Birch Effect” *Tree Physiology Vol. 27 Issue 7, 929-940*

- Katayama, Y., Narahara, Y., Inoue, Y., Amano, F., Kanagawa, T. and Kuraishi, H.: A thiocyanate hydrolase of *Thiobacillus thioparus*. A novel enzyme catalyzing the formation of carbonyl sulfide from thiocyanate., *J. Biol. Chem.*, 267(13), 9170–9175, 1992.
- 5 Kato, H., Saito, M., Nagahata, Y. and Katayama, Y.: Degradation of ambient carbonyl sulfide by *Mycobacterium* spp. in soil, *Microbiology*, 154(1), 249–255, doi:10.1099/mic.0.2007/011213-0, 2008.
- Kesselmeier, J., Teusch, N. and Kuhn, U.: Controlling variables for the uptake of atmospheric carbonyl sulfide by soil, *J. Geophys. Res.*, 104(D9), 11577–11584, doi:10.1029/1999JD900090, 1999.
- 10 Kettle, A. J., Kuhn, U., von Hobe, M., Kesselmeier, J. and Andreae, M. O.: Global budget of atmospheric carbonyl sulfide: Temporal and spatial variations of the dominant sources and sinks, *J.-Geophys.-Res.*, 107(D22), 4658, doi:10.1029/2002JD002187, 2002.
- 15 Kim, S.-J. and Katayama, Y.: Effect of growth conditions on thiocyanate degradation and emission of carbonyl sulfide by *Thiobacillus thioparus* THI115, *Water Research*, 34(11), 2887–2894, doi:10.1016/S0043-1354(00)00046-4, 2000.
- Kitz, F., Gerdel, K., Hammerle, A., Laterza, T., Spielmann, F. M. and Wohlfahrt, G.: In situ soil COS exchange of a temperate mountain grassland under simulated drought, *Oecologia*, 183(3), 851–860, doi:10.1007/s00442-016-3805-0, 2017.
- 20 Kooijmans, L. M. J., Uitslag, N. A. M., Zahniser, M. S., Nelson, D. D., Montzka, S. A. and Chen, H.: Continuous and high-precision atmospheric concentration measurements of COS, CO₂, CO and H₂O using a quantum cascade laser spectrometer (QCLS), *Atmos. Meas. Tech.*, 9(11), 5293–5314, doi:10.5194/amt-9-5293-2016, 2016.
- 25 Launois, T., Belviso, S., Bopp, L., Fichot, C. G. and Peylin, P.: A new model for the global biogeochemical cycle of carbonyl sulfide – Part 1: Assessment of direct marine emissions with an oceanic general circulation and biogeochemistry model, *Atmos. Chem. Phys.*, 15(5), 2295–2312, doi:10.5194/acp-15-2295-2015, 2015.
- 30 Le Mer, J. and Roger, P.: Production, oxidation, emission and consumption of methane by soils: A review, *European Journal of Soil Biology*, 37(1), 25–50, doi:10.1016/S1164-5563(01)01067-6, 2001.
- Lehmann, S. and Conrad, R.: Characteristics of turnover of carbonyl sulfide in four different soils, *J Atmos Chem*, 23(2), 193–207, doi:10.1007/BF00048260, 1996.
- 35 Li, W., Yu, L., Yuan, D., Wu, Y. and Zeng, X.: A study of the activity and ecological significance of carbonic anhydrase from soil and its microbes from different karst ecosystems of Southwest China, *Plant Soil*, 272(1–2), 133–141, doi:10.1007/s11104-004-4335-9, 2005.

- Liu, J., Geng, C., Mu, Y., Zhang, Y., Xu, Z. and Wu, H.: Exchange of carbonyl sulfide (COS) between the atmosphere and various soils in China, *Biogeosciences*, 7(2), 753–762, doi:10.5194/bg-7-753-2010, 2010.
- 5 Maire, V., Alvarez, G., Colombet, J., Comby, A., Despinasse, R., Dubreucq, E., Joly, M., Lehours, A.-C., Perrier, V., Shahzad, T. and Fontaine, S.: An unknown oxidative metabolism substantially contributes to soil CO₂ emissions, *Biogeosciences*, 10(2), 1155–1167, doi:10.5194/bg-10-1155-2013, 2013.
- Manina, G. and McKinney, J.D.: A Single-Cell Perspective on Non-Growing but Metabolically Active (NGMA) Bacteria, in
10 Pathogenesis of *Mycobacterium tuberculosis* and its Interaction with the Host Organism, pp. 135-161, Springer, Berlin, Heidelberg, 2013.
- Masaki, Y., Ozawa, R., Kageyama, K. and Katayama, Y.: Degradation and emission of carbonyl sulfide, an atmospheric trace gas, by fungi isolated from forest soil, *FEMS Microbiol Lett*, 363(18), doi:10.1093/femsle/fnw197, 2016.
- 15 Maseyk, K., Berry, J. A., Billesbach, D., Campbell, J. E., Torn, M. S., Zahniser, M. and Seibt, U.: Sources and sinks of carbonyl sulfide in an agricultural field in the Southern Great Plains, *PNAS*, 111(25), 9064–9069, doi:10.1073/pnas.1319132111, 2014.
- Melillo, J. M., McGuire, A. D., Kicklighter, D. W., Moore, B., Vorosmarty, C. J. and Schloss, A. L.: Global climate change
20 and terrestrial net primary production, *Nature*, 363, 234, 1993.
- Melillo, J. M. and Steudler, P. A.: The effect of nitrogen fertilization on the COS and CS₂ emissions from temperate forest soils, *J Atmos Chem*, 9(4), 411–417, doi:10.1007/BF00114753, 1989.
- 25 de Mello, W. Z. and Hines, M. E.: Application of static and dynamic enclosures for determining dimethyl sulfide and carbonyl sulfide exchange in Sphagnum peatlands: Implications for the magnitude and direction of flux, *J. Geophys. Res.*, 99(D7), 14601–14607, doi:10.1029/94JD01025, 1994.
- 30 Minami, K. and Fukushi, S.: Detection of carbonyl sulfide among gases produced by the decomposition of cystine in paddy soils, *Soil Science and Plant Nutrition*, 27(1), 105–109, doi:10.1080/00380768.1981.10431259, 1981a.
- Minami, K. and Fukushi, S.: Volatilization of carbonyl sulfide from paddy soils treated with sulfur-containing substances, *Soil Science and Plant Nutrition*, 27(3), 339–345, doi:10.1080/00380768.1981.10431288, 1981b.
- 35 Moldrup, P., Yoshikawa, S., Olesen, T., Komatsu, T. and Rolston, D.E.: Gas Diffusivity in Undisturbed Volcanic Ash Soils, *Soil Science Society of America Journal*, 67(1), 41-51, doi: 10.2136/sssaj2033.4100, 2003.

- Montzka, S. A., Calvert, P., Hall, B. D., Elkins, J. W., Conway, T. J., Tans, P. P. and Sweeney, C.: On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO₂, *J. Geophys. Res.*, 112(D9), D09302, doi:10.1029/2006JD007665, 2007.
- 5 Moroney, J. V., Bartlett, S. G. and Samuelsson, G.: Carbonic anhydrases in plants and algae, *Plant, Cell & Environment*, 24(2), 141–153, doi:10.1111/j.1365-3040.2001.00669.x, 2001.
- Ogawa, T., Noguchi, K., Saito, M., Nagahata, Y., Kato, H., Ohtaki, A., Nakayama, H., Dohmae, N., Matsushita, Y., Odaka, M., Yohda, M., Nyunoya, H. and Katayama, Y.: Carbonyl Sulfide Hydrolase from *Thiobacillus thioparus* Strain THI115 Is
10 One of the β -Carbonic Anhydrase Family Enzymes, *J. Am. Chem. Soc.*, 135(10), 3818–3825, doi:10.1021/ja307735e, 2013.
- Ogéé, J., Sauze, J., Kesselmeier, J., Genty, B., Van Diest, H., Launois, T. and Wingate, L.: A new mechanistic framework to predict OCS fluxes from soils, *Biogeosciences*, 13(8), 2221–2240, doi:10.5194/bg-13-2221-2016, 2016.
- Protoschill-Krebs, G. and Kesselmeier, J.: Enzymatic Pathways for the Consumption of Carbonyl Sulphide (COS) by Higher
15 Plants*, *Botanica Acta*, 105(3), 206–212, doi:10.1111/j.1438-8677.1992.tb00288.x, 1992.
- Protoschill-Krebs, G., Wilhelm, C. and Kesselmeier, J.: Consumption of carbonyl sulphide (COS) by higher plant carbonic anhydrase (CA), *Atmospheric Environment*, 30(18), 3151–3156, doi:10.1016/1352-2310(96)00026-X, 1996.
- 20 R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria. [online] Available from: <https://www.R-project.org/>, 2017.
- Raubuch, M., Dyckmans, J., Joergensen, R. G. and Kreutzfeldt, M.: Relation between respiration, ATP content, and Adenylate Energy Charge (AEC) after incubation at different temperatures and after drying and rewetting, *Journal of Plant Nutrition and
25 Soil Science*, 165(4), 435–440, doi:10.1002/1522-2624(200208)165:4<435::AID-JPLN435>3.0.CO;2-3, 2002.
- Rhodes, C., Riddel, S. A., West, J., Williams, B. P. and Hutchings, G. J.: The low-temperature hydrolysis of carbonyl sulfide and carbon disulfide: a review, *Catalysis Today*, 59(3), 443–464, doi:10.1016/S0920-5861(00)00309-6, 2000.
- 30 Roszak, D. B. and Colwell, R. R.: Survival strategies of bacteria in the natural environment., *Microbiol. Rev.*, 51(3), 365–379, 1987.
- Saito, M., Honna, T., Kanagawa, T. and Katayama, Y.: Microbial Degradation of Carbonyl Sulfide in Soils, *Microbes Environ.*, 17(1), 32–38, doi:10.1264/jsme2.2002.32, 2002.

- Sandoval-Soto, L., Stanimirov, M., von Hobe, M., Schmitt, V., Valdes, J., Wild, A. and Kesselmeier, J.: Global uptake of carbonyl sulfide (COS) by terrestrial vegetation: Estimates corrected by deposition velocities normalized to the uptake of carbon dioxide (CO₂), *Biogeosciences*, 2(2), 125–132, doi:10.5194/bg-2-125-2005, 2005.
- 5 Sauze, J., Ogée, J., Maron, P.-A., Crouzet, O., Nowak, V., Wohl, S., Kaisermann, A., Jones, S. P. and Wingate, L.: The interaction of soil phototrophs and fungi with pH and their impact on soil CO₂, CO¹⁸O and OCS exchange, *Soil Biology and Biochemistry*, 115(Supplement C), 371–382, doi:10.1016/j.soilbio.2017.09.009, 2017a.
- Sauze, J., Jones, S., Wingate, L., Wohl, S. and Ogée, J. The role of soil pH on soil carbonic anhydrase activity. *Biogeosciences*
10 Discussions, 2017b.
- Sharma, V.K. & Graham, H.J.D.: Oxidation of amino acids, peptides and proteins by ozone: a review, *Ozone: Science & Engineering*, 32: 81-90 doi: 10.1080/01919510903510507.
- 15 Smith, K. S. and Ferry, J. G.: Prokaryotic carbonic anhydrases, *FEMS Microbiology Reviews*, 24(4), 335–366, doi:10.1111/j.1574-6976.2000.tb00546.x, 2000.
- Stimler, K., Berry, J. A. and Yakir, D.: Effects of Carbonyl Sulfide and Carbonic Anhydrase on Stomatal Conductance, *Plant Physiology*, 158(1), 524–530, doi:10.1104/pp.111.185926, 2012.
20
- Sun, W., Maseyk, K., Lett, C. and Seibt, U.: A soil diffusion–reaction model for surface COS flux: COSSM v1, *Geosci. Model Dev.*, 8(10), 3055–3070, doi:10.5194/gmd-8-3055-2015, 2015.
- Sun, W., Maseyk, K., Lett, C. and Seibt, U.: Litter dominates surface fluxes of carbonyl sulfide in a Californian oak woodland,
25 *J. Geophys. Res. Biogeosci.*, 121(2), 2015JG003149, doi:10.1002/2015JG003149, 2016.
- Suntharalingam, P., Kettle, A. J., Montzka, S. M. and Jacob, D. J.: Global 3-D model analysis of the seasonal cycle of atmospheric carbonyl sulfide: Implications for terrestrial vegetation uptake, *Geophys. Res. Lett.*, 35(19), L19801, doi:10.1029/2008GL034332, 2008.
30
- Ueno, Y., Johnson, M. S., Danielache, S. O., Eskebjerg, C., Pandey, A. and Yoshida, N.: Geological sulfur isotopes indicate elevated OCS in the Archean atmosphere, solving faint young sun paradox, *PNAS*, 106(35), 14784–14789, doi:10.1073/pnas.0903518106, 2009.
- 35 Van Diest, H. and Kesselmeier, J.: Soil atmosphere exchange of carbonyl sulfide (COS) regulated by diffusivity depending on water-filled pore space, *Biogeosciences*, 5(2), 475–483, doi:10.5194/bg-5-475-2008, 2008.

- Van Veldhoven, P. P. and Mannaerts, G. P.: Inorganic and organic phosphate measurements in the nanomolar range, *Analytical Biochemistry*, 161(1), 45–48, doi:10.1016/0003-2697(87)90649-X, 1987.
- Wei, T. and Simko, V.: R package “corrplot”: Visualization of a Correlation Matrix. [online] Available from:
5 <https://github.com/taiyun/corrplot>, 2017.
- Whelan, M. E. and Rhew, R. C.: Carbonyl sulfide produced by abiotic thermal and photodegradation of soil organic matter from wheat field substrate, *J. Geophys. Res. Biogeosci.*, 120(1), 2014JG002661, doi:10.1002/2014JG002661, 2015.
- 10 Whelan, M. E. and Rhew, R. C.: Reduced sulfur trace gas exchange between a seasonally dry grassland and the atmosphere, *Biogeochemistry*, 128(3), 267–280, doi:10.1007/s10533-016-0207-7, 2016a.
- Whelan, M. E., Min, D.-H. and Rhew, R. C.: Salt marsh vegetation as a carbonyl sulfide (COS) source to the atmosphere, *Atmospheric Environment*, 73(Supplement C), 131–137, doi:10.1016/j.atmosenv.2013.02.048, 2013.
- 15 Whelan, M. E., Hilton, T. W., Berry, J. A., Berkelhammer, M., Desai, A. R. and Campbell, J. E.: Carbonyl sulfide exchange in soils for better estimates of ecosystem carbon uptake, *Atmos. Chem. Phys.*, 16(6), 3711–3726, doi:10.5194/acp-16-3711-2016, 2016b.
- 20 Whelan, M. E., Lennartz, S. T., Gimeno, T. E., Wehr, R., Wohlfahrt, G., Wang, Y., Kooijmans, L. M. J., Hilton, T. W., Belviso, S., Peylin, P., Commane, R., Sun, W., Chen, H., Kuai, L., Mammarella, I., Maseyk, K., Berkelhammer, M., Li, K.-F., Yakir, D., Zumkehr, A., Katayama, Y., Ogée, J., Spielmann, F. M., Kitz, F., Rastogi, B., Kesselmeier, J., Marshall, J., Erkkilä, K.-M., Wingate, L., Meredith, L. K., He, W., Bunk, R., Launois, T., Vesala, T., Schmidt, J. A., Fichot, C. G., Seibt, U., Saleska, S., Saltzman, E. S., Montzka, S. A., Berry, J. A. and Campbell, J. E.: Reviews and Syntheses: Carbonyl Sulfide as a Multi-
25 scale Tracer for Carbon and Water Cycles, *Biogeosciences Discuss.*, 2017, 1–97, doi:10.5194/bg-2017-427, 2017.
- Wilhelm, E., Battino, R. and Wilcock, R. J.: Low-pressure solubility of gases in liquid water, *Chem. Rev.*, 77(2), 219–262, doi:10.1021/cr60306a003, 1977.
- 30 Wingate, L., Ogée, J., Cuntz, M., Genty, B., Reiter, I., Seibt, U., Yakir, D., Maseyk, K., Pendall, E., Barbour, M.M., Mortazavi, B., Peylin, P., Miller, J., Mencuccini, M., Burlett, R., Shim, J.H., Hunt, J. & Grace, J. The influence of soil micro-organisms on the oxygen isotope signal of atmospheric CO₂. *Proceedings of the National Academy of Sciences*, Vol. 106, 22411-22415, doi: 10.1073/pnas.0905210106, 2009
- 35 Yi, Z. and Wang, X.: Carbonyl sulfide and dimethyl sulfide fluxes in an urban lawn and adjacent bare soil in Guangzhou, China, *Journal of Environmental Sciences*, 23(5), 784–789, doi:10.1016/S1001-0742(10)60478-0, 2011.

Zoppini, A. and Marxsen, J.: Importance of Extracellular Enzymes for Biogeochemical Processes in Temporary River Sediments during Fluctuating Dry–Wet Conditions, in *Soil Enzymology*, pp. 103–117, Springer, Berlin, Heidelberg., 2010.

SiteID	Country	Site	Biome	Landuse	Latitude	Longitude	Altitude	MAT (°C)	MAP (mm)
FI-Hyy	Finland (FI)	Hyytiala	Boreal	Forest	61.8	24.3	187	3.8	709
FI-Var1	Finland (FI)	Varrio1	Boreal	Forest	67.8	29.6	400	-1	660
FI-Var2	Finland (FI)	Varrio2	Boreal	Forest	67.8	29.6	400	-1	660
SE-Hyl	Sweden (SE)	Hyltemossa	Boreal	Forest	56.1	13.4	111	7	830
SE-Nor	Sweden (SE)	Norunda	Boreal	Forest	60.1	17.5	71	5.6	544
SE-Ros2	Sweden (SE)	Rosinedal_Cont	Boreal	Forest	64.2	19.7	145	1.8	614
SE-Ros1	Sweden (SE)	Rosinedal_Fert	Boreal	Forest	64.2	19.7	145	1.8	614
SE-Sva	Sweden (SE)	Svartberget	Boreal	Forest	64.2	19.8	161	1.8	614
ES-Amo	Spain (ES)	Amoladeras	Mediterranean	Grassland	36.8	-2.3	54	18.5	200
ES-Bal	Spain (ES)	Balsablanca	Mediterranean	Grassland	36.9	-2.0	200	18.5	220
ES-Ube1	Spain (ES)	Ubeda_Veg	Mediterranean	Agrosystem	37.9	-3.2	366	14.8	20.4
ES-Ube2	Spain (ES)	Ubeda_noVeg	Mediterranean	Agrosystem	37.9	-3.2	366	14.8	20.4
IL-Reh	Israel (IL)	Rehovot	Mediterranean	Agrosystem	31.9	34.8	50	19.7	185
IL-Yat	Israel (IL)	Yatir	Mediterranean	Forest	31.3	35.1	661	17.9	292
PT-Cor	Portugal (PT)	Coruche	Mediterranean	Forest	39.1	-8.3	156	17	656
PT-Mit	Portugal (PT)	Mitra	Mediterranean	Forest	38.5	-8.0	240	15.6	656
CH-Cha	Switzerland (CH)	Chamau	Temperate	Grassland	47.2	8.4	393	9.5	1136
CH-Dav	Switzerland (CH)	Davos	Temperate	Forest	46.8	9.9	1639	2.8	1062
CH-Fru	Switzerland (CH)	Fruebuel	Temperate	Grassland	47.1	8.5	982	7.2	1651
CH-Lae	Switzerland (CH)	Laegern	Temperate	Forest	47.1	8.5	866	8.3	1100
CH-Oe2	Switzerland (CH)	Oensingen	Temperate	Agrosystem	47.3	7.7	452	9.2	1100
DE-Hai	Germany (DE)	Hainich	Temperate	Forest	51.1	10.5	463	7.9	646
DE-Lei	Germany (DE)	Leinefelde	Temperate	Forest	51.3	10.4	474	7.6	775
DK-Sor	Denmark (DK)	Soro	Temperate	Forest	55.5	11.6	45	8.5	564
FR-Hes	France (FR)	Hesse	Temperate	Forest	48.7	7.1	313	9.7	650
FR-Lqu1	France (FR)	Laqueuille_Fert	Temperate	Grassland	45.6	2.7	1041	7.9	897
FR-Lqu2	France (FR)	Laqueuille_Cont	Temperate	Grassland	45.6	2.7	1041	7.9	897

Table 1: Locations and names of sites sampled across Europe and Israel describing climatic and land use characteristics. The two names in bold represent the two fertilised parcels in manipulation field experiments. Ubeda_Veg and Ubeda_noVeg represent soils sampled in the same location but with or without vegetation, respectively. Altitude, mean annual temperature (MAT) and precipitation (MAP) are estimated from the literature.

5






Figure 1: Variability in the net COS flux measured at 18°C and at an atmospheric COS concentration of 500 ppt on replicated (n=3) moist (30% WHC) soils sampled from across Europe and Israel (see Table 1). Net COS fluxes were not significantly different between biome or land use.



5 **Figure 2:** Comparison of (a) the soil COS production rates measured on air-dried soils (P_{dry} ; method 1) and moist soils (P_{moist} ; method 2) for the range of total soil nitrogen concentrations (g kg^{-1}) measured at each site and (b) the COS hydrolysis rate of moist soils ($k_{\text{recaI,med}}$ and k_{moist} , respectively) for the range of microbial biomass nitrogen ($\mu\text{g g}^{-1}$) at each site. Each point represents the mean flux \pm SD for each site measured at 18°C ($n=3$). The dashed lines represent the 1:1 slope and the solid black lines represent the slope of the linear models, the grey areas represent the 95% confidence level interval for predictions from the linear models.



5 **Figure 3:** Partitioned mean (\pm SD) gross COS production and COS uptake rates estimated from the measured net COS flux of soils incubated at 18°C and at an atmospheric COS concentration of 500 ppt on moist soils, ranked by biome and by COS production rate in ascending order ($n=3$) for each of the 27 sites sampled in Europe and Israel. Rosinedal_Fert and Laqueuille_Fert are experimental plots intensively fertilised with nitrogen as part of a manipulation experiment. Ubeda_Veg and Ubeda_noVeg represent soils sampled in the same location but with or without vegetation, respectively.




Figure 4: Spearman correlation coefficients (ρ) between variables describing soil COS fluxes and soil properties. F_COS is the net COS flux at 18°C, Source is the gross COS production rate at 18°C, Uptake is the gross COS uptake rate at 18°C, k_{18} is the hydrolysis rate constant at 18°C, Q10P and Q10k are the Q_{10} of the source and the hydrolysis rate, SoilP is the phosphate content, SoilC and SoilN are the soil C and N content, MBC and MBN are the microbial biomass C and N contents, BD is the bulk density, Redox is the potential redox, WFPS is the water filled pore space, Alt is the site altitude, Lon is the site longitude, Lat is the site latitude, MAT and MAP are the mean annual temperature and precipitation for each site. Only significant correlations are shown ($P < 0.05$).




Figure 5: Difference in the COS hydrolysis rate constants of moist soils calculated using the production rates estimated from dried soils (k_{reca} with P_{dry}) and moist soils (k_{moist} with P_{moist}) as a function of three different atmospheric COS concentrations in the outlet. Each grey point represents one soil replicate, with the red box showing the lower quartile, median and upper quartile values, the whiskers indicate the range of variation in the difference, and the box width represents the range of variation in the net COS flux measured for the different soils.




Figure 6: The estimated temperature sensitivity (Q_{10}) of COS production (P_{moist}) and hydrolysis rate (k_{moist}) across 27 sites in Europe and Israel. Each point represents the estimated parameter for each of the 3 replicated microcosms incubated at two temperatures from all sites. The box indicates the lower quartile, median and upper quartile values, the whiskers show the range of variation in the difference and displays the mean (SD) and median value for the Q_{10} parameter.