

Comment of Referee #1:

The work is an extensive analysis of different dust samples, sampled both on land and in the ocean. Samples were analyzed concerning different properties as grain size distributions and mineral composition, aiming at assigning source regions to the samples. Samples on land seem to be more influenced by local sources than oceanic samples, and it is indicated that long range transport decreases the average grain size of the deposited dust. This is all interesting and important information. However, being not a mineralogist, some passages were difficult to follow, and the overall structuring should be improved as well (see below). Also, occasionally statements seem to be too far-fetched. So in total, my judgement is that the manuscript can become fit for publication in ACP, but only after major revisions.

General comments:**Comment of Referee #1:**

Several times in the manuscript, the impression is given that dust settling from the SAL starts to appear at the time when the air mass crosses the shore line, i.e., that SAL is a dust source above the Atlantic, only. This might just be formulated misunderstandingly, but in any case, dust will always settle from the SAL – it is just that on land, where nearby dust sources are present, they might be overwhelming. You have to go through the text and change the respective passages accordingly.

Reply to comment:

This is a good point. Dust already settles on land from the SAL and we formulated this unclearly. We agree, that the passages in which this happened need to be revised. Therefore, we changed all the sentences in which such a misunderstanding could lead to a false view on the dust transport.

Changes made in manuscript:

We modified the following sentences (page and line numbers refer to the revised manuscript with track-changes):

Page 1, lines 19-20:

‘In continental dust source areas dust is also transported in the SAL, however the predominant dust input occurs from nearby dust sources with the low-level trade winds.’

Page 6-7, lines 154-158:

‘Low-level N trade winds blow and transport dust in coastal Mauritania year-round (National Geospatial-Intelligence Agency, 2006). Saharan dust is transported on- and offshore within the ‘Saharan air layer’ (SAL) at

an altitude of about 3 km (Prospero and Carlson, 1970; Carlson and Prospero, 1972; Prospero and Carlson, 1972; Diaz et al., 1976).’

Comment of Referee #1:

Your tables 1 and 2 are helpful, already, but I would have wished for an additional overview, showing which of the different locations contributed data for which kind of analysis. Also, all sites should be introduced together. E.g., Buoy Carmen is treated differently from CB and CBi, appearing for the first time in line 161, but not being shown in Fig. 1 and Fig. 2. Fig. 4 then shows the location of Iwik, CBi and Carmen and also of two new locations, but not the location of CB. This is all confusing, and a table could help to clarify that e.g., one place was only used for its information on meteorology, another one only for size distribution, etc. .

Reply to comment:

We agree that the structure of the manuscript would greatly improve when showing all sites together and when inserting a table showing which analysis has been done at each site. Therefore, we modified figure 1 by inserting all study sites together. Further, we inserted a new table 1 which gives an overview of the study sites with information on the respective location, samples, analyses and data.

Changes made in manuscript:

We inserted the following sentences:

Page 3, line 78:

‘and with the scientific dust-collecting buoy ‘Carmen’

Page 3, lines 90-97:

‘In Fig. 1 the location of the study sites and the North African dust sources are displayed. The dust-collecting buoy ‘Carmen’ (~21°15’ N, ~20°56’ W) and the sediment trap mooring site CB (~21°16’ N, ~20°48’ W) are virtually at the same position ~200 nautical miles offshore Cape Blanc. Sediment-trap station CBi (~20°45’ N, ~18°42’ W) is located ~ 80 nautical miles offshore Cape Blanc. The continental dust collector Iwik (~19°53’ N, ~16° 18’ W) and the meteorological station Arkeiss (~20° 7’ N, ~16° 15’ W) are located in a major potential dust source area (**PSA 2**) in the Parc National de Banc d'Arguin (PNBA) near Iwik and near Arkeiss in Mauritania. A further meteorological station is positioned in the **PSA 2** in Nouadhibou (~20° 55’ N, ~17° 1’ W) in the Western Sahara.’

We shifted the following sentence to:

Page 6, lines 127-128:

‘The local soils surrounding the dust collector site Iwik are composed of sandy deposits often rich in fossil shells and partly cemented by lime (Einsele et al., 1974).’

We modified Fig. 1:

Page 5, lines 107-114:




Figure 1: Map of the study sites under investigation: the scientific buoy Carmen as well as the sediment trap moorings CB and CBi offshore Cape Blanc, the MWAC dust collector onshore near Iwik and the ground stations near Nouadhibou and Arkeiss (shapefile of the surface lithology and the geological provinces: downloaded from the USGS website <http://rmgsc.cr.usgs.gov/ecosystems/africa.shtml#SL> and <http://certmapper.cr.usgs.gov/geoportal/catalog/main/home.page>, major potential dust source areas: redrawn from Scheuvens et al. (2013), ocean currents: redrawn from Mittelstaedt (1991)).

We inserted the following sentences and table:

Page 8, lines 199-208:

‘In Table 1 an overview of the material and methods employed for each study site is presented. Bulk sediment samples were obtained at the sites CB and CBI and dust samples at the sites Carmen and Iwik. All samples were analyzed for particle size and dust flux with the exception of the site Carmen, of which only dust particle size was analyzed. Only the sites CBI and Iwik were analyzed for mineral assemblages and only the samples of the site Iwik were used for microscopic investigation. Meteorological sensors were available for the stations Carmen, Iwik and Arkeiss, while for the site Nouadhibou meteorological data was downloaded online. TRMM precipitation data was downloaded online for all sites except for the site Nouadhibou.

Table 1: Overview of the material and methods employed at each study site. ‘

Study site	Lat./lon.	Samples	Analysis	Meteorological sensor and data	Downloaded meteorological data
Carmen	~21°15' N, ~20°56' W	1 MWAC sample	particle size	Vaisala WXT520: wind direction + speed, precipitation	TRMM 3B42: precipitation
CB	~21°16' N, ~20°48' W	38 sediment trap samples	lithogenic fluxes, particle size	-	TRMM 3B42: precipitation
CBI	~20°45' N, ~18°42' W	38 sediment trap samples	lithogenic fluxes, particle size, mineral assemblages	-	TRMM 3B42: precipitation, HYSPLIT back trajectories
Iwik	~19°53' N, ~16° 18' W	24 MWAC samples	microscopy, dust fluxes, particle size, mineral assemblages	Davis 6250 Vantage Vue: wind direction + speed	TRMM 3B42: precipitation, HYSPLIT back trajectories
Nouadhibou	20° 55' N, 17° 1' W	-	-	-	Wind direction + speed
Arkeiss	~20° 7' N, ~16° 15' W	-	-	Davis 6250 Vantage Vue: precipitation	TRMM 3B42: precipitation

Comment of Referee #1:

Similarly, it is not clear to me what is gained by showing wind directions in Nouadhibou if information from that location is then dismissed due to local effects, unless you want to make it clearer that ground based sampling on

land will not reflect long range transported dust (due to local sources and local winds) – but in this case, this needs to be discussed somewhat stronger. Consequently: if there are these local effects on land, will Iwik not be influenced similarly? This should be discussed, too.

Reply to comment:

We agree that the results on the meteorological data at station Nouadhibou should be mentioned in the discussion. The site served as an example for how local effects may influence the observed wind patterns. At site Iwik the synoptic change in the predominant trade wind direction can be clearly observed, while at other sites like Nouadhibou local effects bias an observation of synoptic wind patterns. This should also be kept in mind when choosing a site for paleoenvironmental reconstructions using sediment core records.

Changes made in manuscript:

We inserted the following sentences:

Page 44, lines 826-829:

‘The variability of the mineralogical composition of dust sampled at site Iwik could be related to the synoptic scale change in the surface trade wind direction. However, meteorological data from nearby sites like e.g., Nouadhibou demonstrate that local effects like the topography exert a strong influence on observed wind directions at ground level. (Fig. 2).’

Comment of Referee #1:

For your determination of possible dust sources, you use trajectories in heights of 10m and 4500m. You chose 10m based on your even lower sampling heights. But still, trajectories in heights as low as 100m or below are always very prone to errors, and 4500m is so height that it might be above the heights of the SAL. Please at least mention that and explain why you chose to calculate the trajectories at these heights, nevertheless.

Alternatively, choose additional different heights (e.g., 100m and 2000m or so) – you could check, if they are similar to those you used. Adding them could add credibility to your work.





Reply to comment:

We chose the altitudes at which back trajectories were calculated on the basis of previous studies to which we compare our data. For example, Skonieczny et al., 2013 use 4500m to study the SAL. Following the reviewer’s suggestion, we now also included additional back trajectories at additional altitudes, which made the observed patterns much clearer. Back-trajectories of the heights 100 m and 3000 m were added to figures 9-12. This improved the detection of the likely dust source areas and the results and interpretation were adjusted.


Changes made in manuscript:

We modified Fig. 9-12:

Page 29, line 602:



Page 32, line 627:



Lithology

Carbonate	Ultramafic	Alluvium - beach, strand, coastal dune
Karst	Extrusive Volcanic	Alluvium - Saline
Non-Carbonate	Colluvium	Alluvium - Gypsum
Metasedimentary	Hydric - Organic	Alluvium - Other
Alkaline Intrusive Volcanic	Aeolian Sediments	Volcanic - Ash, Tuff, Mudflow
Silicic	Alluvium - Fan Deposit	Water
Metaigneous	Alluvium - Fluvial	

Back trajectories

● Iwik ● CBi

— 10 m — 100 m — 10 m — 100 m - - 3000 m - - - 4500 m

PSAs:


□ PSA 1-5




Clay Mineralogy


⊞ No data ⊞ Calcite >8.9 % ⊞ Kaolinite >29 % ⊞ Chlorite >4.1 %

Mineralogy

Kaolinite	Feldspar	Serpentine	Sepiolite	Gibbsite
Chlorite	Mica	Rutile	Anhydrite	Dolomite
Calcite	Amphibole	Zeolite	Smectite	Other minerals
Quartz	Palygorskite	Fluellite	Garnet	★ Ferryglaucophane







We modified the following sentence to:

Page 17, lines 380-384:

‘Four-day back trajectories at altitudes of 10 (following Stuu et al. (2005)), 100, 3000, 4500 (following Skonieczny et al. (2013)) and 5500m were calculated ending at the dust collector site Iwik (19°52’ N, 16°17’ W) and at the proximal marine trap site CBi (20°46’, 18°44’ W) using the Hybrid Single Particle Langrangian

Integrated Trajectory (HYSPLIT) model (Stein et al., 2015) and the reanalysis dataset (2.5° spatial resolution) on the NOAA website at <http://ready.arl.noaa.gov>. ‘

We modified the following sentences:

Page 26, lines 582-583:

‘Four heights, 10, 100, 3000 and 4500 m were chosen to cover both low- (trades) and high-level (SAL) dust transport.’

Page 26, lines 588:

‘The back-trajectories at 5500 m can be found in the supplements. ‘

Page 27, lines 597-598:

‘Thus, the source area of the samples was most likely the chlorite and kaolinite rich sediments located near the Bou Craa phosphate mine in the Western Sahara (Fig. 9b).’

Page 30, lines 620-625:

‘A further source area might be the southern shoreline of the Western Sahara in which chlorite depleted sediments are situated (Fig. 10d). The presence of the mineral kaolinite in this marine winter sample may be explained by a kaolinite-rich source area lying in the southern Senegal-Mauritania Basin (Fig. 10e).’

Comment of Referee #1:

When you discuss the dust sources, the basic assumption seems to be that there is always one source with one mixture of minerals. But couldn't dust be emitted from areas with differing mineral composition in one dust storm (in fact, even the regions you selected as “major potential source regions” are not homogenous concerning the mineral compositions). My understanding of mineralogy might be too limited, here, but for me the attribution to the characteristics of the sampled dusts to the source regions seemed to include a lot of guessing.

Reply to comment:

Of course, your argument is correct. Individual dust outbreaks may result from multiple dust source areas and the source areas are not homogenous with respect to the mineralogical composition. However, in order to be able to distinguish between the different PSA's we chose typical minerals that are characteristic for the different PSA's as defined by Scheuven et al., 2013.

Comment of Referee #1:

Again, tables could help, showing minerals that are present in the different “major potential source regions”, and at the same time showing the minerals found in the different samples. This would certainly have helped me to follow your conclusions. Also, statistically, the number of dust sample cases that could be evaluated and the number of particles analyzed all are rather low. It is, no doubt, a lot of effort to do all this, but the resulting data should not be overrated. In this context, certainly the last sentence in the “summary and conclusion” (line 843) (and some other sentences as well) is formulated too strongly.

Reply to comment:

Yes, indeed it is not easy to follow how the characteristic minerals of the individual samples were assigned to the PSA's and source rocks and deposits. Therefore, we included a table displaying information on the sampling interval and characteristic minerals of each sample. We further inserted information on the bulk mineralogical composition of the chosen PSA according to Scheuven et al., 2013 and information on the characteristic source rocks and deposits of the chosen source areas. When looking at the table, it should be clearer now how the source areas were chosen based on the characteristic minerals of the samples.

We further agree that the last sentence in the summary and conclusion is formulated too strongly based on the fact that the number of samples is rather low. Therefore, we removed the sentence.

Changes made in manuscript:

We inserted a table:

Page 43, lines 819-824:

‘In Table 7 an overview of the chosen dust source areas for the site Iwik and CBI is given together with the characteristic minerals of the samples that may be used as a tracer for the source area. In the following subsections the identification of the source areas and mineralogical tracers is described in detail.

Table 7: Overview of the chosen source areas and the tracer minerals of the individual samples together with the given characteristics of the source areas according to literature. ‘

Sampling interval	Characteristic minerals of sample	Chosen dust source area	Bulk mineralogical composition of chosen PSA ⁽¹⁶⁾	Characteristic source rocks and deposits of chosen source area
Iwik				
15.02.-15.03.14	*Rut, Serp, Cc	PSA 2: Reguibat Shield	C/K = 0.0–1.0 *Pal: 1-30 wt%	Metamorphic and granitic rocks ⁽¹⁾ Serpentinities ⁽²⁾
15.12.14-18.01.15	*Cc, Chl, Pal (8 wt. %)	PSA 2: Senegal-Mauritania Basin	C/K = 0.0–1.0 *Pal: 1-30 wt%	Chalky horizons ⁽³⁾
		PSA 3: Eastern Taoudeni Basin	C/K = 0.2–0.9 *Pal: 1-5 wt%	Carbonate sequences ⁽⁴⁾
24.06.-15.07.13	*Gib	PSA 2: Tidra Island	C/K = 0.0–1.0 *Pal: 1-30 wt%	Gibbsite maximum offshore Cape Blanc ⁽⁵⁾
15.08.-15.09.14	*Cc, Dol, Pal (5 wt. %)	PSA 2: Aaiun-Tarfaya Basin	C/K = 0.0–1.0 *Pal: 1-30 wt%	Limestone deposits ⁽⁶⁾ Outcrops near Laâyoune with dolomites ⁽⁶⁾
CBi				
26.02.-18.03.14	*Chl, Kao (C/K = 1), Pal (11 wt. %), Flu, Anh, Sme, Ga	PSA 2: Aaiun-Tarfaya Basin near Boucraa	C/K = 0.0–1.0 *Pal: 1-30 wt%	Phosphate deposits ⁽⁷⁾
16.12.14-04.01.15	*Kao (C/K = 0), Pal (1 wt. %), Zeo, Se, Sme	dike swarms and sills of northern Taoudeni Basin	-	Basalts with glass ⁽⁹⁾
		PSA 2: Aaiun-Tarfaya Basin	C/K = 0.0–1.0 *Pal: 1-30 wt%	Palygorskite-sepiolite mafic clays ⁽⁷⁾
		Southern Senegal-Mauritania Basin	-	Lateritic soil ⁽⁸⁾ Horizontal layers of palygorskite and sepiolite ⁽⁸⁾
25.06.-16.07.13	*Chl, Pal (2 wt. %), Rut	PSA 2: Mauritanides	C/K = 0.0–1.0 *Pal: 1-30 wt%	Strongly metamorphosed rocks ⁽¹⁰⁾ Greenschist facies ⁽¹¹⁾
01.08.-21.08.14	*Fe-Amf, Zeo	dike swarms and sills of northern Taoudeni Basin	-	Basalts with glass ⁽⁹⁾
		PSA 4: Fezzan uplift	C/K = 0.0–2.6 *Pal: 0 wt%	Zeolite in basaltic rocks ^(12,13)
		PSA 3: Pharusian belt	C/K = 0.2–0.9 *Pal: 1-5 wt%	Blueschists ⁽¹⁴⁾ Glaucofane bearing eclogites ⁽¹⁵⁾
* Amf = amphibole, Pal = palygorskite, Chl = chlorite, Cc = calcite, Dol = dolomite, Gib = gibbsite, Zeo = zeolite, Kao = kaolinite, Sme = smectite, Se = sepiolite, Rut = rutile, Serp = serpentine, Ga = garnet, Anh = anhydrite, Flu = fluellite				
⁽¹⁾ Schofield et al. (2006) and references therein ⁽²⁾ Schlüter (2008) ⁽³⁾ Wissmann (1982) ⁽⁴⁾ Bertrand-Sarfati et al. (1991) ⁽⁵⁾ Biscaye (1964) ⁽⁶⁾ Bosse and Gwosdz (1996) ⁽⁷⁾ Moreno et al. (2006) ⁽⁸⁾ García-Romero et al. (2007) ⁽⁹⁾ Verati et al. (2005) ⁽¹⁰⁾ Villeneuve (2005), ⁽¹¹⁾ Dallmeyer and Lécorché (2012) ⁽¹²⁾ Abdel-Karim et al. (2013) ⁽¹³⁾ Cvetković et al. (2010) ⁽¹⁴⁾ Caby (2014), ⁽¹⁵⁾ Caby et al., 2008) ⁽¹⁶⁾ Scheuven et al. (2013)				

We removed the last sentence in the summary and conclusions:

Page 49, lines 1020-1022:

‘Finally, we have shown how the mineralogical composition of the samples can be used for provenancing of dust particles found in both on-land and marine dust archives.’

Specific comments:

Comment of Referee #1:

Line 67-69: Is this statement taken from one of the papers cited in the text close by? Which one? And in general: How far offshore can regional / local dust emissions be found? (Or, in other words, how large is the “footprint” of a dust source when it does not emit the dust to large heights?)

Reply to comment:

This is a good point. This sentence was formulated unclearly. It was related to the findings of the cited papers in the sentences before so we modified the sentence. Unfortunately, the exact distance to which regional/local dust emissions can be found offshore is not known. We hope to resolve this question in the future by comparing dust that is collected with three boys offshore.

Changes made in manuscript:

We modified the following sentence:

Page 2, lines 69-71:

‘The results of the above mentioned studies imply that dust collected on land is predominantly of local provenance, while the sources of dust sampled offshore NW Africa are of regional and long-distance provenance.’

Comment of Referee #1:

Line 82: Do you mean your Fig. 1 or the Fig. 1 from Scheuven et al.? Clarify.

Reply to comment:

This was related to the Fig. 1 of this manuscript.

Changes made in manuscript:

We modified the sentence to:

Page 3, lines 90 - 100:

'In Fig. 1 the location of the study sites and the North African dust sources are displayed. The buoy 'Carmen' (~21°15' N, ~20°56' W) and the sediment trap mooring sites CB (~21°16' N, ~20°48' W) and CBi (~20°45' N, ~18°42' W) are located ~ 200 and ~ 80 nautical miles offshore Cape Blanc in the north-eastern (NE) equatorial Atlantic Ocean. The continental dust collector Iwik (~19°53' N, ~16° 18' W) and the meteorological station Arkeiss (~20° 7' N, ~16° 15' W) are located in a major potential dust source area (**PSA 2**) in the Parc National de Banc d'Arguin (PNBA) near Iwik and near Arkeiss in Mauritania. A further meteorological station is positioned in the **PSA 2** in Nouadhibou (~20° 55' N, ~17° 1' W) in the Western Sahara.

The major PSA of northern African dust are summarized in a review by Scheuvens et al. (2013).'

Comment of Referee #1:

Line 383: Arkeiss and Iwik are not oceanic, are they? Rephrase.

Reply to comment:

Yes, you are absolutely right. We removed the word 'oceanic'.

Changes made in manuscript:

We modified the sentence to:

Page 18, lines 420-422:

'Moreover, the TRMM satellite product indicated larger rainfall frequencies during the summer season compared to the winter season regarding the stations Carmen, CBi, Iwik and Arkeiss.'

Comment of Referee #1:

Line 432 - 436: This discussion about dust fluxes is confusing, and I suggest to completely remove it, also because (see next comment):

Reply to comment:

Good point. We decided to remove the detailed comparison in the dust fluxes between the continental and oceanic sites as the sampling height and the sampling method complicates the comparison. We only mention the progressive downwind decrease in the dust fluxes from site Iwik to CBi and CB.

Changes made in manuscript:

We removed the following sentences:

Page 21, lines 471-476:

‘The dust fluxes were about 1000 times smaller at the oceanic sites compared to the continental site. A stronger decrease in the fluxes was observed from site Iwik to CB during summer compared to winter. The variation in the seasonal average dust fluxes were well comparable between the continental and oceanic site CBI. A seasonal trend in the dust fluxes could not be observed for the ocean sites CBI and CB. However, a seasonal trend was observed for site Iwik when taking into account the spring and fall samples.’

Comment of Referee #1:

Line 438: Not surprisingly, the sampling heights has influences the results. This prohibits a direct comparison of mass concentrations and such parameters between the different sampling types (on mast and in traps in the water), while comparing shapes of size distributions and dust compositions might be possible. This should be discussed in the text somewhere. It is mentioned at line 607 again, but also not discussed. Based on this, a comparison of fluxes between the stations makes no sense.

Reply to comment:

Yes, good argumentation. The comparison between the fluxes is complicated as described in the answer to the previous comment. Therefore, we excluded the discussion on the dust fluxes. No trend in the modal particle size was observed with sampling height. Therefore, the particle size distributions were most likely not greatly influenced by the sampling height and a comparison between the size distributions of the sampling sites should be possible.

Changes made in manuscript:

We removed the following sentences:

Page 40, lines 709-714:

‘The stronger decrease in the dust fluxes from site Iwik to CB during summer compared to winter (Table 4) may be explained in the following. During summer, dust was additionally transported with the trades to the site Iwik, (Fig. 9-12) leading to anomalously higher dust deposition at site Iwik compared to the oceanic sites. Further, the washout of dust during offshore transport may have depleted the atmospheric dust cloud resulting in strongly decreased dust deposition fluxes at site CB compared to site CBI during summer.’

We inserted and modified the following sentences in the discussion:

Page 40, lines 703-717:

‘The average horizontal fluxes at site Iwik were ~ 1000 times larger with ~ 100000 mgm⁻²d⁻¹ (Table 5) due to the different sampling technique. The MWAC samplers do not measure deposition fluxes but foremost dust concentrations. Only 1% or less drops out of a moving dust cloud within five minutes, hence, the horizontal dust flux is at least ~100 times higher than the dust deposition flux (Goossens, 2008). The fact that the dust fluxes decreased with height (not shown) further complicated a comparison between the sites due to the different sampling heights of the dust collectors (2.90 m at Iwik versus sediment traps in the water). Therefore, the fluxes between the site Iwik and the offshore sediment trap moorings cannot be compared.

Comment of Referee #1:

Line 450: The bi-modal distribution of Iwik 13-14 is shown in Fig. 6c, right? Add this information to the text.

Reply to comment:

One of this bimodal distribution is shown in Fig. 6c. The two other bimodal distributions were measured for a spring and a summer sample and are therefore not included in Fig. 6.

Changes made in manuscript:

We modified the following sentences:

Page 22, lines 490-493:

‘The three bimodal distributions of the Iwik 13-14 time series were characterized by an additional smaller coarse mode population peaking at ~16 μm besides the more pronounced and variable larger coarse mode population peaking at ~42 to 55 μm. The three Iwik dust samples characterized by a smaller coarse mode population were collected during spring, summer (Fig. 6c) and fall.’

Comment of Referee #1:

Fig. 6: There is a change in sequence between the two captions, which I found disturbing.

I’d prefer to always list - CBi - CB - Iwik. Also: maybe put the black data on top of red one, in the plot, otherwise they are hard to see, or make a mix, so that at least some of the black ones are visible a bit better.


Reply to comment:

The comment helped to improve the figure. We put the black data on top of the red one in the plot to improve the visibility. Also, you are right, the order of sites in the legend was not consistent. However, we suggest to present the site CB first – as it is the most distant one – and the site Iwik last – as this is the most proximal sampling site. Or in other words: we displayed the sites in the legend now from West to East.

Changes made in manuscript:

We modified figure 6:

Page 23, line 518:



Comment of Referee #1:

Line 452: Explain what you mean by “the sorting”.

Reply to comment:

The sorting is expressed by the standard deviation of the grain-size distribution: the larger the standard deviation the weaker the sorting. In order to prevent confusion, we removed the term ‘sorting’ from the text and replaced it with the standard deviation.

Changes made in manuscript:

We modified the following sentences:

Page 19, lines 468-469:

‘The average geometric standard deviation was smaller of the CB samples with 2.6 μm compared to the Iwik and CBI time series with 3.1 μm . The lowest mean/mode ratio was recorded for the CBI time-series with ~ 0.5 (Table 5).’

We removed the following sentence:

Page 22, lines 511-512:

‘In other words: the summer samples of sites CBI and Iwik were less well sorted.’

We modified the following sentence:

Page 22, lines 512-513:

‘This seasonal trend was not observed in the CB 24 upper samples (Table 5).’

Comment of Referee #1:

Figure 7 a) and b): Why was a different wind speed chosen in a) (> 5 m/s) and b) (> 6.5 m/s)? Also: there is a frame around Fig. 7 (on 3 of the 4 sides) that has to be removed for the final version.

Reply to comment:

The explanation on why we chose these values is missing in the text. Only the threshold wind speed of the wind events for which the correlation was best was shown in the graph. This optimum threshold wind speed was different between the correlation to the dust fluxes and the correlation to the modal particle size. We added this information to the revised manuscript.

Changes made in manuscript:

We inserted the following sentences:

Page 24, lines 526-527:


‘The correlation was only evident when using a threshold for wind events of 3.5 to 5.5 ms^{-1} and was best for a threshold of 5 ms^{-1} .’

Page 24, lines 532-533:

‘The correlation was only evident when using a threshold for wind events of 6.5 to 7 ms⁻¹ and was best for a threshold of 6.5 ms⁻¹.’

We modified fig. 7:

Page 24, line 540:



Comment of Referee #1:

Line 527: Why are there 2 back trajectories, only? Were they only made for the time of the dust event? Then why 2 and not only 1? This has to be clearly explained (this also is valid for Fig. 10, 11, and 12, where different numbers or trajectories are shown.)

Reply to comment:

A backward trajectory was drawn only for the day with a dust storm event. So in figure 9 of the old version of the manuscript two days with a dust storm event were identified during the sampling interval of each site resulting in two back trajectories per height at each site. We inserted this information to the text.

Changes made in manuscript:

We inserted the following sentences:

Page 26, lines 583-584:

‘A back trajectory was drawn for the day when a dust storm event occurred as depicted on the satellite images.’

Page 26-27, lines 589-592:

‘During the sampling interval of each site at least two days with dust storms occurred (Fig. 9c,d). Therefore, two back trajectories were drawn for each height for the site CBi and CB respectively. The high-level back trajectories ending at site CBi pass either through the major **PSA 2** or point offshore.’

Page 29-30, lines 607-611:

‘During the sampling interval of the site Iwik at least three dust storms occurred and at the site CBi at least two dust storms occurred (Fig. 10f-h). Each dust storm lasted for several days for which we could model as many as 15 back trajectories for the site Iwik and 8 for the site CBi for each height. The large number of back trajectories complicated the determination of the likely source areas.’

Page 32, lines 632-634:

‘Only one dust storm event was observed during the sampling interval at both sites which lasted for one day (Fig. 11c) resulting in only one back trajectory per site and per height.’

Page 35-36, lines 647-651:

‘At least five separate dust events could be identified (Fig. 12f-j) out of which three occurred during the sampling interval of the site Iwik and two during the sampling interval of the site CBi. One of these dust storms occurring during the sampling interval of site CBi lasted for two days (07-08.08.2014), while all other dust storms lasted for only one day. As a result, three back trajectories could be drawn for each site and each height.’

Comment of Referee #1:

Line 538: Just out of curiosity: does sea spray (i.e., the Cl in it) engrave the analysis of chloride?

Reply to comment:

No, because the CBi samples were pre-treated with distilled water before analysis resulting in the washout of Cl-ions.

Comment of Referee #1:

Figure 9: The line for the 4500m back-trajectory for CBi is in the caption, but not in the plot. But it could be interesting to see it in here, as the dust transported from further away at higher altitudes might also contribute. This comment is valid also for Fig. 10.

Reply to comment:

This was a very good comment which helped to determine the likely source areas of the sampled dust. We added the back trajectories at 3000 and 4500 m to Fig. 9 and 10. With the new Fig. 10 we could find a new source area in Senegal which may have supplied the characteristic minerals kaolinite and sepiolite present in the sample.

Changes made in manuscript:

Page 30, lines 623-625:

‘The presence of the mineral kaolinite in this marine winter sample may be explained by a kaolinite-rich source area lying in the southern Senegal-Mauritania Basin (Fig. 10e).’

Comment of Referee #1:

Line 551: Replace “due to the” by “as there are”.

Reply to comment:

We revised the sentence.

Changes made in manuscript:

We modified the sentence to:

Page 30, lines 615-617:

‘Chlorite may have been supplied to Iwik from a source area nearby the Senegal-Mauritania Basin (Fig. 10a) or as far as the eastern Taoudeni Basin (Fig. 10b) as there are anomalously high chlorite content of the soils in these areas.’

Comment of Referee #1:

Line 596 ff: Was this for PM10 or total particulate mass? What exactly do you mean by “annual average dust concentrations”? The average per dust storm, or really the all-time average of dust, or something else? Please explain. Also: It is not clear to me how the monitoring of less dust events in Morocco due to the shorter sampling time can affect an average value, unless this average is "per year".

Reply to comment:

The annual average dust concentrations were calculated for total particulate mass using all dust samples. We modified the sentence to explain this. We also noticed that the sentence on the monitoring of dust events was not adequate also because the average value was calculated by excluding haze-periods and dust-storms at the sampling site. Therefore, the statement was removed and we modified the paragraph.

Changes made in manuscript:

We modified the following paragraph to:

Page 40, lines 687-692:

‘An annual average dust concentration (total suspended particles) of $\sim 214 \mu\text{g m}^{-3}$ and $275 \mu\text{g m}^{-3}$ was estimated for all dust samples of the year 2013 and 2014 respectively regarding the site Iwik (Table 4). These estimates were larger than what has been measured for background dust concentrations (total suspended particles) in Morocco which were in the order of $100 \mu\text{g m}^{-3}$ during spring 2006 (Kandler et al., 2009). However, in Morocco dust was collected at a larger height of 4 m and haze-periods and dust-storms were excluded from the average value.’

Comment of Referee #1:

Line 606-607: If the difference is due to different sampling techniques, does it make sense to do such a comparison as you present it here?

Reply to comment:

You are right. A comparison of the dust fluxes between the onshore and offshore sites is not feasible because of the different sampling techniques. Therefore, we excluded the comparison of the fluxes between site Iwik and the oceanic sites CB and CBi in the discussion.

Changes made in manuscript:

We modified and shifted the following paragraphs to:

Page 40, lines 699-717:

The observed general decrease in the dust flux from the sites CBi and CB can be explained via the increase in the distance to the source area. Decreased dust deposition fluxes offshore NW Africa with increasing distance from the African coast were also observed by Bory and Newton (2000) analysing the lithogenic fluxes in marine sediment traps.

The average horizontal fluxes at site Iwik were ~ 1000 times larger with $\sim 100000 \text{ mg m}^{-2} \text{ d}^{-1}$ (Table 5) due to the different sampling technique. The MWAC samplers do not measure deposition fluxes but foremost dust concentrations. Only 1% or less drops out of a moving dust cloud within five minutes, hence, the horizontal dust flux is at least ~ 100 times higher than the dust deposition flux (Goossens, 2008). The fact that the dust fluxes decreased with height (not shown) further complicated a comparison between the sites due to the different

sampling heights of the dust collectors (2.90 m at Iwik, versus traps in the water). Therefore, the fluxes between the site Iwik and the offshore sediment trap moorings cannot be compared.

Comment of Referee #1:

Line 608: In which time span does 1% drop out of a moving dust cloud? Per day, per year, ever (certainly not), : : :?

Reply to comment:

In a time span of five minutes. We changed the sentence accordingly.

Changes made in manuscript:

We modified the sentence to:

Page 40, lines 705-706:

‘Only 1% or less drops out of a moving dust cloud within five minutes, hence, the horizontal dust flux is at least ~100 times higher than the dust deposition flux (Goossens, 2008).’

Comment of Referee #1:

Line 613-614: Replace “in the following” by “as follows”.

Reply to comment:

We excluded the sentence according to the comments on the fluxes above.

Changes made in manuscript:

We removed the paragraph:

Page 40, lines 710-714:

‘The stronger decrease in the dust fluxes from site Iwik to CB during summer compared to winter (Table 4) may be explained in the following. During summer, dust was additionally transported with the trades to the site Iwik, (Fig. 9-12) leading to anomalously higher dust deposition at site Iwik compared to the oceanic sites. Further, the washout of dust during offshore transport may have depleted the atmospheric dust cloud resulting in strongly decreased dust deposition fluxes at site CB compared to site CBi during summer.’

Comment of Referee #1:

Line 638: You could stress stronger that a bi-modal size distribution (as observed e.g., at Iwik in Fig. 6c) be indicative for nearby sources. Different size modes typically indicate different sources!

Reply to comment:

Yes, of course bimodal distributions are indicative for different sources. Since this has not been stated clearly enough, we further stressed this in the manuscript. Bimodal distributions may further be indicative of precipitation in the study area due to the deposition of fine particles from higher altitude. This is discussed in the text in chapter 4.1.2.

Changes made in manuscript:

We inserted the following sentence:

Page 41, lines 738-739:

‘Bimodal grain-size distributions typically indicate the sampling of different dust sources (Stuut et al. (2009) and references therein).’

Comment of Referee #1:

Line 641: Not only the wind speed, but also the heights into which dust was emitted, will influence how far it can be transported.

Reply to comment:

Absolutely right. We modified this in the manuscript.

Changes made in manuscript:

We modified the following sentence:

Page 41, lines 742-744:

‘Therefore, it may be possible that wind velocities were high enough during the sampling interval to inject dust to higher altitude and transport it from more distant sources (Fig. 12b) to the sampling site resulting in the small peak in the grain-size distributions.’

Comment of Referee #1:

Line 648: I have a hard time imagining how precipitation droplets (with typically high fall velocities) enter your sampler in noticeable amounts. Or do you suggest here that precipitation is formed, falls, evaporates during its fall and leaves these dust particles behind at lower altitudes which can then be sampled? If this is so, mention it. This is only partly true as there were clearly remnants of water in the bottles. So, both are true!

Reply to comment:

Yes, thanks, it could be a good explanation that precipitation evaporates during its fall leaving the dust particles behind at lower altitude. So we added this to the text. However, we also observed that there were clearly remnants of water in the bottles. So also wet deposition into the bottles may have occurred!

Changes made in manuscript:

We inserted the sentence:

Page 41, lines 753-756:

‘The rain droplets may have evaporated during their fall releasing the dust particles at lower altitudes which can then be sampled with the MWAC sampler. However, we also observed some remnants of water in the bottles and therefore wet deposition into the bottles may have also occurred.’

Comment of Referee #1:

Line 692: Prior to this line, you explain that quartz seems to be present “everywhere” in North Africa. If this is the case, a high quartz content cannot be used to assume that it is mainly locally derived dust.

Reply to comment:

Good point. We removed the sentence.

Changes made in manuscript:

We removed the sentences:

Page 43, lines 809-811:

‘Moreover, the continental sampling site is surrounded by sand dunes which are rich in quartz minerals (Schlüter, 2008; Lancaster, 2013). A high quartz content may therefore point to predominantly locally derived dust.’

Comment of Referee #1:

Line 736: This argument can only be correct if only this one nearby source does not have the mineral palygorskite in it, while it would be present in ALL other sources. Also: not finding palygorskite might simply be an issue of low sampling statistics – can you exclude this? So overall: Can this text in the manuscript really be stated like this? This is also connected to what you state in line 709, so check this location for consistency, too.

Reply to comment:

We revised the sentences in the manuscript in which this was stated incorrectly. Palygorskite may have been derived from several local sources in which it was not present in the soils instead of a single localized source.

Changes made in manuscript:

We modified the sentence:

Page 44, lines 837-838:

‘Therefore, we argue that the sampled dust was most likely derived from a localized source of **PSA 2**.’

Page 45, lines 852:

‘This may point to several externally mixed sources of **PSA 2** during transport.’

Page 45, lines 866-867:

‘Again, the absence of the mineral palygorskite is noteworthy which may point to the sampling of a localized dust source.’

Page 45, lines 879-880:

‘Hence, dust may have been supplied from several dust sources of **PSA 2** which were mixed during transport.’

Comment of Referee #1:

Line 784 ff: How fast do dust particles sink in the water – can they travel 500km and still be collected at the sampling site CBi? Or, in other words, how large is the catchment area of a trap? You mention 40km * 40 km above in the text yourself for the upper trap? Therefore, it seems that the here described time delay cannot be used as an explanation.

Reply to comment:

We argued that certain clay minerals may have been transported with the undercurrent to the sampling site. When not incorporated into marine aggregates clay minerals can easily be transported over long distances within nepheloid layers (ca. 500 km) because of their small size.

Comment of Referee #1:

Line 833-835: For continental sampling, rather than using the word “trade wind”, the word “nearby sources” would be more appropriate. Also, the discrimination between dust particles transported in the SAL or by the trade winds is awkward. The trade winds transport the SAL, while the SAL is the source of particles that deposit from air masses that are moved by the trade winds. In this sense, “trade wind” and “SAL” both contribute together and should not be separated. The formulation needs rewording. Also check the whole text to remove respective inconsistencies.

Reply to comment:

We agree that the sentence should be reformulated.

Changes made in manuscript:

We modified the sentence:

Page 49, lines 1007-1010:

- ‘dust deposited on the continent was predominantly transported from near-by local sources (Mauritania, Western Sahara and Mali), while dust deposited in the marine traps was transported from proximal (Mauritania, Western Sahara and Mali) and distal sources (Algeria and Libya)’

Comment of Referee #1:

Line 836-837: This result was only presented in bypassing (line 731 – or am I missing something), and I was quite surprised to see this as one of the main findings. On the other hand, changing grain sizes with distance to sources are not mentioned at all. I suggest to really reconsider which of your results are worth mentioning here, as for some readers, abstract and summary might be all they will ever look at.

Reply to comment:

Yes, you are right, this was not one of the major findings so we removed it from the summary and conclusions. The downwind decrease in particle size is mentioned in the discussion on page 32, lines 676-682 and pages 33-34 lines 752-756.

Changes made in manuscript:

We removed the sentence:

Page 49, lines 1011-1012:

- 'the percentage of mica relative to the quartz content increased in the deposited dust with increasing transport distance, most likely due to the platy shape of these minerals, which reduces settling'

Comment of Referee #1:

Line 842 ff: The wind at the sampling location might differ from the wind in the source region, if the latter is far away, and the sizes of dust particles present in the source region will influence the grain sizes that can be deposited as well. There might still be a connection between wind speed and transported grain size, but it should be discussed in a broader sense, including (or at least mentioning) the points I raise in the previous sentence.

Reply to comment:

We included these considerations in the summary and outlook of the revised manuscript.

Changes made in manuscript:

We inserted the sentence:

Page 49, lines 1022-1024:

'It should be kept in mind, however, that the wind strength in the sampling location might differ from the wind strength in the source region if the source region is further away. Moreover, the sizes of dust particles present in the source region will influence the grain sizes of deposited dust.'

References

- Abdel-Karim, A.-A. M., Ramadan, E.-N. M., and Embashi, M. R.: Multiphase Alkaline Basalts of Central Al-Haruj Al-Abyad of Libya: Petrological and Geochemical Aspects, *Journal of Geological Research*, 2013, 12, 10.1155/2013/805451, 2013.
- Bertrand-Sarfati, J., Moussine-Pouchkine, A., Affaton, P., Trompette, R., and Bellion, Y.: Cover sequences of the West African craton, in: *The West African orogens and circum-Atlantic correlatives*, Springer, 65-82, 1991.
- Biscaye, P. E.: Mineralogy and sedimentation of the deep-sea sediment fine fraction in the Atlantic Ocean and adjacent seas and oceans, Ph.D., Geology, Yale University, Michigan, 86 pp., 1964.
- Bory, A. J. M., and Newton, P. P.: Transport of airborne lithogenic material down through the water column in two contrasting regions of the eastern subtropical North Atlantic Ocean, *Global Biogeochem. Cycles*, 14, 297-315, 10.1029/1999gb900098, 2000.
- Bosse, H.-R., and Gwosdz, W.: Limestone and dolomite resources of Africa, *Geologisches Jahrbuch Reihe D*, 500, 102, 1996.
- Caby, R.: Nature and evolution of Neoproterozoic ocean-continent transition: Evidence from the passive margin of the West African craton in NE Mali, *Journal of African Earth Sciences*, 91, 1-11, <http://dx.doi.org/10.1016/j.jafrearsci.2013.11.004>, 2014.
- Carlson, T. N., and Prospero, J. M.: The Large-Scale Movement of Saharan Air Outbreaks over the Northern Equatorial Atlantic, *Journal of Applied Meteorology*, 11, 283-297, 1972.
- Cvetković, V., Toljić, M., Ammar, N. A., Rundić, L., and Trish, K. B.: Petrogenesis of the eastern part of the Al Haruj basalts (Libya), *Journal of African Earth Sciences*, 58, 37-50, <http://dx.doi.org/10.1016/j.jafrearsci.2010.01.006>, 2010.
- Dallmeyer, R. D., and Lécorché, J.-P.: *The West African orogens and circum-Atlantic correlatives*, Springer Science & Business Media, 2012.
- Diaz, H. F., Carlson, T. N., Prospero, J. M., and Office, E. R. L. W. M. P.: A Study of the Structure and Dynamics of the Saharan Air Layer Over the Northern Equatorial Atlantic During BOMEX, Nr. 32, Weather Modification Program Office, Environmental Research Laboratories, 1976.
- Einsele, G., Herm, D., and Schwarz, H. U.: Sea level fluctuation during the past 6000 yr at the coast of Mauritania, *Quaternary Research*, 4, 282-289, [http://dx.doi.org/10.1016/0033-5894\(74\)90017-9](http://dx.doi.org/10.1016/0033-5894(74)90017-9), 1974.
- García-Romero, E., Suárez, M., Santarén, J., and Alvarez, A.: Crystallochemical Characterization of the Palygorskite and Sepiolite from the Allou Kagne Deposit, Senegal, *Clays and Clay Minerals*, 55, 606-617, 10.1346/CCMN.2007.0550608, 2007.
- Goossens, D.: Relationships between horizontal transport flux and vertical deposition flux during dry deposition of atmospheric dust particles, *Journal of Geophysical Research: Earth Surface*, 113, n/a-n/a, 10.1029/2007JF000775, 2008.
- Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., Jäckel, S., Jaenicke, R., Knippertz, P., Lieke, K., Massling, A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at Tinfou, Morocco, during SAMUM 2006, *Tellus B*, 61, 32-50, 10.1111/j.1600-0889.2008.00385.x, 2009.
- Lancaster, N.: 11.12 Sand Seas and Dune Fields A2 - Shroder, John F, in: *Treatise on Geomorphology*, Academic Press, San Diego, 219-245, 2013.
- Mittelstaedt, E.: The ocean boundary along the northwest African coast: Circulation and oceanographic properties at the sea surface, *Progress In Oceanography*, 26, 307-355, 10.1016/0079-6611(91)90011-A, 1991.
- Moreno, T., Querol, X., Castillo, S., Alastuey, A., Cuevas, E., Herrmann, L., Mounkaila, M., Elvira, J., and Gibbons, W.: Geochemical variations in aeolian mineral particles from the Sahara-Sahel Dust Corridor, *Chemosphere*, 65, 261-270, 2006.
- National Geospatial-Intelligence Agency: North Atlantic, Baltic Sea, North Sea and Mediterranean Sea, 4 ed., Sailing directions (planning guide), ProStar Publications, 2006.
- Prospero, J. M., and Carlson, T. N.: Radon-222 in North Atlantic Trade Winds . Its Relationship to Dust Transport from Africa, *Science*, 167, 974-&, Doi 10.1126/Science.167.3920.974, 1970.

- Prospero, J. M., and Carlson, T. N.: Vertical and areal distribution of Saharan dust over the western equatorial north Atlantic Ocean, *Journal of Geophysical Research*, 77, 5255-5265, 10.1029/JC077i027p05255, 1972.
- Scheuven, D., Schütz, L., Kandler, K., Ebert, M., and Weinbruch, S.: Bulk composition of northern African dust and its source sediments — A compilation, *Earth-Science Reviews*, 116, 170-194, <http://dx.doi.org/10.1016/j.earscirev.2012.08.005>, 2013.
- Schlüter, T.: *Geological atlas of Africa*, Springer, 2008.
- Schofield, D., Horstwood, M., Pitfield, P., Crowley, Q., Wilkinson, A., and Sidaty, H. C. O.: Timing and kinematics of Eburnean tectonics in the central Reguibat Shield, Mauritania, *Journal of the Geological Society*, 163, 549-560, 2006.
- Skonieczny, C., Bory, A., Bout-Roumzeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Diallo, A., and Ndiaye, T.: A three-year time series of mineral dust deposits on the West African margin: Sedimentological and geochemical signatures and implications for interpretation of marine paleo-dust records, *Earth and Planetary Science Letters*, 364, 145-156, <http://dx.doi.org/10.1016/j.epsl.2012.12.039>, 2013.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 10.1175/bams-d-14-00110.1, 2015.
- Stuut, J.-B. W., Zabel, M., Ratmeyer, V., Helmke, P., Schefuß, E., Lavik, G., and Schneider, R. R.: Provenance of present-day eolian dust collected off NW Africa, *Journal of Geophysical Research*, 110, 10.1029/2004JD005161, 2005.
- Stuut, J.-B. W., Smalley, I., and O'Hara-Dhand, K.: Aeolian dust in Europe: African sources and European deposits, *Quaternary International*, 198, 234-245, 10.1016/j.quaint.2008.10.007, 2009.
- Verati, C., Bertrand, H., and Féraud, G.: The farthest record of the Central Atlantic Magmatic Province into West Africa craton: Precise $^{40}\text{Ar}/^{39}\text{Ar}$ dating and geochemistry of Taoudenni basin intrusives (northern Mali), *Earth and Planetary Science Letters*, 235, 391-407, <http://dx.doi.org/10.1016/j.epsl.2005.04.012>, 2005.
- Villeneuve, M.: Paleozoic basins in West Africa and the Mauritanide thrust belt, *Journal of African Earth Sciences*, 43, 166-195, <http://dx.doi.org/10.1016/j.jafrearsci.2005.07.012>, 2005.
- Wissmann, G.: Stratigraphy and structural features of the continental margin basin of Senegal and Mauritania, in: *Geology of the Northwest African continental margin*, Springer, 160-181, 1982.

Anonymous Referee #2

Received and published: 31 March 2017

In the manuscript, "Seasonal provenance changes of present-day Saharan dust collected on- and offshore Mauritania" the authors present data from sediment traps at multiple depths off the coast of Mauritania and surface collection sites close to the coast. They attempt to determine location and seasonality in potential source regions for different case studies based on the mineralogy of the samples and back trajectory analysis.

Comment of Referee #2:

The description of the methodology and measurements is comprehensive and well thought out. I cannot speak to the specifics of the measurement methodology but have added minor comments and clarification requests below. Unfortunately, the broader context and the scientific developments are lacking in the paper. The measurements are clearly valuable and should be published. The analysis of collected samples and the potential source regions is thorough in a qualitative sense. However, the useful scientific conclusions are not clear. This is indicated by the abstract that reads more like an introduction, the long, subjective discussion, and the relatively sparse summary and conclusions. For example, the last paragraph of the manuscript states that sediment records from land and ocean are likely to sample different source regions, based on the measurements showing more local sources over land. This could be an interesting point, but without further analysis (firmer understanding of the sources, dependence on the particular measurement site) the conclusion that sources are more likely to be local on land than at an ocean site further downwind seems common sense. I'm also not convinced that the atmospheric and sediment trap data should be presented side by side based on the difference in collection methodology and catchment. I think the authors need to consider how to better frame the important measurements presented in this manuscript, by presenting the data in a way that is easier to compare with other dust deposition and concentration measurements and a more thorough back trajectory analysis that answers the questions laid out in the introduction.

Comment of Referee #2:

The choice of questions (lines 76-79) is a little strange. For example, (1) why would one not expect there to be seasonality in the deposition when we know that there is seasonality in winds leading to dust emission and transport? (2) This is an interesting question, but how dependent is this on the specific locations chosen? (3) This is very similar to question (1). (4) This is a good question but is only tackled in a qualitative way in the manuscript. I think laying out the questions to be answered is a good format; however, they currently seem like an afterthought and they should be returned to explicitly in the summary/conclusions.

Reply to comment:

Yes, it is correct that the questions were not formulated and addressed adequately in the old version of the manuscript. Therefore, we modified the research questions. Further, we answered them more precisely in the

discussion, summary and conclusions part of the revised manuscript. Especially the newly developed table 6 which was demanded by reviewer 1 should help to answer the revised research question 2-3.

Changes made in manuscript:

We modified the research questions (page and line numbers refer to the revised manuscript with track changes):

Page 3, lines 81-84:

- 1) What is the seasonal variability in particle size of mineral dust deposited on land? How does the variability relate to meteorological parameters (wind speed, precipitation)?
- 2) What are the source regions of dust trapped on land versus dust trapped in the ocean?
- 3) Can we identify characteristic minerals that constitute a tracer for certain source areas?

We inserted the sentences:

Page 41, lines 756-760:

‘During summer, frequent rainfall resulted in a decrease of the particle sizes of deposited Saharan dust at the site Iwik (Fig. 7c). In addition, the seasonal average percentage of PM₁₀ particles was larger during summer compared to winter at the site Iwik and CBI (Tab. 5). These observations may also be explained by the deposition of relatively finer dust particles from higher altitude of the SAL due to precipitation.’

Page 42, lines 763-769:

‘Both at the onshore sampling site Iwik and at the offshore sampling site CBI a clear seasonal trend in the particle sizes of deposited dust could be observed with generally coarser modal particle sizes during summer (Fig. 6b,c). Generally coarser summer modal particle sizes of deposited dust at site CBI were observed before for a three year time series during 2003 to 2006 and related to moist convective dust storm events (Friese et al., 2016). The generally coarser particle sizes during summer at the site Iwik may be explained by the trade wind speed as a positive correlation between the modal grain sizes and surface wind velocities was observed (Fig. 7a).’

Page 43 lines 776-777:

‘To sum up, the seasonal variability in the particle size of deposited dust at the site Iwik was mainly driven by the surface wind speed due to the predominant sampling of nearby dust sources year-round.’

Page 49, lines 1004-1006:

- ‘A clear seasonal variability in the particle size of mineral dust deposited on land could be observed with generally coarser modal grain sizes during summer compared to winter. The modal particle sizes could be related to the trade wind speed.’

- dust deposited on the continent was predominantly transported from near-by local sources (Mauritania, Western Sahara and Mali), while dust deposited in the marine traps was transported from proximal (Mauritania, Western Sahara and Mali) and distal sources (Senegal and Libya).
- Some rare characteristic minerals (e.g. ferryglaucofane, rutile, serpentine) could be related to local outcrops in NW Africa.

We inserted a new table:

Page 43, lines 819-824:

‘In Table 7 an overview of the chosen dust source areas for the site Iwik and CBi is given together with the characteristic minerals of the samples that may be used as a tracer for the source area. In the following subsections the identification of the source areas and mineralogical tracers is described in detail.

Table 7: Overview of the chosen source areas and the tracer minerals of the individual samples together with the given characteristics of the source areas according to literature. ‘

Sampling interval	Characteristic minerals of sample	Chosen dust source area	Bulk mineralogical composition of chosen PSA ⁽¹⁶⁾	Characteristic source rocks and deposits of chosen source area
Iwik				
15.02.-15.03.14	*Rut, Serp, Cc	PSA 2: Reguibat Shield	C/K = 0.0–1.0 *Pal: 1-30 wt%	Metamorphic and granitic rocks ⁽¹⁾ Serpentinities ⁽²⁾
15.12.14-18.01.15	*Cc, Chl, Pal (8 wt. %)	PSA 2: Senegal-Mauritania Basin	C/K = 0.0–1.0 *Pal: 1-30 wt%	Chalky horizons ⁽³⁾
		PSA 3: Eastern Taoudeni Basin	C/K = 0.2–0.9 *Pal: 1-5 wt%	Carbonate sequences ⁽⁴⁾
24.06.-15.07.13	*Gib	PSA 2: Tidra Island	C/K = 0.0–1.0 *Pal: 1-30 wt%	Gibbsite maximum offshore Cape Blanc ⁽⁵⁾
15.08.-15.09.14	*Cc, Dol, Pal (5 wt. %)	PSA 2: Aaiun-Tarfaya Basin	C/K = 0.0–1.0 *Pal: 1-30 wt%	Limestone deposits ⁽⁶⁾ Outcrops near Laâyoune with dolomites ⁽⁶⁾
CBi				
26.02.-18.03.14	*Chl, Kao (C/K = 1), Pal (11 wt. %), Flu, Anh, Sme, Ga	PSA 2: Aaiun-Tarfaya Basin near Boucraa	C/K = 0.0–1.0 *Pal: 1-30 wt%	Phosphate deposits ⁽⁷⁾
16.12.14-04.01.15	*Kao (C/K = 0), Pal (1 wt. %), Zeo, Se, Sme	dike swarms and sills of northern Taoudeni Basin	-	Basalts with glass ⁽⁹⁾
		PSA 2: Aaiun-Tarfaya Basin	C/K = 0.0–1.0 *Pal: 1-30 wt%	Palygorskite-sepiolite mafic clays ⁽¹⁰⁾
		Southern Senegal-Mauritania Basin	-	Lateritic soil ⁽⁸⁾ Horizontal layers of palygorskite and sepiolite ⁽⁸⁾
25.06.-16.07.13	*Chl, Pal (2 wt. %), Rut	PSA 2: Mauritanides	C/K = 0.0–1.0 *Pal: 1-30 wt%	Strongly metamorphosed rocks ⁽¹⁰⁾ Greenschist facies ⁽¹¹⁾
01.08.-21.08.14	*Fe-Amf, Zeo	dike swarms and sills of northern Taoudeni Basin	-	Basalts with glass ⁽⁹⁾
		PSA 4: Fezzan uplift	C/K = 0.0–2.6 *Pal: 0 wt%	Zeolite in basaltic rocks ^(12,13)
		PSA 3: Pharusian belt	C/K = 0.2–0.9 *Pal: 1-5 wt%	Blueschists ⁽¹⁴⁾ Glaucofane bearing eclogites ⁽¹⁵⁾
* Amf = amphibole, Pal = palygorskite, Chl = chlorite, Cc = calcite, Dol = dolomite, Gib = gibbsite, Zeo = zeolite, Kao = kaolinite, Sme = smectite, Se = sepiolite, Rut = rutile, Serp = serpentine, Ga = garnet, Anh = anhydrite, Flu = fluellite				
⁽¹⁾ Schofield et al. (2006) and references therein ⁽²⁾ Schlüter (2008) ⁽³⁾ Wissmann (1982) ⁽⁴⁾ Bertrand-Sarfati et al. (1991) ⁽⁵⁾ Biscaye (1964) ⁽⁶⁾ Bosse and Gwosdz (1996) ⁽⁷⁾ Moreno et al. (2006) ⁽⁸⁾ García-Romero et al. (2007) ⁽⁹⁾ Verati et al. (2005) ⁽¹⁰⁾ Villeneuve (2005), ⁽¹¹⁾ Dallmeyer and Lécorché (2012) ⁽¹²⁾ Abdel-Karim et al. (2013) ⁽¹³⁾ Cvetković et al. (2010) ⁽¹⁴⁾ Caby (2014), ⁽¹⁵⁾ Caby et al., 2008) ⁽¹⁶⁾ Scheuvs et al. (2013)				

Comment of Referee #2:

While the back trajectory analysis is interesting, it is rudimentary. The choices of back trajectory heights appear to be arbitrary, I did not find a reason for the choice of 4500 m and 10 m is understandable, but those trajectories will be highly uncertain. A larger ensemble of back trajectories from different altitudes and start points would better quantify the likelihood of dust sources and also help represent the uncertainties in back trajectories that pass so close to the surface.





Reply to comment:




This comment was raised by reviewer #1 as well. We have chosen the altitudes for back trajectories on the basis of previous studies to which we compared our data. For example, Skonieczny et al. (2013) used 4500m to study the SAL. Following the reviewer's suggestion, we now also included additional back trajectories, which made the observed patterns much clearer. Back-trajectories of the heights 100 m and 3000 m were added to figures 9-12. This improved the detection of the likely dust source areas and the results and interpretations were adjusted accordingly.




Changes made in manuscript:


We modified fig. 9-12:

[Page 29, line 602:](#)









We modified the following sentence to:

Page 17, lines 380-384:

‘Four-day back trajectories at altitudes of 10 (following Stuu et al. (2005)), 100, 3000, 4500 (following Skonieczny et al. (2013)) and 5500m were calculated ending at the dust collector site Iwik (19°52’ N, 16°17’ W) and at the proximal marine trap site CBi (20°46’, 18°44’ W) using the Hybrid Single Particle Langrangian

Integrated Trajectory (HYSPLIT) model (Stein et al., 2015) and the reanalysis dataset (2.5° spatial resolution) on the NOAA website at <http://ready.arl.noaa.gov>. ‘

We modified the following sentences:

Page 26, lines 582-583:

‘Four heights, 10, 100, 3000 and 4500 m were chosen to cover both low- (trades) and high-level (SAL) dust transport.’

Page 26, lines 588:

‘The back-trajectories at 5500 m can be found in the supplement. ‘

Page 27, lines 597-598:

‘Thus, the source area of the samples was most likely the chlorite and kaolinite rich sediments located near the Bou Craa phosphate mine in the Western Sahara (Fig. 9b).’

Page 30, lines 620-625:

‘A further source area might be the southern shoreline of the Western Sahara in which chlorite depleted sediments are situated (Fig. 10d). The presence of the mineral kaolinite in this marine winter sample may be explained by a kaolinite-rich source area lying in the southern Senegal-Mauritania Basin (Fig. 10e).’

Comment of Referee #2:

Further, including surface wind reanalyses (and other meteorology) in a comprehensive analysis of the mineralogy measurements and back trajectories, a more rigorous statistical analysis of likely source regions could be undertaken. A statistical approach would also provide a framework to analyze future measurements, rather than the case-by-case methodology shown. This would also help reduce the speculative nature of the discussion of sources in Section 4.2.

Reply to comment:

Very good point. We followed your advice and executed an analysis of surface wind reanalysis data (20th century reanalysis V2c dataset). This analysis proved anomalously high surface wind velocities for the respective dust storm events and chosen dust source areas. Therefore, the analysis supported the derivation of the source regions based on mineralogy.

Changes made in manuscript:

We inserted the following sentences:

Page 16, lines 369-371:

‘Maps of six hourly mean surface wind vectors and speed (20th century reanalysis V2c dataset) were provided by the NOAA/OAR/ESRL PSD, (Boulder, Colorado, USA) and downloaded from their website (<http://www.esrl.noaa.gov/psd/>). ‘

We inserted the following paragraph and figure:

Page 38-39, lines 667-678:

‘In Fig. 13a-d the mean wind vectors and speed are presented for chosen dust storm events. The individual dust source areas that were identified using the back trajectory of the day with the dust storm as shown in Fig. 9-12 are further displayed in Fig. 13a-d. As can be clearly seen in the subfigures, the mean wind velocities were anomalously large in the chosen dust source areas which enabled dust emission. During winter, six hourly mean wind velocities were larger than 7 ms^{-1} in the chosen dust source areas (Fig. 13a-b). During summer 2013, six hourly mean wind velocities were larger than 6 ms^{-1} in the chosen dust source area (Fig. 13c). During summer 2014 extremely high mean wind velocities were encountered near the study sites and in the dust source area enabling dust emission and transport from a more distant source to the site Iwik (Fig 13d).




Figure 13: Six hourly composite mean wind vectors and speed at 1000 mb for selected days including a dust storm event during winter (a) – (b) and summer (c) – (d). The dust source area that was identified for the individual dust storm event using the back trajectory of the day with the dust storm is further displayed.’

We inserted the sentence:

Page 41, lines 744-746:

‘This interpretation is further supported by the reanalysis wind vector maps showing anomalously high wind velocities between the site Iwik and the proposed distant source area enabling dust emission and transport of dust particles from the more distant source to the site Iwik (Fig. 13d).’

Comment of Referee #2:

Are the distributions really unimodal in Figure 6, as stated on line 620? I can see the finer mode and sometimes a third mode in there.

Reply to comment:

You are right. Some distributions have a finer mode. However, most of the distributions are unimodal. This may have been formulated unclearly in the old version of the manuscript. Therefore, we modified the sentence.

Changes made in manuscript:

We modified the sentence to:

Page 40, lines 719-720:

‘The measured grain-size distributions for dust trapped at 2.90 m on land at site Iwik and for dust settling in the ocean were predominantly unimodal (Fig. 6).’

Comment of Referee #2:

Coming from an atmospheric modeling perspective, the mode between 1-10um is of great interest and it is a shame that this is not discussed more. From an atmospheric perspective, the value of this work could be increased by presenting more information on the finer dust particles. Models are always in need of aerosol size distribution measurements for evaluation of dust emission, transport and deposition.

Reply to comment:

We also agree that we should present more information on the fine particles. We therefore added the spatial and seasonal variation in the PM10 percentages of the samples to Table 5. We also added an explanation on the PM₁₀ variations in the discussion part in chapter 4.1.2.

Changes made in manuscript:

We modified Table 5:

Page 21-22, line 482:

Series	Year	Winter	Summer	Annual
<i>Average dust fluxes [mg.m⁻².d⁻¹] (dust concentration [µg.m⁻³])</i>				
Iwik 13	2013	10000 (30)	113000 (268)	95000 (214)
CBi 11 upper	2013	106	168	99
CB 24 upper	2013	53	44	45
Iwik14	2014	208000 (603)	55000 (127)	102000 (275)
CBi 11+12 upper	2014	98	20	47
<i>Average modal grain size [µm]</i>				
Iwik 13	2013	44	49	48
CBi 11 upper	2013	27	39	29
CB 24 upper	2013	16	17	16

Iwik 14	2014	45	49	48
CBi 11+12 upper	2014	34	44	33
<i>PM₁₀ [vol. %]</i>				
Iwik 13	2013	13	18	17
CBi 11 upper	2013	31	38	34
CB 24 upper	2013	34	31	34
Iwik 14	2014	13	27	19
CBi 11+12 upper	2014	30	37	31
<i>Average mean/mode ratio [μm]</i>				
Iwik 13	2013	0.7	0.6	0.6
CBi 11 upper	2013	0.5	0.3	0.5
CB 24 upper	2013	0.7	0.8	0.7
Iwik14	2014	0.6	0.4	0.6
CBi 11+12 upper	2014	0.5	0.3	0.5
<i>Average standard deviation [μm]</i>				
Iwik 13	2013	2.8	3.1	3.0
CBi 11 upper	2013	3.0	3.3	3.1
CB 24 upper	2013	2.7	2.6	2.6
Iwik 14	2014	2.8	3.5	3.1
CBi 11+12 upper	2014	3.1	3.3	3.0

We inserted the sentences:

Page 22, lines 497-499:

‘The annual average PM₁₀ percentage is about two times larger at the offshore sampling sites CB and CBi compared to the onshore sampling site Iwik. On average, about one third of the dust mass sampled at the sites CB and CBi is composed of particles smaller than 10 μm .‘

Page 22, lines 507-510:

‘On average, the PM₁₀ percentage was larger for dust sampled during summer compared to winter at the sites Iwik and CBi. An opposite, however less pronounced, seasonal trend could be observed for the site CB with generally larger PM₁₀ percentages during winter compared to summer.’

Page 41, lines 757-760:

‘In addition, the seasonal average percentage of PM₁₀ particles was larger during summer compared to winter at the site Iwik and CBi (Table 5). These observations may also be explained by the deposition of relatively finer dust particles from higher altitude of the SAL during summer due to precipitation.’

Comment of Referee #2:

It would also be useful to have the size distributions presented as dM/dlnD or dV/dlnD to allow for comparison with model simulations and other measurements.


Reply to comment:

Good point as many modellers read and publish in the journal ACP. Therefore, we followed your suggestion and revised the figure 6.

Changes made in manuscript:

We modified figure 6:

Page 23, lines 518:



Comment of Referee #2:

To summarize, I think:

- The measurements and presentation of the results are of great value
- Determining dust sources could benefit greatly from an analysis of surface winds from meteorological reanalyses to accompany the back trajectories.
- The summary and conclusions seems like an afterthought in the current presentation. Consider expanding this to crystallize the findings of the research better.
- The questions to be answered that are set out in the introduction are vague. It is useful to set out the motivational questions for the manuscript in this way, but I think more time should be put into the questions to be addressed and then ensuring that you return to these points in the discussion/summary/conclusions.

Reply to comment:

Since this comment summarizes the above mentioned comments, we have already answered them above.

Minor Comments

Comment of Referee #2:

line 14-15, 34 "Environmental parameters" is non-descriptive. Please revise. Line 34 could be deleted.

Reply to comment:

We revised the sentences to clarify what we mean by environmental parameters. Further, we also agree that line 34 could be deleted.

Changes made in manuscript:

We modified the sentences:

Page 1, lines 13-16:

‘Saharan dust has a crucial influence on the earth climate system and its emission, transport, and deposition are intimately related to e.g. wind speed, precipitation, temperature and vegetation cover. The alteration in the physical and chemical properties of Saharan dust due to environmental changes is often used to reconstruct the climate of the past.’

Comment of Referee #2:

line 95 "In the following," - add comma

Reply to comment:

We revised this sentence.

Changes made in manuscript:

Page 5, lines 116:

‘In the following, the lithology of the geological provinces that underlay the major PSA’s is outlined (Fig. 1).’

Comment of Referee #2:

line 131 - Haboobs are normally defined as the dust storm from evaporatively driven cold pool outflow from convective events, not low-level jets

Reply to comment:

This was formulated unclearly. Of course, haboobs are not low-level jets. The sentence included an enumeration of dust emission mechanism. To prevent confusion, we excluded the 'so called' from the sentence.

Changes made in manuscript:

We revised the sentence to:

Page 6, lines 153-154:

'Dust emission is driven by low level jets, 'haboobs', African easterly waves (AEWs) and high surface winds associated with the Saharan heat low (Knippertz and Todd, 2012).'

Comment of Referee #2:

line 161-163 - this paragraph seems disconnected from the rest of the section

Reply to comment:

This comment is similar to the comment of reviewer # 1 who asked for a paragraph and figure in which all study sites are introduced together. Therefore, we shifted this paragraph to the introductory sentences of the other study sites and modified it accordingly.

Changes made in manuscript:

We shifted and modified the sentences to

page 3, lines 90-97:

'In Fig. 1 the location of the study sites and the North African dust sources are displayed. The dust-collecting buoy 'Carmen' (~21°15' N, ~20°56' W) and the sediment trap mooring site CB (~21°16' N, ~20°48' W) are virtually at the same position ~200 nautical miles offshore Cape Blanc. Sediment-trap station CBi (~20°45' N, ~18°42' W) is located ~ 80 nautical miles offshore Cape Blanc. The continental dust collector Iwik (~19°53' N, ~16° 18' W) and the meteorological station Arkeiss (~20° 7' N, ~16° 15' W) are located in a major potential dust

source area (**PSA 2**) in the Parc National de Banc d'Arguin (PNBA) near Iwik and near Arkeiss in Mauritania. A further meteorological station is positioned in the **PSA 2** in Nouadhibou (~20° 55' N, ~17° 1' W) in the Western Sahara. ‘

Comment of Referee #2:

line 222 and 235 repeat

Reply to comment:

We merged the two sentences.

Changes made in manuscript:

We removed the sentence:

page 13, line 254:

‘The masts were aligned to the ambient wind direction via a wind vane (Fig. 2).’

We modified the sentence:

Page 13, lines 267-268:

‘The masts of the buoy Carmen and of the Iwik dust sampler were aligned to the ambient wind direction via a wind vane (Fig. 2).’

Comment of Referee #2:

line 247 - why is 2xCorg removed from the total mass? Is this a general scaling from organic carbon to organic mass?

Reply to comment:

Yes. About 50-60 % of marine organic matter is constituted by organic carbon. Therefore, we estimated the organic matter by multiplying organic carbon by a factor of two.

Comment of Referee #2:

line 350 - I think the long url links should go in the data availability section rather than in the text. Write out the usage but simply reference the data section rather than talking about downloading files in the manuscript.

Reply to comment:

Good point as this will improve the readability of the manuscript. We modified the section '2.8 Mapping with ArcMap' by removing the long url links.

Changes made in manuscript:

The following sentences were modified to:

Page 17, lines 385-390:

'An ArcGIS layer file of the African surface lithology (new_af_lithology_w_glbcrv_waterbdy_90m_dd84_final.lyr) was downloaded from the U.S. Geological survey (USGS) website (<http://rmgsc.cr.usgs.gov>).

An ArcGIS shape file of the African soils (DSMW.shp) was downloaded from the website of the food and agriculture organization of the United Nations (FAO) (<http://www.fao.org>).'

Comment of Referee #2:

line 357-358 - this repeats the previous sentence, condense.

Reply to comment:

We condensed the two sentences into one sentence.

Changes made in manuscript:

The sentence was modified to:

Page 17, lines 392-394:

'The mean percentage of calcite (8.9 %), chlorite (4.1 %) and kaolinite (29%) in the clay fraction of Saharan soils in general and for each soil type is given by Journet et al. (2014).'

Comment of Referee #2:

line 377 - "ground station" could be misinterpreted as on land, consider "surface station"

Reply to comment:

We changed the word 'ground station' into 'surface station' all through the revised manuscript.

Changes made in manuscript:

Page 5, lines 108-110:

Figure 1: Map of the study sites under investigation: the scientific buoy Carmen as well as the sediment trap moorings CB and CBi offshore Cape Blanc, the MWAC dust collector onshore near Iwik and the surface stations near Nouadhibou and Arkeiss

Page 18, lines 416-417:

‘Regarding the surface stations Carmen and Arkeiss, a threshold of $>0.2 \text{ mmd}^{-1}$ was used in order to exclude events which may be related to anomalously high moisture instead of rainfall.’

Page 18, lines 426-428:

‘However, the spatial and seasonal trends observed by the TRMM data were not supported by the sensor on buoy Carmen and by the surface station in Arkeiss.’

Page 18, lines 429-431:

‘Further, disagreements between the surface stations and the TRMM stations maybe caused by the local signal recorded by the respective rain sensor.’

Page 19, line 438:

‘The wind direction and speed for the surface stations Carmen, Nouadhibou and Iwik are displayed in Fig. 4c.’

Comment of Referee #2:

line 461 - define "well sorted"

Reply to comment:

The same comment raised by reviewer # 1. ‘Well sorted’ refers to the standard deviation of the particle size distribution: the smaller the standard deviation the better the sorting. We removed the term ‘sorting’ from the text in order to prevent confusion and replaced it with the standard deviation.

Changes made in manuscript:

We modified the following sentences:

Page 19, lines 468-469:

‘The average geometric standard deviation was smaller of the CB samples with 2.6 μm compared to the Iwik and CBi time series with 3.1 μm . The lowest mean/mode ratio was recorded for the CBi time-series with ~ 0.5 (Table 5).’

We removed the following sentence:

Page 22, lines 511-512:

‘In other words: the summer samples of sites CBi and Iwik were less well sorted.’

We modified the following sentence:

Page 22, lines 512-513:

‘This seasonal trend was not observed in the CB 24 upper samples (Table 5).’

Comment of Referee #2:

Figure 6 - These are not really unimodal, as referred in the text (line 620)

Reply to comment:

Most of them are unimodal, while some are bimodal. That was formulated unclearly. So we modified the sentence on line 620 of the old version of the manuscript.

Changes made in manuscript:

We modified the following sentence:

Page 40, lines 719-720:

‘The measured grain-size distributions for dust trapped at 2.90 m on land at site Iwik and for dust settling in the ocean were predominantly unimodal (Fig. 6).’

Comment of Referee #2:

line 602 - which year?

Reply to comment:

That was for the year 2013.

Changes made in manuscript:

The sentence was modified to:

Page 40, lines 695:

‘At the distal oceanic site CB, the annual average dust deposition flux was $\sim 45 \text{ mgm}^{-2}\text{d}^{-1}$ (2013) (Table 4).’

Comment of Referee #2:

line 608 - is horizontal flux a useful metric to compare with ocean deposition?

Reply to comment:

We also realized that it is not possible to compare the dust fluxes measured at site Iwik with the dust fluxes of the sites CBi and CB due to the different sampling techniques and sampling heights. Therefore, we modified the paragraph on the dust fluxes.

Changes made in manuscript:

We modified the following paragraph:

Page 40, lines 703-717:

‘The average horizontal fluxes at site Iwik were ~ 1000 times larger with $\sim 100000 \text{ mgm}^{-2}\text{d}^{-1}$ (Table 5) due to the different sampling technique. The MWAC samplers do not measure deposition fluxes but foremost dust concentrations. Only 1% or less drops out of a moving dust cloud within five minutes, hence, the horizontal dust flux is at least ~ 100 times higher than the dust deposition flux (Goossens, 2008). The fact that the dust fluxes decreased with height (not shown) further complicated a comparison between the sites due to the different sampling heights of the dust collectors (2.90 m at Iwik, versus traps in the water). Therefore, the fluxes between the site Iwik and the offshore sediment trap moorings cannot be compared.’

Comment of Referee #2:

A minor issue, but there are formatting errors with brackets on the references throughout that need fixing.

Reply to comment:

We modified the references which were characterized by formatting errors.

Changes made in manuscript:

Page 2, lines 44-45:

‘For instance, the particle size of mineral dust in ocean sediment records varies according to the paleo-frequency of dust-storm and rainfall events (e.g. Friese et al. (2016)).’

Page 2, lines 54-59:

‘The source regions of Saharan dust have been studied frequently by analysing the mineralogical composition of dust collected at continental sites (e.g. Skonieczny et al. (2013); Skonieczny et al. (2011); Schütz and Seibert (1987); Kandler et al. (2009); Khiri et al. (2004)), during aircraft flights (e.g. Formenti et al. (2008)), on research ships (Chester et al., 1971; Chester et al., 1972; Stuut et al., 2005; Aston et al., 1973; Chester and Johnson, 1971b; Chester and Johnson, 1971a) and with gravity cores offshore NW Africa (Biscaye, 1964; Biscaye, 1965; Lange, 1982; Rateev et al., 1969; Griffin et al., 1968; Diester-Haass and Chamley, 1978; Meyer et al., 2013).’

References

- Abdel-Karim, A.-A. M., Ramadan, E.-N. M., and Embashi, M. R.: Multiphase Alkaline Basalts of Central Al-Haruj Al-Abyad of Libya: Petrological and Geochemical Aspects, *Journal of Geological Research*, 2013, 12, 10.1155/2013/805451, 2013.
- Bertrand-Sarfati, J., Moussine-Pouchkine, A., Affaton, P., Trompette, R., and Bellion, Y.: Cover sequences of the West African craton, in: *The West African orogens and circum-Atlantic correlatives*, Springer, 65-82, 1991.
- Biscaye, P. E.: Mineralogy and sedimentation of the deep-sea sediment fine fraction in the Atlantic Ocean and adjacent seas and oceans, Ph.D., Geology, Yale University, Michigan, 86 pp., 1964.
- Bosse, H.-R., and Gwosdz, W.: Limestone and dolomite resources of Africa, *Geologisches Jahrbuch Reihe D*, 500, 102, 1996.
- Caby, R.: Nature and evolution of Neoproterozoic ocean-continent transition: Evidence from the passive margin of the West African craton in NE Mali, *Journal of African Earth Sciences*, 91, 1-11, <http://dx.doi.org/10.1016/j.jafrearsci.2013.11.004>, 2014.
- Cvetković, V., Toljić, M., Ammar, N. A., Rundić, L., and Trish, K. B.: Petrogenesis of the eastern part of the Al Haruj basalts (Libya), *Journal of African Earth Sciences*, 58, 37-50, <http://dx.doi.org/10.1016/j.jafrearsci.2010.01.006>, 2010.
- Dallmeyer, R. D., and Lécorché, J.-P.: *The West African orogens and circum-Atlantic correlatives*, Springer Science & Business Media, 2012.
- Friese, C. A., van der Does, M., Merkel, U., Iversen, M. H., Fischer, G., and Stuut, J.-B. W.: Environmental factors controlling the seasonal variability in particle size distribution of modern Saharan dust deposited off Cape Blanc, *Aeolian Research*, 22, 165-179, <http://dx.doi.org/10.1016/j.aeolia.2016.04.005>, 2016.
- García-Romero, E., Suárez, M., Santarén, J., and Alvarez, A.: Crystallochemical Characterization of the Palygorskite and Sepiolite from the Allou Kagne Deposit, Senegal, *Clays and Clay Minerals*, 55, 606-617, 10.1346/CCMN.2007.0550608, 2007.
- Goossens, D.: Relationships between horizontal transport flux and vertical deposition flux during dry deposition of atmospheric dust particles, *Journal of Geophysical Research: Earth Surface*, 113, n/a-n/a, 10.1029/2007JF000775, 2008.
- Journet, E., Balkanski, Y., and Harrison, S. P.: A new data set of soil mineralogy for dust-cycle modeling, *Atmos. Chem. Phys.*, 14, 3801-3816, 10.5194/acp-14-3801-2014, 2014.
- Knippertz, P., and Todd, M. C.: Mineral dust aerosols over the Sahara: Meteorological controls on emission and transport and implications for modeling, *Rev Geophys*, 50, n/a-n/a, 10.1029/2011RG000362, 2012.
- Moreno, T., Querol, X., Castillo, S., Alastuey, A., Cuevas, E., Herrmann, L., Mounkaila, M., Elvira, J., and Gibbons, W.: Geochemical variations in aeolian mineral particles from the Sahara-Sahel Dust Corridor, *Chemosphere*, 65, 261-270, 2006.
- Scheuvs, D., Schütz, L., Kandler, K., Ebert, M., and Weinbruch, S.: Bulk composition of northern African dust and its source sediments — A compilation, *Earth-Science Reviews*, 116, 170-194, <http://dx.doi.org/10.1016/j.earscirev.2012.08.005>, 2013.
- Schlüter, T.: *Geological atlas of Africa*, Springer, 2008.
- Schofield, D., Horstwood, M., Pitfield, P., Crowley, Q., Wilkinson, A., and Sidaty, H. C. O.: Timing and kinematics of Eburnean tectonics in the central Reguibat Shield, Mauritania, *Journal of the Geological Society*, 163, 549-560, 2006.
- Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Diallo, A., and Ndiaye, T.: A three-year time series of mineral dust deposits on the West African margin: Sedimentological and geochemical signatures and implications for interpretation of marine paleo-dust records, *Earth and Planetary Science Letters*, 364, 145-156, <http://dx.doi.org/10.1016/j.epsl.2012.12.039>, 2013.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 10.1175/bams-d-14-00110.1, 2015.
- Stuut, J.-B. W., Zabel, M., Ratmeyer, V., Helmke, P., Schefuß, E., Lavik, G., and Schneider, R. R.: Provenance of present-day eolian dust collected off NW Africa, *Journal of Geophysical Research*, 110, 10.1029/2004JD005161, 2005.

Verati, C., Bertrand, H., and Féraud, G.: The farthest record of the Central Atlantic Magmatic Province into West Africa craton: Precise $^{40}\text{Ar}/^{39}\text{Ar}$ dating and geochemistry of Taoudenni basin intrusives (northern Mali), *Earth and Planetary Science Letters*, 235, 391-407, <http://dx.doi.org/10.1016/j.epsl.2005.04.012>, 2005.

Villeneuve, M.: Paleozoic basins in West Africa and the Mauritanide thrust belt, *Journal of African Earth Sciences*, 43, 166-195, <http://dx.doi.org/10.1016/j.jafrearsci.2005.07.012>, 2005.

Wissmann, G.: Stratigraphy and structural features of the continental margin basin of Senegal and Mauritania, in: *Geology of the Northwest African continental margin*, Springer, 160-181, 1982.

1 Seasonal provenance changes of present-day Saharan dust 2 collected on- and offshore Mauritania

3

4 Carmen A. Friese¹, ~~Johannes A. Hans~~ van Hateren^{2,*}, Christoph Vogt^{1,3}, Gerhard Fischer¹, Jan-
5 Berend W. Stuut^{1,2}

6 ¹Marum-Center of Marine Environmental Sciences, University of Bremen, Bremen, 28359, Germany

7 ²NIOZ-Royal Netherlands Institute for Sea Research, Department of Ocean Systems, and Utrecht University, 1790
8 AB, Den Burg Texel, Netherlands

9 ³ZEKAM, Crystallography, Geosciences, University of Bremen, 28359, Germany

10 *Now at: Vrije Universiteit Amsterdam, Faculty of Earth Sciences, 1081 HV Amsterdam, the Netherlands

11 *Correspondence to:* Carmen A. Friese (cfriese@marum.de)

12 **Abstract.**

13 Saharan dust has a crucial influence on the earth climate system and its emission, transport, and deposition are
14 intimately related to ~~e.g. wind speed, precipitation, temperature and vegetation cover~~ *environmental parameters*.

15 The alteration in the physical and chemical properties of Saharan dust due to ~~environmental changes~~ *changes in*
16 *environmental parameters* is often used to reconstruct the climate of the past. However, to better interpret possible

17 climate changes the dust source regions need to be known. By analysing the mineralogical composition of
18 transported or deposited dust, potential dust source areas can be inferred. Summer dust transport offshore

19 Northwest Africa occurs in the Saharan air layer (SAL). ~~In continental dust source areas dust is also transported~~
20 ~~in the SAL, however the predominant dust input occurs from nearby dust sources with the low-level trade winds.~~

21 ~~In contrast, dust transport in continental dust source areas occurs predominantly with the trade winds.~~ Hence, the
22 source regions and related mineralogical tracers differ with season and sampling location. To test this, dust

23 collected in traps onshore and in oceanic sediment traps offshore Mauritania during 2013 to 2015 was analysed.
24 Meteorological data, particle-size distributions, back-trajectory and mineralogical analyses were compared to

25 derive the dust provenance and dispersal. For the onshore dust samples, the source regions varied according to the
26 seasonal changes in trade-wind direction. Gibbsite and dolomite indicated a Western Saharan and local source

27 during summer, while chlorite, serpentine and rutile indicated a source in Mauritania and Mali during winter. In
28 contrast, for the samples that were collected offshore, dust sources varied according to the seasonal change in the

29 dust transporting air layer. In summer, dust was transported in the SAL from Mauritania, Mali and Libya as
30 indicated by ferrylglauconite and zeolite. In winter, dust was transported with the Trades from ~~the~~ Western Sahara

31 as indicated by e.g. ~~sepiolite and~~ fluellite.

32

33 **Keywords**

34 Saharan dust, MWAC, sediment trap, mineralogy, particle size, major potential source area, provenance

35 1 Introduction

36 Mineral dust influences global climate through many feedback mechanisms and is in turn influenced by variations
37 in environmental parameters. The emission, transport and deposition of mineral dust reacts sensitively to
38 parameters of climate change like rainfall, wind, temperature and vegetation cover (Knippertz and Stuut, 2014).
39 In turn, the emission, transport and deposition of mineral dust has an impact on the atmospheric energy balance
40 (Haywood and Boucher, 2000), precipitation distribution and amplitude (Yoshioka et al., 2007), sea surface
41 temperatures (Lau and Kim, 2007) as well as the oceanic carbon pump (Martin et al., 1991; Martin, 1990; Jickells
42 et al., 2005; Iversen et al., 2010; Iversen and Robert, 2015; Ploug et al., 2008a). The sensitivity of mineral dust to
43 environmental parameters is used to reconstruct the climate of the past (Rea, 1994; Tjallingii et al., 2008; Mulitza
44 et al., 2010; Diester-Haass and Chamley, 1978; Holz et al., 2007; Stein, 1985). For instance, the particle size of
45 mineral dust in ocean sediment records varies according to the paleo-frequency of dust-storm and rainfall events
46 (e.g., Friese et al. (2016)). Further, the mineralogical composition of mineral dust in sediment core records can be
47 used as a qualitative proxy for the paleo-dust source activity (Scheuvens et al., 2013).

48 Every year, about 2000Mt dust are emitted from source areas around the world, of which 75% are deposited on
49 land and 25% into the oceans (Shao et al., 2011). The Saharan Desert is the world's largest source of mineral
50 aerosols with an annual dust transport of ~180 Mt westwards towards the North Atlantic (Yu et al., 2015). About
51 140Mt is actually deposited into the North Atlantic Ocean (Yu et al., 2015). ~~In addition, about 430 Mt is blown~~
52 ~~from the Sahara towards the equatorial Atlantic (Shao et al., 2011) and therefore, Saharan mineral dust~~ constitutes
53 an essential component of the global climate system. The source regions of Saharan dust have been studied
54 frequently by analysing the mineralogical composition of dust collected at continental sites (e.g. Skonieczny et al.
55 (2013); Skonieczny et al. (2011); Schütz and Sebert (1987); Kandler et al. (2009); Khiri et al. (2004)), during aircraft
56 flights (e.g. Formenti et al. (2008)), on research ships (Chester et al., 1971; Chester et al., 1972; Stuut et al.,
57 2005; Aston et al., 1973; Chester and Johnson, 1971b; Chester and Johnson, 1971a) and with gravity cores offshore
58 NW Africa (Biscaye, 1964; Biscaye, 1965; Lange, 1982; Rateev et al., 1969; Griffin et al., 1968; Diester-Haass and
59 Chamley, 1978; Meyer et al., 2013). Continental dust studies in northern Morocco revealed that dust is produced
60 predominantly locally (Khiri et al., 2004; Kandler et al., 2009). For instance, a high percentage of quartz and
61 feldspar and a low amount of micas in the dust samples was interpreted to represent mostly local dust sources and
62 the availability of calcite sources from proximal coastal dunes in Morocco (Khiri et al., 2004). Further, also in
63 Morocco, dust was sampled in Tinfou at a height of 4 m during the SAMUM 2006 field campaign. These samples
64 were analysed for their physical and chemical properties. The particle size correlated to local surface wind speed
65 suggesting the contribution of local dust (Kandler et al., 2009). In contrast, in coastal Senegal dust is sourced by
66 the Sahel during winter as shown by low illite/kaolinite (I/K) ratios and lower palygorskite contents as opposed to
67 the summer samples which were suggested to be originating from the Sahara (Skonieczny et al., 2013). Further,
68 the I/K ratio in dust sampled on the Cape Verde Islands showed that dust was derived from strongly varying
69 sources: north-western Sahara, central and southern Sahara and the Sahel (Caquineau et al., 2002). ~~The results of~~
70 ~~the above mentioned studies imply that dust collected on land is predominantly of local provenance, while the~~
71 ~~sources of dust sampled offshore NW Africa are of regional and long-distance provenance. Hence, dust collected~~
72 ~~on-land is predominantly of local provenance, while the sources of dust sampled offshore NW Africa are of~~
73 ~~regional and long-distance provenance.~~ As a result, a large seasonal difference can be expected in the composition

74 of the marine climate archives, related to the different dominating transport mechanisms of dust in summer and
75 winter (Friese et al., 2016).

76 To test this, we compared the mineralogical composition, the fluxes, and the particle size of Saharan dust sampled
77 from 2013-2015 in Iwik (Mauritania) in on-land dust traps with Saharan dust sampled from 2013-2015 offshore
78 Cape Blanc (Mauritania) in sub-marine sediment traps and with the scientific dust-collecting buoy 'Carmen'. By
79 comparing the data with meteorological data, back trajectories, the African lithology and satellite images we aim
80 to address the following questions:

- 81 1) What is the seasonal variability in the particle size of mineral dust deposited on land? How does the
82 variability relate to meteorological parameters (wind speed, precipitation)?
- 83 2) What are the source regions of dust trapped on land versus dust trapped in the ocean?
- 84 3) Can we identify characteristic minerals that constitute a tracer for certain source areas?
- 85 4) Is there a seasonal variation in the transport patterns of dust deposited on land?
- 86 2) What are the source regions of dust trapped on land versus dust trapped in the ocean?
- 87 3) Do these source regions vary seasonally?
- 88 4) Can we identify characteristic minerals that constitute a tracer for certain source areas?

89 1.1 Study sites and North African dust sources

90 In Fig. 1 the location of the study sites and North African dust sources are displayed.
91 The dust-collecting buoy 'Carmen' (~21°15' N, ~20°56' W) and the sediment trap mooring sites CB (~21°16' N,
92 ~20°48' W) are virtually at the same position ~ 200 nautical miles offshore Cape Blanc. Sediment-trap station
93 and
94 CBi (~20°45' N, ~18°42' W) is located are located ~ 200 and ~ 80 nautical miles offshore Cape Blanc in the
95 north eastern (NE) equatorial Atlantic ocean (Fig. 1). The continental dust collector Iwik (~ 19°53' N, ~ 16°18' W)
96 and the meteorological station Arkeiss (~ 20°7' N, ~ 16°15' W) are located in a major potential dust source area
97 (PSA 2) in the Parc National de Banc d'Arguin (PNBA) near Iwik and near Arkeiss in Mauritania. A further
98 meteorological station is positioned in the PSA 2 in Nouadhibou (~ 20°55' N, ~ 17°1' W) in the Western Sahara.


99 The major PSA potential source areas (PSA) of northern African dust are summarized in a review by Scheuvens
100 et al. (2013), see Fig. 1). Predominant dust transport towards western Africa and offshore the Atlantic Ocean occurs
101 from the foothills of the Atlas mountains, Western Sahara and Western Mauritania (**PSA 2**), southern Algeria and
102 northern Mali (**PSA 3**) and Western Chad including the Bodélé depression (**PSA 5**) (Scheuvens et al., 2013). In
103 contrast, dust sourced from Tunisia and northern Algeria (**PSA 1**) is transported predominantly to the western
104 Mediterranean and Western Europe (Stuut et al., 2009). Central Libya (**PSA 4**) is the most important region for
105 dust transport to the eastern Mediterranean (Scheuvens et al., 2013).


Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Bold





107
 108 **Figure 1: Map of the study sites under investigation: the scientific buoy Carmen as well as the sediment trap moorings**
 109 **CB and CBi offshore Cape Blanc, the MWAC dust collector onshore near Iwik and the surface station near Nouadhibou**
 110 **and Arkeiss. Location of the sediment trap moorings CB and CBi offshore Cape Blanc and the MWAC dust collector on**
 111 **shore near Iwik (shapefile of the surface lithology and the geological provinces: downloaded from the USGS website**
 112 **<http://rmgsc.cr.usgs.gov/ecosystems/africa.shtml#SL> and**
 113 **<http://certmapper.cr.usgs.gov/geoportal/catalog/main/home.page>, major potential dust source areas: redrawn from**
 114 **Scheuven et al. (2013), ocean currents: redrawn from Mittelstaedt (1991)).**

115 **1.2 Geological characterisation of dust-producing areas**

116 In the following, the lithology of the geological provinces that underlay the major PSA's is outlined (Fig. 1).

117 The PSA 1 is underlain by the eastern Atlas chain and the northern Grand erg/Ahnet and Ghadames Basins. The
 118 outcrops in the Atlas uplift are composed of e.g. limestones, sandstones and evaporites (Piqué, 2001). The thick
 119 strata overlying the northern Ahnet and Ghadames Basin consist of e.g. sandstones and mudstones (Selley, 1997b).

120 The **PSA 2** is underlain by the Reguibat Shield, the Mauritanides and the Senegal-Mauritania, Aaiun-Tarfaya,
121 Tindouf and Taoudeni Basins. The western part of the Reguibat Shield is dominated by granitic rocks, while the
122 eastern part is dominated by metamorphic and granitic rocks (Schofield et al. (2006) and references therein). West
123 of the Reguibat Shield, the Mauritanides consist of a metamorphic belt and ophiolite (Villeneuve, 2005). West of
124 the Taoudeni Basin, the Mauritanides are characterized by granites, quartzites and strongly metamorphosed rocks
125 (Villeneuve, 2005). While the Aaiun-Tarfaya Basin features outcrops with dolomites and limestones, the Senegal-
126 Mauritania Basin is characterized by very few carbonate deposits (Bosse and Gwosdz, 1996). The Tindouf Basin
127 is characterized by mainly sandy deposits (Selley, 1997c, b). The local soils surrounding the dust collector site
128 Iwik are composed of sandy deposits often rich in fossil shells and partly cemented by lime (Einsele et al., 1974).

129 The **PSA 3** is underlain by the western Hoggar and parts of the Ahnet, Taoudeni and Iullemeden Basins. The
130 Pharusian belt located in the western Hoggar is characterized by Eburnean granulites, gneiss, graywackes and
131 magmatic rocks (Boullier, 1991). In the southern Ahnet Basin sandstone strata crop out. On the eastern edge of
132 the Taoudeni Basin outcropping sediments are characterized by conglomerates, sandstones and limestones
133 (Bertrand-Sarfati et al., 1991). The outcrops of the Iullemeden Basin are composed of e.g. sandstones,
134 carbonaceous shale, laterites and massive clays (Kogbe, 1973).

135 The **PSA 4** is underlain by parts of the Fezzan and Nubian uplifts and the Sirte and Murzuk Basins. The eastern
136 Fezzan uplift consists of ocean island basalts (Cvetković et al., 2010; Abdel-Karim et al., 2013), while sediments
137 outcropping in the northern Nubian uplift are composed of e.g. sandstones, limestones and gypsiferous horizons
138 (El Makkrouf, 1988). The southern Sirte Basin is covered by sands, gravel and sand seas (Selley, 1997a). Outcrops
139 of the eastern Murzuk Basin are composed of marine limestones and alluvial sandstones (Selley, 1997b, c).

140 The **PSA 5** is underlain by the Chad Basin. During the Holocene, the Chad Basin was filled with fine-grained
141 particles from the drainage of the Tibesti mountains to the north (Prospero et al., 2002). Hence, the sediments that
142 outcrop in the central Chad Basin are characterized by fluvial and alluvial sediments such as laminated diatomites,
143 pelites and coastal sandridges (Schuster et al., 2009).

144 ~~The continental dust collector Iwik (~19°53' N, ~16° 18' W) is located in **PSA 2** in the Parc National de Banc~~
145 ~~d'Arguin (PNBA) near Iwik in Mauritania (Fig. 1). The local soils surrounding the dust collector are composed of~~
146 ~~sandy deposits often rich in fossil shells and partly cemented by lime (Einsele et al., 1974).~~

147 **1.3 Atmospheric setting**

148 Saharan dust emission, transport and deposition are related to seasonal variations in atmospheric circulation
149 (Knippertz and Todd, 2012). The intertropical convergence zone (ITCZ) shifts meridionally from ~12 °N during
150 boreal winter to ~21 °N during boreal summer resulting in a seasonal change in rainfall and winds over the African
151 continent (Nicholson, 2009).

152 During summer, continental rainfall is most intense and the rain belt is positioned near ~10°N with smaller amounts
153 of rainfall near ~21°N. Dust emission is driven by low level jets, ~~so-called~~ 'haboobs', African easterly waves
154 (AEWs) and high surface winds associated with the Saharan heat low (Knippertz and Todd, 2012). Low-level N
155 trade winds blow and transport dust in coastal Mauritania year-round (National Geospatial-Intelligence Agency,
156 2006). Saharan dust is transported on- and offshore. The offshore transport of Saharan dust particles occurs within

157 the ‘Saharan air layer’ (SAL) at an altitude of about 3 km (Diaz et al., 1976; Carlson and Prospero, 1972; Prospero
158 and Carlson, 1972; Prospero and Carlson, 1970).

159 During winter, dust emission is driven by the break-down of nocturnal low-level jets after sunrise, increased surges
160 in Harmattan winds and microscale dust devils and dust plumes (Knippertz and Todd, 2012; Koch and Renno,
161 2005). Dust is transported within the low-level NE and E trade winds to coastal Mauritania (Dobson, 1781) and
162 also offshore to the sediment-trap mooring sites (Stuut et al., 2005).

163 **1.4 Oceanic setting**

164 The surface-water circulation offshore Cape Blanc is influenced by the southward-flowing Canary Current (CC)
165 and the poleward-flowing coastal counter current or Mauritania Current (Fig. 1). Underneath, the undercurrent is
166 flowing poleward in water depths down to 1000 m (Fig. 1). The undercurrent flows along the continental slope
167 and transports water masses originating from ~5-10 °N to latitudes up to 26 °N. The poleward flowing South
168 Atlantic Central Water (SACW) and the southward flowing North Atlantic Central Water (NACW) are situated
169 below the counter current and meet offshore Cape Blanc (Mittelstaedt, 1991). The study area is positioned in a
170 zone of permanent annual upwelling of sub-surface water masses (Cropper et al., 2014). The NACW and SACW
171 may be upwelled and mixed laterally off Cape Blanc (Meunier et al., 2012). The permanent annual upwelling of
172 nutrient-rich subsurface waters results in high phytoplankton concentrations offshore Cape Blanc (Van Camp et
173 al., 1991). As a result, the surface waters are rich in organic detritus, usually referred to as ‘marine snow’, and
174 faecal pellets which are produced by marine zooplankton (Iversen et al., 2010).

175

176 Individual Saharan dust particles which settle at the ocean surface hardly settle to the deep sea. Instead, fine dust
177 particles can be transferred from the ocean surface to the deep sea by being incorporated into marine snow
178 aggregates and faecal pellets (Ternon et al., 2010). The aggregate formation and ballasting of marine snow
179 aggregates and faecal pellets with marine carbonate and opal as well as with Saharan dust particles results in
180 anomalously high sinking velocities (Iversen and Robert, 2015; Fischer and Karakas, 2009; Iversen and Ploug,
181 2010; Iversen et al., 2010; Ploug et al., 2008b). Dust-loaded particles that sink into the deeper water column are
182 assumed to have a mean settling speed of ~ 240 m d⁻¹ at site CB (Fischer and Karakas, 2009).

183

184 The buoy Carmen (–21°15′ N, –20°56′ W) and the sediment trap mooring sites CB (–21°16′ N, –20°48′ W) and
185 CBi (–20°45′ N, –18°42′ W) are located ~200 and ~80 nautical miles offshore Cape Blanc in the north-eastern
186 (NE) equatorial Atlantic ocean (Fig. 1).

187

188

189

190

191

192

193

194

195

196

197 **2. Material and Methods**

198
 199 In Table 1 an overview of the material and methods employed for each study site is presented. Bulk sediment
 200 samples were obtained at the sites CB and CBI and dust samples at the sites Carmen and Iwik. All samples were
 201 analyzed for particle size and dust flux with the exception of the site Carmen, of which only dust particle size was
 202 not analyzed for dust flux. Only the sites CBI and Iwik were analyzed for mineral assemblages and only the samples
 203 of the site Iwik were used for microscopic investigation. Meteorological sensors were available for the stations
 204 Carmen, Iwik and Arkeiss, while for the site Nouadhibou meteorological data was downloaded online. TRMM
 205 precipitation data was downloaded online for all sites except for the site Nouadhibou.

206
 207 **Table 1: Overview of the material and methods employed at each study site.**


<u>Study site</u>	<u>Lat./long.</u>	<u>Samples</u>	<u>Analysis</u>	<u>Meteorological sensor and data</u>	<u>Downloaded meteorological data</u>
<u>Carmen</u>	<u>~21°15' N, ~20°56' W</u>	<u>1 MWAC sample</u>	<u>particle size</u>	<u>Vaisala WXT520: wind direction + speed, precipitation</u>	<u>TRMM 3B42: precipitation</u>
<u>CB</u>	<u>~21°16' N, ~20°48' W</u>	<u>38 sediment trap samples</u>	<u>lithogenic fluxes, particle size</u>	<u>=</u>	<u>TRMM 3B42: precipitation</u>
<u>CBI</u>	<u>~20°45' N, ~18°42' W</u>	<u>38 sediment trap samples</u>	<u>lithogenic fluxes, particle size, mineral assemblages</u>	<u>=</u>	<u>TRMM 3B42: precipitation, HYSPLIT back trajectories</u>
<u>Iwik</u>	<u>~19°53' N, ~16° 18' W</u>	<u>24 MWAC samples</u>	<u>microscopy, dust fluxes, particle size, mineral assemblages</u>	<u>Davis 6250 Vantage Vue: wind direction + speed</u>	<u>TRMM 3B42: precipitation, HYSPLIT back trajectories</u>
<u>Nouadhibou</u>	<u>20° 55' N, 17° 1' W</u>	<u>=</u>	<u>=</u>	<u>=</u>	<u>Wind direction + speed</u>
<u>Arkeiss</u>	<u>~20° 7' N, ~16° 15' W</u>	<u>=</u>	<u>=</u>	<u>Davis 6250 Vantage Vue: precipitation</u>	<u>TRMM 3B42: precipitation</u>

208
 209 **Formatted: Font: Not Bold**


210 **2.1 Sediment traps**

211
 212 Saharan dust was collected in the ocean using marine sediment traps of the type Kiel (model SMT-234/243) which
 213 are conical with an opening of 0.5 m² (Fig. 2). The principle of particle collection is much the same as described

214 by Van der Does et al. (2016) and Korte et al. (2017). At the top of the opening a honeycomb grid is installed to
215 prevent large swimmers (>1 cm) from entering the trap. The sediment traps were equipped with twenty sample
216 cups which rotated according to a pre-programmed sampling interval (Fischer and Wefer, 1991). The sampling
217 interval was chosen depending on the timing of the ship expeditions.



218



219

220 Figure 2: The marine sediment trap moorings CB and CBi offshore Cape Blanc and the dust masts near Iwik,
 221 Mauritania. On the left, a sketch of the sediment trap mooring (sketch of CB 24 copied from Fischer et al. (2013))
 222 together with a photograph of the trap (downloaded from www.kum-kiel.de) is displayed. On the right, a sketch of the
 223 dust mast together with a photograph of the MWAC sampling bottles is depicted.
 224

225 The sampling intervals were synchronized between the two sites. The intervals ranged from 9.5 days to 21.5 days
 226 (Table 24). Deployment and recovery of the sediment-trap samples was performed during the Research Vessel
 227 Poseidon expeditions POS445 (Fischer et al., 2013), POS464 (Fischer et al., 2014) and POS481 (Fischer et al.,
 228 2015a) (Table 24). The working steps related to the trap deployment and treatment are described in Fischer and
 229 Wefer (1991). In order to prevent outflow of water from the cups during sampling, each sampling cup was filled
 230 with 20 ml of filtered (<0.2 μm) seawater with a salinity of 40 ‰. To produce seawater with a salinity of 40 ‰,
 231 100 g NaCl suprapur was added to 1 l of filtered seawater. Microbial and zooplankton activity was inhibited inside
 232 the trap samples by adding 1 ml of a saturated solution of the biocide HgCl_2 per 100 ml of seawater. After recovery,
 233 swimmers <1 cm were removed from the samples by sieving each sample through a 1 mm mesh. A McLane rotary
 234 liquid splitter was used to split the <1 mm fraction of each sample into five equal aliquots.

235 The samples of two sediment-trap deployments during 2013-2015 of the sediment trap mooring stations CB and
 236 CBi were chosen for grain-size analyses (Table 24). The upper traps sampled at an average water depth of ~1300
 237 m and the lower trap sampled at a water depth of ~3600 m (Table 24). Dust which settles at the ocean surface is
 238 advected by ocean currents during settling in the water column. As a result, particles that settle in an area of ~40

239 x 40 km² in the ocean surface above the traps may be collected in a water depth of ~ 1300 m (Friese et al., 2016).
 240 Two winter and two summer samples were chosen for X-ray Diffraction (XRD) measurements (Table 32).

241 **Table 21:** Specifications of the sediment trap samples collected during 2013-2015 chosen for flux and grain-size analysis.

Trap series	Trap type	Sampling period	Cruise deployment	Cruise recovery	Position	Trap depth [m]	Water depth [m]	No. of samples	Sampling intervals
CBi 11 upper (GeoB 18006-2)	SMT 243	29.01.2013 – 25.03.2014	Pos445	Pos464	20°46.4' N 18°44.4' W	1406	2800	18	17x21d, 1x20d
CBi 12 upper (GeoB 19402-01)	SMT 234 NE	14.02.2014 - 23.02.2015	Pos464	Pos481	20°46.4' N 18°44.5' W	1356	2750	20	1x12.5 d, 19x19.5
CB 24 upper (GeoB 18001-1)	SMT 234 NE	24.01.2013 - 05.02.2014	Pos445	Pos464	21°16.9' N 20°50.6' W	1214	4160	18	1x26 d, 16x21 d, 1x15 d
CB 25 lower (GeoB 19401-1)	SMT 234 NE	07.02.2014 – 21.02.2015	Pos464	Pos481	21°17.8' N 20°47.8' W	3622	4160	20	19x19.5 d, 1x9.5 d

242

243 **Table 32:** Sediment trap and MWAC samples chosen for mineralogical investigation.

Sample	Sampling period	Mast	Bottle	Elevation/water depth [m]	Sampling interval
CBi 11 upper # 8	25.06.-16.07.13	-	-	1406	21d
CBi 12 upper # 2	26.02.-18.03.14	-	-	1356	20d
CBi 12 upper # 10	01.08.-21.08.14	-	-	1356	20d
CBi 12 upper # 17	16.12.-04.01.15	-	-	1356	19d
Iwik 13-7-2-3B	24.06.-15.07.13	2	B	1.90	21d
Iwik 14-8-2-5B	15.08.-15.09.14	2	B	2.90	31d
Iwik 14-12-1-4A	15.12.14-18.01.15	1	A	2.40	34d
Iwik 14-2-2-5B	15.02.-15.03.14	2	B	2.90	28d

244

245

246 2.2 Modified Wilson and Cooke (MWAC) samplers

247 Saharan dust was collected on land near Iwik, Mauritania, with a passive dust sampler consisting of two masts (1
 248 and 2) with two sets of five air sampling bottles each (A and B, Fig. 2). The dust sampling bottles are referred to
 249 as modified Wilson and Cooke (MWAC) samplers (Mendez et al., 2011; Wilson and Cooke, 1980) and consist of
 250 a closed Polyethylene bottle through which the wind can pass via two glass tubes of 8 mm openings. Thus, a big
 251 difference between the traps and the MWAC collectors is the much smaller collection area of the MWAC collectors
 252 with 44 mm². The MWAC dust sampler was chosen because it is one of the most common (Zobeck et al., 2003)

253 and most efficient dust samplers (Goossens and Offer, 2000). The sampling bottles were mounted horizontally at
 254 five different heights. ~~The masts were aligned to the ambient wind direction via a wind vane (Fig. 2).~~

255 The samples collected in 2013-2015 were chosen for subsequent flux and grain-size analyses (Table 43). Saltating
 256 dust particles may be collected in the lower sampling bottles at 90 cm. However, the aim was to analyse dust
 257 transported in suspension to enable a better comparison between the continental and marine sites. Therefore, the
 258 highest sampling bottles attached to the mast at 2.90 m height were used for microscope, flux and grain-size
 259 analysis (Table 32). One series of bottles (series B2) of mast 2 were analysed with the microscope. The other three
 260 replicate samples (bottles A1 and B1 of mast 1, bottles A2 of mast 2) were analysed for flux and grain-size analysis.
 261 Out of the three replicate samples, the sample with the highest mass was chosen for the interpretation of the flux
 262 and grain-size data because this bottle was assumed to have sampled most efficiently. Three samples mounted at
 263 a height of 2.40 m of mast 2 were chosen to test the effect of the chemical pre-treatments that we do to isolate the
 264 terrigenous fraction from marine sediments on the resulting grain-size distributions (Fig. 2). Two winter and two
 265 summer samples that contained enough material were chosen for XRD measurements (Table 32).

266 Furthermore, dust was sampled with a MWAC dust sampler mounted on the mast of buoy Carmen, at about 2 m
 267 above the sea surface (Stuut et al., 2015). ~~The masts of the buoy 'Carmen' and of the Iwik dust sampler were
 268 aligned to the ambient wind direction via a wind vane (Fig. 2). A wind vane was attached to the mast which aligned
 269 the sampler to the ambient wind direction.~~ This MWAC dust sample was also analysed for grain-size distribution.

270 **Table 43:** Specifications of the MWAC samples collected during 2013-2015 chosen for flux and grain-size analysis.

Dust collector series	Trap type	Sampling period	Position	Height [m]	No. of samples	Sampling intervals
Iwik 13	MWAC	27.01.2013 – 20.01.2014	19°53.1' N 16° 17.6' W	2.90	11	19 d, 28 d, 32 d, 29 d, 40 d, 21 d, 31 d, 61 d, 31 d, 31 d, 35 d
Iwik 14	MWAC	20.01.2014 - 18.01.2015	19°53.1' N 16° 17.6' W	2.90	13	26 d, 28 d, 31 d, 30 d, 31 d, 30 d, 31 d, 31 d, 30 d, 32 d, 29 d, 34 d
CB-MWAC	MWAC	23.08.2014 – 16.11.2015	21°15.8' N 20°56.1' W	2.00	1	450 d

271

272 2.3 Microscopy

273 The MWAC samples chosen for microscopic investigation were analysed with a Leica M165 C microscope.
 274 Microscope pictures were taken using a Leica DFC420 camera attached to the microscope. The software Leica
 275 application suite 3.8 was used for taking the pictures.

276

277 2.4 Dust and lithogenic fluxes

278 1/5 splits of the sediment trap samples were analysed for dust fluxes and the bulk components following the method
 279 presented in Fischer and Wefer (1991). The lithogenic flux [$\text{mgm}^{-2}\text{d}^{-1}$] was estimated according to Eq. (1):

280
$$\text{lithogenic material} = \text{dust} = \text{total mass} - \text{carbonate} - \text{opal} - 2 \times \text{Corg} \quad (1)$$

281 Organic carbon was measured after the removal of carbonate with 2N HCl using a CHN-Analyser (HERAEUS).
282 Total carbon was estimated by combustion without pre-treatment. Carbonate was determined according to Eq. (2):

$$283 \quad \text{carbonate} = \text{total carbon} - \text{organic carbon} \quad (2)$$

284 Biogenic opal was determined with a sequential leaching technique (Müller and Schneider, 1993).
285 The MWAC samples chosen for dust flux analyses were weighed on a Mettler-Toledo AT261 Delta Range balance
286 with a precision of 0.0001 g. Mean atmospheric dust concentrations were estimated as Eq. (3):

$$287 \quad DL = \frac{MAR}{(v \cdot A)} * \frac{1}{\eta} \quad (3)$$

288 Where DL is the mean dust concentration [μgm^{-3}], MAR is the mass accumulation rate [μgs^{-1}], v is the mean
289 wind speed per sampling month [ms^{-1}], A is the cross-sectional area of the inlet tube of the MWAC sampler [m^2]
290 and η is the estimated sampling efficiency of MWAC bottles. A sampling efficiency of 90 % was assumed based
291 on an efficiency study of Goossens and Offer (2000). Mean horizontal dust fluxes were calculated according to
292 Eq. (4):

$$293 \quad F_h = \frac{MAR}{A} * \frac{1}{\eta} \quad (4)$$


294 where F_h is the horizontal dust flux [$\text{mgm}^{-2}\text{d}^{-1}$], MAR is the mass accumulation rate [mgd^{-1}], A is the cross-
295 sectional area of the inlet tube of the MWAC sampler [m^2] and η is the estimated sampling efficiency of MWAC
296 bottles.

297 **2.5 Particle size**

298 A 1/25 split of the marine sediment trap samples was analysed for particle size of the terrigenous fraction. The
299 samples were pre-treated before measurement in order to isolate this fraction (see also Filipsson et al. (2011); Friese
300 et al. (2016), Meyer et al. (2013) and Stuu (2001) for methodology) with the following steps: (1) removal of
301 organic matter: Addition of 10 ml of H_2O_2 (35%) to the sediment sample and subsequent boiling until the reaction
302 stops, (2) removal of calcium carbonate: Addition of 10 ml HCl (10%) to the sediment sample and subsequent
303 boiling for exactly 1 minute and (3) removal of biogenic silica: Adding 6 g of NaOH pellets to the sediment sample
304 and subsequent boiling for 10 minutes. Before particle-size analysis, 10 drops of $\text{Na}_4\text{P}_2\text{O}_7 \cdot 10\text{H}_2\text{O}$ were added to
305 each sample to assure the full disaggregation of the particles. The pre-treatment of the MWAC samples differed
306 from the pre-treatment of the sediment trap samples as, obviously, these samples did not contain any biogenic
307 material originating from marine plankton. Further, the disaggregation of particles needed to be kept at minimum
308 to allow for the study of dust transport processes, the so-called ‘minimally dispersed’ aeolian fraction (McTainsh
309 et al., 1997). Therefore, the MWAC samples were solely pre-treated with three drops of $\text{Na}_4\text{P}_2\text{O}_7 \cdot 10\text{H}_2\text{O}$ before
310 analysis. The marine sediment-trap samples as well as the MWAC samples were analysed with the laser particle
311 sizer Beckmann Coulter LS13320 at NIOZ using a Micro Liquid Module (MLM). This instrument allows quick,
312 accurate, and precise data acquisition of large size intervals (Bloemsma et al., 2012). An analytical error of ± 1.26
313 μm ($\pm 4.00\%$) was considered for the measurements (Friese et al., 2016).

314 To investigate the comparability of the MWAC samples with the oceanic sediment-trap samples, the particle-size
315 distribution of the MWAC sample attached to buoy Carmen was compared to the averaged particle-size
316 distributions of the upper and lower trap series at site CB (Fig. 3a). The grain-size distribution of the MWAC

317 sample was comparable to both sediment trap time series even though the sampling time period was different. To
 318 ensure that the pre-treatment steps of the traps did not influence the terrigenous fraction itself, tests were made in
 319 which the on-land MWAC samples were exposed to the same pre-treatment steps as the marine samples (Fig. 3b).
 320 One spring sample has been measured with and without a chemical pre-treatment. Two fall dust samples were
 321 obtained from the same height and mast and sampling interval, however from different bottles (A and B) and were
 322 measured with and without pre-treatment. The figure indicates that a pre-treatment of the Iwik dust samples did
 323 not alter the particle distributions of the samples significantly. Further, the particle-size distribution of dust sampled
 324 with different bottles is comparable.



325
 326 **Figure 3: (a) Grain-size distributions for the station CB: Dust sampled with the MWAC sampler 2 m above sea level,**
 327 **with the upper sediment trap at 1214 mbsl and the lower trap at 3622 mbsl. (b) Grain-size distributions of samples of**
 328 **the Iwik 14 time series which have been pre-treated with HCl, H₂O₂ and NaOH (dotted lines) and without pre-**
 329 **treatment (lines).**

330 2.5 Mineral assemblages

331 Two winter and two summer samples of the MWAC dust collector and the sediment-trap series CBi were chosen
 332 for XRD analysis (Table 43). X-Ray Diffraction pattern analyses were carried out in the laboratory of the research
 333 group Crystallography (University of Bremen, Central Laboratory for Crystallography and Applied Material
 334 Sciences, ZEKAM, Dept. of Geosciences).

335 Due to the small amount of material in the available dust samples (< 100 mg), the preparation for the measurement
 336 was done by pipetting a demi-water-sample mixture on glass slides. A thorough preparation commonly increases
 337 reproducibility of the results, however, the standard deviation given by Moore and Reynolds (1989) of ±5% can
 338 be considered as a general guideline for mineral groups with >20% clay fraction. In addition, the determination of

339 well-crystallized minerals like quartz, calcite or aragonite can be done with better standard deviations (Tucker and
340 Tucker, 1988; Vogt et al., 2002). The X-Ray Diffraction was measured on a Philips X'Pert Pro multipurpose
341 diffractometer equipped with a Cu-tube (k_{α} 1.541, 45 kV, 40 mA), a fixed divergence slit of $1/4^{\circ}$, a secondary Ni-
342 Filter and the X'Celerator detector system. The measurements were carried out as a continuous scan from $3 - 85^{\circ}$
343 2θ , with a calculated step size of $0.016^{\circ} 2\theta$ (calculated time per step was 100 seconds). Mineral identification was
344 accomplished using the Philips software X'Pert HighScore™, which, besides the mineral identification, can give
345 a semi-quantitative value for each identified mineral on the basis of Relative Intensity Ratio (R.I.R.)-values. The
346 R.I.R.-values are calculated as the ratio of the intensity of the most intense reflex of a specific mineral phase to the
347 intensity of the most intense reflex of pure corundum (I/Ic) referring to the “matrix-flushing method” after Chung
348 (1974). Unfortunately R.I.R. values are sparse for clay minerals and long chain organic materials hampered the
349 quantification of our samples.

350 **2.6 Meteorological data**

351 The obtained flux and size data were compared to near-by meteorological data (wind speed, wind direction and
352 precipitation).

353 Wind direction, wind speed and precipitation data with a 20 minute resolution were gathered for the sampling site
354 CB ($21^{\circ}17' N - 21^{\circ}12' N, 20^{\circ}56' W - 20^{\circ}54' W$) during the buoy Carmen deployments from November 2013 to
355 September 2015 with a Vaisala WXT520 meteorology sensor. The size of the dataset was reduced by calculating
356 four hour averages. Moreover, wind direction and wind-speed data with a resolution of five minutes to one hour
357 were gathered during sampling at site Iwik ($19^{\circ}53.1' N, 16^{\circ} 17.6' W$) from January 2013 to January 2015 with a
358 Davis 6250 Vantage Vue meteorology sensor. The size of the dataset was reduced by calculating one-hour
359 averages. Further hourly precipitation data were gathered from the station Arkeiss ($20^{\circ} 7' N, -16^{\circ} 15' W$) from
360 December 2013 to March 2015 with another Davis 6250 Vantage Vue meteorology sensor. Continental hourly
361 wind direction and wind-speed data was acquired for the Nouadhibou meteorological station ($20^{\circ} 55' N, 17^{\circ} 1'$
362 W) online from the Cedar Lake Ventures website (<https://weatherspark.com>).

363 Local daily precipitation data (TRMM 3B42 dataset, 0.25° spatial resolution) were derived from the Giovanni
364 online data system, developed and maintained by the NASA GES DISC (<http://gdata1.sci.gsfc.nasa.gov>). Daily
365 precipitation data were downloaded as area-averages around CBI ($20^{\circ} 58' N - 20^{\circ} 34' N, 18^{\circ} 56' W - 18^{\circ} 32' W$),
366 Iwik ($19^{\circ} 41' N - 20^{\circ} 5' N, 16^{\circ} 29' W - 16^{\circ} 05' W$), CB/Carmen ($21^{\circ} 05' N - 21^{\circ} 29' N, 21^{\circ} 02' W - 20^{\circ} 38' W$) and
367 Arkeiss ($20^{\circ} 19' N - 19^{\circ} 55' N, 16^{\circ} 28' W - 16^{\circ} 04' W$) according to the assumed catchment area of the upper trap
368 ($\sim 40 \times 40 \text{ km}^2$).

369 Maps of six hourly mean surface wind vectors and speed (20th century reanalysis V2c dataset) were provided by
370 the NOAA/OAR/ESRL PSD (Boulder, Colorado, USA) and downloaded from their website
371 (<http://www.esrl.noaa.gov/psd/>).

372 **2.7 Mapping with ArcMap**

373 The mapping software ArcMap version 10.3.1 was used to analyze the source regions of the dust samples
374 investigated for mineralogical composition. A map was created with four-day back-trajectories for days with a
375 dust-storm event as depicted on satellite images. In addition, the African surface lithology was included in the map
376 and soils rich in the minerals calcite, kaolinite and chlorite were marked.

377 Satellite quasi-true colour RGB images (MODIS dataset) were retrieved from the NASA [Worldview website](https://worldview.earthdata.nasa.gov)
378 (<https://worldview.earthdata.nasa.gov>). ~~Ocean Biology Distributed Active Archive Centre (OB.DAAC), Goddard~~
379 ~~Space Flight Centre, Greenbelt MD, on their website (<http://oceancolor.gsfc.nasa.gov>).~~

380 Four-day back trajectories (at altitudes of 10 m (following Stuu et al. (2005)), 100 m, 3000 m, 4500 m (following
381 Skonieczny et al. (2013)) and 5500m) were calculated ending at the dust collector site Iwik (19°52' N, 16°17' W)
382 and at the proximal marine trap site CBi (20°46', 18°44' W) using the Hybrid Single Particle Lagrangian
383 Integrated Trajectory (HYSPLIT) model (Stein et al., 2015) and the reanalysis dataset (2.5° spatial resolution) on
384 the NOAA website (at <http://ready.arl.noaa.gov>).

385 An ArcGIS layer file of the African surface lithology
386 (new_af_lithology_w_glbv_r_waterbdy_90m_dd84_final.lyr) was downloaded from the U.S. Geological survey
387 (USGS) website (<http://rmgsc.cr.usgs.gov/outgoing/ecosystems/AfricaData>).

388 An ArcGIS shape file of the African soils (DSMW.shp) was downloaded from the website of the food and
389 agriculture organization of the United Nations (FAO) at
390 (<http://www.fao.org/geonetwork/srv/en/metadata.show?id=14116>). ~~The mean percentage of calcite,~~
391 ~~chlorite and kaolinite in the clay fraction of Saharan soils in general and for each soil type is given by Journet et~~
392 ~~al. (2014). The mean average percentages of calcite (8.9 %), chlorite (4.1 %) and kaolinite (29 %) in the clay~~
393 ~~fraction of Saharan soils in general and for each soil type is given by are 8.9 %, 4.1 % and 29 %, respectively~~
394 Journet et al. (2014). Soils with larger percentages of calcite, chlorite or kaolinite in the clay fraction than the
395 average percentages were marked in the ArcGIS map.

396
397
398
399
400
401
402
403
404
405
406
407
408
409

Field Code Changed

Formatted: Default Paragraph Font, Font: (Default)
+Body (Calibri), 11 pt, German (Germany)

Formatted: Default Paragraph Font, Font: (Default)
+Body (Calibri), 11 pt, German (Germany)


410 **3. Results**

411 **3.1 Meteorology**

412 In Fig. 4 the meteorological data of the sites Carmen (CB), CBi, Iwik, Arkeiss and Nouadhibou during 2013 to
413 2014 are presented (see Fig 4a for location of the sites). The rainfall frequency is given in Fig. 4b for each site.
414 The number of rainfall events were calculated regarding the TRMM stations for precipitation rates $>1 \text{ mmd}^{-1}$
415 because smaller precipitation amounts which were detected by the satellite may not actually reach the ground.
416 Regarding the surfaceground stations Carmen and Arkeiss, a threshold of $>0.2 \text{ mmd}^{-1}$ was used in order to exclude
417 events which may be related to anomalously high moisture instead of rainfall.

418 According to the TRMM satellite product the annual precipitation frequency was larger on the shoreline (station
419 Arkeiss and Iwik) than offshore (station CBi and Carmen) (Fig. 4b). This may be explained by a decrease in
420 atmospheric water vapor content due to precipitation when the winds move westward. Moreover, the TRMM
421 satellite product indicated larger rainfall frequencies during the summer season compared to the winter season
422 regarding the oceanic-stations Carmen, CBi, Iwik and Arkeiss. Larger summer rainfall frequencies can be
423 explained by the summer northward shift of the ITCZ to $\sim 21^\circ \text{ N}$ resulting in more frequent moist convection and
424 rainfall in the study area.

425 The annual rainfall frequency at the site Arkeiss and the summer rainfall frequencies at the sites Arkeiss and
426 Carmen compare quite well between the sensors and the TRMM observations. However, the spatial and seasonal
427 trends observed by the TRMM data were not supported by the sensor on buoy Carmen and by the surfaceground
428 station in Arkeiss. The larger annual and winter rainfall frequency recorded with the sensor on buoy Carmen may
429 be related to water emission from the ocean surface during time periods with strong surface winds. Further,
430 disagreements between the surfaceground stations and the TRMM stations may be caused by the local signal
431 recorded by the respective rain sensor. A larger number of rain sensors would most likely improve the
432 comparability to the TRMM data.



433


434 Figure 4: Meteorological data (a) map showing the study sites Carmen (CB), CBi, and Iwik, Nouadhibou and Arkeiss
 435 the meteorological station in Nouadhibou (b) precipitation at the study sites Carmen (CB), CBi,
 436 and Iwik and Arkeiss (c) wind direction and speed at the study sites Carmen (CB), Nouadhibou and Iwik,
 437 and at the meteorological station in Nouadhibou.

438 The wind direction and speed for the **surfaceground** stations Carmen, Nouadhibou and Iwik are displayed in Fig.
439 4c. The annual average surface wind velocity was maximum offshore at buoy site Carmen(CB) with $\sim 8 \text{ m/s}^1$.
440 The buoy recorded a larger average wind velocity during winter than during summer, which is consistent with this
441 season being dominated by the Trades. On the shoreline, the average wind velocity was slightly larger during
442 summer than during winter. The predominant annual wind direction was NE at site Carmen and Iwik, while
443 predominant NW winds were recorded for the site Nouadhibou. The wind direction changed from predominant
444 NE during winter to predominant NNE direction during summer at site Carmen. A similar, but less pronounced
445 seasonal trend can be observed for the continental site Iwik. In Nouadhibou, the predominant winter wind direction
446 is NNW switching to a predominant NW wind direction during summer. Obviously, with winds originating from
447 the open ocean, not a lot of dust is anticipated. Therefore, we interpret these wind directions as being very local
448 and caused by the shape of the peninsula of Cape Blanc.

Formatted: Superscript

449 3. 2 Microscope findings of the dust samples from Iwik

450 In Fig. 5 the results of the microscopy investigation of the Iwik 2013 time series are presented. In general, the
451 majority of the particles consisted of angular and moderately spherical quartz grains with a diameter of $\sim 50 \mu\text{m}$
452 (Fig. 5a,b). A small percentage of large platy minerals with a diameter of $\sim 200 \mu\text{m}$ were found in all samples (Fig.
453 5b). Large quartz grains with a diameter of ~ 150 to $200 \mu\text{m}$ were detected in 45 % of the samples. An anomalously
454 high percentage of sub-angular and moderately spherical quartz grains with an average diameter of $\sim 200 \mu\text{m}$ was
455 observed in one summer sample (Fig. 5c). Aggregated grains occurred in all samples. However, the percentage
456 and size of the aggregates as well as the size of the aggregated grains differed from sample to sample. Usually, the
457 size of the aggregated grains was $\sim 50 \mu\text{m}$ (Fig. 5a). Two samples were characterized by aggregates composed of
458 particles with a smaller size of $\sim 20 \mu\text{m}$ (Fig. 5d).



459

460 Figure 5: Microscopic photographs of selected dust samples from the Iwik 2013 time series. (a) Spring dust sample with
 461 a ~ 250 x 150 µm aggregate, (b) spring dust sample with a ~ 200 x 100 µm mica chip, (c) summer dust sample with ~
 462 200 x 200 µm quartz grains, (d) fall dust sample with a ~ 600 x 250 µm aggregate.

463 **3.3 Dust fluxes and size on land and in the ocean**

464 In Table 54 the average dust fluxes are given for the sampling sites Iwik, CBi and CB. The dust concentrations at
 465 site Iwik were determined based on the measured wind speed of the meteorological sensor attached to the sampling
 466 mast. For four samples no wind data were available due to a failure of the instrument. For these samples a wind
 467 velocity was assumed based on the seasonal averages calculated from the available wind data of the meteorology
 468 sensor in Iwik (Fig. 4c). The annual average horizontal dust fluxes at site Iwik were of the same order of magnitude
 469 during 2013 and 2014. ~~The PM₁₀ concentration was calculated in order to enable a comparison to other study sites
 470 where only dust particles smaller than 10 µm were sampled.~~ The annual average dust fluxes decreased from the
 471 on-land site Iwik towards the proximal site CBi and the distal site CB. ~~The dust fluxes were about 1000 times
 472 smaller at the oceanic sites compared to the continental site. A stronger decrease in the fluxes was observed from
 473 site Iwik to CB during summer compared to winter. The variation in the seasonal average dust fluxes were well
 474 comparable between the continental and oceanic site CBi. A seasonal trend in the dust fluxes could not be observed
 475 for the ocean sites CBi and CB. However, a seasonal trend was observed for site Iwik when taking into account
 476 the spring and fall samples.~~ At site Iwik (The average dust concentration was maximum during spring plus winter
 477 2013 and 2014 with 393 µgm⁻³ and 341 µgm⁻³, respectively, and minimum in fall 2013 and 2014 with 48 and 68
 478 µgm⁻³, respectively. The dust fluxes generally decreased with collection height in the mast between 90 and 290 cm
 479 (not shown).

480 **Table 54: Seasonal and annual average dust fluxes and average modal grain size, mean/mode ratio and standard**
 481 **deviation of the grain-size distributions from Iwik 13-14, CBi 11-12 upper and CB 24 upper time-series.**

Series	Year	Winter	Summer	Annual
<i>Average dust fluxes [mg·m⁻²·d⁻¹] (dust concentration [µg·m⁻³])</i>				
Iwik 13	2013	10000 (30)	113000 (268)	95000 (214)
CBi 11 upper	2013	106	168	99
CB 24 upper	2013	53	44	45
Iwik 14	2014	208000 (603)	55000 (127)	102000 (275)
CBi 11+12 upper	2014	98	20	47
<i>Average modal grain size [µm]</i>				
Iwik 13	2013	44	49	48
CBi 11 upper	2013	27	39	29
CB 24 upper	2013	16	17	16
Iwik 14	2014	45	49	48
CBi 11+12 upper	2014	34	44	33
<i>PM₁₀ [vol. %]</i>				
Iwik 13	2013	13	18	17
CBi 11 upper	2013	31	38	34
CB 24 upper	2013	34	31	34
Iwik 14	2014	13	27	19
CBi 11+12 upper	2014	30	37	31
<i>Average mean/mode ratio [µm]</i>				
Iwik 13	2013	0.7	0.6	0.6
CBi 11 upper	2013	0.5	0.3	0.5
CB 24 upper	2013	0.7	0.8	0.7
Iwik 14	2014	0.6	0.4	0.6
CBi 11+12 upper	2014	0.5	0.3	0.5

Formatted: Font: Italic

Formatted: Centered

Formatted: Font: Italic, Subscript

Average standard deviation [μm]				
Iwik 13	2013	2.8	3.1	3.0
CBi 11 upper	2013	3.0	3.3	3.1
CB 24 upper	2013	2.7	2.6	2.6
Iwik 14	2014	2.8	3.5	3.1
CBi 11+12 upper	2014	3.1	3.3	3.0

482

483 The statistical values of the measured grain-size distributions for the stations CB, CBi and Iwik are given in Table
 484 [54](#). In addition, the measured grain-size distributions for the time series of the stations CB, CBi and Iwik are
 485 displayed in Fig. 6. In Fig. 6a the average grain-size distribution for the samples of each of the three stations for
 486 the year 2013 are given. The maximum measured particle size decreased from $\sim 223 \mu\text{m}$ on land at site Iwik to
 487 $\sim 169 \mu\text{m}$ at the proximal site CBi and $\sim 140 \mu\text{m}$ at the distal site CB (Fig. 6a). In addition, the average modal grain
 488 size decreased from $\sim 48 \mu\text{m}$ at site Iwik to $16 \mu\text{m}$ at site CB (Table [54](#)). Bimodal grain-size distributions were
 489 encountered for 23 % of the CBi 11-12 samples, 13 % of the Iwik 13-14 samples, and none of the CB 24 samples.
 490 The ~~three~~ bimodal distributions of the Iwik 13-14 time series were characterized by an additional ~~smaller coarse~~
 491 ~~fine~~ mode ~~population~~ peaking at $\sim 16 \mu\text{m}$ besides the more pronounced and variable ~~larger~~ coarse mode ~~population~~
 492 peaking at ~ 42 to $55 \mu\text{m}$. ~~The three~~ ~~Two of the~~ Iwik dust samples characterized by a ~~smaller coarse mode~~
 493 ~~population~~ ~~fine grain size peak~~ were collected during ~~spring, summer (Fig. 6c) and fall–summer and fall~~
 494 ~~respectively~~. ~~The eight~~ bimodal grain-size distributions of the CBi 11-12 time-series were characterized by a
 495 ~~variable~~ larger coarse mode population at ~ 25 to $35 \mu\text{m}$ and a variable smaller coarse mode population at ~ 6 to 16
 496 ~~μm~~ . The bimodal distributions were recorded for three winter, three summer (Fig. 6b,c), one spring and one fall
 497 sample. The annual average PM_{10} percentage is about two times larger at the offshore sampling sites CB and CBi
 498 compared to the onshore sampling site Iwik. On average, about one third of the dust mass sampled at the sites CB
 499 and CBi is composed of particles smaller than $10 \mu\text{m}$. The ~~average geometric standard deviation was smaller~~
 500 ~~of the CB samples~~ ~~–was better than the sorting with $2.6 \mu\text{m}$ compared to the of the Iwik and CBi time series as~~
 501 ~~indicated by the average geometric standard deviations of $2.6 \mu\text{m}$ for CB and with $3.1 \mu\text{m}$ for both Iwik and CBi~~
 502 (Table [54](#)). The lowest average mean/mode ratio was recorded for the CBi time-series with ~ 0.5 ~~due to the weak~~
 503 ~~sorting of the samples~~ (Table [54](#)).

Formatted: Space Before: 12 pt

504 In Fig. 6b-c the measured grain-size distributions for winter and summer samples are displayed. The averaged
 505 modal grain size for the summer samples was coarser grained compared to the winter samples of the respective
 506 grain-size time series (Table [54](#)). The seasonality in modal grain size was largest for the CBi 11 upper trap series
 507 of the year 2013 with a difference of $\sim 12 \mu\text{m}$ (Table [54](#)). ~~On average, the PM_{10} percentage was larger for dust~~
 508 ~~sampled during summer compared to winter at the sites Iwik and CBi. An opposite, however less pronounced,~~
 509 ~~seasonal trend could be observed for the site CB with generally larger PM_{10} percentages during winter compared~~
 510 ~~to summer~~. The average standard deviation was larger and the average mean/mode ratio was smaller in the summer
 511 samples compared to the winter samples regarding the sites Iwik and CBi (Table [54](#)). ~~In other words: the summer~~
 512 ~~samples of sites CBi and Iwik were less well sorted (Fig. 6b and c).~~ This seasonal trend was not observed in the
 513 CB 24 upper samples ~~ples which were generally well sorted~~. (Table [54](#)).


Formatted: Left, Line spacing: Multiple 1,08 li

Formatted: Left, Line spacing: Multiple 1,08 li, Tab stops: 10,32 cm, Left

514

515

516



517

518

519 **Figure 6: Grain-size distributions of the stations Iwik, CBI and CB (a) averaged for the samples of the year 2013 (b)**
 520 **winter samples (c) summer samples.**

521 In Fig. 7a –c the results of the correlation between the characteristics of the dust sampled on land and the local
 522 meteorological data are presented. In Fig. 7a the particle sizes were correlated to the surface wind speed data (N =
 523 13 samples). A correlation above a coefficient of determination (R^2) of 0.3 was considered significant at the 95 %
 524 confidence level for two-tailed probabilities. The modal particle size of the Iwik samples showed a positive linear
 525 correlation with the daily wind speed events with $R^2 = 0.5$, which is significant at the 99.31 % confidence level.

526 The correlation was only evident when using a threshold for wind events of 3.5 to 5.5 ms⁻¹ and was best for a
 527 threshold of 5 ms⁻¹. A better positive linear correlation was obtained when excluding the spring sample resulting
 528 in R² = 0.7 which is significant at the 99.96 % confidence level.

Formatted: Superscript


Formatted: Superscript

529 In Fig. 7b the dust fluxes were correlated to the surface wind-speed data (N = 10 samples). A correlation above R²
 530 = 0.4 was considered significant at the 95 % confidence level for two-tailed probabilities. The horizontal dust flux
 531 of the Iwik samples correlated positively to the daily wind speed events during the sampling interval with R² = 0.7
 532 which is significant at the 99.75 % confidence level. The correlation was only evident when using a threshold for
 533 wind events of 6.5 to 7 ms⁻¹ and was best for a threshold of 6.5 ms⁻¹. Moreover, a significant linear correlation
 534 with R² = 0.6 was observed at the 99.15 % confidence level between the dust fluxes and the mean wind strengths
 535 during the sampling intervals (not shown).

Formatted: Superscript

Formatted: Superscript

536 In Fig. 7c the particle size of the Iwik summer samples was correlated to the local TRMM precipitation data (N = 6
 537 samples). In this case a correlation above R² = 0.7 was considered significant at the 95 % confidence level for two-
 538 tailed probabilities. A good linear negative correlation with R² = 0.9 was observed which is significant at the 99.78
 539 % confidence level.



540
 541 **Figure 7: Correlation between the observed local surface wind speed at site Iwik and the measured (a) modal grain size**
 542 **and (b) flux. (c) Correlation between the observed local precipitation at site Iwik (TRMM data) and the modal grain**
 543 **size of the summer samples.**

544 3.4 Mineral assemblage of dust sampled on land and in the ocean

545 In Table 65 the mineralogical composition averaged over all eight samples, averaged over the four Iwik samples
 546 and the four CBI samples is given. All dust samples contained the minerals quartz and mica. Further minerals that
 547 occurred with significant quantities but which were not present in all dust samples were feldspar, amphibole,
 548 zeolite, chlorite and palygorskite. Calcite, dolomite, gibbsite, kaolinite, smectite, sepiolite, fluellite, anhydrite,
 549 rutile and serpentine occurred only in some samples resulting in a low average abundance ≤ 1%. However, we
 550 argue that these minerals can be used as dust source indicators because of (1) the characteristic distribution of
 551 gibbsite, kaolinite, smectite and sepiolite in North Africa according to different weathering regimes (Biscaye,
 552 1964) and (2) the characteristic occurrence of fluellite, anhydrite, rutile and serpentine according to outcropping
 553 rock type (Deer et al., 1992). Further minerals that occur in low abundances (≤ 3%) were summarized as ‘other

554 minerals' and will not be discussed in the manuscript. While the continental samples were dominated by quartz
 555 and feldspar, the marine samples were dominated by mica, followed by quartz and feldspar.

556 **Table 65: Results of the mineralogical investigation: Mineral assemblage averaged over all samples (Total), the Iwik**
 557 **samples (Iwik) and the CBi samples (CBi).**


*	Qz [%]	Fsp [%]	Mi [%]	Amf [%]	Pal [%]	Chl [%]	Cc [%]	Dol [%]	Gib [%]	Zeo [%]	Kao [%]	Sme [%]	Se [%]	Rut [%]	Serp [%]	Ga [%]	Anh [%]	Flu [%]
Total	25.1	21.5	25.5	5.1	3.4	4.4	0.6	0.1	1.0	3.8	0.9	0.4	1.1	0.5	0.3	0.1	0.1	1.1
Iwik	33.3	30.8	18.0	5.0	3.3	1.8	1.3	0.3	2.0	0.0	0.0	0.0	0.0	0.8	0.5	0.0	0.0	0.0
CBi	17.0	12.3	33.0	5.3	3.5	7.0	0.0	0.0	0.0	7.5	1.8	0.8	2.3	0.3	0.0	0.3	0.3	2.3

*Qz = quartz, Fsp = feldspar, Mi = mica, Amf = amphibole, Pal = palygorsite, Chl = chlorite, Cc = calcite, Dol = dolomite, Gib = gibbsite, Zeo = zeolite, Kao = kaolinite, Sme = smectite, Se = sepiolite, Rut = rutile, Serp = serpentine, Ga = garnet, Anh = anhydrite, Flu = fluellite

558

559 In Fig. 8a-c the results of the mineralogical investigation of the eight chosen dust samples are presented. Figure 8a
 560 depicts again the average composition of the samples per sampling site (N=4). The minerals zeolite, anhydrite,
 561 garnet, sepiolite, fluellite, kaolinite and smectite were only found in the marine samples. Only the continental
 562 sample of 15.08.-15.09.14 contained traces of zeolite. While gibbsite, serpentine, calcite and dolomite were
 563 detected in the continental dust samples, these minerals were absent in all marine samples. The absence of calcite
 564 and gibbsite may have been caused by the pre-treatment of the marine sediment-trap samples with HCl. Although
 565 the concentration of the used acid is fairly low (10%) and the exposure time of the samples was exactly 1 minute,
 566 we cannot exclude that carbonate minerals were dissolved. Therefore, the absence of these minerals in the marine
 567 traps will not be discussed further.

568 In the following, the seasonality in the average mineralogical composition will be outlined for each site as given
 569 in Fig. 8b,c. At site Iwik, the winter dust samples were characterized by the occurrence of chlorite, serpentine and
 570 rutile, while the summer samples were characterized by the minerals gibbsite and dolomite. At site CBi, the winter
 571 dust samples were characterized by the occurrence of the minerals sepiolite, fluellite, kaolinite, smectite, garnet
 572 and anhydrite, while the summer samples were characterized by the mineral rutile. Only for the marine trap
 573 samples an annual average chlorite/kaolinite ratio (C/K = 4) could be derived owing to the occurrence of kaolinite.



574

575 **Figure 8: Mineralogical composition (a) averaged over all samples and for sites Iwik and CBI, (b) averaged for the**
 576 **winter samples at sites Iwik and CBI and for each individual winter sample and (c) averaged for the summer samples**
 577 **at sites Iwik and CBI and for each individual summer sample. The category ‘other minerals’ comprises the minerals**
 578 **todorokite, sodalite, konicklite, gyanaitite, nitratine, urea, bernalite, akermanite, mixed-layer clay and talc.**

579


580 **3.5 Identification of dust source regions using ArcMap**

581 In Fig. 9-12 the results of the four day back-trajectory analysis are presented for each sample which has been
 582 analyzed for mineralogical composition. Four-Two heights, 10 m, 100 m, 3000 m and 4500 m (according to) and
 583 4500 m (according to) were chosen to cover both low- (trades) and high-level (SAL) dust transport. A back
 584 trajectory was drawn for the day when a dust storm event occurred as depicted on satellite images. Only the low-
 585 level back-trajectories were plotted for site Iwik because of the correlation of the measured dust characteristics to
 586 the low-level wind speed. Moreover, the MWAC samplers were designed to only sample dry deposition, whereas
 587 the marine sampling sites collect material settling through the water column, i.e., dust resulting from both dry- and
 588 wet deposition. The back-trajectories at 3000 and 5500 m can be found in the supplements.

589 Figure 9 illustrates a typical late-winter situation. During the sampling interval of each site at least two days with
 590 dust storms occurred (Fig. 9c,d). Therefore, two back trajectories were drawn for each height for the site CBI and

591 Iwik respectively. The high-level back trajectories ending at site CBi pass either through the major PSA 2 or point
592 offshore. Both the low-level back-trajectories ending at the continental trap site Iwik and at the oceanic trap site
593 CBi point to a dust source within the major PSA 2 (Scheuvens et al., 2013). Some calcite was present in the
594 continental dust sample, but no chlorite nor kaolinite was detected. Therefore, the dust source was most likely
595 located in the nearby southwestern Reguibat Shield where sediments are rich in calcite and quartz and depleted in
596 chlorite and kaolinite (Fig. 9a). Dust deposited in the marine traps during the time interval was characterized by
597 the occurrence of chlorite and kaolinite. Thus, the source area of the samples was most likely the chlorite and
598 kaolinite rich sediments located near the Bou Craa phosphate mine in the Western Sahara (Fig. 9b). A small area
599 of chlorite-rich sediments is located in the shoreline of the Western Sahara (Fig. 9b). However, kaolinite is not
600 present in anomalously high amounts in the proposed source area (Fig. 9b).

Formatted: Font: Bold



Lithology

- | | | |
|-----------------------------|------------------------|--|
| Carbonate | Ultramafic | Alluvium - beach, strand, coastal dune |
| Karst | Extrusive Volcanic | Alluvium - Saline |
| Non-Carbonate | Colluvium | Alluvium - Gypsum |
| Metasedimentary | Hydric - Organic | Alluvium - Other |
| Alkaline Intrusive Volcanic | Aeolian Sediments | Volcanic - Ash, Tuff, Mudflow |
| Silicic | Alluvium - Fan Deposit | Water |
| Metigneous | Alluvium - Fluvial | |

Major potential source areas (PSAs)

- PSA 1-5

Study sites and back trajectories



- 10 m 4500 m 10 m


Clay Mineralogy

- No data Calcite >8.9 % Kaolinite >29 % Chlorite >4.1 %

Mineralogy

- | | | | | |
|-----------|--------------|------------|-----------|-----------------|
| Kaolinite | Feldspar | Serpentine | Sepiolite | Gibbsite |
| Chlorite | Mica | Rutile | Anhydrite | Dolomite |
| Calcite | Amphibole | Zeolite | Smectite | Other minerals |
| Quartz | Palygorskite | Fluellite | Garnet | Ferryglaucofane |






602

603 Figure 9: Low-level (10 m) four-day back trajectories of dust events ending during the sampling interval 15.02.-15.03.14
 604 at site Iwik and during the sampling interval 26.02.-18.03.14 at site CBI. The potential dust source areas and the
 605 mineralogy of the samples are given in the subfigures a-b. The dust-storm events occurring during the sampling interval
 606 are indicated in subfigures c-d.


607 Figure 10 represents a typical early-winter situation. During the sampling interval of the site Iwik at least three
 608 dust storms occurred and at the site CBI at least two dust storms occurred (Fig. 10f-h). Each of the dust storms lasted

609 ~~for -and one lasted for~~ several days ~~for which we could model as many as 15 back trajectories for the site Iwik and~~
610 ~~8 for the site CBI for each height. The large number of back trajectories which~~ complicated the determination of
611 the likely dust source areas. All back trajectories pass through the major **PSA 2** and some point to the **PSA 1** and
612 **PSA 3** (Scheuvens et al., 2013). One high-level back trajectory ending at site CBI passes through PSA 2 and two
613 through Mauritania and Senegal. However, most of the high-level back trajectories ending at site CBI point
614 offshore. Dust sampled in the marine traps during this sampling interval did not contain any chlorite, while the
615 dust trapped at Iwik did. Chlorite may have been supplied to Iwik from a source area nearby the Senegal-
616 Mauritania Basin (Fig. 10a) or as far as the eastern Taoudeni Basin (Fig. 10b) ~~as there are due~~ to the anomalously
617 high chlorite content of the soils in these areas. The continental sample is further characterized by the occurrence
618 of calcite and the absence of kaolinite which fits to the soils of the chosen source areas (Fig. 10a,b). The marine
619 sample was characterized by the occurrence of zeolite and absence of chlorite. Therefore, zeolite may have been
620 derived from the extrusive volcanic rocks of the northern Taoudeni Basin (Fig. 10c). A further source area might
621 be the southern shoreline of Western Sahara ~~similar to what was observed for the sample obtained during winter in~~
622 ~~which chlorite depleted sediments are situated 2014 (Fig. 9b, Fig. 10d). Again, the marine winter sample contained~~
623 ~~the mineral kaolinite which cannot be explained with the back trajectories and the soil map. The presence of the~~
624 ~~mineral kaolinite in this marine winter sample may be explained by a kaolinite-rich source area lying in the~~
625 ~~southern Senegal-Mauritania Basin (Fig. 10e).~~

Formatted: Font: Bold



626




627

628 Figure 10: Low-level (10 m) four-day back trajectories of dust events ending during the sampling interval 15.12.14-
 629 18.01.15 at site Iwik and during the sampling interval 16.12.14-04.01.15 at site CBi. The potential source areas and the
 630 mineralogy of the samples are given in the subfigures a-c. The dust storm events occurring during the sampling interval
 631 are indicated in subfigures e-g.

632 In Fig. 11 a typical early-summer situation is presented. Only one dust storm event was observed during the
 633 sampling interval at both sites which lasted for one day (Fig. 11c) resulting in only one back trajectory per site and

634 per height. The low-level back trajectory~~s~~ ending at site CBi runs offshore and the high-level back trajectory
635 passes through the major PSA 2, 3 and 5. The low-level back trajectory ending at site Iwik passes through the
636 major PSA 2, ~~and the high-level back trajectory passes through the major PSA 2 and 3~~ (Scheuven et al., 2013).
637 Dust sampled on land at site Iwik was characterized by the absence of chlorite, kaolinite and calcite which fits to
638 the soils of northern Tidra Island (Fig. 11a) making it a really local phenomenon. In contrast, dust sampled offshore
639 at site CBi was characterized by chlorite and by the absence of kaolinite which fits to the chlorite rich soils in the
640 Mauritanides of Mauritania (Fig. 11b).



Lithology

Carbonate	Ultramafic	Alluvium - beach, strand, coastal dune
Karst	Extrusive Volcanic	Alluvium - Saline
Non-Carbonate	Colluvium	Alluvium - Gypsum
Metasedimentary	Hydric - Organic	Alluvium - Other
Alkaline Intrusive Volcanic	Acolian Sediments	Volcanic - Ash, Tuff, Mudflow
Silicic	Alluvium - Fan Deposit	Water
Metigneous	Alluvium - Fluvial	

Major potential source areas (PSAs)

PSA 1-5

Study sites and back trajectories


10 m 4500 m 10 m
 ● Iwik ● CBi


Clay Mineralogy

⊘ No data Calcite >8.9 % ⊘ Kaolinite >29 % ⊘ Chlorite >4.1 %

Mineralogy

Kaolinite	Feldspar	Serpentine	Sepiolite	Gibbsite
Chlorite	Mica	Rutile	Anhydrite	Dolomite
Calcite	Amphibole	Zeolite	Smectite	Other minerals
Quartz	Palygorskite	Fluellite	Garnet	Ferryglaucophane







642

643 Figure 11: High- (4500 m) and low-level (10 m) four-day back trajectories of a dust event ending during the sampling
 644 interval 24.06.-15.07.13 at site Iwik and during the sampling interval 25.06.-16.07.13 at site CBI. The potential source
 645 areas and the mineralogy of the samples are given in the subfigures a-b. The dust storm event is indicated in subfigure
 646 c.

647 In Fig. 12 a typical late-summer situation is illustrated. At least five separate dust events could be identified (Fig.
 648 12f-j) out of which three occurred during the sampling interval of the site Iwik and two during the sampling interval

649 ~~of the site CBi. One of these dust storms occurring during the sampling interval of the site CBi lasted for two days~~
650 ~~(07-08.08.2014), while all other dust storms lasted for only one day. As a result, three back trajectories could be~~
651 ~~drawn for each site and each height, which one lasted for two days.~~The low-level back trajectories ending at site
652 CBi run offshore. The low-level back trajectories ending at site Iwik pass through the major **PSA 2**. The high-level
653 back trajectories pass through the major **PSA 2**, **PSA3** and **PSA 4** (Scheuvens et al., 2013). Dust deposited in the
654 continental traps was characterized by the presence of calcite and the absence of chlorite and kaolinite. Therefore,
655 the source area of the dust was most likely in the Western Sahara where soils rich in calcite but poor in chlorite
656 and kaolinite are located (Fig. 12a,b). Dust sampled with the oceanic traps during this sampling interval was
657 characterized by the absence of chlorite and kaolinite and by the presence of a high percentage of zeolite (22 %)
658 (Fig. 8c). Therefore, a possible source area may have been extrusive volcanic rocks of the northern Taoudeni Basin
659 (Fig. 12c) and the Fezzan uplift (Fig. 12e). Ferryglaucophane may have been sourced by the Pharusian belt (Fig.
660 12d).





662
 663 **Figure 12: High- (4500 m) and low-level (10 m) four-day back trajectories of dust events ending during the sampling**
 664 **interval 15.08.-15.09.14 at site Iwik and during the sampling interval 01.08.-21.08.14 at site CBI. The potential source**
 665 **areas and the mineralogy of the samples are given in the subfigures a-c. The dust storm events are indicated in**
 666 **subfigures e-i.**

667 In Fig. 13a-d the mean wind vectors and speed are presented for chosen dust storm events. The individual dust
 668 source areas that were identified using the back trajectory of the day with the dust storm as shown in Fig. 9-12 are

684 4. Discussion

685 4.1 Comparison of dust collected on land and in the ocean

686 4.1.1 Dust concentrations

687 ~~An a~~Annual average dust concentration (total suspended particles)s of ~214-~~(2013)~~ and 275 μm^{-3} -~~(2014)~~ ~~wasere~~
688 estimated for all dust samples of the year 2013 and 2014 respectively regarding the site Iwik (Table 54). These
689 estimates were larger than what has been measured for background dust concentrations (total suspended particles)
690 in Morocco which were in the order of 100 μm^{-3} during spring 2006 (Kandler et al., 2009). However, in Morocco
691 dust was collected at a larger height of 4 m and haze-periods and dust-storms were excluded from the average
692 value and the sampling time is much shorter leading to the monitoring of less dust events. The horizontal dust
693 fluxes at site Iwik correlated positively to wind speed (Fig. 7b) and decreased with collection height (not shown).
694 This underscores the proximity of this continental site to the dust emission source.

695 At the distal oceanic site CB, the annual average dust deposition flux was ~ 45 $\text{mgm}^{-2}\text{d}^{-1}$ (2013) (Table 54). The
696 dust flux was slightly larger than the average annual dust flux observed at site CB between 1988 and 2012 with ~
697 30 $\text{mgm}^{-2}\text{d}^{-1}$ (Fischer et al., 2015b). The slightly larger dust fluxes may have been caused by the anomalously high
698 frequency in dust storm events as observed on satellite images occurring during the studied time period (not
699 shown). ~~T~~In addition, the observed general decrease in the dust flux from the site Iwik to the sites CBI and CB can
700 be explained via the increase in the distance to the source area. Decreased dust deposition fluxes offshore NW
701 Africa with increasing distance from the African coast were also observed by Bory and Newton (2000) analysing
702 the lithogenic fluxes in marine sediment traps.

703 The average horizontal fluxes at site Iwik were ~ 1000 times larger with ~ 100000 $\text{mgm}^{-2}\text{d}^{-1}$ (Table 54) due to the
704 different sampling technique. The MWAC samplers do not measure deposition fluxes but foremost dust
705 concentrations. Only 1% or less drops out of a moving dust cloud within five minutes, hence, the horizontal dust
706 flux is at least ~100 times higher than the dust deposition flux (Goossens, 2008). ~~In addition, the observed general~~
707 ~~decrease in the dust flux from the site Iwik to the sites CBI and CB can be explained via the increase in the distance~~
708 ~~to the source area. Decreased dust deposition fluxes offshore NW Africa with increasing distance from the African~~
709 ~~east were also observed by Bory and Newton (2000) analysing the lithogenic fluxes in marine sediment traps.~~
710 The stronger decrease in the dust fluxes from site Iwik to CB during summer compared to winter (Table 4) may
711 be explained in the following. During summer, dust was additionally transported with the trades to the site Iwik,
712 (Fig. 9-12) leading to anomalously higher dust deposition at site Iwik compared to the oceanic sites. Further, the
713 washout of dust during offshore transport may have depleted the atmospheric dust cloud resulting in strongly
714 decreased dust deposition fluxes at site CB compared to site CBI during summer. The fact that the dust fluxes
715 decreased with height (not shown) further complicated a comparison between the sites due to the different
716 sampling heights of the dust collectors (2.90 m at Iwik versus sediment traps in the water). Therefore, the fluxes
717 between the site Iwik and the offshore sediment trap moorings cannot be compared.

718 4.1.2 Dust transport

719 The measured grain-size distributions for dust trapped at 2.90 m on land at site Iwik and for dust settling in the
720 ocean were predominantly nearly all unimodal (Fig. 6). Unimodal grain-size distributions are typical for wind-
721 blown sediments (Pye, 1995). Unimodal grain-size distributions were also measured for dust deposited in a vertical

722 dust sampler in M'Bour (Skonieczny et al., 2011), dust sampled on ship vessels (Stuut et al., 2005) and in other
723 sediment trap samples offshore NW Africa (Ratmeyer et al., 1999b; Van der Does et al., 2016; Friese et al., 2016).

724 The measured annual average modal grain size at site Iwik was 48 μm (Table 54). The obtained average annual
725 modal grain size was close to the large coarse mode population of 44 μm observed by Gillies et al. (1996) for dust
726 trapped at a height of 10 m during spring in Fakarbé (Mali) which is located about 700 km southeast of Iwik.
727 Gillies et al. (1996) conclude that the large coarse mode population in the dust samples points to locally-derived
728 dust. Based on this observation, we argue that also the dust trapped near Iwik was most likely generally of regional
729 instead of long-distance provenance. The distance to the main source area may be, however, not in the direct
730 surrounding of the dust collector since dust sampled with MWAC samplers in the vicinity of barchan dunes of the
731 Bodélé depression at 2.4 m height is characterized by a larger modal particle size of $\sim 100 \mu\text{m}$ (Chappell et al.,
732 2008). The annual average modal and maximum particle size gradually decreased from the on-land site Iwik, to
733 the proximal oceanic site CBi and the distal oceanic site CB (Table 54, Fig. 6a). This decrease in particle size
734 between the stations CB and CBi was observed before and was attributed to the preferred gravitational settling of
735 coarse particles during dust transport (Friese et al., 2016). Moreover, many studies have confirmed a downwind
736 fining of the terrigenous fraction of surface sediments offshore NW Africa (Koopmann, 1981; Holz et al.,
737 2004; Fütterer, 1980; Radczewski, 1939; Lange, 1975), and it is intuitively logical.

738 Bimodal grain-size distributions typically indicate the sampling of different dust sources (Stuut et al. (2009) and
739 references therein). The three samples of the Iwik time series that were characterized by an additional small peak
740 in the grain-size distribution around $\sim 16 \mu\text{m}$ were sampled during sampling intervals of anomalously high wind
741 velocity. The back-trajectories of one of these samples pointed towards a proximal and more distal dust source
742 (Fig. 12a,b). Therefore, it may be possible that wind velocities were high enough during the sampling interval to
743 inject dust to higher altitude and transport ~~it~~ dust from more distant sources (Fig. 12b) to the sampling site resulting
744 in the small peak in the grain-size distributions. This interpretation is further supported by the reanalysis wind
745 vector maps showing anomalously high wind velocities between the site Iwik and the proposed distant source area
746 enabling dust emission and transport of dust particles from more distant source to the site Iwik (Fig. 13d). On the
747 other hand, microscopic examination prior to particle-size analyses of the Iwik samples revealed that the samples
748 included many aggregates (Fig. 5d). Hence, locally derived aggregates may have been sampled during periods of
749 high wind velocities. These aggregates may have been dispersed in the demineralized water during the
750 measurement of the laser resulting in the observed additional smaller coarse mode population fine-peak at $\sim 16 \mu\text{m}$.
751 Further, precipitation was encountered according to the TRMM data during the sampling interval of two of these
752 three samples. Therefore, a further explanation for the bimodal grain-size distributions may be the deposition of
753 ~~finer~~ dust particles with a mode of $\sim 16 \mu\text{m}$ from higher altitude of the SAL due to precipitation. The rain droplets
754 may have evaporated during their fall releasing the dust particles at lower altitudes which can then be sampled
755 with the MWAC sampler. However, we also observed remnants of water in the bottles and therefore wet deposition
756 into the bottles may have also occurred. During summer, frequent rainfall resulted in a decrease of the modal
757 particle size of deposited Saharan dust at site Iwik (Fig. 7c). In addition, the seasonal average percentage of PM10
758 particles was larger during summer compared to winter at the site Iwik and CBi (Tab.5). These observations may
759 also be explained by the deposition of relatively finer dust particles from higher altitude of the SAL during summer
760 due to precipitation. One winter and one summer sample of the~~Two of the three~~ oceanic samples that were

761 characterized by bimodal grain-size distributions have several proposed dust source areas each (Fig. 10, 12). Thus,
762 the sampling of long- as well as short-travelled dust may have resulted in a bimodal grain-size distribution.

763 Both at the onshore sampling site Iwik and at the offshore sampling site CBi a clear seasonal trend in
764 the particle sizes of deposited dust could be observed with generally coarser modal particle sizes during summer
765 compared to winter (Fig. 6b,c). Generally coarser summer modal particle sizes of deposited dust at site CBi were
766 observed before for a three year time series during 2003 to 2006 and related to moist convective events (Friese et
767 al., 2016). The generally coarser particle sizes during summer at the site Iwik compared to winter may be explained
768 by the trade-wind speed as a positive correlation between the modal grain sizes and surface wind velocities was
769 observed (Fig. 7a), a positive correlation between the modal grain sizes and wind velocities was observed (Fig.
770 7a). This implied that dust was transported with the trade winds from sources of a quite constant distance year-
771 round. During dust-storm events particles with a diameter of 40 to 50 μm may be transported ~ 100 km (Tsoar
772 and Pye, 1987). The proposed source areas all fall in this range except for the winter sample of 2014-2015 (Fig.
773 10). The winter sample was characterized by an anomalously low modal grain size of 38 μm and particles of this
774 size may be transported more than 100 km during dust storm events (Tsoar and Pye, 1987). Moreover, Van der
775 Does et al. (2016) observed how particles up to 100 μm were transported ~ 3500 km across the Atlantic Ocean.
776 To sum up, the seasonal variability in the modal particle size of deposited dust at the site Iwik was mainly driven
777 by the surface wind speed due to the predominant sampling of nearby dust sources year-round.

778 4.1.3 Dust mineralogical composition

779 In the dust sampled at Iwik the minerals quartz, feldspar, mica, amphibole, palygorskite, chlorite, calcite, dolomite,
780 gibbsite, rutile and serpentine were present (Fig. 8a). The observed occurrence of the minerals quartz, feldspar,
781 mica, chlorite and calcite has also been described for the bulk size fraction of soil samples and dust samples
782 collected in Mauritania (Schütz and Sebert, 1987). Palygorskite, mica and chlorite have also been detected by
783 Skonieczny et al. (2013) in the PM_{30} size fraction of a three-year time series of dust deposition at M'Bour, Senegal,
784 more than 500km south of Iwik, Mauritania. Smectite and kaolinite, which were absent in the Iwik samples, were
785 the dominant minerals of the dust sampled at M'Bour (Skonieczny et al., 2013). Smectite and kaolinite are
786 considered as indicative for wet tropical soils and their relative abundance in soils increases southwards along the
787 northwest African coast (Lange, 1982; Biscaye, 1964). We argue that the mineralogical differences between the
788 two sites are explained by the >500 km distance between Iwik and M'Bour and the fact that the latter station is
789 surrounded by tropical soils. Gibbsite, rutile and serpentine have not been reported in any continental dust study
790 so far and thus seem to be indicative for locally-derived dust (Fig. 9a, Fig. 11a).

791 The dust sampled at the proximal marine site CBi contained the minerals quartz, feldspar, mica, amphibole,
792 palygorskite, chlorite, zeolite, kaolinite, smectite, sepiolite, rutile, garnet, anhydrite and fluellite (Fig. 8a). The first
793 seven of these minerals were also found in the clay and/or silt and sand fraction of Saharan dust sampled during
794 ship cruises parallel to the coast about 70 km off Cape Blanc (Chester et al., 1971) and perpendicular to the coast
795 about 80 to 180 km off Cape Blanc (Chester and Johnson, 1971b). Analogous to the samples of this study, the
796 PM_{20} fraction of surface sediments of the piston cores RC05-57, RC05-60 and A180-44 also feature zeolites and
797 the surface sediments of core RCRC05-57 also traces of pyrophyllite (sepiolite belongs to the pyrophyllite group)
798 (Biscaye, 1964). Further, rutile was also present in the silt and sand fraction of Saharan dust sampled perpendicular
799 to the coast on the research vessel (Chester and Johnson, 1971b). Palygorskite was found in the clay fraction of

800 the surface sediment of sediment core GIK12329 (19° 22' N, 19°56' W) offshore Cape Blanc and is considered a
 801 characteristic mineral of Saharan dust (Lange, 1975). The observed annual average C/K ratio (C/K=4) recorded
 802 for the bulk size fraction of the trap samples was larger than the C/K ratio (C/K=0.3-1) recorded in the clay fraction
 803 of surface sediment samples offshore Cape Blanc by Lange (1982). The disagreement may be due to the generally
 804 larger percentage of kaolinite in the clay fraction compared to the silt fraction (Journet et al., 2014).

805
 806 The dust samples of the site Iwik were further characterized by a dominance in quartz and feldspar (Fig. 8a). A
 807 dominance in quartz has also been described for continental dust samples and soil samples collected in Mauritania
 808 by Schütz and Sebert (1987). More than 20 papers published XRD data of northern African dust reporting quartz
 809 as the main mineral in most dust samples (Scheuven et al., 2013). ~~Moreover, the continental sampling site is~~
 810 ~~surrounded by sand dunes which are rich in quartz minerals (Schlüter, 2008; Lancaster, 2013). A high quartz~~
 811 ~~content may therefore point to predominantly locally derived dust.~~ The observed increase in micas and decrease
 812 in quartz and feldspar observed for the marine samples relative to the Iwik samples (Fig. 8a) can be explained via
 813 the preferential gravitational settling of the larger dust minerals quartz and feldspar during transport (Delany et al.,
 814 1967; Glaccum and Prospero, 1980; Chester and Johnson, 1971b; Schütz and Sebert, 1987). A strong downwind
 815 decrease in quartz content in Saharan dust was also observed by Korte et al. (2017).

816

817 4.2 Mineralogy as a provenancing tool

818

819 In Table 7 an overview of the chosen dust source areas for the site Iwik and CBi is given together with the
 820 characteristic minerals of the samples that may be used as a tracer for the source area. In the following subsections
 821 the identification of the source areas and mineralogical tracers is described in detail.

822 **Table 7: Overview of the chosen source areas and the tracer minerals of the individual samples together with the given**
 823 **characteristics of the source areas according to literature. ***

Sampling interval	Characteristic minerals of sample	Chosen dust source area	Bulk mineralogical composition of chosen PSA ⁽¹⁶⁾	Characteristic source rocks and deposits of chosen source area
Iwik				
15.02.- 15.03.14	*Rut, Serp, Cc	PSA 2: Reguibat Shield	C/K = 0.0-1.0 *Pal: 1-30 wt%	Metamorphic and granitic rocks ⁽¹⁾ Serpentinites ⁽²⁾
15.12.14- 18.01.15	*Cc, Chl, Pal (8 wt. %)	PSA 2: Senegal-Mauritania Basin	C/K = 0.0-1.0 *Pal: 1-30 wt%	Chalky horizons ⁽³⁾
		PSA 3: Eastern Taoudeni Basin	C/K = 0.2-0.9 *Pal: 1-5 wt%	Carbonate sequences ⁽⁴⁾
24.06.- 15.07.13	*Gib	PSA 2: Tidra Island	C/K = 0.0-1.0 *Pal: 1-30 wt%	Gibbsite maximum offshore Cape Blanc ⁽⁵⁾
15.08.- 15.09.14	*Cc, Dol, Pal (5 wt. %)	PSA 2: Aaiun-Tarfaya Basin	C/K = 0.0-1.0 *Pal: 1-30 wt%	Limestone deposits ⁽⁶⁾ Outcrops near Laäyoune with dolomites ⁽⁶⁾
CBi				
26.02.- 18.03.14	*Chl, Kao (C/K = 1), Pal (11 wt. %), Flu, Anh, Sme, Ga	PSA 2: Aaiun-Tarfaya Basin near Boucraa	C/K = 0.0-1.0 *Pal: 1-30 wt%	Phosphate deposits ⁽⁷⁾

16.12.14- 04.01.15	*Kao (C/K = 0), Pal (1 wt. %), Zeo, Se, Sme	dike swarms and sills of northern Taoudeni Basin	=	Basalts with glass ⁽⁹⁾
		PSA 2: Aaiun-Tarfaya Basin	C/K = 0.0-1.0 *Pal: 1-30 wt%	Palygorskite-sepiolite mafic clays ⁽¹⁰⁾
		Southern Senegal- Mauritania Basin	=	Lateritic soil ⁽⁸⁾ Horizontal layers of palygorskite and sepiolite ⁽⁸⁾
25.06.- 16.07.13	*Chl, Pal (2 wt. %), Rut	PSA 2: Mauritanides	C/K = 0.0-1.0 *Pal: 1-30 wt%	Strongly metamorphosed rocks ⁽¹⁰⁾ Greenschist facies ⁽¹¹⁾
01.08.- 21.08.14	*Fe-Amf, Zeo	dike swarms and sills of northern Taoudeni Basin	=	Basalts with glass ⁽⁹⁾
		PSA 4: Fezzan uplift	C/K = 0.0-2.6 *Pal: 0 wt%	Zeolite in basaltic rocks ^(12,13)
		PSA 3: Pharusian belt	C/K = 0.2-0.9 *Pal: 1-5 wt%	Blueschists ⁽¹⁴⁾ Glaucofane bearing eclogites ⁽¹⁵⁾
* Amf = amphibole, Pal = palygorskite, Chl = chlorite, Cc = calcite, Dol = dolomite, Gib = gibbsite, Zeo = zeolite, Kao = kaolinite, Sme = smectite, Se = sepiolite, Rut = rutile, Serp = serpentine, Ga = garnet, Anh = anhydrite, Flu = fluellite				
⁽¹⁾ Schofield et al. (2006) and references therein ⁽²⁾ Schlüter (2008) ⁽³⁾ Wissmann (1982) ⁽⁴⁾ Bertrand-Sarfati et al. (1991) ⁽⁵⁾ Biscaye (1964) ⁽⁶⁾ Bosse and Gwosdz (1996) ⁽⁷⁾ Moreno et al. (2006) ⁽⁸⁾ García-Romero et al. (2007) ⁽⁹⁾ Verati et al. (2005) ⁽¹⁰⁾ Villeneuve (2005), ⁽¹¹⁾ Dallmeyer and Lécorché (2012) ⁽¹²⁾ Abdel-Karim et al. (2013) ⁽¹³⁾ Cvetković et al. (2010) ⁽¹⁴⁾ Caby (2014), ⁽¹⁵⁾ Caby et al., 2008) ⁽¹⁶⁾ Scheuven et al. (2013)				

Field Code Changed

Formatted: English (United States)

Field Code Changed

Field Code Changed

Formatted: English (United States)

824

825 4.2.1 Dust collected on land

826 The variability of the mineralogical composition of dust sampled at the site Iwik could be related to the synoptic
827 scale change in the surface trade wind direction. However, meteorological data from nearby sites like e.g.
828 Nouadhibou demonstrate that local effects like the topography exert a strong influence on observed wind directions
829 at ground level (Fig. 2). The back trajectories indicate that the dust sources for the dust collected in Iwik during
830 winter were located NE and E of the sampling site (Fig. 9a, Fig. 10a,b), while those during summer were located
831 W (within the PNBA) and NNE of the sampling site (Fig. 11a, Fig. 12a,b). This is in accordance with a change in
832 the dominant local surface wind direction from NE in winter to NNE in summer (Fig. 2) and is also reflected in
833 the clay-mineralogical composition of the samples.

834 Generally, there is not much variability in the clay-mineralogical composition of the Iwik samples. The back
835 trajectories for the winter sample of 2014 indicate that the material was blown from the southwestern Reguibat
836 Shield (PSA 2) (Fig. 9a). The lack of palygorskite in this sample does not fit to the proposed bulk palygorskite
837 content (1-30 %) of PSA 2 (Scheuven et al., 2013) (Table 7). Therefore, we argue that the sampled dust was
838 most likely derived from a single-localized source instead of externally mixed sources of PSA 2, during transport.
839 The sample included the characteristic minerals rutile and serpentine (Table 7 Fig. 8b) which are usually a result
840 of metamorphic processes (Deer et al., 1992). Indeed, the western Reguibat Shield is composed of metamorphic
841 and granitic rocks (Schofield et al. (2006) and references therein) and the rocks are intruded by serpentinites
842 (Schlüter, 2008). The sample was further characterized by the highest quartz percentage among all samples (~ 50
843 %) (Fig. 8b). The sand dunes of the Azefal sand sea which cover part of the southwestern Reguibat Shield might
844 have sourced these quartz grains (Fig. 9a). The sand dunes may have been fed by outcropping carbonate deposits
845 at the northern rim of the Taoudeni Basin via the NE-trade winds leading to anomalously high percentages of
846 calcite in the sand dunes (Fig. 9). Thus, the sand dunes may have also sourced the calcite present in the sample
847 (Fig. 8b).

848

849 The winter sample of 2014-2015 was suggested to be sourced from sediments of the northern Senegal-Mauritania
850 Basin (**PSA 2**) (Fig. 10a) and the eastern rim of the Taoudeni Basin (**PSA 3**) (Fig 10b). The palygorskite content
851 of the sample (8 %) fits to the proposed bulk palygorskite content of **PSA 2** (Scheuven et al., 2013) (**Table 7**).
852 This may points to several externally mixed sources of PSA 2 during transport ~~instead of single local source~~.
853 The sample was further characterized by calcite and chlorite (**Table 7**~~Fig. 8b~~). The sediments in the northern
854 Senegal-Mauritania Basin (Fig. 10a) comprise Quaternary chalky horizons (Wissmann, 1982) which may have
855 sourced the calcite. More likely, calcite may have been derived from the Mesozoic carbonate sequences cropping
856 out in the eastern rim of the Taoudeni Basin (Bertrand-Sarfati et al., 1991) (Fig. 10b). A source area lying at the
857 Algerian/Mali border was also suggested for a chlorite and calcite bearing dust sample collected on the Canary
858 Islands (Alastuey et al., 2005). The winter dust sample trapped at site Iwik was further characterized by the lowest
859 feldspar percentage (~ 5 %), highest mica percentage (~ 40 %) (Fig. 8b) and lowest modal grain size (~ 38 µm)
860 among all Iwik dust samples analysed for mineralogy. The Stokes terminal settling velocity is smaller for platy
861 particles than for spherical particles of similar diameter (Santamarina and Cho, 2004). Therefore, a long-distance
862 transport of dust from the eastern Taoudeni Basin to Iwik may have resulted in a depletion in spherical quartz
863 particles (Fig. 5a,b,c) and an enrichment in platy mica particles (Fig. 5b).

864
865 The summer sample of 2013 was proposed to be sourced from the near-by northern Tidra Island (**PSA 2**) (Fig.
866 11a). Again, the absence of the mineral palygorskite is noteworthy which may points to the sampling of a single
867 localized dust source.

868 The sample was further characterized by the mineral gibbsite (**Table 7**~~Fig. 8e~~). The northern Tidra Island is famous
869 for the local occurrence of west Africa's northernmost mangroves (Proske et al., 2008) which grow in humid and
870 warm climates. Humid and warm conditions are also beneficial for the formation of gibbsite which forms through
871 tropical weathering (Deer et al., 1992). Therefore, we argue that the soils of Tidra Island supplied the gibbsite
872 found in the sample. A localized small gibbsite maximum was outlined for the surface sediments offshore Cape
873 Blanc (Biscaye, 1964) which further supports the view that gibbsite is supplied from a local source. The sample
874 was further characterized by anomalously large moderately spherical quartz grains (Fig. 5c) emphasizing a short
875 travel distance of the dust.

876
877 The summer sample of 2014 was most likely sourced by sediments of the Western Sahara (**PSA 2**) (Fig. 12a,b).
878 The palygorskite content of the sample (5 %) matches with the proposed bulk palygorskite content of **PSA 2**
879 (Scheuven et al., 2013) (**Table 7**). Hence, dust may have been ~~was~~ supplied from several ~~local~~ dust sources of
880 **PSA 2** which were mixed during transport.

881 The sample was further characterized by calcite and dolomite (**Table 7**~~Fig. 8e~~). Sediments outcropping in the
882 Western Sahara are composed of Tertiary sediments (Wissmann, 1982) with limestone deposits (Bosse and
883 Gwosdz, 1996) that may explain the calcite found in the sample (Fig., 12a). Upper cretaceous outcrops in the
884 Aaiun-Tarfaya Basin near Laâyoune comprise dolomites (Bosse and Gwosdz, 1996) and could have sourced the
885 dolomite found in the sample (Fig. 12b). A further evidence for dolomite-bearing dust transport from the Aaiun-
886 Tarfaya Basin is a local dolomite maximum outlined for the surface sediments offshore the Western Sahara
887 (Johnson, 1979). A Saharan dust sample trapped in NE Spain also contained dolomite and calcite and was related
888 to a source area lying in the Western Sahara (Avila et al., 1997).

889

Formatted: Font: Bold

890 4.2.1 Dust collected at the marine sites

891 The seasonal contrast in the dust transport patterns (high-level Saharan Air Layer vs. low-level Trades) potentially
892 led to strongly deviating dust sources for the material deposited in the marine trap samples. During winter, the
893 back trajectories indicated that the potential dust source areas were located NE of the sampling site (Fig. 9b, Fig.
894 10c,d), while those during summer were located NE, E and SE of the sampling site (Fig. 11b, Fig.12a,c,d,e). This
895 large variability in wind patterns can clearly be recognized in the clay-mineralogical compositions of the samples
896 throughout the seasons.

897 Considering the much larger catchment area of the traps, several ~~localized~~ dust sources may have been sampled
898 with the traps. As a result, the composition of the analyzed samples fit well to the bulk composition of the chosen
899 **PSA**. The back trajectories indicate that the winter sample of 2014 originated from the shoreline of the Western
900 Sahara (**PSA 2**) (Fig. 9b). The observed C/K ratio (C/K=1) and the palygorskite content (11 %) are in agreement
901 with the bulk compositional C/K ratio (C/K=0-1) and palygorskite content of **PSA 2** (Scheuvens et al., 2013)
902 (Table 7).

903 The sample was further characterized by the presence of ~~chlorite, kaolinite, smectite,~~ garnet, ~~anhydrite and~~ fluellite
904 ~~and anhydrite~~ (Table 7)(Fig. 8b). The characteristic occurrence of garnet together with the highest quartz content
905 (33 %, Fig. 8b) among all CBi samples confirms a short transport distance of the trapped dust. ~~Chlorite may be~~
906 ~~sourced from a small coastal area where chlorite-rich fluvisols are found (Journet et al., 2014) (Fig. 9b).~~ The
907 mineral fluellite which is a weathering product of phosphate may have been derived from outcropping phosphate
908 deposits near the Bucraa phosphate mine (Moreno et al., 2006) (Fig. 9b). ~~The occurrence of kaolinite in the sample~~
909 ~~is remarkable as this mineral was not observed in high amounts in the soils underlying the back trajectories.~~
910 ~~However, the sample was further characterized by a lack of feldspar (Fig. 8b), which tends to be hydrothermally~~
911 ~~altered to kaolinite (Deer et al., 1992). Anhydrite could. The same process may explain the presence of anhydrite~~
912 ~~in this sample, although this mineral could also originate from evaporites along the coast. Another explanation for~~
913 ~~the presence of kaolinite and smectite may be the transport of these minerals from southern latitudes via the~~
914 ~~poleward flowing undercurrent to the trap site CBi (Fig. 1). Kaolinite and smectite were found in the clay fraction~~
915 ~~of the surface sediments off Senegal (Nizou et al., 2011) and may have been brought into the ocean by the Senegal~~
916 ~~River, and redistributed by ocean currents (Biscaye, 1964). The season of high Senegal River sediment supply is~~
917 ~~between July to October/November (Gac and Kane, 1986). Assuming a mean speed of ~10 cm/s of the~~
918 ~~undercurrent (Mittelstaedt, 1991), it may take about two months for the particles to travel a distance of ~500 km~~
919 ~~to the trap site CBi. This time delay might explain the observed occurrence of these minerals in the trap samples~~
920 ~~during winter, but not during summer.~~

921
922 The back trajectories of the winter sample of 2014 to 2015 lead to the Reguibat Shield (**PSA 2**) (Fig. 10c) and
923 coastal Western Sahara (**PSA 2**) (Fig. 10d). The observed C/K ratio (C/K=0) and palygorskite content (1 %) fall
924 within the ranges of these minerals in **PSA 2** (Scheuvens et al., 2013) (Table 7).

925 The sample was further characterized by the mineral zeolite, ~~kaolinite, sepiolite and smectite~~ (Table 7)(Fig. 8b).
926 Zeolites are formed from volcanic glass and tuff and form well-developed crystals in basalts (Deer et al., 1992).
927 Therefore, the source area of the zeolites may have been outcropping volcanic rocks in the northern Taoudeni
928 Basin (Fig. 10c). These rocks belong to mafic dikes and sills which are commonly basalts with dotted patches of
929 glass (Verati et al., 2005). An additional indication for a distant dust source may be the lowest quartz content (4

930 %) among all samples (Fig. 8b). Palygorskite-sepiolite mafic clays were found in soil samples of the Western
931 Sahara (Moreno et al., 2006) which may supports a Western Saharan source (Fig. 10d).
932 The CBI trap sample was further characterized by the minerals sepiolite and smectite (Fig. 8b). Sepiolite belongs
933 to the pyrophyllites which is a mineral that also may be considered indicative of tropical weathering (Moore and
934 Reynolds, 1989). Similar to the winter dust sample recovered during 2014, sepiolite and smectite may have been
935 derived from humid weathering on the wet shoreline of the Western Sahara (Fig. 10d) or from current transport of
936 clay particles from the Senegal River mouth. Palygorskite-sepiolite mafic clays were found in soil samples of the
937 Western Sahara (Moreno et al., 2006) which may supports a Western Saharan source. Moreover, kaolinite is
938 usually considered indicative of tropical weathering and the laterites of the southern Sahara and Sahel (Lange,
939 1975; Biscaye, 1964; Lange, 1982). Outcrops of quaternary laterites as well as outcrops of lower Eocene horizontal
940 layers of palygorskite and sepiolite were described near Thiès in Senegal (García-Romero et al., 2007). Therefore,
941 the kaolinite-rich soils and outcrops in the southern Senegal-Mauritania basin near Thiès (Fig. 10e) may have
942 served the kaolinite, sepiolite and palygorskite found in the sample.
943 Another explanation for the presence of kaolinite and smectite in the sample may be the transport of these minerals
944 from southern latitudes via the poleward-flowing undercurrent to the trap site CBI (Fig. 1). Kaolinite and smectite
945 were found in the clay fraction of the surface sediments off Senegal (Nizou et al., 2011) and may have been brought
946 into the ocean by the Senegal River, and redistributed by ocean currents (Biscaye, 1964). The season of high
947 Senegal River sediment supply is between July to October/November (Gac and Kane, 1986). Assuming a mean
948 speed of ~10 cm/s. of the undercurrent (Mittelstaedt, 1991), it may take about two months for the particles to travel
949 a distance of ~500 km to the trap site CBI. This time delay might explain the observed occurrence of these minerals
950 in the trap samples during winter, but not during summer.

951
952 Based on the back trajectories, the summer sample of 2013 was suggested to be sourced from the Mauritanides
953 (PSA 2) (Fig. 11c). This is confirmed by the palygorskite content of the sample (2 %) (Scheuven et al., 2013)
954 (Table 7). Outstanding minerals in this sample are chlorite and rutile (Table 7 Fig. 8e). Outcrops in the Mauritanides
955 west of the Taoudeni Basin feature strongly metamorphosed rocks (Villeneuve, 2005) and greenschist facies
956 (Dallmeyer and Lécorché, 2012) which may have been the source of the rutile and chlorite.

957
958 The reconstructed source area of the summer sample of 2014 was the Pharusian belt (PSA 3) (Fig. 12c), ~~and~~ the
959 extrusive volcanics of the northern Taoudeni Basin (PSA 2) (Fig. 12d) ~~and the Fezzan uplift (PSA 4) (Fig. 12e).~~
960 The lack of palygorskite in the sample does corroborate with PSA 4 ('not detected') (Scheuven et al., 2013)
961 suggesting that the provenance of the dust sample may be mainly confined to PSA 4 (Table 7). The sample was
962 further characterized by zeolite and ferryglaucophane (Table 7 Fig. 8e). The dike swarms and sills of the northern
963 Taoudeni Basin (Verati et al., 2005) (Fig. 10c, Fig. 12c) and/or the basalts of the Fezzan uplift (Fig. 12e) may have
964 sourced the zeolite. Indeed, zeolite was described as one of the main secondary minerals in the basaltic rocks of
965 the central Al-Harui Al-Abyas basalt flows (Abdel-Karim et al., 2013) and in vesicles of the east Al Haruj basalts
966 (Cvetković et al., 2010) of the Fezzan uplift. Traces of zeolite were also detected in the Iwik sample during this
967 sampling interval. It may be that the zeolite dropped out of the high-altitude dust cloud and was subsequently
968 transported via the surface trade winds to the continental trap site. The presence of ferryglaucophane and the
969 absence of feldspar and chlorite in the sample indicates highly metamorphous outcrops constituting the dust source.
970 Therefore, the sample may have been additionally sourced by the Pharusian belt (Fig. 12c) because blueschists

Formatted: Font: Bold

971 were observed in Timétrine (Caby, 2014) and glaucophane bearing eclogites in the Gourma fold and thrust belt
972 north of Gao (Caby et al., 2008). The sample was further characterized by the highest mica content (44 %) among
973 all samples (Fig. 8c) supporting a large dust transport distance.

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994


995

996

997

998

999



1000 **5. Summary and conclusions**

1001 The fluxes, grain-size distributions and the mineral assemblages of the continental trap samples and oceanic
1002 sediment trap samples were well comparable to the characteristics of Saharan dust reported for the region. The
1003 following main findings were made:

- 1004 - ~~A clear seasonal variability in the particle size of mineral dust deposited on land could be observed with~~
1005 ~~generally coarser modal grain sizes during summer compared to winter. The modal particle sizes could be~~
1006 ~~related to the trade wind speed.~~
- 1007 - dust deposited on the continent was predominantly transported ~~from near-by local with the trade winds from~~
1008 ~~proximal~~ sources (Mauritania, Western Sahara, Mali), while dust deposited in the marine traps was
1009 transported ~~with both the trade winds (winter, proximal) and in the Saharan Air Layer (summer, distal)~~ from
1010 proximal (Mauritania, Western Sahara, Mali) and distal sources (Senegal and Libya).
- 1011 - ~~the percentage of mica relative to the quartz content increased in the deposited dust with increasing transport~~
1012 ~~distance, most likely due to the platy shape of these minerals, which reduces settling.~~
- 1013 - ~~Some rare characteristic minerals (e.g. ferriyglaucofane, rutile, serpentine) could be related to local outcrops~~
1014 ~~in NW Africa~~

1015 To conclude, the particle size and mineralogy of Saharan dust recorded in continental climate archives should be
1016 interpreted differently with respect to paleo-environmental conditions compared to marine climate archives; the
1017 on-land archive seems to reflect a much more local signal as compared to the regional signal that is recorded in
1018 the marine sediments. Given the relationship between particle size and wind strength, we suggest that the particle
1019 size in the continental archive in NW Africa may indicate the paleo-wind strength of the trade winds. This is an
1020 intuitively logical conclusion, but it has not been demonstrated before so clearly. ~~Finally, we have shown how the~~
1021 ~~mineralogical composition of the samples can be used for provenancing of dust particles found in both on-land~~
1022 ~~and marine dust archives. It should be kept in mind, however, that the wind strength in the sampling location~~
1023 ~~might differ from the wind strength in the source region if the source region is further away. Moreover, the sizes~~
1024 ~~of dust particles present in the source region will influence the grain sizes of deposited dust.~~

Formatted: English (United Kingdom)


Formatted: English (United Kingdom)

1035 **6. Appendices**

1036 **A1 Satellite RGB images**

1037

1038 In Fig. A1-4 satellite RGB true colour images are shown of the identified dust storms occurring during the sampling
1039 interval of the summer samples of site CBi analysed for dust provenance. On 31 July 2014 only few dust can be
1040 observed which overlies the sampling location CBi (Fig. A2). This fits to the observed minor percentage of the
1041 mineral ferrylglaucophane (7 %) in the sample which was suggested to be sourced on 31 July 2014 from PSA 3.
1042 Zeolite, which was more abundant (22 %) in the dust sample, was therefore most likely derived from PSA 4 due
1043 to the major dust storm event occurring on 7 August 2014 (Fig. A3).
1044



1045

1046

1047

Figure A1: Dust storm on 02 July 2013.

1048
1049




Figure A2: Dust storm on 31 July 2014.

1050
1051
1052




Figure A3: Dust storm on 07 August 2014.




Figure A4: Dust storm on 08 August 2014.

1053

1054

1055

1056 **A2 Four day back-trajectories**


1057

1058 In Fig. A5-8 the four day back-trajectories are shown calculated at the heights 3000 m, 4500 m and 5500 m ending
 1059 at site CBI. These high altitude back-trajectories were calculated for the identified summer days with dust storm
 1060 events (shown in Fig A1-4). On the one hand, a height of 4500 m was chosen by Skonieczny et al. (2013) in a dust
 1061 provenance study to represent the Saharan air layer (SAL). On the other hand, a height of 5500 m was chosen by
 1062 Ratmeyer et al. (1999a) in a dust transport study to represent the SAL. Maximum wind velocities within the SAL
 1063 are observed at a height of ~ 3 - 4 km in the area of the Cape Verde Islands during summer according to Carlson
 1064 and Prospero (1972). Therefore, we also plotted the back-trajectories at a height of 3000 m. In order to investigate
 1065 which air layer should be chosen for provenance studies, the back trajectories of the different heights were
 1066 compared.

1067 The back-trajectories deviated slightly from each other regarding their direction and length. The back-trajectories
 1068 at 3000 m showed the most deviation. Further, the back-trajectories at 4500 m showed the best agreement with the
 1069 source areas and the minerals in the samples. Therefore, we chose to use the trajectories at 4500 m for provenance
 1070 studies according to Skonieczny et al. (2013).

1071


NOAA HYSPLIT MODEL
 Backward trajectories ending at 1800 UTC 02 Jul 13
 CDC1 Meteorological Data



1072
 1073
 1074

Figure A5: Four day back-trajectories at a height of 3000 m, 4500 m and 5500 m on 02 July 2013.


NOAA HYSPLIT MODEL
 Backward trajectories ending at 1800 UTC 31 Jul 14
 CDC1 Meteorological Data



1075
 1076
 1077


Figure A6: Four day back-trajectories at a height of 3000 m, 4500 m and 5500 m on 31 July 2014.

NOAA HYSPLIT MODEL
 Backward trajectories ending at 1800 UTC 07 Aug 14
 CDC1 Meteorological Data



1078
 1079

Figure A7: Four day back-trajectories at a height of 3000 m, 4500 m and 5500 m on 07 August 2014.



1080

1081

Figure A8: Four day back-trajectories at a height of 3000 m, 4500 m and 5500 m on 08 August 2014.

1082

1083

7. Supplement link

1084

The data can be accessed on www.pangaea.de.

1085

8. Author contribution

1086

C. Friese carried out the particle size analysis of the sediment trap samples. H. van Hateren carried out the flux

1087

and particle size analysis of the Iwik dust samples. G. Fischer provided the sediment trap samples and supervised

1088

the flux analysis of the sediment trap samples. C. Friese prepared the samples for XRD analysis. C. Vogt carried

1089

out the XRD analysis and was involved in the discussion of the results. J.-B. Stuut managed the projects through

1090 which dust-collecting buoy ‘Carmen’ was constructed and deployed, supervised the particle-size analysis and the
1091 writing of the manuscript. C. Friese prepared the manuscript with contributions from all co-authors.

1092 **9. Competing interests**

1093 The authors declare that they have no conflict of interest.

1094 **10. Acknowledgements**

1095 We thank the captains, crews and scientific teams of the research cruises with RV Poseidon in 2013 (POS445),
1096 RV Poseidon in 2014 (POS464) and RV Poseidon in 2015 (POS481), during which the sediment traps were
1097 deployed and received. Further, we thank Marco Klann for preparing and splitting the sediment trap samples. Jan-
1098 Berend Stuut acknowledges funding from ERC Grant 311152 DUSTTRAFFIC. Funding is acknowledged from
1099 the German Science Foundation (DFG) through the DFG-Research Center/Cluster of Excellence ‘The Ocean in
1100 the Earth System’. We further thank Prof. Dr. Dierk Hebbeln and Dr. Ute Merkel for helpful and productive
1101 scientific discussions.

1102 **11. References**

- 1103 Abdel-Karim, A.-A. M., Ramadan, E.-N. M., and Embashi, M. R.: Multiphase Alkaline Basalts of Central Al-Haruj Al-
1104 Abyad of Libya: Petrological and Geochemical Aspects, *Journal of Geological Research*, 2013, 12, 10.1155/2013/805451,
1105 2013.
- 1106 Alastuey, A., Querol, X., Castillo, S., Escudero, M., Avila, A., Cuevas, E., Torres, C., Romero, P.-M., Exposito, F., and
1107 García, O.: Characterisation of TSP and PM_{2.5} at Izana and Sta. Cruz de Tenerife (Canary Islands, Spain) during a Saharan
1108 Dust Episode (July 2002), *Atmospheric Environment*, 39, 4715-4728, 2005.
- 1109 Aston, S. R., Chester, R., Johnson, L. R., and Padgham, R. C.: Eolian dust from the lower atmosphere of the eastern Atlantic
1110 and Indian oceans, China Sea and Sea of Japan, *Marine Geology*, 14, 15-28, 1973.
- 1111 Avila, A., Queralt-Mitjans, I., and Alarcon, M.: Mineralogical composition of African dust delivered by red rains over
1112 northeastern Spain, *Journal of Geophysical Research*, 102, 21977-21996, 1997.
- 1113 Bertrand-Sarfati, J., Moussine-Pouchkine, A., Affaton, P., Trompette, R., and Bellion, Y.: Cover sequences of the West
1114 African craton, in: *The West African orogens and circum-Atlantic correlatives*, Springer, 65-82, 1991.
- 1115 Biscaye, P.: Mineralogy and sedimentation of recent Deep-sea clay in the Atlantic Ocean and adjacent seas and oceans., *Geol*
1116 *Soc Am Bull*, 76, 803-832, 1965.
- 1117 Biscaye, P. E.: Mineralogy and sedimentation of the deep-sea sediment fine fraction in the Atlantic Ocean and adjacent seas
1118 and oceans, Ph.D., *Geology*, Yale University, Michigan, 86 pp., 1964.
- 1119 Bloemsmma, M. R., Zabel, M., Stuut, J. B. W., Tjallingii, R., Collins, J. A., and Weltje, G. J.: Modelling the joint variability of
1120 grain size and chemical composition in sediments, *Sedimentary Geology*, 280, 135-148,
1121 <http://dx.doi.org/10.1016/j.sedgeo.2012.04.009>, 2012.

- 1122 Bory, A. J. M., and Newton, P. P.: Transport of airborne lithogenic material down through the water column in two
1123 contrasting regions of the eastern subtropical North Atlantic Ocean, *Global Biogeochem. Cycles*, 14, 297-315,
1124 10.1029/1999gb900098, 2000.
- 1125 Bosse, H.-R., and Gwosdz, W.: Limestone and dolomite resources of Africa, *Geologisches Jahrbuch Reihe D*, 500, 102,
1126 1996.
- 1127 Boullier, A.-M.: The pan-African trans-Saharan belt in the Hoggar shield (Algeria, Mali, Niger): A review, in: *The west*
1128 *African orogens and circum-Atlantic correlatives*, Springer, 85-105, 1991.
- 1129 Caby, R., Buscaïl, F., Dembele, D., Diakite, S., Sacko, S., and Bal, M.: Neoproterozoic garnet-glaucophanites and eclogites:
1130 new insights for subduction metamorphism of the Gourma fold and thrust belt (eastern Mali), *Geological Society, London,*
1131 *Special Publications*, 297, 203-216, 2008.
- 1132 Caby, R.: Nature and evolution of Neoproterozoic ocean-continent transition: Evidence from the passive margin of the West
1133 African craton in NE Mali, *Journal of African Earth Sciences*, 91, 1-11, <http://dx.doi.org/10.1016/j.jafrearsci.2013.11.004>,
1134 2014.
- 1135 Caquineau, S., Gaudichet, A., Gomes, L., and Legrand, M.: Mineralogy of Saharan dust transported over northwestern
1136 tropical Atlantic Ocean in relation to source regions, *Journal of Geophysical Research*, 107, 4251, 2002.
- 1137 Carlson, T. N., and Prospero, J. M.: The Large-Scale Movement of Saharan Air Outbreaks over the Northern Equatorial
1138 Atlantic, *Journal of Applied Meteorology*, 11, 283-297, 1972.
- 1139 Chappell, A., Warren, A., O'Donoghue, A., Robinson, A., Thomas, A., and Bristow, C.: The implications for dust emission
1140 modeling of spatial and vertical variations in horizontal dust flux and particle size in the Bodélé Depression, Northern Chad,
1141 *Journal of Geophysical Research: Atmospheres*, 113, n/a-n/a, 10.1029/2007JD009032, 2008.
- 1142 Chester, R., Elderfield, H., and Griffin, J. J.: Dust transported in the North-east and South-east Trade Winds in the Atlantic
1143 Ocean, *Nature*, 233, 474-476, 10.1038/233474a0, 1971.
- 1144 Chester, R., and Johnson, L. R.: Atmospheric dusts collected off the Atlantic coasts of North Africa and the Iberian
1145 Peninsula, *Marine Geology*, 11, 251-260, 1971a.
- 1146 Chester, R., and Johnson, L. R.: Atmospheric dusts collected off the West African Coast, *Nature*, 229, 105-107, 1971b.
- 1147 Chester, R., Elderfield, H., Griffin, J., Johnson, L., and Padgham, R.: Eolian dust along the eastern margins of the Atlantic
1148 Ocean, *Marine Geology*, 13, 91-105, 1972.
- 1149 Chung, F. H.: Quantitative interpretation of X-ray diffraction patterns of mixtures. I. Matrix-flushing method for quantitative
1150 multicomponent analysis, *Journal of Applied Crystallography*, 7, 519-525, 1974.
- 1151 Cropper, T. E., Hanna, E., and Bigg, G. R.: Spatial and temporal seasonal trends in coastal upwelling off Northwest Africa,
1152 1981–2012, *Deep Sea Research Part I: Oceanographic Research Papers*, 86, 94-111,
1153 <http://dx.doi.org/10.1016/j.dsr.2014.01.007>, 2014.
- 1154 Cvetković, V., Toljić, M., Ammar, N. A., Rundić, L., and Trish, K. B.: Petrogenesis of the eastern part of the Al Haruj basalts
1155 (Libya), *Journal of African Earth Sciences*, 58, 37-50, <http://dx.doi.org/10.1016/j.jafrearsci.2010.01.006>, 2010.

- 1156 Dallmeyer, R. D., and Lécorché, J.-P.: The West African orogens and circum-Atlantic correlatives, Springer Science &
1157 Business Media, 2012.
- 1158 Deer, W. A., Howie, R. A., and Zussman, J.: An introduction to the rock-forming minerals, Longman Scientific & Technical,
1159 1992.
- 1160 Delany, A. C., Claire Delany, A., Parkin, D. W., Griffin, J. J., Goldberg, E. D., and Reimann, B. E. F.: Airborne dust
1161 collected at Barbados, *Geochimica et Cosmochimica Acta*, 31, 885-909, 1967.
- 1162 Diaz, H. F., Carlson, T. N., Prospero, J. M., and Office, E. R. L. W. M. P.: A Study of the Structure and Dynamics of the
1163 Saharan Air Layer Over the Northern Equatorial Atlantic During BOMEX, Nr. 32, Weather Modification Program Office,
1164 Environmental Research Laboratories, 1976.
- 1165 Diester-Haass, L., and Chamley, H.: Neogene paleoenvironment off NW Africa based on sediments from DSDP Leg 14,
1166 *Journal of Sedimentary Research*, 48, 1978.
- 1167 Dobson, M.: An account of the Harmattan, a singular African wind, *Philosophical transactions of the Royal Society of*
1168 *London*, 71, 46-57, 1781.
- 1169 Einsele, G., Herm, D., and Schwarz, H. U.: Sea level fluctuation during the past 6000 yr at the coast of Mauritania,
1170 *Quaternary Research*, 4, 282-289, [http://dx.doi.org/10.1016/0033-5894\(74\)90017-9](http://dx.doi.org/10.1016/0033-5894(74)90017-9), 1974.
- 1171 El Makkrouf, A. A.: Tectonic interpretation of Jabal Eghej area and its regional application to Tibesti orogenic belt, south
1172 central Libya (S.P.L.A.J.), *Journal of African Earth Sciences (and the Middle East)*, 7, 945-967,
1173 [http://dx.doi.org/10.1016/0899-5362\(88\)90009-7](http://dx.doi.org/10.1016/0899-5362(88)90009-7), 1988.
- 1174 Filipsson, H. L., Romero, O. E., Stuu, J.-B. W., and Donner, B.: Relationships between primary productivity and bottom-
1175 water oxygenation off northwest Africa during the last deglaciation, *Journal of Quaternary Science*, 26, 448-456,
1176 DOI:10.1002/jqs.1473, 2011.
- 1177 Fischer, G., and Wefer, G.: Sampling, Preparation and Analysis of Marine Particulate Matter, *Geoph Monog Series*, 63, 391-
1178 397, 1991.
- 1179 Fischer, G., and Karakas, G.: Sinking rates and ballast composition of particles in the Atlantic Ocean: implications for the
1180 organic carbon fluxes to the deep ocean, *Biogeosciences*, 6, 85-102, 2009.
- 1181 Fischer, G., Ba, M., Dehning, K., Hefter, J., Iversen, M., Klann, M., Nowald, N., Ploug, H., Ruhland, G., and Witte, Y.:
1182 Report and preliminary results of R/V POSEIDON cruise POS445, Las Palmas–Las Palmas, 19.01. 2013–01.02. 2013, 2013.
- 1183 Fischer, G., Dehning, K., Dia, A., Füssel, J., Hefter, J., Iversen, M., Klann, M., Nowald, N., Olbrich, M., and Ruhland, G.:
1184 Report and preliminary results of RV POSEIDON cruise POS464, Las Palmas (Canary Islands)-Las Palmas (Canary Islands),
1185 03.02. 2014-18.02. 2014, 2014.
- 1186 Fischer, G., Dia, A., Iversen, M., Klann, M., Nowald, N., Markussen, T., Meckel, S., Ruhland, G., Van der Jagt, H., and
1187 Waldmann, C.: Report and preliminary results of R/V POSEIDON cruise POS481, Las Palmas (Canary Islands)-Las Palmas
1188 (Canary Islands), 15.03. 2015-03.03. 2015, 2015a.

- 1189 Fischer, G., Romero, O., Merkel, U., Donner, B., Iversen, M., Nowald, N., Ratmeyer, V., Ruhland, G., Klann, M., and Wefer,
1190 G.: Deep ocean mass fluxes in the coastal upwelling off Mauritania from 1988 to 2012: variability on seasonal to decadal
1191 timescales, *Biogeosciences Discussions*, 12, 2015b.
- 1192 Formenti, P., Rajot, J. L., Desboeufs, K., Caquineau, S., Chevaillier, S., Nava, S., Gaudichet, A., Journet, E., Triquet, S.,
1193 Alfaro, S., Chiari, M., Haywood, J., Coe, H., and Highwood, E.: Regional variability of the composition of mineral dust from
1194 western Africa: Results from the AMMA SOP0/DABEX and DODO field campaigns, *Journal of Geophysical Research:*
1195 *Atmospheres*, 113, n/a-n/a, 10.1029/2008JD009903, 2008.
- 1196 Friese, C. A., van der Does, M., Merkel, U., Iversen, M. H., Fischer, G., and Stuut, J.-B. W.: Environmental factors
1197 controlling the seasonal variability in particle size distribution of modern Saharan dust deposited off Cape Blanc, *Aeolian*
1198 *Research*, 22, 165-179, <http://dx.doi.org/10.1016/j.aeolia.2016.04.005>, 2016.
- 1199 Fütterer, D.: Sedimentation am NW-afrikanischen Kontinentalrand: Quantitative Zusammensetzung und Verteilung der
1200 Siltfraktion in den Oberflächensedimenten, *Meteor-Forschungsergebnisse*, C, 15-60, 1980.
- 1201 Gac, J. Y., and Kane, A.: Le Fleuve Sénégal. I. Bilan hydrologique et flux continentaux de matières particulaires a
1202 l'embouchure, *Sci. Géol. Bull.*, 39, 99-130, 1986.
- 1203 García-Romero, E., Suárez, M., Santarán, J., and Alvarez, A.: Crystallochemical Characterization of the Palygorskite and
1204 Sepiolite from the Allou Kagne Deposit, Senegal, *Clays and Clay Minerals*, 55, 606-617, 10.1346/CCMN.2007.0550608,
1205 2007.
- 1206 Gillies, J. A., Nickling, W. G., and McTainsh, G. H.: Dust concentrations and particle-size characteristics of an intense dust
1207 haze event: inland delta region, Mali, West Africa, *Atmospheric Environment*, 30, 1081-1090, 1996.
- 1208 Glaccum, R. A., and Prospero, J. M.: Saharan aerosols over the tropical North Atlantic - mineralogy, *Marine Geology*, 37,
1209 295-321, 1980.
- 1210 Goossens, D., and Offer, Z. Y.: Wind tunnel and field calibration of six aeolian dust samplers, *Atmospheric Environment*, 34,
1211 1043-1057, 10.1016/s1352-2310(99)00376-3, 2000.
- 1212 Goossens, D.: Relationships between horizontal transport flux and vertical deposition flux during dry deposition of
1213 atmospheric dust particles, *Journal of Geophysical Research: Earth Surface*, 113, n/a-n/a, 10.1029/2007JF000775, 2008.
- 1214 Griffin, J. J., Windom, H., and Goldberg, E. D.: The distribution of clay minerals in the World Ocean, *Deep Sea Research*
1215 *and Oceanographic Abstracts*, 15, 433-459, [http://dx.doi.org/10.1016/0011-7471\(68\)90051-X](http://dx.doi.org/10.1016/0011-7471(68)90051-X), 1968.
- 1216 Haywood, J., and Boucher, O.: Estimates of the direct and indirect radiative forcing due to tropospheric aerosols: A review,
1217 *Rev Geophys*, 38, 513-543, Doi 10.1029/1999rg000078, 2000.
- 1218 Holz, C., Stuut, J.-B. W., and Henrich, R.: Terrigenous sedimentation processes along the continental margin off NW-Africa:
1219 implications from grain-size analyses of surface sediments, *Sedimentology*, 51, 1145-1154, .1111/j.1365-3091.2004.00665.x,
1220 2004.
- 1221 Holz, C., Stuut, J.-B. W., Henrich, R., and Meggers, H.: Variability in terrigenous sedimentation processes off northwest
1222 Africa and its relation to climate changes: Inferences from grain-size distributions of a Holocene marine sediment record,
1223 *Sedimentary Geology*, 202, 499-508, 10.1016/j.sedgeo.2007.03.015, 2007.

- 1224 Iversen, M. H., Nowald, N., Ploug, H., Jackson, G. A., and Fischer, G.: High resolution profiles of vertical particulate organic
 1225 matter export off Cape Blanc, Mauritania: Degradation processes and ballasting effects, *Deep Sea Research Part I:*
 1226 *Oceanographic Research Papers*, 57, 771-784, DOI: 10.1016/j.dsr.2010.03.007, 2010.
- 1227 Iversen, M. H., and Ploug, H.: Ballast minerals and the sinking carbon flux in the ocean: carbon-specific respiration rates and
 1228 sinking velocities of macroscopic organic aggregates (marine snow), *Biogeosciences Discuss.*, 7, 3335-3364, 10.5194/bgd-7-
 1229 3335-2010, 2010.
- 1230 Iversen, M. H., and Robert, M. L.: Ballasting effects of smectite on aggregate formation and export from a natural plankton
 1231 community, *Marine Chemistry*, 175, 18-27, <http://dx.doi.org/10.1016/j.marchem.2015.04.009>, 2015.
- 1232 Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A.,
 1233 Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I.,
 1234 and Torres, R.: Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, *Science*, 308, 67-71,
 1235 2005.
- 1236 Johnson, L. R.: Mineralogical dispersal patterns of North Atlantic deep-sea sediments with particular reference to eolian
 1237 dusts, *Marine Geology*, 29, 335-345, [http://dx.doi.org/10.1016/0025-3227\(79\)90115-4](http://dx.doi.org/10.1016/0025-3227(79)90115-4), 1979.
- 1238 Journet, E., Balkanski, Y., and Harrison, S. P.: A new data set of soil mineralogy for dust-cycle modeling, *Atmos. Chem.*
 1239 *Phys.*, 14, 3801-3816, 10.5194/acp-14-3801-2014, 2014.
- 1240 Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., Jäckel, S., Jaenicke, R., Knippertz, P., Lieke, K., Massling,
 1241 A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass
 1242 concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at
 1243 Tinfou, Morocco, during SAMUM 2006, *Tellus B*, 61, 32-50, 10.1111/j.1600-0889.2008.00385.x, 2009.
- 1244 Khiri, F., Ezaidi, A., and Kabbachi, K.: Dust deposits in Souss-Massa basin, South-West of Morocco: granulometrical,
 1245 mineralogical and geochemical characterisation, *Journal of African Earth Sciences*, 39, 459-464, 2004.
- 1246 Knippertz, P., and Todd, M. C.: Mineral dust aerosols over the Sahara: Meteorological controls on emission and transport and
 1247 implications for modeling, *Rev Geophys*, 50, n/a-n/a, 10.1029/2011RG000362, 2012.
- 1248 Knippertz, P., and Stuut, J.-B. W.: *Mineral dust a key player in the Earth system*, Springer, Dordrecht, 2014.
- 1249 Koch, J., and Renno, N. O.: The role of convective plumes and vortices on the global aerosol budget, *Geophysical Research*
 1250 *Letters*, 32, n/a-n/a, 10.1029/2005GL023420, 2005.
- 1251 Kogbe, C. A.: Geology of the upper cretaceous and tertiary sediments of the Nigerian sector of the Iullemeden Basin (West-
 1252 Africa), *Geol Rundsch*, 62, 197-211, 10.1007/bf01826827, 1973.
- 1253 Koopmann, B.: Sedimentation von Saharastaub im subtropischen Nordatlantik während der letzten 25.000 Jahre, *Meteor*
 1254 *Forschungsergebnisse C*, 35, 23-59, 1981.
- 1255 Korte, L. F., Brummer, G. J., van der Does, M., Guerreiro, C. V., Hennekam, R., van Hateren, J. A., Jong, D., Munday, C. I.,
 1256 Schouten, S., and Stuut, J. B. W.: Downward particle fluxes of biogenic matter and Saharan dust across the equatorial North
 1257 Atlantic, *Atmos. Chem. Phys.*, 17, 6023-6040, 10.5194/acp-17-6023-2017, 2017.

Formatted: Dutch (Netherlands)

- 1258 Lancaster, N.: 11.12 Sand Seas and Dune Fields A2 - Shroder, John F, in: Treatise on Geomorphology, Academic Press, San
1259 Diego, 219-245, 2013.
- 1260 Lange, H.: Herkunft und Verteilung von Oberflächensedimenten des westafrikanischen Schelfs und Kontinentalhanges.⁴,
1261 "Meteor" Forschungsergeb, 22, 61-84, 1975.
- 1262 Lange, H.: Distribution of Chlorite and Kaolinite in Eastern Atlantic Sediments Off North-Africa, *Sedimentology*, 29, 427-
1263 431, DOI 10.1111/j.1365-3091.1982.tb01805.x, 1982.
- 1264 Lau, K. M., and Kim, K. M.: Cooling of the Atlantic by Saharan dust, *Geophys. Res. Lett.*, 34, 10.1029/2007GL031538,
1265 2007.
- 1266 Martin, J. H.: Glacial-interglacial CO₂ change: the iron hypothesis, *Paleoceanography*, 5, 1-13, 1990.
- 1267 Martin, J. H., Gordon, R. M., and Fitzwater, S. E.: The case for iron, *Limnology & Oceanography*, 36, 1793-1802, 1991.
- 1268 McTainsh, G. H., Nickling, W. G., and Lynch, A. W.: Dust deposition and particle size in Mali, West Africa, *Catena*, 29,
1269 307-322, 1997.
- 1270 Mendez, M. J., Funk, R., and Buschiazzo, D. E.: Field wind erosion measurements with Big Spring Number Eight (BSNE)
1271 and Modified Wilson and Cook (MWAC) samplers, *Geomorphology*, 129, 43-48, 10.1016/j.geomorph.2011.01.011, 2011.
- 1272 Meunier, T., Barton, E. D., Barreiro, B., and Torres, R.: Upwelling filaments off Cap Blanc: Interaction of the NW African
1273 upwelling current and the Cape Verde frontal zone eddy field?, *Journal of Geophysical Research: Oceans*, 117, n/a-n/a,
1274 10.1029/2012JC007905, 2012.
- 1275 Meyer, I., Davies, G. R., Vogt, C., Kuhlmann, H., and Stuu, J.-B. W.: Changing rainfall patterns in NW Africa since the
1276 Younger Dryas, *Aeolian Research*, 10, 111-123, <http://dx.doi.org/10.1016/j.aeolia.2013.03.003>, 2013.
- 1277 Mittelstaedt, E.: The ocean boundary along the northwest African coast: Circulation and oceanographic properties at the sea
1278 surface, *Progress In Oceanography*, 26, 307-355, 10.1016/0079-6611(91)90011-A, 1991.
- 1279 Moore, D. M., and Reynolds, R. C.: X-ray Diffraction and the Identification and Analysis of Clay Minerals, Oxford
1280 university press Oxford, 1989.
- 1281 Moreno, T., Querol, X., Castillo, S., Alastuey, A., Cuevas, E., Herrmann, L., Mounkaila, M., Elvira, J., and Gibbons, W.:
1282 Geochemical variations in aeolian mineral particles from the Sahara-Sahel Dust Corridor, *Chemosphere*, 65, 261-270, 2006.
- 1283 Mulitza, S., Heslop, D., Pittauerova, D., Fischer, H. W., Meyer, I., Stuu, J.-B., Zabel, M., Mollenhauer, G., Collins, J. A.,
1284 Kuhnert, H., and Schulz, M.: Increase in African dust flux at the onset of commercial agriculture in the Sahel region, *Nature*,
1285 466, 226-228, 10.1038/nature09213, 2010.
- 1286 National Geospatial-Intelligence Agency: North Atlantic, Baltic Sea, North Sea and Mediterranean Sea, 4 ed., Sailing
1287 directions (planning guide), ProStar Publications, 2006.
- 1288 Nicholson, S. E.: A revised picture of the structure of the "monsoon" and land ITCZ over West Africa, *Climate Dynamics*,
1289 32, 1155-1171, DOI 10.1007/s00382-008-0514-3, 2009.

Formatted: Dutch (Netherlands)

- 1290 Nizou, J., Hanebuth, T. J. J., and Vogt, C.: Deciphering signals of late Holocene fluvial and aeolian supply from a shelf
1291 sediment depocentre off Senegal (north-west Africa), *Journal of Quaternary Science*, n/a-n/a, 10.1002/jqs.1467, 2011.
- 1292 Piqué, A.: *Geology of northwest Africa*, Gebrüder Borntraeger, 2001.
- 1293 Ploug, H., Iversen, M. H., and Fischer, G.: Ballast, sinking velocity, and apparent diffusivity within marine snow and
1294 zooplankton fecal pellets: Implications for substrate turnover by attached bacteria, *Limnology and Oceanography*, 53, 1878-
1295 1886, 2008a.
- 1296 Ploug, H., Iversen, M. H., Koski, M., and Buitenhuis, E. T.: Production, oxygen respiration rates, and sinking velocity of
1297 copepod fecal pellets: Direct measurements of ballasting by opal and calcite, *Limnology and Oceanography*, 53, 469-476,
1298 DOI 10.4319/lo.2008.53.2.0469, 2008b.
- 1299 Proske, U., Hanebuth, T. J. J., Meggers, H., and Leroy, S. A. G.: Tidal flat sedimentation during the last millennium in the
1300 northern area of Tidra Island, Banc d'Arguin, Mauritania, *Journal of African Earth Sciences*, 50, 37-48,
1301 <http://dx.doi.org/10.1016/j.jafrearsci.2007.09.002>, 2008.
- 1302 Prospero, J. M., and Carlson, T. N.: Radon-222 in North Atlantic Trade Winds . Its Relationship to Dust Transport from
1303 Africa, *Science*, 167, 974-&, Doi 10.1126/Science.167.3920.974, 1970.
- 1304 Prospero, J. M., and Carlson, T. N.: Vertical and areal distribution of Saharan dust over the western equatorial north Atlantic
1305 Ocean, *Journal of Geophysical Research*, 77, 5255-5265, 10.1029/JC077i027p05255, 1972.
- 1306 Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental characterization of global sources of
1307 atmospheric soil dust identified with the Nimbus 7 total ozone mapping spectrometer (TOMS) absorbing aerosol product,
1308 *Rev Geophys*, 40, 1-31, 2002.
- 1309 Pye, K.: The nature, origin and accumulation of loess, *Quaternary Science Reviews*, 14, 653-667, 1995.
- 1310 Radczewski, O. E.: *Eolian deposits in marine sediments*, 1939.
- 1311 Rateev, M. A., Gorbunova, Z. N., Lisitzyn, A. P., and Nosov, G. L.: THE DISTRIBUTION OF CLAY MINERALS IN THE
1312 OCEANS, *Sedimentology*, 13, 21-43, 10.1111/j.1365-3091.1969.tb01119.x, 1969.
- 1313 Ratmeyer, V., Balzer, W., Bergametti, G., Chiapello, I., Fischer, G., and Wyputta, U.: Seasonal impact of mineral dust on
1314 deep-ocean particle flux in the eastern subtropical Atlantic Ocean, *Marine Geology*, 159, 241-252, 1999a.
- 1315 Ratmeyer, V., Fischer, G., and Wefer, G.: Lithogenic particle fluxes and grain size distributions in the deep ocean off
1316 northwest Africa: Implications for seasonal changes of aeolian dust input and downward transport, *Deep Sea Research Part I:
1317 Oceanographic Research Papers*, 46, 1289-1337, 1999b.
- 1318 Rea, D. K.: The paleoclimatic record provided by eolian deposition in the deep sea: the geologic history of wind, *Rev
1319 Geophys*, 32, 159-195, 1994.
- 1320 Santamarina, J., and Cho, G.: Soil behaviour: The role of particle shape, *Advances in geotechnical engineering: The
1321 skempton conference*, 2004, 604-617,

- 1322 Scheuvens, D., Schütz, L., Kandler, K., Ebert, M., and Weinbruch, S.: Bulk composition of northern African dust and its
 1323 source sediments — A compilation, *Earth-Science Reviews*, 116, 170-194, <http://dx.doi.org/10.1016/j.earscirev.2012.08.005>,
 1324 2013.
- 1325 Schlüter, T.: *Geological atlas of Africa*, Springer, 2008.
- 1326 Schofield, D., Horstwood, M., Pitfield, P., Crowley, Q., Wilkinson, A., and Sidaty, H. C. O.: Timing and kinematics of
 1327 Eburnean tectonics in the central Reguibat Shield, Mauritania, *Journal of the Geological Society*, 163, 549-560, 2006.
- 1328 Schuster, M., Düringer, P., Ghienne, J.-F., Roquin, C., Sepulchre, P., Moussa, A., Lebatard, A.-E., Mackaye, H. T., Likius,
 1329 A., Vignaud, P., and Brunet, M.: Chad Basin: Paleoenvironments of the Sahara since the Late Miocene, *Comptes Rendus*
 1330 *Geoscience*, 341, 603-611, <http://dx.doi.org/10.1016/j.crte.2009.04.001>, 2009.
- 1331 Schütz, L., and Sebert, M.: Mineral aerosols and source identification, *Journal of Aerosol Science*, 18, 1-10,
 1332 [http://dx.doi.org/10.1016/0021-8502\(87\)90002-4](http://dx.doi.org/10.1016/0021-8502(87)90002-4), 1987.
- 1333 Selley, R. C.: Chapter 3 The sirte basin of libya, in: *Sedimentary Basins of the World*, edited by: Selley, R. C., Elsevier, 27-
 1334 37, 1997a.
- 1335 Selley, R. C.: Chapter 1 The sedimentary basins of northwest africa: stratigraphy and sedimentation, in: *Sedimentary Basins*
 1336 *of the World*, edited by: Selley, R. C., Elsevier, 3-16, 1997b.
- 1337 Selley, R. C.: Chapter 2 The basins of northwest africa: Structural evolution, in: *Sedimentary Basins of the World*, edited by:
 1338 Selley, R. C., Elsevier, 17-26, 1997c.
- 1339 Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Stuut, J. B., Meyer, I.,
 1340 Chiapello, I., Podvin, T., Chatenet, B., Diallo, A., and Ndiaye, T.: The 7-13 March 2006 major Saharan outbreak: Multiproxy
 1341 characterization of mineral dust deposited on the West African margin, *J. Geophys. Res.*, 116, D18210,
 1342 10.1029/2011jd016173, 2011.
- 1343 Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Diallo, A., and Ndiaye, T.: A
 1344 three-year time series of mineral dust deposits on the West African margin: Sedimentological and geochemical signatures and
 1345 implications for interpretation of marine paleo-dust records, *Earth and Planetary Science Letters*, 364, 145-156,
 1346 <http://dx.doi.org/10.1016/j.epsl.2012.12.039>, 2013.
- 1347 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric
 1348 Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 10.1175/bams-
 1349 d-14-00110.1, 2015.
- 1350 Stein, R.: Late neogene changes of paleoclimate and paleoproductivity off northwest africa (D.S.D.P. Site 397),
 1351 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 49, 47-59, [http://dx.doi.org/10.1016/0031-0182\(85\)90004-5](http://dx.doi.org/10.1016/0031-0182(85)90004-5), 1985.
- 1352 Stuut, J.-B. W.: Late Quaternary Southwestern African terrestrial-climate signals in the marine record of Walvis Ridge, SE
 1353 Atlantic Ocean, Faculty of Earth Sciences, Utrecht University, Utrecht, 128 pp., 2001.
- 1354 Stuut, J.-B. W., Zabel, M., Ratmeyer, V., Helmke, P., Schefuß, E., Lavik, G., and Schneider, R. R.: Provenance of present-
 1355 day eolian dust collected off NW Africa, *Journal of Geophysical Research*, 110, 10.1029/2004JD005161, 2005.

- 1356 Stuut, J.-B. W., Smalley, I., and O'Hara-Dhand, K.: Aeolian dust in Europe: African sources and European deposits,
1357 *Quaternary International*, 198, 234-245, 10.1016/j.quaint.2008.10.007, 2009.
- 1358 Stuut, J.-B. W., Bakker, M., Friese, C., Koster, B., Visser, J.-D. d., and Witte, Y.: Cruise Report and preliminary results -
1359 DUSTTRAFFIC: Transatlantic fluxes of Saharan dust - Cruise No. 64PE392 - 19 – 27 August 2014 Las Palmas de Gran
1360 Canaria (Spain) – Mindelo, Sao Vicente (Cape Verdian Islands), 2015.
- 1361 Temon, E., Guieu, C., Loÿe-Pilot, M. D., Leblond, N., Bosc, E., Gasser, B., Miquel, J. C., and Martín, J.: The impact of
1362 Saharan dust on the particulate export in the water column of the North Western Mediterranean Sea, *Biogeosciences*, 7, 809-
1363 826, 10.5194/bg-7-809-2010, 2010.
- 1364 Tjallingii, R., Claussen, M., Stuut, J.-B. W., Fohlmeister, J., Jahn, A., Bickert, T., Lamy, F., and Rohl, U.: Coherent high- and
1365 low-latitude control of the northwest African hydrological balance, *Nature Geoscience*, 1, 670-675, 10.1038/ngeo289, 2008.
- 1366 Tsoar, H., and Pye, K.: Dust transport and the question of desert loess formation, *Sedimentology*, 34, 139-153, 1987.
- 1367 Tucker, M. E., and Tucker, M.: *Techniques in sedimentology*, 552.5. 08 TEC, 1988.
- 1368 Van Camp, L., Nykjaer, L., Mittelstaedt, E., and Schlittenhardt, P.: Upwelling and boundary circulation off Northwest Africa
1369 as depicted by infrared and visible satellite observations, *Progress in Oceanography*, 26, 357-402,
1370 [http://dx.doi.org/10.1016/0079-6611\(91\)90012-B](http://dx.doi.org/10.1016/0079-6611(91)90012-B), 1991.
- 1371 Van der Does, M., Korte, L. F., Munday, C. I., Brummer, G. J. A., and Stuut, J. B. W.: Particle size traces modern Saharan
1372 dust transport and deposition across the equatorial North Atlantic, *Atmos. Chem. Phys.*, 16, 13697-13710, 10.5194/acp-16-
1373 13697-2016, 2016.
- 1374 Verati, C., Bertrand, H., and Féraud, G.: The farthest record of the Central Atlantic Magmatic Province into West Africa
1375 craton: Precise ⁴⁰Ar/³⁹Ar dating and geochemistry of Taoudenni basin intrusives (northern Mali), *Earth and Planetary
1376 Science Letters*, 235, 391-407, <http://dx.doi.org/10.1016/j.epsl.2005.04.012>, 2005.
- 1377 Villeneuve, M.: Paleozoic basins in West Africa and the Mauritanide thrust belt, *Journal of African Earth Sciences*, 43, 166-
1378 195, <http://dx.doi.org/10.1016/j.jafrearsci.2005.07.012>, 2005.
- 1379 Vogt, C., Lauterjung, J., and Fischer, R. X.: Investigation of the Clay Fraction (< 2 µm) of the Clay Minerals Society
1380 Reference Clays, *Clays and Clay Minerals*, 50, 388-400, 2002.
- 1381 Wilson, S., and Cooke, R.: Wind erosion, Soil erosion, 217251, 1980.
- 1382 Wissmann, G.: Stratigraphy and structural features of the continental margin basin of Senegal and Mauritania, in: *Geology of
1383 the Northwest African continental margin*, Springer, 160-181, 1982.
- 1384 Yoshioka, M., Mahowald, N. M., Conley, A. J., Collins, W. D., Fillmore, D. W., Zender, C. S., and Coleman, D. B.: Impact
1385 of Desert Dust Radiative Forcing on Sahel Precipitation: Relative Importance of Dust Compared to Sea Surface Temperature
1386 Variations, Vegetation Changes, and Greenhouse Gas Warming, *Journal of Climate*, 20, 1445-1467,
1387 doi:10.1175/JCLI4056.1, 2007.
- 1388 Yu, H. B., Chin, M., Bian, H. S., Yuan, T. L., Prospero, J. M., Omar, A. H., Remer, L. A., Winker, D. M., Yang, Y. K.,
1389 Zhang, Y., and Zhang, Z. B.: Quantification of trans-Atlantic dust transport from seven-year (2007-2013) record of
1390 CALIPSO lidar measurements, *Remote Sens Environ*, 159, 232-249, 10.1016/j.rse.2014.12.010, 2015.

1391 Zobeck, T. M., Sterk, G., Funk, R., Rajot, J. L., Stout, J. E., and Van Pelt, R. S.: Measurement and data analysis methods for
1392 field-scale wind erosion studies and model validation, *Earth Surface Processes and Landforms*, 28, 1163-1188,
1393 10.1002/esp.1033, 2003.

1394