

Journal: ACP
MS No.: acp-2017-150
Title: Sources of Submicrometre Particles Near a Major International Airport
Author(s): Mauro Masiol et al.

RESPONSE TO REVIEWERS

REFEREE #1

General comments:

This paper presents a study of different components of air pollution at London Heathrow Airport in two periods covering warm and cold environmental conditions. The authors report measurements of both particulate and gaseous pollutants and use k-means clustering and positive matrix factorisation in an attempt to apportion measured pollution to emissions sources and processes.

The dataset is extensive, the measurements are reliable and the analysis methods are appropriate. However, the presentation of the study is poor and therefore, the novel contribution of the paper is unclear. Below are major comments on the paper, after which follow specific line-by-line comments.

The paper has a number of weaknesses, and my opinion is that major revisions are required before it is accepted for publication.

Reply: We would like to thank referee #1 for his/her useful comments. All the points have been considered in this revised version of our manuscript. In particular, the paper was shortened and a number of references were added to the "Introduction" section or moved from other sections (according to the specific referee's comments). The novel contribution of this study with respect to the available literature is now explained and discussed in the "Introduction" section.

We have organised our replies by highlighting the referee comments in bold-italic font; our reply is in normal font.

1. The literature review is incomplete and must be significantly improved. There have been numerous studies investigating the increased concentrations of UFPs close to airports, which must be included in the introduction. Some of these have been referenced in the discussion of results but inclusion of these references in Introduction is required to place this paper in the context of others and define the novel contribution.

34 **Reply:** The “Introduction” section has been substantially amended. We have moved a whole
35 paragraph including a number of up-to-date references from “Results” to “Introduction”. We have
36 also added several other references within the manuscript. We are aware that many other related
37 studies are available in the literature. However, we have cited those that might be more relevant to
38 the goals of our study. Our aim is that the current version of the “introduction” provides an
39 adequate degree of information and allows the reader to understand the state-of-the-art on the
40 topic.

41 In addition, some sentences were added to the “Introduction” to point out the novel contribution of
42 the paper with respect to the available literature. We are confident that our paper represents a
43 novel contribution to the science: the literature offers many studies on the PNC and size
44 distributions measured close or within airports. However, there are few papers using sophisticated
45 chemometric tools to quantify and characterize the airport contribution to UFPs. Air quality close to
46 large airports is potentially affected by many sources. The application of cluster and PMF analyses
47 (and the comparison of their results) allows successful extraction of the profiles of individual
48 sources. The comparison of the two methods (Section 3.3) allows the reader to understand the
49 potentials and limits of each of these source apportionment techniques.

50

51 **2. The discussion of results is not well-structured, statements are not quantified**
52 **sufficiently, and explicit references to figures are not included.**

53 **Reply:** The discussion of results has been extensively improved. Statements are now quantified.
54 Explicit references to figures and/or tables are added throughout the manuscript. We have also
55 shortened the sections related to the cluster analysis (see the referee's points 3 and 4 below).

56

57 **3. The section on the results from the k-means clustering (3.2) does not make any definitive**
58 **conclusions, and it is limited in its contribution to novel science. Most of the results are**
59 **justified by existing literature. Since that seems to be the case, the section is much longer**
60 **than is appropriate for reporting routine results. Discussion should be limited to novel**
61 **results, other results could be discussed in the SI.**

62 **Reply:** Cluster analysis extracts the most common modal spectra. It is not a novel science and it
63 has been extensively applied to PNSD, particularly in London. However, we would like to stress
64 that cluster analysis was never previously applied to 5 min resolved PNSDs in environments
65 potentially affected by airport emissions. It represents the novel application of a well consolidated
66 tool (clustering) to a specific case study (airports). In this way, we believe that its application is
67 important for this study and generates new science useful for the scientific community. In addition,

68 we believe that it is important to report the results of this technique in comparison with PMF
69 analysis (as in manuscript Section 3.3). The PNSD spectra in airport-affected environments are
70 very complex. Cluster analysis offers some simplification but was unable to resolve some of the
71 sources.

72 We agree that the discussion of the cluster analysis results is long. However, we also believe that
73 the cluster analysis gives important information and needs to be carefully discussed in the main
74 text. Following the advice of the referee, the discussion of the cluster results was improved and
75 shortened by about one third. In particular, we have merged the discussion of the warm and cold
76 season results and have limited the main text to the main findings. We have also removed some
77 misleading sentences and some statements not well supported by the data (as pointed out by the
78 other referee). This makes the text shorter and has streamlined the discussion.

79

80 ***4. The discussion of results from the k-mean clustering and PMF analysis is repetitive and***
81 ***many of the same references are used to infer the sources of particular clusters and***
82 ***factors. I would suggest that the discussion of these clusters and factors is combined in***
83 ***order to draw out stronger conclusions from the results (given the discussion in Section 3.3***
84 ***noting good agreement between the two methods in identifying particular source***
85 ***signatures). The paper would be significantly improved by removing repetition within and***
86 ***the length of the Results section.***

87 ***Reply:*** As explained in the previous response to points 2 and 3, we have improved the sections on
88 the cluster analysis. However, these two techniques are very different (as discussed in the text)
89 and generate very different insights. Consequently, we believe that the sections cannot usefully be
90 merged. On the contrary, we prefer to consider the outcomes of each method in turn and then
91 consider what the combined data tells us about particle sources. We would like to point out that
92 referee #2 does not criticise this part. Essentially, we found that in a location with so many particle
93 source influences, clusters will never represent single source categories, whereas a well designed
94 PMF is able to better extract the profiles of individual sources.

95

96 ***5. The paper is overly long and the results are not presented in a concise or coherent***
97 ***manner. There are several instances of repetition.***

98 ***Reply:*** Several sections of the manuscript have been carefully revised according to the comments
99 of the two referees. The "Introduction" section was improved with additional references; the
100 discussion of ancillary variables (particularly for NO_x and O₃) in the Section 3.1 "Overview of Data"
101 was removed; two figures were moved to SI, while other figures were modified following advice

102 from the referees; some statements in the “Results” were moved to the “Introduction”; the section
103 on the results of the cluster analysis was shortened by ~30% by merging the results of the two
104 seasons; statements were quantified (where possible) and references to figures and tables have
105 been added throughout the text; we have removed several instances of repetition, errors/mistakes
106 and misleading sentences.

107 Therefore, the manuscript underwent an overall shortening and improvement, particularly in the
108 discussion of the cluster analysis results. We believe that the manuscript is now well balanced and
109 the discussion streamlined.

110 We recognise that the manuscript is long. However, we would like to stress that most of the
111 literature offers studies based on the application of only one source apportionment method. This
112 study has taken advantage of the synergy of the two methods (clustering and PMF) and their
113 comparison. We believe that the careful discussion of outcomes from both the methods is useful
114 and interesting for the scientific community and, therefore, needs to be preserved.

115
116 ***Specific comments:***

117 ***Line 57: The statement that aviation growth will continue for the next decade cites a study***
118 ***from 8 years ago. Please use a more up to date reference.***

119 **Reply:** Up-to-date reference added. The estimation of the current trend was recently updated by
120 the ICAO and is approx. 5.5%/y.

121

122 ***Line 69: No reference for the ‘indisputable’ role of LHR in driving economic affluence and***
123 ***vitality is given.***

124 **Reply:** Corrected. The whole paragraph was revised and shortened, according to the comments of
125 the referee #2. In particular, the sentences related to the debate over the expansion of London
126 Heathrow have been dropped off.

127

128

129 ***Line 75: Reference(s) for arguments in support of LHR expansion?***

130 **Reply:** Corrected. The whole paragraph was revised and shortened, according to the comments of
131 the referee #2. In particular, the sentences related to the debate over the expansion of London
132 Heathrow have been dropped off.

133

134 **Line 82: Reference for Government approval of 3rd runway?**

135 **Reply:** A reference from the UK Department for Transport has been added.

136

137 **Line 106: Suggest present tense.**

138 **Reply:** Done.

139

140 **Line 108-117: Suggest description of methodology is moved to Section 2.**

141 **Reply:** This paragraph has been modified and shortened. Here, we only present the goals of the
142 study and briefly list the adopted source apportionment methods. In addition, we have added two
143 paragraphs pointing out the novelty of the study compared to the current literature and the
144 secondary goal related to the analysis of the effects of a regional nucleation event.

145

146 **Line 119-121: Move to Acknowledgements.**

147 **Reply:** Done. "Classical" substituted with "Routine". These are pollutants routinely measured at
148 DEFRA air quality sites.

149

150 **Line 150: Repetition.**

151 **Reply:** Done. The whole Materials and Methods section has been revised. In particular, the
152 original subsection 2.1 was modified and moved to the "Introduction". All the details about the
153 sampling campaigns are now in the new subsection 2.1 "Experimental".

154

155 **Line 165: Not all traffic is generated by the airport. Can the proportion be quantified?**

156 **Reply:** Sentence modified accordingly. Unfortunately, this is not quantifiable in this study. Our
157 previous investigations (Masiol and Harrison, AE116, 2015) pointed out that the study of the
158 differences (deltas) among multiple sampling sites across the study area is very useful to apportion
159 the different sources, including traffic. However, just one site was used in this study.

160 The amended text: "The site is also affected by pollutants arising from the large volumes of road
161 traffic in London, from the local road network as well as those generated by the airport. Tunnel Rd.,
162 the main access to LHR from the M4 motorway lies 800 m west, as well as the nearby M4 (640 m
163 north) and M25 (~3.5 km east) motorways, major roads (Bath Rd, part of A4, passes 900 m south;
164 A30 lies 2.8 km SE). The village of Harlington (~400 m west) and advection of air masses from the
165 conurbation of London are other potential external sources."

166 **Line 196: Suggest delete 'Classical', rephrase.**

167 **Reply:** Done.

168

169 **Line 223: Clarify the reason for deletion of data greater than 99.5th percentile. Is this for all**
170 **measured quantities?**

171 **Reply:** Only SMPS data have been handled in this way. Other data from the Harlington site are
172 already checked and ratified by DEFRA. We have added some sentences explaining our choice.
173 Essentially, we recognised some extreme events, which have been interpreted as outliers and/or
174 instrumental errors. Most of them were linked to unusual activities or probable instrumental issues.
175 Anyway, the modal structure of the PMF profiles with or without the removed data does not
176 change. The main reasons driving this choice were: (i) a general improvement in the stability of
177 PMF solutions; (ii) the decrease of uncertainty assessed by BS and DISP; and (iii) the decrease of
178 scaled residuals over ± 3 , which, essentially, are records not well modeled by the PMF.

179 The amended text: "An in-depth analysis of the dataset revealed a few records with anomalously
180 high PNC, which were likely related to probable instrumental issues, extreme weather conditions
181 (e.g., high wind gusts, heavy rain striking the inlet), or infrequent local emissions, e.g.,
182 maintenance, painting and recreational activities (including fires) on the playground where the site
183 is located, road maintenance close the site and probable short-term parking of high-emission
184 vehicles near the site. Since this study aims to investigate the overall contributions of LHR, all data
185 are used for descriptive statistics, but data greater than the 99.5th percentile were further removed
186 for explorative, cluster and PMF analyses. This data exclusion successfully removed the extremely
187 high events occurring during the sampling campaigns and significantly improved the stability and
188 physical meaning of PMF solutions."

189

190 **Line 261: Figures 1 and 2 are not adequately described in the text. If the data is not worth**
191 **mentioning in the text, the figures should be moved to the SI.**

192 **Reply:** Figure 1 (maps of area and sampling site) has been moved to SI (Figure SI1). Figure 2
193 (boxplots and diurnal patterns) was amended: now only some plots of part B (diurnal patterns) are
194 shown in the main text, while part A (boxplots) was moved to SI (Figure SI3) as well as the plots of
195 all the diurnal patterns (Figure SI4). We want to maintain some plots of the part B in the main text:
196 we believe that showing the diurnal patterns of such important variables (PNC over the three size
197 ranges, BC, solar irradiation and airport traffic) is really useful to the reader for helping the
198 interpretation of results. These patterns can be, therefore, quickly compared with those reported for
199 the clusters or the PMF factors.

200

201 **Line 288: This paragraph would be more appropriate in the Introduction.**

202 **Reply:** Done. This part was moved to the “Introduction” (also according to the main point 1 and the
203 referee #2).

204

205 **Line 312: Clarify meaning of ‘intensive sampling’.**

206 **Reply:** Amended: “intensive” removed from the text.

207

208 **Line 330: Rephrase sentence ‘Airport traffic undergoes: : :’**

209 **Reply:** Sentence modified accordingly: “During nighttime, airport traffic is restricted to limit noise
210 and community disturbance: ...”.

211

212 **Line 340: The statement that nucleation particle concentrations ‘drop to near zero**
213 **overnight’ is not substantiated by Figure 3. Statements discussing results must be**
214 **quantified.**

215 **Reply:** Amended. The whole paragraph was amended and the discussion of the patterns is now
216 quantified.

217

218 **Section 3.1: References to figures should be made to aid interpretation.**

219 **Reply:** Amended. The whole text was carefully revised to link sentences and discussion to the
220 appropriate figures and tables.

221

222 **Line 344: From Figure 2 it is not clear that accumulation mode concentrations have a peak**
223 **corresponding to the morning rush hour.**

224 **Reply:** The morning peak is only evident during the warm season. The sentences have been
225 amended consequently: "Accumulation particles also present the morning (6-8 am) and evening (6-
226 11 pm) rush hour peaks during the warm season, but only the evening peak (from 6 pm to the
227 night) was found in the cold season (Figure 2). Generally, the evening peaks start around 6 pm,
228 which is consistent with the peak of traffic (Figure SI5) but they extend late in the evening and night
229 probably because the drop of the mixing layer top and the consequent concentration of pollutants
230 close to the ground level."

231

232 **Line 421-425: Repetition, rephrase.**

233 **Reply:** The discussion of the cluster analysis results was strongly modified and shortened (see
234 reply to the referee's main points 2-5). We believe that there are no repetitions in its current form.

235

236 **Line 471: Clarify which clusters are being compared, 'their' is not sufficient. The**
237 **comparisons between cluster 1 and cluster 5 are confused by the reference to the clusters**
238 **from the warm period.**

239 **Reply:** Done.

240

241 **Line 482: What is the basis for this interpretation? This seems to be pure speculation.**

242 **Reply:** We have removed the sentences and we have improved the further discussion of the two
243 clusters in Section 3.3 (comparison of cluster analysis and PMF). This was added to the text as:
244 "The reasons driving the split of the spectra likely to be affected by LHR into two clusters during the
245 cold season are unclear. A further comparison of the cluster and PMF results will help in
246 interpreting this outcome."

247

248 **Line 498: rephrase**

249 **Reply:** Done. The whole section has been enhanced.

250

251 **Line 514: What is Q?**

252 **Reply:** It is the PMF objective function. In PMF, factor elements are constrained, therefore no
253 sample can have a significantly negative contribution. PMF allows each data value to be
254 individually weighted. This feature allows analysts to adjust the influence of each data point,
255 depending on the confidence in the measurement. The PMF solution minimizes the object function
256 Q via a conjugate gradient algorithm, based upon the estimated data uncertainties (or adjusted
257 data uncertainties). A quick review of the PMF and how the objective function works is reported in
258 Brown et al. (STOTEN 518-519, 2015) or in the PMF user's manual. Factor contributions and
259 profiles are therefore derived by the PMF model minimizing the objective function (Q).

260 This paragraph was moved to the "Materials and Method" section (subsection 2.2), according to
261 the following point of this referee. We have only included its name "objective function" into the text.
262 We did not provide much detail on the science behind the PMF analysis, as it is a very well
263 established technique. However, we noticed that most of the literature on PMF does not provide
264 sufficient data to describe how the model was run and what diagnostics have been taken into
265 account. In our opinion, this is a serious omission, as it is always important to know if the model is
266 performed with the optimal criteria and if the model returns acceptable diagnostics. Consequently,
267 we want to keep this (and other) brief technical information in the main text to provide all the
268 information needed by the readers/PMF analysts who want to fully reproduce/check the method.
269 This information is comparable to a QC/QA applied to the PMF analysis.

270

271 **Line 513-530: These methodological details should be in the Methods section.**

272 **Reply:** Done. Moved to "Materials and Methods" (subsection 2.2).

273

274 **Line 765: Rephrase and quantify 'well agree'.**

275 **Reply:** "well agree" was deleted. The normalised contributions are now quantified in the text and
276 some statistical tests have been also applied in the discussion of results.

277 The new paragraph is: "For the warm period, significantly higher (0.05 significance) PMF
278 contributions of the airport factor (F1) are measured for Cluster 1 (average normalised contribution
279 ~3.5). This result indicates that the airport fingerprint was well captured by both source
280 apportionment methods. During the cold season, the airport factor (F1) is significantly higher for
281 both clusters 1 and 5 (average normalised contributions of ~2 and ~3, respectively). While Cluster
282 5 presents significant high PMF contributions only for factor 1, Cluster 1 also shows high
283 contributions of factor 2 (fresh road traffic). This result indicates that Cluster 5 may be linked as the

284 typical PNSD spectra for airport emissions, while Cluster 2 likely represents mixed emissions from
285 aircraft and airport-related traffic. A close analysis of wind roses for the two clusters in the cold
286 season (Figure 4) reveals that Cluster 5 occurs at significantly higher wind speed regimes than
287 Cluster 1 (Mann-Whitney-Wilcoxon test at 0.05 significance level), i.e. average wind speeds of 8.3
288 and 5.9 m s⁻¹, respectively. As a consequence, the different wind regimes may be likely
289 responsible of the split between the two clusters.”.

290

291 **Line 788-796: Literature review should be in the Introduction.**

292 **Reply:** Done. The whole paragraph was moved to the “Introduction”.

293

294 **Line 813: Quantify ‘fast drop’.**

295 **Reply:** Quantified. The drop was -30 µg/m³ in 3 hours.

296 Amended sentence: “This is also supported by the PM_{2.5} mass, which exhibited a rapid fall in
297 concentrations just a few hours before the event (-30 µg m⁻³ in 3 hours, i.e. from 40 µg m⁻³ at 6 am
298 to 10 µg m⁻³ at 9 am, Figure 9), probably reducing the condensation sink and facilitating
299 nucleation.”.

300

301 **Line 822: Quantify ‘just slightly affected’.**

302 **Reply:** Quantified. The whole paragraph has been re-written and now presents a large number of
303 quantified statements and explicit references to figures.

304 Amended paragraph: “The results of cluster analysis were affected by the event. Before the
305 episode, the PNSD spectra were mostly categorised as Clusters 3 and 4 (urban background and
306 daytime pollution, respectively), i.e. the clusters mostly recorded under north-easterly wind regimes
307 (Figure 3). About 50% and 30% of the clusters were then categorised as “airport” in the first and
308 second hour of the episode, respectively (Figure 9). Since the wind directionality is inconsistent
309 with an origin from the airfield, this categorisation is likely the result of the nucleation event. The
310 growth of particles in the hours after the beginning of the event has further driven the cluster
311 results: (i) about 60-80% of PNSDs were categorised as “fresh road traffic” (Cluster 5) after 2-3
312 hours, and (ii) 80-100% of PNSDs were clustered as “nighttime regional pollution” (Cluster 2) after
313 4-6 hours. In a similar way, PMF results were affected by the event (Figure 9), with a sharp
314 increase of contribution levels for: (i) factor 1 (airport) from 1.5 x 10³ particles cm⁻³ at noon to 13.3
315 x 10³ particles cm⁻³ at 2 pm; (ii) factor 2 (fresh road traffic) from 0.5 x 10³ particles cm⁻³ at 1 pm to

316 21 x 10³ particles cm⁻³ at 3 pm; and (iii) factor 3 (aged road traffic) from 2.1 x 10³ particles cm⁻³ at 2
317 pm to approx. 15 x 10³ particles cm⁻³ at 5-6 pm.”.

318

319 **Line 827: Quantify ‘slightly affected’.**

320 **Reply:** Done. See our reply to the previous point.

321

322 **Line 843: ‘Anomalously’ implies that the measurements are flawed in some way. I do not**
323 **believe that this is the case. Rephrase.**

324 **Reply:** Done. “Anomalously” deleted.

325

326

327 **Line 863: Clarify ‘the fingerprint of London’.**

328 **Reply:** Done. We have amended the sentences in the item: “An urban accumulation mode was
329 found. This source presents a wide mode between 50-150 nm and accounts for around 10% of
330 PNC. The wind directionality is consistent with the advection of air masses from London. It is more
331 evident overnight due to the drop of the mixing layer top, the subsequent increase in air pollutants
332 at ground level and the generation of nighttime secondary nitrate aerosols.”.

333

334 **Figure 2: Check units of particle number concentrations – values are lower than typical**
335 **ambient concentrations of 104 part/cm³ and are inconsistent with other figures.**

336 **Reply:** The unit is 10³ particles/cm³. Corrected in Figure 2 and Figure SI4.

337

338 **Figure 10: This could be moved to the SI as discussion of it is very brief. Supplementary**
339 **material: This is not referenced in the main text. It is also lacking any descriptive text and is**
340 **just a collection of figures. This does not provide the reader with accessible or helpful**
341 **information.**

342 **Reply:** Figure 10 (now Figure 9) is important for the discussion. Most of Section 3.4 refers to the
343 plots shown in this figure. It is referenced in the text several times and allows the reader to interpret

344 the effects of the nucleation event upon the cluster and PMF results. On the contrary, we moved
345 Figure 11 to SI (now Figure SI11), as its discussion is very brief.

346 All the tables and figures in the SI section are referenced in the main text.

347

348 **REFEREE #2**

349 *The manuscript describes the results from a monitoring station close to LHR airport and*
350 *uses two different analytical techniques to apportion or explain the particulate matter*
351 *observed at the site. The data was collected during two intensives, nominally a warm and a*
352 *cold season. The two analysis techniques are used to estimate the influence of the airport*
353 *on the local particulate concentrations. It is clear the authors have analysed the data in*
354 *detail and the use of the two techniques to interpret the data is good, as well as*
355 *acknowledging the limitations.*

356 *Overall, I think the work is suitable for publication. I have 3 main suggestions/questions and*
357 *some detailed comments below.*

358 Reply: We would like to thank anonymous referee #2 for his/her very useful suggestions and the
359 general appraisal of the manuscript. Here are our answers. We have replied point-to-point to all
360 the questions.

361 We have organised our replies by highlighting the referee comments in bold-italic font; our reply is
362 in normal font.

363 We note that a second referee has asked us to shorten the manuscript sections related to the
364 cluster analysis. We have therefore merged the discussion of the warm and cold seasons for the
365 cluster analysis and we have summarised the main findings. We are confident that the whole
366 discussion of results is now lighter, streamlined and easier to follow.

367 **Main suggestions/questions:**

368 *1) I do not think the introduction is suitable as it does not provide a good background to the*
369 *manuscript (see details below). In addition, neither the introduction or the abstract mention*
370 *the use of the regional nucleation event, which is really important for understanding a*
371 *potential limitation in the two analysis techniques*

372 Reply: In response to the points raised by both referees, we have extensively improved the
373 "Introduction" section. In particular, we have (i) improved the literature review by adding a number
374 of up-to-date references, (ii) improved the description of the objectives of the study, (iii) added a

375 discussion over the novel contribution of this study with respect to the available literature, and (iv)
376 moved some references and text from the “results and discussion” to the “introduction”.

377 We are confident that the current version of the “Introduction” provides all the information
378 necessary to summarise the state-of-the-art of the research on the PNC pollution close to airports.
379 We also believe that the improved version of the “Introduction” is able to place our study in the
380 context of others and well defines its novel contribution to the science.

381 We have also added some sentences in the abstract and in the introduction presenting and
382 describing the analysis of the strong nucleation event which occurred during the sampling
383 campaign.

384

385 **2) I think a lot of the NO_x and O₃ details and graphs can be removed. The authors**
386 **acknowledge that NO_x has been studied in detail at the site before and the contributions**
387 **from LHR to the site have been published. The paper itself is about the particulates, not air**
388 **pollution in general. This would streamline the manuscript.**

389 **Reply:** Done. The whole subsection 3.1 was shortened and details on the NO_x and O₃ were
390 removed as well as Figure 2a (completely moved to SI). We have also modified the former Figure
391 2b (now Figure 2): now this figure shows only the variables discussed in the main text and
392 important for the following discussion. The diurnal profiles for all the variables (former Figure 2b)
393 has been moved to SI (new Figure SI4).

394

395 **3) What version of AIM was used to analyse the data? If it was version 10, then no action is**
396 **required. If it was version 9 however, then some work is needed. There is a significant**
397 **difference between V9 and V10 in the diffusional correction algorithms. V9 over estimates**
398 **the size of the correction needed and can lead to an over estimation of the nanoparticles,**
399 **especially below 50nm. As the authors are dividing the particles into Aikten and Ultra fine**
400 **modes (and the characteristics of these modes change with season), there is a need to**
401 **check as it could lead to artefacts in the clustering.**

402 **Reply:** We used the version 8.3, whose outputs we confirmed to be identical to the version 9.
403 Despite the differences in the algorithm used for correcting the diffusion losses between v.10 and
404 previous versions, TSI UK have confirmed that since we used a TSI 3080 electrostatic classifier
405 with a long differential mobility analyser (TSI 3081), which were released with AIM v.9 we were
406 correct to use this version of the software. In addition, since previous studies in London are also
407 based on data extracted with those previous versions of AIM, our results are easily comparable
408 with some of the most relevant literature.

409 **Minor question: *Has the Aethelometer data been corrected according to Collaud Coen et al.,***
410 ***2010? I note the reported protocol as defined in Petzold et al., 2015. If not, it needs***
411 ***correcting especially if the authors want to draw BB aerosol information from it.***

412 **Reply:** No. Data are corrected using the WUAQL AethDataMasher V7.1 to perform data validation
413 and correct data for non-linear loading effects. We cannot apply any further correction to BC data
414 as no reference instrument was run. Anyway, we would like to point out that this study focuses on
415 the PNSDs and black carbon data are only used as ancillary variables to be related with the
416 particle sources. In addition, we would like to report that a large proportion of the black carbon data
417 in the literature are derived from aethalometers with no more than a loading correction.

418 A sentence was added to the text: “eBC was derived from the absorbance at 880 nm wavelength;
419 raw data were post-processed with the Washington University Air Quality Lab AethDataMasher
420 V7.1 to perform data validation and correct data for non-linear loading effects (Virkkula et al., 2007;
421 Turner et al., 2007).”.

422

423 **Details:**

424 ***I question the relevance of some of the introduction to the manuscript. Lines 75 to 83 detail***
425 ***the pros and cons of a third runway, which I do not think adds to the manuscript as the***
426 ***study is about the impacts of Heathrow. It is fine to say the airport is planning to increase***
427 ***capacity, but as to why is not for this manuscript. In line 82, the use of the term ‘Despite***
428 ***this’ could be misinterpreted as questioning the UK government’s decision.***

429 **Reply:** A large number of sentences from this paragraph have been removed, as also
430 recommended by the comments of another referee. As a consequence “despite this” was also
431 deleted. The sentence on the future planned expansion of LHR has now a governmental reference
432 (UK Dept. for Transport).

433

434 ***Lines 85 – 104 Again, I think the context of this paragraph is not appropriate for the***
435 ***manuscript. The manuscript is about characterising the particulates close to Heathrow and***
436 ***not about questioning EU air quality standards as there is nothing in the manuscript that***
437 ***takes this data and compares mortality or morbidity predicted from these measurements***
438 ***with those based on the EU standard method, for example. This section should be that***
439 ***despite the UK meeting the PM standard, it has been show UFP have health issues etc and***
440 ***because there is no network of measurements, campaigns such as this are key to***
441 ***assessing the sources and potential impacts of airports on the UFP. The last sentence***
442 ***should be removed from this section. As the authors point out (line 98), there is limited***

443 **knowledge in this area, so what should be regulated and how? EU has standards on**
444 **particulates from cars. New aircraft will soon be regulated on particulate number and mass.**
445 **EU law requires restricting exposure to nano-particles in the work place. The UK has the**
446 **clean air act. So there are policies in place, but more info is needed before guidelines and**
447 **strategies can be applied.**

448 **Reply:** The section has been comprehensively revised and shortened according to the referees'
449 comments.

450

451 **Section 2.1 – study area and dates. With the exception of the first sentence, this section**
452 **should be in the introduction. It is background and context material.**

453 **Reply:** Done.

454 **Lines 150-151, no need to repeat the dates here as they were just in the previous section.**

455 **Reply:** Done.

456

457 **Line 151-152, data from the site is QA etc. Which data? The authors or the data from the**
458 **AURN site?**

459 **Reply:** AURN site. This sentence was removed. We refer to the QA of DEFRA data later in Section
460 2.1.

461

462 **Lines 288 – 303, this reads like an introduction and is better suited in the introduction.**

463 **Reply:** Done. Moved to "Introduction", as also recommended by Referee #1.

464

465 **Section starting at line 309. This is not an acceptable way to present data and I don't see**
466 **what ranking them in order shows. This also seems to repeat a lot of information already**
467 **contained within figure 2a. Suggest removing most of that section and simply referring to**
468 **fig 2a, M&H 2015 and that the data is representative.**

469 **Reply:** Done. The "ranking" was removed. The whole section was also shortened and improved.
470 Some minor typing errors were also found and corrected.

471

472 ***Line 318, Suggest changing ‘Since’ to ‘Analysis of the data showed it was not: : ...’ and***
473 ***provide a reference for the Kruskal-Wallis analysis.***

474 **Reply:** Done: sentence modified and reference added.

475

476 ***Line 324 and figure 3. Firstly, if the PNSD are shifting towards coarser modes, how can the***
477 ***PVSD be almost constant? The author’s choice of wording is contradictory. Secondly, in***
478 ***the summer it appears the mode during the day could be anywhere between 20 and 50nm***
479 ***and during the night it is around 35nm. Is the statement backed up by fitting a double log-***
480 ***normal to the data? Thirdly, why aren’t the percentiles for the PVSD presented if they are***
481 ***shown for the PNSD? Finally, why are the medians presented and not the means, which is***
482 ***more common? Does the later analysis use the means or medians? If the former, figure 3***
483 ***should show means.***

484 **Reply:** The discussion of the PNSDs and PNVDs has been improved according to the referee’s
485 comments. In addition, the mode ranges presented in the text have been corrected, as highlighted
486 by the referee. We have also fitted the size spectra with lognormal curves, by using 2 to 4 curves.
487 However, we do not believe that this fitting analysis should be included in the manuscript and only
488 served as a preliminary investigation of the PNSDs.

489 This Figure (now Figure 1) was originally planned to be presented as a boxplot (where IQR (25th-
490 75th percentile range) and median are shown). This is the reason for having selected these
491 statistics. However, we recognize that it is also useful to provide the average distributions (more
492 common, but potentially affected by outliers). Therefore, we have added them for PNSD and PVSD
493 as dotted lines.

494 We only showed the IQRs for PNSD because these are the data measured by the SMPS. PNVD
495 are subsequently derived by the simple assumption that all particles are perfectly spherical (which -
496 we know- isn’t true). Therefore, the modeling of PNVD is affected by a large uncertainty and in this
497 study is only used for descriptive purposes (not for quantitative analysis). Consequently, we
498 believe that it makes no sense to go into a deep analysis of the PNVD. In addition, we want to
499 keep the figures as simple as possible and the addition of other curves to the plot can mislead the
500 reader and distract from what is really important here, i.e. the PNSDs.

501 Finally, the PVSD are almost constant because the second mode of the PNSD does not change
502 during the day/night. The text has been improved accordingly.

503 The amended paragraph: “The average PNSDs are shown in Figure 2 as well as their median
504 distributions and interquartile ranges. Spectra are categorised by time of day (7am-7pm and 7pm-
505 7am local time). In addition, the particle volume size distributions (PVSDs) are also provided.

506 Results for the warm season show that the average daytime PNSD is dominated by a main peak in
507 the nucleation range (extending below 14 nm) and a second mode in the Aitken range (between 30
508 and 50 nm). The nocturnal spectrum is characterised by a drop of the nucleation mode to
509 concentration values similar to the Aitken peak (mode around 35 nm). During the cold season, the
510 average diurnal and nocturnal PNSDs present a main peak at 15-25 nm and a second mode at 70-
511 100 nm. In summary, both seasons show reductions of the finest modes during nighttime, while the
512 second mode is almost constant throughout the day. As a consequence, the modal structure of
513 PNVDs is also almost constant throughout the day.”.

514

515 ***Line 330, meteorology plays a role.***

516 **Reply:** A sentence was added to the text.

517

518 ***Line 345, constant to 11pm (same value pre noon in summer)***

519 **Reply:** The sentence has been amended.

520

521 ***Line 349, the accumulation mode peaks at around 10pm, which is inconsistent with the***
522 ***statement that it follows traffic which peaks at 5-6pm, for the M4 at least.***

523 **Reply:** An improved explanation was added. We believe that the dynamics of the mixing layer may
524 play a role in extending the evening peaks up to the night.

525 The amended text: “Generally, the evening peaks start around 6 pm, which is consistent with the
526 peak of traffic (Figure SI5) but they extend late in the evening and night probably because the drop
527 of the mixing layer and the consequent concentration of pollutants close to ground level.”.

528

529 ***Line 351, I don’t think the use of the word intermediate is appropriate. It is either traffic or***
530 ***aircraft or mixed, perhaps. Furthermore, this is interesting. Why is there a difference in***
531 ***winter and summer? If Version 9 of AIM was used, could this change your results if you try***
532 ***version 10?***

533 **Reply:** “Intermediate” has been replaced with “mixed”. We do not attempt a further explanation of
534 the diurnal profiles of the Aitken nuclei in this section. We limit our discussion to presentation of the
535 results. We believe that this “mixed” behavior is related to the influence of different sources and is
536 also complicated by the different meteorological conditions between the two seasons.

537 The amended sentences: "Aitken nuclei exhibit a mixed behaviour between nucleation and
538 accumulation particles (Figure 2): two different patterns can be found, which are more consistent
539 with road traffic in summer and with aircraft traffic in winter."

540 We have not investigated if this pattern is also detected when using AIM10 to extract the data. We
541 believe that the manuscript should only focus on data extracted with AIM V9 (see reply to the
542 referee's main point 3).

543

544 ***Lines 355 – 381. I do not think this section needs to be in the manuscript. The title of the***
545 ***manuscript is about the sources of the particles near an airport. The NO₂ is a) not showing***
546 ***any directionality and b) has been extensively studies already. The same is true for figures***
547 ***2 a and b, the NO_x data can be removed from there to streamline the paper.***

548 **Reply:** We have removed the paragraphs on the NO_x and O₃. The former Figure 2a has been
549 moved to SI. The former Figure 2b (now Figure 2) was modified to show the most important daily
550 patterns in view of the interpretation of the results. The diurnal patterns for all variables are now
551 provided in the SI.

552

553 ***Line 424 – I think cluster 2 can be bi modal as well, it certainly shows in the percentiles.***

554 **Reply:** The whole section on the cluster analysis has been modified. Now, we jointly discuss
555 Cluster 2 and 3 for the warm season and the Cluster 2 for the cold season. All these clusters have
556 bimodal structures and are related to the same source.

557

558 ***Line 429 – suggest changing to ‘the POTENTIAL role: : ...’***

559 **Reply:** Done.

560

561 ***Line 445 - again, is it mono-modal or bi-modal? The small bump at 14nm is in the median***
562 ***and percentiles.***

563 **Reply:** Done. The whole section on the cluster analysis has been modified. Now, we jointly discuss
564 Cluster 5 for the warm season and the Cluster 3 for the cold season. All these clusters have a main
565 peak at 20-35 nm. We added discussion of the second possible peak of Cluster 5 within brackets.

566

567 ***Line 457 & figure 4. The hourly count for cluster 4 is very noisy. Where is the morning***
568 ***peak? The one just after 6am? I also disagree it is the mirror image of clusters 2 & 3 counts.***
569 ***It shows a broadly opposite trend. Suggest rewording.***

570 **Reply:** This was unimportant to the main purpose of the paper and has been deleted.

571 ***Cluster 1 & 5 in the cold season – It seems a little odd that the count profile of cluster 1***
572 ***matches more the profile of LHR than cluster 5, yet the suggestion is that cluster 1 is an***
573 ***aged LHR aerosol, and cluster 5 is fresh because short transit times (high wind speed).***
574 ***Surely a fresh emission will match more the LHR profile, while the process of aging will***
575 ***remove or diminish the effects of source? I don't think the author's conjecture is correct for***
576 ***cluster 5 being fresh emissions.***

577 **Reply:** We have moved our discussion on these clusters from this section to Section 3.3
578 "Comparison of k-means and PMF". This latter analysis has allowed a better interpretation of the
579 Clusters 1 and 5 (cold season). Essentially, Cluster 5 presents high contributions of the "airport"
580 source of PMF and, therefore, may represent the spectra shaped largely by the airport emissions.
581 Cluster 1 also has high contributions for the PMF factor associated with road traffic. Both are mixed
582 spectra.

583

584 ***Cluster 3 cold season – can you back up your BB conjecture with the Aeth delta C data***
585 ***averaged over the same periods? If not, you should remove it.***

586 **Reply:** There is not a significant increase of delta-C. We have therefore removed this
587 "unsupported" interpretation. We have added a revised interpretation for this result: the winter
588 mixing layer effects.

589 The amended text: "However, the diurnal pattern in winter also presents a high number of counts
590 at 3-5 am, i.e. not directly compatible with rush hours. A possible explanation involves the stronger
591 effect of the winter mixing layer dynamics on the air quality due to the presence of more frequent
592 low level thermal inversions, which may build up the pollutants at ground-level especially overnight.
593 This may increase the signal of the less intense, but still significant, nighttime traffic emissions
594 present in the study area."

595

596 ***Line 509 replace hump with mode.***

597 **Reply:** Done.

598

599 ***Line 581 – The contribution to the NO₂ levels at LHR are quoted here as 25-30%, but in the***
600 ***previous section () is was found to be 15-17% from another study. Was this study different***
601 ***to previous ones?***

602 **Reply:** We have removed the sentences in line 367. We recognise that this discussion is
603 misleading. The data refer to two different studies carried out with different apportionment methods
604 and using data from different periods (2002-2004 and 2015-2012). In addition, we have now
605 removed the section discussing the NO_x data (see referee's main point 2).

606 A review of the results available in the literature is presented in the "Introduction": "For example,
607 Carslaw et al. (2006) estimated that airport operations in 2001/4 accounted for ~27% of the annual
608 mean NO_x and NO₂ at the airfield boundary and less than 15% (<10 µg m⁻³) at background
609 locations 2-3 km downwind of the airport. Similar results were found for the 2008/9 period using
610 model evaluation (AEA, 2010) and for the 2005/12 period using experimental data analysis (Masiol
611 and Harrison, 2015). This latter study also reported that PM mass concentrations at eight sites all
612 around LHR were always well below the EU and UK limit."

613 ***Line 637 – 50 – 200nm is not consistent with the authors definitions of UFP (<100nm)***

614 **Reply:** Amended.

615

616 ***Line 694 – 'to' missing from second sentence.***

617 **Reply:** Amended: "to" added.

618

619 ***Line 711 – 3% is not modest, it is minimal.***

620 **Reply:** Amended, "modest" substituted with "minimal".

621

622 ***Line 848 – suggest removing Anomalously . Being downwind of an airport is expected to***
623 ***lead to higher loadings.***

624 **Reply:** Done.

625

626

Sources of Submicrometre Particles Near a Major International Airport

Mauro Masiol^{1,2}, Roy M. Harrison^{1*†},
Tuan V. Vu¹, David C.S. Beddows¹

¹ Division of Environmental Health and Risk Management,
School of Geography, Earth and Environmental Sciences
University of Birmingham
Edgbaston, Birmingham B15 2TT
United Kingdom

² ~~Center for Air Resources Engineering and Science,~~
~~Clarkson~~ Division of Epidemiology, Department of Public
Health Sciences, University, Potsdam of Rochester Medical
Center,
265 Crittenden Boulevard, CU 420644
Rochester, NY 1369914642, United States

Formatted: No Spacing

Formatted: Centered

ABSTRACT

* To whom correspondence should be addressed.

Tele: +44 121 414 3494; Fax: +44 121 414 3709; Email: r.m.harrison@bham.ac.uk

† Also at: Department of Environmental Sciences / Center of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203, Jeddah, 21589, Saudi Arabia

Major airports are often located within or close to large cities; their impacts on the deterioration of air quality at ground level are amply recognised. The international airport of Heathrow is a major source of nitrogen oxides in the Greater London area, but its contribution to the levels of submicrometre particles is unknown, and is the objective of this study. Two sampling campaigns were carried out during warm and cold seasons at a site close to the airfield (1.2 km). Size spectra were largely dominated by ultrafine particles: nucleation particles (<30 nm) were found to be ~10 times higher than those commonly measured in urban background environments of London. A set of chemometric tools was used to discern the pollution arising from aircraft operations and those from other sources within the city or from the traffic generated by the airport. Five clusters and 6 factors were identified by applying *k*-means cluster analysis and positive matrix factorization (PMF) respectively to particle number size distributions; their interpretation was based on their modal structures, wind directionality, diurnal patterns, road and airport traffic volumes and on the relationship with weather and other air pollutants. Airport emissions, fresh and aged road traffic, urban accumulation mode and two secondary sources were then identified and apportioned. The comparison of cluster and PMF analyses allowed extraction of further information. ~~The analysis of a strong regional nucleation event was also performed to detect its effect upon concentrations.~~ The fingerprint of Heathrow has a characteristic modal structure peaking at <20 nm and accounts for 30-35% of total particles in both the seasons. Other main contributors are fresh (24-36%) and aged (16-21%) road traffic emissions and urban accumulation from London (around 10%). Secondary sources accounted for less than 6% in number concentrations but for more than 50% in volume concentration. The analysis of a strong regional nucleation event was also performed to detect its effects upon concentrations and source apportionment methods: results evidenced that both the cluster categorisation and PMF contributions were affected during the first 6 hours of the event. In 2016, the UK government provisionally approved the construction of a third runway; therefore the direct and indirect impact of Heathrow on local air quality is expected to increase unless mitigation strategies are applied successfully.

682 **Keywords:** Airport; black carbon; size distributions; source apportionment; ultrafine particles

683

684 1. INTRODUCTION

685 Emerging markets, developing economies and globalisation ~~are driving~~have driven a fast and
686 continuing growth of civil aviation, ~~which is expected to continue~~ in the ~~next decade~~last decades
687 (Lee et al., 2009); this trend is still growing by ~5.5% y⁻¹ (ICAO, 2017). As a consequence, the
688 aircraft and road traffic at airports is also increasing, but the information available on the impact of
689 airport emissions upon air quality at ground level is still inadequate (Webb et al., 2008; Masiol and
690 Harrison, 2014). The quantification of airport impacts on local air quality is complicated by the
691 complexity of multiple mobile and static emission sources, with many airports being located near to
692 major cities, highways or industrial plants. ~~Under this scenario~~Consequently, the development of
693 successful strategies for emission mitigation and the implementation of measures for air quality
694 ~~control improvement~~ to meet regulatory standards require ~~an exhaustive~~a detailed quantification of
695 the contribution of airport ~~and other~~ emissions to the total air pollution load.

696

697 Biological~~London Heathrow (LHR) is one of the world's busiest international airports: it is ranked~~
698 ~~1st in Europe for total passenger traffic (ACI, 2016). Its role in driving the economic affluence and~~
699 ~~vitality of the Southern UK is indisputable: it accommodates more than 1250 flights every day and~~
700 ~~serves a total of 72.3 million passengers year⁻¹. LHR is composed of 5 terminals and 2 runways:~~
701 ~~northern (3.9 km long) and southern (3.7 km). Currently, runways operate near their maximum~~
702 ~~capacity, with a consequent increase in the potential for delays when flights are disrupted. Since~~
703 ~~2007, the proposal for expanding LHR with a 3rd runway and a 6th terminal has been intensely~~
704 ~~debated in UK. The main reasons supporting its expansion are: (i) the expected increase of~~
705 ~~resilience to disruption caused by congested flight traffic; (ii) the improvement of its connectivity~~
706 ~~with a profitable network of both direct long haul air routes and national flight connections; (iii) the~~
707 ~~potential to directly enhance the economic growth of the London area. On the contrary, opposition~~

708 to LHR expansion highlights the potential increases in air pollution and noise, the community
 709 destruction and argues in favour of alternative options with fewer local impacts, such as the
 710 improvement of other airports in the southern UK or the building of a new airport in the Thames
 711 Estuary (East of London). Despite this, in 2016 the UK government provisionally approved the
 712 construction of a third runway.
 713
 714 Greater London is one of the few UK locations not fully achieving the EU and national air quality
 715 standards: in 2015 nitrogen dioxide breached the hourly and annual limit values for health, while
 716 ozone exceeded the long term objective (DEFRA, 2016). On the other hand, the mass concentration
 717 of particulate matter (PM), which is the standard current metric for measuring and controlling the
 718 exposure to airborne particles, was fully met for both PM₁₀ and PM_{2.5}. However, it has been widely
 719 demonstrated that even PM mass concentrations below guidelines and standards set by legislatures
 720 or international organizations may increase acute and chronic effects and mortality (e.g., Shi et al.,
 721 2015). In this situation, the use of mass concentration as a sole metric for measuring the levels of
 722 airborne particles has the disadvantage of taking greatest account of accumulation and coarse mode
 723 particles, which account for most of the mass. Consequently, the impact of the finest particles is not
 724 accounted for directly. This issue raises serious questions for the air quality standards: biological
 725 evidence associates the exposure to ultrafine particles (UFPs, <100 nm) with adverse effects upon
 726 human health (e.g., Knibbs et al., 2011; Strak et al., 2012; Ostro et al., 2015; Lanzinger et al., 2016).
 727 At the current time, there is still limited knowledge of what specific characteristic or association of
 728 characteristics may dominate the particle toxicity, and the consequent health outcomes (Atkinson et
 729 al., 2010; Strak et al., 2012, Vu et al., 2015a); nevertheless it is well recognised that UFPs can
 730 reach the deepest regions of the lung (Salma et al., 2015) and may have orders of magnitude higher
 731 surface area to mass ratios compared to larger particles. They offer more surface for the absorption
 732 of volatile and semi-volatile species (Kelly and Fussell, 2012; Strak et al., 2012). However, there
 733 are currently no ambient air quality standards or guidelines to drive the regulation of UFP.

734

735 [Several studies have reported large increases of UFPs near airports \(e.g., Westerdahl et al., 2008;](#)
736 [Hu et al., 2009; Klapmeyer et al., 2012; Hsu et al., 2012a;b\). For example, Hsu et al. \(2013\) and](#)
737 [Stafoggia et al. \(2016\) detected substantial increases of total particle number concentration \(PNC\) at](#)
738 [the airports of Los Angeles \(CA, USA\) and Rome Ciampino \(Italy\), respectively, in the few](#)
739 [minutes after take-offs, especially downwind, while landings made only a modest contribution to](#)
740 [ground-level PNC observations. Hsu et al. \(2014\) observed that departures and arrivals on a major](#)
741 [runway of Green International Airport \(Warwick, RI, USA\) had a significant influence on UFP](#)
742 [concentrations in a neighborhood proximate to the end of the runway. In a study carried out at the](#)
743 [Los Angeles international airport \(CA, USA\), Hudda et al. \(2014\) concluded that emissions from](#)
744 [the airport increase PNC by 4- to 5-fold at 8–10 km downwind of the airfield, while](#)
745 [Shirmohammadi et al. \(2017\) reported that the daily contributions of the airport to PNC were](#)
746 [approximately 11 times greater than those from three surrounding freeways. Hudda et al. \(2016\)](#)
747 [reported that average PNC were 2- and 1.33-fold higher at sites 4 and 7.3 km from the Boston \(MA,](#)
748 [USA\) airport when winds were from the direction of the airfield compared to other directions. The](#)
749 [goal of this study was to investigate the impacts of a major airport \(LHR\) serving a megacity](#)
750 [\(London\) upon the levels of submicrometre particles and equivalent black carbon \(eBC\) and to](#)
751 [apportion those impacts to aircraft, road traffic and other sources typical of large cities with](#)
752 [airports. This task was performed by collecting air quality data at a site downwind of LHR and by](#)
753 [applying a series of chemometric tools. The potential sources of submicron particle number](#)
754 [concentrations \(PNC\) are investigated by applying two source apportionment methods: cluster](#)
755 [analysis and positive matrix factorisation \(PMF\). Thus, the origin of the airport plumes was](#)
756 [spatially assessed by matching results with local meteorological data, air mass movements, levels of](#)
757 [common air pollutants, PM_{2.5} mass concentration and its chemical speciation as an indicators of](#)
758 [source location and formation mechanisms. Finally, the disaggregated source profiles are used to](#)

trace the factors affecting the pollutant levels, such as atmospheric dispersion and processing of aircraft emissions as well as of road traffic.

This study was carried out under the Marie Skłodowska-Curie project CHEERS (Chemical and Physical Properties and Source Apportionment of Airport Emissions in the context of European Air Quality Directives, call: FP7 PEOPLE 2012-IEF, project no. 328542).

2. MATERIALS AND METHODS

2.1 Study Area and Dates

The summer (warm season) campaign took place from 13 August to 12 September 2014 and the winter (cold season) campaign from 19 December 2014 to 20 January 2015.

Despite the strong evidence that airports are major sources of UFPs, their fingerprint ~~over~~within the particle number size distribution (PNSD) may be difficult to identify due to: (i) the nature of semi-volatile compounds emitted by aircrafts; (ii) the possible mechanisms of secondary aerosol formation; (iii) the dilution effect; and (iv) the similar modal structures of other emission sources concurrently found in cities, such as road traffic (Masiol and Harrison, 2014). Generally, studies performed within or close to airports ~~have~~ reported increases of particles ranging from 4 to 100 nm in diameter and mostly distributed in the nucleation range (<30 nm). For example, Mazaheri et al. (2009) showed a main nucleation mode and an accumulation mode (40-100 nm) more evident during take-offs; Keuken et al. (2015) reported PNSD dominated by 10-20 nm particles in an area affected by emissions from Schiphol airport (The Netherlands); Hudda and Fruin (2016) found strong increases in particles smaller than 40 nm downwind from the Los Angeles International Airport; Ren et al. (2016) showed that particles peaking at 16 nm dominate the PNSD at various distances from the runway of Tianjin International Airport, China; Masiol et al. (2016) reported that the fingerprint of aircraft emissions sampled under real ambient conditions at the airport of Venice

785 (Italy) ~~have~~has a main mode at approx. 80 nm and a second mode in the nucleation range below 14
786 nm.

787

788 The Greater London area ~~hosts-is~~ home to more than 8.5 million inhabitants and is one of the few
789 UK locations not fully achieving the EU and national air quality standards: in 2015 nitrogen dioxide
790 breached the hourly and annual limit values for health, while ozone exceeded the long-term
791 objective (DEFRA, 2016). However, the standards were fully met for both PM₁₀ and PM_{2.5}.

792

793 London Heathrow (LHR) is one of the world's busiest international airports: it is ranked 1st in
794 Europe for total passenger traffic (ACI, 2016). It accommodates more than 1250 flights every day
795 and serves a total of 72.3 million passengers year⁻¹. LHR is composed of 5 terminals and 2 runways:
796 northern (3.9 km-long) and southern (3.7 km). Currently, runways operate near their maximum
797 capacity, with a consequent increase in the potential for delays when flights are disrupted. Since
798 2007, the proposal for expanding LHR with a 3rd runway and a 6th terminal has been intensely
799 debated in the UK. In 2016 the UK government provisionally approved the construction of a third
800 runway (UK Department for Transport, 2017).

801

802 LHR is located west of London (Figure 4SI1). Consequently, air quality in the surroundings of the
803 airport may be affected by the advection of air masses from the city, with the associated high levels
804 of pollutants emitted from traffic, energy demand for domestic heating and local industries. Airport
805 activities may also contribute to air pollution advected to the city when LHR is upwind, with
806 consequent potential impacts upon public health. In addition, as LHR attracts a large number of
807 passengers and workers, the emissions from large volumes of road traffic generated by the airport
808 and the nearby M4 and M25 motorways are difficult to discriminate from non-airport-related road
809 traffic. Due to this complex scenario, the contribution of LHR is difficult to differentiate from the

810 urban background pollution, as already reported by previous modelling and experimental studies
811 (Farias and ApSimon, 2006; Masiol and Harrison, 2015).

812

813 Various studies have attempted to quantify the effect of LHR upon air quality, mainly focusing on
814 the nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$), which are well-known tracers for aircraft engine exhausts
815 (e.g., Herndon et al., 2008; Masiol and Harrison, 2014 and references therein), but also arise from
816 other combustion sources. For example, Carslaw et al. (2006) estimated that airport operations in
817 2001/4 accounted for ~27% of the annual mean NO_x and NO_2 at the airfield boundary and less than
818 15% ($<10 \mu\text{g m}^{-3}$) at background locations 2-3 km downwind of the airport. Similar results were
819 found for the 2008/9 period using model evaluation (AEA, 2010) and for the 2005/12 period using
820 experimental data analysis (Masiol and Harrison, 2015). This latter study also reported that PM
821 mass concentrations at eight sites all around LHR were always well below the EU and UK limit.

822

823 This study aims to investigate the impacts of a major airport (LHR) serving a megacity (London)
824 upon the levels of submicrometre particles and to apportion those impacts to aircraft, road traffic
825 and other sources typical of large cities with airports. The main particle size distributions modes are
826 simplified by applying cluster analysis; then, the modal structures of the main potential sources are
827 disaggregated and the submicron particle number concentrations (PNC) are quantified through the
828 positive matrix factorisation (PMF). In addition, the origin of the airport plumes was spatially
829 assessed by matching results with local meteorological data, air mass movements, levels of
830 common air pollutants, $\text{PM}_{2.5}$ mass concentration and its chemical speciation as ~~an~~ indicators of
831 source location and formation mechanisms.

832

833 The atmospheric chemistry and physicals properties of UFPs ~~were~~have been extensively
834 investigated in London (e.g., Harrison et al., 2012; Jones et al., 2012; von Bismarck-Osten et al.,
835 2013) with several studies using cluster analysis (Beddows et al., 2009; Brines et al., 2014; 2015)

836 or PMF (Beddows et al., 2015; Vu et al., 2016). However, this study is the first one carried out in
837 South-West London to characterise and quantitatively apportion the impacts of LHR under real
838 ambient conditions. Moreover, only one earlier study (Masiol et al., 2016) jointly has used both
839 cluster analysis and PMF to directly assess the airport contributions to UFPs. In addition, this study
840 also investigated the effects of a regional nucleation event on the results of the two source
841 apportionment methods.

842 ~~Regional photochemical nucleation episodes are regularly recorded in the Southern and Eastern~~
843 ~~UK. Their general characteristics have been reported in a number of studies (e.g., Alam et al., 2003;~~
844 ~~Charron et al., 2007;2008; Beddows et al., 2015; Vu et al., 2016) and can be summarised as~~
845 ~~follows: (i) particle modality at around 20 nm; (ii) higher frequency around noon in association with~~
846 ~~the peak in actinic flux intensities; (iii) clear seasonal cycles (higher average contribution levels in~~
847 ~~the summer, from June to September); (iv) marked directionality from the westerly sectors,~~
848 ~~reflecting maritime atmospheric circulation regimes, with high wind speed and low PM_{2.5}~~
849 ~~concentrations.~~

851 2. MATERIALS AND METHODS

852 2.1 Site Description.1 Experimental

853 Two intensive sampling campaigns (each 1 month-long) were carried out during warm (August-
854 September 2014) and cold (December 2014-January 2015) periods at Harlington (Figure 1). ~~Data~~
855 ~~from the site are quality assured as part of the UK Automatic Urban and Rural Network under the~~
856 ~~auspices of the UK Department for Environment, Food and Rural Affairs (DEFRA; [http://uk-](http://uk-air.defra.gov.uk/)~~
857 ~~[air.defra.gov.uk/](http://uk-air.defra.gov.uk/)) and the site SII1). The site~~ was selected as well located to sample the plumes from
858 the airport emissions. ~~The site;~~ it lies 1.2 km N of the northern runway and is located inside a
859 playground, close to a secondary road and near the village of Harlington. This is the location
860 selected for the construction of the 3rd runway. The site is categorised as “urban industrial” by
861 DEFRA and it is therefore more indicative of community exposure rather than direct fresh aircraft

Formatted: Tab stops: Not at 1.5 cm

emissions. Consequently, it is a good point to quantify the particles generated by the airport after a relatively short ageing and dispersion in the atmosphere, and is more indicative of the fingerprint of aircraft emissions affecting communities than data collected alongside the runway or in the airport apron areas. In addition, previous studies have reported that the site is strongly affected by the plume from the airport (Carslaw et al., 2006; Masiol and Harrison, 2015). Prevailing winds from the 3rd and 4th quadrants are recorded in both summer and winter (Figure [S4S12](#)): under such circulation regimes, Harlington lies just downwind of LHR. ~~However, the~~[The](#) site is also affected by pollutants arising from the large volumes of road traffic ~~of~~[within London, from the local road network as well as those](#) generated by the airport: Tunnel Rd., the main access to LHR from the M4 motorway lies 800 m west, as well as the nearby M4 (640 m north) and M25 (~3.5 km east) motorways, major roads (Bath Rd, part of A4, passes 900 m south; A30 lies 2.8 km SE). The village of Harlington (~400 m west) and [advection of air masses from](#) the conurbation of London are other potential external sources.

~~2.3~~ [Instrumentation Suite](#)

Ultrafine particle counts and their size distributions from 14.3 to 673.2 nm were measured at 5 min time resolution using a SMPS (scanning mobility particle sizer spectrometer) comprising a electrostatic classifier TSI 3080 with a long differential mobility analyser (TSI 3081) and a CPC (condensation particle counter, TSI 3775) based on condensation of *n*-butyl alcohol (Fisher Scientific, ACS). The SMPS operated at a sheath air to aerosol flow ratio of 10:1 (sheath and sample air flow rates were 3.0 and 0.3 L min⁻¹ respectively, voltage 10-9591 V; density 1.2 g/cc; scan time 120 s, retrace 15 s; number of scan 2) while the CPC operated at low flow rate (0.3 L min⁻¹). The use of 5 min resolved spectra has already been used successfully for source apportionment purposes at an airport (Masiol et al., 2016).

eBC was also measured at 5 min resolution using a 7-wavelength aethalometer (Magee Scientific AE31). The aethalometer operated with an inlet cut-off head to collect PM with aerodynamic diameter of $<2.5\ \mu\text{m}$ ($\text{PM}_{2.5}$). eBC was derived from the absorbance at 880 nm wavelength (Petzold et al., 2013); raw data were post-processed with the Washington University Air Quality Lab AethDataMasher V7.1 to perform data validation and correct data for non-linear loading effects (Virkkula et al., 2007; Turner et al., 2007).

Instruments were installed into a plastic/metal case designed for sampling purposes: (i) air inlets were $\sim 1.8\ \text{m}$ ~~over~~ above the ground and were composed of conductive materials to avoid particle losses and sampling artefacts; (ii) the case was cooled by fans in summer and was warmed by an electrical tubular heater in winter for maintaining an indoor air temperature within an acceptable range for running the equipment (temperature inside the case was recorded and periodically checked); (iii) instruments were isolated from vibration using rubber pads and foam foils. Devices were fully serviced, calibrated by authorised companies and underwent internal cross-calibrations with other similar instruments under lab conditions. Moreover, frequent periodic checks, maintenance of instruments and cleaning of inlets was performed throughout the sampling campaign.

~~Classical~~ Routine air pollutants (NO , NO_2 , NO_x , O_3 , PM_{10} , $\text{PM}_{2.5}$) were measured at Harlington with 1 h time resolution ~~by the UK Automatic Urban and Rural Network under the auspices of the UK~~ Department for Environment, Food and Rural Affairs (DEFRA; <http://uk-air.defra.gov.uk/>).

Gaseous species were analysed using automatic instruments according to European standards and National protocols: EN 14211:2012 for nitrogen oxides and EN 14625:2012 for ozone. PM_{10} and $\text{PM}_{2.5}$ were analysed using tapered element oscillating microbalance and filter dynamics measurement system (TEOM-FDMS) to provide measurements accounting for volatile (VPM_{10} , $\text{VPM}_{2.5}$) and non-volatile (NVPM_{10} , $\text{NVPM}_{2.5}$) fractions. Quality assurance and quality control

913 procedures followed the standards applied for the Automatic Urban and Rural Network (AURN)
914 and the London Air Quality Network (LAQN). Instruments were routinely calibrated, and every six
915 months were fully serviced and underwent intercalibration audits.

916

917 Some additional variables are also computed from the air pollutants to help the interpretation of
918 results. The NO_2/NO_x ratio is indicative of the partitioning of nitrogen oxides, while the levels of
919 oxidants ($\text{OX}=\text{O}_3+\text{NO}_2$, expressed in ppbv) can be used to roughly assess the oxidative potential in
920 the atmosphere (Kley et al., 1999; Clapp and Jenkin, 2001). These two new variables are useful in
921 investigating the atmospheric chemistry behind the $\text{NO}-\text{NO}_2-\text{O}_3$ system. Delta-C (the difference
922 between absorbance at 378 and 880 nm, also called UVPM) was also computed. This variable was
923 largely used as a proxy to estimate the fraction of carbonaceous material emitted by biomass
924 burning (e.g., Sandradewi et al., 2008; Wang et al., 2011). However, Delta-C results should be used
925 with caution: Harrison et al. (2013) showed that there are probably other UV absorbing contributors
926 than wood-smoke to the aethalometer signal. ~~This way~~ Consequently, Delta-C is used here only for
927 qualitative purposes.

928

929 Weather data were measured hourly by the Met Office at LHR; met data include wind direction and
930 speed, atmospheric pressure, air temperature, relative humidity (RH), visibility, rain and solar
931 irradiance.

932

933 During the two campaigns, 24-h $\text{PM}_{2.5}$ samples were also collected on quartz filters using a high
934 volume air sampler (TE-6070, Tisch Environmental, Inc.) and analysed for the daily concentrations
935 of major $\text{PM}_{2.5}$ components: organic carbon (OC) and elemental carbon (EC) by thermo-optical
936 analysis (EUSAAR_2 protocol) and major inorganic ions (Na^+ , K^+ , ammonium, nitrate, sulphate,
937 oxalate) by ion chromatography. Analytical methods are reported in detail in Yin et al. (2010). The

938 results of the chemical speciation of PM_{2.5} are presented in a companion paper (in preparation) and
939 are used in this study only to assist the interpretation of PMF results.

940

941

942

943 **2.42 Data Handling and Chemometric Approaches**

944 Data were analysed using R version 3.3.1 (R Core Team, 2015) and a series of supplementary
945 packages, including ‘Openair’ (Carslaw and Ropkins, 2012). Preliminary data handling and clean-
946 up were carried out to check the robustness of the dataset, detect anomalous records and to delete
947 extreme outliers. SMPS data with unreliable behaviour or instrument errors were completely
948 deleted. ~~All remaining~~A deep~~An in-depth analysis of the dataset evidenced~~revealed few records
949 with anomalously high PNC, which were likely related to probable instrumental issues, extreme
950 weather conditions (e.g., high wind gusts, heavy rain ~~stacking~~striking the inlet), or
951 ~~uncommon~~infrequent local emissions, e.g., maintenance, painting and recreational activities
952 (including fires) on the playground where the site is located, road maintenance close the site and
953 probable short-term parking of high-emission vehicles near the site. Since this study aims to
954 investigate the overall contributions of LHR, all data are used for descriptive statistics, but data
955 greater than the 99.5th percentile were further removed for explorative, cluster and PMF analyses.
956 This data exclusion successfully ~~dropped off~~removed the extremely high events occurring~~ed~~during
957 the sampling campaigns and significantly improved the stability and physical meaning of PMF
958 solutions. Missing data for other variables were linearly interpolated between the nearest values of
959 the time series.

960

961 The particle number size distributions (PNSDs) were firstly grouped by applying a *k*-means cluster
962 analysis. The full method is exhaustively discussed in Beddows et al. (2009; 2014) and aims to
963 assemble single spectra into *k* clusters. The clustering groups observations with spectra similar to

964 their cluster centroids (means), i.e. observations that are likely generated by the same set of
965 formation processes or emission sources. The optimum number of clusters (k) was determined by an
966 optimisation algorithm based on the spectral shapes (Beddows et al., 2009). The choice to apply the
967 k -mean clustering method was based on several reasons: (i) Salimi et al. (2014) reported that k -
968 means is the best performing clustering among others methods tested on PNSD data; (ii) k -means is
969 a well-established method which has been largely-widely applied over a number of different sites
970 (e.g., Dall'Osto et al., 2012; Wegner et al., 2012; Beddows et al., 2014; Brines et al., 2014; 2015);
971 and (iii) the method was previously applied successfully to airport data (Masiol et al., 2016).
972 PMF analysis was performed by applying the USEPA PMF5 model. Details of the PMF model are
973 reported elsewhere (Paatero and Tapper, 1994; Paatero, 1997; USEPA, 2014), while the best
974 practice and standards are extensively reviewed in several papers (e.g., Reff et al., 2007; Belis et al.,
975 2014; Brown et al., 2015; Hopke, 2016). SMPS data at 5 min resolution were used as the PMF input
976 matrix. Uncertainties associated with SMPS data were estimated according to the empirical method
977 proposed by Ogulei et al. (2007). Uncertainty for the total variable (total particle number
978 concentration, PNC) was set at 300% of the PNC concentration and also marked as “weak” to avoid
979 it driving the profiles.

980

981 The best PMF solutions were identified: (i) by investigating solutions between 3 and 10 factors; (ii)
982 by considering the minimization of the objective function Q with respect to the expected
983 (theoretical) value and its stability over multiple ($n=100$) runs, (iii) by obtaining low values for the
984 sum of the squares of the differences in scaled residuals for each base run pair by species; (iv) by
985 minimizing the number of absolute scaled residuals over ± 3 and by keeping them symmetrically
986 distributed; (v) by keeping the result uncertainties calculated by bootstrap (BS, $n=200$) and
987 displacement (DISP) methods within an acceptable range (Paatero et al., 2014); (vi) by obtaining
988 modelled total variable (PNC) successfully predicted ($R^2 > 0.9$ and slopes ≈ 1); and (vii) by avoiding

the presence of edges in the G-space plots (Paatero et al., 2002) and, then, the presence of hidden/unresolved sources.

A series of additional tools were used to analyse the raw data, link source apportionment results to other variables, such as local atmospheric circulation and regional/transboundary transport of air masses. Briefly, polar plots aim to map pollutant average concentrations by wind speed and direction as continuous surfaces (Carslaw et al., 2006), while polar annuli plot by wind direction and hours of the day. The potential locations of distant sources were assessed using back-trajectory analysis and a concentration weighted trajectory (CWT) model (Stohl, 1998). Back-trajectories were computed with the HYSPLIT4 model (Stein et al., 2015; Rolph, 2016) using NCEP/NCAR reanalysis gridded meteorological data. Set-up: -96 h with a starting height of 500 m a.g.l. CWT is a method of weighting trajectories with associated concentrations to detect the most probable source areas of long-range transports of pollutants; it has been used and reviewed in a number of prior studies (e.g., Stohl, 1996; Lupu and Maenhaut, 2002; Squizzato and Masiol, 2015).

3. RESULTS AND DISCUSSION

3.1 Overview of Data

The wind roses during the two sampling periods are provided in Figure 1. ~~Descriptive statistics of all collected variables are aggregated as boxplots in Figure 2a. Some additional variables are also computed to help the interpretation of results.~~ SI2. Descriptive statistics of all collected variables are reported as boxplots in Figure SI3. ~~The NO_2/NO_x ratio is indicative of the partitioning of nitrogen oxides, while the levels of oxidants ($\text{OX} = \text{O}_3 + \text{NO}_2$, expressed in ppbv) can be used to roughly assess the oxidative potential in the atmosphere (Kley et al., 1999; Clapp and Jenkin, 2001). These two new variables are useful in investigating the atmospheric chemistry behind the $\text{NO}-\text{NO}_2$~~

~~O₃ system. Delta C (the difference between absorbance at 378 and 880 nm, also called UVPM) was~~
~~also computed. This variable was largely used as a proxy to estimate the fraction of carbonaceous~~
~~material emitted by biomass burning (e.g., Sandradewi et al., 2008; Wang et al., 2011). However,~~
~~Delta C results should be used with caution. Harrison et al. (2013) showed that there are probably~~
~~other UV absorbing contributors than wood smoke to the aethalometer signal. This way, Delta C is~~
~~used here only for qualitative purposes.~~ PNSDs were initially split into 3 ranges: nucleation (14-30
 nm), Aitken nuclei (30-100 nm) and accumulation (>100 nm). On average the total PNC during the
 warm season was 1.9×10^4 particles cm⁻³, of which 1.1×10^4 , 6.4×10^3 and 1.5×10^3 particles cm⁻³
 were classified as nucleation, Aitken and accumulation ranges, respectively. (Figure SI3). During
 the cold season, the total average PNC was 2.2×10^4 particles cm⁻³, composed of 1.4×10^4 , $6.3 \times$
 10^3 and 1.4×10^3 particles cm⁻³ as nucleation, Aitken and accumulation ranges, respectively.
 (Figure SI3). Concentrations lie between those of London, Marylebone Road (kerbside) and
 London, North Kensington (background), and nucleation particles were ~10 times higher than the
 annual average measured in North Kensington as reported by Vu et al. (2016), while Aitken
 particles were 1.9 times higher. It is therefore evident that the main difference lies in the
 concentration of the finest size ranges: in both seasons, spectra were dominated by UFP (D_p<100
 nm) particles (~92% of total PNC), which only accounted for ~12% of total particle volume
 concentration (PVC, computed by approximation to spherical particles). On the other hand,
 accumulation mode particles accounted for ~8% of PNC and ~88% of PVC volume. The high levels
of total PNC are not surprising: several studies carried out into or close to airports (e.g., Hsu et al.,
2013;2014; Hidda et al., 2014; 2016; Stafoggia et al., 2016; Shirmohammadi et al., 2017) reported
significant increases in the concentrations of UFPs.

The high levels of total PNC are not surprising. Several studies have reported large increases in
PNC near airports. For example, Hsu et al. (2013) and Stafoggia et al. (2016) detected substantial
increases of PNC values at the airports of Los Angeles (CA, USA) and Rome Ciampino (Italy).

1041 ~~respectively, in the few minutes after take-offs, especially downwind, while landings made only a~~
1042 ~~modest contribution to ground level PNC observations. Hsu et al. (2014) observed that departures~~
1043 ~~and arrivals on a major runway of Green International Airport (Warwick, RI, USA) had a~~
1044 ~~significant influence on UFP concentrations in a neighborhood proximate to the end of the runway.~~
1045 ~~In a study carried out at the Los Angeles international airport (CA, USA), Hudda et al. (2014)~~
1046 ~~concluded that emissions from the airport increase PNC by 4 to 5 fold at 8–10 km downwind of~~
1047 ~~the airfield, while Shirmohammadi et al. (2017) reported that the daily contributions of the airport~~
1048 ~~to PNC were approximately 11 times greater than those from three surrounding freeways. Hudda et~~
1049 ~~al. (2016) reported that average PNC were 2 and 1.33 fold higher at sites 4 and 7.3 km from the~~
1050 ~~Boston (MA, USA) airport when winds were from the direction of the airfield compared to other~~
1051 ~~directions. The site used in this study is even closer to the airfield (1.2 km) and is also affected by~~
1052 ~~strong non airport sources, such as road traffic emissions due to the presence of two motorways and~~
1053 ~~several busy roads (frequently congested).~~

1054
1055 During the ~~warm season, the average concentrations for other pollutants followed the order (in μg~~
1056 ~~m^{-3}): NO_x (49) > O_3 (31) > NO_2 (31) > PM_{10} (20) > NVPM_{10} (16) > $\text{PM}_{2.5}$ (14) > NO (12) > $\text{NVPM}_{2.5}$~~
1057 ~~(11) > VPM_{10} (4) > $\text{VPM}_{2.5}$ (3.2) > eBC (2.4) > Delta C (<0.1). The average concentrations during the~~
1058 ~~cold season were: NO_x (83) > NO_2 (38) > O_3 (34) > NO (29) > PM_{10} (18) > NVPM_{10} (14) > $\text{PM}_{2.5}$~~
1059 ~~(13) > $\text{NVPM}_{2.5}$ (9.8) > VPM_{10} (4.3) > $\text{VPM}_{2.5}$ (3.4) > eBC (2.1) > Delta C (0.2). These values are two~~
1060 ~~sampling campaigns, air pollutants measured in Harlington (Figure SI3) were~~ similar to the average
1061 concentrations ~~for common air pollutants~~ measured over an 8 year period (2005-2012) in the
1062 vicinity of LHR ~~reported by (Masiol and Harrison (2015) over an 8 year period (2005-2012)).~~
1063 Consequently, despite the ~~intensive sampling campaign~~ two short campaigns carried out in this
1064 study, results may be considered representative of the average levels of air pollution recorded at
1065 Harlington. The average concentrations of eBC were 2.4 and 2.1 $\mu\text{g m}^{-3}$ during the warm and cold

1066 season, respectively. The average concentration of Delta-C was $0.1 \mu\text{g m}^{-3}$ during the warm season
1067 and $0.36 \mu\text{g m}^{-3}$ in winter.
1068 Since
1069 Analysis of the data ~~were generally showed not a non-distributed normally,~~normal distribution for
1070 most of the variables; the nonparametric Kruskal-Wallis one-way analysis of variance was therefore
1071 used to test the difference of concentrations over the two periods; (Kruskal and Wallis, 1952);
1072 almost all variables are different at the 0.05 significance level, except NO, NO_x and O₃. This result
1073 indicates a seasonal effect upon air quality in the LHR area and suggests investigating the sources
1074 over the two periods separately.

1075 _____
1076 The average PNSDs are shown in Figure 31 as well as their median distributions and interquartile
1077 ranges. Spectra are categorised by time of day (7am-7pm and 7pm- 7am local time). In addition, the
1078 particle volume size distributions (PVSDs) are also provided. ~~Results show that in both seasons the~~
1079 ~~nocturnal data are shifted toward coarser modes with respect to the diurnal mean PNSD, while the~~
1080 ~~modal structure of PNVDs is almost constant throughout the day.~~ Results for the warm season show
1081 that the average daytime PNSD is dominated by a main peak in the nucleation range (extending
1082 below 14 nm) and a second mode in the Aitken range (between 30 and 50 nm). The nocturnal
1083 spectrum is characterised by a drop of the nucleation mode to concentration values similar to the
1084 Aitken peak (mode around 35 nm). During the cold season, the average diurnal and nocturnal
1085 PNSDs present a main peak at 15-25 nm and a second mode at 70-100 nm. In summary, both
1086 seasons show ~~drops~~reductions of the finest modes during nighttime, while the second mode is
1087 almost constant throughout the day. As a consequence, the modal structure of PNVDs is also almost
1088 constant throughout the day.

1089
1090 The diurnal cycles of ~~most important variables~~ the 3 particle ranges, eBC, solar irradiation and
1091 airport movements are shown in Figure 2b2. A comprehensive overview of the patterns for all the

Formatted: Font color: Red

variables is provided in Figure SI4. Generally, diurnal cycles derive from the interplay of emissions, dispersion and atmospheric chemical processes. ~~Meteorology also plays a role.~~ Consequently, they need to be investigated along with patterns for airport and motorway traffic (Figure 2b2 and Figure SI2SI5, respectively), and as polar annuli (Figures SI3SI6 and SI4SI7) and polar plots (Figures SI5SI8 and SI5SI9), which give preliminary insights ~~upon~~ into the origin and spatial location of most probable emission sources. ~~Airport~~ During nighttime, airport traffic ~~undergoes to some restrictions~~ is restricted to limit noise and community disturbance: flights are generally constant from 6 am to 8 pm and are kept at minimum overnight, with no departures normally scheduled between 11 pm and 6 am (Figure 2b2). Road traffic is more difficult to define. Data for M4 and M25 motorways are provided by the UK Department for Transport: data for the M4 motorway show typical morning (7-8 am) and evening (5-6 pm) peaks due to rush hours, but this pattern is not well-resolved for the M25 (Figure SI2SI5). In addition, despite it being likely that traffic on minor and local roads also follows patterns dominated by rush hours, traffic generated by the airport is more difficult to characterise, with Tunnel Rd. and other busy roads serving LHR being frequently congested.

Nucleation particles are likely associated with aircraft movements: ~~the~~ The daily pattern shows high and almost constant concentrations between 7 am and ~~1011~~ 11 pm, ~~while levels~~ (Figure 2): hourly averages ranged from 10×10^3 to 15×10^3 particles cm^{-3} during the warm season and from 10×10^3 to 21×10^3 particles cm^{-3} during the cold season. On the contrary, the concentrations of nucleation particles significantly (Kruskal-Wallis at $p < 0.05$) drop ~~to near zero~~ overnight; (hourly averages ranging from 5×10^3 to 6×10^3 particles cm^{-3} and from 1×10^3 to 5×10^3 particles cm^{-3} during the warm and cold season, respectively); the maximum average concentrations are recorded for winds blowing from the SW quadrant, ~~i.e. (polar plots and polar annuli in Figures SI6-9), i.e.~~ the airfield and, in particular, the location of the main LHR terminals- (Figure SI1). As a consequence of the dominance of nucleation particles over size spectra, also total PNC follows ~~this~~ the pattern.

1118 [\(Figures 2\) and wind directionality \(Figures SI8-9\) of nucleation particles.](#) On the contrary,
 1119 accumulation particles appear to be [more](#) associated with road traffic, ~~i.e. daily cycles show typical~~
 1120 ~~rush hour peaks and increases.~~ [These particles increase](#) for winds blowing from northern sectors:
 1121 [\(Figures SI6-9\), i.e. toward the M4. Accumulation particles also present the morning \(6-8 am\) and](#)
 1122 [evening \(6-11 pm\) rush hours peaks during the warm season, but only the evening peak \(from 6 pm](#)
 1123 [to the night\) was found in the cold season \(Figure 2\). Generally, the evening peaks start around 6](#)
 1124 [pm, which is consistent with the peak of traffic \(Figure SI5\) but they extend late in the evening and](#)
 1125 [night probably because the drop of the mixing layer top and the consequent concentration of](#)
 1126 [pollutants close to the ground level.](#) Aitken nuclei exhibit ~~an intermediate~~ [a mixed](#) behaviour
 1127 between nucleation and accumulation particles: [\(Figure 2\)](#); two different patterns can be found,
 1128 which are more consistent with road traffic in summer and with aircraft traffic in winter.
 1129
 1130 ~~Nitrogen oxides are key air pollutants for this study: (i) NO₂ levels do not fully fulfil the air quality~~
 1131 ~~assessment Limit Values for health (1 h and annual mean) in the Greater London urban area~~
 1132 ~~(DEFRA, 2016); (ii) they can be good tracers for airport emissions, since NO₂ is the main species of~~
 1133 ~~nitrogen oxides emitted by turbofan engines at idle, while NO is the dominant species at higher~~
 1134 ~~thrust (Wormhoudt et al., 2007; Masiol and Harrison, 2014); (iii) they are also emitted from road~~
 1135 ~~traffic mainly as NO, although recent non-attainments of NO₂ standards in Europe have been linked~~
 1136 ~~to the growing proportion of diesel powered vehicles, which have higher primary (direct) emissions~~
 1137 ~~of NO₂ (Carslaw et al., 2007; Grice et al., 2009; Anttila et al., 2011; Cyrys et al., 2012). In addition,~~
 1138 ~~nitrogen oxides and atmospheric oxidants are strongly linked by a series of chemical reactions~~
 1139 ~~which are responsible for their partitioning between NO and NO₂ (Finlayson Pitts and Pitts, 2000;~~
 1140 ~~Seinfeld and Pandis, 2006). To date, NO_x has been thoroughly investigated at LHR (Carslaw et al.,~~
 1141 ~~2006; Masiol and Harrison, 2015): it was estimated that the upper limit contribution of LHR~~
 1142 ~~activities to NO₂ at Harlington during the 2001-2012 period was ~15-17%, while that for NO was~~
 1143 ~~~10%. In this study, nitrogen dioxide exhibits two typical rush hour peaks, as previously also~~

1144 observed at the London, North Kensington urban background site (Bigi and Harrison, 2010), and its
1145 concentration increases for winds blowing from all quadrants, suggesting a mix of different sources,
1146 including airport, road traffic and other combustion emissions. Nitric oxide only shows the morning
1147 rush hour peak and northern directionality (toward the M4 motorway) in summer, while in winter it
1148 lacks any significant pattern. The difference between the patterns of NO and NO₂ during the two
1149 periods is also confirmed by the NO₂/NO_x ratio, which shows a morning rush hour minimum in
1150 summer as a consequence of fresh NO emissions, while it is less variable in winter (Figure 2b).
1151
1152 In 2015, ozone met the EU target value, but not the long term objective in the Greater London area
1153 (DEFRA, 2016). In this study, it does not present any wind directionality and exhibits an evident
1154 daily peak in the mid-afternoon, i.e. when the photochemical activity is enhanced by the higher
1155 solar irradiation and the boundary layer depth is greatest, while a second peak in the early morning
1156 corresponds to a minimum in NO (Figure 2b).

1157
1158 Despite some studies indicating that airports are strong sources of black carbon (Dodson et al.,
1159 2009), other studies report no strong relationships with the flight activity (Masiol et al., 2016; Hsu
1160 et al., 2016). Similarly to NO₂, (Figure SI4) and accumulation particles (Figure 2), aethalometer data
1161 also shows typical patterns of road traffic-influenced sites for all wavelengths, with two daily peaks
1162 corresponding to the hours with higher traffic- (Figure 2). However, Delta-C does not present any
1163 evident pattern- (Figure SI4). eBC shows increased concentrations when winds blow from northern
1164 sectors (plus SE in winter, Figure SI7 and SI9); which excludes airport activities as being a
1165 dominant source in the study area.

1166
1167 Particulate matter mass concentration (PM₁₀ and PM_{2.5}) has very weak diurnal patterns- (Figure
1168 SI4). Its wind directionality shows evident increases for northerly winds- (Figure SI8-9). It is
1169 therefore evident that PM mass concentrations are dominated by non-airport sources, i.e. regional

secondary pollutants, traffic from the nearby M4 or background pollution from London. PM_{2.5} concentrations normally do not exceed the Limit Values in the Greater London area (DEFRA, 2016).

3.2 *k*-means Cluster Analysis

The clustering algorithm extracted 5 clusters for both periods. The number of clusters was selected according to the optimisation algorithm, i.e. local maxima in the Dunn indices and silhouette (Beddows et al., 2009). The extraction of 5 clusters represents a good compromise for the interpretation of spectral observations. Hussein et al. (2014) reported that is not prudent to describe the spectra with few clusters (2-4), which are not sufficient to explain variations and detailed differences in the PNSD observed in the urban atmosphere. On the other hand, they also reported that extracting too many (>10) clusters may make the aerosol source attribution more challenging.

The cluster centroids (mean spectra of each cluster), the 10th, 25th, 75th and 90th percentile, the hourly counts patterns and resulting wind roses are shown in [Figure 4](#) [Figures 3](#) and [54](#) for the warm and cold season campaigns, respectively. Despite extracted clusters exhibiting significantly different modal structures for PNC, no differences can be observed for the particle volume size spectra, which all show a unimodal peak at approx. 200-300 nm. [Clusters accounted for 14%-25% of total observations for both the seasons: Table SI1 summarises the percentage of the total observations for each cluster.](#)

~~3.2.1 Three clusters (cluster 1, Warm season~~
~~During~~ during the warm season, ~~20% of total clustered observations were grouped and clusters 1~~
~~and 5 in cluster 1. It presents a winter)~~ are likely shaped by the airport emissions. The modal
~~structures present~~ sharp ~~peak~~ peaks for nucleation particles which ~~extends~~ extend below the SMPS
detection limit (14 nm), ~~and drop at 30-40 nm; no secondary modes are present in the Aitken or~~

Formatted: Font: Not Italic

1196 [accumulation ranges. These clusters show](#) a large increase in frequency during the afternoon [and](#)
 1197 [evening](#) hours (~~noon to 7pm~~) ~~and its wind rose shows that this spectrum shape mostly occurs when~~
 1198 ~~the prevailing wind blows from SW: cluster 1 for the warm season and cluster 5 for the cold season)~~
 1199 [or extended over the daytime \(cluster 1 for the cold season\), similarly to the airport aircraft](#)
 1200 [movement profiles \(Figure 2\).](#) Aircraft are known to emit particles in the nucleation range (e.g.
 1201 Mazaheri et al., 2009;2013; Masiol and Harrison, 2014; and references therein; Lobo et al., 2015)
 1202 and the wind ~~rose is~~[roses are](#) also compatible with an origin from the airfield and the main LHR
 1203 terminals: [\(Figures 3 and 4\).](#) However, ~~a similar PNSD profile and a similar daily pattern was also~~
 1204 ~~reported in North Kensington (daytime regional photochemical nucleation events in London~~
 1205 ~~background) by occur around noon-2 pm and are mostly recorded from June to September (Vu et al.~~
 1206 ~~(., 2016) and was associated with nucleation events. Its interpretation can thus).~~ Consequently, the
 1207 [modal structure of cluster 1 for the warm season could](#) be ~~associated either with airport activities or~~
 1208 [additionally shaped by regional](#) photochemical nucleation.
 1209
 1210 ~~Clusters 2 and 3 account for 19% and 23%~~[The reasons driving the split of the spectra likely shaped](#)
 1211 [by LHR into two clusters during the cold season are unclear. A further comparison of the cluster](#)
 1212 [and PMF results will help in interpreting this outcome.](#)
 1213
 1214 [The modal structures of the clusters 4 for both seasons peak for nucleation particles and extend](#)
 1215 [below 14 nm, but also show probable modes between 50 and 200 nm \(Figures 3 and 4\). They](#)
 1216 [represent the typical spectra associated with aged anthropogenic emissions, mostly due to road](#)
 1217 [traffic. It is recognised that road traffic contributes to a large range \(30-200 nm\) of PNSD in the](#)
 1218 [urban atmosphere \(e.g., Yue et al., 2008; Costabile et al., 2009; Harrison et al., 2011\), which is](#)
 1219 [compatible with observations, respectively. While cluster 2 shows a main peak in number](#)
 1220 [concentrations at 30-40 nm, cluster 3 is bimodal \(14 and 60-70 nm\). Both clusters exhibit similar](#)
 1221 [hourly count profiles with most of the counts occurring overnight these spectra. The directional](#)

analysis for the warm season shows increased levels when air masses move from the sectors more affected by traffic, i.e. London (NE), M4 (N) and M25 (W) motorways and Tunnel Rd (W), while the hour count profile presents a huge maximum during daytime. In winter, this modal structure mostly occurred for westerly winds; the atmospheric circulation during the cold season mostly experienced winds blowing from the SW quadrant, with NE sectors poorly represented (Figure S11). As a consequence, the limited number of observations for air pollution advected from the Greater London area may have affected the detection of the urban background from London. This lack of data is also reflected by diurnal profile, which shows a marked peak in the late afternoon, concurrent to the peak of traffic on M4 and M25 (Figure S15).

Three clusters (*clusters 2 and 3 during the warm season and cluster 2 in winter*) exhibited similar hourly profiles with most of the counts occurring overnight (Figures 3 and 4). This pattern is largely attributable to the dynamics of the mixing layer, since the diurnal cycles are the mirror image of the ambient air temperature (Figure 2b). ~~Because of this, both clusters are strongly affected by the reduced height of the mixing layer occurring overnight. In addition, the~~ SI4). Because of this, these clusters could be potentially affected by the reduced height of the mixing layer occurring overnight. These clusters exhibit bimodal structures with the coarser modes with respect to the remaining clusters: cluster 2 for the warm season shows a main peak in number concentrations at 30-40 nm and a second ~~bp~~ peak in the finest range (<16 nm), clusters 3 for the warm season peaks at 14 and 60-70 nm, and cluster 2 for the cold season extends over a wide size range with two modes around 20-30 nm and 100-150 nm. Consequently, these clusters are likely representative of spectra mostly shaped by the drop of the mixing layer height and the formation of secondary aerosols. ~~Under this view~~ In this context, the potential role of ~~the~~ nighttime nitrate formation through condensation of NH_4NO_3 and the heterogeneous reactions of N_2O_5 and NO_3 on pre-existing particles cannot be ignored (Seinfeld and Pandis, 2006; Bertram and Thornton, 2009; Brown and Stutz, 2012).

~~However, such~~ The wind roses reveal that ~~both~~ clusters 2 occur under ~~different~~ similar westerly

wind regimes, ~~as the wind roses indicate two different potential source locations: cluster 2 shows~~
~~a~~ ~~with a possible origin from W sectors, while cluster 3 indicates the NE. From this we can infer~~
~~that cluster 2 likely represents PNSD shaped by: (i) regional aerosols, since the wind directionality~~
~~suggests an origin from regions west of West London, an area with a lower density of anthropogenic~~
~~sources. This way, regional aerosols is appear to be the most probable source. On the contrary,~~
~~cluster 3 for the warm season occurs~~ ~~for~~ ~~with~~ winds from London (NE) and ~~(ii) emissions from the~~
~~M25 motorway and Tunnel Road, i.e. it can be influenced by aged road traffic emissions. This latter~~
~~interpretation is also supported by the presence of a peak in the hourly counts corresponding to the~~
~~morning rush hours. On the other hand, cluster 3~~ likely represents ~~the~~ particle size spectra mainly
shaped by primary and secondary aerosols advected from the most urbanised areas, i.e. ~~it can be is~~
~~most~~ likely associated ~~to~~ ~~with~~ the urban background of London.

~~The last two clusters are probably associated with road traffic: vehicle exhaust emissions peak in~~
~~the Aitken and accumulation modes with the size ranging from 20 nm to 500 nm (Vu et al., 2015b,~~
~~and references therein). Cluster 5 accounts for 14% of observations and reveals a unimodal~~
~~structure peaking at 25 nm. The hourly count pattern exhibits two maxima (6-8 am and 4-8 pm)~~
~~related to morning and evening rush hours. The wind rose shows that observations in this~~
~~cluster~~ ~~Cluster 5 for the warm season and cluster 3 for the cold season~~ ~~can~~ ~~may~~ be associated with
~~road traffic. They reveal modal structures with a dominant peak around 20-35 nm (cluster 5 also~~
~~shows a possible second peak at 15 nm) and~~ mostly occur when air masses blow from westerly
sectors, which are compatible with the location of motorways and Tunnel Rd, the main roadway
linking LHR to the M4 motorway. In ~~addition, it can be noted that~~ ~~summer,~~ the ~~wind rose hourly~~
~~count pattern~~ exhibits ~~high percentages of high speed winds from W. This pattern is compatible~~
~~with fresh road traffic emissions.~~

1273 ~~Cluster 4 represents 25% of total observations. It peaks at smaller particle sizes, but also shows a~~
1274 ~~wide hump at 50–150 nm.~~~~two. It is recognised that road traffic contributes to a large range (30–200~~
1275 ~~nm) of PNSD in the urban atmosphere (e.g., Yue et al., 2008; Costabile et al., 2009; Harrison et al.,~~
1276 ~~2011), which is compatible with this cluster spectrum. In addition, the hour count profile presents a~~
1277 ~~huge maximum during daytime with possibly 3 maxima (6–8 am and 4–8 pm) related to morning~~
1278 ~~and evening rush hours plus mid-afternoon); this pattern is the mirror image of those for clusters 2~~
1279 ~~and 3. The directional analysis shows increased levels when air masses move from the sectors more~~
1280 ~~affected by traffic: London (NE), M4 (N) and M25 (W) motorways and Tunnel Rd (W). It may~~
1281 ~~represent the typical spectra recorded during daytime and can be associated with aged~~
1282 ~~anthropogenic emissions, mostly due to road traffic.~~

1283 1284 3.2.2 — Cold season

1285 ~~Unfortunately, the atmospheric circulation during the cold season mostly experienced winds~~
1286 ~~blowing from the SW quadrant, while the NE sectors were poorly represented. As a consequence,~~
1287 ~~the limited extent of the wind directionality analysis may blur the interpretation of results. In~~
1288 ~~addition, the limited number of observations for air pollution advected from the Greater London~~
1289 ~~area may have affected the detection of the urban background.~~

1290
1291 ~~Clusters 1 and 5 account for 24% and 17% of total observations, respectively. They occur under~~
1292 ~~comparable wind regimes (from SW) and timing (increased counts during daytime). While~~
1293 ~~compatible with fresh road traffic emissions. However, the diurnal pattern of cluster 1 has the same~~
1294 ~~shape as the LHR aircraft movement profiles (Figure 2), cluster 5 is more comparable with cluster 1~~
1295 ~~for the warm season (maximum in the early afternoon). However, their spectra are quite different:~~
1296 ~~cluster 1 has a main mode at 20–25 nm, while cluster 5 peaks at 15 nm. Based on the prevailing~~
1297 ~~wind directionality, they can both be linked to airport activities. A close analysis of wind roses~~
1298 ~~reveals that cluster 5 occurs at significantly higher wind speed regimes than cluster 1 (Mann-~~

Whitney–Wilcoxon test at 0.05 significance level), in winter also presents a high number of counts at
 3–5 am, i.e. average wind speeds of 8.3 and 5.9 m s^{-1} , respectively, not directly compatible with rush
 hours. A possible interpretation is that cluster 5 represents fresher airport emissions (this may also
 explain the high similarity with the cluster 1 for the warm season), while cluster 1 depicts the
 airport emissions which have undergone more aging. The aging of freshly emitted particles in the
 atmosphere may involve condensation, evaporation and agglomeration processes and has been
 demonstrated to be a major mechanism in altering aerosol PNSD (e.g., Shi et al., 1999; Kim et al.,
 2004; Zhang et al., 2005; Zhou et al., 2005; Zhang et al., 2011; Harrison et al., 2016); this effect
 was also observed for particles emitted by road traffic in London (Dall'Osto et al., 2011). Another
 possible interpretation is that one cluster could represent the PNSD mainly influenced by aircraft
 engine emissions, while the other is related to other on-airport sources, e.g., airport ground service
 equipment, emissions from auxiliary power units (small on-board gas turbine engines) or ground
 power units provided by the airport. However, this latter interpretation is less probable, since the
 spatial extent and temporal pattern of these two sources is the same (airfield) and, thus, they are
 expected to be much better mixed.

Cluster 2 (16% of observations) extends over a wide size range (20 to 150 nm) and presents a daily
 pattern likely attributable to the dynamics of the mixing layer (the pattern is the mirror image of the
 ambient air temperature). In winter, there is an explanation involves the stronger effect of the winter
 mixing layer dynamics on the air quality due to the presence of more frequent low level thermal
 inversions, which may build up the pollutants at ground-level especially overnight. Consequently,
 this cluster cannot be linked to any specific primary anthropogenic source. This may increase the
 signal of the less intense, but still significant, nighttime traffic emissions present in the study area,
 and is likely representative of spectra mostly shaped by the drop of the mixing layer height and the
 formation of secondary aerosols.

Cluster 3 accounts for 20% of data during the cold season. The size spectrum, the wind rose and, partially, the hourly count profile well relates to cluster 5 for the warm season (attributed to fresh road traffic emissions). However, the diurnal pattern also presents a high number of counts at 3–5 am, i.e. not compatible with rush hours. Wood smoke is recognised to peak around 100 nm (e.g., Chandrasekaran et al., 2013; Vu et al., 2015b). A possible interpretation is that observations included in this cluster may represent PNSDs dominated by both traffic but influenced by domestic biomass combustion.

Cluster 4 (22%) peaks at 17 nm and also shows a wide hump at 50–150 nm. Its diurnal pattern shows a marked maximum occurring on the afternoon and is mostly represented under westerly winds regimes. Considering the differences between the two campaigns, it has similar characteristics to cluster 4 for the warm season. Thus, it can be interpreted as typical of spectra recorded during daytime and associated with the aging of anthropogenic emissions, mostly due to road traffic.

3.3 PMF Analysis

The best PMF solutions were identified: (i) by investigating solutions between 3 and 10 factors; (ii) by considering the minimization of the function Q with respect to the expected (theoretical) value and its stability over multiple ($n=100$) runs; (iii) by obtaining low values for the sum of the squares of the differences in scaled residuals for each base run pair by species; (iv) by minimizing the number of absolute scaled residuals over ± 3 and by keeping them symmetrically distributed; (v) by keeping the result uncertainties calculated by bootstrap (BS, $n=200$) and displacement (DISP) methods within an acceptable range (Paatero et al., 2014); (vi) by obtaining modelled total variable (PNC) successfully predicted ($R^2 > 0.9$ and slopes ≈ 1); and (vii) by avoiding the presence of edges in the G space plots (Paatero et al., 2002) and, then, the presence of hidden/unresolved sources.

1351 The interpretation of PMF results was then attempted by considering: (i) the knowledge of sources
 1352 impacting the study area; (ii) the comparison with the results reported by Vu et al. (2016), who
 1353 performed a PMF analysis of SMPS data collected in North Kensington (London urban
 1354 background); (iii) the shape of resulting profiles for both the particle number and volume
 1355 concentrations; (iv) the analysis of diurnal patterns; (v) the directional analysis using the polar plot
 1356 and ~~CBPF~~polar annuli; (vi) the correlations between the source contributions and the other air
 1357 pollutants monitored at the site or with weather variables, and (vii) the analysis of possible remote
 1358 source areas by applying the CWT model.
 1359
 1360 Six-factor solutions were extracted for both the seasons. The resulting factor profiles are presented
 1361 in Figures ~~65~~ and ~~76~~ for the warm and cold season, respectively. The factor profiles are expressed
 1362 as: (i) particle number concentrations and their DISP ranges; (ii) particle volume concentrations,
 1363 and (iii) explained variations showing how much of the variance (from 0 to 1) in the original dataset
 1364 is accounted for by each extracted factor. The ~~figures~~Figures 5 and 6 also show the diurnal patterns
 1365 and the polar plots computed ~~on~~from the hourly-averaged contributions. Table 1 summarises the
 1366 PMF results and spectral characteristics, while Table 2 shows the Pearson correlation matrices with
 1367 weather and air quality variables. Selected PMF solutions were very stable: no errors or unmapped
 1368 factors and few swaps (none in summer and <7% in winter) were found in BS; no swaps or errors
 1369 even at $dQ_{max}=25$ were found for DISP, i.e. solutions were affected by small rotational ambiguity
 1370 and, therefore, their interpretation can be considered robust.
 1371
 1372 DISP analysis is designed to explore the realistic bounds on the optimal (base run) PMF solutions
 1373 that do not result in appreciable increases in the Q values (Brown et al., 2015). In this study, the
 1374 ranges calculated by DISP for the $dQ=4$ were used to assess the uncertainty boundaries associated
 1375 ~~to~~with the final PMF profiles, as suggested in Zikova et al. (2016) and Masiol et al. (2017). This

strategy is useful to better interpret the results, as the regions of spectra affected by high rotational ambiguity are disclosed in the resulting profiles.

3.3.1 Warm season

Factor 1 includes most of the particles in the nucleation range (<20 nm), exhibits a sharp mode in the number distribution below the SMPS detection limit (14 nm) and makes the largest contribution to the total PNC (31.6%, DISP range 31-36%) (Figure 65). However, its contribution to the volume distribution is ~1%. Several studies report that particles in the nucleation range are emitted from the aircraft engines (e.g., Anderson et al., 2005; Herndon et al., 2008; Kinsey et al., 2010; Mazaheri et al., 2009;2013; Masiol and Harrison, 2014; Lobo et al., 2015) as well as from other anthropogenic (e.g., Schneider et al., 2005; Chen et al., 2011; Cheung et al., 2012; Stevens et al., 2012; Kumar et al., 2013;2014; Vu et al., 2015b) and natural (e.g., Kulmala et al., 1998; O'Dowd et al., 1998;1999; Kulmala and Kerminen, 2008; Riccobono et al., 2014) sources. This factor does not show any significant ($p < 0.05$) and strong ($r \geq |0.6|$) correlation with other measured species, but shows a weak ($|0.4| \leq r < |0.6|$) correlation with Factor 2, (Table 2). Its diurnal variation (Figure 65) shows higher concentrations between 6 am and 10 pm, and well agrees with the airport flight movements (Figure 2). The polar plot analysis also indicates enhanced levels when winds $> 2 \text{ m s}^{-1}$ blow from the airfield sectors (SW). All these insights are consistent with the location of Heathrow, i.e. the most plausible interpretation is related to the aircraft engine exhaust emissions. This interpretation is also supported by Keuken et al. (2015), which shows that the PNSD in an area affected by emissions from Schiphol airport (The Netherlands) is dominated by ultrafine (10-20 nm) particles. The large contribution of this factor to the total PNC is not surprising if compared to the results reported for the Los Angeles international airport by Hudda et al. (2014) (emissions from the airport increased PNC 4- to 5-fold at 8–10 km downwind the airfield). Since the airport of Los Angeles and LHR have comparable aircraft traffic, the quite high concentrations found in this study (on annual average nucleation particles are ~10 times higher than those measured in North Kensington urban

background by Vu et al. (2016)) are consistent with the sampling location chosen in this study (~1.2 km to the airfield). In addition, this result also agrees with previous studies on the impacts of LHR on local air quality; Carslaw et al. (2006) and Masiol and Harrison (2015) found comparable percent contributions of LHR emissions on NO₂ levels in the study area (approx. 25-30%). However, the lack of correlations with NO and NO₂ (tracers for aircraft emissions) is probably due to ~~the difference in the time resolution and~~ the presence of several other sources of nitrogen oxides in the area, such as the heavy traffic generated from the airport and from the nearby motorways.

Factor 2 is made up of ultrafine particles in the nucleation-Aitken range (one main peak at 20-35 nm) and accounts for 28% (DISP 25-30%) of PNC; its contribution to the volume distribution is low (~2%) and peaks at 22-45 nm and at 140-220 nm- [\(Figure 5; Table 1\)](#). Several insights seem to link this factor to road traffic emissions: (i) the modal structure; (ii) the strong association with morning and evening rush hours, and (iii) the significant increase for winds in the west and south-westerly sectors consistent with emissions generated from local busy roads close to LHR, Tunnel Rd. and M25 motorway. A similar mode in the nucleation range has been extensively attributed to the size distribution from road traffic (e.g., Vogt et al., 2003; Zhang et al., 2004; Ntziachristos et al., 2007; Vu et al., 2015b) and the growth of nucleation particles from diesel vehicles (Mayer and Ristovski, 2007; Wehner et al., 2009). For example, Charron and Harrison (2003) reported that particles in the range 30–60 nm show a stronger association with light-duty traffic at a traffic hotspot in central London (Marylebone Rd.); Janhäll et al. (2004) reported an average particle size distribution peaking at 15-30 nm during morning peak high traffic intensity in the city of Göteborg (Sweden), which has a car fleet comparable to the UK; Ntziachristos et al. (2007) found a sharp mode at 20-30 nm in sampling from engine exhausts. In addition, PMF factors with similar modal structures were found in other studies and were attributed to road traffic emissions: among others, Harrison et al. (2011) linked a factor peaking at 20 nm to primary road traffic emissions near a major UK highway; Masiol et al. (2016) measured PNSD in an international airport in Northern

Italy during summer and interpreted a factor with a clear mode at 35–40 nm as road traffic from the nearby city; Beddows et al. (2015) and Vu et al. (2016) found traffic factors with modal diameter at around 30 nm in an urban background site in London (North Kensington); Sowlat et al. (2016) reported a factor peaking at 20–40 nm in number concentration and at around 30–40 nm in volume concentration in Los Angeles (US) and interpreted it as traffic tailpipe emissions. However, this factor lacks significant positive correlations with primary road traffic tracers (nitrogen oxides, eBC; [Table 2](#)), while other studies have reported weak positive correlations with such species (Harrison et al., 2011; Masiol et al., 2016; Vu et al., 2016; Sowlat et al., 2016). Similarly to factor 1, this latter result may be due to the difference in the time resolution between chemical species and PNSD and the presence of several sources of nitrogen oxides in the area.

Factor 3 is mostly represented by 25–90 nm particles and contributes about 19% (17–21%) to the total number concentration- ([Figure 5; Table 1](#)). It also shows a second mode below the SMPS detection limit (14 nm), however, the DISP range clearly indicates that this part of the profile is affected by a large amount of rotational ambiguity, so that the presence of this second mode should be interpreted with caution. The volume concentration peaks at around 40–100 nm and 250–450 nm. The factor contribution is higher during rush hours, but the morning peak occurs 1 h later than in factor 2. The wind directionality shows increases for air masses blowing gently ($<4 \text{ m s}^{-1}$) from W and for calm wind periods, suggesting a quite local source; however, also an increase of concentrations is found for higher wind regimes ($>6 \text{ m s}^{-1}$) from the East (London). Factor 3 also shows significant positive correlations with NO (0.43) and NO₂ (0.61) ([Table 2](#)). All these insights seem to point to an aged road traffic source. This interpretation is also supported by Vu et al. (2016), who found a similar factor in London (North Kensington) peaking at ~20–100 nm. In this context, several source apportionment studies on PNSDs have attributed more than one factor to road traffic (e.g. Kasumba et al., 2009; Thimmaiah et al., 2009; Harrison et al., 2011; Liu et al., 2014; Al-Dabbous and Kumar, 2015; Vu et al., 2016; Sowlat et al., 2016). This result is not

surprising in areas where heavy traffic is widespread, as particles may undergo condensation, agglomeration, evaporation and dilution processes and, consequently, they may change modal characteristics in time and space. Such atmospheric processes are the main mechanisms reshaping PNSDs after primary exhausts ~~are~~is emitted into the atmosphere and have been discussed in several studies (Shi et al., 1999; Kim et al., 2004; Zhang et al., 2005; Zhou et al., 2005; Kulmala and Kerminen, 2008; Zhang et al., 2011; Harrison et al., 2016).

Factor 4 is made up of ~~ultrafine~~ particles over a wide range (50-200 nm with a clear mode at ~80 nm for PNC and 60-300 nm for PVC). The factor contributes 14% of PNC, but accounts for the main percentage of the volume concentration (33%). This factor ~~well~~ correlates well with gaseous pollutants linked to combustion sources (mostly road traffic), i.e. NO (0.6), NO₂ (0.76), and non-volatile primary pollutants, such as eBC (0.~~64~~62), NVPM_{2.5} (0.62) and EC (0.75) ([Table 2](#)). The factor also strongly correlates with OC (0.84) and sulphate (0.75). The diurnal pattern shows two main peaks in the morning and evening rush hours, ([Figure 5](#)), but the concentrations recorded between the two maxima are higher overnight than during daytime. This pattern suggests that both local emission sources and the dynamics of the mixing layer may play a key role in shaping its diurnal cycle, i.e. emitted pollutants undergo a wide dispersion within the expanded mixing layer during the daytime, while the drop of the mixing layer top occurring overnight restricts those pollutants to a layer close to ground level. The polar plot indicates increased levels for ~~wind~~ calm wind conditions or winds blowing from London (East sectors); in addition, the factor is strongly negatively correlated with wind speed (-0.64) ([Table 2](#)).

All these insights suggest that Factor 4 represents the fingerprint of the London pollution. Several studies carried out in London (Beddows et al., 2009;2015; Vu et al., 2016) and other megacities (e.g., New York: Masiol et al., 2017) have reported similar results, all interpreting this source profile as urban background (or urban accumulation mode). This source comprises both the solid

1480 particle mode from traffic emissions (Harrison et al., 2011; Pant and Harrison, 2013; Dall'Osto et
1481 al., 2012) and secondary species condensed upon pre-existing particles acting as condensation
1482 nuclei, including secondary sulphate, nitrate and organic aerosols. Secondary sulphate is formed
1483 through the atmospheric processing of local or distant SO₂ emissions (Kerminen et al., 2000) and
1484 neutralisation with ammonia (Benson et al., 2011). Nitrate aerosol is formed through the oxidation
1485 of NO₂ to nitrate and the consequent neutralization with ammonia (Seinfeld and Pandis, 2006) and
1486 occurs during both daytime and night-time; however the semivolatile nature of ammonium nitrate,
1487 makes its partitioning to the condensed-phase very weak. This behaviour also favours the
1488 occurrence of negative artefacts in filter-based sampling, which may explain the lack of significant
1489 correlations between the factor and the PM_{2.5}-bound nitrate (Table 2). On the contrary, the increase
1490 of the intensity of factor 4 during the night-time and the significant association with NO₂ are highly
1491 consistent with the chemistry driving the heterogeneous reactions of N₂O₅ and NO₃ on aerosol
1492 surfaces (Bertram and Thornton, 2009; Brown and Stutz, 2012). In view of this, Dall'Osto et al.
1493 (2009) reported that most nitrate particles in London are: (i) locally produced in urban locations
1494 during nighttime; (ii) mainly present in particles smaller than 300 nm and (iii) internally mixed with
1495 sulphate, ammonium, EC and OC.

1496
1497 Factors 5 and 6 make small contributions to PNC (4-7% and 1-4%, respectively), but are relevant
1498 for the volume concentration (37% and 21%, respectively). Factor 5 shows a main accumulation
1499 mode in number concentration at 110-250 nm and two more modes at ~30-70 nm and below 14 nm;
1500 [\(Figure 5; Table 1\)](#); however, the latter two modes suffer of large rotational ambiguity and should
1501 be interpreted with care. On the contrary, it exhibits a wide mode in volume concentration ranging
1502 from ~100 to ~500 nm. Factor 6 has two relevant modes in number concentration at 55-120 nm and
1503 230-400 nm, and two modes in volume concentration at 260-500 nm and 75-140 nm.

1504

1505 These factors still present two peaks corresponding to the rush hours, but the morning peak occurs
1506 1-2 h earlier than in the road traffic-related factors, i.e. when ambient temperature reaches its daily
1507 minimum. Both factors correlate well with secondary aerosol tracers (nitrate, sulphate, OC) and
1508 non-volatile components (eBC, EC, NVPM_{2.5}), but Factor 6 exhibits much higher correlation
1509 coefficients- (Table 2). Despite the polar plots indicating ~~the~~ main wind directionality toward N-E
1510 sectors, the analysis of air mass histories through the CWT model (Figure 87) clearly indicates likely
1511 continental origin areas rather than local sources.

1512
1513 Vu et al. (2016) observed two factors in North Kensington with very similar modal structures, daily
1514 patterns, correlations with PM_{2.5}-bound species and external source areas maps. Therefore, their
1515 interpretation is confirmed also in this study, i.e. mixed secondary aerosol (Factor 5) and inorganic
1516 secondary aerosol (Factor 6). Both factors are clearly originated from ~~the~~ continental Europe and
1517 are consistent with a previous receptor modelling study carried out in a rural background site
1518 representative of ~~the~~ southern UK (Charron et al., 2013). Similar origin and formation mechanisms
1519 also explain their strong correlation (0.75). ~~Despite~~ ~~Although~~ it is not reasonable ~~to~~ extract ~~much~~
1520 more information from these data due to the short period ~~into account of sampling~~ and the large
1521 uncertainty associated with back-trajectory analysis, it can be observed that Factor 5 shows a wide
1522 source area all over ~~the~~ Central Europe, while Factor 6 exhibits two distinct hotspots (Central and
1523 North-eastern Europe).

1524
1525

1526 3.3.2 Cold season

1527 The 6 factors identified during the cold period (Figure 76) are similar to those for the warm season.
1528 Factor 1 is composed of a high proportion of particles in the nucleation range with a sharp mode at
1529 ~15 nm. It accounts for 33% (32-35%) of PNC and less than 2% of PVC. The polar plot reveals
1530 increased concentrations for moderate winds blowing from the airport sector and the diurnal pattern

1531 is also compatible with the aircraft traffic. No statistically significant correlations are found with
 1532 any other monitored species- [\(Table 3\)](#). Therefore, Factor 1 may be attributed to the airport
 1533 emissions related to ~~the~~ aircraft engine exhausts ~~emissions~~. As in the warm season, factor 1 is
 1534 moderately correlated with factor 2 (fresh road traffic, $r=0.55$), indicating a quite clear relationship
 1535 between the two sources.
 1536
 1537 *Factor 2* represents particles in the 15-35 nm range of number concentration, accounting for 35%
 1538 (33-37%) of total PNC- [\(Figure 6; Table 1\)](#). Its importance for volume concentration is
 1539 ~~modest~~[minimal](#) (3%) with two modes at 30 and 200 nm. The diurnal pattern and the wind
 1540 directionality are compatible with LHR as a source and it shows a weak positive correlation with
 1541 NO₂ (0.42) and a strong correlation with nitrate (0.63) [\(Table 3\)](#). Despite its similarity and
 1542 relationship with Factor 1 and the consequent similar potential origin, Factor 2 may represent a
 1543 different source: Factors 1 and 2 remain clearly separated even at solutions down to 4 factors,
 1544 demonstrating their structural robustness and the lack of potential artefacts ~~upon~~[affecting](#) the PMF
 1545 solution. Consequently, it can be concluded that they do not represent over-resolved solutions (i.e.
 1546 factor splitting). The most plausible interpretation for Factor 2 is therefore the same as for the warm
 1547 season, i.e. fresh road traffic emissions. Furthermore, this factor can be attributed to the road traffic
 1548 generated by the airport and nearby major roads.
 1549
 1550 *Factor 3* includes most of the particles in the Aitken range and accounts for 19% (18-20%) of PNC.
 1551 Its contribution to particle volume concentration is relevant (9%) with a main peak at around 100 nm
 1552 and a secondary peak at 400 nm- [\(Table 1\)](#). It presents two rush hours peaks and the polar plot
 1553 reveals an origin from the SW quadrant. However, as with the warm period, the wind directionality
 1554 suggests increases for slower wind regimes than the fresh road traffic factor and for more westerly
 1555 sectors, which are not compatible with the airfield location. Since factor 3 ~~well~~-correlates [well](#)
 1556 [\(Table 3\)](#) with a number of other pollutants linked to primary emissions from road traffic (NO

(0.51), NO₂ (0.81), [eBC \(0.52\)](#), PM_{2.5} (0.53), OC (0.79) and EC (0.83)), it represents a second road traffic factor, more affected by aging in the atmosphere than factor 2.

Despite the wind regimes from NE sectors being poorly represented during the cold campaign, *Factor 4* is the only one showing a possible origin from London and for calm wind periods. As with the warm season, it is composed of a wide range of particles encompassing the Aitken and accumulation modes (50 to 150 nm), while the peak in volume concentration is at 170 nm- ([Table 1](#)). The diurnal pattern ([Figure 6](#)) is clearly related to the mixing layer dynamics and the correlation analysis reveals strong relationships with many species (NO, NO₂, eBC, [Delta-C](#), NVPM_{2.5}, OC, EC, nitrate, ammonium and potassium; [Table 3](#)). Consequently, it is concluded that it represents the urban accumulation mode, whose contribution to the total volume concentration is also similar to the warm season (33%). It is interesting to note the large similarity with the urban accumulation mode found in the warm season, from which it differs slightly only in the diurnal pattern (higher overnight) and in the presence of a strong correlation with nitrate ($r=0.88$), [possibly](#) due to the lesser extent of negative artefacts on PM_{2.5} filter samples.

The last two factors are interpreted as due to secondary aerosols. Their modal structures, their contributions to total PNC and PVC, and their correlations with PM_{2.5}-bound species ([Table 3](#); [Figure 6](#)) largely reflect the results obtained for the warm period. However, the CWT maps ([Figure 8](#)) highlight different source areas, i.e. the origin of the secondary aerosols is regional (UK and Northern Europe). In addition, the presence of strong positive correlations with chloride may also indicate a contribution from the transport of sea-salt aerosol.

3.3 Comparison of *k*-means and PMF

The cluster analysis revealed the presence of 5 characteristic PNSD shapes during both the seasons. These spectra have been linked to potential sources in the study area, i.e. road traffic, airport

activities, ~~biomass burning~~ and secondary aerosol formation processes. However, the cluster analysis is mostly driven by the ~~size~~-spectral ~~size~~ regions with higher particle number concentrations, i.e. it has the disadvantage of partitioning the single observations predominantly according to the finest region of the size distribution. This limitation is well illustrated by the poor (almost null) separation of clusters based on the particle volume distributions (all clusters showed quite similar particle volume spectra). In addition, cluster analysis also has the disadvantage of linking each cluster to a single source and does not easily account for PNSD resulting from the mix of two or more different sources.

In contrast, the PMF analysis computed over the PNSD also accounts well for the sources with a small impact on the number distribution, but having a larger influence on the particle volume size distributions and, therefore, on the particle mass concentration. Despite the differences in the two methods, some further information can be extracted by combining the results of cluster and PMF analysis. Figure 98 shows the statistics of normalised PMF source contributions relating to each single cluster. ~~Generally, the two methods well agree for the “airport” source, pointing out how much the airport-related emissions may shape the PNSD in the study area.~~

For the warm period, significantly higher (0.05 significance) PMF contributions of the airport factor (F1) are measured for cluster 1, ~~i.e. (average normalised contribution ~3.5). This result indicates that~~ the airport fingerprint was well ~~caught-captured~~ by both source apportionment methods. During the cold season, the airport factor (F1) is ~~significantly higher~~ ~~duringfor~~ both clusters 1 and 5. ~~(average normalised contributions of ~2 and ~3, respectively).~~ While cluster 5 presents significant high PMF contributions only for factor 1, cluster 1 also shows high contributions of factor 2 (fresh road traffic). This result indicates that cluster 5 may be linked as the typical PNSD spectra for airport emissions, while cluster 2 likely represents mixed emissions from aircraft and airport-related traffic. A close analysis of wind roses for the two clusters in the cold season (Figure 4) reveals that

1609 [cluster 5 occurs at significantly higher wind speed regimes than cluster 1 \(Mann-Whitney-Wilcoxon](#)
1610 [test at 0.05 significance level\)](#), i.e. average wind speeds of 8.3 and 5.9 m s⁻¹, respectively. As a
1611 [consequence, the different wind regimes may well be likely responsible for the split between the](#)
1612 [two clusters.](#)

Formatted: Font color: Red

1614 Results for fresh traffic emissions also agree between the two methods. Factors 2 exhibit the higher
1615 normalised contributions to clusters 5 ([normalised contribution 2.5](#)) and 1 ([normalised contribution](#)
1616 [~3](#)) for the warm and cold period, respectively: ([Figure 8](#)). However, in winter it is evident that
1617 PNSDs grouped on cluster 1 are also strongly influenced by airport emissions, probably due to the
1618 lower mixing layer height and, thus, a lesser dispersion in the atmosphere.

1620 Clusters 4 for both the periods show enrichments in the contributions for 4 PMF sources (aged road
1621 traffic, urban accumulation and the two secondary aerosols) ([Figure 8](#)). This further emphasises that
1622 cluster 4 represents the typical PNSD during daytime resulting from the mixing of different sources.
1623 In a similar way, clusters 3 and 2 in the warm and cold periods, respectively, represent the typical
1624 nighttime spectra: ([Figures 3 and 4](#)), i.e. they exhibit similar partitioning over the PMF sources and
1625 similar daily cycles.

1627 3.4 Analysis of a Large Regional Nucleation Event

1628 ~~Regional photochemical nucleation episodes are regularly recorded in the Southern and Eastern~~
1629 ~~UK. Their general characteristics have been reported in a number of studies (e.g., Alam et al., 2003;~~
1630 ~~Charron et al., 2007;2008; Beddows et al., 2015; Vu et al., 2016) and can be summarised as~~
1631 ~~follows: (i) particle modality at around 20 nm; (ii) higher frequency around noon in association with~~
1632 ~~the peak in actinic flux intensities; (iii) clear seasonal cycles (higher average contribution levels in~~
1633 ~~the summer, from June to September); (iv) marked directionality from the westerly sectors;~~

~~reflecting maritime atmospheric circulation regimes, with high wind speed and low PM_{2.5} concentrations.~~
Regional photochemical nucleation episodes are regularly recorded in the Southern and Eastern UK. Their general characteristics have been reported in a number of studies (e.g., Alam et al., 2003; Charron et al., 2007; 2008; Beddows et al., 2015; Vu et al., 2016) and can be summarised as follows: (i) particle modality at around 20 nm; (ii) higher frequency around noon in association with the peak in actinic flux intensities; (iii) clear seasonal cycles (higher average contribution levels in the summer, from June to September); (iv) marked directionality from the westerly sectors, reflecting maritime atmospheric circulation regimes, with high wind speed and low PM_{2.5} concentrations.

A strong regional nucleation event occurred during the warm period sampling campaign (starting on 7th September at 1 pm UTC and lasting for about 12 h). Increases of PNC were almost simultaneously recorded at Harlington and at Harwell, a national network rural background site located approx. 60 km WNW of LHR and representative of the regional background levels of air pollution across the Southern UK. The comparison of PNC time series at the two sites is provided as Figure ~~SI7~~[SI10](#). Figure ~~409~~[409](#) shows the contour plots of SMPS data recorded at Harlington between 7th and 8th September as well as the hourly averaged concentrations of nucleation, Aitken and accumulation particles, TEOM-FDMS PM_{2.5} mass and the contributions of Factors 1 to 4 extracted by the PMF. ~~The figure~~[Figure 9](#) also reports the hourly counts of number of clusters extracted by the *k*-means analysis. The contour plot shows a typical “banana” shape with particle mode growing from ~20 nm (1 pm) to ~100 nm (overnight). The episode strongly influenced the PNSDs until around midnight; however its effect is also visible over the first half of 8th September. The time series (Figure ~~409~~[409](#)) exhibits a clear peak in nucleation particles between 1 pm and 3 pm followed by peaks of Aitken (3-11 pm) and accumulation mode (8 pm-2 am) particles. The back-trajectory analysis (Figure ~~411~~[SI11](#)) ~~has~~ revealed that the event occurred when north-westerly fresh

Formatted: Tab stops: Not at 1.5 cm

(and clean) maritime air masses were advected from the Atlantic. This is also supported by the
 PM_{2.5} mass, which exhibited a fast drop of concentrations just a few hours before the event, ~~(-30 μg~~
 ~~m^{-3} in 3 hours, i.e. from 40 $\mu\text{g m}^{-3}$ at 6 am to 10 $\mu\text{g m}^{-3}$ at 9 am, Figure 9),~~ probably reducing the
 condensation sink and facilitating nucleation.

Both atmospheric nucleation and aircraft engines are recognised to produce particles in the
 nucleation range. The analysis of this single –but strong– episode gives insights into how much the
 source apportionment results can potentially be affected by regional nucleation. This latter analysis
 is possible because the wind directionality during the entire episode was from N sectors, i.e. the
 contribution of LHR can be considered negligible.

The results of cluster analysis were ~~just slightly~~ affected by the event. Before the episode, the PNSD
 spectra were mostly categorised as clusters 3 and 4 (urban background and daytime pollution,
 respectively), ~~while a few i.e. the clusters (less than 1 h of observations) mostly recorded under~~
~~north-easterly wind regimes (Figure 3). About 50% and 30% of the clusters~~ were ~~then~~ categorised
 as “airport” ~~during~~ the ~~beginning~~ first and second hour of the episode, ~~respectively (Figure 9).~~
~~Since the wind directionality is inconsistent with an origin from the airfield, this categorisation is~~
~~likely the result of the nucleation event.~~ The growing of particles in the ~~subsequent~~ hours ~~was then~~
~~identified after the beginning of the event has further driven the cluster results: (i) about 60-80% of~~
~~PNSDs were categorised~~ as “fresh road traffic” (cluster 5) ~~and after 2-3 hours, and (ii) 80-100% of~~
~~PNSDs were clustered as~~ “nighttime regional pollution” (cluster 2) ~~after 4-6 hours.~~ In a similar
 way, PMF results were ~~slightly~~ affected by the event, (Figure 9), with a sharp increase of
 contribution levels for: (i) factor 1 (airport) ~~and, then, for factors~~ from 1.5- $\times 10^3$ particles cm^{-3} at
 noon to 13.3- $\times 10^3$ particles cm^{-3} at 2 pm; (ii) factor 2 (fresh road traffic) ~~and from~~ 0.5- $\times 10^3$ particles
 cm^{-3} at 1 pm to 21- $\times 10^3$ particles cm^{-3} at 3 pm; and (iii) factor 3 (aged road traffic) ~~from~~ 2.1- $\times 10^3$
 particles cm^{-3} at 2 pm to approx. 15- $\times 10^3$ particles cm^{-3} at 5-6 pm.

1686

1687 This episode was the main nucleation event recorded during the two sampling campaigns. Other
1688 possible episodes also occurred (mostly during the warm season), but they were much less
1689 significant and often hard to detect. This qualitative analysis points to some conclusions: (i)
1690 regional photochemical nucleation events may have an effect on clustering and PMF results; (ii) the
1691 effect may lead to an “additive” bias, mostly over the “airport” and “road traffic” factors and
1692 clusters; (iii) the effect of regional nucleation events in the study area is largely overwhelmed by the
1693 strength of local sources, but in other locations with more frequent nucleation events it may be more
1694 important to identify and separate them.

1695

1696 4 CONCLUSIONS

1697 The effect of airport emissions upon the particle number concentration and size distribution was
1698 assessed at a site close to a major European airport (Heathrow) serving a megacity (London). The
1699 conclusions to be drawn are:

- 1700 • ~~Anomalously high~~High particle number concentrations were recorded for the finest sizes
1701 (nucleation <30 nm and Aitken nuclei 30-100 nm) if compared to an urban background site in
1702 London (N. Kensington).
- 1703 • Polar plot analysis indicates that Heathrow is a strong potential source for NO₂, nucleation and
1704 Aitken particles, but its contribution to the mass concentration of PM_{2.5} and eBC is very small.
1705 On the contrary, the urban area of London ~~area seems appears~~ to be ~~a main~~ the main source for
1706 PM and eBC.
- 1707 • The *k*-means cluster analysis has revealed that 20% of PNSDs are mostly shaped by airport
1708 direct emissions, but particle size spectra are also strongly affected by other local sources
1709 (mostly fresh and aged road traffic during daytime) and the reduction of mixing layer depth
1710 (during nighttime). Typical PNSD spectra have been identified for nighttime and daytime
1711 pollution as well. Such spectra are likely the result of multiple source mixtures.

- 1712 • PMF analysis revealed that the fingerprint of Heathrow has a peculiar modal structure peaking
1713 at <20 nm. The direct airport emissions account for 30-35% of total particles in both the
1714 seasons. Such results are in line with percent estimations for NO₂ reported in previous studies.
- 1715 • Other major contributors to PNC are fresh (24-36%) and aged (16-21%) road traffic emissions.
1716 Despite both applied source apportionment methods failing to fully disaggregate the emissions
1717 from the local traffic (including motorway) and traffic generated by the airport, results suggest
1718 that road traffic sources may contribute to the total PNC more than Heathrow (40-56%).
1719 However, making a clear distinction between the influence of traffic generated by the airport
1720 from other road traffic is not feasible from this analysis.
- 1721 • ~~The fingerprint of London has~~An urban accumulation mode was found. This source presents a
1722 wide mode between 50-150 nm.~~This urban accumulation mode and~~ accounts for around 10%
1723 of PNC~~and. The wind directionality is the result of~~consistent with the advection of air masses
1724 from ~~the city~~London. It is more evident overnight due to the drop of the mixing layer top, the
1725 subsequent increase in air pollutants at ground level and the generation of nighttime secondary
1726 nitrate aerosols.
- 1727 • Secondary sources accounted for less than 6% in number concentrations but for more than 50%
1728 in volume concentration. Long-range transport has a key role in advecting polluted air masses
1729 from mainland Europe.

1731 ACKNOWLEDGEMENTS

1732 This study was carried out under the Marie Skłodowska-Curie project ‘CHEERS’ (Chemical and
1733 Physical Properties and Source Apportionment of Airport Emissions in the context of European Air
1734 Quality Directives, call: FP7-PEOPLE-2012-IEF, project no. 328542). The authors gratefully
1735 acknowledge: (i) the European Union for funding the Marie Curie Intra-European Fellowship for
1736 career development to M. Masiol through the project ~~entitled ‘Chemical and Physical Properties and~~
1737 ~~Source Apportionment of Airport Emissions in the context of European Air Quality Directives~~

1738 | ~~(Project CHEERS, call: FP7 PEOPLE 2012 IEF, project no. 328542); 'CHEERS'~~; (ii) Heathrow
1739 | Airport Ltd and Ricardo-AEA for supplying aircraft movement data and for the valuable exchange
1740 | of information and discussion, in particular Katherine Rolfe, Elizabeth Hegarty (Heathrow), Brian
1741 | Stacey (Ricardo-AEA) and David Vowles; (iii) DEFRA Automatic Urban and Rural Network, and
1742 | London Air Quality Network for providing pollutant data; (iv) Met Office and BADC for weather
1743 | data; (v) the NOAA Air Resources Laboratory (ARL) for the provision of the HYSPLIT transport
1744 | and dispersion model used in this publication; ~~and~~ (vi) [the UK Department for Transport, Road](#)
1745 | [Traffic and Road Freight Statistics, for providing M4 and M25 traffic data; and \(vii\)](#) Dr. Stefania
1746 | Squizzato (~~Clarkson~~ [University of Rochester, NY](#), USA) for the valuable exchange of information.

1747

1748

1749 **REFERENCES**

- 1750 ACI (Airport Council International): ACI releases preliminary world airport traffic rankings.
 1751 Airports Council International, Montreal. Available at: [http://www.aci.aero/News/Releases/Most-](http://www.aci.aero/News/Releases/Most-Recent/2016/04/04/ACI-releases-preliminary-world-airport-traffic-rankings-)
 1752 Recent/2016/04/04/ACI-releases-preliminary-world-airport-traffic-rankings- [last accessed: June
 1753 2016].
 1754
 1755 AEA: Heathrow Airport Air Quality Modelling for 2008/9: Results and Model Evaluation. Report
 1756 by AEA Energy & Environment on behalf of BAA, July 2010. AEAT/ENV/R/2948/Issue 1.
 1757
 1758 Al-Dabbous, A. N., Kumar, P.: Source apportionment of airborne nanoparticles in a Middle Eastern
 1759 city using positive matrix factorization, *Environ. Sci. Process Impacts*, 17, 802-812, 2015.
 1760
 1761 Alam, A., Shi, J. P. and Harrison, R. M.: Observations of new particle formation in urban air, *J.*
 1762 *Geophys. Res.*, 108, 4093-4107, 2003. doi:10.1029/2001JD001417
 1763
 1764 Anderson, B. E., Branham, H.-S., Hudgins, C. H., Plant, J. V., Ballenthin, J. O., Miller, T. M.,
 1765 Viggiano, A. A., Blake, D. R., Boudries, H., Canagaratna, M., Miake-Lye, R. C., Onasch, T.,
 1766 Wormhoudt, J., Worsnop, D., Brunke, K. E., Culler, S., Penko P., Sanders, T., Han, H.-S., Lee, P.,
 1767 Pui, D. Y. H., Thornhill, K. L., Winstead, E. L.: Experiment to Characterize Aircraft Volatile
 1768 Aerosol and Trace-Species Emissions (EXCAVATE), NASA/TM-2005-213783, National
 1769 Aeronautics and Space Administration, Hampton, VA., 2005.
 1770
 1771 Anttila, P., Tuovinen, J. P., Niemi, J. V.: Primary NO₂ emissions and their role in the development
 1772 of NO₂ concentrations in a traffic environment, *Atmos. Environ.*, 45, 986-992, 2011.
 1773
 1774 Atkinson, R. W., Fuller, G. W., Anderson, H. R., Harrison, R. M., Armstrong, B.: Urban ambient
 1775 particle metrics and health: a time-series analysis, *Epidemiol.*, 21, 501-511, 2010.
 1776
 1777 Beddows, D. C. S., Dall'Osto, M., Harrison, R. M.: Cluster analysis of rural, urban and curbside
 1778 atmospheric particle size data, *Environ. Sci. Technol.*, 43, 4694-4700, 2009.
 1779
 1780 Beddows, D. C. S., Dall'Osto, M., Harrison, R. M., Kulmala, M., Asmi, A., Wiedensohler, A., Laj,
 1781 P., Fjaeraa, A. M., Sellegri, K., Birmili, W., Bukowiecki, N., Weingartner, E., Baltensperger, U.,
 1782 Zdimal, V., Zikova, N., Putaud, J.-P., Marinoni, A., Tunved, P., Hansson, H.-C., Fiebig, M.,
 1783 Kivekäs, N., Swietlicki, E., Lihavainen, H., Asmi, E., Ulevicius, V., Aalto, P. P., Mihalopoulos, N.,
 1784 Kalivitis, N., Kalapov, I., Kiss, G., de Leeuw, G., Henzing, B., O'Dowd, C., Jennings, S. G., Flentje,
 1785 H., Meinhardt, F., Ries, L., Denier van der Gon, H. A. C., Visschedijk, A. J. H.: Variations in
 1786 tropospheric submicron particle size distributions across the European continent 2008-2009,
 1787 *Atmos. Chem. Phys.*, 14, 4327-4348, 2014.
 1788
 1789 Beddows D. C. S., Harrison R. M., Green D. C. and Fuller G. W.: Receptor modelling of both
 1790 particle composition and size distribution from a background site in London, UK., *Atmos. Chem.*
 1791 *Phys.*, 15, 10107-10125, 2015.
 1792
 1793 Belis, C. A., Larsen, B. R., Amato, F., El Haddad, I., Favez, O., Harrison, R. M., Hopke, P. K.,
 1794 Nava, S., Paatero, P., Prévôt, A., Quass, U., Vecchi, R. and Viana, M.: European guide on air
 1795 pollution source apportionment with receptor models, JRC Reference Reports EUR26080 EN,
 1796 2014.
 1797

1798 Benson, D. R., Yu, J. H., Markovich, A., Lee, S.-H.: Ternary homogeneous nucleation of H₂SO₄,
1799 NH₃, and H₂O under conditions relevant to the lower troposphere, *Atmos. Chem. Phys.*, 11, 4755-
1800 4766, 2011.

1801 Bertram, T. H. and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on
1802 aqueous particles: the competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem.*
1803 *Phys.*, 9, 8351-8363, 2009.

1804
1805 Bigi A and Harrison R. M.: Analysis of the air pollution climate at a central urban background site,
1806 *Atmos. Environ.*, 44, 2004-2012, 2010.

1807
1808 Brines, M., Dall'Osto, M., Beddows, D. C. S., Harrison, R. M. and Querol, X.: Simplifying aerosol
1809 size distributions modes simultaneously detected at four monitoring sites during SAPUSS, *Atmos.*
1810 *Chem. Phys.*, 14, 2973-2986, 2014.

1811
1812 Brines, M., Dall'Osto, M., Beddows, D., Harrison, R., Gómez-Moreno, F., Núñez, L., Artíñano, B.,
1813 Costabile, F., Gobbi, G. And Salimi, F.: Traffic and nucleation events as main sources of ultrafine
1814 particles in high-insolation developed world cities, *Atmos. Chem. Phys.*, 15, 5929-5945, 2015.

1815
1816 Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41,
1817 6405-6447, 2012.

1818
1819 Brown, S. G., Eberly, S., Paatero, P. and Norris, G. A.: Methods for estimating uncertainty in PMF
1820 solutions: Examples with ambient air and water quality data and guidance on reporting PMF results,
1821 *Sci.Total Environ.*, 518, 626-635, 2015.

1822
1823 Carslaw, D. C. and Ropkins, K.: Openair - an R package for air quality data analysis, *Environ.*
1824 *Model. Softw.*, 27-28, 52-61, 2012.

1825
1826 Carslaw, D. C., Beevers, S. D., Ropkins, K. and Bell, M. C.: Detecting and quantifying aircraft and
1827 other on-airport contributions to ambient nitrogen oxides in the vicinity of a large international
1828 airport, *Atmos. Environ.*, 40, 5424-5434, 2006.

1829
1830 Carslaw, D. C., Beevers, S. D. and Bell, M. C.: Risks of exceeding the hourly EU limit value for
1831 nitrogen dioxide resulting from increased road transport emissions of primary nitrogen dioxide,
1832 *Atmos. Environ.*, 41, 2073-2082, 2007.

1833
1834 Chandrasekaran, S. R., Hopke, P. K., Newtown, M. and Hurlbut, A.: Residential-scale biomass
1835 boiler emissions and efficiency characterization for several fuels, *Energy & Fuels*, 27, 4840-4849,
1836 2013.

1837
1838 Charron, A. and Harrison, R. M.: Primary particle formation from vehicle emissions during exhaust
1839 dilution in the roadside atmosphere, *Atmos. Environ.*, 37, 4109-4119, 2003.

1840
1841 Charron, A., Degrendele, C., Laongsri, B. and Harrison, R. M.: Receptor modelling of secondary
1842 and carbonaceous particulate matter at a southern UK site, *Atmos. Chem. Phys.* 13, 1879-1894,
1843 2013.

1844
1845 Charron, A., Birmili, W. and Harrison, R. M.: Factors influencing new particle formation at the rural
1846 site, Harwell, United Kingdom, *J. Geophys. Res.*, 112, D14210, 2007. doi:10.1029/2007JD008425.

1847
1848

1849 Charron, A., Birmili, W. and Harrison, R. M.: Fingerprinting particle origins according to their size
1850 distribution at a UK rural site, *J. Geophys. Res.*, 113, D07202, 2008. doi:10.1029/2007JD008562.
1851

1852 Chen, J. P., Tsai, T. S. and Liu, S. C.: Aerosol nucleation spikes in the planetary boundary layer,
1853 *Atmos. Chem. Phys.*, 11, 7171-7184, 2011.
1854

1855 Cheung, H. C., Morawska, L., Ristovski, Z. D., and Wainwright, D.: Influence of medium range
1856 transport of particles from nucleation burst on particle number concentration within the urban
1857 airshed, *Atmos. Chem. Phys.*, 12, 4951-4962, 2012.
1858

1859 Clapp, L. J. and Jenkin, M. E.: Analysis of the relationship between ambient levels of O₃, NO₂ and
1860 NO as a function of NO_x in the UK, *Atmos. Environ.*, 35, 6391-6405, 2001.
1861

1862 Costabile, F., Birmili, W., Klose, S., Tuch, T., Wehner, B., Wiedensohler, A., Franck, U.,
1863 König, K. and Sonntag, A.: Spatio-temporal variability and principal components of the particle
1864 number size distribution in an urban atmosphere, *Atmos. Chem. Phys.*, 9, 3163-3195, 2009.
1865

1866 Cyrys, J., Eeftens, M., Heinrich, J., Ampe, C., Armengaud, A., Beelen, R., Bellander, T.,
1867 Beregszaszi, T., Birk, M., Cesaroni, G., Cirach, M., de Hoogh, K., De Nazelle, A., de Vocht, F.,
1868 Declercq C., Dedele, A., Dimakopoulou, K., Eriksen, K., Galassi, C., Grauleviciene, R., Grivas, G.,
1869 Gruzieva, O., Hagenbjörk Gustafsson, A., Hoffmann, B., Iakovides, M., Ineichen, A., Krämer, U.,
1870 Lanki, T., Lozano, P., Madsen, C., Meliefste, K., Modig, L., Mölterm, A., Mosler, G.,
1871 Nieuwenhuijsen, M., Nonnemacher, M., Oldenwening, M., Peters, A., Pontet, S., Probst-Hensch,
1872 N., Quass, U., Raaschou-Nielsen, O., Ranzi, A., Sugiri, D., Stephanou, E.G., Taimisto, P., Tsai, M.-
1873 Y., Vaskövi, E., Villani, S., Wang, M., Brunekreef, B. and Hoek, G.: Variation of NO₂ and NO_x
1874 concentrations between and within 36 European study areas: Results from the ESCAPE study,
1875 *Atmos. Environ.*, 62, 374-390, 2012.
1876

1877 Dall'Osto, M., Harrison, R. M., Coe, H., Williams, P. I. and Allan, J.D.: Real time chemical
1878 characterization of local and regional nitrate aerosols, *Atmos. Chem. Phys.*, 9, 3709-3720, 2009.
1879

1880 Dall'Osto, M., Thorpe, A., Beddows, D. C. S., Harrison, R. M., Barlow, J. F., Dunbar, T., Williams,
1881 P.I. and Coe, H.: Remarkable dynamics of nanoparticles in the urban atmosphere, *Atmos. Chem.*
1882 *Phys.*, 11, 6623-6637, 2011.
1883

1884 Dall'Osto, M., Beddows, D. C. S., Pey, J., Rodriguez, S., Alastuey, A., Harrison, R. M. and Querol,
1885 X.: Urban aerosol size distributions over the Mediterranean city of Barcelona, NE Spain, *Atmos.*
1886 *Chem. Phys.*, 12, 10693-10707, 2012.
1887

1888 DEFRA: Air Pollution in the UK 2015. UK Department for Environment, Food and Rural Affairs.
1889 Issue of September 2016. Available at: [https://uk-](https://uk-air.defra.gov.uk/assets/documents/annualreport/air_pollution_uk_2015_issue_1.pdf)
1890 [air.defra.gov.uk/assets/documents/annualreport/air_pollution_uk_2015_issue_1.pdf](https://uk-air.defra.gov.uk/assets/documents/annualreport/air_pollution_uk_2015_issue_1.pdf) (last accessed:
1891 November 2016).
1892

1893 Dodson, R. E., Houseman, E. A., Morin, B. and Levy, J. I.: An analysis of continuous black carbon
1894 concentrations in proximity to an airport and major roadways, *Atmos. Environ.*, 43, 3764-3773,
1895 2009.
1896

1897 Farias, F. and ApSimon, H.: Relative contributions from traffic and aircraft NO_x emissions to
1898 exposure in West London, *Environ. Modell. Softw.*, 21, 477-485, 2006.
1899

1900 Finlayson-Pitts, B. J. and Pitts Jr, J. N.: Chemistry of the upper and lower atmosphere: theory,
 1901 experiments, and applications. Academic press, 2000.

1902

1903 Grice, S., Stedman, J., Kent, A., Hobson, M., Norris, J., Abbott, J., Cooke S.: Recent trends and
 1904 projections of primary NO₂ emissions in Europe, Atmos. Environ., 43, 2154-2167, 2009.

1905

1906 Harrison, R. M., Beddows, D. C. S. and Dall'Osto, M.: PMF Analysis of wide-range particle size
 1907 spectra collected on a major highway, Environ. Sci. Technol., 45, 5522-5528, 2011.

1908

1909 [Harrison, R.M., Dall'Osto, M., Beddows, D.C.S., Thorpe, A.J., Bloss, W.J., Allan, J.D., Coe, H.,](#)
 1910 [Dorsey, J.R., Gallagher, M., Martin, C. and Whitehead, J.: Atmospheric chemistry and physics in](#)
 1911 [the atmosphere of a developed megacity \(London\): an overview of the REPARTEE experiment and](#)
 1912 [its conclusions. Atmos. Chem. Phys., 12\(6\), 3065-3114, 2012.](#)

1913

1914 Harrison, R. M., Beddows, D. C., Jones, A. M., Calvo, A., Alves, C. and Pio, C.: An evaluation of
 1915 some issues regarding the use of aethalometers to measure woodsmoke concentrations, Atmos.
 1916 Environ., 80, 540-548, 2013.

1917

1918 Harrison, R. M., Jones, A. M., Beddows, D. C. S., Dall'Osto, M. and Nikolova, I.: Evaporation of
 1919 traffic-generated nanoparticles during advection from source, Atmos. Environ., 125, 1-7, 2016.

1920

1921 Herndon, S. C., Jayne, J. T., Lobo, P., Onasch, T. B., Fleming, G., Hagen, D. E., Whitefield, P. D.
 1922 and Miake-Lye, R. C.: Commercial aircraft engine emissions characterization of in-use aircraft at
 1923 Hartsfield-Jackson Atlanta International Airport, Environ. Sci. Technol., 42, 1877-1883, 2008.

1924

1925 Hopke, P. K.: Review of receptor modeling methods for source apportionment. JAWMA, 66, 237-
 1926 259, 2016.

1927

1928 [Hsu, H.H., Adamkiewicz, G., Houseman, E.A., Vallarino, J., Melly, S.J., Wayson, R.L., Spengler,](#)
 1929 [J.D. and Levy, J.I.: The relationship between aviation activities and ultrafine particulate matter](#)
 1930 [concentrations near a mid-sized airport, Atmos. Environ., 50, 328-337, 2012a.](#)

1931

1932 [Hsu, H.H., Adamkiewicz, G., Houseman, E.A., Vallarino, J., Melly, S.J., Wayson, R.L., Spengler,](#)
 1933 [J.D. and Levy, J.I.: The relationship between aviation activities and ultrafine particulate matter](#)
 1934 [concentrations near a mid-sized airport, Atmos. Environ., 50, 328-337, 2012b.](#)

1935

1936 Hsu, H. H., Adamkiewicz, G., Houseman, E. A., Zarubiak, D., Spengler, J. D. and Levy, J. I.:
 1937 Contributions of aircraft arrivals and departures to ultrafine particle counts near Los Angeles
 1938 International Airport, Sci. Tot. Environ., 444, 347-355, 2013.

1939

1940 Hsu, H. H., Adamkiewicz, G., Houseman, E. A., Spengler, J. D., Levy and J.I.: Using mobile
 1941 monitoring to characterize roadway and aircraft contributions to ultrafine particle concentrations
 1942 near a mid-sized airport, Atmos. Environ., 89, 688-695, 2014.

1943

1944 [Hu, S., Fruin, S., Kozawa, K., Mara, S., Winer, A.M. and Paulson, S.E.: Aircraft emission impacts](#)
 1945 [in a neighborhood adjacent to a general aviation airport in Southern California, Environ. Sci.](#)
 1946 [Technol., 43\(21\), 8039-8045, 2009.](#)

1947

1948 Hudda, N., Gould, T., Hartin, K., Larson, T. V. and Fruin, S. A.: Emissions from an international
 1949 airport increase particle number concentrations 4-fold at 10 km downwind, Environ. Sci. Technol.,
 1950 48, 6628-6635, 2014.

1951

1952 Hudda, N., Simon, M. C., Zamore, W., Brugge, D. And Durant, J. L.: Aviation emissions impact
 1953 ambient ultrafine particle concentrations in the greater Boston area, Environ.Sci. Technol., 50,
 1954 8514-8521, 2016.

1955

1956 [Hudda, N., Fruin, S.A.: International airport impacts to air quality: size and related properties of](#)
 1957 [large increases in ultrafine particle number concentrations, Environ. Sci. Technol., 50, 3362-3370,](#)
 1958 [2016.](#)

1959

1960 Hussein, T., Molgaard, B., Hannuniemi, H., Martikainen, J., Jarvi, L., Wegner, T., Ripamonti, G.,
 1961 Weber, S., Vesala, T. and Hameri, K.: Fingerprints of the urban particle number size distribution in
 1962 Helsinki, Finland: local vs. regional characteristics, Boreal Env. Res., 19, 1-20, 2014.

1963

1964 [ICAO \(International Civil Aviation Organization\), Annual Report of the ICAO Council: 2014. The](#)
 1965 [World of Air Transport in 2014, Appendix 1: \[https://www.icao.int/annual-report-\]\(https://www.icao.int/annual-report-2014/Documents/Appendix_1_en.pdf\)](#)
 1966 [2014/Documents/Appendix_1_en.pdf, last access: 20 June 2017.](#)

1967

1968 Janhäll S., Jonsson Å. M., Molnár P., Svensson E. A. and Hallquist M.: Size resolved traffic
 1969 emission factors of submicrometer particles, Atmos. Environ., 38, 4331-4340, 2004.

1970

1971 [Jones, A.M., Harrison, R.M., Barratt, B. and Fuller, G.: A large reduction in airborne particle](#)
 1972 [number concentrations at the time of the introduction of “sulphur free” diesel and the London Low](#)
 1973 [Emission Zone, Atmos. Environ., 50, 129-138, 2012.](#)

1974

1975 Kasumba, J., Hopke, P. K., Chalupa, D. C. and Utell, M. J.: Comparison of sources of submicron
 1976 particle number concentrations measured at two sites in Rochester, NY, Sci. Total Environ., 407,
 1977 5071-5084, 2009.

1978

1979 Kelly, F. J. and Fussell, J. C.: Size, source and chemical composition as determinants of toxicity
 1980 attributable to ambient particulate matter, Atmos. Environ., 60, 504-526, 2012.

1981 Kerminen, V. M., Pirjola, L., Boy, M., Eskola, A., Teinilä, K., Laakso, L., Asmi, A., Hienola, J.,
 1982 Lauri, A., Vainio, V. And Lehtinen, K.: Interaction between SO₂ and submicron atmospheric
 1983 particles, Atmos. Res., 54, 41-57, 2000.

1984

1985 Keuken, M. P., Moerman, M., Zandveld, P., Henzing, J. S. and Hoek, G.: Total and size-resolved
 1986 particle number and black carbon concentrations in urban areas near Schiphol airport (the
 1987 Netherlands), Atmos. Environ., 104 132-142, 2015.

1988

1989 Kim, E., Hopke, P. K., Larson, T. V. and Covert, D. S.: Analysis of ambient particle size
 1990 distributions using unmix and positive matrix factorization, Environ. Sci. Technol., 38, 202-209,
 1991 2004.

1992

1993 Kinsey, J. S., Dong, Y., Williams, D. C. and Logan, R.: Physical characterization of the fine
 1994 particle emissions from commercial aircraft engines during the aircraft particle emissions
 1995 experiment (APEX) 1 to 3, Atmos. Environ., 44, 2147-2156, 2010.

1996

1997 [Klapmeyer, M.E. and Marr, L.C.: CO₂, NO_x, and Particle Emissions from Aircraft and Support](#)
 1998 [Activities at a Regional Airport, Environ. Sci. Technol., 46\(20\), 10974-10981, 2012.](#)

1999

2000 Kley, D., Kleinmann, M., Sanderman, H. and Krupa, S.: Photochemical oxidants: state of the
 2001 science, Environ. Pollut., 100, 19-42, 1999.

2002

2003 Knibbs, L. D., Cole-Hunter, T. and Morawska, L.: A review of commuter exposure to ultrafine
2004 particles and its health effects, *Atmos. Environ.*, 45, 2611-2622, 2011.

2005

2006 [Kruskal, W.H. and Wallis, W.A., Use of ranks in one-criterion variance analysis. *J. Amer. Statist.*](#)
2007 [Assoc., 47, 583-621, 1952.](#)

2008

2009 Kulmala, M., Toivonen, A., Mäkelä, J. M. and Laaksonen, A.: Analysis of the growth of nucleation
2010 mode particles observed in Boreal forest, *Tellus B*, 50, 449-462, 1998.

2011

2012 Kulmala, M. and Kerminen, V.-M.: On the formation and growth of atmospheric nanoparticles,
2013 *Atmos. Res.*, 90, 132–150, 2008.

2014

2015 Kumar, P., Morawska, L., Birmili, W., Paasonen, P., Hu, M., Kulmala, M., Harrison, R. M.,
2016 Norford, L. and Britter, R.: Ultrafine particles in cities, *Environ.Int.*, 66, 1-10, 2014.

2017

2018 Kumar, P., Pirjola, L., Ketzel, M. and Harrison, R M.: Nanoparticle emissions from 11 non-vehicle
2019 exhaust sources—A review, *Atmos.Environ.*, 67, 252-277, 2013.

2020

2021 Lanzinger, S., Schneider, A., Breitner, S., Stafoggia, M., Erzen, I., Dostal, M., Pastorkova, A.,
2022 Bastian, S., Cyrys, J., Zscheppang, A. and Kolodnitska, T.: Associations between ultrafine and fine
2023 particles and mortality in five central European cities—Results from the UFIREG study, *Environ.*
2024 *Int.*, 88, 44-52, 2016.

2025

2026 Lee, D. S., Fahey, D. W., Forster, P. M., Newton, P. J., Wit, R. C. N., Lim, L. L., Owen, B., Sausen
2027 and R.: Aviation and global climate change in the 21st century, *Atmos. Environ.*, 43, 3520-3537,
2028 2009.

2029

2030 Liu, X., Wang, W., Liu, H., Geng, C., Zhang, W., Wang, H. and Liu, Z.: Number size distribution
2031 of particles emitted from two kinds of typical boilers in a coal-fired power plant in China, *Eng.*
2032 *Fuels*, 24, 1677-1681, 2010.

2033

2034 Liu, Z. R., Hu, B., Liu, Q., Sun, Y. and Wang, Y. S.: Source apportionment of urban fine particle
2035 number concentration during summertime in Beijing, *Atmos. Environ.*, 96, 359-369, 2014.

2036

2037 Lobo, P., Hagen, D. E. and Whitefield, P. D.: Measurement and analysis of aircraft engine PM
2038 emissions downwind of an active runway at the Oakland International Airport, *Atmos. Environ.*, 61,
2039 114-123, 2012.

2040

2041 Lobo, P., Hagen, D. E., Whitefield, P. D. and Raper, D.: PM emissions measurements of in-service
2042 commercial aircraft engines during the Delta-Atlanta Hartsfield Study, *Atmos. Environ.*, 104, 237-
2043 245, 2015.

2044

2045 Lupu, A. and Maenhaut, W.: Application and comparison of two statistical trajectory techniques
2046 for identification of source regions of atmospheric aerosol species, *Atmos. Environ.*, 36, 5607-5618,
2047 2002.

2048

2049 Masiol, M. and Harrison, R. M.: Aircraft engine exhaust emissions and other airport-related
2050 contributions to ambient air pollution: A review, *Atmos. Environ.*, 95, 409-455, 2014.

2051

2052 Masiol, M. and Harrison, R.M.: Quantification of air quality impacts of London Heathrow Airport
2053 (UK) from 2005 to 2012, *Atmos. Environ.*, 116, 308-319, 2015.

2054

2055 Masiol, M., Vu, V. T., Beddows, D. C. S. and Harrison, R.M.: Source apportionment of wide range
2056 particle size spectra and black carbon collected at the airport of Venice (Italy), *Atmos. Environ.*,
2057 139, 56-74, 2016.

2058

2059 Masiol M., Hopke P. K., Felton H. D., Frank B. P., Rattigan O. V., Wurth M. J. and LaDuke G. H.:
2060 Source apportionment of PM_{2.5} chemically speciated mass and particle number concentrations in
2061 New York City, *Atmos. Environ.*, 148, 215-229, 2017.

2062

2063 Mazaheri, M., Johnson, G. R. and Morawska, L.: Particle and gaseous emissions from commercial
2064 aircraft at each stage of the landing and takeoff cycle, *Environ. Sci. Technol.*, 43, 441-446, 2009.

2065

2066 Mazaheri, M., Bostrom, T. E., Johnson, G. R. and Morawska, L.: Composition and morphology of
2067 particle emissions from in-use aircraft during takeoff and landing, *Environ. Sci. Technol.*, 47, 5235-
2068 5242, 2013.

2069

2070 Meyer, N. K. and Ristovski, Z.: Ternary nucleation as a mechanism for the production of diesel
2071 nanoparticles: experimental analysis of the volatile and hygroscopic properties of diesel exhaust
2072 using the volatilization and humidification tandem differential mobility analyser, *Environ. Sci.*
2073 *Technol.*, 41, 7309-7314, 2007.

2074

2075 Ntziachristos, L., Ning, Z., Geller, M. D. and Sioutas, C.: Particle concentration and characteristics
2076 near a major freeway with heavy-duty diesel traffic, *Environ. Sci. Technol.*, 41, 2223-2230, 2007.

2077

2078 O'Dowd, C. D., Geever, M., Hill, M. K., Smith, M. H. and Jennings, S. G.: New particle formation:
2079 Nucleation rates and spatial scales in the clean marine coastal environment, *Geophys. Res. Lett.*, 25,
2080 1661-1664, 1998.

2081

2082 O'Dowd, C., McFiggans, G., Creasey, D. J., Pirjola, L., Hoell, C., Smith, M. H., Allan, B. J., Plane,
2083 J. M. C., Heard, D. E., Lee, J. D., Pilling, M. J. and Kulmala, M.: On the photochemical production
2084 of new particles in the coastal boundary layer. *Geophys. Res. Lett.*, 26, 1707-1710, 1999.

2085

2086 Ogulei, D., Hopke, P. K., Chalupa, D. C. and Utell, M. J.: Modeling source contributions to
2087 submicron particle number concentrations measured in Rochester, New York, *Aerosol Sci.*
2088 *Technol.*, 41, 179-201, 2007.

2089 Ostro, B., Hu, J., Goldberg, D., Reynolds, P., Hertz, A., Bernstein, L. and Kleeman, M. J.:
2090 Associations of mortality with long-term exposures to fine and ultrafine particles, species and
2091 sources: Results from the California Teachers Study Cohort, *Environ. Health Perspect.*, 123, 549-
2092 556, 2015.

2093

2094 Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemom. Intell. Lab.*,
2095 37, 23-35, 1997.

2096

2097 Paatero, P. and Tapper, U.: Positive matrix factorization: a non-negative factor model with optimal
2098 utilization of error estimates of data values, *Environmetrics*, 5, 111-126, 1994.

2099

2100 Paatero, P., Hopke, P. K., Song, X. H. and Ramadan, Z.: Understanding and controlling rotations in
2101 factor analytic models, *Chemom. Intell. Lab. Syst.*, 60, 253-264, 2002.

2102

2103 Paatero, P., Eberly, S., Brown, S. G. and Norris, G. A.: Methods for estimating uncertainty in
2104 factor analytic solutions, *Atmos. Meas. Tech.*, 7, 781-797, 2014.

2105

2106 Pant, P. and Harrison, R. M.: Estimation of the contribution of road traffic emissions to particulate
 2107 matter concentrations from field measurements: a review, *Atmos. Environ.*, 77, 78-97, 2013.
 2108

2109 Petzold, A., Ogren, J.A., Fiebig, M., Laj, P., Li, S.M., Baltensperger, U., Holzer-Popp, T., Kinne,
 2110 S., Pappalardo, G., Sugimoto, N. and Wehrli, C.: Recommendations for reporting "black carbon"
 2111 measurements. *Atmos. Chem. Phys.*, 13, 8365-8379, 2013.
 2112

2113 R Core Team: R: A language and environment for statistical computing. R Foundation for
 2114 Statistical Computing, Vienna, Austria, 2015. URL <http://www.R-project.org/>.
 2115

2116 Reff, A., Eberly, S. I. and Bhawe, P. V.: Receptor modeling of ambient particulate matter data using
 2117 positive matrix factorization: review of existing methods, *JAWMA*, 57, 146-154, 2007.
 2118

2119 [Ren, J., Liu, J., Li, F., Cao, X., Ren, S., Xu, B. and Zhu, Y.: A study of ambient fine particles at
 2120 Tianjin International Airport, China, *Sci. Total Environ.*, 556, 126-135, 2016.](#)
 2121

2122 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J.,
 2123 Amorim, A., Bianchi, F., Breitenlechner, M. And David, A.: Oxidation products of biogenic
 2124 emissions contribute to nucleation of atmospheric particles, *Science*, 344, 717-721, 2014.
 2125

2126 Rolph, G. D.: Real-time Environmental Applications and Display sYstem (READY) Website,
 2127 <http://www.ready.noaa.gov>, NOAA Air Resources Laboratory, College Park, MD, 2016.
 2128

2129 Salimi, F., Ristovski, Z., Mazaheri, M., Laiman, R., Crilley, L. R., He, C., Clifford, S. and
 2130 Morawska, L.: Assessment and application of clustering techniques to atmospheric particle number
 2131 size distribution for the purpose of source apportionment, *Atmos. Chem. Phys.*, 14, 11883-11892,
 2132 2014.
 2133

2134 Salma, I., Fűri, P., Németh, Z., Balásházy, I., Hofmann, W. and Farkas, Á.: Lung burden and
 2135 deposition distribution of inhaled atmospheric urban ultrafine particles as the first step in their
 2136 health risk assessment, *Atmos. Environ.*, 104, 39-49, 2015.
 2137

2138 Sandradewi, J., Prévôt, A. S., Szidat, S., Perron, N., Alfarra, M. R., Lanz, V. A., Weingartner, E.
 2139 and Baltensperger, U.: Using aerosol light absorption measurements for the quantitative
 2140 determination of wood burning and traffic emission contributions to particulate matter, *Environ.*
 2141 *Sci. Technol.*, 42, 3316-3323, 2008.
 2142

2143 Schneider, J., Hock, N., Weimer, S., Borrmann, S., Kirchner, U., Vogt, R. and Scheer, V.:
 2144 Nucleation particles in diesel exhaust: Composition inferred from in situ mass spectrometric
 2145 analysis, *Environ. Sci. Technol.*, 39, 6153-6161, 2005.
 2146

2147 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics - From Air Pollution to*
 2148 *Climate Change*, second ed., John Wiley & Sons, New York, 2006.
 2149

2150 Shi, J. P. and Harrison, R. M.: Investigation of ultrafine particle formation during diesel exhaust
 2151 dilution, *Environ. Sci. Technol.*, 33, 3730-3736, 1999.
 2152

2153 Shi, L., Zanutti, A., Kloog, I., Coull, B. A., Koutrakis, P., Melly, S. J. and Schwartz, J. D.: Low-
 2154 concentration PM_{2.5} and mortality: Estimating acute and chronic effects in a population-based
 2155 study, *Environ. Health Perspect.*, 124, 46-52, 2015.
 2156

2157 Shirmohammadi, F., Sowlat, M. H., Hasheminassab, S., Saffari, A., Ban-Weiss, G. and Sioutas, C.:
2158 Emission rates of particle number, mass and black carbon by the Los Angeles International Airport
2159 (LAX) and its impact on air quality in Los Angeles, *Atmos. Environ.*, 151, 82-93, 2017.
2160
2161 Sowlat M.H., Hasheminassab S. and Sioutas C.: Source apportionment of ambient particle number
2162 concentrations in central Los Angeles using positive matrix factorization (PMF), *Atmos. Chem.*
2163 *Phys.*, 16, 4849-4866, 2016.
2164
2165 Squizzato, S. and Masiol, M.: Application of meteorology-based methods to determine local and
2166 external contributions to particulate matter pollution: A case study in Venice (Italy), *Atmos.*
2167 *Environ.*, 119, 69-81, 2015.
2168
2169 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D. and Ngan, F.: NOAA's
2170 HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Amer. Meteor. Soc.*, 96,
2171 2059-2077, 2015.
2172
2173 Stevens, R. G., Pierce, J. R., Brock, C. A., Reed, M. K., Crawford, J. H., Holloway, J. S., Ryerson,
2174 T. B., Huey, L. G. and Nowak, J. B.: Nucleation and growth of sulfate aerosol in coal-fired power
2175 plant plumes: sensitivity to background aerosol and meteorology, *Atmos. Chem. Phys.*, 12, 189-
2176 206, 2012.
2177
2178 Stohl A.: Trajectory statistics—a new method to establish source–receptor relationships of air
2179 pollutants and its application to the transport of particulate sulfate in Europe, *Atmos. Environ.*, 30,
2180 579-587, 1996.
2181
2182 Stohl, A.: Computation, accuracy and applications of trajectories- review and bibliography, *Atmos.*
2183 *Environ.*, 32, 947-966, 1998.
2184
2185 Stafoggia, M., Cattani, G., Forastiere, F., di Bucchianico, A. D. M., Gaeta, A. And Ancona, C.:
2186 Particle number concentrations near the Rome-Ciampino city airport, *Atmos. Environ.*, 147, 264-
2187 273, 2016.
2188
2189 Strak, M. M., Janssen, N. A., Godri, K. J., Gosens, I., Mudway, I. S., Cassee, F. R., Lebret, E.,
2190 Kelly, F. J., Harrison, R. M., Brunekreef, B. and Steenhof, M.: Respiratory health effects of
2191 airborne particulate matter: the role of particle size, composition, and oxidative potential-the
2192 RAPTES project, *Environ. Health Perspect.*, 120, 1183-1189, 2012.
2193
2194 Thimmaiah, D., Hovorka, J. and Hopke, P. K.: Source apportionment of winter submicron Prague
2195 aerosols from combined particle number size distribution and gaseous composition data. *Aerosol*
2196 *Air Qual. Res.*, 9, 209-236, 2009.
2197
2198 [Turner, J.R., Hansen, A.D.A., and Allen G.A.: Methodologies to compensate for optical saturation](#)
2199 [and scattering in aethalometer black carbon measurements, in: Proceedings from the Symposium on](#)
2200 [Air Quality Measurement Methods and Technology, San Francisco, CA, USA, 30 April-3 May](#)
2201 [2007, Air and Waste Management Association, 2007.](#)
2202
2203 [UK Department for Transport, Heathrow Airport expansion:](#)
2204 <https://www.gov.uk/government/collections/heathrow-airport-expansion>, last access: 20 June
2205 [2017.](#)
2206
2207 USEPA: EPA Positive Matrix Factorization (PMF) 5.0 - Fundamentals and user guide.
2208 EPA/600/R-14/108, 2014

2209
2210 [Virkkula, A., Mäkelä, T., Hillamo, R., Yli-Tuomi, T., Hirsikko, A., Hämeri, K. and Koponen, I.K.:
2211 A simple procedure for correcting loading effects of aethalometer data, J. Air Waste Manage.
2212 Assoc., 57, 1214-1222, 2007.](#)
2213
2214 Vogt, R., Scheer, V., Casati, R. and Benter, T.: Onroad measurement of particle emission in the
2215 exhaust plume of a diesel passenger car, Environ. Sci. Technol., 37, 4070-4076, 2003.
2216
2217 [von Bismarck-Osten, C., Birmili, W., Ketzel, M., Massling, A., Petäjä, T. and Weber, S.:
2218 Characterization of parameters influencing the spatio-temporal variability of urban particle number
2219 size distributions in four European cities, Atmos. Environ., 77, 415-429, 2013.](#)
2220
2221 Vu, T. V., Delgado-Saborit, J. M. and Harrison, R. M.: A review of hygroscopic growth factors of
2222 submicron aerosols from different sources and its implication for calculation of lung deposition
2223 efficiency of ambient aerosols, Air Quality, Atmos. Health, 8, 429-440, 2015a.
2224
2225 Vu, T. V., Delgado-Saborit, J. M. and Harrison, R. M.: Review: Particle number size distributions
2226 from seven major sources and implications for source apportionment studies, Atmos. Environ., 122,
2227 114-132, 2015b.
2228
2229 Vu, T. V., Beddows, D. C. S., Delgado-Saborit, J. M. and Harrison, R. M.: Source Apportionment
2230 of the Lung Dose of Ambient Submicrometre Particulate Matter, Aerosol Air Quality Res., doi:
2231 10.4209/aaqr.2015.09.0553, 2016
2232
2233 Yin, J., Harrison, R. M., Chen, Q., Rutter, A. and Schauer, J. J.: Source apportionment of fine
2234 particles at urban background and rural sites in the UK atmosphere, Atmos. Environ., 44, 841-851,
2235 2010.
2236
2237 Yue, W., Stolzel, M., Cyrys, J., Pitz, M., Heinrich, J., Kreyling, W. G., Wichmann, H.-E., Peters, A.,
2238 Wang, S. and Hopke, P.K.: Source apportionment of ambient fine particle size distribution using
2239 positive matrix factorization in Erfurt, Germany, Sci. Total Environ., 398, 133-144, 2008.
2240
2241 Wang, Y., Hopke, P. K., Rattigan, O. V., Xia, X., Chalupa, D. C., Utell, M. J.: Characterization of
2242 residential wood combustion particles using the two-wavelength aethalometer, Environ.Sci.
2243 Technol., 45, 7387-7393, 2011.
2244
2245 Webb, S., Whitefield, P. D., Miake-Lye, R. C., Timko, M. T. and Thrasher, T. G.: Research needs
2246 associated with particulate emissions at airports, ACRP Report 6, Transportation Research Board,
2247 Washington, D.C., 2008.
2248
2249 Wehner, B., Uhrner, U., Von Löwis, S., Zallinger, M. and Wiedensohler, A.: Aerosol number size
2250 distributions within the exhaust plume of a diesel and a gasoline passenger car under on-road
2251 conditions and determination of emission factors, Atmos. Environ., 43, 1235-1245, 2009.
2252
2253 Wegner, T., Hussein, T., Hämeri, K., Vesala, T., Kulmala, M. and Weber, S.: Properties of aerosol
2254 signature size distributions in the urban environment as derived by cluster analysis, Atmos.
2255 Environ., 61, 350-360, 2012.
2256
2257 [Westerdahl, D., Fruin, S.A., Fine, P.L. and Sioutas, C.: The Los Angeles International Airport as a
2258 source of ultrafine particles and other pollutants to nearby communities, Atmos. Environ., 42\(13\),
2259 3143-3155, 2008.](#)
2260

2261 Wormhoudt, J., Herndon, S. C., Yelvington, P. E., Lye-Miake, R. C. and Wey, C.: Nitrogen oxide
 2262 (NO/NO₂/HONO) emissions measurements in aircraft exhausts, *J. Propul. Power*, 23, 906-911,
 2263 2007.
 2264
 2265 Zhang, K. M., Wexler, A. S., Zhu, Y. F., Hinds, W. C. and Sioutas, C.: Evolution of particle
 2266 number distribution near roadways. Part II: the 'Road-to-Ambient' process, *Atmos. Environ.*, 38,
 2267 6655-6665, 2004.
 2268
 2269 Zhang, K. M., Wexler, A. S., Niemeier, D. A., Zhu, Y. F., Hinds, W. C. and Sioutas, C.: Evolution
 2270 of particle number distribution near roadways. Part III: Traffic analysis and on-road size resolved
 2271 particulate emission factors, *Atmos. Environ.*, 39, 4155-4166, 2005.
 2272
 2273 Zhang, R., Khalizov, A., Wang, L., Hu, M. and Xu, W.: Nucleation and growth of nanoparticles in
 2274 the atmosphere, *Chem. Rev.*, 112, 1957-2011, 2011.
 2275
 2276 Zhou, L., Hopke, P. K., Stanier, C. O., Pandis, S. N., Ondov, J. M. and Pancras, J. P.: Investigation
 2277 of the relationship between chemical composition and size distribution of airborne particles by
 2278 partial least squares and positive matrix factorization, *J. Geophys. Res.-Atmos.*, 110, D07S18, 2005,
 2279 doi:10.1029/2004JD005050.
 2280
 2281 Zíková, N., Wang, Y., Yang, F., Li, X., Tian, M. and Hopke, P. K.: On the source contribution to
 2282 Beijing PM 2.5 concentrations, *Atmos. Environ.*, 134, 84-95, 2016.
 2283
 2284

TABLE LEGENDS:

Table 1. Summary of PMF results for both seasons.

Table 2. Results of Pearson's correlation analysis among extracted factor contributions and other measured variables recorded at different time resolutions. Only correlations significant at $p < 0.05$ are reported, strong correlations ($\rho > |0.6|$) are highlighted in bold.

FIGURE LEGENDS:

Figure 1. Map of LHR and sampling site (left) and map of the Greater London area (upper right). Wind roses calculated over the two sampling periods are also provided (bottom right). The location of some main potential sources is also highlighted: T1, T2, T3, T4 and T5 are the Heathrow terminals; TR= Tunnel Rd.

Figure 2. Boxplots (a) and diurnal patterns (b) of the most important measured variables (and derived variables) during the two sampling periods. All valid data are used for computing boxplot statistics: Boxplot lines= medians, crosses= arithmetic means, boxes= 25th-75th percentile ranges, whiskers= ± 1.5 *inter-quartile ranges. Diurnal variations **Figure 1** report the average levels as a filled line and the associated 95th confidence interval calculated by bootstrapping the data ($n = 200$). Outliers (data > 99.5 th percentile) were removed for computing the diurnal patterns. Hours are given in UTC. LHR traffic movements (bottom right plot) are reported as arrivals (dotted lines) and departures (solid lines). The offset between the seasons is largely due to daylight saving time (BST= UTC + 1) in the summer data.

Figure 3. Statistics of size distribution spectra for particle number (red) and volume (blue) concentrations categorised by sampling periods and time of the day (daytime= 7am-7pm and nighttime=7pm- 7am local time). For the particle number spectra, solid lines represent the median concentrations, while shaded areas report the 1st-3rd quartile intervals. (interquartile range, IQR). For the particle volume spectra, only medians are reported (dotted lines).

Figure 2. Diurnal patterns of PNC, LHR traffic, solar irradiance and eBC. Plots report the average levels as a filled line and the associated 95th confidence interval calculated by bootstrapping the data ($n = 200$). Outliers (data > 99.5 th percentile) were removed for computing the diurnal patterns. Hours are given in UTC. LHR traffic movements (bottom right plot) are reported as arrivals (dotted lines) and departures (solid lines). The offset between the seasons is largely due to daylight saving time (BST = UTC + 1) in the summer data. **Figure 4** The diurnal patterns of all the measured variables in reported in Figure S1X4.

Figure 3. Results of cluster analysis for the warm season data. Average cluster PNSD spectra (left) are reported as solid red lines along with: (i) their 10th, 25th, 75th and 90th percentile spectrum as shaded areas; (ii) the volume size distributions (dotted blue line); (iii) the hourly counts and (iv) the wind roses associated with each cluster.

2336 | **Figure 4.** Results of cluster analysis for the cold season data. Average cluster PNSD spectra
2337 | (left) are reported as solid red lines along with: (i) their 10th, 25th, 75th and 90th
2338 | percentile spectrum as shaded areas; (ii) the volume size distributions (dotted blue
2339 | line); (iii) the hourly counts and (iv) the wind roses associated with each cluster.
2340 |
2341 | **Figure 5.**
2342 |
2343 |
2344 |
2345 |
2346 |
2347 |
2348 |
2349 |
2350 |
2351 |
2352 | **Figure 5.** ~~Results of cluster analysis for the cold season data. Average cluster PNSD spectra~~
2353 | ~~(left) are reported as solid red lines along with: (i) their 10th, 25th, 75th and 90th~~
2354 | ~~percentile spectrum as shaded areas; (ii) the volume size distributions (dotted blue~~
2355 | ~~line); (iii) the hourly counts and (iv) the wind roses associated with each cluster.~~
2356 |
2357 | **Figure 6.** Results of PMF analysis for the warm season data. Factor profiles are reported on the
2358 | left as: (i) number concentration in solid red lines; (ii) their DISP ranges in shaded red
2359 | areas; (iii) volume concentrations in dotted blue lines; (iv) explained variation in
2360 | dashed grey lines. The plots on the centre report the normalised daily patterns
2361 | calculated on the hourly-averaged factor contributions along with their 95th
2362 | confidence intervals (n=200 bootstrap). The plots on the right show the polar plot
2363 | analysis (normalised average factor contributions). SA=secondary aerosol.
2364 |
2365 | **Figure 76.** Results of PMF analysis for the cold season data. Factor profiles are reported on the
2366 | left as: (i) number concentration in solid red lines; (ii) their DISP ranges in shaded red
2367 | areas; (iii) volume concentrations in dotted blue lines; (iv) explained variation in
2368 | dashed grey lines. The plots on the centre report the normalised daily patterns
2369 | calculated on the hourly-averaged factor contributions along with their 95th
2370 | confidence intervals (n=200 bootstrap). The plots on the right show the polar plot
2371 | analysis (normalised average factor contributions). SA=secondary aerosol.
2372 |
2373 | **Figure 7.** CWT maps of the secondary aerosol-related factors for both the seasons. Map scales
2374 | refer to the average factor contributions to the total variable (PNC).
2375 |
2376 |
2377 |
2378 |
2379 | **Figure 8.** ~~CWT maps of the secondary aerosol-related factors for both the seasons. Map scales~~
2380 | ~~refer to the average factor contributions to the total variable (PNC).~~
2381 |
2382 | **Figure 9.** ~~Comparison of k-means and PMF for the warm (upper plots) and cold (bottom plots)~~
2383 | ~~seasons. Boxplot statistics: lines= medians, crosses= arithmetic means, boxes= 25th-~~
2384 | ~~75th percentile ranges, whiskers= ± 1.5 *inter-quartile ranges.~~
2385 |
2386 |

2387 | **Figure 10.** Analysis of the regional nucleation episode occurring on September 7th. The selected
2388 period is from 7 September midnight to 8 September 4 pm. The plots represent (from
2389 upper to the bottom): (a) contour plots of SMPS data; (b) Concentrations of some
2390 measured species (Nucl= particles in the nucleation range 14-30 nm; Ait= particles in
2391 the Aitken Nuclei range 30-100 nm; Acc= particles in the accumulation range >100
2392 nm; mass of PM_{2.5}); (c) Source contributions from PMF for the Factors 1, 2, 3 and 4;
2393 (d) hourly counts of number of clusters. The arrows in the (b) and (c) plots show the
2394 wind direction (arrow direction) and speed (proportional to arrow length).
2395

Formatted: Not Superscript/ Subscript

2396 | **Figure 11.** Backward air mass trajectories during the nucleation event. Dots indicate 24 h step
2397 times
2398
2399

Formatted: Indent: Left: 0 cm, Hanging: 2.25 cm, Tab stops: 2.25 cm, Left

2400
2401

Table 1. Summary of PMF results for both seasons.

Factor number and interpretation	Particle Number Concentration		Particle Volume Concentration	
	No. modes ^a (peak ranges ^b)	Percent contribution (DISP range)	No. modes ^a (peak ranges ^b)	Percent contribution
Warm season (Aug-Sep 2014)				
Factor 1: Airport	1 (<20 nm)	31.6 (30.8–36.2)	2 (60–160 nm; <25 nm)	1.2
Factor 2: Fresh road traffic	1 (20–35 nm)	27.9 (24.7–30.2)	2 (22–45 nm; 140–220 nm)	1.7
Factor 3: Aged road traffic	1 (30–60 nm)	18.9 (16.6–21.1)	2 (40–100 nm; 250–450 nm)	5.6
Factor 4: Urban accumulation	1 (50–150 nm)	14.4 (13.8–18)	1 (80–250 nm)	33.2
Factor 5: Mixed SA^c	1 (110–250 nm)	5.2 (3.6–6.9)	1 (160–350 nm)	37.4
Factor 6: Inorganic SA	2 (55–120 nm; 230–400 nm)	2.1 (1.1–3.5)	2 (260–500 nm; 75–140 nm)	20.8
Cold season (Dec 2014-Jan 2015)				
Factor 1: Airport	1 (<20 nm)	33.1 (31.7–34.8)	2 (160–350 nm; 15–25 nm)	1.7
Factor 2: Fresh road traffic	1 (18–35 nm)	35.2 (33.4–36.9)	2 (22–45 nm; 150–300 nm)	3.1
Factor 3: Aged road traffic	1 (28–60 nm)	18.9 (17.9–19.7)	2 (40–150 nm; 330–450 nm)	8.7
Factor 4: Urban accumulation	1 (55–170 nm)	7.6 (7.3–8.3)	1 (100–250 nm)	32.5
Factor 5: Mixed SA	2 (130–280 nm, <17 nm)	2.3 (2.1–3.3)	1 (170–400 nm)	30.8
Factor 6: Inorganic SA	3 (17–28 nm; 55–100 nm, 250–400 nm)	2.9 (2.4–3.9)	2 (280–550 nm; 90–140 nm)	23.3

(a) Only modes above the DISP ranges are shown; (b) Range endpoints are taken at approx. half the mode height; (c) SA = secondary aerosol.

2402

Table 2. Results of Pearson's correlation analysis among extracted factor contributions and other measured variables recorded at different time resolutions. Only correlations significant at $p < 0.05$ are reported, strong correlations ($\rho > |0.6|$) are highlighted in bold.

Variables	Warm period					
	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6
	Airport	Fresh road traffic	Aged road traffic	Urban accumulation	Mixed SA	Inorganic SA
<i>Weather parameters (1 h-resolution time)</i>						
Solar irr.	0.12	-0.15	-0.24	-0.26	-0.24	-0.28
Air temp.	0.25	-0.21	-0.37	-0.1	0.1	
RH		0.1	0.32	0.22	0.26	0.33
Wind speed	0.38		-0.47	-0.64	-0.45	-0.49
<i>5 min-resolution time</i>						
Factor 1	–					
Factor 2	0.46	–				
Factor 3	0.03	0.28	–			
Factor 4	-0.17	-0.04	0.47	–		
Factor 5	-0.15	-0.06	0.21	0.56	–	
Factor 6	-0.17	-0.14	0.15	0.56	0.75	–
eBC	-0.1	-0.03	0.3233	0.6162	0.5452	0.5553
Delta-C			-0.0613	-0.0907	-0.12	-0.1306
<i>1 h-resolution time</i>						
NO			0.43	0.6	0.32	0.33
NO ₂		0.18	0.61	0.76	0.52	0.52
NO _x		0.11	0.58	0.77	0.48	0.48
O ₃	0.14	-0.19	-0.57	-0.54	-0.37	-0.43
PM _{2.5}	-0.23	-0.24	0.13	0.61	0.63	0.77
NVPM _{2.5}	-0.22	-0.22	0.17	0.62	0.61	0.75
VPM _{2.5}	-0.17	-0.24		0.42	0.54	0.65
<i>1 day-resolution time PM_{2.5}-bound species</i>						
OC				0.84	0.74	0.83
EC	-0.47	-0.54		0.75	0.51	0.67
TC	-0.45	-0.44		0.85	0.69	0.82
Chloride						
Nitrate		-0.45			0.83	0.85
Sulphate		-0.57		0.75	0.5	0.67
Oxalate		-0.47		0.59	0.89	0.93
Sodium						
Ammonium	-0.44	-0.52		0.57	0.54	0.71
Potassium		-0.47		0.46	0.5	0.66
Magnesium	0.5			-0.53		
Calcium						

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

Formatted: Font color: Auto

2408 **Table 2.** Continued.

2409

Variables	Cold period					
	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6
	Airport	Fresh road traffic	Aged road traffic	Urban accumulation	Mixed SA	Inorganic SA
<i>Weather parameters (1 h-resolution time)</i>						
Solar irr.				-0.11		
Air temp.	0.38		-0.43	-0.67	-0.5	-0.59
RH			0.23	0.38	0.46	0.46
Wind speed	0.3		-0.49	-0.67	-0.54	-0.61
<i>5 min-resolution time</i>						
Factor 1	–					
Factor 2	0.55	–				
Factor 3	0.24	0.54	–			
Factor 4	-0.11	0.08	0.53	–		
Factor 5	-0.05	0.15	0.38	0.65	–	
Factor 6	-0.09	0.08	0.39	0.7	0.81	–
eBC		0.4916	0.5452	0.7577	0.5760	0.6163
Delta-C			0.435	0.2462	0.2255	0.4952
<i>1 h-resolution time</i>						
NO	-0.14		0.51	0.81	0.62	0.63
NO ₂	0.13	0.42	0.81	0.82	0.61	0.66
NO _x		0.17	0.63	0.85	0.64	0.68
O ₃		-0.29	-0.71	-0.78	-0.65	-0.7
PM _{2.5}	-0.1	0.16	0.53	0.82	0.88	0.88
NVPM _{2.5}	-0.11	0.16	0.53	0.82	0.85	0.85
VPM _{2.5}			0.19	0.39	0.49	0.48
<i>1 day-resolution time PM_{2.5}-bound species</i>						
OC			0.79	0.79	0.76	0.8
EC			0.83	0.8	0.64	0.66
TC			0.81	0.8	0.73	0.77
Chloride				0.58	0.82	0.85
Nitrate		0.63	0.73	0.88	0.93	0.9
Sulphate					0.92	0.88
Oxalate					0.87	0.81
Sodium		-0.58	-0.74	-0.64		
Ammonium			0.63	0.78	0.99	0.97
Potassium				0.71	0.98	0.97
Magnesium						
Calcium						

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

2410

2411

2412




Figure 1. Map of LHR and sampling site (left) and map of the Greater London area (upper right). Wind roses calculated over the two sampling periods are also provided (bottom right). The location of some main potential sources is also highlighted: T1, T2, T3, T4 and T5 are the Heathrow terminals; TR= Tunnel Rd.




Figure 2. Boxplots (a) and diurnal patterns (b) of the most important measured variables (and derived variables) during the two sampling periods. All valid data are used for computing boxplot statistics: Boxplot lines= medians, crosses= arithmetic means, boxes= 25th-75th percentile ranges, whiskers= ± 1.5 inter-quartile ranges. Diurnal variations report the average levels as a filled line and the associated 95th confidence interval calculated by bootstrapping the data ($n=200$). Outliers (data ≥ 99.5 th percentile) were removed for computing the diurnal patterns. Hours are given in UTC. LHR traffic movements (bottom right plot) are reported as arrivals (dotted lines) and departures (solid lines). The offset between the seasons is largely due to daylight saving time (BST = UTC + 1) in the summer data.




Figure 3. Statistics of size distribution spectra for particle number (red) and volume (blue) concentrations categorised by sampling periods and time of the day (daytime= 7am-7pm and nighttime=7pm- 7am local time). For the particle number spectra, solid lines represent the median concentrations, while shaded areas report the 1st-3rd quartile intervals- ([interquartile range, IQR](#)). For the particle volume spectra, only medians are reported (dotted lines).





Figure 2. Diurnal patterns of PNC, LHR traffic, solar irradiance and eBC. Plots report the average levels as a filled line and the associated 95th confidence interval calculated by bootstrapping the data ($n=200$). Outliers (data >99.5th percentile) were removed for computing the diurnal patterns. Hours are given in UTC. LHR traffic movements (bottom right plot) are reported as arrivals (dotted lines) and departures (solid lines). The offset between the seasons is largely due to daylight saving

2445 time (BST = UTC + 1) in the summer data.



2446
2447 The diurnal patterns of all the measured variables in reported in Figure SIX.

2448





Figure 3. Results of cluster analysis for the warm season data. Average cluster PNSD spectra (left) are reported as solid red lines along with: (i) their 10th, 25th, 75th and 90th percentile spectrum as shaded areas; (ii) the volume size distributions (dotted blue line); (iii) the hourly counts and (iv) the wind roses associated with each cluster.



Wind roses over the clusters

Figure 54. Results of cluster analysis for the cold season data. Average cluster PNSD spectra (left) are reported as solid red lines along with: (i) their 10th, 25th, 75th and 90th percentile spectrum as shaded areas; (ii) the volume size distributions (dotted blue line); (iii) the hourly counts and (iv) the wind roses associated with each cluster.




Figure 65. Results of PMF analysis for the warm season data. Factor profiles are reported on the left as: (i) number concentration in solid red lines; (ii) their DISP ranges in shaded red areas; (iii) volume concentrations in dotted blue lines; (iv) explained variation in dashed grey lines. The plots on the centre report the normalised daily patterns calculated on the hourly-averaged factor contributions along with their 95th confidence intervals ($n=200$ bootstrap). The plots on the right show the polar plot analysis (normalised average factor contributions). SA=secondary aerosol.




Figure 76. Results of PMF analysis for the cold season data. Factor profiles are reported on the left as: (i) number concentration in solid red lines; (ii) their DISP ranges in shaded red areas; (iii) volume concentrations in dotted blue lines; (iv) explained variation in dashed grey lines. The plots on the centre report the normalised daily patterns calculated on the hourly-averaged factor contributions along with their 95th confidence intervals ($n=200$ bootstrap). The plots on the right show the polar plot analysis (normalised average factor contributions). SA=secondary aerosol.





Figure 87. CWT maps of the secondary aerosol-related factors for both the seasons. Map scales refer to the average factor contributions to the total variable (PNC).

2489



2490

2491 **Figure 98.** Comparison of k-means and PMF for the warm (upper plots) and cold (bottom plots)
2492 seasons. Boxplot statistics: lines= medians, crosses= arithmetic means, boxes= 25th-75th percentile
2493 ranges, whiskers= ± 1.5 inter-quartile ranges.




Figure 499. Analysis of the regional nucleation episode occurring on September 7th. The selected period is from 7 September midnight to 8 September 4 pm. The plots represent (from upper to the bottom): (a) contour plots of SMPS data; (b) Concentrations of some measured species (Nucl= particles in the nucleation range 14-30 nm; Ait= particles in the Aitken Nuclei range 30-100 nm; Acc= particles in the accumulation range >100 nm; mass of $PM_{2.5}$); (c) Source contributions from PMF for the Factors 1, 2, 3 and 4; (d) hourly counts of number of clusters. The arrows in the (b) and (c) plots show the wind direction (arrow direction) and speed (proportional to arrow length).

7 September 2014




Figure 11. Backward air mass trajectories during the nucleation event. Dots indicate 24 h step times