

Interactive comment on “Impacts of aerosol direct effects on tropospheric ozone through changes in atmospheric dynamics and photolysis rates” by Jia Xing et al.

Jia Xing et al.

xingjia@tsinghua.edu.cn

Received and published: 23 June 2017

We thank the referee for a thoughtful and detailed review of our manuscript. Incorporation of the reviewer’s suggestions has led to a much improved manuscript. Below we provide a point-by-point response to the reviewer’s comments and summarize the changes that have been incorporated in the revised manuscript.

[Comment]: In this paper, the authors have applied the WRF-CMAQ model to analyze the impact of aerosols on tropospheric ozone through their impacts on atmospheric dynamics and photolysis rates. Their results indicate that reducing aerosols may have negative impacts on ozone which need to be considered for air quality management

Printer-friendly version

Discussion paper



in China. The topic is of general interest given the focus on reducing PM_{2.5} pollution. The simulations have been designed appropriately to address the goals of the study.

[Response]: We thank the reviewer for the overall positive assessment of the manuscript and recognition of the implications of the results of the analysis presented.

[Comment]: However, the authors have not considered the prominent way aerosols impact tropospheric ozone formation - via heterogeneous reactions - which leads me to question the conclusions of this study. Several studies have highlighted the role of aerosols in modulating ozone via heterogeneous reactions (eg., Liao and Seinfeld, 2005; Ti et al., 2005; Pozzoli et al., 2008; Xu et al., 2012; Lou et al., 2014), which have largely been ignored in this study. The authors need to provide a strong justification for ignoring the impact of aerosols on ozone via heterogeneous reactions, before I can recommend this paper for publication.

[Response]: We agree with the reviewer that the heterogeneous reactions associated with aerosols have substantial impacts on ozone, including hydrolysis of N₂O₅, irreversible absorption of NO₂ and NO₃ as well as the uptake of HO₂, and are well documented in the literature (Tang et al., 2004; Tie et al., 2005; Liao and Seinfeld, 2005; Pozzoli et al., 2008; Li et al., 2011; Xu et al., 2012; Lou et al., 2014). Our model contains comprehensive treatment of heterogeneous hydrolysis of N₂O₅ (Davis et al., 2008; Sarwar et al., 2012; Sarwar et al., 2014). While our model accounts for such a heterogeneous reaction, we have not quantified its impacts on ozone in this study.

However, in this study, we focused our analysis on another important aspect of aerosol influence (ADE, the aerosol direct effects), i.e., scattering and absorption of incoming solar radiation and how the subsequent effects of the associated cooling suppresses atmospheric ventilation. The assessment of impacts of aerosol direct effects (ADE) is an important aspect of designing emission reduction strategies that seek co-benefits associated with reductions in both particulate matter and ozone. In this study, we examine the ADE impacts which were not well quantified in previous studies. It may

[Printer-friendly version](#)[Discussion paper](#)

be noted that all model calculations analyzed in this study included the heterogeneous N₂O₅ hydrolysis pathway. We agree with the reviewer that all influences of aerosols on ozone need to be addressed before definite policy-relevant conclusions regarding their overall impact can be reached. China plans to implement stringent control actions aimed at lowering the ambient concentrations of PM_{2.5} in the next two decades. It will be necessary to quantify all the possible influences resulting from this expected reduction of aerosols, including changes in heterogeneous reactions associated with aerosols, as well as the changes expected in ADE discussed in this manuscript. In addition, secondary aerosol and ozone comes from both NO_x and VOCs emissions. Reducing aerosols by controlling gaseous precursors will also have substantial impacts on ozone (Xing et al., 2017), which needs further evaluation.

To address the reviewer's concern, we have clarified the scope of our analysis in the revised manuscript as below:

(Page 2 Line 21-25) "Many studies suggest that aerosols may have substantial impacts on ozone through heterogeneous reactions including hydrolysis of N₂O₅, irreversible absorption of NO₂ and NO₃, as well as the uptake of HO₂ (Tang et al., 2004; Tie et al., 2005; Li et al., 2011; Lou et al., 2014). While our model contains comprehensive treatment of the heterogeneous hydrolysis of N₂O₅ (Davis et al., 2008; Sarwar et al., 2012; Sarwar et al., 2014), we have not quantified its impacts on ozone in this study. However, ADE impacts on ozone have not been well evaluated previously."

(Page 10 Line 15-19) "Reducing aerosols will have substantial impacts on ozone. Quantification of the aerosol influence on ozone is important to understand co-benefits associated with reductions in both particulate matter and ozone. This study focused on the evaluation of ADE impacts which were not well quantified previously. However, the heterogeneous reactions associated with aerosols, as well as the impacts of emission controls of gaseous precursors on both aerosols and ozone also need to be studied in order to fully understand the influence from reducing aerosols on ambient ozone."

[Printer-friendly version](#)[Discussion paper](#)

Reference

Li J., Wang Z., Wang X., Yamaji K., et al. Impacts of aerosols on summertime tropospheric photolysis frequencies and photochemistry over Central Eastern China. *Atmospheric Environment*, 2011, 45: 1817-1829.

Liao, H., Seinfeld, J.H., Global impacts of gas-phase chemistry–aerosol interactions on direct radiative forcing by anthropogenic aerosols and ozone, *J. Geophys. Res.*, 2005, 110, D18208

Tang Y. H., Carmichael G. R., Kurata G., Uno I., et al. Impacts of dust on regional tropospheric chemistry during the ACE-Asia experiment: A model study with observations. *Journal of Geophysical Research-Atmospheres*, 2004, 109

Tie X. X., Madronich S., Walters S., et al. Assessment of the global impact of aerosols on tropospheric oxidants. *Journal of Geophysical Research-Atmospheres* 2005, 110

Lou S. J., Liao H., Zhu B. Impacts of aerosols on surface-layer ozone concentrations in China through heterogeneous reactions and changes in photolysis rates. *Atmospheric Environment*, 2014, 85: 123-138.

Pozzoli, L., Bey, I., Rast, S., Schultz, M.G., Stier, P., Feichter, J., Trace gas and aerosol interactions in the fully coupled model of aerosol-chemistry-climate ECHAM5-HAMMOZ: 1. Model description and insights from the spring 2001 TRACE-P experiment, *J. Geophys. Res.*, 2008, 113, D07308

Xu, J., Zhang, Y.H., Zheng, S.Q., He, Y.J., Aerosol effects on ozone concentrations in Beijing: a model sensitivity study, *J. Environ. Sci.*, 2012, 24 (4), 645–656

Davis, J. M., Bhave, P. V., and Foley, K. M.: Parameterization of N₂O₅ reaction probabilities on the surface of particles containing ammonium, sulfate, and nitrate, *Atmos. Chem. Phys.*, 8, 5295–5311, doi:10.5194/acp-8-5295-2008, 2008.

Sarwar, G., Simon, H., Bhave, P., and Yarwood, G.: Examining the impact of heteroge-

neous nitryl chloride production on air quality across the United States, *Atmospheric Chemistry & Physics*, 12, 1-19, 2012.

Sarwar, G., Simon, H., Xing, J., Mathur, R.: Importance of tropospheric ClNO₂ chemistry across the Northern Hemisphere, *Geophysical Research Letters*, 41, 4050-4058, 2014.

Xing, J., Wang, S. X., Jang, C., et al. Overview of ABaCAS: an air pollution control cost-benefit and attainment assessment system and its applications in China, EM of Air & Waste Management Association, 2017 April.

[Comment]: P2L10: Please provide a reference for “ADE modulate the temperature, atmospheric ventilation, cloud and rainfall”.

[Response]: We have provided the references for this sentence as below:

(Page 2 Line 10-12) “ADE modulate the temperature (e.g., Hansen et al., 1997; Mitchell et al., 1995), atmospheric ventilation (e.g., Jacobson et al., 2007; Mathur et al., 2010;), cloud and rainfall (e.g., Albrecht, 1989; Liou and Ou, 1989; Twomey, 1977)”

Mitchell JFB, Davis RA, Ingram WJ, Senior CA. On Surface Temperature, Greenhouse Gases, and Aerosols: Models and Observations. *Journal of Climate* 1995; 8: 2364-2386.

Hansen J, Sato M, Ruedy R. Radiative forcing and climate response. *J Geophys Res-Atmos* 1997; 102: 6831-6864

Albrecht BA. Aerosols, Cloud Microphysics, and Fractional Cloudiness. *Science* 1989; 245: 1227-1230.

Liou K, Ou S. The role of cloud microphysical processes in climate - an assessment from a one-dimensional perspective. *J Geophys Res-Atmos* 1989; 94: 8599-8607

Twomey S. Influence of pollution on shortwave albedo of clouds. *Journal of the Atmospheric Sciences* 1977; 34: 1149-1154

Mathur, R. et al. The WRF-CMAQ Integrated On-Line Modeling System: Development, Testing, and Initial Applications. in Air Pollution Modeling and its Applications XX, D. G. Steyn, S. T. Rao, Eds. Springer, Netherlands, Netherlands, 2010, 155-159.

Jacobson, M. Z.; Kaufman, Y. J.; Rudich, Y. Examining feedbacks of aerosols to urban climate with a model that treats 3-D clouds with aerosol inclusions, J. Geophys. Res. 2007, 112, D24205

[Comment]: P3 Section 2.1: What meteorological fields are used to drive WRF-CMAQ?

[Response]: The meteorological inputs for WRF simulations were derived from the NCEP FNL (Final) Operational Global Analysis data which has 1 degree spatial and 6-hour temporal resolution. NCEP ADP Operational Global Surface Observations were used for surface reanalysis which is used for indirect soil moisture and temperature nudging in the Pleim-Xiu Land Surface Model (PX LSM). The strength of nudging coefficients for four dimensional data assimilation and indirect soil temperature nudging employed in WRF have been tested and chosen to improve model performance for meteorological variables without dampening the effects of radiative feedbacks. The nudging coefficient for both u/v-wind and potential temperature is set to 0.00005 sec⁻¹, while 0.00001 sec⁻¹ is used for nudging of water vapor mixing ratio.

We have clarified in the revised manuscript as below:

(Page 3 Line 8-13) “The meteorological inputs for WRF simulations were derived from the NCEP FNL (Final) Operational Global Analysis data which has 1 degree spatial and 6-hour temporal resolution. NCEP ADP Operational Global Surface Observations were used for surface reanalysis and four dimensional data assimilation. We have tested and chosen proper strength of nudging coefficients, i.e., 0.00005 sec⁻¹ is used for nudging of both u/v-wind and potential temperature, 0.00001 sec⁻¹ is used for nudging of water vapor mixing ratio, to improve model performance without dampening the effects of radiative feedbacks (Hogrefe et al., 2015; Xing et al., 2015).”

[Printer-friendly version](#)[Discussion paper](#)

[Comment]: P3L5,6: Please define acronyms (e.g., AERO6, BHCOAT) before using them.

[Response]: We have clarified the acronyms in the revised manuscript as below

(Page 3 Line 13-18) “In the model version used here, concentrations of gaseous species and primary and secondary aerosols are simulated by using Carbon Bond 05 (CB05) gas-phase chemistry (Sarwar et al., 2008) and the sixth-generation CMAQ modal aerosol model (AERO6) (Appel et al., 2013). The aerosol optical properties were estimated by the coated-sphere module (i.e., BHCOAT, Bohren and Huffman, 1983) based on simulated aerosol composition and size distribution (Gan et al., 2015).”

[Comment]: P4L20: Please clarify if this is for observations. The observations are hardly visible in Figure 2.

[Response]: Yes, this is for observations. We have replotted Figure 2 in the revised manuscript to make the comparison more evident, as below

(Page 4 Line 28 - Page 5 Line 3) “In January, higher DM1O3 concentrations are seen in PRD where solar radiation is stronger than in the north. The model generally captured the spatial pattern with highest DM1O3 in PRD over the simulated domain. Simulated DM1O3 in YRD, SCH and HUZ are higher than observations. Such overestimation might be associated with the relative coarse spatial resolution in the model. NO titration effects in urban areas were not well represented in the model.”

[Comment]: P4L24: Please provide a reference for “In January, O3 production in north China is VOC-limited regime”

[Response]: Following reviewer’s suggestion, a reference has been added as below:

(Page 5 Line 6-8) “In January, O3 production in north China is occurring in a VOC-limited regime (e.g., Liu et al., 2010), thus increases in NOx at the surface stemming from the stabilized atmosphere by ADE inhibit O3 formation due to enhanced titration by NO.”

[Printer-friendly version](#)[Discussion paper](#)

Liu, X.H., Zhang, Y., Xing, J., Zhang, Q., Wang, K., Streets, D.G., Jang, C., Wang, W.X. and Hao, J.M., 2010. Understanding of regional air pollution over China using CMAQ, part II. Process analysis and sensitivity of ozone and particulate matter to precursor emissions. *Atmospheric Environment*, 44(30), pp.3719-3727.

[Comment]: P4: It would also be helpful to see maps of PM_{2.5} to assess if the reductions in O₃ due to aerosol feedbacks are co-located with PM concentrations.

[Response]: Following the reviewer's suggestion, we made the plot of PM_{2.5} concentrations as Figure R1. It clearly shows that the ADE effects on O₃ are co-located with PM concentrations. Besides, we calculated the O₃ responses to ADE under low- and high-PM_{2.5} levels, summarized in Table 2. Mostly, the O₃ responses to ADE are larger under high PM_{2.5} levels, indicating the positive correlations between O₃ responses and PM levels.

We have added corresponding discussions to the revised manuscript, as below:

(Page 6 Line 3-5) "Comparing the O₃ responses to ADE (see Δ -ADE in Table 2) under low- and high- PM_{2.5} levels, reveals that O₃ responses to ADE are larger under high PM_{2.5} levels, indicating the positive correlations between O₃ responses and PM_{2.5} levels."

[Comment]: P4-5: How significant are the changes in O₃ in response to Dynamics and Photolysis

[Response]: Our results suggest that ADE reduced surface daily maxima 1h O₃ (DM1O₃) in China by up to 39 $\mu\text{g m}^{-3}$ through the combination of Δ Dynamics and Δ Photolysis in January, but enhanced surface DM1O₃ by up to 4 $\mu\text{g m}^{-3}$ in July.

The enhancement in peak O₃ in summer found in this study is different from the traditional expectation of a negative O₃ response to ADE when solely considering the reduction in solar radiance.

[Printer-friendly version](#)[Discussion paper](#)

Please also note the supplement to this comment:
<http://www.atmos-chem-phys-discuss.net/acp-2017-198/acp-2017-198-AC2-supplement.pdf>

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2017-198>, 2017.

ACPD

Interactive
comment

Printer-friendly version

Discussion paper




Figure R1 spatial distribution of PM_{2.5} concentration and O₃ response to ADE

Fig. 1.

Printer-friendly version

Discussion paper

