



Effects of atmospheric transport and trade on air pollution mortality in China

Hongyan Zhao¹, Xin Li¹, Qiang Zhang¹, Xujia Jiang^{1,2}, Jintai Lin³, Glen G. Peters⁴, Meng Li¹, Guannan Geng¹, Bo Zheng², Hong Huo⁵, Lin Zhang³, Steven J. Davis⁶ and Kebin He^{1,2}

¹Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science, Tsinghua University, Beijing, China;
²State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing, China;
³Laboratory for Climate and Ocean-Atmosphere Studies, Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, China;
⁴Center for International Climate and Environmental Research—Oslo, N-0318 Oslo, Norway;
⁵Institute of Energy, Environment and Economy, Tsinghua University, Beijing, China;
⁶Department of Earth System Science, University of California, Irvine, CA 92697, USA

15 *Correspondence to:* Qiang Zhang (qiangzhang@tsinghua.edu.cn) and Steven J. Davis (sjdavis@uci.edu)

Abstract. Air quality is a major environmental concern in China, where premature deaths due to air pollution exceed 1 million people per year in recent years. Here, using a novel coupling of economic, physical and epidemiological models, we estimate the premature mortality in seven regions of China related to emissions of $PM_{2.5}$ and precursor gases in 2010 and show for the first time how the distribution of these deaths in China is determined by a combination of economic activities and physical

- 20 transport of pollution in the atmosphere. We find that 33% of China's air pollution deaths in 2010 were caused by pollutants emitted in a different region of the country and transported in the atmosphere, especially from north to south and from east to west. Similarly, 38% of deaths were related to goods and services consumed in a different region from where they were produced. For example, 36% of the deaths number associated to emissions in Central region of China were caused by consumption in other regions. As a combined result from atmospheric transport and trade, 56% of pollution deaths in China
- 25 were related to consumption in a different region. Among these, 14% of China's pollution deaths were caused by international export. Our results indicate that multilateral and multi-stage cooperation under a regional sustainable development framework is in urgent need to mitigate air pollution and health impacts, and efforts to reduce the health impacts of air pollution in China should be prioritized according to the source and location of emissions, the type and economic value of the emitting activities, and the related patterns of consumption.

30 1 Introduction

Atmospheric pollution is a major environmental problem in China, with substantial adverse health effects (Yang et al.2013; Apte et al., 2015). Between 2006 and 2012, approximately 1.1 billion people (82% of the nation's population) live in areas

⁵





5

where the yearly average mass concentrations of fine particulate matter (PM_{2.5}) exceeds 35 µg m⁻³ (Geng et al., 2015)—above the interim target-1 for annual average exposure set by the World Health Organization (WHO, 2005). In turn, this magnitude of exposure has had large impacts on public health and economic output. In 2010, PM_{2.5} pollution alone was linked to 1.2 million premature deaths in China (Yang et al. 2013), or ~35% of all such deaths worldwide (Apte et al., 2015; Brauer et al., 2016), with associated economic losses equivalent to more than 10% of China's GDP (Hamilton, 2015). The distribution of air pollution and attendant impacts vary across Chinese provinces due to differences in physical geography, meteorology, population density, level of economic development, production structure, and available technologies (Geng et al., 2015; Jiang et al., 2015; Ma et al., 2014). For example, annual average PM_{2.5} concentrations in northern China are roughly 1.5 times greater than the national average and two times greater than concentrations in southern China (Geng et al., 2015). In light of these

10 differences, the central and local governments have established various goals, strategies and measures for reducing air pollution, with varying degrees of success (Lin et al., 2010).

Effective and efficient control of air pollution relies upon an understanding of the pollution sources and their relative environmental impacts. This has led to an increasing number of studies aimed at attributing pollution to sources at high spatial, temporal, and sectoral resolutions (e.g., Chambliss et al., 2014; Zhang et al., 2015; Turner et al., 2015; Li et al., 2015). An

- 15 important finding of these studies is that regional air quality is in many cases strongly influenced by pollution produced in other regions and transported in the atmosphere across regional boundaries. For example, a recent study found that during the month of January 2013-2015, roughly half of the PM_{2.5} present in Beijing and Tianjin (47% and 55%, respectively) was due to emissions produced in other regions (Li et al., 2015). Other recent work has highlighted the effect of interprovincial trade on the spatial distribution of emissions, with emissions increasingly outsourced to less-developed interior provinces (Feng et
- 20 al., 2013; Zhao et al., 2015). Here, we build on these studies to assess for the first time the health impacts of trans-boundary PM_{2.5} pollution and trade within China. Our results reveal with greater detail than previously the health impacts of specific economic activities (e.g., the production of raw materials and intermediate goods, production of final goods, and consumption of final goods) by region in China. This information may be used by policymakers in the design and evaluation of control strategies that account for cross-regional pollution.
- 25 Our analysis entails a novel coupling of physical, economic, and epidemiological models that use the latest available data, from 2010. Together, these models allow us to estimate premature deaths in China due to local and trans-boundary anthropogenic PM_{2.5} pollution associated with three different economic activities (production of raw materials/intermediate goods, production of final goods, and consumption of final goods) for each of seven regions (North, Yangtze River Delta, Southeast, Central, Northwest, Southwest, and Northeast China; see Table A1 for region definitions).

30 2 Materials and Methods

In this study, four state-of-the-art models were integrated to analyse the drivers of $PM_{2.5}$ -related deaths across seven regions in China. Figure 1 depicts the integrated assessment framework in 4 steps. Below we describe in details for each step.





5

2.1 Estimation of PM_{2.5} related premature deaths

Satellite-based ground-level $PM_{2.5}$ mass concentrations at a 0.5 °×0.667 γ esolution used in this study were derived from our previous research (Geng et al., 2015). It was estimated by using the aerosol optical depth (AOD) derived from satellite instruments and conversion factors between AOD and $PM_{2.5}$, simulated by a chemical transport model. The satellite-based AOD were generated by combining results from MODIS and MISR instruments onboard the Terra satellite, after being filtered

- by using ground-based AOD measurements. The conversion factors between AOD and $PM_{2.5}$ were calculated by the nested GEOS–Chem model over China at a resolution of $0.5^{\circ} \times 0.667^{\circ}$. And for this simulation, anthropogenic emissions over China were taken from the Multi-resolution Emission Inventory of China (MEIC: http://www.meicmodel.org/), a updated version of the technology-based, bottom-up pollution inventory developed by Zhang et al. (2009) , Lei et al. (2011) and Liu et al (2015).
- 10 Note that, in this study the simulated proportion of mineral dust in surface PM2.5 was subtracted to exclude the impact of natural mineral dusts on premature deaths, and we assumed that the contribution of dust to PM2.5 concentrations was proportional to their previously-estimated air pollution disease burden (Chafe et al., 2015; Bhalla et al., 2014).

The IER model developed by Burnett et al. (2015) describes the concentration-response relationship between long term exposure to PM2.5 (annual mean values in this study) and premature deaths for various leading causes. It is fitted by incorporating information from cohort studies of ambient air pollution, second hand tobacco smoke, household solid, cooking fuel, and active smoking (Burnett et al., 2015). Its relative risk (*RR*) was calculated as:

$$RR_{l}(C) = \begin{cases} 1 + \alpha_{l} \left(1 - e^{-\gamma_{l}(C - C_{0})^{\delta_{l}}} \right), & \text{if } C > C_{0} \\ 1 & \text{, else} \end{cases}$$
(1)

where *C* is the annual mean PM_{2.5} concentrations in 2010; C_0 is the counterfactual concentration; *l* represents a given health effect; and α_l , γ_1 , and δ_l are parameters used to describe the shape of the concentration-response curve.

20 The RR then was can be converted to the attributable fraction (AF):

$$AF = \frac{RR-1}{RR}$$
(2)

The health outcomes or mortality attributable to PM_{2.5} was then estimated:

$$\mathbf{M} = \mathbf{AF} \times \mathbf{B} \times \mathbf{P} \tag{3}$$

where B is the death incidence of a given health effect derived from the national average data in GBD2013 (Forouzanfar et al.,
2015); P is the size of the exposed population obtained from the LandScan global population database (Bright et al., 2011).

Following the previous studies (Jiang et al., 2015; Lee et al., 2015), in this work we mainly focus on four leading causes of the PM_{2.5} related premature deaths: ischemic heart disease (IHD), stroke, chronic obstructive pulmonary disease (COPD) and lung cancer (LC). And according to the Global Burden of Disease (GBD) projects (Forouzanfar et al., 2015), we assume that these PM_{2.5} related health impacts are source and composition independent.





2.2 Contribution of individual source emissions to regional premature deaths

Following Lee et.al (2015), the GEOS-Chem adjoint (version 3.5, driven by MEIC inventory) combined with the IER model were applied over East Asia (11 S-55 N, 70 E-150 E) at a resolution of 0.5 ×0.667 ° to calculate the contributions of location- and species-specific emissions to PM_{2.5}-related premature deaths in individual regions (see region definitions in Table

5 A1).

Firstly, we defined eq.3 as the adjoint cost function (or concentration-dependent function) of a given region (e.g. region *r*); and the total value of M^r is with respect to the satellite-based PM_{2.5} in region *r* obtained from Geng et.al (2015). We then used the adjoint model to calculate partial derivatives of this cost function with respect to anthropogenic emissions of individual species $(\frac{\partial M^r}{\partial E_{i,j,k}})$, which we referred to as the sensitivity of region *r*'s premature deaths (M^r) to gridded emissions ($E_{i,j,k}$; *i*, *j* and

k are indices for longitude, latitude and species) in the simulation domain. A single adjoint simulation provided sensitivities of M^r with respect to emissions at all species, locations and times (Lee et al., 2015; Pappin and Hakami, 2013;Turner et al., 2015). After computing the model sensitivities, we multiplied the emission sensitivity by the amount of emissions to obtain a semi-normalized sensitivity (SS) (Henze et al., 2007;Henze et al., 2009;Turner et al., 2015; Dedoussi and Barrett, 2014), which means the contribution of species- and location-specific emissions to the premature deaths (Turner et al., 2015; Dedoussi and Barrett, 2014):

$$SS_{i,j,k}^r = \frac{\partial M^r}{\partial E_{i,j,k}} \times E_{i,j,k}$$
(4)

Then, a normalized *SS* (hereafter P), which represents the percentage contribution of source- specific emissions to premature deaths was calculated as:

$$P_{i,j,k}^{r} = \frac{SS_{i,j,k}^{r}}{\sum_{i}\sum_{j}\sum_{k}SS_{i,j,k}^{r}} \times 100\%$$
(5)

20 The normalization process minimizes the effects of nonlinear relation between emissions and pollutant concentrations and between concentrations and mortality. A similar approach was taken by Li et.al. (2016) for attributing ozone radiative forcing to individual countries.

For this work, we calculated responses to absolute changes in NH_3 , SO_2 , NO_x , BC, OC and anthropogenic $PM_{2.5}$ dust (Zhang et al., 2015). And a total of seven groups of GEOS-Chem adjoint model simulations were conducted; one group for each

receptor region. In order to reduce the computation costs, four months (January, April, July and October of 2010) of simulations

25

2.3 Regional pollutant emissions attributed to regions producing final goods and regions consuming the final goods.

were conducted for each group. Results for these four months are averaged to represent the annual mean SS in 2010.

The production of a specific product or service represents only one stage in a supply chains because such production requires material and energy inputs and may in turn supply other production processes (i.e., the products are intermediate) or final sales





10

(i.e., the products are finished goods ready for final consumption) (Davis et al., 2011). Using the 30-province, 30-sector MRIO table of China compiled by Liu et.al (2014), we attribute the emissions released in a region (i.e., the producer) to both final produces in supply chains (who produced the finished products using intermediate inputs made locally or imported from other regions, here we call this regions as "assembler") and final consumers (who ultimately consume the finished products).

5 The MRIO analysis starts with the monetary flows between sectors and regions:

$$\begin{pmatrix} \mathbf{x}^{1} \\ \mathbf{x}^{2} \\ \mathbf{x}^{3} \\ \vdots \\ \mathbf{x}^{m} \end{pmatrix} = \begin{pmatrix} \mathbf{A}^{1,1} & \mathbf{A}^{1,2} & \mathbf{A}^{1,3} & \cdots & \mathbf{A}^{1,m} \\ \mathbf{A}^{2,1} & \mathbf{A}^{2,2} & \mathbf{A}^{2,3} & \cdots & \mathbf{A}^{2,m} \\ \mathbf{A}^{3,1} & \mathbf{A}^{3,2} & \mathbf{A}^{3,3} & \cdots & \mathbf{A}^{3,m} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{A}^{m,1} & \mathbf{A}^{m,2} & \mathbf{A}^{m,3} & \cdots & \mathbf{A}^{m,m} \end{pmatrix} \begin{pmatrix} \mathbf{x}^{1} \\ \mathbf{x}^{2} \\ \mathbf{x}^{3} \\ \vdots \\ \mathbf{x}^{m} \end{pmatrix} + \begin{pmatrix} \sum_{r} \mathbf{y}^{1,s} \\ \sum_{r} \mathbf{y}^{2,s} \\ \sum_{r} \mathbf{y}^{3,s} \\ \vdots \\ \sum_{r} \mathbf{y}^{m,s} \end{pmatrix}$$
(6)

where x^r is a vector of the total economic output of each sector in province *r*; $y^{r,s}$ is a vector of the finished products by each sector produced in region *r* and consumed in region *s* A^{*r*,*s*} is a normalized matrix of intermediate coefficients in which the columns reflect the input from the sectors in region *r* required to produce one unit of output from each sector in region *s*; and *m* is the total province number (here *m*=30). Solving for total output, eq can be written as:

$$\mathbf{X} = (\mathbf{I} - \mathbf{A})^{-1} \mathbf{y} \tag{7}$$

where I is identity matrix, A is the block matrix in eq.6, and $(I-A)^{-1}$ is the Leontief inverse matrix.

Under this framework, pollutant emissions embodied in the trade flow can be calculated as:

$$\mathbf{E} = \hat{f} (\mathbf{I} - \mathbf{A})^{-1} \mathbf{y}$$
(8)

15 where \hat{f} is the diagonalization of the vector of region-specific pollutant emissions for unit output of each sector.

Region- and sector-specific emissions attributed to assembler region s can be calculated as follows:

$$\mathbf{e}_{asse}^{s} = \hat{f} (\mathbf{I} - \mathbf{A})^{-1} \begin{pmatrix} 0 \\ \vdots \\ \sum_{r} \mathbf{y}^{s,r} \\ \vdots \\ 0 \end{pmatrix}$$
(9)

where $\mathbf{e}_{asse}^{s} = (\mathbf{e}_{asse}^{1,s} \mathbf{e}_{asse}^{2,s} \mathbf{e}_{asse}^{3,s} \dots \mathbf{e}_{asse}^{m,s})$; and $\mathbf{e}_{asse}^{r,s}$ is a sector-specific vector for emissions occurred in region *r* caused by producing intermediate products to be assembled (as finished products) in region *s*.

20 Region- and sector-specific emissions attributed to consumer region *s* can be calculated as:

1 c



(10)



$$\mathbf{e}_{cons}^{s} = \hat{f}(\mathbf{I} - \mathbf{A})^{-1} \begin{pmatrix} \mathbf{y}^{1,s} \\ \mathbf{y}^{2,s} \\ \mathbf{y}^{3,s} \\ \vdots \\ \mathbf{y}^{m,s} \end{pmatrix}$$

where $\mathbf{e}_{cons}^{s} = (\mathbf{e}_{cons}^{1,s} \ \mathbf{e}_{cons}^{2,s} \ \mathbf{e}_{cons}^{3,s} \dots \ \mathbf{e}_{cons}^{m,s})^{2}$; and $\mathbf{e}_{cons}^{r,s}$ is a sector-specific vector for emissions occurred in region *r* caused by production of intermediate or final products to be consumed in region *s*.

In this section, sector specific emissions to produce f in eq.8 to eq.10 were derived from mapping process between MEIC 5 model and sectors defined in the MRIO model for each provinces, which can be found in our previous studies (Huo et al., 2014; Zhao et al., 2015). Within each region, emissions attributed to each activator (assembler or consumer) can be allocated to individual locations (grid cells) based on the sector-spatial distribution in MEIC and the attributed ratios:

$$R_{k,asse}^{r,s} = \mathbf{e}_{k,asse}^{r,s} / \mathbf{e}_{k}^{r}$$
(11)

$$R_{k,cons}^{r,s} = \mathbf{e}_{k,cons}^{r,s} / \mathbf{e}_{k}^{r}$$
(12)

10 where \mathbf{e}_k^r is sector-specific emissions vector (species k) produced in region r; $R_{k,asse}^{r,s}$ and $R_{k,cons}^{r,s}$ are sector-specific ratios of emissions occurred in region r but allocated to region s from assembler and consumer perspectives, respectively. As part of our calculation, we aggregated the interregional emission impact of 30 provinces into 7 regions, as defined in Table A1.

2.4 Premature deaths attributed to regions producing final goods and regions consuming the final goods

Results from above three steps were integrated to attribute regional- and source-specific PM_{2.5} deaths to specific economic activities (i.e. the production of final goods by the "assembler", and the ultimate consumption of those goods) in specific regions along supply chains as:

$$M_{asse}^{s} = \sum_{r} M^{r} \sum_{t} \sum_{k} (P_{(i,j)\in t,k}^{r} \times R_{(i,j)\in t,k,asse}^{t,s})$$

$$M_{cons}^{s} = \sum_{r} M^{r} \sum_{t} \sum_{k} (P_{(i,j)\in t,k}^{r} \times R_{(i,j)\in t,k,cons}^{t,s})$$

$$(13)$$

$$(14)$$

where M_{asse}^s and M_{cons}^s mean premature deaths attributed to region *s* from assembler and consumer perspectives, respectively; 20 $R_{(i,j)\in t,k,asse}^{t,s}$ and $R_{(i,j)\in t,k,cons}^{t,s}$ are sector average ratios of emission occurred in grid (i,j) relocated to region *s* from assembler and consumer perspectives, respectively.

3 Results

3.1 National and regional mortality attributed to anthropogenic PM_{2.5}

In 2010, China's population-weighted PM_{2.5} concentrations caused by anthropogenic emissions reached 53 μg m⁻³, leading to 1.13 (95% CI: 0.7-1.35) million premature deaths, which account for about 35% of the global total mortality from ambient





PM_{2.5} (Apte et al., 2015). Adding another 0.12 (95% CI: 0.08-0.14) million premature deaths from windblown natural dusts, our estimate of premature deaths is within 2% of the result of GBD 2010 for China (1.27 million deaths; Brauer et al., 2015).

Table 1 and Figure 2A show in detail about regional anthropogenic $PM_{2.5}$ concentrations and related mortality. As figures and table shown, atmosphere pollution and related health impact vary substantially across the seven Chinese regions. Dominated

- 5 by heavy industries, North region bear the most severe pollution, and its population-weighted mean PM_{2.5} concentrations reached 82 μ g m⁻³, followed by the Central (67 μ g m⁻³), Southwest (52 μ g m⁻³) and Yangtze River Delta (50 μ g m⁻³). However, considering the total population exposed to pollution, the Central region has the highest mortality [342,662 (208,950-414,669) premature deaths] and a high mortality ratio [102 (62-123) deaths per 10⁵ people], followed by the Southwest [211,016 (131,518-253,869) premature deaths; 83 (52-100) deaths per 10⁵ people), and the North [203,278 (126,611-239,255) premature
- 10 deaths; 106 (66-125) deaths per 10^5 people].

Regional atmospheric pollution and related health impacts can be attributed to emissions from both local and other regions as a result of atmospheric transport (figure 2B). Further, emissions in a given region can also be attributed to regions who consuming the emission related products due to trade, thus pollution deaths can finally be attributed to the consuming regions (figure C). In section 3.2 and 3.3 we show in detail about regional pollution deaths and its affecting regions from these two perspectives, respectively.

15 perspectives, respectively.

3.2 Effects of atmospheric transport of air pollution on regional mortality

As figure 2B shown, in 2010 33% of all premature deaths due to air pollution were caused by trans-boundary pollution, and ratios for specific regions vary from 30% in Northeast to 40% in Northwest. Among these, less than 1% were caused by pollution transported from region out of the seven regions.

- 20 Figure 3A further shows the effect of atmospheric transport on deaths in each Chinese region due to air pollution produced in other regions, with particularly large interregional impacts highlighted by arrows. The red shading in figure 3A corresponds to regions (e.g., the North, Yangtze River Delta and Northwest) whose emissions caused a greater number of deaths in other regions than pollution in other regions caused in that region—a net export of air pollution deaths. In contrast, the blue shaded regions (the Southwest, Southeast and Central) experienced greater numbers of deaths due to extra-regional emissions than
- 25 their emissions caused in other regions. Regionally, pollution from the North region that is transported in the atmosphere to the populous Central and Yangtze River Delta regions is particularly harmful and causes the most deaths, with 43,218 (95% CI : 26,354-52,300) and 19,620 (95% CI : 12,003-24,102) deaths related to these trans-boundary flows, respectively. Perhaps due to its substantial emissions and central location in the country, deaths caused in the Central region by emissions produced elsewhere and deaths caused by Central region's emissions transported elsewhere are approximately equal [114,959 (95% CI : 112,003-24,102)
- 30 69,481-137,887) and 101,252 (95% CI :63,431-121,995), respectively], and Southwest bear the most pollution deaths from emission in Central. Nationally, the net flows of trans-boundary PM2.5 related health impact mainly caused by pollution transported from north to south and from east to west.





5

3.3 Effects of interregional trade on regional mortality

Compare to the physical atmospheric transport, trade enable the cross-regional impact more broad (figure 2B vs. 2C), as the production of emissions can occurred far from where the products were finally consumed. However, for a finished products, it may experience different stages before being sold to final consumers, such as material production and products assemble, these may occurred in different regions. Figure 3B shows the effect of trade between the region producing a raw material or intermediate good and the region producing the final good ready for consumption. This is important because in many cases the region assembling or otherwise preparing the final good is able to capture a large fraction of the final good's value without undertaking more energy- and pollution-intensive processes that were required to produce the raw materials and intermediate goods (Prell et al., 2014;Liu et al., 2016). This distinction is particularly relevant in China because previous studies have shown

- 10 that more affluent coastal provinces in China are increasingly importing intermediate goods and materials from less-developed provinces (Feng et al., 2013; Jiang et al., 2015; Zhao et al., 2015). Here, we find that deaths related to final goods produced in red-shaded regions are substantially greater than the deaths due to the emission produced in those regions. In particular, final goods assembled/manufactured in the Yangtze River Delta and Southeast regions led to 62,670 (95% CI : 38,972-75,536) and 37,269 (95% CI : 23,106-44,983) deaths due to emission in other regions, respectively. In contrast, blue-shaded regions like
- 15 the Central, Northwest and Southwest are those which disproportionately produce and export raw materials and intermediate goods (e.g., mineral ores and metals), and therefore suffer air pollution deaths to support the manufacture of final goods in other regions. For example, 15% of deaths caused by emission in the Central region are related to final goods manufactured in the North, Yangtze River Delta, and Southeast regions.

With supply chains or trade extending, finished products may finally be consumed by another region. Figure 3C further shows

- 20 the full effect of trade from the producer of emissions related deaths to the final consumer in map. As this study doesn't include premature deaths caused by international imports, in this figure we only present regional production related deaths caused by domestic consumption, deaths induced by goods and services produced in China for international export are shown in figure 4 separately. As figure 3C shows, deaths related to consumption in red-shaded regions are substantially greater than the deaths due to the emission produced in those regions. Note that, in this figure Central region shows net export of production related
- 25 premature deaths with all other six regions, this can be attributed to its abundance interregional export, severe pollution and high population density. Including international export, 36% of Central region's production related deaths were induced by consumption in other regions. Moreover, figure 3C also highlight the case of Northeast. Even though Northwest shows net pollution export with other regions (Zhao et al., 2015), but its exported emissions cause less deaths than the relative small emissions occurred on other regions to support consumption in Northeast, due to population and production intensities in this
- 30 region are far less than other regions, such as Central and North regions. Nationally, in 2010 24% of pollution deaths in China were related to goods and services consumed in a different region from where they were produced.

In 2010, international export accounted for approximately 14% of $PM_{2.5}$ -related deaths in China (cf. 12% in 2007 reported by Jiang et.al., 2015). Figure 4 attributed these deaths to regions where the emission were produced and where the final products





5

were exported. As shown, of these international exports, roughly three-quarters (76%) of the related deaths are associated with the exports from the east coast regions (North, Yangtze River Delta and Southeast). However, only 58% of deaths related to exports from these coastal regions are caused by emissions actually produced in those regions, and even fewer (48%) of associated deaths actually occurred in those regions due to atmospheric transport. These results emphasize that international exports commonly entail intermediate inputs from less-developed regions of China (e.g., the Central region; (Feng et al., 2013;

Zhao et al., 2015)).

Figure 3D finishes the comparison, and showing the combined effect of atmospheric transport and trade on each region. Same as figure 3C, deaths related to international export were exclude from this map. Here, deaths related to final goods consumed in red-shaded regions are substantially greater than the deaths occurred in those regions. Even though similar with figure 3C,

10 figure 3D highlight that atmospheric transport aggravated the pollution deaths outsourcing from North and Y.R.D to Central and Southwest. As a combined result from atmospheric transport and trade, nationally 56% of pollution deaths in China were related to consumption in a different region (14% for international export).

Summing from the previous sections, figure 5 integrates the results related to both atmospheric transport and trade to show air pollution deaths in each region due to local and other regions' manufacturing and consumption activities, and separating the

- 15 effects of locally-produced and atmospherically-transported pollution. For a given region, emissions produced in the region to supply either local or other regions' consumption account for the largest share of deaths in the region (60-70%; purple and light blue bars in figure 5A), followed by the 'spillover impact' of emissions produced in other regions that are not related to the local region's manufacturing or consumption activities (27-37%; gray bars in figure 5A). Emissions produced in other regions and related to the local region's consumption contributed 1-3% of each regions' mortality, via atmospheric trans-
- 20 boundary transport (dark blue bars in figure 5A). Finally, the effect of atmospheric transport from other countries contributed only 1-2% of deaths in any Chinese region (light purple bars in figure 5A).

Figure 5B further breakdown the regions involved in each region's spillover impacts according to where the emissions were produced and where related goods were ultimately consumed, respectively. The magnitude of spillover deaths depends largely on a given region's population and the extent of emissions in their upwind regions. For instance, 75% and 71% of spillover

25 deaths in the Yangtze River Delta and Southwest regions are linked to emissions in upwind regions (primarily the Central and North regions), respectively. As the most populated region, the Central region suffered the most spillover deaths [108,963 (95% CI: 66,444-131,860)], 69% of which were related to emission produced in the North and Yangtze River Delta regions.

3.4 Uncertainties discussion

As evident from the analysis above, the calculation of pollution deaths caused by atmospheric transport and trade is subject to 30 uncertainties in emission inventories, chemical transport model, satellite-based PM_{2.5} concentration, heath impact model and MRIO model.





5

Bottom-up emission inventories are uncertain due to incomplete knowledge about activity, technology distribution and emission factors. The uncertainties in China's emission inventory were estimated to be -14%~13%, -13%~37%, -17%~54%, -25%~136% and -40%~121% for SO₂, NO_x, PM_{2.5}, black carbon (BC), and organic carbon (OC), respectively (Zhao et al., 2011). Although the quantitative uncertainties are not provided by the MEIC inventory, it has been widely used in chemical transport models and validated against surface and satellite observations (Zheng et al., 2015;Li et al., 2015;Geng et al., 2015;Hu

et al., 2016;Chen et al., 2015).

Uncertainties from the simulation of GEOS-Chem and its adjoint subject to their limitations or errors in chemical and physical representation, such as the chemical conversion, diffusion, deposition, and advection transport. Here we conduct a comparison of the modelled and the satellite-derived $PM_{2.5}$ concentration, and use the normalized mean error (NME) between these two

- 10 datasets over China seven regions to represent the overall model errors. As figure 6 shown, the NME various among seven regions, and range from the lowest in North (30%) to the highest in Northwest (71%). Note that the two datasets agree reasonably well, the R range from 0.67 to 0.95 for seven regions. This provides confidence that the results of this study are based on realistic simulation. The adjoint model may introduce additional uncertainties due to lack consideration of the nonlinear response of the predicted concentration to perturbation of emission input. However, due to its complex in backward
- 15 calculation and integration with the forward model, there are very few statistical quantification for its uncertainties so far. Lee et.al (2015) used $\pm 40\%$ to represent the total uncertainties caused by the GEOS-Chem adjoint model.

Uncertainties in satellite-derived $PM_{2.5}$ map is $\pm 5\%$ on average according to GBD 2013 (Brauer et al., 2016), as it has been calibrated by satellite-based and surface observations. Uncertainties in IER model are relatively high, mainly arising from the model itself, as it is fitted by limited information on actual exposure to $PM_{2.5}$ for source-specific relative risks. Burnett et.al

- 20 (Burnett et al., 2015) estimated the uncertainties from IER model by using simulation approach, and they fitted out 1000 sets parameters for the IER function to represent its possible shape. Additionally, the IER model is limited to several assumptions, e.g. PM_{2.5} related health impact is independent of exposure period, PM_{2.5} composition and toxicity for particles from different sources (Burnett et al., 2015; Jiang et al., 2015; Lee et al., 2015).
- Additional uncertainties originate from MRIO analysis when linking trade among different regions. MRIO model inherit all uncertainties in its source (survey) data and data manipulation (Peters, 2007; Weber, 2008; Wiedmann et al., 2011; Wiedmann, 2009). In addition, MRIO analysis is limited to sector detail, region coverage, and the number of environmental extensions (Tukker and Dietzenbacher, 2013). The study conducted by Lin et al (2014) found that the uncertainty in Chinese input-output model are relative small, it contributes ~10% of total errors in export-related pollutant emissions.

A comprehensive uncertainty analysis combining all affecting factors above is impossible due to the limitations of the

30 computational loads. The uncertainty ranges presented in this study only represent the uncertainties in IER function, which is obtained by 1000 sets runs of IER parameters fitted by Burnett et.al (2015) to calculate the possible distribution of regional pollution deaths.





4 Discussion and Conclusions

Patterns of atmospheric PM_{2.5} pollution and resulting premature deaths in China are the result of complex interacting physical transport processes and economic activities (Lin et al., 2014). As we have shown, large numbers of air pollution deaths are caused by economic activities in a different region from where the deaths occur. Moreover, more economically-developed
regions (e.g., the Yangtze River Delta, Southeast and North regions) tend to externalize their emissions and related health impacts by importing goods from less economically-developed regions (e.g., the Central region). Pollution abatement is thus not simply local problem, but one in which multi-regional coordination may be necessary to effectively and efficiently reduce emissions and their health impacts. This is especially true because, as we have shown, relocating emissions within the nation will not completely alleviate the environmental and health burden; atmospheric transport of pollution often leads to health

10 impacts in other, downwind regions, including both regions where the related goods are ultimately consumed and regions with no connection to the economic activities at all (i.e. spillover).

As a main driver of China's production, international export accounted for 14% of China's anthropogenic PM2.5 -related premature deaths in 2010. Moreover, its impact was not evenly distributed among regions, as the developed east coastal regions partly transfer their export related pollution deaths to the less developed central and west regions by importing raw material

- 15 from the less developed central and west regions (figure 3B and figure 4). This exerts disproportionally life loss and economic gains from international exports among regions (Jiang et al., 2015). Besides, the added pollution results from international export can further affect other countries atmospheric environment (Lin et al., 2016) or even pollution deaths (Takahashi et al., 2014) through cross-continental atmospheric transport. Thus a jointed pollution mitigation action among regions, nations and even production chains is in urgent needed, not only for domestic equality in development, but also for global human health.
- 20 Our results represent the most detailed analysis of air pollution mortality in China, its sources, and its underlying economic drivers. Based on these findings, future measures to alleviate these health impacts could be prioritized according to the source and location of emissions as well as the type and economic value of the emitting activities and related patterns of consumption.

Acknowledgements. This study was supported by the National Science Foundation of China (41625020, 41629051, 41541039 and 41222036) and China's National Basic Research Program (2014CB441301). G.P. Peters was funded by the Norwegian

25 Research Council (235523). Q. Zhang and K. B. He are supported by the Collaborative Innovation Center for Regional Environmental Quality.





Appendix A

Table A1. Region Definitions

Region	Provinces/municipalities included in each region
North	Beijing, Tianjin, Hebei and Shandong
Yangtze River Delta	Shanghai, Jiangsu and Zhejiang
Southeast	Fujian, Guangdong and Hainan
Central	Henan, Anhui, Hubei, Hunan and Jiangxi
Northwest	Shaanxi, Shanxi, Gansu, Qinghai, Ningxia, Xinjiang and Inner Mongolia
Southwest	Sichuan, Chongqing, Guizhou, Yunnan and Guangxi
Northeast	Liaoning, Jilin and Heilongjiang





References:

25

30

- Apte, J. S., Marshall, J. D., Cohen, A. J., and Brauer, M.: Addressing Global Mortality from Ambient PM2.5, Environ. Sci. Technol., 49, 8057-8066, doi:10.1021/acs.est.5b01236, 2015.
- Bhalla, K., Shotten, M., Cohen, A., Brauer, M., Shahraz, S., Burnett, R., Leach-Kemon, K., Freedman, G., and Murray, C.:
- 5 Transport for health: the global burden of disease from motorized road transport. Global Road Safety Facility, Institute for Health Metrics and Evaluation and World Bank, Washington DC, 2014.
 - Brauer, M., Freedman, G., Frostad, J., van Donkelaar, A., Martin, R. V., Dentener, F., Dingenen, R. v., Estep, K., Amini, H., Apte, J. S., Balakrishnan, K., Barregard, L., Broday, D., Feigin, V., Ghosh, S., Hopke, P. K., Knibbs, L. D., Kokubo, Y., Liu, Y., Ma, S., Morawska, L., Sangrador, J. L. T., Shaddick, G., Anderson, H. R., Vos, T., Forouzanfar, M. H., Burnett,
- R. T., and Cohen, A.: Ambient Air Pollution Exposure Estimation for the Global Burden of Disease 2013, Environ. Sci. Technol., 50, 79-88, doi:10.1021/acs.est.5b03709, 2016.
 - Burnett, R. T., Pope III, C. A., Ezzati, M., Olives, C., Lim, S. S., Mehta, S., Shin, H. H., Singh, G., Hubbell, B., and Brauer, M.: An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure, Environ. Health Perspect., 122, 397-403, doi: :10.1289/ehp.1307049, 2014.
- 15 Chafe, Z. A., Brauer, M., Klimont, Z., Van Dingenen, R., Mehta, S., Rao, S., Riahi, K., Dentener, F., and Smith, K. R.: Household cooking with solid fuels contributes to ambient PM_{2.5} air pollution and the burden of disease, Environ. Health Perspect., 122, 1314-1320,doi: 10.1289/ehp.1206340, 2014.
 - Chambliss, S. E., Silva, R., West, J. J., Zeinali, M., and Minjares, R.: Estimating source-attributable health impacts of ambient fine particulate matter exposure: global premature mortality from surface transportation emissions in 2005, Environ. Res.
- 20 Lett., 9, 104009, doi:10.1088/1748-9326/9/10/104009,2014.
 - Chen, Y., Zhang, Y., Fan, J., Leung, L.-Y., Zhang, Q., and He, K.: Application of an Online-Coupled Regional Climate Model, WRF-CAM5, over East Asia for Examination of Ice Nucleation Schemes: Part I. Comprehensive Model Evaluation and Trend Analysis for 2006 and 2011, Climate, 3, 627-667, 2015.
 - Davis, S. J., Peters, G. P., and Caldeira, K.: The supply chain of CO₂ emissions, P. Natl. Acad. Sci. USA, 108, 18554-18559, doi:10.1073/pnas.1107409108, 2011.
 - Dedoussi, I. C., and Barrett, S. R. H.: Air pollution and early deaths in the United States. Part II: Attribution of PM_{2.5} exposure to emissions species, time, location and sector, Atmospheric Environment, 99, 610-617, doi:0.1016/j.atmosenv.2014.10.033, 2014.
 - Feng, K., Davis, S. J., Sun, L., Li, X., Guan, D., Liu, W., Liu, Z., and Hubacek, K.: Outsourcing CO2 within China, P. Natl. Acad. Sci. USA, 110, 11654-11659, doi: 10.1073/pnas.1219918110, 2013.
 - Forouzanfar, M. H., Alexander, L., Anderson, H. R., Bachman, V. F., Biryukov, S., Brauer, M., Burnett, R., Casey, D., Coates, M. M., and Cohen, A.: Global, regional, and national comparative risk assessment of 79 behavioural, environmental and





5

occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013, The Lancet, 386, 2287-2323, doi: 10.1016/S0140-6736(15)00128-2, 2015.

- Geng, G., Zhang, Q., Martin, R. V., van Donkelaar, A., Huo, H., Che, H., Lin, J., and He, K.: Estimating long-term PM_{2.5} concentrations in China using satellite-based aerosol optical depth and a chemical transport model, Remote Sens. Environ, 166, 262-270, doi:10.1016/j.rse.2015.05.016, 2015.
- Hamilton, K.: Calculating PM_{2.5} Damages as % of GDP for Top CO₂ Emitters: A Technical Note, New Climate Economy Contributing Paper. http://newclimateeconomy. report, 2015.
 - Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, Atmos. Chem. Phys., 7, 2413-2433, doi:10.5194/acp-7-2413-2007, 2007.
- Henze, D. K., Seinfeld, J. H., and Shindell, D. T.: Inverse modeling and mapping US air quality influences of inorganic PM
 2.5 precursor emissions using the adjoint of GEOS-Chem, Atmos. Chem. Phys., 9, 5877-5903, doi:10.5194/acp-9-5877-2009, 2009.
 - Hu, J., Chen, J., Ying, Q., and Zhang, H.: One-year simulation of ozone and particulate matter in China using WRF/CMAQ modeling system, Atmos. Chem. Phys., 16, 10333-10350, doi:10.5194/acp-16-10333-2016, 2016.
- 15 Huo, H., Zhang, Q., Guan, D., Su, X., Zhao, H., and He, K.: Examining Air Pollution in China Using Production- And Consumption-Based Emissions Accounting Approaches, Environ. Sci. Technol., 48, 14139-14147, doi:10.1021/es503959t, 2014.
 - Jiang, X., Zhang, Q., Zhao, H., Geng, G., Peng, L., Guan, D., Kan, H., Huo, H., Lin, J., Brauer, M., Martin, R. V., and He, K.: Revealing the Hidden Health Costs Embodied in Chinese Exports, Environ. Sci. Technol., 49, 4381-4388, doi:10.1021/es506121s, 2015.
 - Lee, C. J., Martin, R. V., Henze, D. K., Brauer, M., Cohen, A., and Donkelaar, A. v.: Response of global particulate-matterrelated mortality to changes in local precursor emissions, Environ. Sci. Technol., 49, 4335-4344, doi: 10.1021/acs.est.5b00873, 2015.
 - Lei, Y., Zhang, Q., He, K. B., and Streets, D. G.: Primary anthropogenic aerosol emission trends for China, 1990–2005, Atmos.

Chem. Phys., 11, 931-954, doi:10.5194/acp-11-931-2011, 2011.

25

20

- Li, B., Gasser, T., Ciais, P., Piao, S., Tao, S., Balkanski, Y., Hauglustaine, D., Boisier, J.-P., Chen, Z., Huang, M., Li, L. Z., Li, Y., Liu, H., Liu, J., Peng, S., Shen, Z., Sun, Z., Wang, R., Wang, T., Yin, G., Yin, Y., Zeng, H., Zeng, Z., and Zhou, F.: The contribution of China's emissions to global climate forcing, Nature, 531, 357-361, doi:10.1038/nature17165, 2016.
- 30 Li, X., Zhang, Q., Zhang, Y., Zheng, B., Wang, K., Chen, Y., Wallington, T. J., Han, W., Shen, W., Zhang, X., and He, K.: Source contributions of urban PM_{2.5} in the Beijing–Tianjin–Hebei region: Changes between 2006 and 2013 and relative impacts of emissions and meteorology, Atmos. Environ., 123, Part A, 229-239, doi:10.1016/j.atmosenv.2015.10.048, 2015.





20

Lin, J., Nielsen, C. P., Zhao, Y., Lei, Y., Liu, Y., and McElroy, M. B.: Recent Changes in Particulate Air Pollution over China Observed from Space and the Ground: Effectiveness of Emission Control, Environ. Sci. Technol., 44, 7771-7776, doi:10.1021/es101094t, 2010.

Lin, J., Pan, D., Davis, S. J., Zhang, Q., He, K., Wang, C., Streets, D. G., Wuebbles, D. J., and Guan, D.: China's international

- 5 trade and air pollution in the United States, P. Natl. Acad. Sci. USA, 111, 1736-1741, doi: 10.1073/pnas.1312860111, 2014.
 - Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., and He, K. B.: High-resolution inventory of technologies, activities, and emissions of coal-fired power plants in China from 1990 to 2010, Atmos. Chem. Phys., 15, 13299-13317, 10.5194/acp-15-13299-2015, 2015.
- 10 Liu, W., Tang, Z., Chen, J., and Yang, B.: Multi-regional Input-output model for 30 provinces of China in 2010, China Statistics Press, Beijing, 2014.
 - Liu, Z., Davis, S. J., Feng, K., Hubacek, K., Liang, S., Anadon, L. D., Chen, B., Liu, J., Yan, J., and Guan, D.: Targeted opportunities to address the climate-trade dilemma in China, Nature Clim. Change, 6, 201-206, doi: 10.1038/nclimate2800, 2016.
- 15 Ma, Z., Hu, X., Huang, L., Bi, J., and Liu, Y.: Estimating Ground-Level PM2.5 in China Using Satellite Remote Sensing, Environ. Sci. Technol., 48, 7436-7444, doi:10.1021/es5009399, 2014.

National Bureau of Statistics (NBS): China Statistical Yearbook, China Statistics Press, Beijing, China, 2011.

World Health Organization (WHO) Air quality guidelines global update 2005. WHO/Europe, Copenhagen, Denmark, 2005.

Pappin, A. J., and Hakami, A.: Source attribution of health benefits from air pollution abatement in Canada and the United States: an adjoint sensitivity analysis, Environ. Health Perspect., 121, 572–579, doi:10.1289/ehp.1205561, 2013

- Peters, G. P.: Opportunities and challenges for environmental MRIO modeling: illustrations with the GTAP database, 16th International Input-Output Conference, 2007.
 - Prell, C., Feng, K., Sun, L., Geores, M., and Hubacek, K.: The Economic Gains and Environmental Losses of US Consumption: A World-Systems and Input-Output Approach, Social Forces, 93: 405-428, doi:10.1093/sf/sou048, 2014.
- 25 Takahashi, K., Nansai, K., Tohno, S., Nishizawa, M., Kurokawa, J.-i., and Ohara, T.: Production-based emissions, consumption-based emissions and consumption-based health impacts of PM2.5 carbonaceous aerosols in Asia, Atmospheric Environment, 97, 406-415, doi:10.1016/j.atmosenv.2014.04.028, 2014.
 - Tukker, A., and Dietzenbacher, E.: Global Multiregional Input-output frameworks: An introduction and outlook, Econ Syst Res, 25, 1-19, 10.1080/09535314.2012.761179, 2013.
- 30 Turner, M. D., Henze, D. K., Capps, S. L., Hakami, A., Zhao, S., Resler, J., Carmichael, G. R., Stanier, C. O., Baek, J., and Sandu, A.: Premature deaths attributed to source-specific BC emissions in six urban US regions, Environ. Res. Lett., 10, 114014, doi:10.1088/1748-9326/10/11/114014, 2015.
 - Weber, C. L.: Uncertainties in constructing environmental multiregional input-output models, International input-output meeting on managing the environment, 2008,





- Wiedmann, T.: A review of recent multi-region input–output models used for consumption-based emission and resource accounting, Ecol Econ, 69, 211-222, doi:10.1016/j.ecolecon.2009.08.026, 2009.
- Wiedmann, T., Wilting, H. C., Lenzen, M., Lutter, S., and Palm, V.: Quo Vadis MRIO? Methodological, data and institutional requirements for multi-region input–output analysis, Ecol Econ, 70, 1937-1945, doi:10.1016/j.ecolecon.2011.06.014, 2011.

5

- Yang, G., Wang, Y., Zeng, Y., Gao, G. F., Liang, X., Zhou, M., Wan, X., Yu, S., Jiang, Y., Naghavi, M., Vos, T., Wang, H., Lopez, A. D., and Murray, C. J. L.: Rapid health transition in China, 1990-2013: findings from the Global Burden of Disease Study 2010, The Lancet, 381, 1987-2015, 10.1016/S0140-6736(13)61097-1,
- Zhang, L., Liu, L., Zhao, Y., Gong, S., Zhang, X., Henze, D., K., L. Capps, S., Fu, T.-M., Zhang, Q., and Wang, Y.: Source
- 10attribution of particulate matter pollution over North China with the adjoint method, Environ. Res. Lett., 10, 084011, doi:10.1088/1748-9326/10/8/084011, 2015.
 - Zhang, Q., Streets, D. G., Carmichael, G. R., He, K. B., Huo, H., Kannari, A., Klimont, Z., Park, I. S., Reddy, S., Fu, J. S., Chen, D., Duan, L., Lei, Y., Wang, L. T., and Yao, Z. L.: Asian emissions in 2006 for the NASA INTEX-B mission, Atmos. Chem. Phys., 9, 5131-5153, doi:10.5194/acp-9-5131-2009, 2009.
- 15 Zhao, H. Y., Zhang, Q., Guan, D. B., Davis, S. J., Liu, Z., Huo, H., Lin, J. T., Liu, W. D., and He, K. B.: Assessment of China's virtual air pollution transport embodied in trade by using a consumption-based emission inventory, Atmos. Chem. Phys., 15, 5443-5456, doi:10.5194/acp-15-5443-2015, 2015.
 - Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B., and Hao, J.: Quantifying the uncertainties of a bottom-up emission inventory of anthropogenic atmospheric pollutants in China, Atmos. Chem. Phys., 11, 2295-2308, doi:10.5194/acp-11-2295-2011, 2011.
 - Zheng, B., Zhang, Q., Zhang, Y., He, K. B., Wang, K., Zheng, G. J., Duan, F. K., Ma, Y. L., and Kimoto, T.: Heterogeneous chemistry: a mechanism missing in current models to explain secondary inorganic aerosol formation during the January 2013 haze episode in North China, Atmos. Chem. Phys., 15, 2031-2049, doi:10.5194/acp-15-2031-2015, 2015.

25

20





Table 1. China seven regions' populations, PM_{2.5} concentrations, mortality and mortality ratios in 2010

Region	Population (millions)	Population weighted PM _{2.5} concentrations (µg	Mortality (thousands of deaths)	Mortality ratio (deaths per 10 ⁵
		m ⁻³)		persons)
North	192	82	203 (95%CI:127-239)	106 (95%CI :66-125)
Yangtze River Delta	144	50	125 (76-153)	87 (53-107)
Southeast	143	27	80 (52-98)	56 (36-69)
Central	337	67	343 (209-415)	102 (62-123)
Northwest	157	34	98 (62-118)	63 (40-76)
Southwest	254	52	211(132-254)	83 (52-100)
Northeast	114	28	66 (42-82)	58 (37-72)
National average/ total	1354	53	1126 (701-1349)	84 (52-101)







Figure 1. Schematic of the methodology used in this study.







Figure 2. Anthropogenic PM_{2.5}-related premature deaths in China for 2010 at the 0.5 °×0.667 ° horizontal resolution (A), and regional premature deaths attributed to regions where emissions were produced (B) and regions where products were ultimately consumed (C). Data sets at the end of each bar mean the percentages of regional pollution death attributed to local source and the percentages attributed to other regions .Y.R.D. is the Yangtze River Delta.

5







Figure 3. The effect of atmospheric transport (A) and trade (B-D) on each region's pollution deaths. Panel A compares the number of pollution deaths related to emissions produced in each region and deaths that occurred in that region. Panel B and C compare regional production related premature deaths with deaths related to production of final goods

- 5 in that region, and deaths related to consumption of goods and services in that region, respectively. Panel D compares the number of pollution deaths occurred in each region with deaths related to consumption of goods and services in that region. Deaths in other regions due to Chinese pollution and deaths due to emissions in other nations are not included in any of the maps, and international export on regional pollution deaths are not included in map C and D. Arrows between regions denote the largest interregional transfers, with numbers of displaced air pollution deaths
- 10 shown in thousands.







Figure 4. Flow map of premature deaths connecting producers and international exporter of finished products. The percentage value are relative to national total anthropogenic PM_{2.5} related premature deaths (1.13 millions).







Figure 5. Source of pollution deaths in each region (A) and their "spillover" source by producing regions (i.e. the gray bars in A: deaths due to emissions in other regions related to goods and services consumed in other regions).







Figure 6. Comparisons between the simulated and satellite-derived PM_{2.5} concentrations over the seven China regions. 5