Modeling of the chemistry in oxidation flow reactors with high initial NO

Zhe Peng and Jose L. Jimenez

Cooperative Institute for Research in Environmental Sciences and Department of Chemistry, University of Colorado, Boulder, CO 80309, USA
Correspondence to: J.L. Jimenez (jose.jimenez@colorado.edu)

Abstract. Oxidation flow reactors (OFRs) are increasingly employed in atmospheric chemistry research because of their high efficiency of OH radical production from low-pressure Hg lamp emissions at both 185 and 254 nm (OFR185) or 254 nm only (OFR254). OFRs have been thought to be limited to studying low-NO chemistry (where peroxy radicals (RO_2) react preferentially with HO_2) because NO is very rapidly oxidized by the high concentrations of O_3, HO_2, and OH in OFRs. However, many groups are performing experiments aging combustion exhaust with high NO levels, or adding NO in the hopes of simulating high-NO chemistry (where RO_2 + NO dominates). This work systematically explores the chemistry in OFRs with high initial NO. Using box modeling, we investigate the interconversion of N-containing species and the uncertainties due to kinetic parameters. Simple initial injection of NO in OFR185 can result in more RO_2 reacted with NO than with HO_2 and minor non-tropospheric photolysis, but only under a very narrow set of conditions (high water mixing ratio, low UV intensity, low external OH reactivity (OHR_{ext}), and initial NO concentration (NO^{in}) of tens to hundreds of ppb) that account for a very small fraction of the input parameter space. These conditions are generally far away from experimental conditions of published OFR studies with high initial NO. In particular, studies of aerosol formation from vehicle emissions in OFR often used OHR_{ext} and NO^{in} several orders of magnitude higher. Due to extremely high OHR_{ext} and NO^{in}, some studies may have resulted in substantial non-tropospheric photolysis, strong delay to RO_2 chemistry due to peroxynitrate formation, VOC reactions with NO_3 dominating over those with OH, and faster reactions of OH-aromatic adducts with NO_2 than those with O_2, all of which are irrelevant to ambient VOC photooxidation chemistry. Some of the negative effects are worst for alkene and aromatic precursors. To avoid undesired chemistry, vehicle emissions generally need to be diluted by a factor of >100 before being injected into OFR. However, sufficiently diluted vehicle emissions generally do not lead to high-NO chemistry in OFR, but are rather dominated by the low-NO RO_2+HO_2 pathway. To ensure high-NO conditions without substantial atmospherically irrelevant chemistry in a more controlled fashion, new techniques are needed.

1 Introduction

The oxidation of gases that are emitted into the atmosphere, in particular volatile organic compounds (VOCs), is one of the most important atmospheric chemistry processes (Haagen-Smit, 1952; Chameides et al., 1988). VOC oxidation is closely related to radical production and consumption (Levy II, 1971), O_3 production, and formation of secondary aerosols (Odum et al., 1996; Hoffmann et al., 1997; Volkamer et al., 2006; Hallquist et al., 2009), which have impacts on air quality and climate (Lippmann, 1991; Nel, 2005; Stocker et al., 2014).

Chemical reactors are critical tools for research of VOC oxidation. Oxidation reactions of interest often have typical timescales of hours to weeks. Studying these processes in ambient air can be confounded by dispersion and changes in ambient conditions, which often occur on similar timescales. Chemical reactors allow for decoupling of these two types of processes. Also, they should be able to simulate the different regimes of reactions occurring in the atmosphere, e.g., VOC oxidation under low and high-NO conditions (peroxy radical fate dominated by reaction with HO_2 or with NO) representing remote and urban areas, respectively (Orlando and Tyndall, 2012).

Large environmental chambers are a commonly used reactor type (Carter et al., 2005; Wang et al., 2011). They typically employ actinic wavelength (>300 nm) light sources (e.g., outdoor solar radiation and UV blacklights) to produce oxidants and radicals and have large volumes (on the order of several cubic meters or larger). However, the capability of generating sustained elevated levels of OH, the most important tropospheric oxidant, is usually limited in chambers, resulting in OH concentrations similar to those in the atmosphere (10^6–10^7 molecules cm^{-3}; Mao et al., 2009; Ng et al., 2010), and consequently, long simulation times (typically hours) to reach OH equivalent ages of atmospheric relevance (George et al., 2007; Kang et al., 2007; Carlton et al., 2009; Seakins, 2010; Wang et al., 2011). The partitioning of gases and aerosols to chamber walls (usually made of Teflon) in timescales of tens of minutes to hours makes it difficult to conduct very long experiments that simulate high atmospherically-relevant photochemical ages (Cocker et al., 2001; Matsunaga and Ziemann, 2010; Zhang et al., 2014; Krechmer et al., 2016). In addition, the long simulation times and large size of chambers and auxiliary equipment are logistically difficult for field deployment, and their cost limits the number of laboratories equipped with them.

Given the limitations of environmental chambers, a growing number of experimenters have instead employed oxidation flow reactors (OFRs). OFRs have a much smaller size (of the order of 10 L), efficiently generate OH via photolysis of H_2O and/or O_3 by more energetic 185 and 254 nm photons from low-pressure Hg lamps, and overcome the abovementioned shortcomings of chambers due to a much shorter residence time (George et al., 2007; Kang et al., 2007, 2011; Lambe et al., 2011). Moreover, OFRs are able to rapidly explore a wide range of OH equivalent ages within a short period (~2 hr), during which significant changes of ambient conditions can usually be avoided in the case of field deployment (Ortega et al., 2016; Palm et al., 2016, 2017). Because of these advantages, OFRs have recently been widely used to study atmospheric chemistry, in particular secondary organic aerosol (SOA) formation and aging, in both the laboratory and the field (Kang et al., 2011; Li et al., 2013; Ortega et al., 2013,

2016; Tkacik et al., 2014; Palm et al., 2016).

In addition to experimental studies using OFRs, there has also been some progress in the characterization of OFR chemistry by modeling. Li et al. (2015) and Peng et al. (2015) developed a box model for OFR HO_x chemistry that predicts measurable quantities [e.g., OH exposure (OH_{exp}, in molecules cm^{-3} s] and O_3 concentration (abbr. O_3 hereinafter, in ppm)] in good agreement with experiments. This model has been used to characterize HO_x chemistry as a function of H_2O mixing ratio (abbr. H_2O hereinafter, unitless), UV light intensity (abbr. UV hereinafter, in photons cm^{-2} s^{-1}), and external OH reactivity [in s^{-1}, $OHR_{ext}=\sum k_i c_i$, i.e., the sum of the products of concentrations of externally introduced OH-consuming species (c_i) and rate constants of their reactions with OH (k_i)]. Based on this characterization, Peng et al. (2015) found that OH suppression, i.e., reduction of OH concentration caused by OHR_{ext}, is a common feature under many typical OFR operation conditions. Peng et al. (2016) systematically examined the relative importance of non-OH/non-tropospheric reactants on the fate of VOCs over a wide range of conditions, and provided guidelines for OFR operation to avoid non-tropospheric VOC photolysis, i.e., VOC photolysis at 185 and 254 nm.

In previous OFR modeling studies, NO_x chemistry was not investigated in detail, since in such in typical OFR experiments with large amounts of oxidants (e.g., OH, HO_2, and O_3), NO would be very rapidly oxidized and thus unable to compete with HO_2 for reaction with peroxy radicals (RO_2). Li et al. (2015) estimated an NO (NO_2) lifetime of ~0.5 (~1.5) s under a typical OFR condition. From these estimates, OFRs processing ambient air or laboratory air without large addition of NO_x were assumed to be not suitable for studying oxidation mechanisms relevant to polluted conditions under higher NO concentrations. OFRs have recently been used to conduct laboratory experiments with very high initial NO_x levels (Liu et al., 2015) and deployed to an urban tunnel, where NO_x was high enough to be a major OH reactant (Tkacik et al., 2014). The former study reported evidence for the incorporation of nitrogen into SOA. Besides, OFRs have been increasingly employed to process emissions of vehicles, biomass burning, and other combustion sources (Table 1), where NO can often be hundreds of ppm (Ortega et al., 2013; Martinsson et al., 2015; Karjalainen et al., 2016; Link et al., 2016; Schill et al., 2016; Alanen et al., 2017; Simonen et al., 2017). It can be expected that such a high NO input together with very high VOC concentrations would cause a substantial deviation from good OFR operation conditions identified in Peng et al. (2016). Very recently, N_2O injection has been proposed by Lambe et al. (2017) as a way to study oxidation of VOCs under high NO conditions in OFR. As more OFR studies at high NO_x level are conducted, there is growing need to understand the chemistry of N-containing species in OFRs and whether it proceeds along atmospherically-relevant channels.

In this study, we present the first comprehensive model of OFR NO_y chemistry. We extend the model of Li et al. (2015) and Peng et al. (2015) by including a scheme for NO_y species. Then this model is used to investigate i) if an OFR with initial NO injection results in NO significantly reacting with RO_2 under any conditions, ii) if previously published OFR experiments with high initial NO concentrations led to RO_2+NO being dominant in VOC oxidation without negative side effects (e.g., non-tropospheric reactions), iii) how to avoid undesired chemistry in future studies. The results can provide insights into

the design and interpretation of future OH-oxidation OFR experiments with large amounts of NO_x injection.

2 Methods

The physical design of the OFR modeled in the present work, the chemical kinetics box model, and the method of propagating and analyzing the parametric uncertainties on the model have already been introduced previously (Kang et al., 2007; Li et al., 2015; Peng et al., 2015). We only provide brief descriptions for them below.

2.1 Potential Aerosol Mass flow reactor

The OFR modeled in this study is the "Potential Aerosol Mass" (PAM) flow reactor, firstly introduced by Kang et al. (2007). The PAM OFR is a cylindrical vessel with a volume of ~13 L, equipped with low-pressure Hg lamps (model no. 82-9304-03, BHK Inc.) to generate 185 and 254 nm UV light. This popular design is being used by many atmospheric chemistry research groups, particularly those studying SOA (Lambe and Jimenez, 2017 and references therein). When the lamps are mounted inside Teflon sleeves, photons at both wavelengths are transmitted and contribute to OH production ("OFR185 mode"). In OFR185, H_2O photolyzed at 185 nm produces OH and HO_2, while O_2 photolyzed at the same wavelengths results in O_3 formation. $O(^1D)$ is produced via O_3 photolysis at 254 nm and generates additional OH through its reaction with H_2O. 185 nm lamp emissions can be filtered by mounting the lamps inside quartz sleeves, leaving only 254 nm photons to produce OH ("OFR254 mode"). In this mode, injection of externally formed O_3 is necessary to ensure OH production. As the amount of O_3 injected is a key parameter under some conditions (Peng et al., 2015), we adopt the notation OFR254-X to denote OFR254 experiments with X ppm initial O_3 ($O_{3,in}$). In this study, we investigate OFR experiments with NO injected and thus utilize "OFR185-iNO" to describe the OFR185 mode of operation with initially (at the reactor entrance) injected NO. The same terminology is used for the OFR254 mode. For instance, the initial NO injection into OFR254-7 is denoted as OFR254-7-iNO.

2.2 Model description

The basic framework of the box model used in this study, a standard chemical kinetics model, is the same as in Peng et al. (2015). Plug flow is assumed in the model, since approximately taking residence time distribution into account leads to similar results under most conditions but at much higher computational expense (Peng et al., 2015). In addition to the reactions in the model of Peng et al. (2015), including all HO_x reactions available in the JPL Chemical Kinetic Data Evaluation (Sander et al., 2011), all gas-phase NO_y reactions available in the JPL database except those of organic nitrates and peroxynitrates are also considered in the current reaction scheme. An updated JPL evaluation was published recently (Burkholder et al., 2015), with slightly different (~20%) rate constants for $NO_2+HO_2+M \rightarrow HO_2NO_2+M$ and $NO_2+NO_3 \rightarrow N_2O_5$. The updated rate constants only result in changes of ~10–20% of the concentrations of the species directly consumed/produced by these reactions. These changes are smaller than the parametric uncertainties of the model (see Section 3.1.3). For other species, concentration changes are negligible. $HO_2NO_2+M \rightarrow HO_2+NO_2+M$ and $N_2O_5+M \rightarrow NO_2+NO_3+M$, are also included in the scheme, with kinetic parameters from the IUPAC Task Group on Atmospheric

Chemical Kinetic Data Evaluation (Ammann et al., 2016). As in Peng et al. (2015, 2016), SO_2 is used as a surrogate of external OH reactants (e.g., VOCs). NO_y species, although also external OH reactants, are explicitly treated in the model and *not* counted in OHR_{ext} in this work. Therefore, OHR_{ext} stands for *non*-NO_y OHR_{ext} only hereinafter, unless otherwise stated.

Also, particle-phase chemistry and physical and chemical interactions of gas-phase species with particles are not considered in this study. We have made this assumption because:

i) The presence of aerosols has typically negligible impacts on the gas-phase chemistry of radicals, NO_x/NO_y, and OH reactants studied here. Condensational sink (CS) of ambient aerosols can rarely exceed 1 s^{-1} even in polluted areas and is usually 1-3 orders of magnitude lower (Donahue et al., 2016; Palm et al., 2016). Thus, even under the assumption of unity uptake coefficient, CS cannot compete with OHR_{ext} (usually on the order of 10 s^{-1} or higher) in OH loss. Uptake of NO onto aerosols only occurs through the reaction with RO_2 on particle surface (Richards-Henderson et al., 2015), which is formed very slowly (see below) compared to gas-phase HO_x and NO_x chemistry. Uptake of HO_2, O_3, NO_3 etc. is even more unlikely to be of importance due to lower uptake coefficients (Moise and Rudich, 2002; Moise et al., 2002; Hearn and Smith, 2004; Lakey et al., 2015). Combustion exhausts can have high aerosol loadings with condensational sinks on the order of 10^2–10^3 s^{-1} (Matti Maricq, 2007). Even if these exhausts are directly injected into the reactor without any pre-treatment, uptake onto the particles still cannot play a major role in the fate of gas-phase radical and NO_x species, since VOCs and NO_x in raw exhausts, which are proportionally orders-of-magnitude higher, still dominate the fate of oxidants. Dilution of combustion emissions simultaneously lowers condensational sinks and the sinks of oxidants due to chemical reactions, with their relative importance remaining the same as in undiluted emissions.

ii) Gas-phase radical and NO_x/NO_y species only has limited impacts on OA *chemistry* in this study. Heterogeneous oxidation of OA by OH is generally slow. Significant OA loss due to heterogeneous oxidation can only be seen at photochemical ages as high as weeks (Hu et al., 2016). The enhancement of heterogeneous oxidation due to NO is remarkable only at OH concentration close to the ambient values but not at typical values in OFR (Richards-Henderson et al., 2015).

It is well known that the aerosol concentration can have a major impact on the physical uptake of semivolatile and low-volatility gas-phase species. However this process is not explicitly modeled in this study.

As OHR_{ext} plays a major and even dominant role in OH loss, it is an important approximation that the *real* OHR_{ext} decay (due to primary VOC oxidation and subsequent oxidation of higher generation products) is surrogated by that of SO_2 (see Fig. S2 of Peng et al. 2015). Gas-phase measurements in literature laboratory studies revealed that there is a large variability of the evolution of total OHR_{ext} during oxidation of primary VOCs and subsequent oxidation of their intermediate products, depending on the type of precursors (Nehr et al., 2014; Schwantes et al., 2017). This variability is obviously mainly

due to the formation of different types and amounts of oxidation intermediates/products contributing to OHR_{ext}. This variation is highly complex due to the large number of possible oxidation intermediates and the limited knowledge of detailed higher-generation mechanisms, and thus is difficult to accurately capture even if modeling with a mechanism as explicit as Master Chemical Mechanism is performed (Schwantes et al., 2017). Therefore, it is justified to use a lumped surrogate to model the OHR_{ext} decay for simplicity and efficiency. This approximation is a substantial contributor to uncertainty of our model. The uncertainties due to the types of oxidation intermediates/products are very likely larger than those due to mass transfer processes between gas and particle phases, wall losses, etc., which are not considered in this study.

A residence time of 180 s and typical temperature (295 K) and atmospheric pressure (835 mbar) in Boulder, CO, USA are assumed for all model cases. The lower-than-sea level pressure only leads to minor differences in the outputs (Li et al., 2015). We explore physical input cases evenly spaced in a logarithmic scale over very wide ranges: H_2O of 0.07%–2.3%, i.e., relative humidity (RH) of 2–71% at 295 K; 185 nm UV of $1.0x10^{11}$–$1.0x10^{14}$ and 254 nm UV of $4.2x10^{13}$–$8.5x10^{15}$ photons cm^{-2} s^{-1}; OHR_{ext} of 1–16000 s^{-1}; $O_{3,in}$ of 2.2–70 ppm for OFR254; initial NO mixing ratio (NO^{in}) from 10 ppt to 40 ppm. Besides, conditions with OHR_{ext}=0 are also explored. UV at 254 nm is estimated from that at 185 nm according to the relationship determined by Li et al. (2015). Several typical cases within this range as well as their corresponding 4 or 2-character labels (e.g., MM0V and HL) are defined in Table 2. Literature studies are modeled by adopting all reported parameters (e.g., residence time, H_2O, and $O_{3,in}$) and estimating any others that may be needed (e.g., UV) from the information provided in the papers.

In this study, OH equivalent ages are calculated under the assumption of an ambient OH concentration of $1.5x10^6$ molecules cm^{-3} (Mao et al., 2009). Conditions leading to a ratio of RO_2 reacted with NO over the entire residence time [$r(RO_2+NO)$] to that with HO_2 [$r(RO_2+HO_2)$] larger than 1 are regarded as "high NO" (under the assumption of constant OHR_{ext} from VOCs, see Section S1 for more details), where [r(X)] is the total reactive flux for reaction X over the entire residence time. $F185_{exp}/OH_{exp}$ and $F254_{exp}/OH_{exp}$ are used as measures of the relative importance of VOC photolysis at 185 and 254 nm to their reactions with OH, respectively [$F185_{exp}$ ($F254_{exp}$) are 185 (254) nm photon flux exposure, i.e., product of 185 (254) nm photon flux and time]. Readers may refer to Figs. 1 and 2 of Peng et al. (2016) for the determination of the relative importance of non-tropospheric (185 and 254 nm) photolysis of individual VOCs. Although the relative importance of non-tropospheric photolysis depends on individual VOCs, in the present work, we set criteria on $F185_{exp}/OH_{exp}<3x10^3$ cm/s and $F254_{exp}/OH_{exp}<4x10^5$ cm/s to define "good" conditions and $F185_{exp}/OH_{exp}<1x10^5$ cm/s and $F254_{exp}/OH_{exp}<1x10^7$ cm/s (excluding good conditions) to define "risky" conditions. Conditions with higher $F185_{exp}/OH_{exp}$ or $F254_{exp}/OH_{exp}$ are defined as "bad". Under good conditions, photolysis of most VOCs has a relative contribution <20% to their fate; under bad conditions, non-tropospheric photolysis is likely to be significant in all OFR experiments, since it can hardly be avoided for oxidation intermediates, even if the precursor(s) does not photolyze at all. Under risky conditions, some species photolyzing slowly and/or reacting with OH rapidly (e.g., alkanes, aldehydes, and most biogenics) still

have a relative contribution of photolysis <20% to their fates, while species photolyzing more rapidly and/or reacting with OH more slowly (e.g., aromatics and other highly conjugated species and some saturated carbonyls) will undergo substantial non-tropospheric photolysis. Note that these definitions are slightly different than in Peng et al. (2016). All definitions of the types of conditions are summarized in Table 3.

2.3 Uncertainty analysis

We apply the same method as in Peng et al. (2014, 2015) to calculate and analyze the output uncertainties due to uncertain kinetic parameters in the model. Random samples following log-normal distributions are generated for all rate constants and photoabsorption cross sections in the model using uncertainty data available in the JPL database (Sander et al., 2011) or estimated based on IUPAC data (Ammann et al., 2016). Then, Monte Carlo Uncertainty Propagation (BIPM et al., 2008) is performed for these samples through the model to obtain the distributions of outputs. Finally, we compute squared correlation coefficients between corresponding input and output samples and apportion the relative contributions of individual kinetic parameters to the output uncertainties based on these coefficients (Saltelli et al., 2005).

3 Results and discussion

In this section, we study the NO_y chemistry in OFR while considering relevant experimental issues. Based on these results, we propose some guidelines for OFR operation for high-NO OH oxidation of VOCs.

3.1 NO_y chemistry in typical OFR cases with initial NO injection

NO was thought to be unimportant (i.e., unable to significantly react with RO_2) in OFRs with initial NO injection (OFR-iNO) based on the argument that its lifetime is too short due to large amounts of O_3 OH, and HO_2 to compete with RO_2+HO_2 (Li et al., 2015). We evaluate this issue below by calculating NO effective lifetime (τ_{NO}, in s), defined as NO exposure (NO_{exp}, in molecules cm^{-3} s) divided by initial NO concentration, under various conditions. This definition cannot effectively capture the true NO average lifetime if it is close to or longer than the residence time. In this case, τ_{NO} close to the residence time will be obtained, which is still long enough for our characterization purposes.

3.1.1 OFR185-iNO

In OFR185-iNO, NO is *not* oxidized extremely quickly under *all* conditions. For instance, under a typical condition in the midrange of the phase space shown in Fig. 1a, τ_{NO} ~13 s. This lifetime is much shorter than the residence time, but long enough for OH_{exp} to reach ~$3x10^{10}$ molecules cm^{-3} s, which is equivalent to an OH equivalent age of ~6 hrs. Such an OH equivalent age is already sufficient to allow some VOC processing and even SOA formation to occur (Lambe et al., 2011; Ortega et al., 2016). Within τ_{NO}, NO suppresses HO_2 through the reaction NO+HO_2→NO_2+OH, leading to NO_{exp}/HO_{2exp} of ~700 during this period, high enough for RO_2 to dominantly react with NO. Meanwhile, NO+HO_2→NO_2+OH enhances OH production, which helps OH_{exp} build up in a relatively short period. In addition, non-tropospheric photolysis of VOCs at 185 and 254 nm is minor ($F185_{exp}$/OH_{exp} ~ 600 cm/s, Fig. 1a), because of enhanced OH production and moderate UV. Therefore, such an OFR condition may be of some interest for high-

NO VOC oxidation. We thus analyze the NO_y chemistry in OFR185-iNO in more detail below, by taking the case shown in Fig. 1a as a representative example.

In OFR185-iNO, HO_x concentrations are orders-of-magnitude higher than in the atmosphere while the amount of O_3 produced is relatively small during the first several seconds after the flow enters the reactor. As a result, NO is not oxidized almost exclusively by O_3 as in the troposphere, but also by OH and HO_2 to form HONO and NO_2, respectively (Fig. 1a). The large concentration of OH present then oxidizes HONO to NO_2, and NO_2 to HNO_3. Photolysis only plays a negligible role in the fate of HONO and NO_2 in OFRs, in contrast to the troposphere, where it is the main fate of these species. This is because the reactions of HONO and NO_2 with OH are greatly accelerated in OFR compared to those in the troposphere, while photolysis not (Peng et al., 2016). The interconversion between NO_2 and HO_2NO_2 is also greatly accelerated (Fig. 1a), since a large amount of HO_2 promotes the formation of HO_2NO_2, whose thermal decomposition and reaction with OH in turn enhance the recycling of NO_2. Though not explicitly modeled in this study, RO_2 are expected to undergo similar reactions with NO_2 to form reservoir species, i.e., peroxynitrates (Orlando and Tyndall, 2012). Peroxynitrates that decompose on timescales considerably longer than OFR residence times may serve as effectively permanent NO_y sinks in OFRs (see Section 3.4.1).

Interestingly but not surprisingly, the NO_y chemistry shown in Fig. 1a is far from temporally uniform during the OFR residence time (Fig. S1a). Within τ_{NO}, NO undergoes an e-fold decay as it is rapidly converted into NO_2 and HONO, whose concentrations reach maxima around that time. After most NO is consumed, HONO and NO_2 also start to decrease, but significantly more slowly than NO, since they do not have as many and efficient loss pathways as NO. The reaction of OH with HONO, the dominant fate of HONO, is slower than that with NO (Fig. 1a). The net rate of the NO_2-to-HO_2NO_2 conversion becomes low because of the relatively fast reverse reaction (Fig. 1a). Besides, the total loss of NO_2 is partially offset by the production from HONO. The generally stable concentrations of HONO and NO_2 (Fig. S1a) result in their respective reaction rates with OH that are comparable during and after τ_{NO} (Fig. 1a), as OH variation is also relatively small during the entire residence time (Fig. S1b). However, the NO_2-to-HO_2NO_2 conversion after τ_{NO} is much faster than during it (Fig. 1a), resulting from substantially decreased NO and HO_2 concomitantly increasing >1 order of magnitude after τ_{NO} (Fig. S1a,b). HNO_3 and HO_2NO_2, which are substantially produced only after NO_2 is built up, have much higher concentrations later than within τ_{NO}.

Under other OFR185-iNO conditions than in Fig. 1a, the major reactions interconverting NO_y species are generally the same, although their relative importance may vary. At lower NO^{in}, the perturbation of HO_x chemistry caused by NO_y species is smaller. Effects of NO^{in} less than 1 ppb (e.g., typical non-urban ambient concentrations) are generally negligible regarding HO_x chemistry. Regarding NO_y species, the pathways in Fig. 1a are still important under those conditions. At higher NO^{in} (e.g., >1 ppm), one might expect NO_3 and N_2O_5 to play a role (as in OFR254-iNO; see Section 3.1.2 below), since high NO_y concentrations might enhance self/cross reactions of NO_y. However, this would not occur unless OH production is high, since relatively low O_3 concentrations in OFR185-iNO cannot oxidize NO_2

297 to NO_3 rapidly. Also, a large amount of NO_y can lead to significant OH suppression. That would in turn
298 slow down the NO_3 production from HNO_3 by OH. This is especially true when an OFR is used to oxidize
299 the output of highly concentrated sources (e.g., from vehicle exhausts). When sources corresponding
300 to OHR_{ext} of thousands of s^{-1} and NO^{in} of tens of ppm are injected into OFR185 (Fig. 1b), they essentially
301 inhibit active chemistry except NO consumption, as all subsequent products are much less abundant
302 compared to remaining NO (Fig. S1c).

303 **3.1.2 OFR254-iNO**

304 The ppm-level $O_{3,in}$ used in the OFR254-iNO mode of operation has a strong impact on its NO_y
305 chemistry. An $O_{3,in}$ of 2.2 ppm (lowest in this study) is already enough to shorten τ_{NO} to ~1 s, preventing
306 NO from playing a role in the chemistry under most explored conditions. The reaction fluxes under a
307 typical $O_{3,in}$ of 7 ppm are shown in Fig. 1c. A reactive flux from $NO+O_3 \rightarrow NO_2$ makes the reaction of NO
308 with other oxidants (OH, HO_2 etc.) negligible. The HNO_3 production pathway from NO_2 is similar to that
309 in OFR185-iNO. The interconversion between NO_2 and HO_2NO_2 is also fast over the residence time, and
310 even faster than in OFR185-iNO during τ_{NO}, since a high concentration of O_3 also controls the OH-HO_2
311 interconversion and makes HO_2 more resilient against suppression due to high NO (Fig. S1f; Peng et al.,
312 2015). A major difference in the NO_y chemistry in OFR254-iNO (Fig. 1c) compared to OFR185-iNO (Fig.
313 1a) is significant NO_3/N_2O_5 chemistry due to high O_3 in OFR254-iNO, which accelerates the oxidation of
314 NO_2 to NO_3. Interconversion between NO_2+NO_3 and N_2O_5 also occurs to a significant extent because of
315 high NO_2. Under the conditions of Fig. 1c, NO_3 can also be significantly consumed by HO_2. Unlike
316 OFR185-iNO, OFR254-iNO can substantially form NO_3 from HNO_3 under conditions that are not on the
317 extremes of the explored physical condition space, e.g., at higher UV and lower NO^{in} (e.g., Fig. S2). In
318 the case of very high NO^{in} (equal to or higher than $O_{3,in}$), all O_3 can be rapidly destroyed by NO. As a
319 consequence, OH production is shut down and these cases are of little practical interest (Fig. S3h).

320 **3.1.3 Uncertainty analysis**

321 The results of uncertainty propagation confirm that the output uncertainties due to uncertain
322 kinetic parameters are relatively low compared to other factors (e.g., non-plug flow in OFR; Peng et al.,
323 2015) and the overall model accuracy compared to experimental data (a factor of 2–3; Li et al., 2015).
324 For OFR185-iNO, NO, NO_3, and OH exposures have relative uncertainties of ~0–20%, ~40–70%, and ~15–
325 40%, respectively. The uncertainties in OH exposure are very similar to those in the cases without NO_x
326 (Peng et al., 2015). The contribution of NO_y reactions to OH_{exp} uncertainty is negligible, except for some
327 contribution of $OH+NO \rightarrow HONO$ in a few cases with high NO^{in} (Fig. 2). The uncertainties on NO_{exp} are
328 dominated by the reactions producing HO_x and O_3, i.e., the major consumers of NO. For NO_3 exposure,
329 a few major production and loss pathways (e.g., $NO_2+NO_3 \rightarrow N_2O_5$, $N_2O_5 \rightarrow NO_2+NO_3$, and
330 $HO_2+NO_3 \rightarrow OH+NO_2+O_2$) dominate its uncertainties. OFR254-iNO has a simpler picture of parametric
331 uncertainties in terms of composition. O_3 controls the NO oxidation under most conditions and this
332 reaction contributes most of output uncertainties for NO exposures. $HO_2+NO_3 \rightarrow OH+NO_2+O_2$ dominates
333 the uncertainty on NO_3 exposure. The levels of those uncertainties are lower than in OFR185-iNO (<2%
334 for NO exposure; <60% in all cases and <25% in most cases for NO_3 exposure). Thus, model uncertainties

in OFR254-iNO are not shown in detail.

3.2 Different conditions types

Having illustrated the main NO_y chemical pathways for typical cases, we present the results of the exploration of the entire physical parameter space (see Section 2.2). Note that the explored space is indeed very large and gridded logarithmically uniformly in every dimension. Therefore, the statistics of the exploration results can be useful to determine the relative importance of the conditions types defined in Section 2.2 and Table 3.

It has been shown that during τ_{NO}, RO_2 can react dominantly with NO (Section 3.1.1), while to determine if a condition is high-NO (see Table 3), the entire residence time is considered. This is done because for VOC oxidation systems of interest, there will be significant oxidation of the initial VOC and its products under low-NO conditions, if τ_{NO} is shorter than the reactor residence time. After most NO is consumed, the longer the remaining residence time, the more RO_2 will react with HO_2 and the more likely that an input condition is classified as low-NO. For a condition to be high-NO, a significantly long τ_{NO} is required. Figure 3 shows the fractional occurrence distribution of good/risky/bad conditions in the entire explored condition space over logarithm of $r(RO_2+NO)/r(RO_2+HO_2)$, which distinguishes high- and low-NO conditions. In OFR254-iNO, τ_{NO} is so short that no good high-NO condition is found in the explored range in this study (Fig. 3a). A fraction of explored conditions are bad high-NO. These conditions result from a full consumption of O_3 by NO. Then very little HO_x is produced (right panels in Fig. S3h), but the fate of any RO_2 formed is dominated by RO_2+NO (right panels in Fig. S3i). However, also due to negligibly low OH concentration, little RO_2 is produced and non-tropospheric photolysis of VOCs is also substantial compared to their reaction with OH under these conditions, classifying all of them as “bad” (Fig. 3a).

In OFR185-iNO, in addition to the typical case shown in Fig. 1a, many other cases have a τ_{NO} of ~10 s or longer (Figs. S3b and S4), which allow the possibility of high-NO conditions. Indeed, ~1/3 of explored conditions in OFR185-iNO with a residence time of 3 min are high-NO (Fig. 3b). Most of these high-NO conditions are also classified as bad, similar with those in OFR254-iNO. More importantly, in contrast to OFR254-iNO, good and risky high-NO conditions also comprise an appreciable fraction of the OFR185-iNO conditions. It is easily expected that very high OHR_{ext} and NO^{in} lead to bad high-NO conditions (all panels in Fig. 4), since they strongly suppress HO_x, which yields bad conditions and in turn keep NO destruction relatively low. Besides, the occurrence of bad high-NO conditions is reduced at high UV (bottom panels in Fig. 4), which can be explained by lowered NO due to high O_3 production and fast OH reactant loss due to high OH production. Good high-NO conditions are rare in the explored space. They are only 1.1% of total explored conditions (Fig. 3b) and present under very specific conditions, i.e., higher H_2O, lower UV, lower OHR_{ext}, and NO^{in} of tens to hundreds of ppb (Figs. 4 and S5). Since a very high NO can suppress OH, to obtain both a significant NO level and a good conditions, NO^{in} can only be tens to hundreds of ppb. As NO^{in} is lower and OH is higher than under bad high-NO conditions, UV should be lower than bad high-NO conditions to keep a sufficiently long presence of NO. Thus, UV at 185 nm for good high-NO conditions are generally lower than 10^{12} photons cm^{-2} s^{-1} (Fig. S5).

In addition, a low OHR_{ext} (generally <50 s^{-1}) and a higher H_2O (the higher the better, although there is no apparent threshold) are also required for good high-NO conditions (Fig. S5), as Peng et al. (2016) pointed out. Risky high-NO conditions often occur between good and bad high-NO conditions, e.g., at lower NO^{in} than bad conditions (e.g., Cases ML, MM, HL, and HM in Fig. 4, see Table 2 for the typical case label code), at higher OHR_{ext} and/or NO_{in} than good conditions (e.g., Cases ML and MM), and at lower H_2O than good conditions (e.g., Case LL).

The trend of the distributions of good, risky, and bad low-NO conditions is generally in line with the analysis in Peng et al. (2016). For low-NO conditions, NO_y species can be simply regarded as external OH reactants, as in Peng et al. (2016). As H_2O decreases and/or OHR_{ext} or NO^{in} increases, a low-NO condition becomes worse (good→risky→bad) (Figs. 4 and 5). In OFR185-iNO, increasing UV generally makes a low-NO condition better because of an OH production enhancement (Fig. 4); while in OFR254-iNO, increasing UV generally makes a low-NO condition worse (Fig. 5), since at a higher UV, more O_3 is destroyed and the resilience of OH to suppression is reduced.

As discussed above, the fraction of high-NO conditions also depends on OFR residence time. A shorter residence time is expected to generally lead to a larger fraction of high-NO conditions, since the time spent in the reaction for $t > \tau_{NO}$ is significantly smaller. Thus, we also investigate an OFR185-iNO case with a residence time of 30 s. In Fig. 3b, compared to the case with a residence time of 3 min, the distributions of all condition types (good/risky/bad) of the 30 s residence time case shift toward higher $r(RO_2+NO)/r(RO_2+HO_2)$. Nevertheless, shortening the residence time also removes the period when the condition is better (i.e., less non-tropospheric photolysis), when external OH reactants have been partially consumed and OH suppression due to OHR_{ext} has been reduced later in the residence time. As a result, the fractions of good and risky conditions decrease (Fig. 3b). With the two effects (higher $r(RO_2+NO)/r(RO_2+HO_2)$ and more significant non-tropospheric photolysis) combined, the fraction of good high-NO conditions increases by a factor of ~3. An even shorter residence time does not result in a larger good high-NO fraction, since the effect of enhancing non-tropospheric photolysis is even more apparent.

3.3 Effect of non-plug flow

We performed model runs where the only change with respect to our box model introduced in Section 2.2 is that the plug-flow assumption is replaced by the residence time distribution (RTD) measured by Lambe et al. (2011) (also see Fig. S8 of Peng et al. (2015)). The chemistry of different air parcels with different residence times is simulated by our box model and outputs are averaged over the RTD. Lateral diffusion between different air parcels is neglected in these simulations.

OH_{exp} calculated from the mode with RTD ($OH_{exp,RTD}$) is higher than that calculated from the plug-flow model ($OH_{exp,PF}$) in both OFR185-iNO and OFR254-iNO (Table 4 and Fig. S6). Under most explored conditions deviations are relatively small, which leads to an overall positive deviation of $OH_{exp,RTD}$ from $OH_{exp,PF}$ by ~x2 (within the uncertainties of the model and its application to real experimental systems). For OFR185-iNO, most conditions (~90%) in the explored space lead to <x3 differences between $OH_{exp,PF}$ and $OH_{exp,RTD}$, while for a small fraction of cases the differences can be larger (Fig. S6). The larger

deviations are mainly present at high UV, OHR_{ext}, and NO^{in}, where conditions are generally "bad" and in which experiments are of little atmospheric relevance. Under these specific conditions, external OH reactants and NO_y can be substantially destroyed for the air parcels with residence times longer than the average, while this is not the case for the average residence time. This feature was already described by Peng et al. (2015) (see Fig. S10 of that study). Although only non-NO_y external OH reactants were considered in that study, the results are the same. In the present study, a higher upper limit of the explored OHR_{ext} range (compared to Peng et al., 2015, due to trying to simulate extremely high OHR_{ext} used in some recent literature studies) large amounts of NO_y and cause somewhat larger deviations. In OFR254-iNO, OH is less suppressed at high OHR_{ext} and NO^{in} than in OFR185-iNO because of high O_3 (Peng et al., 2015), $OH_{exp,RTD}$ deviations from $OH_{exp,PF}$ are also smaller (Table 4).

Based on the outputs of the model with RTD, similar mapping of the physical input space as Figs. 4 and 5 can be done (Figs. S7 and S8). Overall, the mapping of the RTD model results is very similar with that of the plug-flow model. The conditions appear to be only slightly better in a few places of the explored space than those from the plug-flow model, which can be easily explained by the discussions above. Besides, the mapping in Figs. S7 and S8 also appear to be slightly more low-NO, for the same reasons discussed above. After NO is destroyed at long residence times, HO_2, suppressed by NO, also recovers as OH. $r(RO_2+NO)/r(RO_2+HO_2)$ is obviously expected to be smaller than in the plug-flow model in general.

Note that most conditions that appear to be better in the RTD model results are already identified as bad by the plug-flow model. Those conditions look slightly better only because of their better *RTD-averaged* $F185_{exp}/OH_{exp}$ and $F254_{exp}/OH_{exp}$. However, each of those cases is actually composed of both a better part at longer residence times and also a worse part at shorter residence times. Under those conditions, the reactor simultaneously works in two distinct regimes, one of which is bad due to heavy OH suppression. Such conditions are obviously not desirable for OFR operation.

3.4 Possible issues related to high-NO_x levels

In the discussion above, we focused on obtaining high-NO conditions and considered only one experimental issue (non-tropospheric photolysis) that had been previously investigated in Peng et al. (2016) and is not specific for experiments with high NO injection. We discuss additional potential reasons why the OFR-iNO chemistry can deviate strongly from tropospheric conditions, as specifically related to high-NO_x level in this subsection.

3.4.1 NO_2

NO_2 reacts with RO_2 to form peroxynitrates, generally regarded as reservoir species in the atmosphere as most of them thermally decompose very quickly compared to atmospheric time scales. However, in OFRs, with residence times on the order of minutes, some peroxynitrates may no longer be considered as fast decomposing. This is especially true for acylperoxy nitrates, whose lifetimes can be hours at room temperature (Orlando and Tyndall, 2012). Acylperoxy nitrates are essentially sinks instead of reservoirs in OFRs for both NO_2 and RO_2. RO_2 is estimated to be as high as several ppb in OFRs by our model (e.g., ~6 ppb RO_2 in OFR185 at H_2O=1%, UV at 185 nm=$1x10^{13}$ photons cm^{-2} s^{-1}, OHR_{ext}=1000 s^{-1},

and NO^{in}=0), while high-NO experiments can yield far higher NO_2. If all RO_2 were acylperoxy, the RO_2 chemistry could be rapidly shut down by NO_2, as rate constants of these $RO_2 + NO_2$ reactions are around 10^{-11} cm^3 $molecule^{-1}$ s^{-1} (Orlando and Tyndall, 2012). Nevertheless, acylperoxy nitrates are not expected to typically be the dominant component of peroxynitrates, since acyl radicals are not a direct oxidation product of most common VOCs and can only be formed after several steps of oxidation (Atkinson and Arey, 2003; Ziemann and Atkinson, 2012). Most alkylperoxy nitrates retain their short-lived reservoir characteristics in OFRs due to their relatively short thermal decomposition time scales (on the order of 0.1 s; Orlando and Tyndall, 2012). Even so, OFR experiments can be seriously hampered at extremely high NO_2. If NO_2 reaches ppm levels, the equilibrium between RO_2+NO_2 and alkylperoxy nitrate (RO_2+NO_2↔RO_2NO_2) is greatly shifted toward the alkylperoxy nitrate side, as the forward and reverse rate constants are on the order of 10^{-12} cm^3 $molecule^{-1}$ s^{-1} and 1 s^{-1}, respectively (Orlando and Tyndall, 2012). This results in a substantial decrease in effective RO_2 concentration, or in other words, a substantial slow-down of RO_2 chemistry.

Parts per million levels of NO_2 may impose an additional experimental artifact in the oxidation chemistry of aromatic precursors. OH-aromatic adducts, i.e., the immediate products of aromatic oxidation by OH, undergo addition of O_2 and NO_2 at comparable rates under ppm levels of NO_2 (rate constants of the additions of O_2 and NO_2 are on the order of 10^{-16} and 10^{-11} molecules cm^{-3} s^{-1}, respectively ;Atkinson and Arey, 2003). However, only the former addition is atmospherically relevant (Calvert et al., 2002). Liu et al. (2015) performed OFR254-iNO experiments with toluene over a range of NO^{in} of 2.5–10 ppm, encompassing the NO concentration range at which the reactions of OH-toluene adduct with O_2 and with NO_2 are of equal importance (~5 ppm; Atkinson and Arey, 2003). This suggests that nitroaromatics, whose formation was reported in the study of Liu et al. (2015), might have been formed in substantial amounts in that study through the addition of NO_2 to the OH-toluene adduct.

3.4.2 NO_3

As discussed in Section 3.1, NO_3 can be formed in significant amounts in OFRs with high NO injection. Although NO_3 is also present in the atmosphere, especially during nighttime, significant VOC oxidation by both OH and NO_3 results in more complex chemistry that may complicate the interpretation of experimental results. NO_3 oxidation-only OFR has been previously realized experimentally via thermal dissociation of injected N_2O_5 (Palm et al., 2017). We discuss below how to avoid significant VOC oxidation by NO_3 and achieve OH-dominated VOC oxidation in OFRs with high NO injection.

If $NO_{3exp}/OH_{exp} > 0.1$, NO_3 can be a competitive reactant for biogenic alkenes and dihydrofurans, which have a C=C bond for NO_3 addition, and phenols, which have activated hydroxyl for fast hydrogen abstraction by NO_3 (Atkinson and Arey, 2003), while for lower NO_{3exp}/OH_{exp}, OH is expected to dominate the oxidation of all VOCs, as shown in Fig. 6. Oxidation for VOCs without alkene C=C bonds and phenol hydroxyl (such as alkanes and (alkyl)benzenes) is dominated by OH unless $NO_{3exp}/OH_{exp} > 1000$. Despite its double bond, ethene reacts as slowly with NO_3 as alkanes, likely due to lack of alkyl groups enriching electron density on the C=C bond, which slows NO_3 addition. We calculate NO_{3exp}/OH_{exp} for OFR185-

iNO and OFR254-iNO and plot histograms of this ratio in Fig. 6. Many experimental conditions lead to high enough NO_{3exp}/OH_{exp} that NO_3 is a competitive sink for alkenes, while only under very extreme conditions can NO_3 be a competitive sink for species without C=C bonds. High-NO conditions in OFR185-iNO have lower NO_{3exp}/OH_{exp} (~10^{-2}–10^{2}) than in OFR254-iNO (~10^{1}–10^{5}) (Figs. 6 and S3d,g,j). This difference in NO_{3exp}/OH_{exp} is due to the different levels of O_3 in the two modes, as high O_3 promotes NO_2-to-NO_3 oxidation. Note that low-NO conditions in both OFR185-iNO and OFR254-iNO can also reach high NO_{3exp}/OH_{exp} as some high-NO conditions have. This is because in OFR185-iNO a large part of NO_3 is formed by OH oxidation, resulting in NO_{3exp}/OH_{exp} being largely influenced by NO^{in} but not by other factors mainly governing OH (Fig. S3d); and under low-NO conditions in OFR254-iNO, NO_3 can form rapidly from NO_2+O_3, while OH can be heavily suppressed by high OHR_{ext} (Fig. S3g,j).

Most of the species shown in Fig. 6 are primary VOCs, except phenols and a dihydrofuran, which can be intermediates of the atmospheric oxidation of (alkyl)benzenes (Atkinson and Arey, 2003) and long-chain alkanes (Aimanant and Ziemann, 2013; Strollo and Ziemann, 2013; Ranney and Ziemann, 2016), respectively. Nevertheless, only the phenol production may occur in high-NO OFRs, as the particle-phase reaction in the photochemical formation of dihydrofurans from alkanes is too slow compared to typical OFR residence times (Ranney and Ziemann, 2016). Therefore, the impact of NO_3 oxidation on VOC fate needs to be considered only if the OFR input flow contains high NO mixed with biogenics and/or aromatics [(alkyl)benzenes and/or phenols]. However, (alkyl)benzenes were likely to be major SOA precursors in, to our knowledge, the only few literature OFR studies with high NO levels (Ortega et al., 2013; Tkacik et al., 2014; Liu et al., 2015). In the study of the air in a traffic tunnel (OFR185-iNO mode; Tkacik et al., 2014), where toluene is usually a major anthropogenic SOA precursor as in other urban environments (Dzepina et al., 2009; Borbon et al., 2013; Hayes et al., 2015; Jathar et al., 2015), NO_x was several hundreds of ppb. This resulted in an estimated NO_{3exp}/OH_{exp} range of ~0.1–1, where up to ~30% of cresols (intermediates of toluene oxidation) may have been consumed by NO_3. Dihydrofurans may also have formed in the tunnel air (but outside the OFR) in the presence of NO_x (Aimanant and Ziemann, 2013; Strollo and Ziemann, 2013) and, after entering the OFR, they would have been substantially (up to ~50%) consumed by NO_3. In the laboratory experiment of Liu et al. (2015) with toluene, the injection of as much as 10 ppm NO elevated NO_{3exp}/OH_{exp} to ~100, where cresols from toluene oxidation reacted almost exclusively with NO_3 in addition to being photolyzed.

3.4.3 A case study

We use a case study of an OFR254-13-iNO laboratory experiment with a large amount of toluene (5 ppm) and NO^{in} (10 ppm) to illustrate how very high VOC and NO concentrations cause multiple types of atmospherically irrelevant reactions in OFR. Due to very high OHR_{ext} and NO^{in}, photolysis of toluene at 254 nm may have been important (Peng et al., 2016). In case of a high (close to 1) quantum yield, up to ~80% of the consumed toluene in their experiments could have been photolyzed (Scheme 1). Of the rest of reacted toluene, ~10% undergoes H-abstraction by OH from the methyl group in the model, leading to an RO_2 similar to alkyl RO_2 and likely proceeding with normal RO_2 chemistry. ~90% of the toluene formed an OH-adduct (Calvert et al., 2002). As discussed above, 70% of this adduct (depending

on NO^{in}) is predicted to recombine with NO_2 producing nitroaromatics because of the ppm-level NO_x. The adduct could also react with O_2 via two types of pathways, of which one was addition forming a special category of RO_2 (OH-toluene-O_2 adducts) potentially undergoing ring-opening (Atkinson and Arey, 2003; Orlando and Tyndall, 2012; Ziemann and Atkinson, 2012), the other H-elimination by O_2 producing cresols. Again, like toluene, cresols may have been substantially photolyzed. As a result of NO_{3exp}/OH_{exp} ~100, only a minor portion of cresol could have undergone OH addition and then H-elimination again. This pathway leads to the formation of methyldihydroxybenzenes and other OH-oxidation products (Atkinson and Arey, 2003). The rest of cresols may have formed methylphenoxy radicals, nevertheless, dominantly via H-abstraction by NO_3, since H-abstraction by OH was even a minor pathway compared to the OH-addition one (Atkinson et al., 1992). In summary, the model results suggest that there were two possible routes leading to nitroaromatic formation. However, one of them (recombination of OH-aromatic adducts with NO_2) is likely of little atmospheric relevance due to very high NO_x needed, and the other (H-abstraction from cresol) occurs in the atmosphere but is not a major fate of aromatics (Calvert et al., 2002).

3.5 Implications for OFR experiments with combustion emissions as input

Emissions from combustion sources, e.g., vehicles and biomass burning, usually contain VOCs and NO_x at very high concentrations (Table 1). An injection of this type of emissions (typically with OHR_{ext} of thousands of s^{-1} or larger and NO^{in} of tens of ppm or larger) in OFRs without any pretreatment is likely to cause all experimental issues discussed in Peng et al. (2016) and this paper, i.e., strong OH suppression, substantial non-tropospheric photolysis, strong RO_2 suppression by NO_2 whether RO_2 is acyl RO_2 or not, fast reactions of NO_2 with OH-aromatic hydrocarbon adducts, substantial NO_3 contribution to VOC fate, and even a near-total inhibition of OFR chemistry due to complete titration of O_3 by NO in the case of OFR254. We take the study of Karjalainen et al. (2016), who used an OFR to oxidize diluted car exhaust in real-time, as an case study to investigate the extent to which these issues may affect typical combustion source studies and to explore approaches to mitigate the problems.

During the first 200 s of their experiment (defined as the “cold start” period when the catalyst is cold and emissions are high), NO and total hydrocarbon in the emissions of the test vehicle reached ~400 and ~600 ppm, respectively. We first simulate the oxidation of those emissions without any dilution (even though x12 dilution was used in their experiments) to explore the most extreme conditions. Our model simulation indicates that such an extremely concentrated source would generally lead to bad high- or low-NO conditions (depending on NO concentration) in their OFR (Fig. 7), even though it was run at relatively high H_2O and UV. OH suppression can be as high as 3 orders of magnitude; VOC fates by non-tropospheric photolysis and reactions of alkenes and phenols with NO_3 can be nearly 100%; up to ~1/3 of OH-toluene adduct may be recombined with NO_2 instead of forming an adduct with O_2. After the test vehicle entered the “hot stabilized” stage (200–1000 s), its VOC emissions (on the order of ppm) were still too high for an undiluted OFR to yield a good condition (Fig. S9). OH suppression can still reach 2 orders of magnitude; non-tropospheric photolysis, and sometimes reactions with NO_3, can still dominate over reactions with OH in VOC fates; reactions of OH-toluene adduct with NO_2 can

still be substantial at some small NO emission spikes. Moreover, although NO emissions were roughly at ppm level even during the hot stabilized period, NO effective lifetime may be very short during that period, leading to low-NO conditions in their OFR.

As suggested in Peng et al. (2016) for low-NO OFR, dilution of sources can also mitigate strong deviations on OFR-iNO chemistry vs. atmospherically-relevant conditions. A dilution by a factor of 12, as actually used by Karjalainen et al. (2016), appears to be sufficient to bring most of the hot stabilized period under good conditions (Fig. S9). However, most VOC, or in other words, most SOA formation potential, was emitted during the cold start period, when risky and bad conditions still prevailed (Figs. 7 and 8). Even if the emissions are diluted by x100, the cold-start emission peak (Fig. 7) is still under risky conditions. Although bad conditions are eliminated and good condition is present during most of time, this emission peak under risky condition may contribute >50% to total SOA formation potential (Fig. 8). For SOA formed under good condition to be dominant, a dilution factor >400 would be needed. Note that a strong dilution lowers aerosol mass loading in vehicle emissions. As a result, condensation of gases onto particles is slower than in raw exhausts. However, condensational sinks after dilution may still be significantly higher than typical ambient values (Matti Maricq, 2007; Donahue et al., 2016).

Note that the emissions of the test vehicle of Karjalainen et al. (2016) are rather clean compared to the typical 2013 US on-road fleet (i.e., all at the hot stabilized stage) measured by Bishop and Stedman (2013) (Figs. 9 and S10). For emissions of an average on-road fleet, a dilution by a factor of 100 or larger would be necessary to ensure that most emissions would be processed in OFR185 under good conditions at the highest H_2O and UV in this study (Figs. 9b and S10b,e,h). In the case of lower H_2O and/or UV, an even larger dilution factor would be required.

Conducting OFR185-iNO experiments at high UV lowers the dilution factor needed for good conditions. However, it also renders good high-NO condition impossible (see Section 3.2 and Fig. S4). If one wants to oxidize vehicle exhausts in a high-NO environment in OFR, as in an urban atmosphere, OFR185 at low UV is necessary. Consequently, a much stronger dilution is in turn necessary to keep the operation condition still good. Nevertheless, not all vehicle emissions can be moved into good high-NO region through a simple dilution (Figs. 9c and S10c,f,i). Furthermore, a low UV would seriously limit the highest OH_{exp} that OFR can achieve (~3×10^{11} molecules cm^{-3} s for modeled good high-NO conditions in this study), while a much higher OH_{exp} would be desirable to fully convert SOA formation potential into measurable SOA mass. If both good high-NO condition and high OH_{exp} are required, new techniques (e.g., injection of N_2O at percent level proposed by Lambe et al. (2017)) may be necessary.

4 Conclusions

In this study, OFR chemistry involving NO_y species was systematically investigated over a wide range of conditions. NO initially injected into the OFR was found to be rapidly oxidized under most conditions. In particular, due to high O_3 concentrations, NO lifetime in OFR254-iNO was too short to result in a significant RO_2 consumption by NO compared to that by HO_2 under all conditions with active chemistry. Nevertheless, it is not completely impossible for OFR185-iNO to have a significant RO_2 fate by NO and minor non-tropospheric photolysis at the same time ("good high-NO conditions"). According

to our simulations, these conditions are most likely present at high H_2O, low UV, low OHR_{ext}, and NO^{in} of tens to hundreds of ppb.

However, many past OFR studies with high NO injection were conducted under conditions remarkably different from the abovementioned very narrow range. NO^{in} and/or OHR_{ext} in those studies were often much higher than good high-NO conditions require (particularly, >3 orders of magnitude in some OFR studies using combustion emissions as input). In addition to non-tropospheric organic photolysis, OFR oxidation of highly concentrated sources can cause multiple large deviations from tropospheric OH oxidation, i.e., RO_2 suppression by high NO_2, substantial nitroaromatic formation from the recombination of NO_2 and OH-aromatic adducts, and fast reactions of VOCs with NO_3 compared to those with OH.

Working at lower NO_x (sub-ppm level) and VOC concentrations or dilution can mitigate these experimental problems. In general, a strong dilution (by a factor of >100) is needed for OFR that process typical on-road vehicle emissions. Humidification can also make good conditions more likely. By these measures, good conditions can be guaranteed, as long as NO and/or precursor concentrations are sufficiently low, while high-NO conditions cannot be ensured. To aid design and interpretation of OFR experiments with high NO injection, we provide our detailed modeling results in a visualized form (Fig. S3). For OFR users in need for both high OH_{exp} and high NO, simple NO injection is not a good option. New techniques (e.g., injection of N_2O proposed by Lambe et al. (2017) or other innovations) may be necessary to meet this need.

Acknowledgements

This work was partially supported by DOE (BER/ASR) DE-SC0011105 & DE-SC0016559, EPA STAR 83587701-0, and NSF AGS-1360834. We thank Pengfei Liu, Andrew Lambe, and Daniel Tkacik for providing some OFR experimental data, the authors of Karjalainen et al. (2016) and their project IEA-AMF Annex 44 for providing the data and information for the vehicle tests, Gary Bishop for providing on-road vehicle emission data, and Andrew Lambe and William Brune for useful discussions.

References

Aimanant, S. and Ziemann, P. J.: Chemical Mechanisms of Aging of Aerosol Formed from the Reaction of n-Pentadecane with OH Radicals in the Presence of NOx, Aerosol Sci. Technol., 47(9), 979–990, doi:10.1080/02786826.2013.804621, 2013.

Alanen, J., Simonen, P., Saarikoski, S., Timonen, H., Kangasniemi, O., Saukko, E., Hillamo, R., Lehtoranta, K., Murtonen, T., Vesala, H., Keskinen, J. and Rönkkö, T.: Comparison of primary and secondary particle formation from natural gas engine exhaust and of their volatility characteristics, Atmos. Chem. Phys. Discuss., (February), 1–27, doi:10.5194/acp-2017-44, 2017.

Ammann, M., Cox, R. A., Crowley, J. N., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., Wallington, T. J., Cox, B., Atkinson, R., Baulch, D. L. and Kerr, J. A.: IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, [online] Available from: http://iupac.pole-ether.fr/#, 2016.

Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds., Chem. Rev., 103(12), 4605–38, doi:10.1021/cr0206420, 2003.

Atkinson, R., Aschmann, S. M. and Arey, J.: Reactions of hydroxyl and nitrogen trioxide radicals with phenol, cresols, and 2-nitrophenol at 296 .+-. 2 K, Environ. Sci. Technol., 26(7), 1397–1403, doi:10.1021/es00031a018, 1992.

BIPM, IEC, IFCC, ILAC, ISO, IUPAC and IUPAPOIML: JCGM 101: 2008 Evaluation of measurement data — Supplement 1 to the " Guide to the expression of uncertainty in measurement " — Propagation of distributions using a Monte Carlo method., 2008.

Bishop, G. A. and Stedman, D. H.: Fuel Efficiency Automobile Test: Light-Duty Vehicles, [online] Available from: http://www.feat.biochem.du.edu/light_duty_vehicles.html (Accessed 1 February 2017), 2013.

Borbon, A., Gilman, J. B., Kuster, W. C., Grand, N., Chevaillier, S., Colomb, A., Dolgorouky, C., Gros, V., Lopez, M., Sarda-Esteve, R., Holloway, J., Stutz, J., Petetin, H., McKeen, S., Beekmann, M., Warneke, C., Parrish, D. D. and De Gouw, J. A.: Emission ratios of anthropogenic volatile organic compounds in northern mid-latitude megacities: Observations versus emission inventories in Los Angeles and Paris, J. Geophys. Res. Atmos., 118(4), 2041–2057, doi:10.1002/jgrd.50059, 2013.

Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Wilmouth, D. M. and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies: Evaluation Number 18, Pasadena, CA, USA. [online] Available from: http://jpldataeval.jpl.nasa.gov/, 2015.

Calvert, J. G., Atkinson, R., Becker, K. H., Kamens, R. M., Seinfeld, J. H., Wallington, T. H. and Yarwood, G.: The Mechanisms of Atmospheric Oxidation of the Aromatic Hydrocarbons, Oxford University Press, USA. [online] Available from: https://books.google.com/books?id=P0basaLrxDMC, 2002.

Carlton, A. G., Wiedinmyer, C. and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, Atmos. Chem. Phys., 9(14), 4987–5005, doi:10.5194/acp-9-4987-2009, 2009.

Carter, W. P. L., Cocker, D. R., Fitz, D. R., Malkina, I. L., Bumiller, K., Sauer, C. G., Pisano, J. T., Bufalino, C. and Song, C.: A new environmental chamber for evaluation of gas-phase chemical mechanisms and secondary aerosol formation, Atmos. Environ., 39(40), 7768–7788, doi:10.1016/j.atmosenv.2005.08.040, 2005.

Chameides, W., Lindsay, R., Richardson, J. and Kiang, C.: The role of biogenic hydrocarbons in urban photochemical smog: Atlanta as a case study, Science (80-.)., 241(4872), 1473–1475, doi:10.1126/science.3420404, 1988.

Cocker, D. R., Flagan, R. C. and Seinfeld, J. H.: State-of-the-Art Chamber Facility for Studying Atmospheric Aerosol Chemistry, Environ. Sci. Technol., 35(12), 2594–2601, doi:10.1021/es0019169, 2001.

Donahue, N. M., Posner, L. N., Westervelt, D. M., Li, Z., Shrivastava, M., Presto, A. A., Sullivan, R. C., Adams, P. J., Pandis, S. N. and Robinson, A. L.: Where Did This Particle Come From? Sources of Particle Number and Mass for Human Exposure Estimates, in Airborne Particulate Matter: Sources, Atmospheric Processes and Health, edited by R. M. Harrison, R. E. Hester, and X. Querol, pp. 35–71, Royal Society of Chemistry., 2016.

Dzepina, K., Volkamer, R. M., Madronich, S., Tulet, P., Ulbrich, I. M., Zhang, Q., Cappa, C. D., Ziemann, P.

J. and Jimenez, J. L.: Evaluation of recently-proposed secondary organic aerosol models for a case study in Mexico City, Atmos. Chem. Phys., 9(15), 5681–5709, doi:10.5194/acp-9-5681-2009, 2009.

George, I. J., Vlasenko, A., Slowik, J. G., Broekhuizen, K. and Abbatt, J. P. D.: Heterogeneous oxidation of saturated organic aerosols by hydroxyl radicals: uptake kinetics, condensed-phase products, and particle size change, Atmos. Chem. Phys., 7(16), 4187–4201, doi:10.5194/acp-7-4187-2007, 2007.

Haagen-Smit, A. J.: Chemistry and Physiology of Los Angeles Smog, Ind. Eng. Chem., 44(6), 1342–1346, doi:10.1021/ie50510a045, 1952.

Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prevot, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R. and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, Atmos. Chem. Phys., 9(14), 5155–5236, 2009.

Hayes, P. L., Carlton, a. G., Baker, K. R., Ahmadov, R., Washenfelder, R. a., Alvarez, S., Rappenglück, B., Gilman, J. B., Kuster, W. C., de Gouw, J. a., Zotter, P., Prévôt, a. S. H., Szidat, S., Kleindienst, T. E., Offenberg, J. H., Ma, P. K. and Jimenez, J. L.: Modeling the formation and aging of secondary organic aerosols in Los Angeles during CalNex 2010, Atmos. Chem. Phys., 15(10), 5773–5801, doi:10.5194/acp-15-5773-2015, 2015.

Hearn, J. D. and Smith, G. D.: Kinetics and Product Studies for Ozonolysis Reactions of Organic Particles Using Aerosol CIMS †, J. Phys. Chem. A, 108(45), 10019–10029, doi:10.1021/jp0404145, 2004.

Hoffmann, T., Odum, J. R., Bowman, F., Collins, D., Klockow, D., Flagan, R. C. and Seinfeld., J. H.: Formation of Organic Aerosols from the Oxidation of Biogenic Hydrocarbons, J. Atmos. Chem., 26(2), 189–222, doi:10.1023/A:1005734301837, 1997.

Hu, W., Palm, B. B., Day, D. A., Campuzano-Jost, P., Krechmer, J. E., Peng, Z., de Sá, S. S., Martin, S. T., Alexander, M. L., Baumann, K., Hacker, L., Kiendler-Scharr, A., Koss, A. R., de Gouw, J. A., Goldstein, A. H., Seco, R., Sjostedt, S. J., Park, J.-H., Guenther, A. B., Kim, S., Canonaco, F., Prévôt, A. S. H., Brune, W. H. and Jimenez, J. L.: Volatility and lifetime against OH heterogeneous reaction of ambient isoprene-epoxydiols-derived secondary organic aerosol (IEPOX-SOA), Atmos. Chem. Phys., 16(18), 11563–11580, doi:10.5194/acp-16-11563-2016, 2016.

Jathar, S. H., Cappa, C. D., Wexler, a. S., Seinfeld, J. H. and Kleeman, M. J.: Multi-generational oxidation model to simulate secondary organic aerosol in a 3-D air quality model, Geosci. Model Dev., 8(8), 2553–2567, doi:10.5194/gmd-8-2553-2015, 2015.

Kang, E., Root, M. J., Toohey, D. W. and Brune, W. H.: Introducing the concept of Potential Aerosol Mass (PAM), Atmos. Chem. Phys., 7(22), 5727–5744, doi:10.5194/acp-7-5727-2007, 2007.

Kang, E., Toohey, D. W. and Brune, W. H.: Dependence of SOA oxidation on organic aerosol mass concentration and OH exposure: experimental PAM chamber studies, Atmos. Chem. Phys., 11(4), 1837–1852, doi:10.5194/acp-11-1837-2011, 2011.

Karjalainen, P., Timonen, H., Saukko, E., Kuuluvainen, H., Saarikoski, S., Aakko-Saksa, P., Murtonen, T., Bloss, M., Dal Maso, M., Simonen, P., Ahlberg, E., Svenningsson, B., Brune, W. H., Hillamo, R., Keskinen, J. and Rönkkö, T.: Time-resolved characterization of primary particle emissions and secondary particle formation from a modern gasoline passenger car, Atmos. Chem. Phys., 16(13), 8559–8570, doi:10.5194/acp-16-8559-2016, 2016.

Krechmer, J. E., Pagonis, D., Ziemann, P. J. and Jimenez, J. L.: Quantification of Gas-Wall Partitioning in Teflon Environmental Chambers Using Rapid Bursts of Low-Volatility Oxidized Species Generated in Situ, Environ. Sci. Technol., 50(11), 5757–5765, doi:10.1021/acs.est.6b00606, 2016.

Lakey, P. S. J., George, I. J., Whalley, L. K., Baeza-Romero, M. T. and Heard, D. E.: Measurements of the HO2 Uptake Coefficients onto Single Component Organic Aerosols, Environ. Sci. Technol., 49(8), 4878–4885, doi:10.1021/acs.est.5b00948, 2015.

Lambe, A., Massoli, P., Zhang, X., Canagaratna, M., Nowak, J., Daube, C., Yan, C., Nie, W., Onasch, T., Jayne, J., Kolb, C., Davidovits, P., Worsnop, D. and Brune, W.: Controlled nitric oxide production via O(¹D) + N₂O

reactions for use in oxidation flow reactor studies, Atmos. Meas. Tech., 10(6), 2283–2298, doi:10.5194/amt-10-2283-2017, 2017.

Lambe, A. T., Ahern, A. T., Williams, L. R., Slowik, J. G., Wong, J. P. S., Abbatt, J. P. D., Brune, W. H., Ng, N. L., Wright, J. P., Croasdale, D. R., Worsnop, D. R., Davidovits, P. and Onasch, T. B.: Characterization of aerosol photooxidation flow reactors: heterogeneous oxidation, secondary organic aerosol formation and cloud condensation nuclei activity measurements, Atmos. Meas. Tech., 4(3), 445–461, doi:10.5194/amt-4-445-2011, 2011.

Lambe, A. T. and Jimenez, J. L.: PAM Wiki: Publications Using the PAM Oxidation Flow Reactor, [online] Available from: https://sites.google.com/site/pamwiki/publications (Accessed 10 February 2017), 2017.

Levy II, H.: Normal atmosphere: large radical and formaldehyde concentrations predicted., Science, 173(3992), 141–143, doi:10.1126/science.173.3992.141, 1971.

Li, R., Palm, B. B., Borbon, A., Graus, M., Warneke, C., Ortega, a M., Day, D. a, Brune, W. H., Jimenez, J. L. and de Gouw, J. a: Laboratory Studies on Secondary Organic Aerosol Formation from Crude Oil Vapors, Environ. Sci. Technol., 47(21), 12566–12574, doi:10.1021/es402265y, 2013.

Li, R., Palm, B. B., Ortega, A. M., Hu, W., Peng, Z., Day, D. A., Knote, C., Brune, W. H., de Gouw, J. and Jimenez, J. L.: Modeling the radical chemistry in an Oxidation Flow Reactor (OFR): radical formation and recycling, sensitivities, and OH exposure estimation equation, J. Phys. Chem. A, 119(19), 4418–4432, doi:10.1021/jp509534k, 2015.

Link, M. F., Friedman, B., Fulgham, R., Brophy, P., Galang, A., Jathar, S. H., Veres, P., Roberts, J. M. and Farmer, D. K.: Photochemical processing of diesel fuel emissions as a large secondary source of isocyanic acid (HNCO), Geophys. Res. Lett., 43(8), 4033–4041, doi:10.1002/2016GL068207, 2016.

Lippmann, M.: Health effects of tropospheric ozone, Environ. Sci. Technol., 25(12), 1954–1962, doi:10.1021/es00024a001, 1991.

Liu, P. F., Abdelmalki, N., Hung, H.-M., Wang, Y., Brune, W. H. and Martin, S. T.: Ultraviolet and visible complex refractive indices of secondary organic material produced by photooxidation of the aromatic compounds toluene and m-Xylene, Atmos. Chem. Phys., 15(3), 1435–1446, doi:10.5194/acp-15-1435-2015, 2015.

Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, a., Huey, L. G., Cohen, R. C., Heikes, B., Singh, H. B., Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R. and Shetter, R. E.: Airborne measurement of OH reactivity during INTEX-B, Atmos. Chem. Phys., 9(1), 163–173, doi:10.5194/acp-9-163-2009, 2009.

Martinsson, J., Eriksson, A. C., Nielsen, I. E., Malmborg, V. B., Ahlberg, E., Andersen, C., Lindgren, R., Nyström, R., Nordin, E. Z., Brune, W. H., Svenningsson, B., Swietlicki, E., Boman, C. and Pagels, J. H.: Impacts of Combustion Conditions and Photochemical Processing on the Light Absorption of Biomass Combustion Aerosol, Environ. Sci. Technol., 49(24), 14663–14671, doi:10.1021/acs.est.5b03205, 2015.

Matsunaga, A. and Ziemann, P. J.: Gas-Wall Partitioning of Organic Compounds in a Teflon Film Chamber and Potential Effects on Reaction Product and Aerosol Yield Measurements, Aerosol Sci. Technol., 44(10), 881–892, doi:10.1080/02786826.2010.501044, 2010.

Moise, T. and Rudich, Y.: Reactive Uptake of Ozone by Aerosol-Associated Unsaturated Fatty Acids: Kinetics, Mechanism, and Products, J. Phys. Chem. A, 106(27), 6469–6476, doi:10.1021/jp025597e, 2002.

Moise, T., Talukdar, R. K., Frost, G. J., Fox, R. W. and Rudich, Y.: Reactive uptake of NO3 by liquid and frozen organics, J. Geophys. Res., 107(D2), 4014, doi:10.1029/2001JD000334, 2002.

Nehr, S., Bohn, B., Fuchs, H., Häseler, R., Hofzumahaus, A., Li, X., Rohrer, F., Tillmann, R. and Wahner, A.: Atmospheric photochemistry of aromatic hydrocarbons: OH budgets during SAPHIR chamber experiments, Atmos. Chem. Phys., 14(13), 6941–6952, doi:10.5194/acp-14-6941-2014, 2014.

Nel, A.: Air Pollution-Related Illness: Effects of Particles, Science (80-.)., 308(5723), 804–806, doi:10.1126/science.1108752, 2005.

Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. a., Prévôt, a. S. H., Dinar, E., Rudich, Y., Worsnop, D. R., Prevot, A. S. H., Dinar, E., Rudich, Y. and

780 Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol
781 Mass Spectrometry, Atmos. Chem. Phys., 10(10), 4625–4641, doi:10.5194/acp-10-4625-2010, 2010.

782 Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan Richard, C. and Seinfeld John, H.: Gas particle
783 partitioning and secondary organic aerosol yields, Environ. Sci. Technol., 30(8), 2580–2585,
784 doi:10.1021/es950943+, 1996.

785 Orlando, J. J. and Tyndall, G. S.: Laboratory studies of organic peroxy radical chemistry: an overview with
786 emphasis on recent issues of atmospheric significance, Chem. Soc. Rev., 41(19), 6294,
787 doi:10.1039/c2cs35166h, 2012.

788 Ortega, A. M., Day, D. A., Cubison, M. J., Brune, W. H., Bon, D., de Gouw, J. A. and Jimenez, J. L.:
789 Secondary organic aerosol formation and primary organic aerosol oxidation from biomass-burning
790 smoke in a flow reactor during FLAME-3, Atmos. Chem. Phys., 13(22), 11551–11571, doi:10.5194/acp-
791 13-11551-2013, 2013.

792 Ortega, A. M., Hayes, P. L., Peng, Z., Palm, B. B., Hu, W., Day, D. A., Li, R., Cubison, M. J., Brune, W. H.,
793 Graus, M., Warneke, C., Gilman, J. B., Kuster, W. C., de Gouw, J., Gutiérrez-Montes, C. and Jimenez, J. L.:
794 Real-time measurements of secondary organic aerosol formation and aging from ambient air in an
795 oxidation flow reactor in the Los Angeles area, Atmos. Chem. Phys., 16(11), 7411–7433,
796 doi:10.5194/acp-16-7411-2016, 2016.

797 Palm, B. B., Campuzano-Jost, P., Day, D. A., Ortega, A. M., Fry, J. L., Brown, S. S., Zarzana, K. J., Dube, W.,
798 Wagner, N. L., Draper, D. C., Kaser, L., Jud, W., Karl, T., Hansel, A., Gutiérrez-Montes, C. and Jimenez, J.
799 L.: Secondary organic aerosol formation from in situ OH, O3, and NO3 oxidation of ambient forest air in
800 an oxidation flow reactor, Atmos. Chem. Phys., 17(8), 5331–5354, doi:10.5194/acp-17-5331-2017, 2017.

801 Palm, B. B., Campuzano-Jost, P., Ortega, A. M., Day, D. A., Kaser, L., Jud, W., Karl, T., Hansel, A., Hunter, J.
802 F., Cross, E. S., Kroll, J. H., Peng, Z., Brune, W. H. and Jimenez, J. L.: In situ secondary organic aerosol
803 formation from ambient pine forest air using an oxidation flow reactor, Atmos. Chem. Phys., 16(5),
804 2943–2970, doi:10.5194/acp-16-2943-2016, 2016.

805 Peng, Z., Carrasco, N. and Pernot, P.: Modeling of synchrotron-based laboratory simulations of Titan's
806 ionospheric photochemistry, GeoResJ, 1–2, 33–53, doi:10.1016/j.grj.2014.03.002, 2014.

807 Peng, Z., Day, D. A., Ortega, A. M., Palm, B. B., Hu, W., Stark, H., Li, R., Tsigaridis, K., Brune, W. H. and
808 Jimenez, J. L.: Non-OH chemistry in oxidation flow reactors for the study of atmospheric chemistry
809 systematically examined by modeling, Atmos. Chem. Phys., 16(7), 4283–4305, doi:10.5194/acp-16-
810 4283-2016, 2016.

811 Peng, Z., Day, D. A., Stark, H., Li, R., Lee-Taylor, J., Palm, B. B., Brune, W. H. and Jimenez, J. L.: HOx radical
812 chemistry in oxidation flow reactors with low-pressure mercury lamps systematically examined by
813 modeling, Atmos. Meas. Tech., 8(11), 4863–4890, doi:10.5194/amt-8-4863-2015, 2015.

814 Ranney, A. P. and Ziemann, P. J.: Kinetics of Acid-Catalyzed Dehydration of Cyclic Hemiacetals in Organic
815 Aerosol Particles in Equilibrium with Nitric Acid Vapor, J. Phys. Chem. A, 120(16), 2561–2568,
816 doi:10.1021/acs.jpca.6b01402, 2016.

817 Richards-Henderson, N. K., Goldstein, A. H. and Wilson, K. R.: Large Enhancement in the Heterogeneous
818 Oxidation Rate of Organic Aerosols by Hydroxyl Radicals in the Presence of Nitric Oxide, J. Phys. Chem.
819 Lett., 6, 4451–4455, doi:10.1021/acs.jpclett.5b02121, 2015.

820 Saltelli, A., Ratto, M., Tarantola, S. and Campolongo, F.: Sensitivity Analysis for Chemical Models, Chem.
821 Rev., 105(7), 2811–2828, doi:10.1021/cr040659d, 2005.

822 Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Wine, P. H., Abbatt, J. P. D., Burkholder,
823 J. B., Kolb, C. E., Moortgat, G. K., Huie, R. E. and Orkin, V. L.: Chemical Kinetics and Photochemical Data
824 for Use in Atmospheric Studies Evaluation Number 17, Pasadena, CA, USA. [online] Available from:
825 http://jpldataeval.jpl.nasa.gov/pdf/JPL 10-6 Final 15June2011.pdf, 2011.

826 Schill, G. P., Jathar, S. H., Kodros, J. K., Levin, E. J. T., Galang, A. M., Friedman, B., Link, M. F., Farmer, D.
827 K., Pierce, J. R., Kreidenweis, S. M. and DeMott, P. J.: Ice-nucleating particle emissions from
828 photochemically aged diesel and biodiesel exhaust, Geophys. Res. Lett., 43(10), 5524–5531,
829 doi:10.1002/2016GL069529, 2016.

830 Schwantes, R. H., Schilling, K. A., McVay, R. C., Lignell, H., Coggon, M. M., Zhang, X., Wennberg, P. O. and

831 Seinfeld, J. H.: Formation of highly oxygenated low-volatility products from cresol oxidation, Atmos.
832 Chem. Phys., 17(5), 3453–3474, doi:10.5194/acp-17-3453-2017, 2017.

833 Seakins, P. W.: A brief review of the use of environmental chambers for gas phase studies of kinetics,
834 chemical mechanisms and characterisation of field instruments, EPJ Web Conf., 9, 143–163,
835 doi:10.1051/epjconf/201009012, 2010.

836 Simonen, P., Saukko, E., Karjalainen, P., Timonen, H., Bloss, M., Aakko-Saksa, P., Rönkkö, T., Keskinen, J.
837 and Dal Maso, M.: A new oxidation flow reactor for measuring secondary aerosol formation of rapidly
838 changing emission sources, Atmos. Meas. Tech., 10(4), 1519–1537, doi:10.5194/amt-10-1519-2017,
839 2017.

840 Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and
841 Midgley, P. M.: Climate Change 2013 - The Physical Science Basis, edited by Intergovernmental Panel on
842 Climate Change, Cambridge University Press, Cambridge., 2014.

843 Strollo, C. M. and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from
844 the reaction of 3-methylfuran with OH radicals in the presence of NOx, Atmos. Environ., 77, 534–543,
845 doi:10.1016/j.atmosenv.2013.05.033, 2013.

846 Tkacik, D. S., Lambe, A. T., Jathar, S., Li, X., Presto, A. A., Zhao, Y., Blake, D., Meinardi, S., Jayne, J. T.,
847 Croteau, P. L. and Robinson, A. L.: Secondary Organic Aerosol Formation from in-Use Motor Vehicle
848 Emissions Using a Potential Aerosol Mass Reactor, Environ. Sci. Technol., 48(19), 11235–11242,
849 doi:10.1021/es502239v, 2014.

850 Volkamer, R., Jimenez, J. L., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L. T., Worsnop,
851 D. R. and Molina, M. J.: Secondary organic aerosol formation from anthropogenic air pollution: Rapid
852 and higher than expected, Geophys. Res. Lett., 33(17), L17811, doi:10.1029/2006GL026899, 2006.

853 Wang, J., Doussin, J. F., Perrier, S., Perraudin, E., Katrib, Y., Pangui, E. and Picquet-Varrault, B.: Design of
854 a new multi-phase experimental simulation chamber for atmospheric photosmog, aerosol and cloud
855 chemistry research, Atmos. Meas. Tech., 4(11), 2465–2494, doi:10.5194/amt-4-2465-2011, 2011.

856 Zhang, X., Cappa, C. D., Jathar, S. H., McVay, R. C., Ensberg, J. J., Kleeman, M. J. and Seinfeld, J. H.:
857 Influence of vapor wall loss in laboratory chambers on yields of secondary organic aerosol., Proc. Natl.
858 Acad. Sci. U. S. A., 111(16), 5802–7, doi:10.1073/pnas.1404727111, 2014.

859 Ziemann, P. J. and Atkinson, R.: Kinetics, products, and mechanisms of secondary organic aerosol
860 formation, Chem. Soc. Rev., 41(19), 6582, doi:10.1039/c2cs35122f, 2012.

861



OFR185-iNO

(H_2O=1.5%, UV at 185 nm=1.5x10^{12} photons cm^{-2} s^{-1}, OHR_{ext}=15 s^{-1}, NO^{in}=150 ppb)
($F185_{exp}/OH_{exp}$~600 cm/s, $F254_{exp}/OH_{exp}$~1x10^5 cm/s)

862



OFR185-iNO

(H_2O=1%, UV at 185 nm=1x10^{13} photons cm^{-2} s^{-1}, OHR_{ext}=3000 s^{-1}, NO^{in}=30 ppm, T=295 K)
($F185_{exp}/OH_{exp}$~2x10^5 cm/s, $F254_{exp}/OH_{exp}$~2x10^7 cm/s)

863



OFR254-7-iNO

(H_2O=1.5%, UV at 254 nm=3.4x10^{14} photons cm^{-2} s^{-1}, OHR_{ext}=15 s^{-1}, NO^{in}=150 ppb, T=295 K)

864

Figure 1. Schematics of main N-containing species and their major interconversion pathways under typical input conditions for (a) OFR185-iNO with NO^{in}=150 ppb, (b) OFR254-7-iNO with NO^{in}=150 ppb, and (c) OFR185-iNO with NO^{in}=30 ppm. Species average concentrations (in molecules cm^{-3}) are shown in black beside species names. Arrows denote directions of the conversions. Average reaction fluxes (in units of 10^9 molecules cm^{-3} s^{-1}) are calculated according to the production rate, and shown on or beside the corresponding arrows and in the same color. Within each schematic, the thickness of the arrows is a measure of their corresponding species flux. Multiple arrows in the same color and pointing to the same species should be counted only once for reaction flux on a species. Note that all values in these schematics are average ones over the residence time, except for those in square brackets in panels a and b, which are average values within approximate NO effective lifetime (τ_{NO}, or more accurately, an integer multiple of the model's output time step closest to NO effective lifetime). All concentrations and fluxes have two significant digits.


NO exposure
NO3 exposure
OH exposure
Relative variance
Rel. uncertainty (%)
Case
MLVH LHLH HM0L MHHL MHVH HMLH HLHL MMVV MLHH LLLL HL0H LMHV HHVL LMLH LH0L
N2O5→NO2+NO3
HO2NO2→NO2+HO2
NO2+NO3→N2O5
HO2+NO3→OH+NO2+O2
OH+NO→HONO
O(3P)+NO2→NO3
2HO2→H2O2+O2
H2O+hv185→OH+H
O3+hv254→O2+O(1D)
O2+hv185→2O(3P)
OH+HNO3→H2O+NO3
OH+NO2→HNO3
NO2+O3→NO3+O2
NO+O3→NO2+O2
OH+HO2→H2O+O2
HO2+O3→OH+2O2
HO2+NO→OH+NO2
OH+O3→HO2+O2
O(1D)+N2→O(3P)+N2
Other effects

877

878 **Figure 2.** Relative variances (left axes)/uncertainties (right axes) of several outputs (i.e., NO, NO_3, and OH exposures) of Monte Carlo uncertainty propagation, and relative
879 contributions of key reactions to these relative variances in several typical cases (denoted in 4-character labels, see Table 2 for the typical case label code) in OFR185-iNO.
880 Relative variances are shown in linear scales (left axis), while corresponding relative uncertainties, equal to relative variances' square roots, are indicated by the non-linear
881 right axis. Only the reactions with a contribution of no less than 0.04 to at least one relative variance are shown.



882

883 **(a) OFR254-iNO**



884

885 **(b) OFR185-iNO**

886 **Figure 3.** Frequency occurrence distributions of good, risky, and bad conditions (see Table 3) over
887 logarithm of the ratio between RO_2 reacted with NO and with HO_2 (see Section S1 for more detail) for
888 (a) OFR254-iNO (only the case with a residence time of 180 s) and (b) OFR185-iNO (including two cases
889 with residence times of 180 and 30 s). Low and high-NO regions (see Table 3) are colored in light blue
890 and grey, respectively.


Increasing UV
Increasing H2O
Bad low-NO
Risky low-NO
Good low-NO
Bad high-NO
Risky high-NO
Good high-NO
OHRext (s-1)
Initial NO (ppb)
Case LL
Case LM
Case LH
Case ML
Case MM
Case MH
Case HL
Case HM
Case HH

899 **Figure 4.** Image plots of the condition types defined in Table 3 vs. external OH reactivity (excluding N-containing species) and initial NO for several typical cases in OFR185-
900 iNO (see Table 2 for the case label code).
901


Increasing UV
Increasing H2O
Bad low-NO
Risky low-NO
Good low-NO
Bad high-NO
Risky high-NO
Good high-NO
OHRext (s-1)
Initial NO (ppb)
Case LL
Case LM
Case LH
Case ML
Case MM
Case MH
Case HL
Case HM
Case HH

910 **Figure 5.** Same format as Fig. 4, but for OFR254-22-iNO.



Figure 6. Fractional importance of the reaction rate of several species of interest with NO_3 vs. that with OH, as a function of the ratio of exposure to NO_3 and OH. The curves of biogenics and phenols are highlighted by solid dots and squares, respectively. The turquoise and orange markers show the ranges of modeled exposure ratios between NO_3

and OH of a source study in an urban tunnel (Tkacik et al., 2014) and a laboratory study (Liu et al., 2015) using OFR, respectively. In the upper part of the figure, the modeled frequency distributions of ratios of NO_3 exposure to OH exposure under good/risky/bad high/low-NO conditions for OFR185-iNO and OFR254-iNO are also shown. See Table 3 for the definitions of the three types of conditions. All curves, markers, and histograms in this figure share the same abscissa.


Bad low-NO
Risky low-NO
Good low-NO
Bad high-NO
Risky high-NO
Good high-NO
No dilution
Diluted by x12
Diluted by x100
NO (ppm)
Total hydrocarbon (ppm)
Time (s)
OH exposure (molecules cm^{-3} s)
Percentage of remaining OH after suppression
Relative importance of non-OH fate of benzene (%)
NO_3 exposure /OH exposure
Risky
Bad
NO effective lifetime (s)
No dilution
Diluted by x12
Diluted by x100
Relative importance of R+NO_2 in OH-benzene adduct fate (%)

Figure 7. (left) NO and total hydrocarbon during the first 200 s of the test of Karjalainen et al. (2016) in the cases of no dilution, dilution by a factor of 12 (as actually done in that study), and dilution by a factor of 100. Different periods of time are colored according to corresponding emissions (i.e., input conditions for OFR), classified as good/risky/bad high/low-NO. (right) OH exposure/percentage of remaining OH after suppression, relative importance of non-OH fate of benzene, exposure ratio of NO_3 to OH, NO effective lifetime, and relative importance of reaction of OH-toluene adduct with NO_2 in the fate of this adduct in the OFR of Karjalainen et al. (2016) during the first 200 s of their test in the cases of no dilution, dilution by a factor of 12, and dilution by a factor of 100. Horizontal orange and red dashed lines in the middle right panel denote "risky" and "bad" regions for exposure ratio of NO_3 to OH, respectively. Above the orange (red) dashed line, reaction with NO_3 contributes >20% to the fate of phenol (isoprene).



Figure 8. Secondary organic aerosol (SOA) potential (estimated from the total hydrocarbon measurement) in the OFR of Karjalainen et al. (2016) formed during periods of time in the OFR corresponding to good/risky/bad high/low-NO conditions, as a function of dilution factor. Vertical lines denoting dilution factors of 1, 12 (as actually used in that study), and 100 are also shown.



931

932 **(a) No dilution (background: Case HH)**



933

934 **(b) Dilution by a factor of 100 (background: Case HH)**



935

936 **(c) Dilution by a factor of 100 (background: Case HL)**

937 **Figure 9.** Location of individual 1 s datapoints vs. OFR185-iNO reaction conditions. Datapoints are shown
938 from the test vehicle of Karjalainen et al. (2016), as well as average exhaust from gasoline vehicle on-
939 road emissions measured by Bishop and Stedman (2013). On-road emissions are classified by vehicle
940 year and the distribution of each category is shown as a cross representing 1 standard deviation (with
941 log-normal distribution assumed). The X and Y axes are NO and external OH reactivity (excluding N-
942 containing species) due to vehicle emissions in OFR in the cases of (a) no dilution and (b,c) dilution by a
943 factor of 100. The Karjalainen et al. (2016) points are classified as cold start (during first 200 s) and hot
944 stabilized (during 200–1000 s). In addition, the same image plots as the panels of Cases HH (high H_2O
945 and high UV, see Table 2 for the case label code) and HL in Fig. 4 (OFR185-iNO) are shown as background
946 for comparison.

947

Scheme 1. Possible major reactions in an OFR254-13-iNO with 5 ppm toluene and 10 ppm initial NO. Branching ratios in red are estimated by the model and/or according to Calvert et al. (2002), Atkinson and Arey (2003), Ziemann and Atkinson (2012), and Peng et al. (2016). Note that addition/substitution on the aromatic ring may occur at other positions. Intermediates/products shown here are the isomers that are most likely to form. Branching ratios shown in red are not overall but from immediate reactant.



Table 1. Experimental conditions of several OFR studies with high NO injection.

Study	Source type	Temperature (K)	Relative humidity (%)	Dilution factor	External OH reactivity of undiluted source (s^{-1})	Source NO_x concentration (ppm)
Link et al. (2016)	Diesel vehicle emission		50	45–110	~5000[*1]	436[*1]
Martinsson et al. (2015)	Biomass burning emission			1700	156400[*1]	154
Karjalainen et al. (2016)	Gasoline vehicle emission	295	60	12	~73000[*2,a]	~400[*1,b]
Liu et al. (2015)	Purified gas	293	13	1	~1400[*1,a]	10[*1,b]
Tkacik et al. (2014)	Tunnel air	293	42	1	~60[*1,a]	~0.8[*1]
Ortega et al. (2013)	Biomass burning emission	290	30	~500	~15-500	~0.2

[*1] maximum value in the study
[*2] value at the moment of maximum NO emission
[*a] NO_y species excluded
[*b] NO only

Table 2. Code of the labels of typical cases. A case label can be composed of four characters denoting the water mixing ratio, the photon flux, the external OH reactivity excluding N-containing species, and the initial NO mixing ratio, respectively. A case label can also be composed of two characters denoting the water mixing ratio and the photon flux.

	Water mixing ratio	Photon flux	External OH reactivity (no ON)	Initial NO mixing ratio
Options	L=low (0.07%)	L=low (10^{11} photons cm^{-2} s^{-1} at 185 nm; $4.2x10^{13}$ photons cm^{-2} s^{-1} at 254 nm)	0	0
	M=medium (1%)	M=medium (10^{13} photons cm^{-2} s^{-1} at 185 nm; $1.4x10^{15}$ photons cm^{-2} s^{-1} at 254 nm)	L=low ($10\ s^{-1}$)	L=low (10 ppb)
	H=high (2.3%)	H=high (10^{14} photons cm^{-2} s^{-1} at 185 nm; $8.5x10^{15}$ photons cm^{-2} s^{-1} at 254 nm)	H=high ($100\ s^{-1}$)	H=high (316 ppb)
			V=very high ($1000\ s^{-1}$)	V=very high (10 ppm)
Example	LH0V:	low water mixing ratio, high photon flux, no external OH reactivity (excluding ON), very high initial NO mixing ratio		
	ML:	medium water mixing ratio, low photon flux		

963 **Table 3.** Definition of condition types in this study (good/risky/bad high/low-NO).

Condition	Good	Risky	Bad
Criterion	$F185_{exp}/OH_{exp}<3x10^3$ cm s^{-1} and $F254_{exp}/OH_{exp}<4x10^5$ cm s^{-1}	$F185_{exp}/OH_{exp}<1x10^5$ cm s^{-1} and $F254_{exp}/OH_{exp}<1x10^7$ cm s^{-1} (excluding good conditions)	$F185_{exp}/OH_{exp}\geq1x10^5$ cm s^{-1} or $F254_{exp}/OH_{exp}\geq1x10^7$ cm s^{-1}
Condition	High-NO		Low-NO
Criterion*	$\frac{r(RO_2+NO)}{r(RO_2+HO_2)} > 1$		$\frac{r(RO_2+NO)}{r(RO_2+HO_2)} \leq 1$

964 * See Section S1 for detail.
965

966 **Table 4.** Statistics of the ratio between OH exposures calculated in the model with the Lambe et al. (2011) residence time distribution ($OH_{exp,RTD}$) and in the plug-flow model
967 ($OH_{exp,PF}$). The geometric mean, uncertainty factor (geometric standard deviation), and percentage of outlier cases (>3 or <1/3) are shown for OFR185-iNO, OFR254-70-iNO,
968 and OFR254-7-iNO.

	Geometric mean	Uncertainty factor	Outlier cases (%)
OFR185-iNO	1.91	1.64	11
OFR254-7-iNO	1.59	1.51	7
OFR254-70-iNO	1.48	1.29	3

969