

Author's response to the comments of Anonymous Referee #1 and #2 and the track changes version.

Author's response to the comments of Anonymous Referee #1. - below

Author's response to the comments of Anonymous Referee #2. - from page 11

The track changes version. - from page 14

5

Author's response to the comments of Anonymous Referee #1.

Authors comments are written in bold.

MAIN ISSUES

10

The paper cannot be published as is, and major revisions are required. I do not mean that the main outcomes of the work (trends of ozone and aerosol optical depths) are wrong, but several issues should be settled, or at least discussed, before the paper can be accepted.

15 1/ The determination of the ETC appears as the main problem, along with the estimate of its uncertainty:

1a/ the uncertainty of the mean, better taking into account the number of ETC estimates, should be used instead of the standard deviation of the ETC estimates. Otherwise, as said in the paper, the uncertainty can be lower for a time interval with few ETC estimates than for one with several estimates;

20 **The problem with a few, namely 2 ETC estimates was only in the first intercalibration period. This problem was solved by using additional data from the year 1993. Thanks to it the number of ETC estimates is sufficient now. The minimum number of ETC estimates for all intercalibration periods and all wavelengths is 7 now. Therefore I think that the standard deviation of the ETC estimates is a good tool for the estimate of the uncertainty of AOD.**

25 1b/ in his reply to my review, the author states that "the values of the uncertainties of the annual averages do not enter the calculation of the uncertainty of the linear trend". Instead, I think that the uncertainties should enter the calculation of both the value and the uncertainty of the linear trend;

Excuse me, but I do not know how to take correctly into account the uncertainties of the annual averages in the trend calculations.

30

1c/ minimum data coverage thresholds (e.g., minimum measurement days per month, minimum measurement months per year) should be fixed before averaging the data (p. 11 l. 9-15), especially in case of large gaps (due to maintenance of the

instrument or clouds) and presence of a seasonal cycle. Maybe this is not an issue, but it should be specified how long is the longest period without data, if any;

This sentences was added to the manuscript: „The longest period without data is 12 days for the ozone optical depth and 26 days for AOD. There was no month without data. In both cases, February 2013 was the month with the absolute lowest number of days with the determined optical depth. In the first case, it was 10 days, and it was 3 days in the second case. With regard to the long-term average, December is the month with the lowest number of days with the determined AOD with 13 days. The opposite extreme falls on August with 24 days.“

1d/ I could have missed this information, but I do not understand why "an advantage of the LPM is that it takes into account the change of ETCs during the intercalibration period" (p. 16 l. 19-20). From Fig. 4, I can understand that the ETCs estimated by the LPM are averaged and fixed for an intercomparison period, aren't they?

Maybe the problem was my bad formulation and therefore I have changed related text in the manuscript a little bit: „An advantage of the presented LPM is that it partially takes into account the potential change in the sensitivity of the instrument during the intercalibration period, because ETCs are determined based on real measurements in a given period. The BSM utilizes ETCs determined during calibration. Therefore, the potential change in the sensitivity of the instrument during the intercalibration period is not taken into account at all“

1e/ is the Langley plot performed by correcting S_{λ} (Eq. 3) for O₃ absorption or by including O₃ in τ_{λ} ? In the first case, why the ozone cycle is mentioned as a source of error (p. 5)? In the second case, how is μ_w calculated (including ozone)?

O₃ is including in τ_{λ} . This explanation is included in the manuscript: „ μ_w is the air mass factor for atmosphere as a whole. Its value was calculated as a weighted arithmetic average of individual aforementioned components, while the optical depth of a given component was the weighting factor.“ Moreover, the text located after equation (4) was enriched about an information related to calculation of μ_w : The final values of ETCs used to calculate the AOD were available only after two iterations. At the beginning, initial values of AOD were not available yet. For this reason, it was not possible to apply the second and third criterion and it was also not possible to take into account the effect of aerosols in the calculation of μ_w and to determine the correction for the diffuse radiation. It was possible to do so on the first and second iteration.

1f/ since the BSM method is basically a calibration transfer from a reference instrument, the author should explain how Brewer #017 was calibrated;

This explanation was added to the manuscript: " ETCs for the reference instrument are determined by LPM at the Mauna Loa or Izaña observatory."

2/ the AOD differences between the LPM and the BSM methods can be easily attributed to the ETC difference ($\text{AOD difference} = 1/\mu * \log(\text{ETC1}/\text{ETC2})$). Therefore, the deviations between the two methods are mainly not "random" (contrary to what is written at p. 15 l. 14 and at p. 16 l. 12); the observed variability between summer and winter (p. 15 l. 21-23) is an obvious consequence of air mass variation during the year; the AOD differences, always positive in the first period and always negative in the second period, are dictated by the difference in ETC shown in Fig. 5. I would thus recommend to plot the AOD differences as a function of air mass to check that the ETC difference is the main cause of deviation;

Note in the bracket at p. 15 l. 14: „so-called large random error" was removed. Word "random" was removed in related sentence (p. 16 l. 12) and the sentence was reword. I have investigated the AOD differences as a function of air mass for the longest wavelength. Possible causes are discussed in the manuscript, included ETCs difference. It has been found that: " The values of the determined ETCs affect the observed differences the most. However, it is true only if their difference between LPM and BSM is sufficiently large." It was added to the manuscript.

3/ as I mentioned in my previous review, an additional source of errors can be the missing SL correction in the BSM, which is not mentioned among the sources listed at p. 16. However, according to the maintainer of the Brewer operating software: "Using the real-time AOD (or even ozone) data from the operating software for anything other than a rough idea about the AOD doesn't make any sense since no corrections are applied in real time for the SL ratio changes. It's likely worse than real time data for ozone due to potential changes in absolute sensitivity in Brewers (that do not affect ozone)" (V. Savastiouk, personal communication, 2017). This issue should be discussed and its potential impact estimated;

Using the real-time AOD (or even ozone) data from the operating software for anything other than a rough idea about the AOD doesn't make any sense since no corrections are applied in real time for the SL ratio changes. **My comment to this sentence: I know that BSM has a lot of disadvantages, but it was used also for calculation of AOD in the study Kumharn et al., 2012.**

It's likely worse than real time data for ozone due to potential changes in absolute sensitivity in Brewers (that do not affect ozone). **My comment to this sentence: I know (I asked IOS) that SL correction can be used for ozone calculation but not for AOD calculation (for counts rates corrections) because standard lamp has been losing its intensity over time. For example, we observe an decrease in count rates, but we don't know if the reason is a change in absolute sensitivity in Brewer or a change in the intensity of standard lamp.**

Deep analysis of the influence of missing SL correction in case of BSM is outside the scope of this paper. But it was mentioned as a potential source of error. This sentence was added to the manuscript: " The disadvantage of BSM is also the absence of the SL (standard lamp) test that leads to inaccurate TCO values, which consequently affects the size of AOD."

4/ even though it is probably a minor issue (especially at low air masses), the spectral stray light effect is inaccurately taken into account, in my opinion, for two reasons:

5 4a/ both the slits employed for UV scans and the grating positions are different from those of DS measurements, therefore the ratio between the irradiance at longer and shorter wavelengths in the "scanning mode" may not be representative of stray light in "slitmask" mode (DS measurements). The cited study of Arola & Koskela (2004), using this method, only aimed at a rough estimate of the effect;

**This sentence was added to the manuscript: " The potential impact of a different measurement mode was neglected
10 for the sake of simplification."**

4b/ the assumption of spectrally constant stray light is simplistic, as pointed out in Garane et al. (2006), and will lead to some uncertainty.

**The end of part "2.4.1 Correction for the diffuse radiation and stray-light effect" was changed: " In view of
15 simplification, it is also assumed that the value of stray light is constant for all wavelengths. The said simplifications may lead to some uncertainties when estimating the stray light effect. These uncertainties were not investigated further."**

Therefore, if the author wants to keep the analysis of wavelengths shorter than 320 nm, they should point out that they could be affected by the stray light, rather than trying to correct them and jump to the conclusion that the Angstrom coefficient is
20 negative;

**I have decided to let the stray light correction in the manuscript. Now, I know that is not perfect, but I think that it is better like no correction. In the Conclusions is stated that the Ångström exponent is probably negative mainly due to the size and accuracy of ETC determination: "It was also determined that the application of corrections to the diffuse radiation, stray-light effect and polarization decreased the difference between the value of AOD for the shortest and
25 longest of the examined wavelengths. Although the difference has been reduced, it still reaches negative values, which results in the negative values of the Ångström exponent as well. With regard to the long-term average for 23 years, the difference in AOD is only –0.006. The application of the correction for the polarization had the most significant influence on the reduction of the given difference. On the contrary, the lowest contribution was achieved by the correction with regard to the diffuse radiation. The key factor influencing the value of the examined difference is
30 probably the size and accuracy of ETC determination."**

5/ most of all, language needs major revisions, as well as the structure of the paper:

5a/ check the whole text for the correct use of articles "a" and "the";

Several sections of the text have been repaired.

5b/ remove all technical details that are not fundamental for the paper (e.g., discussion about wavelengths, p. 4) and cite some reference where the information can be found elsewhere (e.g., instead of the data reduction explained at p. 8);

Some parts (not only technical) of the manuscript has been removed, namely:

- 5 **Page 2: „The direct impact means scattering and absorption of shortwave and longwave radiation. Absorption of radiation subsequently leads to warming of those atmospheric parts, where aerosols are present (primarily in the boundary layer of the atmosphere) and higher temperature consequently leads to evaporation of cloud layers. The last sentence concisely described the semidirect impact, resulting in a higher density of solar radiation flow reaching the Earth’s surface (Cazorla et al., 2009). The higher temperature may lead to a change of thermal stratification of**
- 10 **the atmosphere, which consequently affects vertical and horizontal movements of air in the atmosphere. The indirect impact pertains to the ability of aerosols to act as condensation nuclei or ice nuclei, which affects microphysical and optical properties of clouds. Yet, it concerns a change of their radiation characteristics, change of atmospheric precipitation characteristics, and alteration in cloud lifetime as well. An increase in the number of condensation nuclei leads to the rise of cloud droplet count and to the reduction of their size under the given conditions of water**
- 15 **content in the atmosphere, causing an increase of albedo and extension of cloud lifetime (Lohmann and Feichter, 2005; Unger et al., 2009).“**

- Page 4: „The spectral separation of solar radiation is carried out by means of a modified Ebert f/6 spectrometer, which uses a holographic diffraction grating with a resolution of 1,200 lines per one mm (Sci-Tec, 1999). The instrument stability of Brewer spectrophotometer measurements is ± 0.01 nm, namely throughout the temperature**
- 20 **range (Sci-Tec, 1999). The value of the smallest wavelength increment (microstep) is 0.006 ± 0.002 nm (Sci-Tec, 1999).“**

This sentence has been added instead of the removed text : „More informations about the instrument and the measurement process can be found in Kerr (2010).“

- Page 4: „During the monitored 23 years, the sizes of wavelengths were refined five times in total. Their accurate sizes**
- 25 **were used at the beginning of the examined period. In 1994, the sizes were 306.276 nm, 310.04 nm, 313.494 nm, 316.799 nm and 319.999 nm. The end of the period, that is 2015, was characterized by values of 306.276 nm, 310.035 nm, 313.476 nm, 316.755 nm and 319.989 nm. Observed deviations are only minimal. Therefore, the aforementioned long-term average is generally applicable to the whole period.“**

- 30 5c/ split the theory ("Methods") from the results. E.g., p. 11 l. 19-26 are theory, as well as p. 12 l. 5-14;

It was accepted. This new part of subchapter 2.4 has been created: "2.4.1 Correction for the diffuse radiation and stray-light effect". Moreover, the main structure of subchapter 2.4 has been changed. It means that, also other parts of subchapter 2.4 has been created, namely:

2.4.2 Calculation of extraterrestrial constants

2.4.3 Quality control process

2.4.4 Additional notes

5d/ use letters, e.g. (a), (b), etc., not "right/left", to identify the subfigures in the panels, according to the

5 ACP guidelines

(https://www.atmospheric-chemistry-and-physics.net/for_authors/manuscript_preparation.html).

It was accepted.

TECHNICAL CORRECTIONS

10 - replace "Operational Brewer Program" with "Brewer Operating Software";

It was accepted.

- better use "AERONET" instead of "Cimel", since the data are provided by AERONET;

I did not accept it because "Cimel sunphotometer" is common in another scientific publications, for example:

15 **De Bock, V., De Backer, H., Mangold, A., and Delcloo, A.: Aerosol Optical Depth measurements at 340 nm with a Brewer spectrophotometer and comparison with Cimel sunphotometer observations at Uccle, Belgium, Atmos. Meas. Tech., 3, 1577-1588, <https://doi.org/10.5194/amt-3-1577-2010>, 2010.**

López-Solano, J., Redondas, A., Carlund, T., Rodriguez-Franco, J. J., Diémoz, H., León-Luis, S. F., Hernández-Cruz, B., Guirado-Fuentes, C., Kouremeti, N., Gröbner, J., Kazadzis, S., Carreño, V., Berjón, A., Santana-Díaz, D.,

20 **Rodríguez-Valido, M., De Bock, V., Moreta, J. R., Rimmer, J., Boulkelia, L., Jepsen, N., Eriksen, P., Bais, A. F., Shiroto, V., Vilaplana, J. M., Wilson, K. M., and Karppinen, T.: Aerosol optical depth in the European Brewer Network, Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2017-1003>, in review, 2017.**

- p. 1: use the long form "spectral stray light" instead of just "stray light" at the first occurrence, to differentiate it from other sources of stray light;

25 **It was accepted.**

- p. 3: latitude °N and longitude °E (NW/EW?);

It was accepted.

30 - p. 3: order the pollution sources by relevance;

The local pollution sources are mentioned by relevance. It means solid fuel combustion and the agriculture as first and second. Industrial activities in near city as third.

- p. 4: unless very old instruments are considered, UV, SO₂ and O₃ estimates are performed by all instruments. NO₂ instruments are performed only by MKIV Brewers, which are no longer manufactured;

The problem sentence has been corrected: The model MKIV allow moreover measuring the vertical column of SO₂, NO₂ (for this purpose, measurements of solar radiation are made in the visible region of the spectrum) and global UV radiation as well (from 290 to 325 nm, with an increment of 0.5 nm).

- p. 5: it is said that LPM is not recommended for low latitude stations, however Poprad-Gánovce (706 m a.s.l.) cannot be considered a high altitude station. Please, clarify this point;

I agree, that the Poprad-Gánovce station (706 m a.s.l.) can not by considered as a high altitude station but also is not a typical low altitude stations in urbanized area.

Related text in the manuscript was enriched. This description in the manuscript clarify your remark: „In the light of the listed errors, it is not recommended to determine extraterrestrial constants (ETCs) using the Langley plot method (LPM) for low altitude stations in urbanized areas, unless corrections for the impact of diffuse radiation and the daily cycle of ozone are known. Poprad-Gánovce is not a typical station in a low-altitude urbanized area and is not even a typical high mountain station. It is between these two cases. The impact of daily ozone cycle and NO₂ can be neglected considering the rural location and higher altitude of the Poprad-Gánovce station. The diffuse radiation impact and stray-light effect were not neglected.“

- p. 5: how were the air mass thresholds of 3 (line 12) and 4 (line 32) established?

The scientific reference was added to the manuscript: „A recommendation to ensure the air mass factor does not exceed the value of 3 in the calculation of ETCs was taken into account as well (Arola and Koskela, 2004; De Bock et al., 2010).“

Also the change related to threshold 4 has occurred in the manuscript: „A DS measurement is accepted only with the relative optical mass of less than 4 (as recommended by IOS) and it takes approximately 2.5 minutes.“

- p. 5: explain better the "polarization effect" and use the form "internal polarization effect";

The related part of the manuscript was enriched about this explanation: „It occurs due to the combined effect of two polarization sensitive elements of the Brewer spectrophotometer. The entrance window is the first of mentioned elements and the grating is the second. Polarization effect depends on the zenith angle of the Sun and is almost wavelength independent.“

The form "internal polarization effect" has used at the first occurrence instead of the form "polarization effect".

- p. 5: "corrections published in Cede at al. (2006)" -> which one of the methods described in the paper was used?

The name of the used method was added in to the brackets. „As a result, corrections (the field measurements method) published in Cede et al. (2006) could be applied to the instrument No. 97.“

- p. 6: include the physical units for the Loschmidt constant;

5 It was accepted: „ n is the molecule count in the volume determined by 1 DU and 1 cm²; for O₃, it is a constant with the value of $n = 2.687 * 10^{16} \text{ cm}^{-3}$ “

- p. 7: remove repetitions in lines 1-3;

Some changes has occurred in the manuscript. But I'm not sure if I understand your request properly (for more information about the changes in the text, please see page 6, lines 22-27).

10

- p. 7: explain why the method works best at lower latitudes;

The reasons are mentioned in the previous text: „It is also necessary to avoid an impact of cloudiness on direct solar radiation (DS) measurements and to ensure a sufficient scope of zenith angles of individual DS measurements during the day, which is needed for the given method.

15 For the said reasons, this method is most appropriate for lower latitudes (especially in mountainous regions near the tropics), and it has certain limitations in the middle and particularly higher latitudes (Nieke et al. 1999; Marengo 2007).“

- p. 7: "implemented", not "developed" by Vladimir Savastiouk;

20 It was accepted.

- p. 7: are the alternative Komhyr (1980) and Kasten and Young (1989) formulations used for O₃ as well?

25 This explanation was added to the manuscript: „Brewer Operating Software and O3Brewer software use the same formula to calculate an airmass factor for ozone. The same applies to Rayleigh scattering. The given formulas are defined in the Brewer MKIV Spectrophotometer Operator's Manual (Sci-Tec, 1999). Different, yet on the other hand more accurate formulas, were used to calculate the AOD by LPM – for ozone according to Komhyr (1980) and for Rayleigh scattering according to Kasten and Young (1989).“

- p. 8: why was the stray light correction applied at that stage?

30 This explanation was added to the manuscript: „After the deadtime compensation, a correction was applied to the stray-light effect, in the same order as Garane et al. (2006).“

- p. 8: explain if ND filter spectral transmittance was taken into account for AOD calculations;

A change has occurred in the manuscript. Related text including explanation is: „In the fourth step, a correction for the temperature dependence was applied, including a spectral transmittance correction for the utilized neutral density (ND) filter. These filters are automatically selected by the Brewer spectrophotometer with respect to the current density of the solar radiation flow. There are 5 ND filters and 5 wavelengths as well, so 25 attenuation values are needed in total. The attenuation values of given filters are determined during calibration of the instrument.“

- p. 8: what are the "practical purposes"?

The sentence has been changed: „Theoretically, it is feasible to determine the unknowns already from two measurements, but for the ETC determination, it is advisable to acquire as many measurements as possible.“

- p. 9: the conditions 1-3 are related to the single measurement, while conditions 4-9 concern the whole day. They should be splitted;

It was accepted. The conditions has been splitted.

- p. 11: "more relevant" compared to what?

The sentence was enriched about the explanation: „The main goal of this calculation was to acquire a comparison of AOD for the same wavelength, which is more relevant than a direct comparison of AOD for two different wavelengths.“

- p. 11: include a formula for the correction factor;

I think that a formula is unnecessary, because the correction factor is good explained in the text (one sentences was added): „A ratio of the circumsolar radiation to the direct solar radiation was calculated using the SMARTS 2.9.5 programme (available at <https://www.nrel.gov/rredc/smarts/>). The calculations in SMARTS were implemented for rural aerosol conditions, which are characterized by the Ångström exponent equal to 0.96. The ratio of the circumsolar radiation to the direct solar radiation was determined for all five wavelengths. This ratio is hereinafter referred to as the correction factor. The values of zenith angle, aerosol optical depth and atmospheric air pressure were taken into account in the calculation of the correction factor. It follows that this factor was determined for specific conditions at a given time.“

- p. 14: what criterium was followed to define "similar" the two ln(ETC)'s, sometimes deviating by 0.1 (l. 19)?

Related sentence was reworted: " It can be seen that ETC values show some differences."

Similarly, how can you quantitatively tell that the agreement is "excellent" (p. 15)

Related sentence was reworted: „A good agreement of both methods was seen in the first period.“
and "high level"?

Related sentence was reworded: „On the other hand, good agreement is proven by a median and standard deviation (1σ) of the differences in AOD values for the LPM and BSM with the values of -0.004 and 0.01 .“

Compare your results to the ones obtained in a recent paper about Brewer AOD in the UV range (of López-Solano et al., 2017, <https://www.atmos-chem-phys-discuss.net/acp-2017-1003/>);

- 5 **As you recommended, I have inspired by the article López-Solano et al., 2017. Instead of average of differences now I use the median of differences together with standard deviation (1σ) of differences.**

- p. 16: "it is not known that the BSM would take into account a change in the distance of Earth from the Sun". Does it mean that the Brewer Operating Software does not take into account the Earth-Sun distance or that you do not know whether it takes this factor into account? In the latter case, examine the source code or ask to the maintainer of the Brewer operating software!

Now I have confirmation from IOS (International Ozone Services) and therefore I have changed the sentence: „The first cause is that the BSM does not take into account a change in the distance of Earth from the Sun.“

- 15 - p. 18: since only direct sun measurements are considered in the paper, only report the ozone trend for DS;

It was accepted. The ozone trend and also 5-years averages was corrected only for DS measurements. Related text in the manuscript has been changed: „If only DS measurements are taken into consideration, the average value of total ozone was 326 DU in the last five years (2012–2016), which is only 1 DU more than in the first five years of the monitored period (1994–1998). The linear trend for the period of 1994–2016 has the value of 0.8 ± 2.2 DU for 10 years.“

- p. 21: explain what agricultural activities can determine the increase of AOD (wildfires?).

- Related text in the manuscript has been changed and enriched by an explanation: „Higher values of AOD in the months of April to September are presumably related to agricultural activities in the vicinity of the station. The location is windy and a bare dry soil or even plant products are often blown away.“**

Author's response to the comments of Anonymous Referee #2.

Authors comments are written in bold.

This new version of the paper shows many improvements in its organization and readability. I have some remaining
5 questions and suggestions, but they are fairly minor. Once they have been taken into consideration by the author, I believe
the paper will be suitable for publication.

1) General questions:

a) Could you insert in the manuscript a brief explanation for the approximation of the aerosol air mass presented on page 7,
10 line 30 (that is, $\mu_a \sim \mu_{H_2O}$)?

A brief explanation was added to the manuscript: „ μ_a is the air mass factor of aerosols ($\mu_a < 4$ for AOD), while due to the similar vertical profile of aerosols and water vapor it holds that $\mu_a \cong \mu_{H_2O}$, where μ_{H_2O} is the air mass factor of water vapor determined according to Kasten (1966).“

b) Could you provide some justification (a reference would suffice) for the sentence on page 8, line 2, stating that the SO₂
15 has a low impact on the AOD determination and its determination is not accurate?

A reference was added and also a little change was made in the sentence: „The contribution of sulfur dioxide was neglected, namely due to its low impact (Arola and Koskela, 2004) and due to its inaccurate determination at the Poprad-Gánovce station as well.“

c) Why do you call the algorithm presented in page 10 a “cloud screening”? It clearly removes a lot of bad measurements,
20 but are these measurements only produced by clouds?

The algorithm presented in the Figure 1 was renamed to quality control process. Related text was therefore corrected.

d) On page 16, line 14, it is stated that “it is not known if the BSM would take into account a change in the distance of Earth
25 from the Sun”. Have you asked IOS?

Now I have confirmation from IOS and therefore I have changed the sentence: „The first cause is that the BSM does not take into account a change in the distance of Earth from the Sun.“

2) Figures:

a) On Fig.4, the line used for the average ETC shows some sloped segments. However, the average ETC are constant values in every two-year period. The sloped segments should be removed, leaving a blank space if necessary. This would also help to identify the calibration periods.

5 **It was accepted.**

b) Figs. 6, 7, and 8: I suggest adding the date range above on top of the figure for easier reference.

It was accepted.

10 c) Figs. 7 and 8: the label on the vertical axes should be “AOD – CSP” and not “AOD – fotometer”

It was accepted.

3) English: please check the following

a) Page 1, line 7: instead of “presented”, it should be “present”

15 **It was accepted**

b) P.1, l.10: remove the blank space between “MK” and “IV”, as in the rest of the paper

It was accepted

c) P.7, l.6: I think “mountainous” is more usual than “montane”

It was accepted

20 d) P.7, l.10: remove “a” from “is a part”

It was accepted

e) P.7, l.13: change “size” to “value”

It was accepted

f) P.7, l.14: change “reference instrument” to “reference Brewer instrument”

25 **It was accepted**

g) P.8, l.5: change “An NOAA” to “A NOAA”

It was accepted

h) P.8, l.23: change “Eq. (2) is adjusted” to “Eq. (2) is linearized”

It was accepted

30 i) P.8, l.31: change “The inclination of obtained line a” to “The slope a of the obtained line”

It was accepted

j) P.8, l.32: change “The natural logarithm of ETC $\ln(S0, \lambda)$ ” to “The natural logarithm of the ETC, $\ln(S0, \lambda)$ ”

It was accepted

k) P.11, l.19: change “was made pursuant to the recommendations” to “was made following the recommendations”

It was accepted

l) P.14, l.5: change “as well ETCs” to “as well as the ETCs”

It was accepted

m) P.18, l.15: change “by means of an average value of ETC” to “by means of the LPM method using an average ETC value
5 for every 2-year period shown in Fig.4”

It was accepted

n) P.18, l.17: change “by analogy” to “in the same way but adding the value of the standard deviation”

It was accepted

o) P.20, l.5: change “In case of AOD” to “In the case of the AOD”

10 **It was accepted**

p) P.21, l.4: change “Figure 13 presents the same characteristics as in Fig. 12” to “Fig. 13 is the same as Fig. 12”

It was accepted

q) P.21, l.17: remove the commas enclosing “for which it was possible to determine the AOD”

It was accepted

15 r) P.22, l.21: change “Cloud screening employed” to “The cloud screening method employed”

It was accepted

s) P.23, l.5: remove “of the two compared methods”

It was accepted

t) P.23, l.7: change “longest of examined wavelengths” to “longest of the examined wavelengths”

20 **It was accepted**

u) P.23, l.9: change “application of correction” to “application of the correction”

It was accepted

v) P.23, l.14: change “The slightly” to “A slightly”

It was accepted

25

30

Comparison of the optical depth of total ozone and atmospheric aerosols in Poprad-Gánovce, Slovakia

Peter Hrabčák¹

¹Slovak Hydrometeorological Institute, Poprad-Gánovce, 058 01, Slovakia

5 Correspondence to: Peter Hrabčák (peter.hrabcak@shmu.sk)

Abstract. The amount of ultraviolet solar radiation reaching the Earth's surface is significantly affected by atmospheric ozone along with aerosols. The presented paper is focused on a comparison of the total ozone and atmospheric aerosol optical depth in the area of Poprad-Gánovce, which is situated at the altitude of 706 metres above sea level in the vicinity of the highest mountain in the Carpathian mountains. The direct solar ultraviolet radiation has been measured here continuously since ~~August 1994-1993~~ using a Brewer MK-IV ozone spectrophotometer. These measurements have been used to calculate the total amount of atmospheric ozone and, subsequently, its optical depth ~~as well~~. They have also been used to determine the atmospheric aerosol optical depth (AOD) using the Langley plot method. Results obtained by this method were verified by means of comparison with a method that is ~~a~~ part of the ~~Operational Brewer Program~~ Brewer Operating Software, as well as with measurements made by a Cimel sunphotometer. Diffuse radiation, stray-light effect and polarization corrections were applied to calculate the AOD using the Langley plot method. In this paper, two factors that substantially attenuate the flow of direct ultraviolet solar radiation to the Earth's surface are compared. The paper presents results for 23 years of measurements, namely from 1994 to 2016. Values of optical depth were determined for the wavelengths of 306.3 nm, 310 nm, 313.5 nm, 316.8 nm and 320 nm. A statistically significant decrease in the total optical depth of the atmosphere was observed with all examined wavelengths. Its root cause is the statistically significant ~~drop~~ in the total optical depth of aerosols.

1 Introduction

It is known that anthropogenic changes in the amount of total ozone and atmospheric aerosols have a considerable impact on the solar UV radiation reaching the Earth's surface (De Bock et al., 2014; Czerwińska et al., 2016). An increased transmittance of UV radiation through Earth's atmosphere has a ~~manifest~~ influence on human health and natural ecosystems. Higher doses of UV radiation have adverse effects mainly on terrestrial plants that are exposed to it on a long-term basis (Jansen et al., 1998). Exposure of human organism to excessive UV radiation doses may cause premature ageing of the skin, weakening of immune system, and damage to cells and DNA, which may consequently lead even to skin cancer and other health ~~conditions~~ (Greinert et al., 2015). Positive effects of UV radiation are known as well, which include vitamin D production in the skin in particular. This vitamin is much needed for proper functioning of a human organism (Kimlin and Schallhorn, 2004). In the past, anthropogenic impact led to the increased transmittance of solar UV radiation through Earth's

atmosphere as a result of a decrease in the total amount of ozone. Global ozone layer depletion began to show more significantly in the 1980s, reaching its peak with the value of ca 5 % (with respect to the average for the period of 1964–1980) in early 1990s. The depletion has already been reduced in recent years, amounting to ca 3 % on average for the whole Earth (WMO, 2014). In the middle latitudes of the Northern Hemisphere (35° N–60° N), the depletion of ozone layer
5 reached 3.5 % around 2010 (2008–2012). The depletion of up to 6 % was peculiar to the middle latitudes of the Southern Hemisphere (35° S–60° S) in the same period (WMO, 2014).

On the other hand, anthropogenic emission of aerosols into the atmosphere causes reduction of solar UV radiation reaching the Earth's surface, especially in industrialized areas. In the early 1990s, it was determined that solar UV-B radiation had decreased by ca 5–18 % in non-urbanized areas of industrialized countries since the Industrial Revolution as a result of air
10 pollution (Liu et al., 1991). Anthropogenic aerosols may reduce the UV radiation reaching the Earth's surface by more than 50 % even in highly polluted urban areas (Krotkov et al., 1998; Sellitto et al., 2006). Anthropogenic emissions of aerosols have been gradually reduced in the developed countries, and a drop in the aerosol optical depth (AOD) has been observed in several locations (Kazadzis et al., 2007; Mishchenko and Geogdzhayev, 2007; Alpert et al., 2012; de Meij et al., 2012; Zerefos et al., 2012). Aerosols have a substantial effect on other physical and chemical processes taking place in the
15 atmosphere as well (Seinfeld and Pandis, 2006; Raghavendra Kumar et al., 2010). They affect predominantly the chemical composition of the troposphere and, in certain cases, of the stratosphere as well, primarily with regard to major solar eruptions or flights of aircraft (Finlayson and Pitts, 2000; Seinfeld and Pandis, 2006). They can reduce the visibility (Lyamani et al., 2010) and also have a significant effect on human health in many cases (WHO, 2006). The presence of aerosols in the atmosphere has an impact on the energy balance of the Earth as well, namely directly, semi-directly and
20 indirectly (De Bock et al., 2010).

~~The direct impact means scattering and absorption of shortwave and longwave radiation. Absorption of radiation subsequently leads to warming of those atmospheric parts, where aerosols are present (primarily in the boundary layer of the atmosphere) and higher temperature consequently leads to evaporation of cloud layers. The last sentence concisely described the semidirect impact, resulting in a higher density of solar radiation flow reaching the Earth's surface (Cazorla et al., 2009).~~
25 ~~The higher temperature may lead to a change of thermal stratification of the atmosphere, which consequently affects vertical and horizontal movements of air in the atmosphere. The indirect impact pertains to the ability of aerosols to act as condensation nuclei or ice nuclei, which affects microphysical and optical properties of clouds. Yet, it concerns a change of their radiation characteristics, change of atmospheric precipitation characteristics, and alteration in cloud lifetime as well. An increase in the number of condensation nuclei leads to the rise of cloud droplet count and to the reduction of their size under~~
30 ~~the given conditions of water content in the atmosphere, causing an increase of albedo and extension of cloud lifetime (Lohmann and Feichter, 2005; Unger et al., 2009). For the said reasons,~~ the anthropogenically emitted aerosol particles considerably contribute to the ongoing global climate change, while their influence on radiation balance is still uncertain to a great extent (IPCC, 2014, and references therein).

This paper is focused primarily on the aerosol optical depth obtained by measurements made using the Brewer ozone spectrophotometer and presents one of the possible methodical approaches to its calculation. The presented method is verified by means of an alternative method of AOD calculation and is also verified by Cimel sunphotometer measurements. In the paper, obtained values of AOD are compared with the impact of total atmospheric ozone optical depth on the reduction of solar UV radiation. The Brewer spectrophotometer allows determining the optical depth in the UV region of the spectrum for the wavelengths of 306.3 nm, 310 nm, 313.5 nm, 316.8 nm and 320 nm. It was determined for the period of 1994–2016. The examined series is 23 years long, which enabled to quantify a linear trend of examined optical depth characteristics.

The employment of Brewer spectrophotometer measurements to determine the AOD **as well** has already been published in **multiple** studies. Some of them presented only short series of measurements. Therefore, they do not include multiannual trends of AOD (Carvalho and Henriques, 2000; Kirchhoff et al., 2001; Marenco et al., 2002). It is mainly more recent papers that present also multiannual measurements, but they do not always present trend information as well. It is stated in Jaroslowski et al. (2003) that the development of AOD for the Polish Belsk station in the period of 1992–2002 appears to have no trend. A quantification of the trend for the Belgian Uccle station can be found in Cheymol and De Backer (2003). It indicates a significant negative trend at the level of 2σ for all wavelengths for Brewer MKII spectrophotometer measurements in the period of 1989–2002. For example, the trend value for 320.1 nm is -2.13 ± 0.39 %/year. It also states that equally significantly negative trend was not observed for the period of 1984–2002. The trend value for the **above** wavelength was only -0.78 ± 0.42 %/year in that period. The trend is also mentioned in Kazadzis et al. (2007), namely for the Greek Thessaloniki station in the period of 1997–2005. Data analysis for the Brewer MKIII spectrophotometer adjusted for the seasonal cycle using a linear regression showed the trend of -2.9 ± 0.92 %/year for the wavelength of 320.1 nm. It was also determined by a Student's t-test that the statistical significance of this change was more than 99 %.

2 Methodology

2.1 Location of experiment

The Brewer ozone spectrophotometer (MKIV) and the Cimel sunphotometer are located on the roof of a building of the Aerological and Radiation Centre, Slovak Hydrometeorological Institute (SHMI), in Gánovce near the city of Poprad. **Their** coordinates are 49.03°N ~~lat.~~ and 20.32°E ~~long.~~ with **the** altitude of 706 m above sea level. The aerosol content in the air, both total amount and composition by types, is determined by local sources on one hand, and by atmospheric circulation on the other hand, which can move a certain air mass together with aerosols even for several thousand kilometres. In isolated instances, it may also **concern a** transport of Sahara dust from Africa. The Sahara dust was present over the Slovak Republic at least for 20 days in 2016 (Hrabčák, 2016). Major local sources include products of solid fuel combustion in adjacent municipalities and the agriculture. A bare dry soil or even plant products are often blown away by wind, as the location is rather windy. The proximity of the city of Poprad (ca 1.5 km) with the population of ca 53,000 and various industrial

activities plays a certain role as well. On the other hand, it is a submontane location, since the highest mountain of the Carpathian mountains (2,655 metres above sea level) is situated only 20 km away from the station. In spite of the proximity of the mentioned city, the area can be deemed rural in general with respect to the anthropogenic impact.

2.2 Instrumentation

- 5 The Brewer ozone spectrophotometer (No. 97, a single monochromator – model MKIV) is a scientific instrument operating in the ultraviolet and visible region of the solar spectrum. The Brewer spectrophotometer was originally designed to measure the vertical column of ozone in the atmosphere (Brewer, 1973). ~~More recent instruments (including the~~ The model MKIV) allow moreover measuring the vertical column of SO₂, NO₂ (for this purpose, measurements of solar radiation are made in the visible region of the spectrum) and global UV radiation as well (from 290 to 325 nm, with an increment of 0.5 nm). The
- 10 instrument breaks down the solar radiation reaching the Earth's surface using its optical system and selects predetermined wavelengths with higher and lower absorption of O₃ and SO₂ from the ultraviolet part of its spectrum. On the basis of different radiation absorption for the selected wavelengths, it is possible to derive the total amount of given gases in the vertical column of the atmosphere. ~~(using the DS (direct sun) or ZS (zenith sky) measurements). The spectral separation of solar radiation is carried out by means of a modified Ebert f/6 spectrometer, which uses a holographic diffraction grating with a resolution of 1,200 lines per one mm (Sci Tec, 1999). The instrument stability of Brewer spectrophotometer measurements is ±0.01 nm, namely throughout the temperature range (Sci Tec, 1999). The value of the smallest wavelength increment (microstep) is 0.006 ±0.002 nm (Sci Tec, 1999). More information about the instrument and the measurement process can be found in Kerr (2010).~~
- 15 It performs standard measurements of direct solar radiation (DS) in the UV region of the Brewer spectrophotometer at five selected wavelengths, namely 306.3 nm, 310 nm, 313.5 nm, 316.8 nm and 320 nm. They are not completely identical for individual instruments, which is given by minor differences in the position of a measurement slit and other minute mechanical and optical differences (Savastiouk and McElroy, 2005). The aforementioned five wavelengths represent their long-term average, as very small changes in their size occur over the years. ~~During the monitored 23 years, the sizes of wavelengths were refined five times in total. Their accurate sizes were used at the beginning of the examined period. In 1994, the sizes were 306.276 nm, 310.04 nm, 313.494 nm, 316.799 nm and 319.999 nm. The end of the period, that is 2015, was characterized by values of 306.276 nm, 310.035 nm, 313.476 nm, 316.755 nm and 319.989 nm. Observed deviations are only minimal. Therefore, the aforementioned long term average is generally applicable to the whole period.~~
- 20 The instrument No. 097 has undergone regular 2-year calibrations and daily tests using internal lamps (mercury and standard lamp) from the start of the measurements (18 August 1993). The calibrations are provided by International Ozone Services (IOS). The
- 25 instrument is calibrated against the World Brewer Reference Triad (World Meteorological Organization standards), maintained by Environment Canada, by means of a portable reference instrument No. 017.
- Direct solar radiation measurements can be used to determine the aerosol optical depth as well. This optical property can be determined in the ultraviolet region of the solar spectrum for the five aforementioned wavelengths, for which the

DS measurement is performed by default. It is known that the utilization of Brewer spectrophotometer to calculate the AOD encompasses particular sources of potential systematic errors. If these errors are neglected, it may lead to negative values of the Ångström exponent for the examined wavelengths (Arola and Koskela, 2004). The first source of errors is the impact of undesired diffuse radiation ~~transmittance on~~ the DS measurements. The second source is the daily cycle of ozone in urbanized areas. The third source is ~~the neglect~~ of NO₂ absorption impact. In addition, the spectral stray-light effect is the fourth source for the single monochromator. It is stated in Arola and Koskela (2004) that ~~the neglect~~ of diffuse radiation impact is potentially the biggest source of errors. On the contrary, ~~the neglect of~~ stray-light effect has the lowest impact. Unlike the AOD for the shortest and the longest wavelength, its impact in the cited paper is approximately 7-fold lower than the diffuse radiation impact. In the light of the listed errors, it is not recommended to determine extraterrestrial constants (ETCs) using the Langley plot method (LPM) for low altitude stations in urbanized areas, unless corrections for the impact of diffuse radiation and the daily cycle of ozone are known. Poprad-Gánovce is not a typical station in a low-altitude urbanized area and is not even a typical high mountain station. It is between these two cases.

The impact of daily ozone cycle and NO₂ can be neglected considering the rural location and higher altitude of the Poprad-Gánovce station. The diffuse radiation impact and stray-light effect were not neglected. A recommendation to ensure the air mass factor does not exceed the value of 3 in the calculation of ETCs was taken into account ~~as well~~ (Arola and Koskela, 2004; De Bock et al., 2010). It is important to mention that the internal polarization effect is another potential source of systematic errors as well (Cede et al., 2006). It occurs due to the combined effect of two polarization sensitive elements of the Brewer spectrophotometer. The entrance window is the first of mentioned elements and the grating is the second. Polarization effect depends on the zenith angle of the Sun and is almost wavelength independent. The polarization effect for the instrument No. 97 was not determined directly. It is assumed that the dependence of polarization effect on the zenith angle is similar for all types of Brewer spectrophotometers (Cede et al., 2006). As a result, corrections (the field measurements method) published in Cede et al. (2006) could be applied to the instrument No. 97. The impact of temperature changes was not also ~~neglected~~. The corrections resulting from the changes in temperature of the instrument are allowed for in the adjustment of raw data.

The CE 318 NE dP automatic Cimel sunphotometer was used to verify the AOD values obtained by the Brewer spectrophotometer ~~(Cimel – advanced monitoring, 2015)~~. It is an instrument that enables to measure the direct, diffused and polarized solar radiation. It performs measurements of direct solar radiation for the selected wavelengths in the ultraviolet, visible and infrared region of the spectrum (Cimel – advanced monitoring, 2015). AOD values obtained for the wavelengths of 340 nm and 380 nm were used for comparison with the measurements of Brewer spectrophotometer. The Cimel sun photometer began to perform first measurements at the Poprad-Gánovce station on 12 December 2014. If the conditions allow, it is in automatic operation every day except for the calibration period. For that reason, no continuous measurements are available. The calibration period is approximately 2 months long, and the instrument is away from the station during that time. The calibration is provided within the AERONET (AErosol RObotic NETwork) global network and is carried out by

Service National d'Observation PHOTONS/AERONET, Laboratoire d'Optique Atmosphérique, CNRS-Université de Lille. Only the level 2.0 data were used in this paper. It is the highest possible level defined within the AERONET.

2.3 Calculation of total ozone optical depth

Measured values for the total column ozone amount (TCO) obtained by the direct sun (DS) procedure were used to calculate the total ozone optical depth. A DS measurement is ~~performed-accepted~~ only ~~with the relative optical mass-airmass factor of the ozone layer~~ of less than 4 (as recommended by IOS) and it takes approximately 2.5 minutes. During that time, the density of solar radiation flow is measured five times for each of the five wavelengths. Thus five values of total ozone in Dobson units (DU) are obtained from a single DS measurement, which are consequently used to calculate an average and a standard deviation. Only the measurements that meet the standard deviation criterion ($STDEV \leq 2.5$ DU) are selected for further data analysis. The total ozone was calculated using the Brewer Spectrophotometer B Data Files Analysis Program software v. 5.0 (O3Brewer) by Martin Stanek (<http://www.o3soft.eu/o3brewer.html>). The optical depth was calculated for each accepted value of the total ozone. The calculation is represented by the following equation:

$$\tau_{\lambda, O_3} = \Omega_{O_3} \alpha(\lambda, T) = \Omega_{O_3} \sigma(\lambda, T) n, \quad (1)$$

where τ_{λ, O_3} is the total ozone optical depth, Ω_{O_3} is the total ozone in Dobson units, $\alpha(\lambda, T)$ is the absorption coefficient for ozone, $\sigma(\lambda, T)$ is the effective absorption cross-section of the ozone molecule (it is usually quantified for 1 cm^2) and n is the molecule count in the volume determined by 1 DU and 1 cm^2 ; for O_3 , it is a constant with the value of $n = 2.687 * 10^{16} \text{ cm}^{-3}$ (Schwartz and Warneck, 1995). It is recommended in Carlund et al. (2017) to utilize the same effective absorption cross-sections of the ozone molecule to calculate both the TCO and its optical depth, which is required to determine the AOD. The ~~Operational Brewer Program~~ Brewer Operating Software for the Brewer spectrophotometer utilizes the effective absorption cross-sections of the ozone molecule, which are determined on the basis of Bass and Paur measurements (Bass and Paur, 1985). This scale is employed to calculate the total amount of ozone within the network of Brewer spectrophotometers according to the recommendations of the International Ozone Commission (<http://www.esrl.noaa.gov/gmd/ozwv/dobson/papers/coeffs.html>).

Today, more recent and accurate values of effective absorption cross-sections are already available based on measurements of the Molecular Spectroscopy Lab, Institute of Environmental Physics (IUP), University of Bremen, (<http://www.iup.physik.uni-bremen.de/gruppen/molspec/databases/index.html>; Gorshelev et al., 2014; Serdyuchenko et al., 2014).

More recent values according to the IUP were used to calculate the ozone optical depth and the AOD. In order to preserve the consistency between the calculation of optical depth and the calculation of the TCO, it was necessary to calculate the TCO values according to a more recent set of IUP. This calculation was carried out as per the recommendations in Redondas et al. (2014). The dependence of effective cross-section on the temperature is very important. A so-called effective temperature for the given gas is usually used. In case of ozone measurements using the Brewer spectrophotometer, an

average standard effective temperature of -45°C (228.15 K) is defined (Redondas et al., 2014). Another difference in the calculation of the TCO compared to the ~~Operational Brewer Program~~ Brewer Operating Software pertained to the used absorption coefficients for Rayleigh scattering. Instead of coefficients used by default, the value of TCO was determined with coefficients according to Bodhaine et al. (1999), which is in line with coefficients used to calculate the optical depth.

- 5 Brewer Operating Software and O3Brewer software use the same formula to calculate an airmass factor for ozone. The same applies to Rayleigh scattering. The given formulas are defined in the Brewer MKIV Spectrophotometer Operator's Manual (Sci-Tec, 1999). Different, yet on the other hand more accurate formulas, were used to calculate the AOD by LPM – for ozone according to Komhyr (1980) and for Rayleigh scattering according to Kasten and Young (1989).

2.4 Calculation of aerosol optical depth

- 10 The ~~Langley plot method~~ (LPM) was employed to calculate the AOD. It is a traditional method employed to calculate the AOD by the Brewer spectrophotometer (Carvalho and Henriques, 2000; Kirchhoff et al., 2001; Silva and Kirchhoff 2004; Cheymol et al., 2006; Sellitto et al., (2006). This method requires stable atmospheric conditions in order to determine the ~~extraterrestrial constant~~ (ETC). It needs mainly a low variability of the total ozone and atmospheric aerosols during the day, ~~for which the ETC is determined~~. It is also necessary to avoid an impact of cloudiness on direct solar radiation (DS) measurements and to ensure a sufficient scope of zenith angles of individual DS measurements during the day, which is needed for the given method.

- For the said reasons, this method is most appropriate for lower latitudes (especially in ~~montane~~ mountainous regions near the tropics), and it has certain limitations in the middle and particularly higher latitudes (Nieke et al. 1999; Marengo 2007). This method of AOD calculation has already been applied to the examined location of Poprad-Gánovce in the past (Pribullová, 2002). An alternative method of AOD calculation, ~~developed~~ implemented by Vladimir Savastiouk, has been available since 2005 (Savastiouk and McElroy, 2005; Savastiouk, 2006; Kumharn et al., 2012). In this case, an algorithm of AOD calculation is ~~a~~ part of the ~~Operational Brewer Program~~ Brewer Operating Software. Hence, the given method will be referred to as the Brewer software method (BSM) in this paper. The fundamental difference against the previous method is that ETCs for individual wavelengths are not determined using the LPM. They are obtained during calibration of the instrument, hence once in 2 years. Their ~~size value is~~ determined during calibration based on a comparison with the portable reference Brewer instrument No. 017. ETCs for the reference instrument are determined by LPM at the Mauna Loa or Izaña observatory. Neglected changes in the sensitivity of the instrument over shorter time periods represent a disadvantage of both methods, as the ETCs are fixed for a longer period – 2 years in both cases. The LPM applied in this paper employs fixed ETCs for a 2-year intercalibration period, which is identical with the standard intercalibration period for the measurement of ozone. It is assumed that any significant service modifications to the Brewer spectrophotometer during calibration may affect both the calculation of ozone and the calculation of AOD. For that reason, the period not exceeding 2 years was used.

To calculate the AOD, it was necessary to apply the Beer–Bouguer–Lambert law:

$$S_{\lambda} = S_{0,\lambda} e^{-\mu_w \tau_{\lambda}} = S_{0,\lambda} e^{-\mu_{O_3} \tau_{\lambda,O_3} - \mu_r \tau_{\lambda,r} - \mu_a \tau_{\lambda,a}} = S_{0,\lambda} e^{-\mu_{O_3} \alpha(\lambda,T) \Omega_{O_3} - \mu_r \frac{\beta(\lambda) P}{P_{std}} - \mu_a \tau_{\lambda,a}}, \quad (2)$$

where S_{λ} is the flux density of solar radiation flow for the selected wavelength expressed by photon count per unit of time on Earth's surface, $S_{0,\lambda}$ is the flux density of solar radiation flow for the selected wavelength expressed by photon count per unit of time above Earth's atmosphere (extraterrestrial constant – ETC), τ_{λ} is the total optical depth of atmosphere, τ_{λ,O_3} is the optical depth for ozone, $\tau_{\lambda,r}$ is the optical depth for Rayleigh scattering, $\tau_{\lambda,a}$ is the optical depth for aerosols, and μ_w is the air mass factor for atmosphere as a whole. Its value was calculated as a weighted arithmetic average of individual aforementioned components, while the optical depth of a given component was the weighting factor.

Furthermore, μ_{O_3} is the airmass factor of the ozone layer determined according to Komhyr (1980), μ_r is the air mass factor for Rayleigh scattering determined according to Kasten and Young (1989), μ_a is the air mass factor of aerosols ($\mu_a \leq 4$ for AOD), while due to the similar vertical profile of aerosols and water vapor it holds that $\mu_a \cong \mu_{H_2O}$, where μ_{H_2O} is the air mass factor of water vapor determined according to Kasten (1966), assuming that $\mu_a \cong \mu_{H_2O}$. Next element $\alpha(\lambda,T)$ is the absorption coefficient for ozone, Ω_{O_3} is the total amount of ozone in Dobson units, $\beta(\lambda)$ is the normalized optical depth for Rayleigh scattering (for the standard atmospheric air pressure and the vertical column), P is the atmospheric air pressure in the location of observation (the daily average was used), and P_{std} is the standard atmospheric air pressure (101,325 Pa). The contribution of sulfur dioxide was neglected, namely due to its low impact (Arola and Koskela, 2004) and due to its inaccurate determination at the Poprad-Gánovce station as well.

The normalized optical depth for Rayleigh scattering $\beta(\lambda)$ was calculated according to Bodhaine et al. (1999). The use of coefficients according to Bodhaine et al. (1999) is in compliance with the recommendations of the NOAA (National Oceanic and Atmospheric Administration). An NOAA document (<https://www.esrl.noaa.gov/gmd/grad/neubrew/docs/RayleighInBrewer.pdf>) states that standard coefficients for the Operational Brewer Program-Brewer Operating Software must not be used to calculate the AOD. It is stated in Carlund et al. (2017) that the total amount of ozone calculated using the standard coefficients is higher compared to the use of coefficients according to Bodhaine et al. (1999).

The value of S_{λ} was obtained by adjustment of raw data (raw counts). It is essential to keep the sequence of the following steps in their adjustment. In the first step, raw counts saved in a B-file were converted to count rates. In the second step, deadtime compensation was applied. After the deadtime compensation, a correction was applied to the stray-light effect in the same order as in Garane et al. (2006). In the fourth step, a correction for the temperature dependence was applied, including a spectral transmittance correction for the utilized neutral density (ND) filter. These filters are automatically selected by the Brewer spectrophotometer with respect to the current density of the solar radiation flow. There are 5 ND filters and 5 wavelengths as well, so 25 attenuation values are needed in total. The attenuation values of given filters are determined during calibration of the instrument. The impact of polarization was compensated in the fifth step. The last sixth step involved the correction due to the impact of diffuse radiation on DS measurements. When the above criteria are applied,

five initial values of S_{λ} will be obtained from a single DS measurement, and subsequently, the AOD will be calculated from each of them. The final AOD for the given DS measurement is calculated as an arithmetic average of five values.

2.4.1 Correction for the diffuse radiation and stray-light effect

The correction of diffuse radiation impact on DS measurements was made ~~pursuant~~ following to the recommendations in Arola and Koskela (2004). The fact that the full field of view reached the value of 2.6 for the Brewer spectrophotometer was taken into consideration. A ratio of the circumsolar radiation to the direct solar radiation was calculated using the SMARTS 2.9.5 programme (available at <https://www.nrel.gov/rredc/smarts/>). The calculations in SMARTS were implemented for rural aerosol conditions, which are characterized by the Ångström exponent equal to 0.96. The ratio of the circumsolar radiation to the direct solar radiation was determined for all five wavelengths. This ratio is hereinafter referred to as the correction factor. The values of zenith angle, aerosol optical depth and atmospheric air pressure were taken into account in the calculation of the correction factor. It follows that this factor was determined for specific conditions at a given time. ~~Detection of photons by the Brewer spectrophotometer is affected by noise photons received outside the analyzed wavelength.~~ Another undesired phenomenon is the influence of noise photons (received outside the range of the analyzed wavelength) on the photon detection. The noise photons are caused by radiation that does not follow the required optical path inside the monochromator due to the scattering on the grating, mirror, or housing. This problem is called the stray-light effect and is negligible for the double monochromators. The Brewer spectrophotometer at the Poprad-Gánovce station is classified as a single monochromator. For that reason, a correction for the stray-light effect was necessary (Arola and Koskela, 2004; Garane et al., 2006). The correction was determined by analysing the spectral global UV radiation measured on cloudless days. It was feasible because the optical light path is the same for both the UV and DS measurement in the monochromator. The potential impact of a different measurement mode was neglected for the sake of simplification. In general, it is assumed that at the lowest wavelengths, i.e., below 292 nm, there is no transmittance of radiation to the Earth's surface due to the absorption in the atmosphere, and any signal measured must, therefore, be stray light. In view of simplification, it is also assumed that the value of stray light is constant for all wavelengths. The said simplifications may lead to some uncertainties when estimating the stray light effect. These uncertainties were not investigated further.

2.4.2 Calculation of extraterrestrial constants

The ETC $S_{0,\lambda}$ was determined using the Langley plot method. The Langley plot method employs multiple measurements of the direct solar radiation at various zenith angles of the Sun in the sky. Its fundamental principle is as follows: The above Eq. (2) is ~~adjusted~~ linearized by applying a natural logarithm:

$$\ln(S_\lambda) = \ln(S_{0,\lambda}) - \mu_w \tau_\lambda . \quad (3)$$

There is one equation available for every single measurement of the direct solar radiation, while μ_w and $\ln(S_\lambda)$ are the knowns, and $\ln(S_{0,\lambda})$ and τ_λ are the unknowns. A number of equations equal to the number of measurements will be acquired by applying multiple measurements at various zenith angles of the Sun. Theoretically, it is feasible to determine the unknowns already from two measurements, but for the ~~practical purposes~~ ETC determination, it is advisable to acquire as many measurements as possible. That will guarantee a higher accuracy of the result. It is essential to linearly interpolate the obtained dependence of natural logarithm of the solar radiation flow density $\ln(S_\lambda)$ on the total air mass factor of the atmosphere μ_w using the method of least squares. The ~~inclination of obtained line~~ slope a of the obtained line (from the equation of a straight line $y = ax + b$) equals τ_λ . The natural logarithm of the ETC $\ln(S_{0,\lambda})$ is obtained when x (in the equation of a straight line) equals 0 (x represents μ_w). The ETC for the given wavelength is valid for the entire intercalibration period, hence for 2 years. Its determination follows the following procedure: The ETCs for individual wavelengths can be determined solely ~~on days from DS measurements~~ that meet the three following conditions:

1. The air mass factor for the atmosphere as a whole is less than 3.
2. The AOD calculated as an average of five values within a single DS measurement for the wavelength of 320 nm is less than **0.5**.
3. The difference between the maximum and minimum value of AOD within a single DS measurement is less than 0.03.

Additionally, the ETCs are determined only for days that meet the conditions 4 to 9:

4. The number of direct solar radiation measurements is at least 50 (i.e., 10 DS measurements).
5. The difference between the maximum and minimum air mass factor for the atmosphere as a whole is greater than 1.
6. The standard deviation from the measured values of the total ozone on the given day is less than 2.5.
7. The standard deviation from the measured AOD values on the given day is less than 0.07.
8. $|\ln(S_\lambda)_i - \ln(S_\lambda)_j| < 1.75 * \frac{\sum_{i,j=1}^n |\ln(S_\lambda)_i - \ln(S_\lambda)_j|}{n}$, where $\ln(S_\lambda)_i$ is the value of $\ln(S_\lambda)$ obtained from the equation of linear interpolation after the substitution of a specific μ_w , $\ln(S_\lambda)_j$ is the actually measured value of $\ln(S_\lambda)$ at the given value of μ_w , and n is the total number of direct solar radiation measurements on the given day. The fundamental principle is to exclude measurements that have too high values of residues of the linear interpolation (greater than or equal to 1.75 times the average of the residues on the given day). The threshold value of 1.75 in the above equation was selected based on the results of an optimization test. The goal of the test was to acquire the maximum possible number of days with very good linear interpolation (the selected days had to meet the ninth condition).
9. The determination coefficient for the linear interpolation is greater than 0.98.

The conditions defined above were applied respectively. The following criterion was applied to all determined ETCs within the given intercalibration period:

$$\frac{|ETC - AVERAGE(ETCs)|}{STDEV(ETCs)} < 1.5, \quad (4)$$

where AVERAGE (ETCs) is an average of the determined ETCs and STDEV (ETCs) is a standard deviation. An average is calculated from the ETCs that meet the said criterion. This average is valid for the entire intercalibration period. The final values of ETCs used to calculate the AOD were available only after two iterations. At the beginning, ~~it was not possible to apply the second and third criterion. initial values of AOD were not available yet. For this reason, it was not possible to apply the second and third criterion and it was also not possible to take into account the effect of aerosols in the calculation of μ_w and to determine the correction for the diffuse radiation.~~ It was possible to do so on the first and second iteration. The ETC varies around its mean value with regard to the distance of Earth from the Sun. Its correction was carried out according to the recommendation of the Guide to Meteorological Instruments and Methods of Observation (WMO, 2008). The calculation of apparent elevation and apparent zenith angle of the Sun in the sky was also made as per the recommendations of the WMO (WMO, 2008). The apparent elevation of the Sun constitutes the fundamental principle in the calculation of air mass factor.

2.4.3 Quality control process

All final values of AOD have undergone a ~~cloud screening quality control process~~, which ~~also includes cloud screening. The quality control process~~ is illustrated schematically in Fig. 1. ~~Prior to the cloud screening process, all negative values of AOD were deleted.~~ The first step of ~~cloud screening is the quality control was~~ to ~~delete~~ the DS measurements with a standard deviation for $O_3 > 2.5$ DU (described in more detail in Sect. 2.3). ~~Next, all negative values of AOD were deleted.~~ In the ~~third~~~~second~~ step, all AOD values ≥ 1.5 were ~~deleted~~. Comparison of AOD values above 1.5 obtained from the Brewer spectrophotometer for the Belgian Uccle station determined that they could not be paired with Cimel sunphotometer measurements at all (De Bock et al., 2010). It proves that the limit of 1.5 is substantiated for Poprad-Gánovce station. The ~~third-fourth~~ step was to ~~delete~~ the DS measurements with a difference between the maximum and minimum AOD value greater or equal to 0.03. ~~_____~~ The international Cimel sunphotometer network AERONET uses a limit for ~~measurement triplet~~ (three measurements per minute) with the value of 0.02 (https://aeronet.gsfc.nasa.gov/new_web/Documents/Cloud_scr.pdf). We decided to apply a less stringent criterion because our case involved five measurements performed in ca 2.5 minutes. The attached diagram shows that the limit of 0.03 has caused a considerable reduction in the total number of DS measurements. Application of the second and third criterion has resulted in a reduction of the total number of measurements by up to 43 %. The DS measurements that satisfied all ~~three-four~~ previous criteria were included in a so-called daily database. A daily standard deviation of AOD (SDAOD) and a daily average of AOD (AAOD) were determined from these measurements. When SDAOD < 0.015, all AOD

values in the daily database were transferred to a final database (Good AOD). When $SDAOD \geq 0.015$ in the ~~fourth~~ fifth step, all DS measurements on the given day underwent the last ~~fifth-sixth~~ step. The following applied to the ~~fifth-sixth~~ step: When the difference between the AOD for a given DS measurement and the daily average AAOD was ≥ 0.5 , the DS measurement was not included in the final database. A total of ~~36,497~~ 37,451 DS measurements were included in the final database. That represents 57 % of the total number of measurements that entered the ~~second-third~~ step of ~~cloud screening the~~ quality control.




Figure 1: Schematic illustration of the DS measurement quality control process that includes cloud screening as well. The number of DS measurements for the period 18.8.1993–13.5.2017 that satisfy the respective criterion is indicated on the left side.

2.4.4 Additional notes

Comparison of AOD values obtained by the Cimel sunphotometer with AOD values from the Brewer spectrophotometer was not ~~executed~~ directly. First, the value of Ångström exponent was determined using pairs of wavelengths and their corresponding AOD values:

$$\alpha = -\frac{\log(\tau_{\lambda_1}/\tau_{\lambda_2})}{\log(\lambda_1/\lambda_2)}, \quad (5)$$

where $\lambda_1 = 340$ nm and $\lambda_2 = 380$ nm. On the basis of Ångström exponent and the known value of AOD (for the shorter wavelength), an AOD value for the wavelength of 320 nm was determined in the next step. This calculation was carried out

for all individual measurements. The main goal of this calculation was to acquire a comparison of AOD for the same wavelength, which is more relevant than a direct comparison of AOD for two different wavelengths.

The following procedure was employed to calculate individual characteristics of the total ozone and AOD optical depth: Daily averages were calculated as an arithmetic average of all values for the given day (minimum of one value). Monthly averages were calculated as an arithmetic average of such days, for which an average daily value was available. Annual averages were calculated as an arithmetic average of individual monthly values. The longest period without data is 12 days for the ozone optical depth and 26 days for AOD. There was no month without data. In both cases, February 2013 was the month with the absolute lowest number of days with the determined optical depth. In the first case, it was 10 days, and it was 3 days in the second case. With regard to the long-term average, December is the month with the lowest number of days with the determined AOD with 13 days. The opposite extreme falls on August with 24 days. A linear trend was calculated by means of a linear regression using the least squares method. An autocorrelation was not confirmed. Therefore, it was possible to determine the linear trend. The uncertainty of the linear trend is defined by the standard deviation ($\pm\sigma$) of the slope of the obtained linear dependence. The value of linear trend and the standard deviation was determined for a period of 10 years. The seasonal cycle was eliminated by annual averages.

3 Results and discussion

3.1 Correction for the diffuse radiation, stray-light effect and polarization

~~The results of calculation and application of individual correction factors are presented in the introduction to this part of the paper. A factor that eliminates the influence of diffuse radiation is the first of these factors. The correction of diffuse radiation impact on DS measurements was made pursuant to the recommendations in Arola and Koskela (2004). The fact that the full field of view reached the value of 2.6 for the Brewer spectrophotometer was taken into consideration. A ratio of the circumsolar radiation to the direct solar radiation was calculated using the SMARTS 2.9.5 programme (available at <https://www.nrel.gov/rrecc/smart/>). The calculations in SMARTS were implemented for rural aerosol conditions, which are characterized by the Ångström exponent equal to 0.96. The ratio of the circumsolar radiation to the direct solar radiation was determined for all five wavelengths. The values of zenith angle, aerosol optical depth and atmospheric air pressure were taken into account in the calculation of the correction factor. It follows that this factor was determined for specific conditions at a given time.~~ In Fig. 2, there is a demonstration of correction factor values for five wavelengths and the selected sequence of zenith angles of the Sun. This demonstration characterizes the conditions that are close to the ones that may normally occur at the examined station. It can be seen that the value of the factor is primarily dependent on the zenith angle. It is also true that the value of the factor is inversely proportional to the size of a wavelength. The factor for the longest wavelength of 320 nm is by ca 7 % lower than the factor for the shortest wavelength of 306.3 nm.




Figure 2: Value of correction factor for the elimination of diffuse radiation impact for the selected zenith angles of the Sun, for five wavelengths for the conditions of AOD value of 0.3 (for 320 nm) and air pressure of 930 hPa.

- 5 Another correction factor was determined to eliminate the stray-light effect. ~~Detection of photons by the Brewer spectrophotometer is affected by noise photons received outside the analyzed wavelength. The noise photons are caused by radiation that does not follow the required optical path inside the monochromator due to the scattering on the grating, mirror, or housing. This problem is called the stray light effect and is negligible for the double monochromators. The Brewer spectrophotometer at the Poprad Gánovec station is classified as a single monochromator. For that reason, a correction for~~
- 10 ~~the stray light effect was necessary (Arola and Koskela, 2004; Garane et al., 2006). The correction was determined by analysing the spectral global UV radiation measured on cloudless days. It was feasible because the optical light path is the same for both the UV and DS measurement in the monochromator. In general, it is assumed that at the lowest wavelengths, i.e., below 292 nm, there is no transmittance of radiation to the Earth's surface due to the absorption in the atmosphere, and any signal measured must, therefore, be stray light. It is also assumed that the value of stray light is constant for all~~
- 15 ~~wavelengths.~~
- Hence, a ratio of average count rates for four wavelengths in the region from 290 to 291.5 nm to the count rates for the monitored wavelength (one out of five) was determined for 3,386 spectral analyses, in total, as well as for various zenith angles of the Sun within them. The value of determined correction factor is a function of the zenith angle. This dependence was described for the monitored wavelengths by a polynomial of the fourth degree. A demonstration is presented in Fig. 3.
- 20 As a general rule, lower wavelengths are characterized by higher values of the correction factor. The observed difference between the wavelength of 306.3 nm and 320 nm was approximately 10-fold. Furthermore, it was determined that the dependence on the zenith angle was not directly proportional for all monitored wavelengths. The direct proportion was seen only with the three shortest wavelengths.







Figure 3: Size of the correction factor for the elimination of stray-light effect depending on the apparent zenith angle (a) for the wavelength of 306.3 nm (on the left); (b) and for the wavelength of 320 nm (on the right).

The impact of individual correction factors was tested on a shorter period, namely 2014. When the stray-light effect was allowed for, a drop in the difference in the AOD for the wavelengths of 306.3 nm and 320 nm was seen. Before the correction, the difference reached the value of -0.029 , and after the correction, it was decreased by ca 0.0076 . This value is similar to the results in Arola and Koskela (2004). A correction eliminating the polarization of radiation followed after the correction for the stray-light effect. A drop in the difference with the value of 0.008 was observed again. It was then followed by a correction for the impact of diffuse radiation. A drop in the difference was observed again, but it was lower, and its value was only 0.0004 . This result is significantly lower compared to the result presented in Arola and Koskela (2004). In conclusion, it can be stated that application of all three corrections resulted in the drop in the difference in AOD for the shortest and the longest wavelength by 0.015 to the final value of -0.014 . The negative value of Ångström exponent for the given pair of wavelengths was reduced, but it still persists.

3.2 Extraterrestrial constants

Strict conditions for the selection of appropriate days for the determination of ETCs and other mentioned criteria significantly eliminate obtaining of non-representative values. To calculate the ETC characterizing the entire 2-year period, 157 values of individual ETCs were employed with respect to the long-term average. This number is the same in case of all wavelengths. If the conditions were less strict, there would have been more days, for which it was possible to determine the ETC. On the other hand, the spread of determined ETCs would be wider, which would have a negative effect on the required accuracy. Therefore, the chosen criteria represent an optimum compromise. The inaccuracy of ETC determination has a direct impact on the final values of AOD. On one hand, the root cause of such inaccuracy is weather influences, which considerably eliminate the number of days that are suitable for the determination of ETC. On the other hand, additional factors may include instrument instability.





Figure 4: Trend of ETCs values for the wavelength of 320 nm during 12 intercalibration periods, 1994-2016 18.8.1993-13.5.2017.

The trend of ETCs (their natural logarithm) for the wavelength of 320 nm is illustrated in Fig. 4. It shows ETC values that characterize the entire 2-year intercalibration period, as well as the ETCs determined on individual days in the given period. The ETC for the monitored wavelength did not remain constant during the monitored period of 23 years from 18 August 1993 to 13 May 2017. Particular changes occurred in case of all five examined wavelengths. A total of 12 ETC values were determined for every single wavelength, 1 for each intercalibration period. The highest instability was observed in case of the longest wavelength. The value of its variation coefficient is 158.12 %. By contrast, the lowest instability was observed for the wavelength of 3103.5 nm. The value of its variation coefficient is 115.62 %. In Fig. 4, you can notice that a more significant change in the value of ETC for the longest wavelength occurred four three times in total. The first change occurred between the first and second intercalibration period. There is no specific explanation for this change. It is presumably related to the instability of ETCs during the first intercalibration period. A shorter intercalibration period might have also caused the mentioned change. It resulted in lower number of days that were suitable for the determination of ETC. Additional changes occurred between the fifth, sixth, seventh and eighth intercalibration period. They were caused by issues with the instrument. For that reason, a secondary power supply board had to be replaced in January 2005. In February 2007, a micrometer was replaced, and during calibration in May 2007, an optical filter No. 3 was replaced and a BM-E80 high-frequency source was repaired as well.

Figure 5 presents a comparison of ETC values for individual wavelengths for the LPM and BSM in the last two intercalibration periods. It can be seen that the ETC values are similar show some differences. The first period is characterized by the fact that ETCs for all five wavelengths for the LPM exceed the values for the BSM. It is exactly the reverse in the second period. In case of both methods, there was a decrease in ETCs between the given two periods. The decrease for the BSM was more marked.






Figure 5: Comparison of ETC values for individual wavelengths for the LPM and BSM ~~in the period from 25 May 2013 to 19 May 2015 (on the left) and in the period from 20 May 2015 to 31 December 2016 (on the right).~~

5 3.3 Comparison of AOD values obtained by LPM, BSM and the Cimel sunphotometer

The ~~plausibility of~~ results for the presented methodology of AOD calculation by means of the Langley plot method (LPM) was validated in two ways. In the first place, it was confronted with the results acquired by means of the Brewer software method (BSM). The given comparison is illustrated in Fig. 6. As has already been mentioned above, the BSM has been available since 2005. Its ~~output~~ has been plausible for the Poprad-Gánovce station only since calibration in 2013. Figure 6 consists of two graphs. The graph ~~on the left side (a)~~ represents a comparison of AOD for the wavelength of 320 nm for the intercalibration period from 25 May 2013 to 19 May 2015. The graph ~~on the right side (b)~~ covers the comparison of results for the same wavelength in the next intercalibration period from 20 May 2015 to the end of 2016. This division is necessary to point out the impact of used ETCs on the obtained results. In both intercalibration periods, only the measurements that had undergone the ~~cloud screening quality control process~~ were selected. The ~~mutual~~ difference of AOD ~~at in the absolute value~~ did not exceed the value of 0.1 ~~(so-called large random error) in any of monitored periods. the first period. In the second period, it happened 133 times.~~ The total number of DS measurements in both intercalibration periods was 5,7889. A total of 3,246 DS measurements were compared in the first period and 2,5423 in the second period. ~~An excellent good~~ agreement of both methods was seen in the first period. On one hand, it is proven by very high values of determination coefficient and correlation coefficient. On the other hand, ~~the high level of good~~ agreement is proven by ~~an average a median and standard deviation (1σ) of the~~ differences in AOD values for the LPM and BSM ~~that reached 0.004 with the values of -0.004 and 0.01.~~ The following intercalibration period is also characterized by very high values of determination coefficient and correlation coefficient, yet they are somewhat lower than in the first period. A more significant disagreement ~~was seen in case of mutual difference in AOD between the LPM and BSM that reached 0.054 is proven by the median with the value of -0.06 and the standard deviation with the value of 0.02.~~ The said differences between the LPM and BSM are not constant

within the intercalibration period, as certain variability has been observed for them. The biggest differences within a year are observed in summer months.

Results for the remaining four wavelengths were compared in a similar fashion as well. For the wavelengths of 310 nm, 313.5 nm, 316.8 nm and 320 nm, the value of correlation coefficient reached at least 0.99. This value was not reached only in case of the lowest wavelength of 306.3 nm in the first intercalibration period, where it was at the level of 0.97. The ~~average-median~~ of ~~mutual~~ differences in AOD between the LPM and BSM reached the absolute value of maximum 0.0654 in the already mentioned case for the wavelength of 320 nm. For the lower wavelengths, the ~~average-of-differences-median~~ in the first period was always positive (0.0364 on average). On the contrary, it was always negative in the second period (-0.0328 on average). The average of standard deviations for the given four wavelengths was 0.024 in the first period and 0.017 in the second period. Comparison of the AOD value and wavelength shows no unambiguous dependence for neither of the two methods. It applies to the LPM that the highest AOD was reached for the wavelength of 316.8 nm and the lowest AOD was reached for the wavelength of 320 nm with regard to the long-term average in the monitored period. It applies to the BSM that the highest AOD was reached for the lowest wavelength of 306.3 nm and the lowest AOD was reached for the wavelength of 310 nm with regard to the long-term average. Pribullová (2002) does not mention the unambiguous dependence of AOD on the wavelength as well. It indicates the lowest AOD in case of the lowest wavelength and presents the highest values of AOD for the wavelength of 310 nm.

It follows that the LPM and BSM correlate nicely. On the other hand, ~~random-observed~~ differences constitute a certain problem. They result in AOD values for the LPM that are lower or higher compared to the AOD values for the BSM. There are several causes of differences. ~~It is not known that the BSM would~~ The first cause is that the BSM does not take into account ~~a~~ change in the distance of Earth from the Sun. The presented LPM takes the given change into account. Another ~~distinction~~ between the two methods is the fact that the BSM takes into consideration the total measured amount of SO₂ in the calculation of AOD. The presented LPM neglects the impact of SO₂. It may seem at a glance that the BSM is thus advantaged. However, the reverse is true. The Brewer spectrophotometer at the Poprad-Gánovce station fails to measure the total amount of SO₂ accurately. Even negative values are seen often. ~~The disadvantage of BSM is also the absence of the SL~~ (standard lamp) test that leads to inaccurate TCO values, which consequently affects the size of AOD. Another drawback of the BSM is the fact that it does not take into consideration corrections for the diffuse radiation, polarization and stray-light effect. ~~The Values-values~~ of ~~the~~ determined ETCs affect the observed differences ~~probably~~ the most. However, it is true only if their difference between LPM and BSM is sufficiently large. These constants are fixed for both methods during the entire 2-year intercalibration period. An advantage of the presented LPM is that it partially takes into account the potential change ~~of ETCs in the sensitivity of the instrument~~ during the intercalibration period, because ETCs are determined based on real measurements in a given period. The BSM utilizes ETCs determined during calibration. ~~They are valid for 2 years and a potential change during the intercalibration period is not taken into account.~~ Therefore, the potential change in the sensitivity of the instrument during the intercalibration period is not taken into account at all.




Figure 6: Comparison of AOD values for the wavelength of 320 nm obtained by means of the LPM and BSM ~~in the period from~~
~~25 May 2013 to 19 May 2015 (on the left) and in the period from 20 May 2015 to 31 December 2016 (on the right).~~

Both presented methods were further compared with Cimel sunphotometer (CSP) measurements. These comparisons are illustrated in Fig. 7 (for the LPM) and Fig. 8 (for the BSM). The figures consist of two graphs just like in the previous case. In both intercalibration periods, only the DS measurements that had undergone the ~~cloud screening quality control process~~ were selected. The comparison was made as follows: A CSP measurement was selected for an individual DS measurement from the Brewer spectrophotometer with the maximum allowed difference of 5 minutes between the two measurements. If there were several suitable measurements, the nearest one was selected. In case of comparison between the LPM and the CSP, there was no difference in AOD with the absolute value greater than 0.1. On the contrary, in case of comparison between the BSM and the CSP, there were as much as 22 ~~12~~ instances of such DS measurements in total. A total of 199 measurements were compared in the first intercalibration period and 1,11 ~~56~~ in the second period.

It is apparent from the results of the comparison that both methods correlate very well with the CSP measurements. Yet the LPM shows higher values of the correlation coefficient in both intercalibration periods than the BSM. It was determined in the calculation of ~~mutual~~ differences that both methods matched the CSP measurements ~~very~~ well in the first intercalibration period. The ~~median of~~ differences between ~~the~~ LPM and ~~the~~ CSP was ~~-0.0162~~, and the ~~median of~~ differences between ~~the~~

BSM and ~~the~~ CSP was -0.0094 . The standard deviation was 0.011 and 0018. In the following period, a good match was observed only for LPM. In the second monitored period, ~~t~~the average median of differences between the LPM and the CSP reached the value of 0.0132, and the average median of differences between ~~the~~ BSM and ~~the~~ CSP reached the value of 0.0758. In this case, the standard deviation was 0.013 and 0.025. The mentioned ~~These~~ average differences are the primary reason for observed offsets in attached graphs. As a result of the offset, the intersection of the linear fit is not the same as the intersection of the main axes of the graph. It is illustrated the best by the graph ~~(b) on the right~~ in Fig. 8, because in this case, the average median of differences has the highest absolute value of all presented comparisons. In the light of the results, it can be stated that the LPM is the more reliable of the two methods.





Figure 7: Comparison of AOD values for the wavelength of 320 nm obtained by means of the LPM and the Cimel sunphotometer (CSP) in the period from 12 December 2014 to 19 May 2015 (on the left) and in the period from 20 May 2015 to 31 December 2016 (on the right).



5 **Figure 8: Comparison of AOD values for the wavelength of 320 nm obtained by means of the BSM and the Cimel sunphotometer (CSP)-in the period from 12 December 2014 to 19 May 2015 (on the left) and in the period from 20 May 2015 to 31 December 2016 (on the right).**

3.4 Total ozone optical depth and AOD

- 10 Measurements of total ozone from the Brewer spectrophotometer for the Poprad-Gánovce station are available from 1994. ~~Over the last five years (2012–2016), the average value was 330 DU, which is 5 DU more than in the first five years of the monitored period (1994–1998). If only DS measurements are taken into consideration, the average value of total ozone was 326 DU in the last five years (2012–2016), which is only 1 DU more than in the first five years of the monitored period (1994–1998).~~ The linear trend for the period of 1994–2016 has the value of 2.6 ± 2.3 DU for 10 years. ~~If only DS~~
- 15 ~~measurements are taken into consideration, the value of the trend will be 0.8 ± 2.2 DU for 10 years. The total ozone optical depth was determined only from the DS measurements.~~ For that reason, the rising trend of total ozone optical depth for the monitored wavelengths in the period of 1994–2016 was equally statistically insignificant. For the wavelength of 306.3 nm,

the value of the trend is 0.005 ± 0.009 for 10 years, and the value of the trend for the wavelength of 320 nm is 0.001 ± 0.002 for 10 years.

Figure 9 shows annual averages of AOD for the wavelength of 320 nm together with the uncertainty of their determination. The annual averages of AOD were calculated in a standard manner, i.e., by means of [the LPM method using](#) an average ~~value of ETC~~ [ETC value for every 2-year period shown in Fig.4.](#) The lower limit of uncertainty was calculated by means of an average value of ETC, from which its standard deviation for the given intercalibration period was deducted. The upper limit of uncertainty was determined ~~by analogy in the same way but adding the value of the standard deviation~~. The range of uncertainty interval depends primarily on **suitable** weather conditions in the given intercalibration period, as well as on the stability and homogeneity of measurements on days when it was possible to determine the ETC. The number of days, when it was possible to determine the ETC, plays its role too. [With regard to the long-term average, the number of these days was 15. The lowest number was 7 and the highest number was 24.](#)~~For instance, there were only two measurements in the first intercalibration period (it covers 1994 and a smaller part of 1995), and for that reason, inter alia, the uncertainty interval for 1994 is narrow and has a very low relevance. In other intercalibration periods, there were at least 8 ETCs. Therefore, the following reliability intervals can be deemed trustworthy.~~





Figure 9: Annual averages of AOD and their uncertainty for the wavelength of 320 nm, 1994–2016.

Figure 10 shows a comparison of annual averages for the total ozone and aerosol optical depth for the selected wavelengths. It is obvious already at a glance that the AOD, unlike the total ozone optical depth, exhibits an apparent decline for the monitored period. For the wavelength of 306.3 nm, the value of the trend is -0.067 ± 0.01 for 10 years, and the value of the trend for the wavelength of 320 nm is -0.056 ± 0.01 for 10 years. ~~If we omit first 2 years due to their lower reliability, then the value of the trend for the wavelength of 306.3 nm is -0.05 ± 0.01 for 10 years, and the value of the trend for the wavelength of 320 nm is -0.04 ± 0.01 for 10 years.~~ In view of the above, a conclusion can be made that the transmittance of the atmosphere in the region of examined wavelengths has increased in the area of Poprad-Gánovce over the last 23 or 21 years. The total optical depth of the atmosphere for the wavelength of 306.3 nm has a trend with the value of -0.076 ± 0.01 for 10 years, and its trend for the wavelength of 320 nm has the value of -0.056 ± 0.01 for 10 years. ~~If the first 2 years are omitted, the value of the trend for the wavelength of 306.3 nm is -0.06 ± 0.01 for 10 years, and the value of the trend for the wavelength of 320 nm is -0.04 ± 0.01 for 10 years.~~ The negative trend has been caused particularly by the drop in the AOD.




Figure 10: Comparison of values of average annual optical depth for the selected wavelengths for ozone and aerosols, 1994–2016.

Figure 11 presents a comparison of multiannual averages for the ozone optical depth, optical depth of Rayleigh scattering and AOD for all five examined wavelengths. It can be seen that ozone is dominant only in case of the shortest wavelength. It is exactly the reverse for the pair of the longest wavelengths, and ozone has the lowest impact among the examined factors. Rayleigh scattering has a dominant position in attenuation of direct solar radiation for all wavelengths except for the shortest one. In the case of the AOD, no unambiguous dependence of its size on the wavelength was observed. With respect to the long-term average, the AOD reached the highest value for the wavelengths of 310 nm and 316.8 nm, namely 0.32. On the contrary, the AOD reached the lowest value for the wavelengths of 306.3 nm and 320 nm, namely 0.289. The middle wavelengths of 313.5 nm and 320 nm are characterized by the value of 0.293. With regard to the long-term average for 23 years, the difference in AOD for the shortest and the longest wavelengths is only -0.004. For the test year of 2014, this difference was -0.014. In conclusion, it can be stated that values of ETC have probably the greatest impact on the observed difference, while the value of the observed difference in AOD is consequently derived from them. It is also substantiated by Fig. 10. It can be seen that the AOD is greater for the wavelength of 306.3 nm than for the wavelength of 320 nm in certain years, while it is the reverse in other years.




Figure 11: Comparison of long-term averages for the total ozone optical depth, the optical depth of Rayleigh scattering and AOD, 1994–2016.

Figure 12 illustrates an annual cycle of total ozone optical depth and its variability for individual months for the wavelength of 320 nm. The optical depth reaches its maximum in April with the value of 0.263 and the minimum in October with the value of 0.2032. The highest variability is attributed to the month of February and the lowest one to the month of July. It applies to all the months of the year that the variability is considerably lower compared to the average. The month of February has the maximum variation coefficient amounting to 5.9 %. All the aforementioned characteristics of the total ozone optical depth depend only on the TCO value. It is peculiar to the central European location of the station that the annual maximum of the TCO occurs in April and the annual minimum occurs in October. It is also confirmed by the measurements from the nearby station in Hradec Králové (Vaníček et al., 2012). Therefore, it can be stated that the observed annual course of the total ozone optical depth is typical for the central European location of the station.




Figure 12: Average monthly characteristics of the total ozone optical depth for the wavelength of 320 nm, 1994–2016.

Figure 13 presents the same characteristics as in Fig. 12, but they refer to the AOD in this case. The annual cycle of monthly averages is characterized by two peaks. The primary peak, that is the annual maximum, occurs in the month of August with the value of 0.41. The secondary peak occurs in the month of April with the value of 0.4 a slightly lower value.

The minimum is in the month of December with the value of 0.16. The variability is also characterized by a two-peak annual cycle. The primary and secondary peaks are in the same months as in case of the average. The annual minimum occurs in the month of November. The variation coefficient reached significantly higher values compared to ozone. The minimum is attributed to the month of May amounting to 179.1 %. The maximum is attributed to the month of December amounting to 42.42 %. The location of the station at a higher altitude in the submontane area has the primary impact on the said characteristics, which explains lower values of AOD in winter months. Higher values of AOD in ~~spring and summer months~~ the months of April to September are presumably related to agricultural activities in the vicinity of the station. The location is windy and a bare dry soil or even plant products are often blown away.




Figure 13: Average monthly characteristics of the AOD for the wavelength of 320 nm, 1994–2016.

Figure 14 shows the relative distribution of daily AOD averages for individual months for the monitored period of 1994–2016. It is important to point out that only the days, for which it was possible to determine the AOD, were taken into consideration. In the month of December, there is the highest percentage of days with AOD within the interval from 0 to 0.1, specifically up to 419 %. On the contrary, the lowest percentage was seen in the month of ~~August-June~~ with the value of only 1 %. Days with a daily average of AOD above 0.3 dominate in the months of April to August. Days with a daily average of

AOD within an interval below 0.3 dominate in the remaining months of the year. The AOD above 0.6 occurs the most often in the month of August, that is in 18 % of days. The lowest percentage of such days is in the month of November and December, namely only 1 %. In conclusion, it can be stated that the presented percentages of individual days correlate very well with the annual cycle of AOD in Fig. 13.




Figure 14: Relative distribution of daily average values of AOD for the wavelength of 320 nm, 1994–2016.

4 Conclusions

10 The total ozone and aerosol optical depth were determined by the Brewer ozone spectrophotometer in Gánovce near Poprad. The AOD was calculated by means of ETCs obtained using the Langley plot method. The analysed data of the direct solar radiation were available from 1994 to 2016, which is a 23-year series of measurements. In the 1990s, the ETCs were not

determined during calibrations. Therefore, the LPM is the only ~~practicable~~ method for the determination of AOD for the whole series of measurements. The ETCs used in this paper were determined for ~~the~~ 2-year intercalibration period. Use of such a long time period has both ~~its~~ advantage and ~~disadvantage~~. The advantage lies in the fact that the ETC determined as an average of a higher number of days prevents the accidental impact of fluctuations that have no particular explanation.

5 The disadvantage lies in the suppression of weather influences (especially the change of temperate) and modifications of the instrument in time intervals of less than 2 years. In addition to the methodology of ETC determination, the employed cloud screening affects the values of AOD as well. If the cloud screening was not employed, the values of AOD would have been distinctly higher. ~~The Cloud-cloud~~ screening ~~method~~ employed in this paper is relatively simple. Hence, it can be argued that the actual AOD is even somewhat lower. It pertains particularly to the presented monthly and annual averages. In the future, 10 the cloud screening methodology is planned to be improved. As a result, it will be possible to obtain more representative values of AOD.

On the basis of a comparison of the Brewer spectrophotometer measurements using the Langley plot method and the Brewer software method, which is a part of the ~~Operational Brewer Program~~ ~~Brewer Operating Software~~, with Cimel sunphotometer measurements, it was determined that the LPM achieved better results ~~of the two compared methods~~. It was also determined 15 that the application of corrections to the diffuse radiation, stray-light effect and polarization decreased the difference between the value of AOD for the shortest and longest of ~~the~~ examined wavelengths. Although the difference has been reduced, it still reaches negative values, which results in the negative values of the Ångström exponent as well. With regard to the long-term average for 23 years, the difference in AOD is only ~~-0.0064~~. The application of ~~the~~ correction for the polarization had the most significant influence on the reduction of the given difference. On the contrary, the lowest contribution was achieved by 20 the correction with regard to the diffuse radiation. The key factor influencing the value of the examined difference is probably the size and accuracy of ETC determination.

The obtained results clearly show the decrease in average annual values of the total optical depth of the atmosphere for the monitored wavelengths from 1994 to 2016. ~~The A statistically insignificant~~ slightly upward trend of the total ozone has been observed, which is exhibited by a statistically insignificant increase in the total ozone optical depth. This insignificant 25 increase has only a ~~minimum-minimal~~ effect on the trend of the total optical depth of the atmosphere. The root cause of the decrease in the total optical depth of the atmosphere is the statistically significant ~~drop~~ in the AOD. It follows that the transmittance of the atmosphere in the UV region of the spectrum from 306.3 nm to 320.1 nm has increased in the location of Poprad-Gánovce.

30 *Data availability*

The data used in the research and presented in this paper are available from the author upon agreement. It is recommended to send a request by email.

Competing interests. The author declares that he has no competing interests.

Acknowledgements. The author of the paper thanks the Slovak Hydrometeorological Institute for the provision of data. A special acknowledgement goes to Anna Pribullová for running the station of the Aerological and Radiation Centre, SHMI, in Poprad-Gánovce. The author also thanks the Faculty of Mathematics, Physics and Informatics of Comenius University in Bratislava for the possibility of external PhD study in meteorology and climatology, during which this paper was prepared. Many thanks also to the organizing team of international Quadrennial Ozone Symposium held in Edinburgh on 4–9 September 2016, namely, inter alia, for the provision of financial support. Finally, the author thanks the European Aerosols, Clouds, and Trace gases Research Infrastructure (ACTRIS) for providing the calibration of the Cimel sunphotometer.



ACTRIS-2 project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 654109.

10

References

- Alpert, P., Shvainshtein, O., and Kishcha, P.: AOD trends over megacities based on space monitoring using MODIS and MISR, *Am. J. Clim. Change*, 12, 117–131, doi:10.4236/ajcc.2012.13010, 2012.
- Arola, A., and Koskela, T.: On the sources of bias in aerosol optical depth retrieval in the UV range, *J. Geophys. Res.*, 109, D08209, doi:10.1029/2003JD004375, 2004.
- Bass, A. M. and Paur, R. J.: The ultraviolet cross-sections of ozone. I. The measurements, II. Results and temperature dependence, in: *Atmospheric ozone; Proceedings of the Quadrennial ozone symposium*, 3–7 September 1984, Halkidiki, Greece, 606–616, 1985.
- Bodhaine, B. A., Wood, N. B., Dutton, E. G., and Slusser, J. R.: On Rayleigh Optical Depth Calculations, *J. Atmos. Oceanic Technol.*, 16, 1854–1861, 1999.
- Brewer, A.: A replacement for the Dobson spectrophotometer?, *Pure and Applied Geophysics*, 106, 919–927, 1973.
- Carlund, T., Kouremeti, N., Kazadzis, S., and Gröbner, J.: Aerosol optical depth determination in the UV using a four-channel precision filter radiometer, *Atmos. Meas. Tech.*, 10, 905–923, <https://doi.org/10.5194/amt-10-905-2017>, 2017.
- Carvalho, F., and Henriques, D.: Use of Brewer ozone spectrophotometer for aerosol optical depth measurements on ultraviolet region, *Adv. Space Res.*, 25, 997–1006, 2000.
- ~~Cazorla, A., Shields, J. E., Karr, M. E., Olmo, F. J., Burden, A., and Alados-Arboledas, L.: Technical Note: Determination of aerosol optical properties by a calibrated sky imager, *Atmos. Chem. Phys.*, 9, 6417–6427, doi:10.5194/acp-9-6417-2009, 2009.~~
- Cede, A., Kazadzis, S., Kowalewski, M., Bais, A., Kouremeti, N., Blumthaler, M., and Herman J.:

- Correction of direct irradiance measurements of Brewer spectrophotometers due to the effect of internal polarization, *Geophys. Res. Lett.*, 33, L02806, doi:10.1029/2005GL024860, 2006.
- Cheyamol, A. and De Backer, H.: Retrieval of the aerosol optical depth in the UV-B at Uccle from Brewer ozone measurements over a long time period 1984-2002. *J. Geophys. Res.*, 108, 4800, doi:10.1029/2003JD003758, 2003.
- 5 Cheymol, A., De Backer, H., Josefsson, W., and Stubli, R.: Comparison and validation of the aerosol optical depth obtained with the Langley plot method in the UV-B from Brewer Ozone Spectrophotometer measurements, *J. Geophys. Res.*, 111, D16202, doi:10.1029/2006JD007131, 2006.
- Cimel – advanced monitoring: CE318 Photometer User's Operation Manual, 2015.
- Czerwińska, A. E., Krzyścin, J. W., Jarosławski, J., and Posyniak, M.: Effects of urban agglomeration on surface-UV doses: a comparison of Brewer measurements in Warsaw and Belsk, Poland, for the period 2013–2015, *Atmos. Chem. Phys.*, 16, 13641–13651, doi:10.5194/acp-16-13641-2016, 2016.
- De Bock, V., De Backer, H., Mangold, A., and Delcloo, A.: Aerosol Optical Depth measurements at 340 nm with a Brewer spectrophotometer and comparison with Cimel sunphotometer observations at Uccle, Belgium, *Atmos. Meas. Tech.*, 3, 1577–1588, doi:10.5194/amt-3-1577-2010, 2010.
- 15 De Bock, V., De Backer, H., Van Malderen, R., Mangold, A., and Delcloo, A.: Relations between erythemal UV dose, global solar radiation, total ozone column and aerosol optical depth at Uccle, Belgium, *Atmos. Chem. Phys.*, 14, 12251–12270, doi:10.5194/acp-14-12251-2014, 2014.
- de Meij, A., Pozzer, A., and Lelieveld, J.: Trend analysis in aerosol optical depths and pollutant emission estimates between 2000 and 2009, *Atmos. Environ.*, 51, 75–85, doi:10.1016/j.atmosenv.2012.01.059, 2012.
- 20 Finlayson-Pitts, B. J., Pitts, J. N.: *Chemistry of the upper and lower atmosphere: theory, experiments, and applications*, Academic Press, San Diego, USA, 2000.
- Garane, K., Bais, A. F., Kazadzis, S., Kazantzidis, A. and Meleti, C.: Monitoring of UV spectral irradiance at Thessaloniki (1990–2005): Data re-evaluation and quality control, *Ann. Geophys.*, 24, 3215–3228, 2006.
- Gorshelev, V., Serdyuchenko, A., Weber, M., Chehade, W., and Burrows, J. P.: High spectral resolution ozone absorption cross-sections – Part 1: Measurements, data analysis and comparison with previous measurements around 293 K, *Atmos. Meas. Tech.*, 7, 609–624, 2014.
- 25 Greinert, R., de Vries, E., Erdmann, F., Espina, C., Auvinen, A., Kesminiene, A., and Schuz, J.: *European Code against Cancer 4th edition: Ultraviolet radiation and cancer*, *Cancer Epidem. Biomar.*, 39, S75–S83, 2015.
- Hrabčák, P.: Saharan dust over Slovakia, *Meteorological Journal*, 19/2, 83–91, 2016.
- 30 IPCC: *Climate change 2013: the physical science basis. Working Group I contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Tech. rep., Intergovernmental Panel on Climate Change, 2014.
- Jansen, M. A., Gaba, V., and Greenberg, B. M.: Higher plants and UV-B radiation: balancing damage, repair and acclimation, *Trends in plant science*, 3(4), 131–135, 1998.
- Jaroslawski, J., J. W. Krzyścin, S. Puchalski, and Sobolewski, P.: On the optical thickness in the UV range: Analysis of the

- ground-based data taken at Belsk, Poland, *J. Geophys. Res.*, 108(D23), 4722, doi:10.1029/2003JD003571, 2003.
- Kasten, F.: A new table and approximation formula for the relative optical air mass, *Arch. Met. Geophys. Biokl., Ser. B*, 14 (2), 206–223, 1966.
- Kasten, F. and Young, A. T.: Revised optical air mass tables and approximation formula, *Appl. Opt.* 28, 4735–4738, 1989.
- 5 Kazadzis, S., Bais, A., Kouremeti, N., Gerasopoulos, E., Garane, K., Blumthaler, M., Schallhart, B., and Cede, A.: Direct spectral measurements with a Brewer spectroradiometer: Absolute calibration and aerosol optical depth retrieval, *Appl. Opt.*, 44, 1681–1690, 2005.
- Kazadzis, S., Bais, A., Amiridis, V., Balis, D., Meleti, C., Kouremeti, N., and Tzoumaka, P.: Nine years of UV aerosol optical depth measurements at Thessaloniki, Greece, *Atmospheric Chemistry and Physics*, 7(8), 2091–2101, 2007.
- 10 Kerr, J. B.: The Brewer Spectrophotometer, in: *UV Radiation in Global Climate Change*, pp. 160–191, Springer, 2010.
- Kimlin, M. G., and Schallhorn, K. A.: Estimations of the human ‘vitamin D’UV exposure in the USA, *Photochemical & Photobiological Sciences*, 3(11-12), 1067–1070, 2004.
- Kirchhoff, V. W. J. H., Silva, A. A., Costa, C. A., Lem, N. P., Pavao, H. G., and Zaratti, F.: UV-B optical thickness
15 observations of the atmosphere, *J. Geophys. Res.*, 106, 2963–2973, 2001.
- Komhyr, W.: Operations Handbook - Ozone Observations with a Dobson Spectrophotometer, WMO Global Ozone Research and Monitoring Project, report no. 6 edition, 1980.
- Krotkov, N. A., Bhartia, P. K., Herman, J. R., Fioletov, V., and Kerr, J.: Satellite estimation of spectral surface UV irradiance in the presence of tropospheric aerosols 1. Cloud-free case, *J. Geophys. Res.*, 103 (D8), 8779–8793,
20 1998.
- Kumharn, W., Rimmer, J. S., Smedley, A. R. D., Ying, Y. T., and Webb, A. R.: Aerosol Optical Depth and the Global Brewer Network: A Study Using U.K. - and Malaysia-Based Brewer Spectrophotometers, *J. Atmos. Oceanic Technol.*, 29, 857–866, 2012.
- Liu, S. C., McKeen, S. A., and Madronich, S.: Effect of anthropogenic aerosols on biologically active ultraviolet radiation,
25 *Geophys. Res. Lett.*, 18, 2265–2268, 1991.
- ~~Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715–737, doi:10.5194/acp-5-715-2005, 2005.~~
- Lyamani, H., Olmo, F. J., and Alados-Arboledas, L.: Physical and optical properties of aerosols over an urban location in Spain: seasonal and diurnal variability, *Atmos. Chem. Phys.*, 10, 239–254, doi:10.5194/acp-10-239-2010, 2010.
- 30 Marengo, F., di Sarra, A., and Di Luisi, J.: Methodology for determining aerosol optical depth from Brewer 300–320nm ozone measurements, *Appl. Optics*, 41(9), 1805–1814, doi:10.1364/AO.41.001805, 2002.
- Marengo, F.: On Langley plots in the presence of a systematic diurnal aerosol cycle centered at noon: A comment on recently proposed methodologies, *J. Geophys. Res.*, 112, D06205, doi:10.1029/2006JD007248, 2007.

- Mishchenko, M. I. and Geogdzhayev, I. V.: Satellite remote sensing reveals regional tropospheric aerosol trends, *Opt. Express*, 15, 7423–7438, doi:10.364/OE.15.007423, 2007.
- Nieke, J., Pflug, B., and Zimmermann, G.: An aureolecorrected Langley-plot method developed for the calibration of HiRES grating spectrometers, *J. Atmos. Sol. Terr. Phys.*, 61, 739–744, 1999.
- 5 Pribullová, A.: Spectral UV aerosol optical thickness determined from the Poprad-Gánovce Brewer spectrophotometer observations, *Contributions to Geophysics and Geodesy*, 32/3, 291–307, 2002.
- Raghavendra Kumar, K., Narasimhulu, K., Balakrishnaiah, G., Suresh Kumar Reddy, B., Rama Gopal, K., Reddy, R. R., Satheesh, S. K., Krishna Moorthy, K., and Suresh Babu, S.: A study on the variations of optical and physical properties of aerosols over a tropical semi-arid station during grassland fire, *Atmos. Res.*, 95, 77–87, doi:10.1016/j.atmosres.2009.08.012, 2010.
- 10 Redondas, A., Evans, R., Stuebi, R., Köhler, U., and Weber, M.: Evaluation of the use of five laboratory-determined ozone absorption cross sections in Brewer and Dobson retrieval algorithms, *Atmos. Chem. Phys.*, 14, 1635–1648, 2014.
- Serdyuchenko, A., Gorshelev, V., Weber, M., Chehade, W., and Burrows, J. P.: High spectral resolution ozone absorption cross-sections – Part 2: Temperature dependence, *Atmos. Meas. Tech.*, 7, 625–636, 2014.
- 15 Savastiouk, V., and McElroy, C.: Brewer spectrophotometer total ozone measurements made during the 1998 Middle Atmosphere Nitrogen Trend Assessment (MANTRA) campaign, *Atmos.–Ocean*, 43, 315–324, 2005.
- Savastiouk, V.: Improvements to the direct-sun ozone observations taken with the Brewer spectrophotometer. Ph.D. thesis, York University, Toronto, Canada. 2006.
- Schwartz, S. E., and Warneck, P.: Units for use in atmospheric chemistry, *Pure and Applied Chemistry*, 67, 1377–1406, 20 1995.
- Sci-Tec: Brewer MKIV Spectrophotometer Operator’s Manual, Saskatoon, Canada, 1999.
- Seinfeld, J. H., Pandis, S. N.: *Atmospheric chemistry and physics: From air pollution to climate change*, second edition, John Wiley & sons, inc., New Jersey, USA, 2006.
- Sellitto, P., d. Sarra, A., and Siani, A. M.: An improved algorithm for the determination of aerosol optical depth in the 25 ultraviolet spectral range from Brewer spectrophotometer observations, *J. Opt.*, 10A, 849–855, 2006.
- Silva, A. A., and Kirchhoff, V. W. J. H.: Aerosol optical thickness from Brewer spectrophotometers and an investigation into the stray-light effect, *Appl. Opt.*, 43, 2484–2489, 2004.
- ~~Unger, N., Menon, S., Koch, D. M., and Shindell, D. T.: Impacts of aerosol cloud interactions on past and future changes in tropospheric composition, *Atmos. Chem. Phys.*, 9, 4115–4129, doi:10.5194/acp-9-4115-2009, 2009.~~
- 30 Vaníček, K., Metelka, L., Skřivánková, P., Stanek, M.: Long-term changes of the ozone layer over the Czech republic, 1st edition, Transaction of the Czech Hydrometeorological Institute, Volume 56, Published by the Czech Hydrometeorological Institute, Praha, Czech republic, 68 pp., 2012.
- WHO: Air quality guidelines: global update 2005: particulate matter, ozone, nitrogen dioxide, and sulfur dioxide, Tech. rep., World Health Organization, Geneva, Switzerland, 2006.

WMO: Guide to meteorological instruments and methods of observation, World Meteorological Organization, WMO - No 8, 7th Edn., Geneva, Switzerland, 2008.

WMO: Scientific Assessment of Ozone Depletion: 2014, World Meteorological Organization, Global Ozone Research and Monitoring Project-Report No. 55, Geneva, Switzerland, 2014.

- 5 Zerefos, C. S., Tourpali, K., Eleftheratos, K., Kazadzis, S., Meleti, C., Feister, U., Koskela, T., and Heikkilä, A.: Evidence of a possible turning point in solar UV-B over Canada, Europe and Japan, *Atmos. Chem. Phys.*, 12, 2469-2477, <https://doi.org/10.5194/acp-12-2469-2012>, 2012.