

Interactive comment on “Combined Impacts of Nitrous Acid and Nitryl Chloride on Lower Tropospheric Ozone: New Module Development in WRF-Chem and Application to China” by Li Zhang et al.

Anonymous Referee #2

Received and published: 26 May 2017

Zhang et al. describe an updated WRF-Chem model with a new chemical mechanism (CBMZ_ReNOM) developed to improve predictions of photochemical O₃ production in Eastern China, incorporating revised source chemistry of HONO and photolyzable chlorine species.

The paper is written well and is publishable. Ultimately, I wasn't sure how much an improvement this work actually represents. One might hope that a more explicit representation of chemistry within a model does improve its accuracy. Perhaps this part of the manuscript could be strengthened, for instance, through a more reasonable

[Printer-friendly version](#)

[Discussion paper](#)



comparison of predicted with observed concentrations: instead of comparing averages (Tables 4 and 5), why not compare selected time series of measurements and model predictions, in particular for transient species such as HONO and CINO₂ whose concentrations are highly variable.

Specific comments

pg 3 line 17 "To the best of our knowledge, no global or regional models, however, have simultaneously considered the sources/processes of HONO and CINO₂ and evaluated their regional impacts on the formation of O₃ pollution in the boundary layer of the atmosphere." I am not sure the statement as written is true. Many studies have simultaneously considered HONO and CINO₂ as radical sources and showed how these species affect O₃. For instance, Sarwar et al 2004 [GRL, 2014] studied O₃ formation using CMAQV5.02 which contains the RACM2 mechanism [Goliff et al., AE 2013] and HONO chemistry. Also, Ahmadov et al. [ACP, 2015] and Edwards et al. [Nature, 2014] used models and HONO/CINO₂ data to investigate high wintertime ozone pollution events in an oil- and natural gas-producing region of the western US. There have also been numerous papers using 0D box models examining this chemistry.

pg 4 line 9 "reproduced the observed HONO by 85% on average" Not sure what this means.

pg 5 line 17. What is FMCI? A fluorine-metal compound?

pg 7 line 7. "has been proved" A model cannot be proven, at least not in a mathematical sense. Why not simply say that this model has made reasonable predictions of PM_{2.5} and O₃?

pg 7, section "2.2.3 O₃ and NO₂ measurement data" Please state how accurate these measurements are.

pg 7, last line. The Mo converter also "detects" NO₃, 2N₂O₅, HONO, CINO₂, PAN, and HNO₃ to some degree as if it were NO₂. The model should give some indication

[Printer-friendly version](#)[Discussion paper](#)

as what fraction of NO_y is in the form of NO_z (as a function of time of day); consider a sensitivity run in which the NO₂ reported by the routine measurements is scaled down by this factor.

pg 8 " 3.1.2 Spatial and vertical distributions of N₂O₅ and ClNO₂ ". These predicted concentrations are considerably lower than recent measurements in the HK area (see sections further down).

pg 9, line 11 "The model very well captured the measured HONO at Wangdu in the NCP region during a matching simulation period, reproducing 86% of the observations (0.81 vs. 0.94 ppb)" I don't understand this sentence. How does a model reproduce 86% of observations? Consider instead a scatter plot of model concentrations vs observations.

pg 9, line 25 onward. Brown, S. S., et al. (2016), Nighttime chemistry at a high altitude site above Hong Kong, J. Geophys. Res.-Atmos., 121(5), 2457-2475, doi: 10.1002/2015jd024566 observed much higher concentrations than the model predicts. Please discuss.

pg 18, Table 1. Please state the units of the reaction rates.

rxn 7. Water should have a subscript.

rxn 26. What are PAR and X? Also, define the other terms, such as OLI, ALD2 etc.

Interactive comment on Atmos. Chem. Phys. Discuss., doi:10.5194/acp-2017-389, 2017.

Printer-friendly version

Discussion paper

