

Canopy uptake dominates nighttime carbonyl sulfide fluxes in a boreal forest

Linda M.J. Kooijmans¹, Kadmiel Maseyk², Ulli Seibt³, Wu Sun³, Timo Vesala^{4,5}, Ivan Mammarella⁴, Pasi Kolari⁴, Juho Aalto^{4,6}, Alessandro Franchin^{4,8}, Roberta Vecchi⁷, Gianluigi Valli⁷, and Huilin Chen^{1,8}

¹Centre for Isotope Research, University of Groningen, Groningen, The Netherlands.

²School of Environment, Earth and Ecosystem Sciences, The Open University, Milton Keynes, United Kingdom.

³Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA.

⁴Department of Physics, University of Helsinki, Helsinki, Finland.

⁵Department of Forest Sciences, University of Helsinki, Helsinki, Finland.

⁶SMEAR II, Hyytiälä Forestry Field Station, University of Helsinki, Korkeakoski, Finland.

⁷Department of Physics, Università degli Studi di Milano and INFN, Milan, Italy.

⁸Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado, Boulder, Colorado, USA.

Correspondence to: Huilin Chen (Huilin.Chen@rug.nl)

Abstract. Nighttime vegetative uptake of carbonyl sulfide (COS) can exist due to the incomplete closure of stomata and the light-independence of the enzyme carbonic anhydrase, which complicates the use of COS as a tracer for gross primary productivity (GPP). In this study we derived nighttime COS fluxes in a boreal forest (the SMEAR II station in Hyytiälä, Finland; 61°51' N, 24°17' E, 181 m ASL) from June to November 2015 using two different methods: eddy-covariance (EC) measurements ($F_{\text{COS-EC}}$) and the radon-tracer method ($F_{\text{COS-Rn}}$). The total nighttime COS fluxes averaged over the whole measurement period were -6.8 ± 2.2 and -7.9 ± 3.8 $\text{pmol m}^{-2} \text{s}^{-1}$ for $F_{\text{COS-Rn}}$ and $F_{\text{COS-EC}}$, respectively, which is 33–38 % of the average daytime fluxes and 21 % of the total daily COS uptake. The correlation of ²²²Radon (of which the source is the soil) with COS (average $R^2 = 0.58$) was lower than with CO_2 (0.70), suggesting that the main sink of COS is not located at the ground. These observations are supported by soil chamber measurements that show that soil contributes to only 34–40 % of the total nighttime COS uptake. We found a decrease of COS uptake with decreasing night-time stomatal conductance and increasing VPD and air temperature, driven by stomatal closure in response to a warm and dry period in August. We also discuss the effect that canopy layer mixing can have on the radon-tracer method and the sensitivity of ($F_{\text{COS-EC}}$) to atmospheric turbulence. Our results suggest that the nighttime uptake of COS is mainly driven by the tree foliage and is significant in a boreal forest, such that it needs to be taken into account when using COS as a tracer for GPP.

15 *Copyright statement.* TEXT

1 Introduction

The global budget of carbonyl sulfide (COS) is of interest for both stratospheric and tropospheric chemistry (Watts , 2000; Kettle et al., 2002; Berry et al., 2013; Launois et al., 2015). COS contributes to the formation of the sulfate aerosol layer in the stratosphere (Crutzen, 1976; Chin and Davis, 1995) and thereby also plays a role in ozone depletion (Brühl et al., 2012). In the troposphere COS is linked to the carbon cycle because it follows the same diffusion pathway into plant stomata as CO₂ during photosynthesis. After COS has entered a plant cell it is hydrolyzed by the enzyme carbonic anhydrase (CA) to form H₂S and CO₂ (Protoschill-Krebs et al., 1996). As this reaction is practically irreversible, COS is not re-emitted by plants, in contrast to CO₂. The close coupling of COS and CO₂ uptake fluxes by vegetation makes COS a potentially powerful tracer for estimates of gross primary production (GPP) (Sandoval-Soto et al., 2005; Montzka et al., 2007; Campbell et al., 2008; Wohlfahrt et al., 2012; Asaf et al., 2013).

Besides the difference in re-emission, the COS and CO₂ uptake processes differ in the sense that the consumption of COS by the CA enzyme is light-independent. This means that vegetative uptake of COS can continue during the night if stomata are not completely closed (Maseyk et al., 2014). Caird et al. (2007) showed that nighttime stomatal conductance exists in a wide variety of plant species and several studies report nighttime depletion of COS mole fractions (White et al., 2010; Belviso et al., 2013; Commane et al., 2013; Berkelhammer et al., 2014; Billesbach et al., 2014; Maseyk et al., 2014; Commane et al., 2015; Wehr et al., 2017). The measurements presented in White et al. (2010), Maseyk et al. (2014), Berkelhammer et al. (2014) and Wehr et al. (2017) indicated that nighttime ecosystem COS fluxes were indeed dominated by the vegetation, and not by the soil. In these studies, nighttime vegetative fluxes varied between 25 and 50 % of average daytime fluxes. A correlation between nighttime COS fluxes and stomatal conductance is expected when the nighttime sink of COS is primarily driven by vegetative uptake. The relation between H₂O and COS fluxes shown by Seibt et al. (2010), Wohlfahrt et al. (2012) and Berkelhammer et al. (2014) underpins the likely relation between stomatal conductance and COS fluxes. However, the relation between COS fluxes and stomatal conductance measurements has not been studied under field conditions. Instead, Wehr et al. (2017) used COS ecosystem fluxes to estimate stomatal conductance. This relation can especially be useful for estimating nighttime stomatal conductance, which cannot be accurately determined under humid conditions as the concentration gradient of water vapor in leaf chambers gets too small (Maseyk et al., 2014).

Although COS is not used as a GPP tracer during nighttime conditions (when GPP is zero), nighttime COS fluxes may interfere with the use of COS for GPP estimates (Berry et al., 2013; Berkelhammer et al., 2014). To analyze the role of nighttime COS fluxes on the total COS budget and study correlations with environmental drivers, it is key to determine nighttime COS fluxes accurately. Eddy-covariance (EC) is a well-established technique to determine ecosystem fluxes (Aubinet et al., 2012); however, stable nighttime conditions complicate the measurements due to non-turbulent processes like canopy-layer storage and advection (Papale et al., 2006; Wohlfahrt et al., 2012; Aubinet et al., 2012). A method that has been used to derive specifically nighttime fluxes of trace gases, including COS, is the radon-tracer method (Schmidt et al., 1996; Van der Laan et al., 2009; Belviso et al., 2013). This method relates the nighttime buildup of trace gas concentrations to that of ²²²Radon

(^{222}Rn) concentrations and the ^{222}Rn flux, which is solely driven by the soil. Both the EC and radon-tracer methods can complement each other to help understand and reduce uncertainties of nighttime flux measurements.

The aim of this study is to quantify nighttime COS fluxes to determine the role of these fluxes in the ecosystem COS budget, and to understand the driving parameters of nighttime COS uptake. In the summer of 2015, we conducted a field campaign in a Finnish boreal forest using a combination of COS measurements: atmospheric concentration profiles, and EC and soil chamber measurements. We use both the EC and radon-based fluxes to quantify nighttime COS fluxes and infer information about the sink apportionment within the canopy. We also investigate the correlation of nighttime COS fluxes with stomatal conductance and environmental parameters and discuss the implications of nighttime COS fluxes for large-scale GPP estimates.

2 Field measurements and data

2.1 Measurement site

The field campaign was held from June to November 2015 at the Station for Measuring Forest Ecosystem-Atmosphere Relations (SMEAR II) in Hyytiälä, Finland ($61^{\circ}51' \text{ N}$, $24^{\circ}17' \text{ E}$, 181 m ASL). The forest represents boreal coniferous forest and the measurement site is covered by 50–60 year old Scots pine (*Pinus sylvestris*) up to 1 km towards the North from the measurement site and for about 200 m in all other directions (Rannik et al., 1998, 2004). The forest outside this area covers younger pine and spruce. About 700 m southwest of the measurement site is an oblong lake of about 200 m wide. The dominant canopy height is 17 m with the base at 7 m and the site is characterized by modest height variation. At this latitude, the daylight duration has a maximum in June with 19 hours and 40 minutes and is 7 hours in November.

2.2 Instrumentation for measurements of COS, CO_2 , and H_2O

Two quantum cascade laser spectrometers (QCLS) manufactured by Aerodyne Research Inc. (BillERICA, MA, USA) were deployed in the field for simultaneous measurements of COS, CO_2 , CO, and H_2O and are described separately in the following two sections.

2.2.1 QCLS for vertical profile and soil flux measurements

From June 1 until November 4, one QCLS was operated at 1 Hz for concentration measurements of sampled air at 4 heights: 125 m (tall tower), 23 m, 14 m, and 4 m (small tower at 30 m distance from the tall tower). An additional height of 0.5 m was measured as part of the soil chamber measurement routine from June 28 onwards. A multi-position Valco valve (VICI; Valco Instruments Co. Inc.) was used to switch between the sample tubing from the different profile heights, soil chambers and calibration cylinder gases. A cycle of 1 hour during the night and during the day is shown in Figs. S1 and S2 in supplementary material. The sample tubing was continuously flushed. For the profile measurements, the flow rates were set such that there was a time delay between 30 and 60 s from the moment that the air enters the inlet at different heights until it reaches the cell of the QCLS, which is 17 L min^{-1} for 125 m and 2 L min^{-1} for 4 m. The flow rate from the Valco valve through the sample

cell was set at 0.15 L min^{-1} where the sample cell has a volume of 0.5 L. The following measurements were made during each hour: 3 minutes for each of the four heights, 16 minutes for each of the two soil chambers, two times 3 minutes for one calibration cylinder to correct for instrument drift, 3 minutes for each of two other calibration cylinders to assess the accuracy of the measurements. The first 60 s of each 3-minute measurement were discarded to account for cell flushing time. The three
5 cylinders were filled with ambient air and calibrated against two NOAA/ESRL standards for COS (NOAA-2004 scale) and CO_2 (WMO-X2007 CO_2 scale) at the University of Groningen. A 'zero' air spectrum was measured once every six hours using high-purity nitrogen ($\text{N } 5.0$). The overall uncertainty including scale transfer, water vapor corrections, and measurement precision of this analyzer was determined to be 7.5 ppt for COS and 0.23 ppm for CO_2 (Kooijmans et al., 2016). More detailed information about the calibration and correction methods can be found in (Kooijmans et al., 2016).

10 2.2.2 QCLS for eddy covariance measurements

A second QCLS was used to measure COS, CO_2 , CO, and H_2O concentrations at 10 Hz from June 28 onwards. The air is sampled with a flow of $9\text{--}10 \text{ L min}^{-1}$ at 23 m height at a small tower that is at 30 m distance from the 125 m tall tower. Wind velocity components were measured by a sonic anemometer (Solent Research HS1199, Gill Ltd., Lymington, Hampshire, England) to derive ecosystem fluxes through the EC method. For this analyzer a 'zero' air spectrum was measured once every
15 30 minutes. This QCLS was calibrated against a standard on the same scale as the first QCLS. The CO_2 and H_2O fluxes from the QCLS were compared with those obtained at the nearby tall tower as quality control. The instrumentation in the tall tower is a Gill Solent 1012R anemometer and a Li-Cor LI-6262 gas analyzer (Mammarella et al., 2009).

2.3 Soil chambers

Two soil flux chambers (LI8100-104C; Li-Cor) modified for analysis of COS were used in combination with the concentration
20 measurements of the QCLS at 1 Hz to derive soil fluxes. The modifications included operation in an open flow configuration, replacing the chamber bowl and soil collar with stainless steel components, and removing or replacing other COS-producing material. Each chamber was closed once per hour for 9 or 10 minutes. For supply flow into the chambers, air was sampled at 0.5 m height in the vicinity of the soil chambers and was measured for 3 minutes before and after chamber closure. The air was pumped into the chambers with flow rates between 1.5 and 2.1 L min^{-1} through a diaphragm pump (KNF 811) for which we
25 found no interference with COS. More details on the soil measurements can be found in Sun et al. (2017).

2.4 Auxiliary data

2.4.1 ^{222}Rn

^{222}Rn concentrations were obtained by measurement of its short-lived decay products attached to aerosol particles (i.e. ^{214}Bi). Detection of short lived decay products concentration in outdoor air was done by continuous on-line alpha spectroscopy during
30 aerosol sampling. Aerosol particles were collected at 8 m height as part of the ongoing aerosol monitoring at the site (Hari and Kulmala, 2005; Nieminen et al., 2014) about 50 m away from the tower where COS and CO_2 was sampled. Particles were

collected on a glass micro fibre filter (Whatman GF/A, 47-mm diameter) with an average flowrate of 17.4 l min^{-1} . Alpha particles emitted by Radon decay products were recorded by a silicon surface barrier detector (ULTRA™ Alpha Detector by ORTEC, with F.W.H.M. of 42 keV) placed a few millimeters in front of the filter in order to optimize the efficiency and to allow the detection of alpha particles in air. The hourly alpha energy spectra were continuously recorded. The concentration of radon daughters is calculated by taking into account radioactive decay equations, the accumulation of decay products on the filter during the sampling and the hypothesis of equilibrium in the progeny after subtraction of the ^{220}Rn daughter contribution. Following (Schmidt et al., 1996), ^{222}Rn and its decay products were considered in secular radioactive equilibrium in this work. Further details on the experimental procedure are reported in (Marcazzan et al., 2003) and (Sesana et al., 2003).

2.4.2 Stomatal conductance

The stomatal conductance to water vapor (g_{sw}) was determined from transpiration measurements obtained through shoot chamber measurements at a pine shoot at the top of the canopy crown (Altimir et al., 2006). The conductance is derived from the vapor pressure deficit at leaf temperature assuming that the resistance due to the leaf boundary layer is negligible due to ventilation of the air in the shoot chambers. The leaf temperature is calculated following a leaf energy balance model that incorporated heating by incoming shortwave radiation, cooling by transpiration and convection, and thermal radiation balance. Conductances measured under humid conditions (relative humidity (RH) > 80 %) were rejected due to the underestimation of transpiration at higher RH levels. The stomatal conductance to COS (g_{sCOS}) is derived based on the relationship between COS and H_2O conductance: $g_{\text{sCOS}} = g_{\text{sw}} / R_{\text{wCOS}}$ (Seibt et al., 2010) where R_{wCOS} is the ratio of H_2O and COS diffusivities and is derived by (Seibt et al., 2010) to be 2.0 ± 0.2 .

2.4.3 Meteorological data

Meteorological data such as the friction velocity (u_*), air temperature (T_{air}), relative humidity (RH), soil water content (SWC) and wind direction were available through the SmartSMEAR database which contains continuous data records from the SMEAR sites (available at <http://avaa.tdata.fi>). The vapour-pressure deficit (VPD) was calculated from RH and T_{air} .

3 Flux derivations

3.1 The EC-based method

3.1.1 Eddy covariance fluxes

The EC technique is based on turbulence measurements above the canopy and fluxes are derived from the covariance between a scalar (in this case COS or CO_2) and the vertical wind speed (e.g., Aubinet et al., 2012; Mammarella et al., 2007). The fluxes derived through this method represent the net exchange of gases between the canopy layer and the air above. The EC technique requires turbulent conditions, otherwise gases that accumulate or get depleted due to sources and/or sinks within the canopy do not reach the sensors above the canopy. As soon as turbulence is enhanced in the early morning, these gases are released to

levels above the canopy and are only then being captured by the EC system. This so called storage change within the canopy can be significant and should be added to the turbulence flux to account for the delayed capture of fluxes by the EC system (Aubinet et al., 2012). In this study we refer to the storage-corrected COS and CO₂ EC flux as F_{COS-EC} and NEE_{EC}, respectively. The calculation of storage fluxes is discussed in the next section. In this study the EC fluxes were calculated using the EddyUH software package developed at the University of Helsinki (Mammarella et al., 2016). In short, the high-frequency EC data were despiked according to standard approach (Vickers and Mahrt, 1997). The spectroscopic correction due to H₂O impact on the absorption line shape was accounted for along with the dilution correction in the QCLS acquisition software. A 2D rotation of sonic anemometer wind components was performed, and 30 min covariances between the scalars and vertical wind velocity were calculated using linear detrending method. Short-term drift in the QCLS high-frequency concentration data was negligible and there was no need to apply more sophisticated approach for detrending the data, e.g. high pass recursive filters (Mammarella et al., 2010). The time lag between the concentration and wind measurements induced by the sampling line was determined by maximizing the covariance. Due to better signal-to-noise ratio, the lag for COS was determined by maximizing the covariance for QCLS CO₂, and the same lag was assigned to COS. Finally, spectral correction was done according to Mammarella et al. (2009). Total random uncertainty of the fluxes (Rannik et al., 2016) was calculated according to the method implemented in EddyUH, the method proposed by (Finkelstein and Sims, 2001). The uncertainties of NEE_{EC} and F_{COS-EC} are estimated from the standard deviation of data points per night, where night is defined as the time when the sun elevation angle is below -3°. A general observation that is seen with EC measurements is that nighttime NEE_{EC} decreases with lower u_{*}, whereas respiration is not expected to depend on atmospheric turbulence. For this reason we filtered out (storage-corrected) fluxes with u_{*} values below a threshold of 0.3 m s⁻¹ (Mammarella et al., 2007). A difference between COS and CO₂ fluxes is, however, that the uptake of COS by leaves is concentration dependent (Berry et al., 2013) and the leaf boundary layer may get depleted in COS under low turbulence conditions, slowing uptake rates. It is unknown to what extent this affects COS fluxes in practice, but it has to be kept in mind that the u_{*} filtering may be an overstated filtering to COS fluxes. To determine the fraction that nighttime COS fluxes contribute to total daily COS uptake we gapfilled COS fluxes with a rectangular hyperbola light response function that is based on the measured data. Missing COS data under dark conditions were filled based on the average nighttime flux obtained from this study.

CO₂ and H₂O ecosystem fluxes from the QCLS were compared with those from the nearby tall tower. During nighttime, the QCLS CO₂ flux is a factor 0.73 smaller than the tall-tower fluxes at the same height and the underestimation has been observed with another EC-system at the small tower as well. Kolari et al. (2009) found that the tall tower NEE_{EC} agrees well with upscaled soil and branch chamber measurements. As we rely on the accuracy of NEE_{EC} in the radon-tracer method (Section 3.2) we use NEE_{EC} from the tall tower instead of the QCLS at the smaller tower throughout the manuscript. The underestimation is not the same for all gases, e.g. the evapotranspiration flux is only a factor 0.97 smaller. It is therefore unknown by how much the F_{COS-EC} flux is affected by the general underestimation at the small tower.

3.1.2 Storage fluxes

Storage fluxes (F_{stor}) are defined as the integral of concentration changes over height up to the height of the EC measurements (h_{EC}):

$$F_{stor} = \frac{P}{RT_{air}} \int_0^{h_{EC}} \frac{dC(z)}{dt} dz, \quad (1)$$

5 with P the atmospheric pressure, R the molar gas constant and C(z) the COS or CO₂ concentrations (ppt for COS or ppm for CO₂) along a profile (Aubinet et al., 2001; Papale et al., 2006). The integral was determined from hourly measured profile concentrations at 0.5, 4, 14, and 23 m in two ways: (1) by integrating an exponential fit through the data, and (2) using trapezoidal areas (Winderlich et al., 2014). The concentration at ground level that is used for the second calculation method is estimated by extrapolating the gradient between 0.5 and 4 m to the ground level. A third calculation was done assuming a
10 constant profile from the EC measurement height (23 m) to the ground level, to test the bias in storage fluxes when no profile measurements are available. The results of the different calculation methods will be discussed in Section 4.1. To reduce the error due to the random noise of COS concentration measurements, a running average over a 5 hour window was applied to the COS concentration data before the storage fluxes were calculated.

3.2 The radon-tracer method

15 ²²²Rn is a natural radioactive gas that is formed by the decay of ²²⁶Radium, which is uniformly distributed in soils (Van der Laan et al., 2009). Once in the atmosphere, ²²²Rn is affected by radioactive decay and atmospheric mixing. As the exhalation rate of ²²²Rn by the soil (F_{Rn}) is considered constant and uniformly distributed, and ²²²Rn is mixed through the atmosphere in the same way as other trace gases, the surface fluxes of these trace gases (F_C) can be determined from the concentration change of these gases over time (ΔC) relative to that of ²²²Rn ($\Delta^{222}Rn$) (Schmidt et al., 1996; Van der Laan et al., 2009; Belviso et
20 al., 2013):

$$F_C = F_{Rn} \frac{\Delta C}{\Delta^{222}Rn}. \quad (2)$$

²²²Rn generally builds up in the boundary layer when it gets shallower during the night. Fig 1. shows an example of one night during the measurement campaign where ²²²Rn concentrations increase in the evening and reach a maximum in the night, while at the same time CO₂ increases and COS decreases. This nighttime buildup of gases and the constant surface flux of ²²²Rn
25 make the radon-tracer method appropriate to derive nighttime fluxes of trace gases. Requirements for this method are that the ²²²Rn concentrations are corrected for radioactive decay, that F_{Rn} is known, and that a high correlation exists between the trace gas and ²²²Rn concentrations. Moreover, when the spatial distribution of sources and sinks of a trace gas are similar to the source of ²²²Rn at the ground, a high correlation between the trace gas and ²²²Rn can be expected. Therefore, the correlation between COS and ²²²Rn concentrations may give insight into the distribution of sinks of COS within the ecosystem.

One of the main uncertainties of the radon tracer method is the magnitude of F_{Rn} . In Szegvary et al. (2007), F_{Rn} was measured at a site 46 km away from the SMEAR II site, which resulted in $F_{Rn} = 15.3 \text{ mBq m}^{-2} \text{ s}^{-1}$. Model studies have estimated F_{Rn} in Europe from 4.0 to 12.4 $\text{mBq m}^{-2} \text{ s}^{-1}$, (summarized in Table S1 in supplementary material), leading to an overall average of $9.6 \pm 4.1 \text{ mBq m}^{-2} \text{ s}^{-1}$. The exhalation rates depend on the uranium content and soil properties that affect diffusive transport such as the soil texture and soil moisture (Karsens et al., 2015). The F_{Rn} values of 4.0 and 11.4 $\text{mBq m}^{-2} \text{ s}^{-1}$ that were modelled by Karsens et al. (2015) for two different soil moisture maps indicate that the uncertainty of F_{Rn} is in large part caused by different soil moisture.

As the uncertainty of the COS and CO_2 ecosystem fluxes derived through the radon-tracer method ($F_{\text{COS-Rn}}$ and NEE_{Rn} respectively) is in large part determined by the uncertainty of F_{Rn} , it is key to further limit the F_{Rn} range between 4.0 and 15.3 $\text{mBq m}^{-2} \text{ s}^{-1}$ in Table S1. For that reason we inverted the radon-tracer method to derive F_{Rn} from CO_2 and ^{222}Rn concentrations with a known ecosystem CO_2 flux (NEE_{EC}), instead of a known F_{Rn} to derive NEE, which is normally used in the radon-tracer method (Van der Laan et al., 2016). The advantage of this method is that F_{Rn} is obtained from actual measurements at the site, and we will therefore use this F_{Rn} to determine $F_{\text{COS-Rn}}$. We derived F_{Rn} over the period from July to November and found an average of 5.9 $\text{mBq m}^{-2} \text{ s}^{-1}$ with a standard deviation of 3.9 $\text{mBq m}^{-2} \text{ s}^{-1}$ and a standard error of 0.8 $\text{mBq m}^{-2} \text{ s}^{-1}$. This value of F_{Rn} is within the range listed in Table S1, but is lower than the average of 9.6 $\text{mBq m}^{-2} \text{ s}^{-1}$. We will discuss in Section 5.2 what the effect of canopy layer mixing can be on the derivation of F_{Rn} and COS fluxes. Temporal variation of F_{Rn} can be expected due to the changes in SWC that affects the soil permeability; however, no temporal change or correlation with SWC was found ($R^2 = 0.07$) throughout the season (see Fig. S3 in supplementary material).

In Hyytiälä, ^{222}Rn measurements were made at 8 m, and COS and CO_2 concentrations from the same height need to be used to derive their surface fluxes. We derived concentrations at 8 m from an exponential fit through the profile concentrations at 0.5, 4, 14 and 23 m. A linear fit between 4 and 14 m was used in cases where the algorithm for the exponential fit did not converge. The factor $\Delta C/\Delta^{222}\text{Rn}$ is determined from a linear regression of concentrations of COS or CO_2 against ^{222}Rn for data when the sun elevation is below 0° (see Fig. 1 for an example). Per night, a minimum of 5 data points need to be available and R^2 between ^{222}Rn and CO_2 and COS should be at least 0.5 (for CO_2) and 0.3 (for COS). Uncertainties of NEE_{Rn} and $F_{\text{COS-Rn}}$ are determined from the linear regression as the standard error of the slope.

3.3 Soil fluxes

Soil fluxes (F_{soil}) were calculated from least square fits of the concentrations during chamber closure and by considering mass balance equations within the chamber (Sun et al., 2017). At the start of the campaign we did blank tests by placing FEP foil over the soil and calculated fluxes through the standard measurement procedure. Soil fluxes were corrected for blank chamber effects of $0.66 \pm 0.48 \text{ pmol m}^{-2} \text{ s}^{-1}$ for COS, blanks for CO_2 were negligible ($-0.05 \pm 0.15 \text{ } \mu\text{mol m}^{-2} \text{ s}^{-1}$). Further details about the soil flux measurements can be found in (Sun et al., 2017).

4 Results

4.1 COS and CO₂ storage fluxes

The storage fluxes of COS (Fig. 2) are slightly negative during nighttime with an average nighttime value of $-0.9 \text{ pmol m}^{-2} \text{ s}^{-1}$ in Jul–Aug and $-0.5 \text{ pmol m}^{-2} \text{ s}^{-1}$ in Sep–Nov (Fig. S4 in supplementary material). The average nighttime gradient between 23 and 0.5 m corresponding to these storage fluxes is 63 ppt for COS and -45 ppm for CO₂ (23 - 0.5 m concentration) in Jul–Aug and is 57 ppt and -17 ppm in Sep–Nov. Early in the morning when turbulence is enhanced, the storage fluxes become positive and have an average maximum of 2.1 (1.8) $\text{pmol m}^{-2} \text{ s}^{-1}$ at 09:00 (10:00) in Jul–Aug (Sep–Nov). The storage fluxes of CO₂ follow a similar pattern but have the opposite sign. Storage fluxes of COS calculated from trapezoidal areas are on average 25 % larger than when an exponential fit through the profile is integrated. When the concentration profile is assumed to be constant from the EC measurement height to the ground level, the storage flux is on average 7 % smaller compared to a profile with an exponential fit. These differences are small compared to the size of the ecosystem fluxes. Neglecting storage fluxes would not influence the long-term budget of COS and CO₂, as it only corrects for the delay in release of accumulated gases from within the canopy (Aubinet et al., 2012); however, it does affect the diurnal variability of fluxes, and any attempt at flux partitioning, particularly if storage fluxes are large. In this dataset, storage fluxes of both COS and CO₂ are small compared to the EC flux, i.e. storage fluxes are on average 5 % of $F_{\text{COS-EC}}$ and 7 % of NEE_{EC} , with variation between summer and autumn from 4 % (Jul–Aug) to 6 % (Sep–Nov) for $F_{\text{COS-EC}}$.

4.2 COS and CO₂ nighttime fluxes through the radon-tracer and EC-based method

The linear correlation between the concentrations of ²²²Rn and the scalar (COS or CO₂) is key in interpreting the fluxes derived from the radon-tracer method. Fig. 3 shows the distribution of R² values for the correlation between ²²²Rn and COS or CO₂. The correlation between ²²²Rn and CO₂ peaks at R² values in the range 0.9–1.0 and has a median value of 0.70. The R² for COS is generally lower with a median of 0.58. The lower R² values for COS can partly be explained by the lower precision of COS measurements compared to those of CO₂. However, the average R² only slightly increases to 0.64 when the noise of COS is diminished by taking a running average of a 5 hour window over the COS measurements. This indicates that the lower precision of COS is not the main aspect influencing the correlation with ²²²Rn. Another aspect that influences the correlation with ²²²Rn is the similarity in vertical distribution of sources and sinks between the scalar and ²²²Rn, which will be further discussed in Section 5.1.

The radon-based nighttime fluxes of COS and CO₂ are compared with the EC-based fluxes in Fig. 4. $F_{\text{COS Rn}}$ (NEE_{Rn}) was determined for 69 (66) out of 128 nights during the campaign that passed the criteria of a minimum R² and a minimum number of available data. Nighttime fluxes derived with the EC method were determined for 56 nights following removal of 43 % of the data due to u_* filtering. F_{Rn} was derived from ²²²Rn concentrations in relation to NEE_{EC} and CO₂ concentrations in order to limit the uncertainty of F_{Rn} on $F_{\text{COS-Rn}}$. This means that the average NEE_{EC} and NEE_{Rn} values are close (3.30 ± 0.62 and $3.34 \pm 0.82 \mu\text{mol m}^{-2} \text{ s}^{-1}$ respectively) as they are not independent from each other. Both NEE_{EC} and NEE_{Rn} show a decreasing

trend from summer towards autumn. However, the R^2 value between NEE_{EC} and NEE_{Rn} is only 0.03, which is likely due to the low signal-to-noise ratio of both flux techniques.

Both the EC-based and radon-tracer methods show negative nighttime COS fluxes with an average of $-7.9 \pm 3.8 \text{ pmol m}^{-2} \text{ s}^{-1}$ (F_{COS-EC}) and $-6.8 \pm 2.2 \text{ pmol m}^{-2} \text{ s}^{-1}$ (F_{COS-Rn}). In comparison, nighttime soil fluxes of COS are on average $-2.7 \text{ pmol m}^{-2} \text{ s}^{-1}$ (-2.8 ± 1.0 and $-2.5 \pm 1.2 \text{ pmol m}^{-2} \text{ s}^{-1}$ for the two chambers) and soil fluxes do not show a clear diurnal (Fig. 2) or seasonal cycle. An overview of the soil fluxes is presented in Sun et al. (2017). Similar to NEE, a decreasing trend is visible in both F_{COS-Rn} and F_{COS-EC} with an average of $-10.9 \text{ pmol m}^{-2} \text{ s}^{-1}$ in July and $-4.6 \text{ pmol m}^{-2} \text{ s}^{-1}$ in October as obtained from F_{COS-EC} . The nighttime uptake is 33–38 % of the average daytime fluxes (defined as when sun elevation is above 20°) and 21 % of the total daily COS uptake (obtained from gapfilled data). When the soil flux is subtracted from the ecosystem flux, the nighttime uptake is 17 % of the total daily uptake.

4.3 F_{COS} correlation with g_{sCOS} , VPD, T_{air} and u_*

Fig. 5 shows F_{COS} against nighttime averaged g_{sCOS} , VPD, T_{air} and u_* with their respective uncertainties. Soil fluxes did not show a seasonal or daily cycle (Sun et al., 2017) and are therefore not subtracted from the ecosystem-scale fluxes, as this would only add noise to the fluxes. The nights shown in Fig. 5 only cover summer nights between June 28 and August 25, 2015, as g_{sCOS} data did not pass the RH filter criteria after this period due to higher RH. The month August was characterized by a dry period with SWC decreasing from about 20 % down to 7 %, the average nighttime temperature increased and RH decreased. Over the same time period, nighttime g_{sCOS} decreased from $0.02 \text{ mol m}^{-2} \text{ s}^{-1}$ to $0.006 \text{ mol m}^{-2} \text{ s}^{-1}$ (see Fig. S3 in supplementary material for an overview of the meteorological conditions).

Weak correlations are found between F_{COS-Rn} and g_{sCOS} ($R^2 = 0.32$), T_{air} ($R^2 = 0.22$), VPD ($R^2 = 0.22$) and u_* ($R^2 = 0.33$), where fluxes decrease under lower g_{sCOS} and u_* , and higher VPD and T_{air} . The same comparison was made for F_{COS-EC} (Fig. S5 in supplementary material), which gave correlations $R^2 = 0.36$ (g_{sCOS}), 0.30 (T_{air}), 0.56 (VPD) and 0.50 (u_*) and showed that also F_{COS-EC} decreased under lower g_{sCOS} and u_* , and higher VPD and T_{air} . However, these correlations were only found when no u_* filter was applied, as only a few data points remained after the u_* filtering.

g_{sCOS} was on average $0.016 \text{ mol m}^{-2} \text{ s}^{-1}$ during nighttime and $0.117 \text{ mol m}^{-2} \text{ s}^{-1}$ during daytime. The average nighttime g_{sCOS} showed a correlation with the average nighttime VPD ($R^2 = 0.54$, not shown) and g_{sCOS} was negatively correlated with T_{air} ($R^2 = 0.60$; not shown).

5 Discussion

5.1 Vertical distribution of sinks and sources of COS and CO_2 compared to that of ^{222}Rn

The benefit of stable conditions within the canopy layer is that the correlation of COS or CO_2 with ^{222}Rn can shed light on the spatial distribution of sources and sinks of these gases in comparison to the only source of ^{222}Rn , which is the soil. When the source or sink of COS or CO_2 is focused at the ground level, a high correlation between ^{222}Rn and these gases can be expected.

The fact that CO₂ shows a high correlation with ²²²Rn indicates that the main source of CO₂ is located near the surface, which is confirmed by the magnitude of nighttime soil chamber measurements relative to branch chamber measurements in Kolari et al. (2009), who found that respiration of the tree foliage was 1.5–2 μmol m⁻² s⁻¹ during summer nights and soil respiration was 5–6 μmol m⁻² s⁻¹. In contrast, we find that the correlation between ²²²Rn and COS is lower, which suggests that the main sink of COS is not near the surface, but rather at higher levels in the canopy layer. This is also supported by the soil chamber measurements that were on average -2.7 pmol m⁻² s⁻¹ with only little variation between the two chambers, which suggest that the soil contributes to 34–40 % of the total nighttime COS uptake.

5.2 The effect of canopy layer mixing on flux derivations

When the canopy air is fully mixed, the flux obtained through the radon-tracer method represents the net exchange flux in that canopy layer, regardless of the potential difference in the spatial distribution of the tracer fluxes, e.g. CO₂ and ²²²Rn. In this study, however, the ²²²Rn concentrations are measured within the canopy layer at 8 m and decoupling of canopy layers may exist (Alekseychik et al., 2013). Fluxes derived from concentrations within the canopy may therefore not represent the exchange of these gases in the whole canopy. To discuss the effect of decoupling on radon-flux calculations we have to distinguish between two decoupling situations; (1) when the 8 m air is decoupled from the air close to the ground, and (2) when the 8 m air is decoupled from the canopy layer above:

1. When the 8 m canopy layer is decoupled from the air close to the ground, the different flux distribution of CO₂ and ²²²Rn can become apparent. In the case of decoupling, the respiration of the tree foliage would influence the 8 m concentration, while the CO₂ respiration and radon flux at the surface do not influence the air at 8 m. The 8 m concentration is then not representative for the canopy layer CO₂ flux and would lead to a lower F_{Rn}. This would explain why the F_{Rn} that we find (5.9 mBq m⁻² s⁻¹) is lower than the average F_{Rn} reported in other literature (9.6 ± 4.1 mBq m⁻² s⁻¹). At the same time, when COS fluxes do not entirely take place at the surface but within the canopy, this would lead to a higher F_{COS-Rn}.
2. When the 8 m layer is decoupled from the canopy layer above, the air that is depleted in COS due to the sinks within the canopy may not reach the lower canopy layers on which F_{COS-Rn} is based and leads to an underestimation of F_{COS-Rn}. Furthermore, the decoupled layer at the surface is more susceptible to horizontal advection which may affect the concentration profile as well.

Alekseychik et al. (2013) identified decoupling of different canopy levels at the Hyytiälä site based on changing wind directions at different heights. They observed a decrease in NEE_{EC} under decoupled circumstances, which occurred in at least 18.6 % of all nighttime periods. We did not observe a correlation with F_{COS-Rn} and the difference in wind direction between 16.8 and 8.4 m. However, a limitation is that we can only compare nighttime averages, whereas decoupling does not have to last throughout the whole night and can also exist during only a fraction of a night. Furthermore, we do not have wind direction data at other heights within the canopy to be able to determine if the decoupling takes place below or above 8 m.

5.3 Sensitivity of $F_{\text{COS-EC}}$ to u_* .

It is well accepted that NEE_{EC} underestimates the true NEE under low u_* , as nighttime NEE (respiration only) is not expected to depend on atmospheric turbulence. By applying a u_* filter to COS fluxes, we assume the same independence of COS uptake to atmospheric turbulence. However, a negative correlation between F_{COS} and u_* can be expected when the leaf boundary layer gets depleted in COS under low turbulence conditions and the uptake of COS gets limited by the COS gradient at the leaf boundary layer. If this is the case, that means that by applying the u_* filtering to $F_{\text{COS-EC}}$ we bias to higher $F_{\text{COS-EC}}$ data. The correlation between u_* and nighttime COS and CO_2 fluxes that is observed with the EC method ($R^2 = 0.50$ for $F_{\text{COS-EC}}$ and 0.30 for NEE_{EC} , not shown) is also observed with the radon-tracer method for $F_{\text{COS-Rn}}$ ($R^2 = 0.33$) but not for NEE_{Rn} ($R^2 = 0.003$, not shown). This suggests that nighttime COS uptake by plants is limited by the reduced COS concentrations at the leaf boundary layer, which is not the case for CO_2 . This means that the u_* filtering that is applied to $F_{\text{COS-EC}}$ is possibly an overstated filtering and leads to an overestimated $F_{\text{COS-EC}}$, which could explain the difference between $F_{\text{COS-EC}}$ and $F_{\text{COS-Rn}}$.

Similar to the limitation on leaf uptake by depleted COS concentrations, soil COS uptake may also be limited by the depleted COS at the soil-atmosphere interface. In contrast, soil emissions of CO_2 and ^{222}Rn do not depend on atmospheric concentrations. This may explain the stronger similarity between CO_2 and ^{222}Rn emissions, which is reflected in the higher correlation between CO_2 and ^{222}Rn concentrations than that between COS and ^{222}Rn (Fig. 3). However, Sun et al. (2017) found no correlation between soil COS fluxes and COS concentrations ($R^2 < 0.001$) for ambient concentrations between 200 and 450 ppt. This implies that the soil COS flux is not limited by the low ambient concentration at night, and a correlation between u_* and soil COS uptake is not warranted.

5.4 Stomatal control of nighttime F_{COS}

A correlation between nighttime F_{COS} and g_{sCOS} was expected due to stomatal diffusion and the light-independence of the CA enzyme. A weak correlation of g_{sCOS} with F_{COS} was indeed observed for both the radon-tracer and EC method, although the latter was only found when no u_* filtering was applied to the data, as only a few data points remained when the u_* filtering was included. The decrease in F_{COS} when g_{sCOS} decreases and VPD increases is likely related to the dry and warm period in August to which plants respond by closing their stomata to prevent excessive water loss. This would also explain why F_{COS} is lower under high T_{air} . In general we do not find strong correlations between the COS flux and the nighttime environmental parameters, which can be explained by the low signal-to-noise ratio of the flux measurements and the fact that $F_{\text{COS-Rn}}$ may not represent the full canopy layer due to decoupling (see Section 5.2). Moreover, we compare ecosystem fluxes with leaf-level g_{sCOS} within enclosed chambers, which may not represent the full canopy dynamics. Nevertheless, the fact that both the radon-tracer and the EC methods confirm that the COS uptake decreases with decreasing g_{sCOS} indicates that the nighttime uptake of COS is indeed driven by vegetation. Moreover, soil fluxes were found to be $-2.7 \text{ pmol m}^{-2} \text{ s}^{-1}$ on average. With the total nighttime COS uptake being -6.8 to $-8.1 \text{ pmol m}^{-2} \text{ s}^{-1}$, soil fluxes contribute to only 34–40 % of the nighttime COS uptake. Besides uptake of COS by the soil and leaf stomatal diffusion there is no other process to our knowledge that would lead to uptake of COS in the ecosystem. This leads to the conclusion that the nighttime COS uptake is predominantly driven

by vegetative uptake and supports the use of COS to estimate g_{sCOS} (Wehr et al., 2017). Assuming that the soil is the only sink besides the vegetation, we can say that the nighttime vegetative uptake contributes to 17 % of the total daily COS uptake. Moreover, this study has confirmed that nighttime stomatal conductance exists at the Hyytiälä site.

5.5 Effect of nighttime COS fluxes on GPP derivation

5 The measurements in this study showed that, unlike the uptake of CO_2 , the COS uptake continues during the night, which agrees with the light-independence of the CA enzyme. We showed that the nighttime plant COS fluxes cover 17 % of the total daily COS plant uptake, which indicates that nighttime COS uptake is a significant sink in the total COS budget. Including this nighttime sink is essential in regional COS models and will affect COS-based GPP model simulations as well. The relationships that we found between F_{COS} , g_{sCOS} , VPD and T_{air} will aid in implementing nighttime F_{COS} in models. Furthermore, the light-
10 independence of COS uptake should be taken into account when COS is being used as tracer for GPP. Besides restricting COS as GPP-tracer to light conditions, the leaf relative uptake ratio (LRU), which is the normalized ratio between COS and CO_2 fluxes, can be expected to increase when GPP becomes zero around sunrise and sunset while at the same time COS is continuously being taken up by vegetation. So far, only Stimler et al. (2011) showed the light-dependence of LRU from leaf-scale measurements and Maseyk et al. (2014) and Commane et al. (2015) observed a light-dependence in the ratio of
15 ecosystem fluxes of COS and CO_2 . Other studies have focused on LRU values under high light conditions (e.g., Sandoval-Soto et al., 2005; Berkelhammer et al., 2014). More leaf-level COS flux measurements should be made to accurately parameterize the light-dependence of LRU in the field.

6 Conclusions

In this study we quantified nighttime COS fluxes in a boreal forest using both the EC and the radon-tracer methods, and found
20 that nighttime F_{COS} between June and November 2015 was on average $-7.9 \pm 3.8 \text{ pmol m}^{-2} \text{ s}^{-1}$ and $-6.8 \pm 2.2 \text{ pmol m}^{-2} \text{ s}^{-1}$ according to the two different methods, respectively. A high correlation between CO_2 and ^{222}Rn indicates that the sources of these gases have a similar spatial distribution, namely at the soil. A lower correlation of ^{222}Rn with COS suggests that the main sink of COS is not located at the surface, but rather at higher levels in the canopy. This is supported by soil chamber measurements, which show that the soil flux is on average $-2.7 \text{ pmol m}^{-2} \text{ s}^{-1}$ and only contributes to 34–40 % of the total
25 nighttime COS uptake.

Our estimates for nighttime F_{COS} are 33–38 % of the size of daytime average NEE_{EC} fluxes. Based on the EC method, the nighttime COS uptake is 21 % of the total daily COS uptake and is mostly driven by aboveground vegetation. Furthermore, we investigated the relation of the nighttime COS fluxes with stomatal conductance (g_{sCOS}) and environmental parameters. Measurements of both F_{COS-Rn} and F_{COS-EC} pointed to a decrease of COS uptake with decreasing g_{sCOS} and increasing VPD
30 and T_{air} , which is likely related to a dry and warm period in August to which plants responded by closing their stomata to prevent excessive water loss. Our results suggest that the nighttime uptake of COS is mainly driven by the tree foliage and the relationships that we find between F_{COS} , g_{sCOS} , VPD and T_{air} will aid in implementing nighttime COS uptake in models. Both

the EC and the radon-tracer methods indicate that the nighttime sink of COS plays an important role in the total COS budget in a boreal forest and needs to be taken into account when using COS as a tracer for GPP estimates

Data availability. The nighttime ecosystem fluxes of COS and CO₂ obtained through the radon-tracer and eddy-covariance method can be accessed at <https://doi.org/10.5281/zenodo.858625>.

5 *Author contributions.* U.S., K.M., H.C., T.V. and L.M.J.K designed the research. L.M.J.K., K.M., J.A. conducted the field work, W.S., I.M., P.K., A.F, R.V and G.V provided data. L.M.J.K performed data analysis. L.M.J.K. and H.C. wrote the paper with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

Disclaimer. TEXT

10 *Acknowledgements.* We greatly appreciate the maintenance and help of the technical staff at SMEAR II in Hyytiälä, in particular Helmi Keskinen and Janne Levula. We also thank Bert Kers and Marcel de Vries for their help during preparations of the campaign at the University of Groningen. We would like to thank Ute Karstens and Navin Manohar for making F_{Rn} simulations and corresponding data available. We also thank the anonymous reviewers for their comments on the manuscript. The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) in the InGOS project (n°284274), the NOAA contract
15 NA13OAR4310082, the Academy of Finland Centre of Excellence (n°118780), Academy Professor projects (n°284701 and 282842), ICOS-Finland (n°281255) and CARB-ARC (n°286190).

References

- Alekseychik, P., Mammarella, I., Launiainen, S., Rannik, Ü., and Vesala, T.: Evolution of the nocturnal decoupled layer in a pine forest canopy, *Agric. Forest Meteorol.*, 174-175, 15-27, 2013.
- Altimir, N., Kolari, P., Tuovinen, J.-P., Vesala, T., Bäck, J., Suni, T., Kulmala, M., and Hari, P.: Foliage surface ozone deposition: a role for surface moisture?, *Biogeosciences*, 3, 209-228, doi: doi:10.5194/bg-3-209-2006, 2006.
- Asaf, D., Rotenberg, E., Tatarinov, F., Dicken, U., Montzka, S. A., and Yakir, D.: Ecosystem photosynthesis inferred from measurements of carbonyl sulphide flux, *Nat. Geosci.*, 6, 186-190, doi: 10.1038/ngeo1730, 2013.
- Aubinet, M., Chermanne, B., Vandenhaute, M., Longdoz, B., Yernaux, M., and Laitat, E.: Long term carbon dioxide exchange above a mixed forest in the Belgian Ardennes, *Agric. For. Meteorol.*, 108, 293-315, doi: [http://dx.doi.org/10.1016/S0168-1923\(01\)00244-1](http://dx.doi.org/10.1016/S0168-1923(01)00244-1), 2001.
- Aubinet, M., Vesala, T., and Papale, D.: *Eddy Covariance: A Practical Guide to Measurement and Data Analysis*, Springer, Dordrecht Heidelberg London New York, 2012.
- Belviso, S., Schmidt, M., Yver, C., Ramonet, M., Gros, V., and Launois, T.: Strong similarities between night-time deposition velocities of carbonyl sulphide and molecular hydrogen inferred from semi-continuous atmospheric observations in Gif-sur-Yvette, Paris region, *Tellus B*, 65, 2013.
- Berkelhammer, M., Asaf, D., Still, C., Montzka, S., Noone, D., Gupta, M., Provencal, R., Chen, H., and Yakir, D.: Constraining surface carbon fluxes using in situ measurements of carbonyl sulfide and carbon dioxide, *Global Biogeochem. Cy.*, 28, 161-179, doi: 10.1002/2013GB004644, 2014.
- Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., Seibt, U., Stimler, K., Yakir, D., and Zhu, Z.: A coupled model of the global cycles of carbonyl sulfide and CO₂: A possible new window on the carbon cycle, *J. Geophys. Res. Biogeo.*, 118, 842-852, doi: 10.1002/jgrg.20068, 2013.
- Billesbach, D. P., Berry, J. A., Seibt, U., Maseyk, K., Torn, M. S., Fischer, M. L., Mohammad Abu-Naser, and Campbell, J. E.: Growing season eddy covariance measurements of carbonyl sulfide and CO₂ fluxes: COS and CO₂ relationships in Southern Great Plains winter wheat, *Agric. For. Meteorol.*, 184, 48-55, 2014.
- Brühl, C., Lelieveld, J., Crutzen, P. J., and Tost, H.: The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate, *Atmos. Chem. Phys.*, 12, 1239-1253, doi: 10.5194/acp-12-1239-2012, 2012.
- Caird, M. A., Richards, J. H., and Donovan, L. A.: Nighttime Stomatal Conductance and Transpiration in C₃ and C₄ Plants, *Plant Physiology*, 143, 4-10, doi: 10.1104/pp.106.092940, 2007.
- Campbell, J. E., Carmichael, G. R., Chai, T., Mena-Carrasco, M., Tang, Y., Blake, D. R., Blake, N. J., Vay, S. A., Collatz, G. J., Baker, I., Berry, J. A., Montzka, S. A., Sweeney, C., Schnoor, J. L., and Stanier, C. O.: Photosynthetic Control of Atmospheric Carbonyl Sulfide During the Growing Season, *Science*, 322, 1085-1088, doi: 10.1126/science.1164015, 2008.
- Chin, M. and Davis, D.: A Reanalysis of Carbonyl Sulfide as a Source of Stratospheric Background Sulfur Aerosol, *Journal of Geophysical Research-Atmospheres*, 100, 8993-9005, doi: 10.1029/95JD00275, 1995.
- Commane, R., Herndon, S. C., Zahniser, M. S., Lerner, B. M., McManus, J. B., Munger, J. W., Nelson, D. D., and Wofsy, S. C.: Carbonyl sulfide in the planetary boundary layer: Coastal and continental influences, *J. Geophys. Res. Atmos.*, 118, 8001-8009, doi: 10.1002/jgrd.50581, 2013.
- Commane, R., Meredith, L. K., Baker, I. T., Berry, J. A., Munger, J. W., Montzka, S. A., Templer, P. H., Juice, S. M., Zahniser, M. S., and Wofsy, S. C.: Seasonal fluxes of carbonyl sulfide in a midlatitude forest, *P. Natl. Acad. Sci. USA*, doi: 10.1073/pnas.1504131112, 2015.

- Crutzen, P. J.: The possible importance of CSO for the sulfate layer of the stratosphere, *Geophys. Res. Lett.*, 3, 73-76, doi: 10.1029/GL003i002p00073, 1976.
- Finkelstein, P. L. and Sims, P. F.: Sampling error in eddy correlation flux measurements, *J. Geophys. Res.*, 106, 3503-3509, 2001.
- Hari, P. and Kulmala, M.: Station for Measuring Ecosystem - Atmosphere Relations (SMEAR II), *Boreal Environ. Res.*, 10, 315-322, 2005.
- 5 Karstens, U., Schwingshackl, C., Schmithüsen, D., and Levin, I.: A process-based 222radon flux map for Europe and its comparison to long-term observations, *Atmos. Chem. Phys.*, 15, 12845-12865, doi: 10.5194/acp-15-12845-2015, 2015.
- 1 Kettle, A., Kuhn, U., von Hobe, M., Kesselmeier, J., and Andreae, M.: Global budget of atmospheric carbonyl sulfide: Temporal and spatial variations of the dominant sources and sinks, *J. Geophys. Res. Atmos.*, 107, 4658, doi: 10.1029/2002JD002187, 2002.
- Kolari, P., Kulmala, L., Pumpanen, J., Launiainen, S., Ilvesniemi, H., Hari, P., and Nikinmaa, E.: CO₂ exchange and component CO₂ fluxes of a boreal Scots pine forest. *Boreal Environ. Res.*, 14, 761-783, 2009.
- 10 Kooijmans, L. M. J., Uitslag, N. A. M., Zahniser, M. S., Nelson, D. D., Montzka, S. A., and Chen, H.: Continuous and high-precision atmospheric concentration measurements of COS, CO₂, CO and H₂O using a quantum cascade laser spectrometer (QCLS), *Atmos. Meas. Tech.*, 9, 5293-5314, doi: 10.5194/amt-9-5293-2016, 2016.
- Launois, T., Belviso, S., Bopp, L., Fichot, C. G., and Peylin, P.: A new model for the global biogeochemical cycle of carbonyl sulfide - Part 1: Assessment of direct marine emissions with an oceanic general circulation and biogeochemistry model, *Atmos. Chem. Phys.*, 15, 2295-2312, doi: 10.5194/acp-15-2295-2015, 2015.
- 15 López-Coto, J., Mas, J. L., and Bolívar, J. P.: A 40-year retrospective European radon flux inventory including climatological variability, *Atmos. Environ.*, 73, 22-33, doi: doi:10.1016/j.atmosenv.2013.02.043, 2013.
- Mammarella, I., Kolari, P., Rinne, J., Keronen, P., Pumpanen, J., and Vesala, T.: Determining the contribution of vertical advection to the net ecosystem exchange at Hyytiälä forest, Finland, *Tellus B*, 59, 900-909, doi: 10.1111/j.1600-0889.2007.00306.x, 2007.
- 20 Mammarella, I., Launiainen, S., Gronholm, T., Keronen, P., Pumpanen, J., Rannik, Ü., and Vesala, T.: Relative humidity effect on the high frequency attenuation of water vapour flux measured by a closed-path eddy covariance system, *J. Atmos. Ocean. Tech.*, 26, 1856-1866, 2009.
- Mammarella, I., Werle, P., Pihlatie, M., Eugster, W., Haapanala, S., Kiese, R., Markkanen, T., Rannik, Ü., and Vesala, T.: A case study of eddy covariance flux of N₂O measured within forest ecosystems: quality control and flux error analysis, *Biogeosciences*, 7, 427-440, doi: 10.5194/amt-9-4915-2016, 2010.
- 25 Mammarella, I., Peltola, O., Nordbo, A., Järvi, L., and Rannik, Ü.: Quantifying the uncertainty of eddy covariance fluxes due to the use of different software packages and combinations of processing steps in two contrasting ecosystems, *Atmos. Meas. Tech.*, 9, 4915-4933, doi: 10.5194/amt-9-4915-2016, 2016.
- 30 Manohar, S. N., Meijer, H. A. J., and Herber, M. A.: Radon flux maps for the Netherlands and Europe using terrestrial gamma radiation derived from soil radionuclides, *Atmos. Environ.*, 81, 399-412, doi: http://dx.doi.org/10.1016/j.atmosenv.2013.09.005, 2013.
- Marcazzan, G. M., Caprioli, E., Valli, G., and Vecchi, R.: Temporal variation of ²¹²Pb concentration in outdoor air of Milan and a comparison with ²¹⁴Bi, *J. Environ. Radioact.*, 65, 77-90, doi: http://dx.doi.org/10.1016/S0265-931X(02)00089-9, 2003.
- Maseyk, K., Berry, J. A., Billesbach, D., Campbell, J. E., Torn, M. S., Zahniser, M., and Seibt, U.: Sources and sinks of carbonyl sulfide in an agricultural field in the Southern Great Plains, *P. Natl. Acad. Sci. USA*, 2015 112 (46) 14162-14167, doi: 10.1073/pnas.1319132111, 2014.
- 35

- Montzka, S. A., Calvert, P., Hall, B. D., Elkins, J. W., Conway, T. J., Tans, P. P., and Sweeney, C.: On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO₂, *J. Geophys. Res. Atmos.*, 112, D09302, doi: 10.1029/2006JD007665, 2007.
- Nieminen, T., Asmi, A., Dal Maso, M., Aalto, P.P., Keronen, P., Petäjä, T., Kulmala, M., Kerminen, V.-M.: Trends in atmospheric new-particle formation: 16 years of observations in a boreal-forest environment, *Boreal Environ. Res.*, 19, 191-214, 2014.
- Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., Longdoz, B., Rambal, S., Valentini, R., Vesala, T., and Yakir, D.: Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation, *Biogeosciences*, 3, 571-583, doi: 10.5194/bg-3-571-2006, 2006.
- Protoschill-Krebs, G., Wilhelm, C., and Kesselmeier, J.: Consumption of carbonyl sulphide (COS) by higher plant carbonic anhydrase (CA), *Atmos. Environ.*, 30, 3151-3156, doi: 10.1016/1352-2310(96)00026-X, 1996.
- Rannik, Ü.: On the surface layer similarity at a complex forest site, *Journal of Geophysical Research: Atmospheres*, 103, 8685-8697, doi: 10.1029/98JD00086, 1998.
- Rannik, Ü., Keronen, P., Hari, P., and Vesala, T.: Estimation of forest-atmosphere CO₂ exchange by eddy covariance and profile techniques, *Agric. Forest Meteorol.*, 126, 141-155, 2004.
- Rannik, Ü., Peltola, O., and Mammarella, I.: Random uncertainties of flux measurements by the eddy covariance technique, *Atmos. Meas. Tech.*, 9, 5181, doi: 10.5194/amt-9-5163-2016, 2016.
- Sandoval-Soto, L., Stanimirov, M., von Hobe, M., Schmitt, V., Valdes, J., Wild, A., and Kesselmeier, J.: Global uptake of carbonyl sulfide (COS) by terrestrial vegetation: Estimates corrected by deposition velocities normalized to the uptake of carbon dioxide (CO₂), *Biogeosciences*, 2, 125-132, 2005.
- Schmidt, M., Graul, R., Sartorius, H., Levin, I.: Carbon dioxide and methane in continental Europe: A climatology, and ²²²Rn-based emission estimates, *Tellus B*, 48B, 457-473, 1996.
- Seibt, U., Kesselmeier, J., Sandoval-Soto, L., Kuhn, U., and Berry, J. A.: A kinetic analysis of leaf uptake of COS and its relation to transpiration, photosynthesis and carbon isotope fractionation, *Biogeosciences*, 7, 333-341, doi: 10.5194/bg-7-333-2010, 2010.
- Sesana, L., Caprioli, E., and Marazzan, G. M.: Long period study of outdoor radon concentration in Milan and correlation between its temporal variations and dispersion properties of atmosphere, *J. Environ. Radioact.*, 65, 147-160, doi: [http://dx.doi.org/10.1016/S0265-931X\(02\)00093-0](http://dx.doi.org/10.1016/S0265-931X(02)00093-0), 2003.
- Stimler, K., Berry, J. A., Montzka, S. A., and Yakir, D.: Association between Carbonyl Sulfide Uptake and $\delta^{13}C$ during Gas Exchange in C-3 and C-4 Leaves, *Plant Physiol.*, 157, 509-517, doi: 10.1104/pp.111.176578, 2011.
- Sun, W., Kooijmans, L.M.J., Maseyk, K., Chen, H., Mammarella, I., Vesala, T., Levula, J., Keskinen, H., Seibt, U.: Soil fluxes of carbonyl sulfide (COS), carbon monoxide, and carbon dioxide in a boreal forest in southern Finland, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2017-180, in review, 2017.
- Szegvary, T., Leuenberger, M. C., and Conen, F.: Predicting terrestrial ²²²Rn flux using gamma dose rate as a proxy, *Atmos. Chem. Phys.*, 7, 2789-2795, doi: 10.5194/acp-7-2789-2007, 2007.
- Szegvary, T., Conen, F., and Ciais, P.: European ²²²Rn inventory for applied atmospheric studies, *Atmos. Environ.*, 43, 1536-1539, doi: 10.1016/j.atmosenv.2008.11.025, 2009.
- Van der Laan, S., Neubert, R. E. M., and Meijer, H. A. J.: Methane and nitrous oxide emissions in The Netherlands: ambient measurements support the national inventories, *Atmos. Chem. Phys.*, 9, 9369-9379, doi: 10.5194/acp-9-9369-2009, 2009.

- Van der Laan, S., Manohar, S., Vermeulen, A., Bosveld, F., Meijer, H., Manning, A., van, d. M., and van, d. L.: Inferring ^{222}Rn soil fluxes from ambient ^{222}Rn activity and eddy covariance measurements of CO_2 , *Atmos. Meas. Tech.*, 9, 5523-5533, doi: 10.5194/amt-9-5523-2016, 2016.
- 5 Vickers, D. and Mahrt, L.: Quality control and flux sampling problems for tower and aircraft data, *J. Atmos. Ocean. Tech.*, 14, 512-526, 1997.
- Watts, S. F.: The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide, *Atmos. Environ.*, 34, 761-779, 2000.
- Wehr, R., Commane, R., Munger, J. W., McManus, J. B., Nelson, D. D., Zahniser, M. S., Saleska, S. R., and Wofsy, S. C.: Dynamics of canopy stomatal conductance, transpiration, and evaporation in a temperate deciduous forest, validated by carbonyl sulfide uptake, *Biogeosciences*, 14, 389-401, doi: 10.5194/bg-14-389-2017, 2017.
- 10 White, M. L., Zhou, Y., Russo, R. S., Mao, H., Talbot, R., Varner, R. K., and Sive, B. C.: Carbonyl sulfide exchange in a temperate loblolly pine forest grown under ambient and elevated CO_2 , *Atmos. Chem. Phys.*, 10, 547-561, doi: 10.5194/acp-10-547-2010, 2010.
- Winderlich, J., Gerbig, C., Kolle, O., and Heimann, M.: Inferences from CO_2 and CH_4 concentration profiles at the Zotino Tall Tower Observatory (ZOTTO) on regional summertime ecosystem fluxes, *Biogeosciences*, 11, 2055-2068, doi: 10.5194/bg-11-2055-2014, 2014.
- 15 Wohlfahrt, G., Brilli, F., Hoernagl, L., Xu, X., Bingemer, H., Hansel, A., and Loreto, F.: Carbonyl sulfide (COS) as a tracer for canopy photosynthesis, transpiration and stomatal conductance: potential and limitations, *Plant Cell Environ.*, 35, 657-667, doi: 10.1111/j.1365-3040.2011.02451.x, 2012.




Figure 1. COS, CO₂ and ²²²Rn concentrations, u* and the storage flux of COS and CO₂ (F_{stor-COS} and F_{stor-CO₂}) during 12-13 July 2015 where the data with sun elevation below 0° are used to derive nighttime fluxes of COS and CO₂ (black, filled). The bottom figures show the linear regression between ²²²Rn and COS (left) and CO₂ concentrations (right) on which F_{COS-Rn} and NEE_{Rn} are based.




Figure 2. Storage fluxes F_{stor} (green), ecosystem fluxes NEE_{EC} and F_{COS-EC} (red) and soil fluxes F_{soil} (blue) of COS (left) and CO_2 (right) in summer (July and August) 2015. Thick lines indicate the median values of the data over the whole measurement period, and the shaded areas specify the 25th-75th percentiles. The median values of NEE_{EC} and F_{COS-EC} without storage correction are shown in gray. The ecosystem fluxes are filtered for low u_* values with a threshold of 0.3 m s^{-1} .




Figure 3. Relative frequency of R^2 values of the correlation between concentrations of ^{222}Rn and CO_2 (top) and COS (bottom).




Figure 4. Left: comparison of EC- and radon-based fluxes for average nighttime CO₂ (top) and COS (bottom) fluxes. Middle and right: time series of EC based fluxes (middle) and radon-based fluxes (right). The uncertainty bars of the EC and radon-based fluxes are not directly comparable due to the different ways of determining these uncertainties.




Figure 5. Correlations of $F_{\text{COS-Rn}}$ with g_{SCOS} , T_{air} , VPD, and u_* . All data (except $F_{\text{COS-Rn}}$) are averages over individual nights (with nighttime defined as sun elevation below -3°). Data in this plot largely represent a period in August 2015 with dry conditions (i.e. decreasing SWC, and increasing T_{air} and VPD).