

Response to Anonymous referee #1

- *It is still unclear what the (revised) description of the surface layer model humidity means ["soaking wet (30% water volume mixing ratio)"], lines 195-196. Is that what is sometimes referred to as surface moisture availability (e.g., 1 if the surface corresponds to the ocean and smaller value over land)? In a simple parameterization of the surface vapor flux, the surface temperature prescribes the water vapor mixing ratio as corresponding to saturation at the surface temperature and smaller a value over land. If I want to repeat the authors experiment I need to know what to use and currently I am left guessing. The explanation what C_{skin} is still confusing, line 203, table 1. I looked at Heus et al. GMD 2010 and there I see C_{sk} that is different to what the authors use in the paper under review. I think the units and values given in the table imply that C_{skin} is simply the depth of a uniform-temperature layer of water for which the energy balance equation is given by $\rho_w c_w H dT/dt = \text{net surface energy flux}$. That is, C_{skin} is the same as H in my formula. A complication is that Heus et al. also state that the surface model uses 4 layers. I think the authors need to be very specific what they use [1 layer, 4 layers? constant temp and moisture availability (or whatever variable is used)?]. Again, I would not be able to setup my own simulation following the authors' explanations.*

I rewrote the explanation to further clarify the setup. We use four, saturated with respect to water, initially uniform layers, plus a skin layer on top with a varying heat capacity. Our C_{skin} is the C_{sk} of Heus et al. (2010). The latent heat release is solved by the land surface model in UCLA-LES, as described in Heus et al. (2010). Furthermore, for a full reference, I added the input files for the experiments to the code repository which is now mentioned in a Code-Availability section.

- *Additional analysis: I feel that the paper has been improved and it is close to be accepted. However, I still feel a couple details in the explanation of the model setup need to be corrected. I provide below a short list what I feel would be needed in association with Fig. 1. Figure 1 is quite impressive, but I wonder if the difference in cloud sizes (smaller in the lower panel to my eye) comes from difference in the boundary depth and/or radiative cooling (BTW, the radiative cooling is the "destabilization rate" in 5th bullet in specific comments section of my previous review; the authors were confused with what I meant). I feel that at least the lower-tropospheric moisture and temperature profiles, cloud fraction profiles, as well as mean radiative cooling should be shown together with the two panels. If they are really close, than please say so in the text to emphasize that other factors (like the BL depth) are practically the same in both simulations. But showing a figure would be better.*

Yes, you are correct, the two simulations are not the same. The simulations start with the same initial profiles but then diverge. 3D radiative transfer illuminates the surface for an elongated period of time and tends to strengthen the updraft, leading to an invigorated growth of clouds. To compare our simulations with varying parameters, I think, we have two choices. One, we try to homogenize the forcing that drives the simulations (e.g. renormalize the surface fluxes to a fixed value) or, secondly, we don't evaluate the individual course of each simulation but rather look at the qualitative distribution of say, cloud streets.

I feel that the approach of homogenizing the fluxes would be rather difficult. Renormalized surface fluxes for simulations with large zenith angles for example would certainly introduce very high (physically unrealistic) peaks in illumination. I am not saying that these types of simulations are not interesting but I prefer to keep the interactions of the radiative transfer methods as physically sound as possible. While I think that it would be great if one could disentangle the various mechanisms (including the dependency on boundary layer height etc.), we feel that this study does already a great job in showing that there is a link between radiative transfer and the organization of convection through dynamic heterogeneities in surface fluxes. Irrespective of boundary layer characteristics, the fact that 1D simulations do not produce cloud streets, remains. Another example would be the fact that 3D simulations with a solar azimuth of $\phi = 90^\circ$ and $\phi = 180^\circ$ have the same boundary layer characteristics and cloud statistics but show the opposite sign in the autocorrelation ratio (i.e. orientation of streets). If you are interested in the evolution of individual simulations of shallow cumulus convection, pertaining 1D and 3D radiative transfer, maybe I can excite your interest in Jakub (2016, sec 3.2).

Response to Anonymous referee #2

- *I my first review I requested discussion of advected boundary layers. This was not included in the revised version. The advection of boundary layers adds significant complexity to the problem. Please add a few words in the closing paragraph (Moving forward...) along with an appropriate reference.*

Indeed, the interplay of static and dynamic heterogeneities in conjunction with horizontal advection is a fascinating topic. We tried to keep the focus purely on the radiative effects (and as simple as possible) and did not delve into these aspects but we agree that it would be great to revisit earlier studies that tackled static heterogeneities in windy conditions. I added a paragraph to the outlook steering towards that line of research.

- *A technical editor should decide on orthogonal vs orthogonally and perpendicular vs. perpendicularly. I suspect it is the former in both cases.*

I would appreciate that. I am not sure which is more appropriate either. I asked a native colleague and he suspects that the adverb form is the grammatically correct but that in case of perpendicular, he deems the adjective form to sound better.

- *Line 177 (marked version) change infer to cause*
done.

- *Line 203 situations*
done.

- *Line 204 a well mixed ocean*
done.

- *Line 244 liquid*
expanded.

- *Line 295 first develops*

Indeed, what I meant was that first clouds develop after half an hour... I rewrote the sentence.

- *Caption figure 4: remove "Shown is"*

Yes, done.

- *Line 441, remove ; and replace with)*

The ; between citations is the default style. The ; at the end of the line in the *diff* version is apparently a hiccup of latexdiff. It is fixed in the normal version.

- *Line 543 remove typically (the uncertainty is already conveyed by 'order of')*

done.

The Role of 1D and 3D Radiative Heating in the Organization of Shallow Cumulus Convection and the Formation of Cloud Streets

Jakub Fabian¹ and Mayer Bernhard¹

¹Meteorological Institute, Ludwig Maximilian Universität München, LMU

Correspondence to: Jakub Fabian (fabian.jakub@physik.uni-muenchen.de)

Abstract. The formation of shallow cumulus cloud streets was historically attributed primarily to dynamics. Here, we focus on the interaction between radiatively induced surface heterogeneities and the resulting patterns in the flow. Our results suggest that solar radiative heating has the potential to organize clouds ~~perpendicularly~~ perpendicular to the sun's incidence angle.

5 To quantify the extent of organization, we performed a high resolution LES parameter study. We varied the horizontal wind speed, the surface heat capacity, the solar zenith and azimuth angles, as well as radiative transfer parameterizations (1D and 3D). As a quantitative measure we introduce a simple algorithm that provides a scalar quantity for the degree of organization and the alignment. We find that, even in the absence of a horizontal wind, 3D radiative transfer produces cloud streets per-

10 pendicular to the sun's incident direction, whereas the 1D approximation or constant surface fluxes produce circular, randomly positioned, clouds. Our reasoning for the enhancement or reduction of organization is the geometric position of the cloud's shadow and its corresponding surface fluxes. Furthermore, when increasing horizontal wind speeds to 5 or 10 m s⁻¹, we observe the development of dynamically induced cloud streets. If in addition, solar radiation illuminates the surface beneath

15 the cloud, i.e. when the sun is positioned orthogonally to the mean wind field and the solar zenith angle is larger than 20°, the cloud-radiative feedback has the potential to significantly enhance the tendency to organize in cloud streets. In contrast, in the case of the 1D approximation (or overhead sun), the tendency to organize is weaker or even prohibited because the shadow is cast directly be-

20 neath the cloud. In a land-surface type situation, we find the organization of convection happening on a timescale of half an hour. The radiative feedback, creating surface heterogeneities is generally diminished for large surface heat capacities. We therefore expect radiative feedbacks to be strongest over land surfaces and weaker over the ocean. Given the results of this study we expect that sim-

25 ulations including shallow cumulus convection will have difficulties producing cloud streets if they employ 1D radiative transfer solvers or may need unrealistically high wind speeds to excite cloud street organization.

1 Introduction

The advent of airborne and satellite observations allow for a bird's eye view of the atmosphere and, ever since, meteorologists have been fascinated by the striped patterns often evident in cloud systems. Kuettner (1959) presented some early pictures of cloud streets from rocket and aircraft instruments. Descriptions of cloud streets, date back as far as Steinhoff (1935), who gave a detailed description of a long-distance glider flight, or Woodcock (1942) who investigated the soaring patterns of seagulls. Scientific literature documenting the existence and explaining the prerequisites for the formation of cloud streets is plentiful. Brown (1980); Etling and Brown (1993); Weckwerth et al. (1997); Houze Jr (2014) provide a thorough review of past observations and theoretical frameworks. The above literature suggests two prominent effects to be responsible for such vortices, namely inflection-point instabilities (e.g. from cross-roll wind components in a Ekman boundary layer) and thermal instabilities (buoyancy driven). Purely buoyancy driven convection, without any horizontal wind or shear, produces a random pattern of updrafts. Introducing a linear wind shear, the convective elements become stretched out along-wind. Following Grossman (1982): "*At some point (increasing the wind speed/shear) the shearing becomes strong enough so that dynamic instability may interact with buoyancy to produce a hybrid roll vortex/convective cell mechanism. As the shear becomes stronger, shearing instability or roll vortex motion is predominant.*" In this work, we will focus on the radiative impact, with the most prominent effect being cloud shadows which modulate surface fluxes and consequently build up surface heterogeneities. These induced surface heterogeneities are the link between radiative transfer and buoyancy driven convection (Lohou and Patton, 2014; Horn et al., 2015; Gronemeier et al., 2016). Our focus is therefore more on buoyancy driven roll vortices in a linear shear environment (Asai, 1970) and less so on inflection-point instabilities. To that end, we omit cross-wind shear by neglecting Coriolis force and correspondingly neglect the horizontal turning of the wind as it would be the case in an Ekman boundary layer. Several studies investigated the role of surface fluxes on the development of such boundary layer circulations. Here the literature distinguishes between static heterogeneities, i.e. differences in land-surface parameters such as vegetation, surface roughness or surface albedo and dynamic heterogeneities, such as moisture budget or temperature fluctuations. Static heterogeneities in conjunction with shallow cumulus clouds and cloud streets have been examined for example by Avissar and Schmidt (1998); Patton et al. (2005); Rieck et al. (2014). In contrast, Schumann et al. (2002); Wapler (2007); Frame et al. (2009); Gronemeier et al. (2016) investigated the influence of dynamic heterogeneities in surface shading and even considered 3D radiative effects (i.e. the displacement of the shadow). However, they did not include a realistic surface model, but rather adjusted the surface fluxes instantaneously. This does not allow to study the timescales on which radiation and dynamics may interact. Others investigated the influence of shading coupled to an interactive surface model (Vilà-Guerau de Arellano et al., 2014; Lohou and Patton, 2014; Horn et al., 2015). However, one particularly questionable issue with those studies was the application of 1D radiative transfer solvers, which are known to introduce large spa-

tial error in surface heating rates (O’Hirok and Gautier, 2005; Wapler and Mayer, 2008; Wissmeier et al., 2013; Jakub and Mayer, 2015).

65 Overall, we can summarize that the formation of cloud streets has been extensively explored from theoretical and observational perspectives. The above mentioned studies shed light on the various aspects of interaction with the cloud field but either lack a realistic representation of surface processes, neglect 3D radiative transfer effects or do not examine the relationship concerning the background wind speed.

70 In this study we strive to overcome these shortcomings and determine the prerequisites for the formation of cloud streets, while our main focus lies on dynamic heterogeneities and (3D) radiative transfer. We try to disentangle the underlying processes with a rigorous parameter study using Large-Eddy-Simulations (LES).

Section 2 briefly outlines the LES model, explains the setup of the simulations and introduces a scalar metric to quantify the organization in cloud streets. In section 3 we interpret the magnitude of cloud street formations in the parameter space spanning surface properties, background wind speeds and the sun’s angles. Section 4 finally summarizes key findings of the parameter study.

2 Methods and Experiments

2.1 LES Model

80 The Large-Eddy-Simulations (LES) were performed with the UCLA–LES model. A description and details of the LES model can be found in Stevens et al. (2005). The land surface model included in the UCLA–LES follows the implementation of the Dutch Atmospheric Large-Eddy Simulation code Heus et al. (2010). The simulations presented here use warm micro-physics formulated in Seifert and Beheng (2001) where the formation of rain is turned off to prevent any further complications such as cold pool dynamics. The radiative transfer calculations are performed with the

85 TenStream package (Jakub and Mayer, 2015), which includes a 1D Schwarzschild (thermal only), a δ -Eddington two-stream (solar and thermal), as well as the 3D TenStream (solar and thermal) solver.

The TenStream is a MPI-parallelized solver for the full 3D radiative transfer equation. In analogy to a two-stream solver, the TenStream solver computes the radiative transfer coefficients for up- and downward fluxes and additionally for sideward streams. The coupling of individual boxes leads to

90 a linear equation system which is written as a sparse matrix and is solved using parallel iterative methods from the “Portable, Extensible Toolkit for Scientific Computation”, PETSc (Balay et al., 2014) framework. In Jakub and Mayer (2015, 2016), we extensively validated the TenStream by comparison with the exact Monte Carlo code *MYSTIC* (Mayer, 2009).

95 The most pronounced differences between 1D and 3D radiative transfer solvers, pertaining the setup here, is the displacement of the sun’s shadow at the surface. In the case of 1D radiative transfer, the shadow of a cloud is by definition always directly beneath it (so called independent pixel



Figure 1. Virtual photograph of LES simulations at a cruising altitude of 15 km. Top panel: cloud formation of a simulation driven by 3D radiation (TenStream with sun in the east, i.e. right ($\varphi = 90^\circ$)). Lower-panel: cloud formation of a simulation which was performed with 1D radiation (Two-stream). The specific model setup is the same as referenced in fig. 2, i.e., no background wind and a continental land surface. The simulations differ with respect to cloud size distributions and the organization in cloud streets, the cloud fraction though is the same (27%). The visualization was performed with a physically correct rendering with MYSTIC (MonteCarlo solver in libRadtran (Mayer, 2009; Emde et al., 2015)).

or independent column approximation). Contrarily, 3D radiative transfer allows the propagation of energy horizontally and correctly displaces the clouds shadow depending on the sun's position. The features of 3D radiative transfer in the thermal spectral range are an increased cooling on cloud edges and a smoothed distribution of surfaces fluxes. While we compute thermal radiative transfer in a 3D fashion, we expect these effects to be less important for this setup because feedbacks on the dynamics appear to happen only longer timescales of a day (Klinger et al., 2017) and more importantly because it does not infer-cause any asymmetries in the heating or cooling pattern.

The spectral integration is performed using the correlated-k method following Fu and Liou (1992). The coupling of the TenStream solver to the UCLA-LES follows the description in Jakub and Mayer (2016). One exception is the use of the Monte-Carlo-Spectral-Integration (Pincus and Stevens, 2009) which we do not use because of limitations with regards to computations involving interactive surface models (Pincus and Stevens, 2013).

2.2 Model Experiment Setup

The base setup of the UCLA-LES simulates a domain of $50 \text{ km} \times 50 \text{ km}$ with a horizontal grid length of 100 m and 50 m vertically. The simulations start from a well-mixed initial background profile with a constant virtual potential temperature (292 K) in the lower 700 m and increases by

Table 1. Parameter space for the LES simulations: the mean west wind \mathbf{u} , the solar azimuth and zenith angle φ, θ , the surface skin heat capacity C_{skin} as a water column equivalent and three settings for the computation of net radiative surface fluxes (Q_{net}). The radiative transfer computations are done either with a 1D δ -Eddington two-stream , with the 3D TenStream solver or simulations with constant mean net irradiance. Variations of the solar azimuth φ are only applied for 3D radiative transfer. Values of Q_{net} in case of simulations without interactive radiative transfer were set to the mean surface irradiance of the 1D simulations. In total a number of 192 simulations.

\mathbf{u}	0, 5, 10	m s^{-1}
φ	90, 180	$^{\circ}$
θ	20, 40, 60, 75	$^{\circ}$
C_{skin}	1, 10, 100, 1000	cm
Q_{net}	constant, 1D, 3D	

6 K km^{-1} above. Water vapor near the surface amounts to 9.5 g kg^{-1} , decreasing with $-1.3 \text{ g kg}^{-1} \text{ km}^{-1}$.

115 The ~~layers of the surface model~~ surface model has four layers which have the same initial temperature of 291 K, are stripped of vegetation and are soaking wet (~~30% water volume mixing ratio~~) and are stripped of vegetation with an initial temperature of 291 K saturated clay with 30% water volume mixing ratio. The surface albedo for shortwave radiation is set to 7%. The land-surface model solves the surface energy balance equation for an imaginary skin layer which often has no heat capacity. We ~~manipulate~~
 120 vary the heat capacity of the surface skin layer C_{skin} to mimic a water layer covering the surface. The heat capacities are chosen to be representative for ~~situation~~ situations ranging from continental land surfaces to a well mixed ocean. The thickness of this imaginary water layer lends the simulations and the radiative transfer a memory on the surface. All other parameters of the land-surface model such as surface resistances or roughness lengths for momentum or heat are kept constant in order to
 125 focus on these memory effects.

The focus of this study is to determine the interplay of radiation with the atmosphere, the surface and the clouds, and finally take a closer look on the formation of cloud streets. To that end we run the simulations with five free parameters, namely the heat capacity of the surface skin layer (C_{skin}), the background wind (\mathbf{u} , i.e. west-winds), the solar zenith (θ) and azimuth (φ) angle as well as with
 130 different radiative transfer approximations (see table 1). The coupling of radiative transfer to the land-surface model is realized in four ways. We either compute the net surface irradiance Q_{net} with a 1D δ -eddington two-stream solver , or employ the 3D TenStream solver, with two azimuth angles. Additionally, we conducted the experiments where Q_{net} is set to a prescribed constant value (spatial and temporal average of the surface irradiance of the corresponding 1D simulation).

135 The time it takes the simulations to form the first clouds depends on the choice of the parameters. Foremost the solar zenith angle determines the energy input into the atmosphere and the sur-

face (lower positioned sun hence leads to a later onset of cloud development). To compare the heterogeneous simulations we limit the following analysis to the time-steps (output every 5 min) where the cloud fraction is between 10 % and 50 % (typical for shallow cumulus convection, e.g. Seifert and Heus (2013)). Most simulations produce clouds after about one hour and show an increase in cloud cover up to and beyond 50 % in the first 6 h. Simulations with low positioned sun took longer and were hence run for a longer period of 12 h. Our analysis is mostly independent of the specific, individual course of each simulation as we find robust signals across the various groups of parameters. The interested reader, however, is referred to Jakub (2016, sec. 3.2) for further details (e.g. [Fig. liquid](#) water path, cloud fraction, mean cloud size distribution) on the evolution of a typical simulation.

Figure 1 shows a photo rendering of the LES cloud field for two simulations with differing options for the radiative transfer solver. In the top panel, 3D radiative transfer is considered with the sun positioned in the east (zenith $\theta = 60^\circ$) and in the bottom panel panel 1D solver is applied where the shadow is by definition always cast directly beneath the clouds. In the former the organization in cloud streets perpendicular to the sun’s incident angle is evident whereas the latter (1D) does not seem to organize in any way. Figure 2 presents the liquid water content and the surface heat flux for the same two simulations plus one 3D simulation where the sun is in the south. This time we look at volume rendered liquid water content and surface heat fluxes for the full domain. In figs. 1 and 2, simulations with 3D radiative transfer show organization in cloud streets with length scales of up to 20 km, perpendicular to the sun’s incident angle. We can clearly identify these coherent cloud structures with the naked eye. However, to solidify our claims, we present a quantitative measure for the cloud distribution.

2.3 Correlation Ratio

Since we do not deal with towering and tilted or multilayer clouds we can use the cloud mask as a proxy to separate individual clouds. We derive the cloud mask as the binary field of the liquid water path ($LWP > 0$). We then use the normalized 2D auto correlation function of the cloud mask to analyze the spatial distribution of cloudy and clear-sky patches. The three upper panels of fig. 3 illustrate the 2D correlation coefficient for the three simulations presented in fig. 2.

Next, we use the transects of the correlation coefficient along the x- and y-axis (indicated as a black line). The lower panels in fig. 3, respectively, show the linearly interpolated line-cuts of the discrete auto-correlation function. The location where the normalized correlation coefficients goes to zero defines the mean distance from a cloudy pixel where it is more likely to find a clear-sky pixel. We use the north-south and the east-west distances d_{NS} and d_{EW} , respectively, to define the correlation ratio R_c as:

$$R_c = d_{NS}/d_{EW}$$

This definition would miss cloud streets in diagonal direction which, however, is no limitation for our analysis. For one, we know that the background wind induces cloud streets along the mean wind




Figure 2. Volume rendered liquid water mixing ratio (LWC) and surface latent and sensible heat flux ($L + H$) for three simulations. The cloud scene of the left and mid panel have already been presented in fig. 1. In the left panel, radiative transfer calculations are performed with the TenStream solver and the sun is positioned in the east ($\varphi = 90^\circ$). The simulation in the mid-panel is driven by a 1D two-stream solver, whereas the right panel simulation also employs the TenStream solver but the sun shining from the south ($\varphi = 180^\circ$). The solar zenith angle is in all three simulations $\theta = 60^\circ$, the mean background wind speed is 0 ms^{-1} and the surface skin heat capacity set to an equivalent of 1 cm water depth (representative for continental land surface). The snapshot shows the simulations after 3 h model time at a cloud fraction of 27%. Volume rendered plots were created with VISIT (Childs et al., 2012).

direction, i.e. here in the west-east component (see e.g. Weckwerth et al. (1997)). At the same time we hypothesize that radiatively induced effects will be either along or perpendicular to the incident solar beam, i.e. follow the surface inhomogeneities (see, e.g., Gronemeier et al. (2016)). The two major directions should therefore capture the dominant effects of dynamically and radiatively induced cloud dynamics.

The correlation ratio reduces a cloud field snapshot into a scalar which yields $R_c = 1$ for symmetrically distributed clouds, $R_c < 1$ for organized cloud fields along the north-south direction and $R_c > 1$ if cloud features are arranged east to west.

3 Results and Discussion

As an example for the evolution of convective organization, fig. 4 illustrates the correlation ratio R_c over time for one of the earlier introduced simulations (depicted in fig. 2). **The simulation develops** In the simulation, first cumulus clouds occur after about half an hour with the clouds being oriented randomly. The resulting shadowing of these clouds introduces surface temperature heterogeneities which in turn act on the flow through changes in latent and sensible heat fluxes. About one hour after the onset of clouds, we find the convection to organize into bands from north to south ($R_c < 1$).




Figure 3. The panels exemplarily depict the auto-correlation coefficients of the cloud distribution in the three simulations presented in fig. 2. The upper panels show the normalized 2D autocorrelation coefficient with two intersection lines in the North-South (vertical) and the East-West (horizontal) direction. The markers pinpoint the distance in N-S (red) and E-W (blue) direction, respectively, where the auto-correlation coefficient reaches a zero value and therefore denote the distance where it becomes more likely not to find a cloud. The lower panels follow the black line-cuts and further describe the two transects depicting the correlation function’s root points from which we derive the correlation ratio. Simulations with 3D radiative transfer (left and right panels) shows in contrast to 1D radiative transfer (mid panel) a distinct asymmetry perpendicular to the solar incidence angle. The organization of clouds and their alignment is represented in values of the correlation ratio R_c being less than or greater than one for alignment along the y- or x-axis, respectively.

To further highlight the involved timescales, we restart the simulation from 2 h onwards with a 90° rotated sun and find that convection changes from a north to south orientation to bands from east to west in approximately one hour. This example yields a $1/e$ timescale for convective organization of half an hour. This timescale will however, depend on several factors, most certainly on the solar zenith angle and the surface heat capacity which determine the timescales at which surface heterogeneities can be introduced.

To reduce the information of convective organization into a single scalar value, we compute the mean correlation ratio $\overline{R_c}$ as the arithmetic mean of R_c calculated at all output time-steps (every 5 minutes) where the cloud fraction is between 10 % and 50 %. The aim of the cloud fraction filtering is to allow a comparison of simulations with varying temporal evolutions due to different energy inputs (solar zenith angles) and heat sinks (C_{skin}).




Figure 4. Shown is the ~~Time~~ evolution of the correlation ratio R_c (e.g. as in fig. 2). The solar zenith angle is $\theta = 40^\circ$, there is no mean background wind speed ($\mathbf{u} = 0 \text{ m s}^{-1}$) and the surface skin heat capacity is set to an equivalent of 1 cm water depth (representative for continental land surface). The radiative transfer is computed with the TenStream solver and the sun is positioned in the east ($\varphi = 90^\circ$). The first shallow cumulus clouds develop with a random orientation ($R_c = 1$). The radiative response (i.e. surface shadows) changes the organization of convection to bands from north to south $R_c < 1$ in about one hour. Additionally, to examine the timescales of radiatively induced organization of convection, we perform a restart of the simulation with the sun positioned in the south ($\varphi = 180^\circ$). Once the sun is rotated, it takes the simulation again about one hour to change the orientation of convection into bands from east to west ($R_c > 1$).

The basis for the following analysis is the evaluation of mean correlation ratios as a function of the five free parameters, \mathbf{u} , φ , θ , C_{skin} , and the radiative transfer solver (for details, see table 1). Figure 5 shows the mean correlation ratio $\overline{R_c}$ for each of the 192 simulations. The three panels show
 195 results for different horizontal background wind speeds, 0 m s^{-1} , 5 m s^{-1} and 10 m s^{-1} . Each panel's x-axis is divided into four categories for the surface skin heat capacity and the colorbar describes the solar zenith angle. Additionally, four different markers denote the various options concerning the radiative transfer solvers while the rotation of triangle markers (3D RT) denote the azimuth angle.

We will first focus on the left panel which shows the correlation ratios for the simulations without
 200 any background wind and later move on to simulations with wind. In other words, we start by focusing on purely radiative effects and their influence on the organization of convection and eventually add dynamically induced cloud streets to the discussion.

3.1 Without Wind: $\mathbf{u} = 0 \text{ m s}^{-1}$

The three simulations presented in section 2 are located on the far left panel of fig. 5 with a surface
 205 skin heat capacity equivalent of 1 cm water column (furthest to the left shaded area). Correspondingly, the markers for 3D radiative transfer are shown as triangle markers in light blue (zenith angle of 60°). The upward triangle represents the sun positioned in the south and yields a mean correlation ratio of 1.5 (rolls produced west to east). In contrast, the left rotated triangle presents a sun positioned in the east and shows a mean correlation ratio of .7 (rolls produced south to north). The simulation
 210 with 1D radiative transfer is presented with a diamond shaped marker and shows a mean correlation ratio of ≈ 1 (no organization).




Figure 5. Correlation ratio for simulations with a variable surface skin heat capacity (C_{skin}), solar zenith angle (θ), and three wind velocities (panels left to right). Shaded areas group simulations with a constant C_{skin} according to their respective values, while the horizontal spread inside a group is merely to separate data-points visually. Wind-component u is always from west to east while the individual markers denote simulations where the surface irradiance Q_{net} is set to a constant value, or is computed either with a 1D two-stream solver, or with the 3D TenStream where the sun is either shining from the south (180°) or from the east (90°). The correlation ratio is averaged over all time-steps where the cloud fraction is between 10 % and 50 %.

To explain the concept of why 3D RT creates rolls, we may setup a quick thought experiment. First start with the assumption that there already is a single cloud which will cast the shadow along the sun's incident angle. The surface fluxes for latent sensible heat will be smaller in the shadowy area and hence we expect the next convective plume to rise in sun-lit areas. Figure 6 illustrates the concept for a single cloud and the resulting pattern for surface fluxes. The schematic only constrains convection to be less favorable on the shadowy side but it does not necessarily favor the perpendicular directions over the direction towards the sun. However, if a cloud would evolve on the sun-facing side, the resulting shadow would in turn lead to a faster dissipation of the initial cloud and is thereby an unstable environment for persistent cloud patterns. Following this, we expect the convection to occur favorably perpendicular to the sun's incident angle purely from geometric reasoning.

It is also clear from the horizontal axis of fig. 5 that higher heat capacities lead to less pronounced formations of cloud streets which is to be expected because it weakens the radiative impact and consequently reduces the dynamically induced surface heterogeneities. Yet, though weaker, we still




Figure 6. Sketch from an aerial view depicting surface fluxes in the vicinity of a cloud with a tilted solar incidence. The cloud casts a shadow on the westward surface pixels (blue dots). The available convective energy is directly proportional to latent and sensible heat release of the surface in the vicinity of the convective updraft. Arrows illustrate the confluence of near surface air masses from adjacent pixels in a thermally driven updraft event. Convective tendencies will be weaker on pixels that are adjacent to shaded patches, e.g. at *a*). In contrast, pixels that are surrounded by sun-lit patches, e.g. *b*), are likely to show enhanced convective motion. This pattern favors the organization of cumulus convection in stripes perpendicular to the sun’s incident.

225 find an impact in 3D radiative transfer simulations even for a water column equivalent of 10 m. In this case with such high surface heat capacities, the simulations do not exhibit any variability in surface fluxes and radiation solely acts through atmospheric heating. We recover this behavior also in simulations with a fixed sea surface temperature or with constant latent and sensible surface fluxes (not shown). In Jakub (2016, fig. 3.22), we show that the asymmetric heating of the cloud
 230 sides (or similarly in Wapler (2007); Gronemeier et al. (2016) for displaced surface shadows) introduces a secondary circulation by lifting the sun-lit side and enhancing subsidence on the shadowy side. This asymmetry introduces a wind shear component consisting of a horizontal wind away from the sun at cloud height and towards the sun near the surface. Given that the effects of atmospheric heating is much smaller and happens on longer timescales compared to the surface feedback we put
 235 the interpretation aside for another time.

Simulations with one-dimensional radiative transfer or constant Q_{net} do not produce cloud streets which is reflected by correlation ratios $R_c \approx 1$. If we apply the same geometric reasoning from fig. 6 for these simulations, where the shadow is either directly beneath the cloud or with no heterogeneity at all, it is clear that there can be no preferential direction for convective organization.

240 Three-dimensional radiation calculations with high or low solar zenith angles also show a reduced production of cloud streets. This is, (a) because low zenith angles (sun above head) practically behave just as 1D radiative transfer, and (b), because large zenith angles (low sun, smaller heating rates) have a weaker potential to create surface heterogeneities.

3.2 Medium Wind: $u = 5 \text{ m s}^{-1}$

245 The middle panel of fig. 5 presents the correlation ratios for simulations with a horizontal background
wind of 5 m s^{-1} . If we first shift our attention to the simulations with constant surface irradiance
 Q_{net} (round markers), it is evident that the introduction of a mean wind profile leads to the formation
of cloud streets ($\overline{R}_c > 1$), irrespective of radiatively induced surface heterogeneities. The fact that we
find cloud streets also without any radiation is not surprising and is expected from the literature on the
250 formation of buoyancy driven cloud streets (introduced in section 1. Furthermore, we find a spread
in the development of cloud streets depending on the magnitude of the prescribed Q_{net} , with correla-
tion ratios ranging from 1 to 5. The fact that buoyancy driven cloud street organization is favored in
slightly unstable conditions (low sun) compared to stronger instabilities (high sun) agrees well with
observations (e.g. [Woodcock \(1942\); ?](#); [Grossman \(1982\)](#); [Weckwerth et al. \(1997\)](#) [Woodcock \(1942\)](#); [Priestley \(1957\)](#); [Grossman \(](#)

255 So far we discussed only the simulations with constant Q_{net} . When we look at land surfaces that
are coupled to radiative transfer calculations (1D and 3D markers in fig. 5), we find that radiative
heating may either enhance the organization (\overline{R}_c up to 13) or counter-act it ($\overline{R}_c < 1$). The following
paragraph examines the superposition of dynamically and radiatively induced tendencies to organize
the clouds.

260 Let's consider the case that there is a dynamically induced cloud street along the mean background
wind, i.e. from west to east. Quasi 1D radiation (1D and 3D if sun is close to zenith) casts a shadow
onto the cloud's updraft region and therefore hinders further development of the cloud. This results
in values for the correlation ratio of $\overline{R}_c \approx 1$. Similarly, 3D radiation where the azimuth is in the same
direction as the wind (here east, $\varphi = 90^\circ$, left-rotated markers) also inhibits the formation of cloud
265 streets or may even oppose the dynamically induced organization and produce correlation ratios
 $\overline{R}_c < 1$.

In contrast, for 3D radiative transfer with solar incidence perpendicular to the mean wind, i.e.
sun from south or north, and permitted that the sun's zenith angle allows to illuminate the surface
beneath the cloud ($\theta > 20^\circ$), we find that the radiative tendency to organize the clouds amplifies the
270 dynamically one. This synergistic behaviour results in correlation ratios \overline{R}_c between 5 and 13.

As mentioned previously in section 3.1, we again find a generally diminished influence of surface
radiative heating in simulations with larger surface heat capacities.

3.3 Strong Wind: $u = 10 \text{ m s}^{-1}$

A stronger background wind profile of 10 m s^{-1} principally shows similar behavior as the case that
275 was presented with medium wind speeds (see right panel of fig. 5). The mean correlation ratios
of purely dynamically induced cloud streets (simulations with constant Q_{net} , i.e. circle markers)
cover an increasingly large range of ratios from 2 to 14. Strong solar radiation coupled with small
surface heat capacities still manage to efficiently suppress the formation of cloud streets (i.e. \overline{R}_c

consistently smaller than purely dynamic values). Whereas illumination perpendicular to the wind
280 direction ($\varphi = 180$ and $\theta > 20^\circ$) again greatly amplifies the occurrence of cloud streets. This might
be surprising if we consider that horizontal wind should indeed smooth out the impact of radiative
surface heating. Lohou and Patton (2014) for example also suggest that wind speeds of 10 ms^{-1} may
decouple the effects of dynamically induced surface heterogeneities from the evolution of clouds.
However, if we consider that the dynamically induced cloud streets have typical length scales of
285 50 km (Kuettner, 1959), then, as far as radiative heating at the surface is concerned, the cloud appears
to be standing still. In other words, when a dynamically induced cloud feature aligns in such a
way that it persistently shades a surface region for an extended period of time, we expect that the
radiatively induced surface heterogeneities in turn interact with the flow. It is this intricate linkage
between dynamically induced cloud structures and (3D) radiative transfer that may enable or prohibit
290 the formation of cloud streets.

4 Summary & Conclusions

The formation of cumulus cloud streets was historically attributed primarily to dynamics. This work
aims to document and quantify the generation of radiatively induced cloud street structures. To that
end we performed 192 LES simulations with varying parameters (see table 1) for the horizontal wind
295 speed, the surface heat capacity, the solar zenith and azimuth angle, as well as for different radiative
transfer solvers (section 2.2). As a quantitative measure for the development of cloud streets, we
introduce a simple algorithm using the autocorrelation function on the cloud mask (section 2.3),
which provides a scalar quantity for the degree of organization in cloud streets and the alignment
along the cardinal directions.

300 We find that, in the absence of a horizontal wind, 3D radiative transfer produces cloud streets per-
pendicular to the sun's incident direction whereas the 1D approximation or constant surface irradi-
ance produce circular, randomly positioned, clouds. Our reasoning for this is the geometric position
of the cloud's shadow and the corresponding feedback on surface fluxes which enhances or dimin-
ishes convective tendencies (see fig. 6 for details). While the simulations indicate that there exists
305 an influence due to atmospheric heating rates, we find that the differences between 1D and 3D radi-
ation stem predominantly from surface heating, i.e. the horizontal displacement of cloud shadows.
Furthermore, with increasing horizontal wind speeds of 5 or 10 ms^{-1} , we observe the development
of dynamically induced cloud streets. The dynamical formation of cloud streets is not particularly
surprising, but leads to the question if and how radiative transfer interacts with the organization of
310 convection.

We find that if solar radiation illuminates the surface beneath the cloud, i.e. when the sun is
positioned orthogonal to the mean wind field and the solar zenith angle is larger than 20° , the cloud-
radiative feedback may significantly enhance the tendency to organize in cloud streets. In contrast, in

the case of the 1D approximation (or also 3D if the sun is aligned with the mean wind), the tendency
315 to organize in cloud streets is weaker or even prohibited because the shadow is cast directly beneath
the cloud, weakening the cloud's updraft. The timescale of the convective organization through
radiative transfer is found to happen ~~typically~~ on the order of one hour (see fig. 4). The radiative
feedback, creating surface heterogeneities is generally diminished for large surface heat capacities.
We therefore expect radiative feedbacks to be strongest over land surfaces and less so over the ocean.
320 Given the results of this study we expect that simulations including shallow cumulus convection will
have difficulties to produce cloud streets if they employ 1D radiative transfer solvers or, may need
unrealistically high wind speeds to excite cloud street organization.

An interesting future topic would be the influence atmospheric heating rates on the evolution of
cloud shapes, particularly the corresponding timescales and how the introduced asymmetry and shear
325 changes the local flow. Moving forward, we will examine if the relationship between radiative trans-
fer and convective cloud streets also applies to the real world with all the complexities of a diurnal
cycle or static surface heterogeneities and combined with complex wind fields. Several studies
perform detailed analyses on the footprint of static surface heterogeneities in windy conditions,
i.e. how upstream heterogeneities influence the characteristics of boundary layer dynamics (e.g.
330 (Raasch and Harbusch, 2001; Prabha et al., 2007; Courault et al., 2007; Maronga and Raasch, 2013; Chen et al., 2015; Gronemeier
It may very well be worth to revisit their analyses and particularly focus on static and dynamic(radiative)
heterogeneities. A promising start is an analysis of the simulations within the HDCP² project (Heinze
et al., 2017) which will allow us to test the here proposed interpretations in a more realistic setup.

5 Code availability

335 The UCLA-LES model is publicly available at <https://github.com/uclales>. The calculations were
done with the modified radiation interface which is available at [git-revision "56587a6"](#).

To obtain a copy of the TenStream code, please contact one of the authors. This study used
the TenStream model at [git-revision "5e0a2d5"](#). For the sake of reproducibility we provide the input
parameters for the here mentioned UCLA-LES computations along with the TenStream sources.

340 *Acknowledgements.* This work was funded by the Federal Ministry of Education and Research (BMBF) through
the High Definition Clouds and Precipitation for Climate Prediction (HD(CP)2) project (FKZ: 01LK1208A,
01LK1507D). Many thanks to Cathy Hohenegger, Bjorn Stevens and the DKRZ, Hamburg for fruitful discus-
sions and for providing us with the computational resources to conduct our studies. Special thanks are also
due to Alois Dirnhaichner and the anonymous reviewers who improved this manuscript by proofreading and
345 commenting on the original manuscript.

References

- Asai, T.: Stability of a plane parallel flow with variable vertical shear and unstable stratification, *Journal of the Meteorological Society of Japan*. Ser. II, 48, 129–139, doi:10.2151/jmsj1965.48.2_129, 1970.
- 350 Avissar, R. and Schmidt, T.: An evaluation of the scale at which ground-surface heat flux patchiness affects the convective boundary layer using large-eddy simulations, *Journal of the Atmospheric Sciences*, 55, 2666–2689, doi:10.1175/1520-0469(1998)055<2666:AEOTSA>2.0.CO;2, 1998.
- Balay, S., Abhyankar, S., Adams, M. F., Brown, J., Brune, P., Buschelman, K., Eijkhout, V., Gropp, W. D., Kaushik, D., Knepley, M. G., McInnes, L. C., Rupp, K., Smith, B. F., and Zhang, H.: PETSc Users Manual, Tech. Rep. ANL-95/11 - Revision 3.5, Argonne National Laboratory, 2014.
- 355 Brown, R. A.: Longitudinal instabilities and secondary flows in the planetary boundary layer: A review, *Reviews of Geophysics*, 18, 683–697, doi:10.1029/RG018i003p00683, 1980.
- Chen, G., Zhu, X., Sha, W., Iwasaki, T., Seko, H., Saito, K., Iwai, H., and Ishii, S.: Toward improved forecasts of sea-breeze horizontal convective rolls at super high resolutions. Part II: The impacts of land use and buildings, *Monthly Weather Review*, 143, 1873–1894, 2015.
- 360 Childs, H., Brugger, E., Whitlock, B., Meredith, J., Ahern, S., Pugmire, D., Biagas, K., Miller, M., Harrison, C., Weber, G. H., Krishnan, H., Fogal, T., Sanderson, A., Garth, C., Bethel, E. W., Camp, D., Rübél, O., Durant, M., Favre, J. M., and Navrátil, P.: VisIt: An End-User Tool For Visualizing and Analyzing Very Large Data, in: *High Performance Visualization—Enabling Extreme-Scale Scientific Insight*, pp. 357–372, 2012.
- Courault, D., Drobinski, P., Brunet, Y., Lacarrere, P., and Talbot, C.: Impact of surface heterogeneity on a buoyancy-driven convective boundary layer in light winds, *Boundary-Layer Meteorology*, 124, 383–403, doi:10.1007/s10546-007-9172-y, <https://doi.org/10.1007/s10546-007-9172-y>, 2007.
- 365 Emde, C., Buras-Schnell, R., Kylling, A., Mayer, B., Gasteiger, J., Hamann, U., Kylling, J., Richter, B., Pause, C., Dowling, T., et al.: The libRadtran software package for radiative transfer calculations (Version 2.0), *Geoscientific Model Development Discussions*, 8, 10 237–10 303, doi:10.5194/gmd-9-1647-2016, 2015.
- 370 Etling, D. and Brown, R.: Roll vortices in the planetary boundary layer: A review, *Boundary-Layer Meteorology*, 65, 215–248, doi:10.1007/BF00705527, 1993.
- Frame, J. W., Petters, J. L., Markowski, P. M., and Harrington, J. Y.: An application of the tilted independent pixel approximation to cumulonimbus environments, *Atmospheric Research*, 91, 127–136, doi:10.1016/j.atmosres.2008.05.005, 2009.
- 375 Fu, Q. and Liou, K.: On the correlated k-distribution method for radiative transfer in nonhomogeneous atmospheres, *Journal of the Atmospheric Sciences*, 49, 2139–2156, doi:10.1175/1520-0469(1992)049<2139:OTCDMF>2.0.CO;2, 1992.
- Gronemeier, T., Kanani-Sühring, F., and Raasch, S.: Do Shallow Cumulus Clouds have the Potential to Trigger Secondary Circulations Via Shading?, *Boundary-Layer Meteorology*, pp. 1–27, doi:10.1007/s10546-016-0180-7, 2016.
- 380 Grossman, R. L.: An analysis of vertical velocity spectra obtained in the BOMEX fair-weather, trade-wind boundary layer, *Boundary-Layer Meteorology*, 23, 323–357, doi:10.1007/BF00121120, <https://doi.org/10.1007/BF00121120>, 1982.
- Heinze, R., Dipankar, A., Henken, C. C., Moseley, C., Sourdeval, O., Trömel, S., Xie, X., Adamidis, P., Ament, F., Baars, H., Barthlott, C., Behrendt, A., Blahak, U., Bley, S., Brdar, S., Brueck, M., Crewell, S., Deneke, H.,
- 385

- Di Girolamo, P., Evaristo, R., Fischer, J., Frank, C., Friederichs, P., Göcke, T., Gorges, K., Hande, L., Hanke, M., Hansen, A., Hege, H.-C., Hoose, C., Jahns, T., Kalthoff, N., Klocke, D., Kneifel, S., Knippertz, P., Kuhn, A., van Laar, T., Macke, A., Maurer, V., Mayer, B., Meyer, C. I., Muppa, S. K., Neggers, R. A. J., Orlandi, E., Pantillon, F., Pospichal, B., Röber, N., Scheck, L., Seifert, A., Seifert, P., Senf, F., Siligam, P., Simmer, C., Steinke, S., Stevens, B., Wapler, K., Weniger, M., Wulfmeyer, V., Zängl, G., Zhang, D., and Quaas, J.: Large-eddy simulations over Germany using ICON: a comprehensive evaluation, *Quarterly Journal of the Royal Meteorological Society*, 143, 69–100, doi:10.1002/qj.2947, <http://dx.doi.org/10.1002/qj.2947>, 2017.
- 390 Heus, T., Van Heerwaarden, C., Jonker, H., Pier Siebesma, A., Axelsen, S., Dries, K., Geoffroy, O., Moene, A., Pino, D., De Roode, S., et al.: Formulation of the Dutch Atmospheric Large-Eddy Simulation (DALES) and overview of its applications, *Geoscientific Model Development*, 3, 415–444, doi:10.5194/gmd-3-415-2010, 2010.
- 395 Horn, G., Ouwersloot, H., De Arellano, J. V.-G., and Sikma, M.: Cloud Shading Effects on Characteristic Boundary-Layer Length Scales, *Boundary-Layer Meteorology*, 157, 237–263, doi:10.1007/s10546-015-0054-4, 2015.
- 400 Houze Jr, R. A.: *Cloud dynamics*, vol. 104, Academic press, <https://www.elsevier.com/books/cloud-dynamics/houze-jr/978-0-12-374266-7>, 2014.
- Jakub, F.: On the impact of three dimensional radiative transfer on cloud evolution, <http://nbn-resolving.de/urn:nbn:de:bvb:19-197226>, 2016.
- Jakub, F. and Mayer, B.: A three-dimensional parallel radiative transfer model for atmospheric heating rates for use in cloud resolving models—The TenStream solver, *Journal of Quantitative Spectroscopy and Radiative Transfer*, pp. –, doi:<http://dx.doi.org/10.1016/j.jqsrt.2015.05.003>, <http://www.sciencedirect.com/science/article/pii/S0022407315001727>, 2015.
- 405 Jakub, F. and Mayer, B.: 3-D radiative transfer in large-eddy simulations—experiences coupling the TenStream solver to the UCLA-LES, *Geoscientific Model Development*, 9, 1413–1422, doi:10.5194/gmd-9-1413-2016, 2016.
- 410 Klinger, C., Mayer, B., Jakub, F., Zinner, T., Park, S.-B., and Gentine, P.: Effects of 3-D thermal radiation on the development of a shallow cumulus cloud field, *Atmospheric Chemistry and Physics*, 17, 5477–5500, doi:10.5194/acp-17-5477-2017, <https://www.atmos-chem-phys.net/17/5477/2017/>, 2017.
- Kuettner, J.: The Band Structure of the Atmosphere, *Tellus*, 11, 267–294, doi:10.3402/tellusa.v11i3.9319, 1959.
- 415 Lohou, F. and Patton, E. G.: Surface energy balance and buoyancy response to shallow cumulus shading, *Journal of the Atmospheric Sciences*, 71, 665–682, doi:10.1175/JAS-D-13-0145.1, 2014.
- Maronga, B. and Raasch, S.: Large-Eddy Simulations of Surface Heterogeneity Effects on the Convective Boundary Layer During the LITFASS-2003 Experiment, *Boundary-Layer Meteorology*, 146, 17–44, doi:10.1007/s10546-012-9748-z, <https://doi.org/10.1007/s10546-012-9748-z>, 2013.
- 420 Mayer, B.: Radiative transfer in the cloudy atmosphere, in: *EPJ Web of Conferences*, vol. 1, pp. 75–99, EDP Sciences, doi:10.1140/epjconf/e2009-00912-1, 2009.
- O’Hirok, W. and Gautier, C.: The impact of model resolution on differences between independent column approximation and Monte Carlo estimates of shortwave surface irradiance and atmospheric heating rate., *Journal of the atmospheric sciences*, 62, doi:10.1175/JAS3519.1, 2005.

- 425 Patton, E. G., Sullivan, P. P., and Moeng, C.-H.: The influence of idealized heterogeneity on wet and dry planetary boundary layers coupled to the land surface, *Journal of the Atmospheric Sciences*, 62, 2078–2097, doi:10.1175/JAS3465.1, 2005.
- Pincus, R. and Stevens, B.: Monte Carlo spectral integration: A consistent approximation for radiative transfer in large eddy simulations, *Journal of Advances in Modeling Earth Systems*, 1, doi:10.3894/JAMES.2009.1.1, 430 2009.
- Pincus, R. and Stevens, B.: Paths to accuracy for radiation parameterizations in atmospheric models, *Journal of Advances in Modeling Earth Systems*, 5, 225–233, doi:10.1002/jame.20027, 2013.
- Prabha, T. V., Karipot, A., and Binford, M. W.: Characteristics of secondary circulations over an inhomogeneous surface simulated with large-eddy simulation, *Boundary-Layer Meteorology*, 123, 239–261, 435 doi:10.1007/s10546-006-9137-6, <https://doi.org/10.1007/s10546-006-9137-6>, 2007.
- Priestley, C.: Convection from the earth's surface, in: *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, vol. 238, pp. 287–304, The Royal Society, doi:10.1098/rspa.1957.0001, 1957.
- Raasch, S. and Harbusch, G.: An Analysis Of Secondary Circulations And Their Effects Caused By Small-Scale Surface Inhomogeneities Using Large-Eddy Simulation, *Boundary-Layer Meteorology*, 101, 31–59, 440 doi:10.1023/A:1019297504109, <https://doi.org/10.1023/A:1019297504109>, 2001.
- Rieck, M., Hohenegger, C., and van Heerwaarden, C. C.: The influence of land surface heterogeneities on cloud size development, *Monthly Weather Review*, 142, 3830–3846, doi:10.1175/MWR-D-13-00354.1, 2014.
- Schumann, U., Dörnbrack, A., and Mayer, B.: Cloud-shadow effects on the structure of the convective boundary 445 layer, *Meteorologische Zeitschrift*, 11, 285–294, doi:10.1127/0941-2948/2002/0011-0285, 2002.
- Seifert, A. and Beheng, K. D.: A double-moment parameterization for simulating autoconversion, accretion and selfcollection, *Atmospheric research*, 59, 265–281, doi:10.1016/S0169-8095(01)00126-0, 2001.
- Seifert, A. and Heus, T.: Large-eddy simulation of organized precipitating trade wind cumulus clouds, *Atmospheric Chemistry and Physics*, 13, 5631–5645, doi:10.5194/acp-13-5631-2013, 2013.
- 450 Steinhoff, E.: Excerpt from flight report on the first glider flight in cloud-streets, *Tellus*, 11, 267–294, doi:10.3402/tellusa.v11i3.9319, excerpt is attached in the Appendix of J. Kuettner, 1959, *The Band Structure of the Atmosphere*, 1935.
- Stevens, B., Moeng, C.-H., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J.-C., Jiang, H., Khairoutdinov, M., et al.: Evaluation of large-eddy simulations via observations of nocturnal 455 marine stratocumulus, *Monthly weather review*, 133, 1443–1462, doi:10.1175/MWR2930.1, 2005.
- Vilà-Guerau de Arellano, J., Ouwersloot, H. G., Baldocchi, D., and Jacobs, C. M.: Shallow cumulus rooted in photosynthesis, *Geophysical Research Letters*, 41, 1796–1802, doi:10.1002/2014GL059279, 2014.
- Wapler, K.: Der Einfluss des dreidimensionalen Strahlungstransportes auf Wolkenbildung und -entwicklung, <http://nbn-resolving.de/urn:nbn:de:bvb:19-68948>, 2007.
- 460 Wapler, K. and Mayer, B.: A Fast Three-Dimensional Approximation for the Calculation of Surface Irradiance in Large-Eddy Simulation Models., *Journal of Applied Meteorology & Climatology*, 47, doi:10.1175/2008JAMC1842.1, 2008.

- Weckwerth, T. M., Wilson, J. W., Wakimoto, R. M., and Crook, N. A.: Horizontal convective rolls: Determining the environmental conditions supporting their existence and characteristics, *Monthly weather review*, 125, 505–526, doi:10.1175/1520-0493(1997)125<0505:HCRDTE>2.0.CO;2, 1997.
- 465
- Wissmeier, U., Buras, R., and Mayer, B.: paNTICA: A Fast 3D Radiative Transfer Scheme to Calculate Surface Solar Irradiance for NWP and LES Models., *Journal of Applied Meteorology & Climatology*, 52, doi:10.1175/JAMC-D-12-0227.1, 2013.
- Woodcock, A. H.: Soaring over the Open Sea, *The Scientific Monthly*, 55, 226–232, 1942.