

# Aerosol Surface Area Concentration: a Governing Factor for New Particle Formation in Beijing

Runlong Cai<sup>1,#</sup>, Dongsen Yang<sup>2,#</sup>, Yueyun Fu<sup>1</sup>, Xing Wang<sup>2</sup>, Xiaoxiao Li<sup>1</sup>, Yan Ma<sup>2</sup>, Jiming Hao<sup>1</sup>, Jun Zheng<sup>2\*</sup> and Jingkun Jiang<sup>1\*</sup>

5 <sup>1</sup>State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing, 100084, China

<sup>2</sup>Collaborative Innovation Center of Atmospheric Environment and Equipment Technology, Nanjing University of Information Science & Technology, Nanjing 210044, China

#: Runlong Cai and Dongsen Yang contribute equally to this work

10 \*: *Correspondence to:* J. Jiang (jiangjk@tsinghua.edu.cn) and J. Zheng (zheng.jun@nuist.edu.cn)

**Abstract.** The predominating role of aerosol Fuchs surface area,  $A_{\text{Fuchs}}$ , in determining the occurrence of new particle formation (NPF) events in Beijing was elucidated in this study. The analysis was based on a field campaign from March 12<sup>th</sup> to April 6<sup>th</sup>, 2016, in Beijing, during which aerosol size distributions down to ~1 nm and sulfuric acid concentrations were simultaneously monitored. The 26 days were classified into 11 typical NPF days, 2 undefined days, and 13 non-event days. A dimensionless factor,  $L_{\Gamma}$ , characterized by the relative ratio of the coagulation scavenging rate over the condensational growth rate (Kuang et al., 2010), was applied in this work to reveal the governing factors for NPF events in Beijing. The three parameters determining  $L_{\Gamma}$  are sulfuric acid concentration, the growth enhancement factor characterized by contribution of other gaseous precursors to particle growth,  $\Gamma$ , and  $A_{\text{Fuchs}}$ . Different from other atmospheric environments such as in Boulder and Hyytiälä, daily maximum sulfuric acid concentration and  $\Gamma$  in Beijing varied in a narrow range with geometric standard deviations of 1.40 and 1.31, respectively. A positive correlation between the estimated new particle formation rate,  $J_{1.5}$ , and sulfuric acid concentration was found with a mean fitted exponent of 2.4. However, the maximum sulfuric acid concentrations on NPF days were not significantly higher (even lower, sometime) than those on non-event days, indicating that the abundance of sulfuric acid in Beijing was high enough to initiate nucleation, but may not necessarily lead into NPF events. Instead,  $A_{\text{Fuchs}}$  in Beijing varied greatly among days with a geometric standard deviation of 2.56, while that in Tecamac, Atlanta, and Boulder were reported to be much less variable. In addition, there was a good correlation between  $A_{\text{Fuchs}}$  and  $L_{\Gamma}$  in Beijing ( $R^2 = 0.88$ ). Therefore, it was  $A_{\text{Fuchs}}$  that fundamentally determined the occurrence of NPF events. Among 11 observed NPF events, 10 events occurred when  $A_{\text{Fuchs}}$  was smaller than  $200 \mu\text{m}^2/\text{cm}^3$ . NPF events were suppressed due to the coagulation scavenging when  $A_{\text{Fuchs}}$  was greater than  $200 \mu\text{m}^2/\text{cm}^3$ . Measured  $A_{\text{Fuchs}}$  in Beijing was in good correlation with its  $\text{PM}_{2.5}$  mass concentration ( $R^2 = 0.85$ ) since  $A_{\text{Fuchs}}$  in Beijing was mainly determined by particles in the size range of 50 – 500 nm that also contribute to the  $\text{PM}_{2.5}$  mass concentration.

## 1 Introduction

New particle formation (NPF) is closely related to atmospheric environment. It is a common atmospheric phenomenon, which has been observed all over the world (Kulmala et al., 2004). High concentrations of ultrafine particles are formed intensively during NPF events. It has been illustrated through both theoretical modelling and field observations that these ultrafine particles can grow and serve as cloud condensation nuclei (Kuang et al., 2009; Spracklen et al., 2008), and thus affect climate (IPCC, 2013). The increased number concentration of ultrafine particles also raises concerns on human health (HEI, 2013).

New particles are formed by nucleation from gaseous precursors such as sulfuric acid, ammonia, and organics. Newly formed particles either grow by condensation or are lost by coagulation with other particles (McMurry, 1983). Aerosol Fuchs surface area,  $A_{\text{Fuchs}}$ , is a parameter that describes the coagulation scavenging effect quantitatively. In addition to gaseous precursors participating in nucleation and subsequent condensational growth, it has been a consensus that the occurrence of a NPF event is also limited by  $A_{\text{Fuchs}}$ , because the survival possibility of nucleated particles is suppressed when the coagulation scavenging effect is significant (Weber et al., 1997; Kerminen et al., 2001; Kuang et al., 2012). Reported average  $A_{\text{Fuchs}}$  (or in the form of condensation sink) on NPF days was found to be lower than that on non-event days at several locations (Dal Maso et al., 2005; Gong et al., 2010; Qi et al., 2015).

A dimensionless criterion,  $L_{\Gamma}$ , was proposed to characterize the ratio of particle scavenging loss rate over condensational growth rate, and to predict the occurrence of NPF events in diverse atmospheric environments (Kuang et al., 2010). By definition,  $L_{\Gamma}$  is determined by three factors, i.e., sulfuric acid concentration, the growth enhancement factor representing contribution of other gaseous precursors in addition to sulfuric acid,  $\Gamma$ , and  $A_{\text{Fuchs}}$ . Diurnal sulfuric acid can vary drastically diurnal variation due to the substantial change in radiation (e.g., from several thousand to  $\sim 1.5 \times 10^6$  #/cm<sup>3</sup> in this campaign) and the increase in sulfuric acid concentration after the sunrise can potentially lead to nucleation. The values of  $A_{\text{Fuchs}}$ , however, were usually reported within a narrow range at locations such as Tecamac, Atlanta and Boulder (Kuang et al., 2010). Sulfuric acid concentration in Atlanta and Hyytiälä can differ significantly among days (Eisele et al., 2006; Petäjä et al., 2009). Therefore, sulfuric acid often governs nucleation and subsequent growth in the sulfur-rich atmosphere such as in Atlanta (McMurry et al., 2005). The growth enhancement factor,  $\Gamma$ , at Hyytiälä varied in a wide range while those at Tecamac and Boulder were found in a relatively narrow range.

Aerosol concentrations in Beijing are usually much higher than those in clean environments. The annual average PM<sub>2.5</sub> mass concentration in 2016 was 73  $\mu\text{g}/\text{m}^3$  (reported by Beijing Municipal Environmental Protection Bureau), and the average  $A_{\text{Fuchs}}$  measured in Beijing by this campaign was 381.5  $\mu\text{m}^2/\text{cm}^3$ , which is approximately a magnitude higher than those measured in clean environments such as in Hyytiälä (Dal Maso et al., 2002). Different from comparatively slow

accumulation and depletion process of aerosol concentrations in clean environments,  $A_{\text{Fuchs}}$  in Beijing may change rapidly because of changes in air mass origins (Wehner et al., 2008) or accumulation of pollutants.

Sulfuric acid concentration is needed to estimate  $L_{\Gamma}$  and direct measurement of particle size distribution down to  $\sim 1$  nm will help to better quantify NPF events. Although sulfuric acid has been measured around the world (Erupe et al., 2010) and the analysis based on sub-3 nm size distributions have been conducted sporadically since the development of diethylene glycol scanning mobility particle spectrometer (DEG-SMPS, Jiang et al., 2011a; Jiang et al., 2011b; Kuang et al., 2012) and particle size magnifier (PSM, Vanhanen et al., 2011; Kulmala et al., 2013), there are limited data on atmospheric sulfuric acid concentrations and directly measured sub-3 nm particle size distributions in China. A campaign in Beijing during 2008 Olympic Games (Yue et al., 2010; Zheng et al., 2011) characterized atmospheric sulfuric acid concentration and its correlation with new particle formation rate. The exponent in the correlation of formation rate,  $J_3$ , with sulfuric acid was found to be 2.3. The exponent for correlating derived  $J_{1.5}$  with sulfuric acid was 2.7 (Wang et al., 2011). They were different from the exponents between 1 and 2 often reported in other places around the world (Riipinen et al., 2007; Sihto et al., 2006; Kuang et al., 2008). The same instrument used in the Beijing campaign was also deployed in Kaiping to measure sulfuric acid concentration during a one-month campaign in 2008 (Wang et al., 2013a). Sub-3 nm particle size distributions have not been reported previously in China, except for 1-3 nm particle number concentration in Shanghai in Winter 2013 inferred by a PSM (Xiao et al., 2015). Due to the limitation of observation data, although good correlation between new particle formation rate and sulfuric acid concentration in Beijing was found and the ratio of sulfuric acid concentration over  $A_{\text{Fuchs}}$  was reported to positively correlate with number concentration of 3-6 nm particles (Wang et al., 2011), the roles of sulfuric acid concentration and  $A_{\text{Fuchs}}$  in determining the occurrence of NPF events have not been quantitatively illustrated.

In this study, we aimed to examine the roles of  $A_{\text{Fuchs}}$  and sulfuric acid in determining whether a NPF event will occur on a particular day in Beijing. The data analysis was based on simultaneous measurement of particle size distributions down to  $\sim 1$  nm and sulfuric acid. The correlation between particle formation rate,  $J_{1.5}$ , and sulfuric acid concentration was examined.  $L_{\Gamma}$  was used to predict the occurrence of NPF events. Daily variations of the three parameters determining  $L_{\Gamma}$ , i.e., sulfuric acid concentration,  $\Gamma$ , and  $A_{\text{Fuchs}}$ , were compared. A nominal value of  $A_{\text{Fuchs}}$  was suggested to predict the occurrence of NPF events in Beijing. The relationship between the  $\text{PM}_{2.5}$  mass concentration and NPF events was also examined.

## 2 Experiments

A field campaign studying NPF in Beijing was carried out from Mar. 7<sup>th</sup> to Apr. 7<sup>th</sup>, 2016. The campaign site was located on the campus of Tsinghua University. Details of this site can be found elsewhere (Cai & Jiang, 2017; He et al., 2001).

A home-made DEG SMPS was used to measure sub-5 nm particle size distributions and a particle size distribution system (including a TSI aerodynamic particle sizer and two parallel SMPSs, equipped with a TSI nanoDMA and a TSI long DMA, respectively) was used to measure size distributions of particles from 3 nm to 10  $\mu\text{m}$  (Liu et al., 2016). A specially  
95 designed miniature cylindrical differential mobility analyzer (mini- cyDMA) for effective classification of sub-3 nm aerosol was equipped with the DEG-SMPS (Cai et al., 2017). A cyclone was used at the sampling inlet to remove particles larger than 10  $\mu\text{m}$ . The sampled aerosol was subsequently dried by a silica-gel diffusion drier. The diameter change due to drying was neglected when calculating  $A_{\text{Fuchs}}$  since the mean daytime relative humidity during the campaign period was  $\sim 25\%$ . Diffusion losses, charging efficiency, penetration efficiencies through the DMAs, detection efficiencies of  
100 particle counters, and multi-charging effect were considered during data inversion. Particle density was assumed to be 1.6  $\text{g}/\text{cm}^3$  according to local observation results (Hu et al., 2012).

Sulfuric acid was measured by a modified high-resolution time-of-flight chemical ionization mass spectrometer (HR-ToF-CIMS, Aerodyne Research Inc.). Instead of using radioactive ion source, a home-made corona discharge (CD) ion source was utilized with the HR-TOF-CIMS. The CD ion source was designed to be able to operate from a few Torr up  
105 to near atmospheric pressure and has been successfully implemented in measuring ambient amine (Zheng et al., 2015a) and formaldehyde (Ma et al., 2016). In this study, nitrate reagent ions were used to measure gaseous sulfuric acid (Zheng et al., 2010). The detailed ion chemistry to generate nitrate ions and the calibration procedure for sulfuric acid measurement have been reported in Zheng et al. (2015b). Ambient sulfuric acid concentration in Beijing has been reported only once in a field campaign conducted in 2008 (Zheng et al., 2011; Wang et al., 2011). Compared to that campaign,  
110 sulfuric acid concentration measured in this study displayed similar diurnal variations, but with lower daily maximum values. This might be caused by the relatively weak solar radiation intensity encountered in this springtime observation compared with the previous summertime campaign. To verify the precision of sulfuric acid measurement, the instrument was calibrated daily at night and background checks were performed for  $\sim 3$  minutes each hour during daytime.

A meteorological station (Davis 6250) measuring temperature, relative humidity, wind speed, wind direction, and  
115 precipitation was located at  $\sim 10$  m away from the sampling inlet. The  $\text{PM}_{2.5}$  mass concentration measured in the nearest national monitoring station (Wanliu station,  $\sim 5$  km away on the southwest of our campaign site) was also used for analysis. Backward trajectories were obtained from online HYSPLIT server of national oceanic and atmospheric administration (NOAA).

### 3 Theory

120 Nucleation is only the first step of new particle formation. Gaseous precursors form clusters by random collisions and bound together by Van der Waals force and/or chemical bond. These clusters become particles if they are more likely to

grow by condensation rather than evaporate. However, particles formed by nucleation may be scavenged through coagulation with larger particles before they grow large enough to be detected. Nucleation only refers to the process that stable molecular clusters formed spontaneously from gaseous precursors. New particle formation also requires subsequent condensational growth of freshly nucleated particles. That is, the occurrence of nucleation is mainly determined by gaseous precursors (e.g., sulfuric acid and organics) in atmospheric environment while new particle formation is also influenced by the coagulation scavenging effect of pre-existing aerosols. A possibility exists that nucleation occurs while NPF events are not observed because of the short lifetime of nucleated particles due to a strong coagulation scavenging. In fact, nucleation can also be suppressed when the aerosol concentration is high since vapours and clusters may also be scavenged by aerosol surface.

Aerosol Fuchs surface area,  $A_{\text{Fuchs}}$ , is a representative parameter of coagulation scavenging based on kinetic theory. It is corrected for particles whose size falls in the transition regime (Davis et al., 1980; McMurry, 1983). The formula assuming unity mass accommodation coefficient (sticking probability) is shown in Eq. (1),

$$A_{\text{Fuchs}} = \frac{4\pi}{3} \int_{d_{\min}}^{\infty} d_p^2 \cdot \left( \frac{Kn + Kn^2}{1 + 1.71Kn + 1.33Kn^2} \right) \cdot n \cdot dd_p, \quad (1)$$

where  $d_p$  is particle diameter,  $d_{\min}$  is the smallest particle diameter in theory and the smallest detected one in practice,  $Kn$  is Knudsen number and  $n$  is particle size distribution function,  $dN/dd_p$ . The condensation sink and coagulation sink can also describe how rapidly gaseous precursors and particles are scavenged by pre-existing aerosols, respectively (Kerminen et al., 2001; Kulmala et al., 2001). Since the condensation sink is proportional to  $A_{\text{Fuchs}}$  (McMurry et al., 2005) and the coagulation sink can be approximately converted to the condensation sink using a simple formula (Lehtinen et al., 2007), only  $A_{\text{Fuchs}}$  is used in this study to describe the coagulation scavenging effect. Condensation sink values reported in previous studies are referred in the form of  $A_{\text{Fuchs}}$ . The diffusion coefficient of sulfuric acid was assumed to be  $0.117 \text{ cm}^2 \text{ s}^{-1}$  (Gong et al., 2010) when converting the condensation sink into  $A_{\text{Fuchs}}$ .

A dimensionless criterion,  $L_{\Gamma}$ , was proposed to predict the occurrence of NPF events (Kuang et al., 2010). It is defined as,

$$L_{\Gamma} = \frac{\bar{c} \cdot A_{\text{Fuchs}}}{4\beta_{11}N_1} \cdot \frac{1}{\Gamma}, \quad (2)$$

where  $\bar{c}$  is the mean thermal speed of sulfuric acid that can be calculated from molecular kinetic theory;  $\beta_{11}$  is the coagulation coefficient between sulfuric acid monomers that can be calculated using Eq. 13.56 in Seinfeld & Pandis (2006);  $N_1$  is the number concentration of sulfuric acid;  $\Gamma$  is a growth enhancement factor and is defined as,

$$\Gamma = \frac{2GR}{v_1 N_m c}, \quad (3)$$

150 where  $GR$  is the observed mean growth rate;  $v_1$  is the corresponding volume of sulfuric acid monomer and was estimated to be  $1.7 \times 10^{-28} \text{ m}^3$  (the volume of a hydrated sulfuric acid molecule, Kuang et al., 2010);  $N_m$  is the maximum number concentration of sulfuric acid during a whole NPF event period. Since other gaseous precursors in addition to sulfuric acid might also contribute to the condensational growth of particles formed by nucleation (O'Dowd et al., 2002; Ristovski et al., 2010) and only concentration of sulfuric acid is used in Eq. (2), the ratio of measured growth rate over the sulfuric acid condensational growth rate (Weber et al., 1997), i.e.,  $\Gamma$ , was used for correction. It should be clarified that  $L_\Gamma$  in Eq. (2) is defined similar to that in McMurry et al (2005) but slightly different from that in Kuang et al (2010), since  $L_\Gamma$  in this study are time-resolved values rather than event specific ones. Theoretically,  $\Gamma$  can also be time and size-resolved if using time and size-resolved GR and time-resolved sulfuric acid (Kuang et al., 2012). However,  $\Gamma$  during each NPF event is assumed to be constant in Eq. (3) because further evaluations are needed for this time and size-resolved model. Note that in Eq.(2) the absolute sulfuric acid concentrations were effectively normalized by the corresponding daily sulfuric acid maximum concentrations and thus has no influence on  $L_\Gamma$  values and conclusions based on  $L_\Gamma$  reported in this study.

A new balance formula to estimate new particle formation rate was proposed recently (Cai & Jiang, 2017) and is given below,

$$J_k = \frac{dN_{[d_k, d_u]}}{dt} + \sum_{d_g=d_k}^{d_u-1} \sum_{d_i=d_{\min}}^{+\infty} \beta_{(i,g)} N_{[d_i, d_{i+1}]} N_{[d_g, d_{g+1}]} - \frac{1}{2} \sum_{d_g=d_k}^{d_u-1} \sum_{\substack{d_i^3 + d_{j+1}^3 = d_g^3 \\ d_{i+1}^3 + d_j^3 = d_g^3 \\ d_i, d_j \geq d_{\min}}} \beta_{(i,j)} N_{[d_i, d_{i+1}]} N_{[d_j, d_{j+1}]} + n_u \cdot GR_u, \quad (4)$$

165 where  $J_k$  is the formation rate of particles at the size of  $d_k$ ,  $N_{[d_k, d_u]}$  is the total number concentration of particles from  $d_k$  to  $d_u$  (not included),  $d_u$  is the upper bound of the size range for calculation (25 nm in this study),  $d_{\min}$  is the size of the smallest cluster in theory and the smallest detected size in practice (1.3 nm in this study). The second and third terms in the right hand side of Eq. (4) are the coagulation sink term (*CoagSnk*) and the coagulation source term (*CoagSrc*), respectively. The difference between *CoagSnk* and *CoagSrc* is the net *CoagSnk* representing the net rate of particles from  $d_k$  to  $d_u$ , i.e., lost by coagulation scavenging. The last term is often negligible according to the determination criterions for  $d_u$ .  $dN/dt$  is the balance result of  $J_k$  and net *CoagSnk*.

## 4 Results and Discussion

A total of 26 days from Mar. 12<sup>th</sup> to Apr. 6<sup>th</sup> was classified by the occurrence of daytime NPF event. A typical NPF day is featured with distinct and persisting increases in sub-3 nm particle number concentration and subsequent growth of these nucleated particles. A non-event day means that neither of these two features was observed. As shown in Fig. 1, there are 11 typical NPF days and 13 non-event days. The rest 2 days, i.e., Mar. 19<sup>th</sup> and Mar. 30<sup>th</sup>, were classified as undefined days. On these days, the increase in sub-3 nm particle number concentration and subsequent growth were both

observed. However, sub-3 nm particle number concentration was relatively low and the evolution of particle size distributions was not continuous. NPF events mainly occurred when wind came from northwest of Beijing and non-event days were associated with air masses from southwest (as summarized in Table 1). Air masses coming from north usually experience less influence from urban pollution (Wehner et al., 2008; Wang et al., 2013b), i.e.,  $A_{\text{Fuchs}}$  is likely to be lower than that on days dominated by southwest wind.

The occurrence of NPF events in most days can be predicted by  $L_{\Gamma}$  if unity was empirically chosen as the threshold value. Greater  $L_{\Gamma}$  indicates higher possibilities of nucleated particles to be scavenged by coagulation before they can continue to grow. Growth rates on non-event days were assumed to 2.4 nm/h, the mean value of observed growth rates on NPF days (the range is 1.2 nm/h to 3.3 nm/h). A threshold value of  $L_{\Gamma}$  can not be theoretically predicted but can be empirically estimated. 0.7 was suggested as the threshold value by Kuang et al. (2010). However, unity suggested by McMurry et al. (2005) appeared to work better for results from this campaign in Beijing. As shown in Table 1, the median and mean values of  $L_{\Gamma}$  on NPF days observed in this campaign were 0.55 and 0.71 (with a standard deviation of 0.40), respectively, comparing to 3.05 and 3.45 on non-event days (with a standard deviation of 1.79), respectively. However, some exceptions were also observed. On the two undefined days,  $L_{\Gamma}$  were 1.40 and 0.64, respectively, and weak nucleation was observed. Although the estimated  $L_{\Gamma}$  value on Mar. 18<sup>th</sup> was 1.75, a comparatively weak but still distinct NPF event was observed. Despite these few exceptions,  $L_{\Gamma}$  works well in most days in this campaign and were verified in other places (Kuang et al., 2010). Following discussion is focused on the contribution of different factors, i.e., sulfuric acid concentration,  $\Gamma$ , and  $A_{\text{Fuchs}}$ .

#### 4.1 The Role of Gaseous Precursors

There was a positive correlation between the estimated new particle formation rate,  $J_{1.5}$ , and sulfuric acid concentration during most NPF periods (typically 8:00-16:00 when the estimated  $J_{1.5}$  was greater than zero). On NPF days, an increase in sub-3 nm particle number concentration was often accompanied with an increase in sulfuric acid concentration (as shown in Fig. 2). Considering the possible sensitivity of the fitted parameters to the fitting time period (Kuang et al., 2008), the correlation between  $J_{1.5}$  and sulfuric acid concentration was only examined for NPF periods. We found that the mean coefficient of determination ( $R^2$ ) in this campaign was 0.53. The exponents for correlating the  $J_{1.5}$  and sulfuric acid concentration ranged from 1.5 to 4.0 in the 10 days with a mean value of 2.4 (Mar. 29<sup>th</sup> was not included because of insignificant correlation). This is in consensus with previously reported mean exponent of 2.3 using  $J_3$  in Beijing (Wang et al., 2011). However, it is quite different from those exponents no greater than 2 observed in North America and Europe (Kuang et al., 2008; Riipinen et al., 2007; Sihto et al., 2006) indicating activation or kinetic nucleation alone can not explain all NPF events observed in this campaign.

Although the correlation between sulfuric acid concentration and the particle formation rate was significant, sulfuric acid appeared not to be the determining factor for whether a NPF event would occur in Beijing. As illustrated by the temporal trend of sulfuric acid concentration in Fig. 2, a significant diurnal variation was observed every day. However, the differences among daily maximum sulfuric acid concentrations were small. The variations of daily maximum sulfuric acid concentration were significantly less than those of  $A_{\text{Fuchs}}$ . The geometrical standard deviation and relative standard deviation of maximum sulfuric acid concentration on each day were 1.40 and 0.34, respectively, while those of daily averaged  $A_{\text{Fuchs}}$  values were 2.56 and 0.82, respectively. Sulfuric acid concentrations during NPF periods were not significantly higher than those between 8:00 - 16:00 on non-event days ( $p=1$ ). In addition, comparatively high concentrations of sulfuric acid, e.g., on Apr. 4<sup>th</sup> - 6<sup>th</sup>, did not necessarily lead to NPF events.

The influence of growth enhancement factor,  $\Gamma$ , on the occurrence of NPF events also needs to be addressed because sulfuric acid alone may not explain the observed growth rates. Estimated  $\Gamma$  value for each event was normalized by the geometric mean  $\Gamma$  value for the whole campaign to make it comparable with those obtained from previous studies (Kuang et al., 2010): MILAGRO in Tecamac (Iida et al., 2008); ANARChE (McMurry et al., 2005) in Atlanta; Boulder (Iida et al., 2006); QUEST II (Sihto et al., 2006), QUEST IV (Riipinen, et al., 2007), and EUCAARI (Manninen et al., 2009) at the SMEAR II station in Hyytiälä. It should be clarified that the relative value of  $\Gamma$  can improve the comparability by overcoming some uncertainties in measuring sulfuric acid concentration in different studies. Fig. 4 indicates that  $\Gamma$  values observed in this study distribute in a relatively narrow range, similar to those observed in Tecamac, Atlanta, and Boulder, while different from the wide-spreading characteristics of  $\Gamma$  values in Hyytiälä. Geometric standard deviations of  $\Gamma$  values were 1.31, 1.75, 2.23, 1.87, 1.62, 2.77, and 2.87 in this campaign, MILAGRO, ANARChE, Boulder, QUEST II, QUEST IV, and EUCAARI, respectively. The daily variations of  $\Gamma$  values in Beijing were less than those observed in other places. They were also less than the daily variations of  $A_{\text{Fuchs}}$  values measured in this campaign. Considering the small daily variations of both sulfuric acid concentration and  $\Gamma$  values, it is reasonable to conclude that the abundance of gaseous precursors such as sulfuric acid in Beijing during the campaign period was sufficiently high for nucleation to occur but the occurrence of NPF events appeared to be governed by  $A_{\text{Fuchs}}$ .

#### 4.2 Relationship between $A_{\text{Fuchs}}$ and NPF Events

Comparatively lower  $A_{\text{Fuchs}}$  values were found during most of the NPF days while sulfuric acid concentrations on NPF days were not significantly higher than those on non-event days. NPF events mainly occurred when  $A_{\text{Fuchs}}$  was smaller than  $200 \mu\text{m}^2/\text{cm}^3$ . Non-event days mainly corresponded to a real-time  $A_{\text{Fuchs}}$  value greater than  $200 \mu\text{m}^2/\text{cm}^3$  and an average  $A_{\text{Fuchs}}$  value greater than  $350 \mu\text{m}^2/\text{cm}^3$  (Fig. 5). The value of  $200 \mu\text{m}^2/\text{cm}^3$  appeared to be an empirical division between NPF days and non-event days. If  $A_{\text{Fuchs}}$  was lower than this value, a NPF event tended to occur. Otherwise, the occurrence of NPF events was suppressed because of the predominant coagulation scavenging effect.



The variation of  $L_{\Gamma}$  in Beijing was governed by  $A_{\text{Fuchs}}$ . The measured  $L_{\Gamma}$  and  $A_{\text{Fuchs}}$  values were in good correlation with a coefficient of determination ( $R^2$ ) of 0.88. The mean relative error of fitted  $L_{\Gamma}$  using  $A_{\text{Fuchs}}$  was 11.4% compared to the measured ones (Fig. 6(a)). It should be clarified that  $GR$  on non-event days in this campaign was assumed to be the same (2.4 nm/h, an average of the fitted values on NPF days). The correlation between  $L_{\Gamma}$  and  $A_{\text{Fuchs}}$  on NPF days alone had the  $R^2$  of 0.89. The  $A_{\text{Fuchs}}$  of  $200 \mu\text{m}^2/\text{cm}^3$  corresponds to an  $L_{\Gamma}$  of approximate unity in this campaign. Since  $L_{\Gamma}$  has been verified as a proper nucleation criterion in diverse atmospheric environments, it is reasonable to conclude that  $A_{\text{Fuchs}}$  was the governing factor of the occurrence of NPF events observed in this campaign.

The characteristics of  $A_{\text{Fuchs}}$  dominated NPF events in Beijing are different from those at other locations. As shown in Fig. 6(b),  $L_{\Gamma}$  and  $A_{\text{Fuchs}}$  in most other places do not correlate well, indicating that  $A_{\text{Fuchs}}$  alone can not predict the occurrence of NPF events at these locations. The variations of these parameters at various locations are illustrated in Fig. 7. In Atlanta and Boulder,  $A_{\text{Fuchs}}$  values fluctuated within relatively narrow ranges while the concentrations of gaseous precursors participating in nucleation differed significantly. The variations of  $L_{\Gamma}$  at these locations were mainly caused by the relatively large variations in the concentrations of gaseous precursors. However, the contribution of gaseous precursors to  $L_{\Gamma}$  in Beijing was relatively stable and the variations of  $L_{\Gamma}$  were mainly caused by the variations in  $A_{\text{Fuchs}}$  values.

The predominant role of  $A_{\text{Fuchs}}$  in Beijing can also be explained using the balance formula shown as Eq. (4). It is  $dN/dt$  rather than the formation rate,  $J$ , that directly reflects whether a NPF event has occurred or not.  $dN/dt$  is the balanced result of the formation rate and the net *CoagSnk*. Different from  $L_{\Gamma}$  that is the ratio of the particle loss rate over the growth rate, the ratio of the net *CoagSnk* over  $J$  represents how many nucleated particles are lost due to the coagulation scavenging. The surviving particles are accounted for by the increment in number concentration of particles in the nucleation mode (1-25 nm). The nucleation mode was used in this study to estimate  $dN/dt$  caused by nucleation because newly formed particles seldom grew beyond 25 nm in the evaluated time period. Surviving possibilities of nucleated particles can also be inferred using the growth rate and  $A_{\text{Fuchs}}$  (Weber et al., 1997; Kerminen & Kulmala, 2002; Kuang et al., 2012). However, the ratio of the net *CoagSnk* over  $J$  was used because it is based on measured particle size distributions. Note that theoretically the ratio of the net *CoagSnk* over  $J$  can be greater than unity. This would correspond to a negative  $dN/dt$  value. For better description of the occurrence of NPF events rather than the whole process including termination, only NPF periods when  $dN/dt$  was positive were considered here. On average, 70% of particles formed by nucleation were lost due to coagulation scavenging on NPF days (as shown in Fig. 8) indicating high coagulation loss in Beijing even on NPF days. When the  $A_{\text{Fuchs}}$  value was much greater, most nucleated particles were lost due to the coagulation scavenging rather than grew into to larger sizes such that NPF events were less likely to be observed.

It should be clarified that although with much less possibility, NPF events may also occur in Beijing when  $A_{\text{Fuchs}}$  was greater than  $200 \mu\text{m}^2/\text{cm}^3$ . In this campaign, a distinct NPF event was observed with a comparatively high  $A_{\text{Fuchs}}$  value of

270  $329 \mu\text{m}^2/\text{cm}^3$  (on Mar. 18<sup>th</sup>). It was significantly higher than the suggested threshold value of  $200 \mu\text{m}^2/\text{cm}^3$ . As indicated by Table 1, this exception was caused by the failure of  $L_{\Gamma}$  rather than  $A_{\text{Fuchs}}$  alone, i.e., NPF events occurred when estimated  $L_{\Gamma}$  was greater than unity (the empirical threshold value). The comparatively low number concentration of sub-3 nm particles together with the moderate particle formation rate indicated that the NPF event was suppressed. In addition, previous studies in Beijing also observed some NPF events when  $A_{\text{Fuchs}}$  values were relatively high (Wu et al., 2007; Wang et al., 2013c; Wang et al., 2017), e.g., an  $A_{\text{Fuchs}}$  value of  $\sim 555 \mu\text{m}^2/\text{cm}^3$  (Kulmala et al., 2016). These reported  $A_{\text{Fuchs}}$  values might be overestimated since the daily average rather than the average only over NPF event periods was used.  $A_{\text{Fuchs}}$  in Beijing during non-event periods can be significantly higher. Nevertheless,  $A_{\text{Fuchs}}$  can be considered as the major determining factor of the occurrence of NPF events in Beijing while admitting that exceptions can occasionally occur at a medium  $L_{\Gamma}$  value greater than unity (corresponding to the  $A_{\text{Fuchs}}$  value of  $200 \mu\text{m}^2/\text{cm}^3$ ).

### 280 4.3 A case Study of 3 Days

Three continuous days including two NPF days and one non-event day are shown in Fig. 9 to further illustrate the roles of  $A_{\text{Fuchs}}$  and sulfuric acid (together with other gaseous precursors) in affecting the occurrence of NPF events in Beijing. On Apr. 2<sup>nd</sup>,  $A_{\text{Fuchs}}$  remained at a relatively low level. A NPF event occurred after sunrise (together with an increase in sulfuric acid concentration) and ended in the afternoon when sulfuric acid concentration decreased to a low level. The whole NPF event began at approximately 7:30 and ended at approximately 14:30 that was also the typical time period for other NPF events observed in this campaign. However, when wind direction changed from northwest to southwest at the noon of on Apr. 3<sup>rd</sup>, sulfuric acid concentration decreased and  $A_{\text{Fuchs}}$  increased rapidly because of particles transported from south. This leads to an increase in  $L_{\Gamma}$ . The ongoing NPF event was interrupted and no newly nucleated particles was detected even when sulfuric acid concentration increased again later. On Apr. 4<sup>th</sup>,  $A_{\text{Fuchs}}$  stayed at a high level.  $L_{\Gamma}$  was always greater than unity. Maximum sulfuric acid concentrations on Apr. 4<sup>th</sup> were even higher than those on Apr. 2<sup>nd</sup> and 3<sup>rd</sup>. However, no NPF event was observed. It supports the argument that the abundance of gaseous precursors in Beijing are often high enough for nucleation to happen, however, whether or not a NPF event occurs is mainly governed by  $A_{\text{Fuchs}}$ .

### 290 4.4 Predicting NPF Days Using $\text{PM}_{2.5}$ Mass Concentration

The  $\text{PM}_{2.5}$  mass concentration in Beijing serves as a rough but simple parameter to predict whether a NPF event can happen. The value of  $A_{\text{Fuchs}}$  is affected by particle size distributions. Accumulation mode particles ranging from 50 nm to 500 nm in Beijing were the major contribution to  $A_{\text{Fuchs}}$ . Normalized size distributions of accumulation mode particles were relative stable at various  $A_{\text{Fuchs}}$  levels (as shown in Fig. 10). On NPF days when  $A_{\text{Fuchs}}$  were relatively low, nucleation mode particles formed by nucleation and subsequent growth also contributed to  $A_{\text{Fuchs}}$ , although  $A_{\text{Fuchs}}$  was still governed by accumulation mode particles. Thus,  $A_{\text{Fuchs}}$  should show better correlation with the particle mass concentration rather

300 than the particle number concentration. Figure 11 indicates that there was a good correlation between  $A_{\text{Fuchs}}$  and the  $\text{PM}_{2.5}$  mass concentration in Beijing with a  $R^2$  of 0.85, although the correlation at high  $A_{\text{Fuchs}}$  level was generally better than that at low  $A_{\text{Fuchs}}$  level because particles formed by nucleation significantly changed the shape of particle size distribution functions on NPF days. Measured  $\text{PM}_{2.5}$  mass concentrations in the 26 days ranged from 3 to  $420 \mu\text{g}/\text{m}^3$ , wide enough to represent both relative clean days and severe polluted days in Beijing. The  $\text{PM}_{2.5}$  mass concentrations during NPF event periods were mostly lower than  $30 \mu\text{g}/\text{m}^3$  except for the event on Mar. 18<sup>th</sup>. On non-event days, the  $\text{PM}_{2.5}$  mass concentrations between 8:00 and 16:00 were typically greater than  $30 \mu\text{g}/\text{m}^3$ . Note that this threshold  $\text{PM}_{2.5}$  value of  $30 \mu\text{g}/\text{m}^3$  may not be valid for the whole year. This campaign was in March and early April. Emissions and radiation intensity are different in different seasons such that the concentrations of gaseous precursors can vary with seasons as well.

The criterion of  $\text{PM}_{2.5}$  mass concentration was applied to predict NPF events measured at the same site in Beijing in April and May, 2014. Among 38 days in that campaign, 11 typical NPF events were identified. For 9 NPF events, average  $\text{PM}_{2.5}$  mass concentrations during event periods were lower than  $30 \mu\text{g}/\text{m}^3$ . For the other 2 events, it was 49.8 and  $40.5 \mu\text{g}/\text{m}^3$ , respectively. In another campaign in Beijing during January 2016 (Jayaratne et al., 2017), 14 NPF events were observed. Among them, 12 events occurred when daily average  $\text{PM}_{2.5}$  mass concentration was lower than  $30 \mu\text{g}/\text{m}^3$ . Daily average  $\text{PM}_{2.5}$  mass concentrations on 16 non-event days were all greater than  $40 \mu\text{g}/\text{m}^3$ .

## 315 5 Conclusions

Factors governing the occurrence of NPF events in Beijing were examined using data from a field campaign during Mar. 12<sup>th</sup> to Apr. 6<sup>th</sup>, 2016. In these 26 days, 11 typical NPF events were observed. The rest were 2 undefined days and 13 non-event days. The new particle formation rate,  $J_{1.5}$ , was in positive correlation with sulfuric acid concentration with a fitted mean exponent of 2.4. However, sulfuric acid concentrations on NPF days were not significantly higher than those on non-event days. A dimensionless criterion proposed by Kuang et al. (2010),  $L_{\Gamma}$ , was found to be applicable to predict NPF events in most days. Theoretically,  $L_{\Gamma}$  was determined by sulfuric acid concentration, the enhancement factor,  $\Gamma$ , and aerosol Fuchs surface area,  $A_{\text{Fuchs}}$ , together. In Beijing, however,  $A_{\text{Fuchs}}$  alone was found to be in good correlation with  $L_{\Gamma}$  ( $R^2 = 0.88$ ). Different from NPF events observed at other locations such as Hyytiälä, daily maximum sulfuric acid concentration and the enhancement factor in Beijing only varied in a narrow range with geometric standard deviations of 1.40 and 1.31, respectively, while  $A_{\text{Fuchs}}$  varied significantly among days with a geometric standard deviation of 2.56. It was inferred that the concentrations of gaseous precursors such as sulfuric acid in Beijing were high enough to initiate nucleation while it was  $A_{\text{Fuchs}}$  that determined whether a NPF event would occur or not. An  $A_{\text{Fuchs}}$  value of  $200 \mu\text{m}^2/\text{cm}^3$  was proposed as the empirical threshold in Beijing below which NPF events are highly likely to occur. NPF events will be suppressed when  $A_{\text{Fuchs}}$  is higher than this threshold value. The  $A_{\text{Fuchs}}$  dominated characteristics in Beijing are different

330 from those at other locations such as Atlanta, Boulder, and Hyytiälä. Since  $A_{\text{Fuchs}}$  in Beijing was mainly governed by  
accumulation mode particles (50 to 500 nm) and the normalized  $dA_{\text{Fuchs}}/d\log d_p$  in this size range was relatively stable at  
different  $A_{\text{Fuchs}}$  levels in Beijing, measured  $A_{\text{Fuchs}}$  was in good correlation with the  $\text{PM}_{2.5}$  mass concentration ( $R^2 = 0.85$ ).  
Accordingly, the  $\text{PM}_{2.5}$  mass concentration may also serve as a rough and simple parameter to predict the occurrence of  
NPF events in Beijing. An empirical  $\text{PM}_{2.5}$  threshold value of  $30 \mu\text{g}/\text{m}^3$  was proposed based on data from this field  
335 campaign and was found to also work well for other field campaigns in Beijing.

### Acknowledgement

Financial supports from the National Science Foundation of China (21422703, 41227805, 21521064, 21377059 &  
41575122) and the National Key R&D Program of China (2014BAC22B00, 2016YFC0200102 & 2016YFC0202402)  
are acknowledged.

### 340 References

- Cai, R., Chen, D.-R., Hao, J., & Jiang, J.: A Miniature Cylindrical Differential Mobility Analyzer for sub-3 nm Particle  
Sizing. *Journal of Aerosol Science*, 106, 111-119, doi:10.1016/j.jaerosci.2017.01.004, 2017.
- Cai, R., & Jiang, J.: A new balance formula to estimate particle formation rate: reevaluating effect of coagulation  
scavenging, *Atmospheric Chemistry and Physics Discussion*, doi:10.5194/acp-2017-199, in review, 2017.
- 345 Dal Maso, M., Kulmala, M., Lehtinen, K.E., Mäkelä, J.M., Aalto, J., & O'Dowd, C.D.: Condensation and coagulation  
sinks and formation of nucleation mode particles in coastal and boreal forest boundary layers. *Journal of  
Geophysical Research*, 107, doi:10.1029/2001jd001053, 2002.
- Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P., & Lehtinen, K.E.: Formation and growth of  
fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II, Hyytiälä, Finland.  
350 *Boreal Environment Research*, 10, 323-336, 2005.
- Davis, E.J., P., R., & K., R.A.: A Review of Theory and Experiments on Diffusion from Submicroscopic Particles.  
*Chemical Engineering Communications*, 5, 251-268, 1980.
- Eisele, F.L., Lovejoy, E.R., Kosciuch, E., Moore, K.F., Mauldin, R.L., Smith, J.N., McMurry, P.H., & Iida, K.: Negative  
atmospheric ions and their potential role in ion-induced nucleation. *Journal of Geophysical Research*, 111,  
355 doi:10.1029/2005jd006568, 2006.
- Erupe, M.E., Benson, D.R., Li, J., Young, L.-H., Verheggen, B., Al-Refai, M., Tahboub, O., Cunningham, V., Frimpong,  
F., Viggiano, A.A., & Lee, S.-H.: Correlation of aerosol nucleation rate with sulfuric acid and ammonia in Kent,  
Ohio: An atmospheric observation. *Journal of Geophysical Research*, 115, doi:10.1029/2010jd013942, 2010.
- Gong, Y., Hu, M., Cheng, Y., Su, H., Yue, D., Liu, F., Wiedensohler, A., Wang, Z., Kalesse, H., & Liu, S.: Competition  
360 of coagulation sink and source rate: New particle formation in the Pearl River Delta of China. *Atmospheric  
Environment*, 44, 3278-3285, doi:10.1016/j.atmosenv.2010.05.049, 2010.
- He, K., Yang, F., Ma, Y., Zhang, Q., Yao, X., Chan, C.K., Cadel, S., Chan, T., & Mulawa, P.: The characteristics of  $\text{PM}_{2.5}$   
in Beijing, China. *Atmospheric Environment*, 35, 4959-4970, 2001.
- HEI Review Panel on Ultrafine Particles: Understanding the Health Effects of Ambient Ultrafine Particles. HEI  
365 Perspectives 3. Health Effects Institute, Boston, MA, 2013.

- Hu, M., Peng, J., Sun, K., Yue, D., Guo, S., Wiedensohler, A., & Wu, Z.: Estimation of size-resolved ambient particle density based on the measurement of aerosol number, mass, and chemical size distributions in the winter in Beijing. *Environmental science & technology*, 46, 9941-9947, doi:10.1021/es204073t, 2012.
- 370 Iida, K., Stolzenburg, M., McMurry, P., Dunn, M.J., Smith, J.N., Eisele, F., & Keady, P.: Contribution of ion-induced nucleation to new particle formation: Methodology and its application to atmospheric observations in Boulder, Colorado. *Journal of Geophysical Research: Atmospheres*, 111, doi:10.1029/2006jd007167, 2006.
- Iida, K., Stolzenburg, M.R., McMurry, P.H., & Smith, J.N.: Estimating nanoparticle growth rates from size-dependent charged fractions: Analysis of new particle formation events in Mexico City. *Journal of Geophysical Research: Atmospheres*, 113, doi:10.1029/2007jd009260, 2008.
- 375 IPCC.: *Climate Change 2013: IPCC Fifth Assessment Report (AR5)*. Cambridge University Press, 2013.
- Jayarathne, R., Pushpawela, B., He, C., Gao, J., Hui, L., Morawska, L.: Observations of Particles at their Formation Sizes in Beijing, China. *Atmospheric Chemistry and Physics Discussions*, doi:10.5194/acp-2017-156, in review, 2017.
- Jiang, J., Chen, M., Kuang, C., Attoui, M., & McMurry, P.H.: Electrical Mobility Spectrometer Using a Diethylene Glycol Condensation Particle Counter for Measurement of Aerosol Size Distributions Down to 1 nm. *Aerosol Science and Technology*, 45, 510-521, doi:10.1080/02786826.2010.547538, 2011.
- 380 Jiang, J., Zhao, J., Chen, M., Eisele, F.L., Scheckman, J., Williams, B.J., Kuang, C., & McMurry, P.H.: First Measurements of Neutral Atmospheric Cluster and 1–2 nm Particle Number Size Distributions During Nucleation Events. *Aerosol Science and Technology*, 45, ii-v, doi:10.1080/02786826.2010.546817, 2011.
- Kerminen, V.M., & Kulmala, M.: Analytical formulae connecting the "real" and the "apparent" nucleation rate and the nuclei number concentration for atmospheric nucleation events. *Journal of Aerosol Science*, 33, 609-622, 2002.
- 385 Kerminen, V.-M., Pirjola, L., & Kulmala, M.: How significantly does coagulative scavenging limit atmospheric particle production? *Journal of Geophysical Research: Atmospheres*, 106, 24119-24125, doi:10.1029/2001jd000322, 2001.
- Kuang, C., Chen, M., Zhao, J., Smith, J., McMurry, P.H., & Wang, J.: Size and time-resolved growth rate measurements of 1 to 5 nm freshly formed atmospheric nuclei. *Atmospheric Chemistry and Physics*, 12, 3573-3589, doi:10.5194/acp-12-3573-2012, 2012.
- 390 Kuang, C., McMurry, P. H., & McCormick, A.V.: Determination of cloud condensation nuclei production from measured new particle formation events, *Geophysical Research Letters*, 36, doi:10.1029/2009gl037584, 2009.
- Kuang, C., McMurry, P.H., McCormick, A.V., & Eisele, F.L.: Dependence of nucleation rates on sulfuric acid vapor concentration in diverse atmospheric locations. *Journal of Geophysical Research*, 113, doi:10.1029/2007jd009253, 2008.
- 395 Kuang, C., Riipinen, I., Sihto, S.L., Kulmala, M., McCormick, A.V., & McMurry, P.H.: An improved criterion for new particle formation in diverse atmospheric environments. *Atmospheric Chemistry and Physics*, 10, 8469-8480, doi:10.5194/acp-10-8469-2010, 2010.
- 400 Kulmala, M., Dal Maso, M., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen, P., Hämeri, K., and O'Dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus*, 53, 479-490, 2001.
- Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H.E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Jarvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P.P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R.L., 3rd, Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M.V., Smith, J.N., Ehn, M., Mentel, T.F., Lehtinen, K.E., Laaksonen, A., Kerminen, V.-M., & Worsnop, D.R.: Direct observations of atmospheric aerosol nucleation. *Science*, 339, 943-946, doi:10.1126/science.1227385, 2013.
- 405 Kulmala, M., Petäjä, T., Kerminen, V.-M., Kujansuu, J., Ruuskanen, T., Ding, A., Nie, W., Hu, M., Wang, Z., Wu, Z., Wang, L., & Worsnop, D.R.: On secondary new particle formation in China. *Frontiers of Environmental Science & Engineering*, 10, doi:10.1007/s11783-016-0850-1, 2016.
- 410

- Kulmala, M., Vehkamäki, H., Petäjä, T., Dal Maso, M., Lauri, A., Kerminen, V.M., Birmili, W., & McMurry, P.H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations. *Journal of Aerosol Science*, 35, 143-176, doi:10.1016/j.jaerosci.2003.10.003, 2004.
- 415 Lehtinen, K.E.J., Dal Maso, M., Kulmala, M., & Kerminen, V.-M.: Estimating nucleation rates from apparent particle formation rates and vice versa: Revised formulation of the Kerminen–Kulmala equation. *Journal of Aerosol Science*, 38, 988-994, doi:10.1016/j.jaerosci.2007.06.009, 2007.
- Liu, J., Jiang, J., Zhang, Q., Deng, J., & Hao, J.: A spectrometer for measuring particle size distributions in the range of 3 nm to 10  $\mu\text{m}$ . *Frontiers of Environmental Science & Engineering*, 10, 63-72, doi:10.1007/s11783-014-0754-x, 2016.
- 420 Ma, Y., Diao, Y., Zhang, B., Wang, W., Ren, X., Yang, D., Wang, M., Shi, X., and Zheng, J.: Detection of formaldehyde emissions from an industrial zone in the Yangtze River Delta region of China using a proton transfer reaction ion-drift chemical ionization mass spectrometer. *Atmospheric Measurement Techniques*, 9, 6101-6116, 2016.
- Manninen, E.H., Petäjä, T., Asmi, E., Riipinen, I., Nieminen, T., Mikkilä, J., Hörrak, U., Mirme, A., Mirme, S., Laakso, L., Kerminen, V.-M., & Kulmala, M.: Long-term field measurements of charged and neutral clusters using Neutral cluster and Air Ion Spectrometer (NAIS). *Boreal Environment Research*, 2009, 591-605, 2009.
- 425 McMurry, P.H. New particle formation in the presence of an aerosol: rates, time scales, and sub-0.01  $\mu\text{m}$  size distributions *Journal of Colloid and Interface Science*, 95, 72-80, 1983.
- McMurry, P.H., Fink, M., Sakurai, H., Stolzenburg, M.R., Mauldin, R.L., Smith, J., Eisele, F., Moore, K., Sjostedt, S., Tanner, D., Huey, L.G., Nowak, J.B., Edgerton, E., & Voisin, D.: A criterion for new particle formation in the sulfur-rich Atlanta atmosphere. *Journal of Geophysical Research*, 110, doi:10.1029/2005jd005901, 2005.
- 430 O'Dowd, C.D., Aalto, P., Hmeri, K., Kulmala, M., & Hoffmann, T.: Aerosol formation: atmospheric particles from organic vapours. *Nature*, 416, 497-498, doi:10.1038/416497a, 2002.
- Petäjä, T., Mauldin, R.L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., & Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site. *Journal of Aerosol Science*, 9, 7435-7448, 2009.
- 435 Qi, X.M., Ding, A.J., Nie, W., Petäjä, T., Kerminen, V.-M., Herrmann, E., Xie, Y.N., Zheng, L.F., Manninen, H., Aalto, P., Sun, J.N., Xu, Z.N., Chi, X.G., Huang, X., Boy, M., Virkkula, A., Yang, X.-Q., Fu, C.B., & Kulmala, M.: Aerosol size distribution and new particle formation in the western Yangtze River Delta of China: 2 years of measurements at the SORPES station. *Atmospheric Chemistry and Physics*, 15, 12445-12464, doi:10.5194/acp-15-12445-2015, 2015.
- 440 Riipinen, I., Sihto, S.-L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinilä, K., Kerminen, V.-M., Laaksonen, A., & Lehtinen, K.E.J.: Connections between atmospheric sulphuric acid and new particle formation during QUEST III-IV campaigns in Heidelberg and Hyytiälä. *Atmospheric Chemistry and Physics*, 7, 1899-1914, doi:10.5194/acp-7-1899-2007, 2007.
- 445 Ristovski, Z.D., Suni, T., Kulmala, M., Boy, M., Meyer, N., K., Duplissy, J., Turnipseed, A., Morawska, L., & Baltensperger, U.: The role of sulphates and organic vapours in growth of newly formed particles in a eucalypt forest. *Atmospheric Chemistry and Physics*, 10, 2919-2926, 2010.
- Seinfeld, J.H., & Pandis, S.N.: *Atmospheric Chemistry and Physics* (2nd ed.). John Wiley & Sons, Inc., New Jersey, 2006.
- 450 Sihto, S.L., Kulmala, M., Kerminen, V.M., Dal Maso, M., Petäjä, T., Riipinen, I., Korhonen, H., Arnold, F., Janson, R., Boy, M., Laaksonen, A., & Lehtinen, K.E.: Atmospheric sulphuric acid and aerosol formation: implications from atmospheric measurements for nucleation and early growth mechanisms. *Atmospheric Chemistry and Physics*, 6, 4079-4091, 2006.
- 455 Spracklen, D.V., Carslaw, K.S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I., Merikanto, J., Mann, G.W., Chipperfield, M.P., Wiedensohler, A., Birmili, W., & Lihavainen, H.: Contribution of particle formation to global cloud condensation nuclei concentrations. *Geophysical Research Letters*, 35, doi:10.1029/2007gl033038, 2008.

- 460 Vanhanen, J., Mikkilä, J., Lehtipalo, K., Sipilä, M., Manninen, H.E., Siivola, E., Petäjä, T., & Kulmala, M.: Particle Size Magnifier for Nano-CN Detection. *Aerosol Science and Technology*, 45, 533-542, doi:10.1080/02786826.2010.547889, 2011.
- Wehner, B., Birmili, W., Ditas, F., Wu, Z., Hu, M., Liu, X., Mao, J., Sugimoto, N., and Wiedensohler, A.: Relationships between submicrometer particulate air pollution and air mass history in Beijing, China, 2004 - 2006, *Atmospheric Chemistry and Physics*, 8, 6155–6168, 2008.
- 465 Wang, Z., Wu, Z., Yue, D., Shang, D., Guo, S., Sun, J., Ding, A., Wang, L., Jiang, J., Guo, H., Gao, J., Cheung, H.C., Morawska, L., Keywood, M., & Hu, M.: New particle formation in China: Current knowledge and further directions. *The Science of the total environment*, 577, 258-266, doi:10.1016/j.scitotenv.2016.10.177, 2017.
- 470 Wang, Z. B., Hu, M., Yue, D. L., He, L. Y., Huang, X. F., Yang, Q., Zheng, J., Zhang, R. Y., and Zhang, Y. H.: New particle formation in the presence of a strong biomass burning episode at a downwind rural site in PRD, China, *Tellus B*, 65, 10.3402/tellusb.v65i0.19965, 2013a.
- Wang, Z. B., Hu, M., Wu, Z. J., Yue, D. L., He, L. Y., Huang, X. F., Liu, X. G., and Wiedensohler, A.: Long-term measurements of particle number size distributions and the relationships with air mass history and source apportionment in the summer of Beijing, *Atmos Chem Phys*, 13, 10159-10170, 10.5194/acp-13-10159-2013, 2013b.
- 475 Wang, Z.B., Hu, M., Sun, J.Y., Wu, Z.J., Yue, D.L., Shen, X.J., Zhang, Y.M., Pei, X.Y., Cheng, Y.F., & Wiedensohler, A.: Characteristics of regional new particle formation in urban and regional background environments in the North China Plain. *Atmospheric Chemistry and Physics*, 13, 12495-12506, doi:10.5194/acp-13-12495-2013, 2013c.
- 480 Wang, Z.B., Hu, M., Yue, D.L., Zheng, J., Zhang, R.Y., Wiedensohler, A., Wu, Z.J., Nieminen, T., & Boy, M.: Evaluation on the role of sulfuric acid in the mechanisms of new particle formation for Beijing case. *Atmospheric Chemistry and Physics*, 11, 12663-12671, doi:10.5194/acp-11-12663-2011, 2011.
- Weber, R.J., Marti, J.J., McMurry, P.H., Eisele, F.L., Tanner, D.J., & Jefferson, A.: Measurements of new particle formation and ultrafine particle growth rates at a clean continental site. *Journal of Geophysical Research: Atmospheres*, 102, 4375-4385, doi:10.1029/96jd03656, 1997.
- 485 Wu, Z., Hu, M., Liu, S., Wehner, B., Bauer, S., Maßling, A., Wiedensohler, A., Petäjä, T., Dal Maso, M., & Kulmala, M.: New particle formation in Beijing, China: Statistical analysis of a 1-year data set. *Journal of Geophysical Research*, 112, 2007.
- Xiao, S., Wang, M.Y., Yao, L., Kulmala, M., Zhou, B., Yang, X., Chen, J.M., Wang, D.F., Fu, Q.Y., Worsnop, D.R., & Wang, L.: Strong atmospheric new particle formation in winter in urban Shanghai, China. *Atmospheric Chemistry and Physics*, 15, 1769-1781, 10, doi:10.29/2006jd007406, 2015.
- 490 Yue, D. L., Hu, M., Zhang, R. Y., Wang, Z. B., Zheng, J., Wu, Z. J., Wiedensohler, A., He, L. Y., Huang, X. F., and Zhu, T.: The roles of sulfuric acid in new particle formation and growth in the mega-city of Beijing. *Atmospheric Chemistry and Physics*, 10, 4953-4960, doi:10.5194/acp-10-4953-2010, 2010.
- 495 Zheng, J., Khalizov, A., Wang, L., and Zhang, R.: Atmospheric pressure-ion drift chemical ionization mass spectrometry for detection of trace gas species. *Analytical Chemistry*, 82, doi:10.1016/j.atmosenv.2014.05.024, 7302-7308, 2010.
- Zheng, J., Hu, M., Zhang, R., Yue, D., Wang, Z., Guo, S., Li, X., Bohn, B., Shao, M., He, L., Huang, X., Wiedensohler, A., and Zhu, T.: Measurements of gaseous H<sub>2</sub>SO<sub>4</sub> by AP-ID-CIMS during CAREBeijing 2008 Campaign. *Atmospheric Chemistry and Physics*, 11, 7755-7765, doi:10.5194/acp-11-7755-2011, 2011.
- 500 Zheng, J., Ma, Y., Chen, M., Zhang, Q., Wang, L., Khalizov, A. F., Yao, L., Wang, Z., Wang, X., and Chen, L.: Measurement of atmospheric amines and ammonia using the high resolution time-of-flight chemical ionization mass spectrometry. *Atmospheric Environment*, 102, 249-259, doi:10.1016/j.atmosenv.2014.12.002, 2015a.
- Zheng, J., Yang, D., Ma, Y., Chen, M., Cheng, J., Li, S., and Wang, M.: Development of a new corona discharge based ion source for high resolution time-of-flight chemical ionization mass spectrometer to measure gaseous H<sub>2</sub>SO<sub>4</sub> and aerosol sulfate. *Atmospheric Environment*, 119, 167-173, doi:10.1016/j.atmosenv.2015.08.028, 2015b.
- 505

**Table 1: Characteristics of each campaign day.**

Date (mm/dd)	Classification	Max $J_{1.5}$ ( $\text{cm}^{-3} \text{s}^{-1}$ )	$N_{1-3}$ ( $\#/\text{cm}^3$ )	$A_{\text{Fuchs}}$ ( $\mu\text{m}^2/\text{cm}^3$ )	$L_{\Gamma}$	Wind direction*
03/12	Non-event	–	0	919.5	3.63	SW
03/13	NPF	156.0	26347.5	119.7	0.71	NW
03/14	Non-event	–	0	632.7	3.05	NW
03/15	Non-event	–	0	733.9	3.73	SW
03/16	Non-event	–	0	796.2	4.15	WSW
03/17	Non-event	–	0	1140.1	9.04	WSW
03/18	NPF	33.8	741.2	329.0	1.75	WNW
03/19	Undefined	Weak**	1643.67	240.8	1.40	SE
03/20	Non-event	–	137.9	348.8	1.74	NNW
03/21	Non-event	–	0	512.0	2.76	SSW
03/22	Non-event	–	0	457.6	2.58	E
03/23	NPF	30.1	3846.3	76.1	0.57	NNW
03/24	NPF	46.8	5576.7	145.2	0.76	NNW
03/25	NPF	57.0	4637.7	126.7	0.52	NNE
03/26	NPF	41.5	9640.9	100.4	0.71	N
03/27	NPF	31.2	2806.2	90.6	0.44	NW
03/28	Non-event	–	0	508.1	2.86	W
03/29	NPF	32.3	2449.8	121.0	0.69	NW
03/30	Undefined	17.7	2885.7	88.8	0.64	NW
03/31	Non-event	–	0	767.0	4.21	SW
04/01	NPF	50.9	5477	51.7	0.22	WNW
04/02	NPF	46.9	10002	63.1	0.31	NW
04/03	NPF	21.6	10962.9	105.7	0.24	NW
04/04	Non-event	–	442	398.2	3.09	SW
04/05	Non-event	–	185	391.2	2.33	NW
04/06	Non-event	–	0	365.5	1.71	SW

\*: Indicated by 12-hour backward trajectory (starting at noon, 500 m in altitude).

\*\* : Difficult to estimate.






Figure 1 Contour of measured particle size distributions during Mar. 12<sup>th</sup> to Apr. 6<sup>th</sup>. Identified thirteen non-event days and two undefined days are shadowed by grey and yellow background, respectively.






Figure 2: Time series for Fuchs surface area ( $A_{Fuchs}$ ), sulfuric acid concentration, and number concentration of 1-3 nm particles.

Typical NPF days and undefined days are shadowed by light blue and light green background, respectively.



520 **Figure 3:** The correlations between the estimated new particle formation rate,  $J_{1.5}$ , and sulfuric acid concentration during NPF event period on each NPF day. The regression line of  $J_{1.5}$  versus sulfuric acid concentration was exponentially fitted.  $n$  is the exponent. Data on Mar. 29<sup>th</sup> was not included because the correlation was not significant ( $p = 0.34$ ).



525 **Figure 4: Normalized growth enhancement factor,  $\Gamma$ , in this campaign in comparison to those reported for other campaigns.  $\Gamma$  was normalized by the geometric mean value in each campaign.**




Figure 5: (a) The relationship between Fuchs surface area and number concentration of 1-3 nm particles,  $N_{1-3}$ . The relative concentration of measured sulfuric acid is represented by symbol size, i.e., the higher the relative concentration, the bigger the symbol size. Data points are 5-minute-resolved. (b) Frequencies of observed NPF days, undefined days and non-event days in comparison to daily averaged Fuchs surface area. On typical NPF days and undefined days,  $A_{\text{Fuchs}}$  was averaged during NPF event periods. On non-event days, it was averaged between 8:00 and 16:00.  $A_{\text{Fuchs}}$  values were binned in logarithmic scale ranging from 45 to 1150.

530

535




Figure 6: (a) The correlation between  $L_T$  and  $A_{Fuchs}$  (data from Table 1) in this campaign. NPF days, non-event days, and undefined days are shown as different symbols. The regression was based on all campaign days. (b) The correlation between  $L_T$  and  $A_{Fuchs}$  estimated for this study in comparison to other campaigns.




Figure 7: The schematic of governing factors for  $L_\Gamma$  at different locations. Concentration of growth relevant gaseous precursors is represented by  $\Gamma \cdot N_1$ , where  $\Gamma$  is the growth enhancement factor and  $N_1$  is sulfuric acid number concentration. Background colour represents the magnitude of  $L_\Gamma$ . Data for each location are shown as different symbols (circle: Beijing; square: Atlanta; diamond: Boulder; triangle: Hyytiälä). The ellipse and the boxes were artificially drawn to illustrate the variations. Tecamac was not included due to the lack of data on non-event days. Both axes are in log scale.

545






Figure 8: Average contribution of the net CoagSnk,  $dN/dt$ , and the condensational growth term (GR term) to the estimated new particle formation rate,  $J_{1.5}$ , on identified typical NPF days. The percentage presented in each column is the relative ratio of the net CoagSnk compared to  $J_{1.5}$  of that NPF event. Note that only the time period when  $dN/dt$  was positive during a NPF event was taken in to account when calculating average contribution.

550



555 Figure 9: (a) Contour of measured particle size distributions on Apr. 2<sup>nd</sup>, 3<sup>rd</sup>, and 4<sup>th</sup>. (b) Representative parameters on these three NPF days. Time periods when  $L_{\Gamma}$  was lower than 1.0 are shadowed by light blue background. When wind speed was close to zero, the corresponding wind direction data were not included in the plot.






560 **Figure 10:** Normalized distribution of cumulative Fuchs surface area,  $\overline{A_{\text{Fuchs}}}$ , as a function of the particle diameter,  $d_p$ , on two

NPF days (red circle) and two non-event days (blue diamond).  $\overline{A_{\text{Fuchs}}}$  is equal to  $A_{\text{Fuchs}}$  when  $d_p$  is approaching positive infinity.

$d\overline{A_{\text{Fuchs}}}/d\log d_p$  is normalized by  $A_{\text{Fuchs}}$ .



565 **Figure 11: Relationship between hourly averaged  $A_{Fuchs}$  and the  $\text{PM}_{2.5}$  mass concentration in Beijing. Data when  $A_{Fuchs}$  changed rapidly was not included to avoid potential influence caused by the distance between Wanliu station and our campaign site. NPF period, daytime (8:00-16:00) on non-event days and undefined days, and other time are shown as different symbols. The regression of  $A_{Fuchs}$  versus the  $\text{PM}_{2.5}$  mass concentration was based on all the data. The proposed criterion for the occurrence of NPF events, i.e.,  $A_{Fuchs}$  is lower than  $200 \mu\text{m}^2/\text{cm}^3$  (the  $\text{PM}_{2.5}$  mass concentration is lower than  $30 \mu\text{g}/\text{m}^3$ ), is shadowed by**  
 570 **light green background.**