



Climate Impact of Polar Mesospheric and Stratospheric Ozone Losses due to Energetic Particle Precipitation

Katharina Meraner¹ and Hauke Schmidt¹

¹Max Planck Institute for Meteorology, Bundesstraße 53, 20146 Hamburg, Germany *Correspondence to:* Katharina Meraner (katharina.meraner@mpimet.mpg.de)

Abstract. Energetic particles enter the polar atmosphere and enhance the production of nitrogen oxides and hydrogen oxides in the winter stratosphere and mesosphere. Both components are powerful ozone destroyers. Recently, it has been inferred from observations that the direct effect of energetic particle precipitation (EPP) causes significant long-term mesospheric ozone variability. Satellites observe a decrease in mesospheric ozone by up to 34 % between EPP maximum and EPP minimum.

5 Here, we analyze the climate impact of polar mesospheric and polar stratospheric ozone losses due to EPP in the coupled climate model MPI-ESM. Using radiative transfer modeling, we find that the radiative forcing of a mesospheric ozone loss during polar night is small. Hence, climate effects of a mesospheric ozone loss due to energetic particles seem unlikely. A stratospheric ozone loss due to energetic particles warms the winter polar stratosphere and subsequently weakens the polar vortex. However, those changes are small, and few statistically significant changes in surface climate are found.

10 1 Introduction

Energetic particles enter the Earth's atmosphere near the magnetic poles altering the chemistry of the middle and upper atmosphere. Energetic particle precipitation (EPP) is the major source of nitrogen oxides (NO_x) and hydrogen oxides (HO_x) in the polar middle and upper atmosphere (Crutzen et al., 1975; Solomon et al., 1981). Both chemical components catalytically deplete ozone; NO_x mainly below and HO_x mainly above 45 km.

- 15 HO_x is short-lived in the middle atmosphere and depletes mainly the ozone in the mesosphere. In contrast, NO_x persists up to several months in the polar winter middle atmosphere. Inside the polar vortex, NO_x can be transported downward from the lower thermosphere to the stratosphere where it depletes ozone (Funke et al., 2007; Sinnhuber et al., 2014). Ozone loss influences stratospheric temperature and the polar vortex. The Northern Annual Mode (NAM) index is often used to describe the strength of the polar vortex, with positive NAM values indicating a strong polar vortex and negative NAM values indicating
- 20 a weak polar vortex. Observations indicate that anomalous weather regimes associated with the NAM index can propagate from the stratosphere down to the surface (Baldwin and Dunkerton, 2001). Hence, energetic particle precipitation may provide a link from space weather to surface climate. Here, we study the impact of an ozone loss due to EPP on the circulation and subsequently on climate. Discussed are both a polar mesospheric and a polar stratospheric ozone loss.

Since the discovery of the ozone hole in the mid-1980s, the climate impact of a stratospheric ozone loss has been intensively studied (e.g., Shine, 1986; Randel and Wu, 1999; Lubis et al., 2016). Most studies concentrated on the climate impact of





the ozone hole during austral spring and reported a cooling in the spring Southern Hemispheric stratosphere due to reduced absorption if solar radiation and a strengthening of the polar vortex. In contrast, our study concentrates on an ozone loss during the boreal polar night. During polar night reduced ozone slightly decreases the infrared cooling of the polar stratosphere resulting in a net stratospheric warming (Graf et al., 1998; Langematz et al., 2003).

5 Several studies suggested a significant influence of EPP on the surface air temperature during winter. Seppälä et al. (2009) analyzed surface temperature changes in reanalysis data for years with various strengths of EPP. They found a warming over Eurasia and a cooling over Greenland for winters with enhanced EPP, but could not rule out that the estimated changes are induced by NAM variability independent of EPP.

Other studies relied on atmospheric chemistry models, which showed similar surface temperature change patterns as found 10 in the reanalysis data (e.g., Rozanov et al., 2005; Baumgaertner et al., 2011; Arsenovic et al., 2016). They reported a small cooling in the polar winter stratosphere due to EPP, although the radiative effect of a polar night ozone loss should be a warming. This stratospheric cooling is attributed to a dynamical, adiabatic cooling caused by a decrease in the mean meridional circulation (Schoeberl and Strobel, 1978; Christiansen et al., 1997). Langematz et al. (2003) suggested that the weaker mean meridional circulation is caused by a decrease in midlatitude tropospheric wave forcing. The aforementioned model studies

- 15 analyzing the climate impact of EPP relied on relatively few simulation years and applied complex forcings. Instead of prescribing ozone, these studies simulated EPP effects by changing the production of NO_x and HO_x and modeling the effects on ozone interactively. This could potentially be more realistic than simulations with prescribed ozone anomalies but introduces uncertainties related to the representation of chemistry and transport in the model, and renders the understanding of the effects more complicated as the ozone forcing varies in space and time. To avoid these difficulties and to obtain a clear signal-to-noise
- 20 ratio, we use an idealized ozone forcing and a long simulation period.

Commonly, the effects of EPP are classified into direct and indirect effects (Randall et al., 2006, 2007). Direct effects are the effects of the local production of NO_x and HO_x , whereas indirect effects are the effects of the NO_x transport from the thermosphere to the stratosphere. Whereas most of the above mentioned studies discuss a mainly stratospheric ozone loss due to the indirect EPP effect, Andersson et al. (2014) suggested a potential climate influence of a mesospheric ozone loss due to

- the direct EPP effect. By using satellite observations they showed that HO_x causes long-term variability in mesospheric ozone up to 34 % between EPP maximum and EPP minimum. Arsenovic et al. (2016) were the first to include the direct effect of HO_x local production due to EPP in a chemistry-climate model. They found a similar mesospheric ozone loss as Andersson et al. (2014) and ultimately, reported a cooling over Greenland and a warming over Eurasia. Arsenovic et al. (2016) also considered the indirect effect of the NO_x descent. Hence, the sole impact of a mesospheric ozone loss due to the direct EPP effect as suggested by Anderson et al. (2014) remains unclear
- 30 suggested by Andersson et al. (2014) remains unclear.

This paper studies the circulation and climate impact of idealized mesospheric and stratospheric ozone losses that could be attributed to energetic particle precipitation. The idealized mesospheric ozone loss that we prescribe may be considered to be mostly a direct EPP effect, whereas the prescribed stratospheric ozone loss should be considered indirect. We use simulations with the Max Planck Institute Earth System Model (MPI-ESM) applying an idealized ozone forcing in either the mesosphere

35 or the stratosphere. Additionally, we use a radiative transfer model to quantify the radiative forcing of ozone at different





altitudes and months. Ultimately, we discuss whether an ozone loss in the middle atmosphere due to EPP has the potential to significantly alter the surface climate. Section 2 describes the MPI-ESM as well as the radiative transfer model. Section 3 links mesospheric and stratospheric ozone losses to changes in the atmospheric temperatures and winds. Finally, Section 4 summarizes and discusses the main outcomes and limitations of this study.

5 2 Models and numerical experiments

2.1 MPI-ESM: The Max Planck Institute Earth System Model

The Max Planck Institute Earth System Model (MPI-ESM; Giorgetta et al. (2013)) consists of the coupled atmospheric and ocean general circulation models, ECHAM6 (Stevens et al., 2013) and MPIOM (Jungclaus et al., 2013) as well as of the land and vegetation model JSBACH (Reick et al., 2013) and of the model for marine bio-geochemistry HAMOCC (Ilyina et al., 2013). We use the 'mixed-resolution' configuration of the model (MPI-ESM-MR). The ocean model uses a tripolar quasi-isotropic grid with a nominal resolution of 0.4° and 40 vertical layers. ECHAM6 is run with a triangular truncation at wave number 63 (T63), which corresponds to 1.9° in latitude and longitude. The vertical grid contains 95 hybrid sigma-pressure levels resolving the atmosphere from the surface up to 0.01 hPa. The vertical resolution is nearly constant (700 m) from the upper troposphere to the middle stratosphere and less than 1000 m at the stratopause. The time steps in the atmosphere and 15 ocean are 450 and 3600 s, respectively.

The model has been used for many simulations within the CMIP5 (Coupled Model Intercomparison Project Phase 5) framework (Taylor et al., 2012). An overview of the dynamics of the middle atmosphere in these simulations is given by Schmidt et al. (2013). In this study, the preindustrial CMIP5 simulation (piControl) is used as reference. The forcing is constant in time and uses pre-industrial conditions (1850 AD) for the greenhouse gases. Solar irradiance and ozone concentrations are averaged over a solar cycle (1844 – 1856 for the solar irradiance and 1850 – 1860 for ozone concentrations). No volcanic

20

forcing is applied. A period of 150 years of this simulation is used.

In order to analyze the impact of ozone changes on the model climate, two additional experiments with reduced ozone concentration are carried out. In one experiment, the mesospheric ozone is reduced by 40 % between 0.01 hPa and 0.1 hPa polewards of 60° N (this is called "meso-O₃"). In the other experiment, stratospheric ozone is reduced by 20 % between

- 25 1 hPa and 10 hPa polewards of 60° N (this is called "strato-O₃"). The magnitude of the ozone losses is based on the results from previous studies for boreal winter. Andersson et al. (2014) showed that the direct HO_x effect induces a variability in mesospheric ozone up to 34 % from November to February in satellite data. Baumgaertner et al. (2011) simulated polar winter losses of up to 20 % in the upper stratosphere due to EPP in a model. Note that the applied ozone losses are slightly larger than the EPP influence diagnosed from observations. We use the stronger forcing to obtain a clear signal from which to diagnose
- 30 the sensitivity of the climate to an ozone loss due to EPP. However, this implies a potentially overestimated climate response. To facilitate the experiment design, we applied the ozone losses constant over time. Although we concentrate our analysis on boreal winter high latitudes, this allows us to gain insights on boreal spring (i.e., the transition time from polar night to polar day). Observed ozone losses in summer are in general smaller than during winter, but this idealized setting allows an easy





comparison of potential effects during the different seasons. It is unlikely that signals in summer affect the climate of the next winter.

Both experiments are forced by the same conditions as the piControl experiment. Moreover, the simulations are restarted from the same year in the piControl experiment. This ensures that the ocean state is similar in all experiments. For both

5 simulations 150 years are simulated. In the Sections 3.2 and 3.3 the differences between the experiments and the control simulation (i.e., piControl) are analyzed. Statistical significance is calculated using the 95% confidence intervals assuming normally-distributed regression errors and using the 0.975 percentile of Student's t-distribution with the appropriate degrees of freedom. Properties of two simulations are considered statistically significantly different if the mean value of the control simulation is outside 95% confidence interval of the experiment.

10 2.2 The radiative transfer model PSrad

The radiative transfer scheme of MPI-ESM is based on the rapid radiation transfer suite of models optimized for general circulation models (RRTMG; (Mlawer et al., 1997; Iacono et al., 2008)). The RRTMG is widely used and its ability to calculate radiative forcing has been evaluated by Iacono et al. (2008). In its stand-alone version, it is used here to study the impact of ozone on heating rates. The radiative transfer scheme is further described in Pincus and Stevens (2013) and Stevens et al. (2013) and onwards we will refer to it as the radiative transfer model "PSrad".

The shortwave and longwave components are calculated separately. Furthermore, optical properties for gases, clouds and aerosols are computed separately and then combined to compute heating rates. PSrad expects profiles of gases (H₂O, N₂O, CH₄, CO, O₃), profile of cloud parameters as well as additional parameters (e.g., albedo and zenith angle) as input. For gases, we use multi-year monthly means representative for the late 20^{th} century provided by the atmospheric and chemistry

- 20 model HAMMONIA (Hamburg Model for Neutral and Ionized Atmosphere; Schmidt et al. (2006)). For the albedo and cloud properties (e.g., cloud fraction, cloud water/ice content), multi-year monthly means from the piControl experiment are used. All quantities are extracted for 75° N. The zenith angle is calculated for 12 UTC at 75° N/0° E for the 15th of each month. The latitude of 75° N is chosen as it represents a mean polar latitude. The results are insensitive to the actual latitude, the main difference at other polar latitudes is the length of the polar night.
- To quantify the impact of ozone on the heating rates, we perform multiple runs in which for each run the ozone concentration of a single layer is set to 0 once. Then the differences between each single run and a control run are calculated and finally, added up.

3 Results

15

3.1 Ozone effects on the heating rates

30 An ozone loss directly alters the atmospheric energy transfer. Before analyzing circulation and climate impacts due to ozone losses, we study the heating rate response using the radiative transfer model PSrad. The heating rates are calculated for the





polar latitude of 75° N (see Figure 1). As the effect of EPP is most important at the winter polar cap, we will concentrate our analysis on boreal winter high latitudes.

In the shortwave part of the spectrum, ozone strongly absorbs solar radiation and heats the whole atmosphere. The strongest heating (about 12 K/day) occurs at the altitudes near the maximum volume mixing rate of ozone. An ozone loss would, hence,

5 result in a relative cooling due to reduced heating. The ozone heating and, hence, the cooling caused by an ozone reduction are getting smaller for larger zenith angles and vanish in polar night.

In the longwave part of the spectrum, the radiative effect of ozone is highly temperature dependent. Ozone cools the atmosphere via infrared emission in the stratosphere and in warm regions of the mesosphere below 0.1 hPa (see Figure 1b). The strongest cooling (about -2 K/day) occurs at the stratopause. In the troposphere and in the cold regions of the mesosphere

10 above 0.1 hPa, the absorption of outgoing radiation exceeds the infrared emission resulting in a heating of the atmosphere due to ozone.

In total, the shortwave heating dominates all sunlit months. During polar night, ozone cools the atmosphere between 0.1 and 100 hPa and, hence, an ozone loss in the stratosphere and lower mesosphere results in a warming. Near the terminator (e.g., at 75° N in November and February), the net influence of ozone is more complex: At some altitudes ozone heats and at some it

15 cools the atmosphere. The net radiative forcing of an ozone loss depends on when and where ozone is reduced. For example, in November, a stratospheric (1 hPa) ozone loss leads to a heating, but a mesospheric (0.1 hPa) ozone loss to a cooling.

These results are in line with previous work. It is widely accepted that an ozone loss in spring and summer leads to a stratospheric cooling (e.g., Shine, 1986; Randel and Wu, 1999). Some studies analyzed the radiative forcing of a winter stratospheric ozone loss. Graf et al. (1998) showed that the observed stratospheric ozone loss in the late 20th century led to a winter warm-

20 ing and a summer cooling in a GCM. Using a radiative transfer model with fixed dynamical heating, Langematz et al. (2003) confirmed that a stratospheric ozone loss over the winter pole results in a stratospheric warming. Shine (1986) showed that the shortwave cooling of the stratosphere due to an ozone loss dominates in all sunlit months the infrared heating due to an ozone loss.

The above stated results are confirmed by the actual heating rate anomalies induced by the applied ozone losses in the 25 experiments 'meso-O₃' and 'strato-O₃' (not shown). The heating rates are calculated at the first time step of the model at which 26 the radiation is updated (1 January) excluding any feedbacks occurring only at later time steps. Compared to the total heating 27 rates of piControl, the changes in heating rates caused by a 40 % reduction of mesospheric ozone in polar night is very small 28 (below -2 %) and of a 20 % reduction of stratospheric ozone small (about -10 %).

3.2 Climate effects of a mesospheric ozone loss

30 As changes in heating rates due a reduced ozone during polar night are small, one might reason that climate impact of a winter polar ozone loss is small. But large effects may occur in regions slightly outside the polar night. Furthermore, several studies suggested that changes in the heating rates due to a winter polar ozone loss leads to a dynamical cooling (e.g., Baumgaertner et al., 2011; Arsenovic et al., 2016), whereas the initial radiative forcing suggests a warming. Hence, we further analyze the climate impact of a winter polar ozone loss. As large variations in the polar vortex can propagate downward and







Figure 1. Monthly mean heating rates of ozone [K/day] for 75° N calculated by the radiative transfer model PSrad for (a) shortwave, (b) longwave and (c) total (shortwave + longwave) radiation.

affect the surface climate, we first concentrate on the circulation changes of the middle atmosphere due to an ozone loss, which are a prerequisite for a potential climate impact of EPP. In the following, we analyze the climate effect of an idealized polar mesospheric ozone loss, while in Section 3.3 we analyze the climate effect of an idealized polar stratospheric ozone loss.

Figures 2a and 2d show the zonal mean temperature and zonal mean wind simulated for boreal winter (December - February).
Main observed characteristics of the zonal mean temperature, e.g., the stratopause tilt from the summer towards the winter pole, are well reproduced. The changes in the zonal mean zonal wind are consistent with the temperature changes via the thermal wind balance. In most regions, the difference between meso-O₃ and piControl is very small (see Figures 2b and 2e).

Near the winter pole, a dipole structure emerges with cooling in the upper stratosphere and warming in the mesosphere. According to our radiative transfer calculations a mesospheric winter polar ozone loss should lead to a cooling. However, the temperature differences are small (below 1 K) and not significant at the 95 % level. As the applied forcing is very small, small and low significant values are expected. At the winter pole, the polar vortex slightly weakens, whereas the mesospheric winds strengthen: these differences are not significant. As stated above, large variations in the winter polar vortex can propagate to the surface influencing the surface climate. However, the changes reported here are small. The anomalies reaching the troposphere are statistical artifacts. Indeed, the surface temperature reveals no statistically significant change (not shown).

- 15 Although, the temperature and wind signals are not statistically significant after 150 simulated years, nevertheless, it makes sense to analyze if the signals could have a physical explanation and not be purely accidental. Note that with fewer simulation years apparently very different results can be obtained. Analyzing different simulation periods we obtain mesospheric warming and cooling of apparent significance. Particularly, we calculated a statistically significant weakening of the polar vortex when using only the first 80 simulation years. The high degree of internal variability of the winter polar stratosphere can obviously
- 20 create wrong apparent signals. The most dramatic demonstration of this variability are major sudden stratospheric warmings





5



Figure 2. (upper row) Zonal mean temperature [K] and (lower row) zonal mean zonal wind [m/s] averaged over December - February (DJF) for (a,d) piControl, (b,e) the difference between meso- O_3 and piControl and (c,f) the difference between strato- O_3 and piControl. Shaded areas are statistically significant at the 95% confidence interval. The black, dashed boxes highlight the regions where ozone is reduced.

(SSW), which occur on average about 6 times per decade in the Northern Hemisphere (see Charlton and Polvani (2007) for more information on SSW). A short simulation period may lead to an over-representation or under-representation of SSWs. Over our whole simulation period (150 years) the number of major SSWs is balanced in all three experiments. In total, there are 102 events in piControl, 99 events in meso- O_3 and 109 events in strato- O_3 (using a reversal of the zonal wind at 60° N and 10 hPa as criterion of a major SSW occurrence).





5

3.3 Climate effects of a stratospheric ozone loss

In this section, we analyze the climate effect of an idealized polar stratospheric ozone loss. Figures 2a and 2d show the zonal mean temperature and zonal wind simulated for boreal winter (December - February) for piControl, and Figures 2c and 2f the difference between strato- O_3 and piControl. The winter stratosphere warms due to an ozone loss as expected from the calculations with the radiative transfer model. As a consequence of the warming, the stratospheric winds weaken. The small mesospheric cooling likely results from enhanced eastward momentum deposition from gravity waves as shown by Lossow et al. (2012). The statistically significant warming of the summer mesopause is an indication of inter-hemispheric coupling as discussed by Karlsson and Becker (2016).

Figure 2 shows only changes for the mean over December to February, while the radiative transfer model suggests that the month-to-month variability of the forcing is large. To study whether the impact of a stratospheric ozone loss differs over the course of the winter, we analyze the monthly means of temperature and zonal wind (see Figure 3). An ozone loss during most of the polar night (except December) leads to a warming, whereas at all other times and locations it leads to a cooling. This agrees with the calculations of the radiative transfer model and with our assumption that the winter cooling is not affected by a strong summer warming. However, the cooling in December is unexpected from the radiative transfer modeling. Kodera and Kuroda

- 15 (2002) argued that the polar winter atmosphere transits from a radiatively controlled state in early to a dynamically controlled state in late winter. Given the opposite sign of the diabatic forcing, the simulated cooling must be dynamically caused already in December. This is in agreement with early model studies which showed that uniform ozone losses lead to dynamical cooling at the boreal winter polar latitudes (e.g., Schoeberl and Strobel, 1978; Kiehl and Boville, 1988). Langematz et al. (2003) suggested that the dynamical cooling is due to a weakening of the mean meridional circulation related to reduced wave forc-
- 20 ing caused by a reduction of mid-latitude wave flux into the stratosphere. Similarly, in our simulations we find a (albeit not significant) reduction of the zonal mean eddy heat flux at 100 hPa in the midlatitudes from December to March (not shown). However, the origin of the changes in the wave activity remains unexplained. Also Baumgaertner et al. (2011) reported a dynamical cooling in the winter polar stratosphere due to EPP. However, in their model the cooling dominates the winter (DJF) signal, whereas we obtain a small warming for the DJF average (see Figure 2).
- The zonal wind changes consistently with the temperature changes via the thermal wind balance. Simultaneously with the warming (cooling), the polar wind weakens (strengthens). Anomalies in the polar vortex occasionally reach the troposphere (e.g., the strengthening in November or the weakening in December or February). Although, most of those changes are not significant, some disturbances in the polar vortex may still force the surface temperature (see Figure 4). In our simulations for boreal winter, stratospheric ozone loss cools large parts of the northern high latitudes from northern Europe to Eurasia and over
- 30 northern America. Over Greenland and the pole, the surface warms. This is consistent with the weakening of the zonal wind in December (see Figure 3 d). However, most changes are small and not significant.







Figure 3. Monthly mean (upper row) temperature averaged between 60° N and 90° N [K] and (lower row) zonal wind [m/s] for 60° N for (a,c) piControl and (b,d) the difference between strato-O₃ and piControl. Shaded areas are statistically significant at the 95 % confidence interval.







Figure 4. Surface temperature [K] for the difference between strato- O_3 and piControl (a) averaged over December - February (DJF) for the Northern Hemisphere and (b) averaged over September - November (SON) for the Southern Hemisphere. Shaded areas are statistically significant at the 95 % confidence interval.

4 Summary and conclusion

In this study, we analyzed the climate impact of mesospheric and stratospheric ozone losses. Although this study is motivated from the enhancement of NO_x due to energetic particle precipitation (EPP), the results presented here could also be applied to other processes causing ozone destruction. The radiative forcing of polar ozone is calculated by the radiative transfer model PSrad. In sensitivity studies with the Max Planck Institute Earth System Model (MPI-ESM), we reduced ozone either by 40 % in the winter polar mesosphere or by 20 % in the winter polar stratosphere.

Recently, Andersson et al. (2014) showed that the direct EPP-HO_x effect induces large long-term variability in winter mesospheric ozone. They suggested that these large changes may have an impact on climate. Following their idea, we analyzed the atmospheric response to a mesospheric ozone loss. We found that the winter atmospheric changes due to a mesospheric ozone

5

ozone is very small during polar night, which makes the small dynamic response plausible. Several studies analyzed the climate effect of a stratospheric ozone loss due to EPP. Seppälä et al. (2009) calculated a

loss in our model are negligible. Calculations with a radiative transfer model showed that the radiative forcing of mesospheric

correlation of the winter surface temperature and energetic particle precipitation in reanalysis data. However, they could not rule out an accidental occurrence of the correlation. Since then several model studies tried to establish a physical link between EPP





and climate (Baumgaertner et al., 2011; Rozanov et al., 2012; Arsenovic et al., 2016). In all these model studies, a dynamical cooling of the winter polar stratosphere due to energetic particle precipitation was simulated. In our model, a stratospheric ozone loss during polar night (except December) results in a warming, whereas at all other times and locations it leads to a cooling. This agrees with the calculations of the radiative transfer model. We obtained a cooling during December due to

- 5 stratospheric ozone loss caused by a reduced vertical wind. However, the changes in the polar winter stratosphere are small and not significant in our model. Consequently, also the impact on the simulated winter surface temperature is weak. In contrast to the above mentioned studies, in our experiment the dynamical feedback leading to the stratospheric cooling is not dominant throughout the boreal winter. However, the earlier model studies were based on only a few simulation years. Using only the first 80 years of our simulations we obtained false positives. The high degree of internal variability of the polar vortex can
- 10 create wrong apparent signals.

Our results suggest that the climate impact of an ozone loss due to EPP is small. As the radiative forcing of mesospheric ozone loss is negligible, a significant climate impact of a mesospheric ozone change as suggested by Andersson et al. (2014) seems unlikely. However, we cannot rule out that our experimental design with an ozone loss constant in time is too simplified, and that the time and altitude dependent loss caused by the downward propagation of NO_x concentrations would create a

15 different result. Furthermore, the catalytic destruction of ozone by NO_x works only effectively if sunlight is available. The influence of EPP induced NO_x may be larger near the terminator.

Finally, although previous studies have shown that MPI-ESM reproduces stratospheric temperature responses to forcings reasonably well (e.g., Bittner et al., 2016; Schmidt et al., 2013), the possibility remains that the model's sensitivity to ozone loss is biased low. To address this, we would like to encourage multi-model studies on EPP climate impact as currently suggested

20

for the third phase of the SOLARIS-HEPPA project, which investigates solar influences on climate as part of the 'Stratospheretroposphere Processes And their Role in Climate' (SPARC) project.

5 Code and data availability

Primary data and scripts used in the analysis and other supplementary information that may be useful in reproducing the author's work are archived by the Max Planck Institute for Meteorology and can be obtained by contacting publications@mpimet.mpg.de.

25 Acknowledgements. The authors acknowledge scientific and practical input from Matthias Bittner and Elisa Manzini. This study was supported by the Max-Planck-Gesellschaft (MPG) and computational resources were made available by Deutsches Klimarechenzentrum (DKRZ) through support from Bundesministerium für Bildung und Forschung (BMBF).





References

- Andersson, M. E., Verronen, P. T., Rodger, C. J., Clilverd, M. A., and Seppälä, A.: Missing driver in the Sun–Earth connection from energetic electron precipitation impacts mesospheric ozone, Nature Communications, 5, doi:10.1038/ncomms6197, 2014.
- Arsenovic, P., Rozanov, E., Stenke, A., Funke, B., Wissing, J. M., Mursula, K., Tummon, F., and Peter, T.: The influence of Middle Range Energy Electrons on atmospheric chemistry and regional climate, Journal of Atmospheric and Solar-Terrestrial Physics, doi:10.1016/j.jastp.2016.04.008, http://www.sciencedirect.com/science/article/pii/S1364682616301080, 2016.
 - Baldwin, M. P. and Dunkerton, T. J.: Stratospheric Harbingers of Anomalous Weather Regimes, Science, 294, 581–584, doi:10.1126/science.1063315, http://science.sciencemag.org/content/294/5542/581, 2001.
- Baumgaertner, A. J. G., Seppälä, A., Jöckel, P., and Clilverd, M. A.: Geomagnetic activity related NOx enhancements and polar surface air temperature variability in a chemistry climate model: modulation of the NAM index, Atmos. Chem. Phys., 11, 4521–4531, doi:10.5194/acp-11-4521-2011, 2011.
 - Bittner, M., Timmreck, C., Schmidt, H., Toohey, M., and Krüger, K.: The impact of wave-mean flow interaction on the Northern Hemisphere polar vortex after tropical volcanic eruptions, Journal of Geophysical Research: Atmospheres, 121, 2015JD024603, doi:10.1002/2015JD024603, http://onlinelibrary.wiley.com/doi/10.1002/2015JD024603/abstract, 2016.
- 15 Charlton, A. J. and Polvani, L. M.: A New Look at Stratospheric Sudden Warmings. Part I: Climatology and Modeling Benchmarks, Journal of Climate, 20, 449–469, doi:10.1175/JCLI3996.1, http://journals.ametsoc.org/doi/abs/10.1175/JCLI3996.1, 2007.
 - Christiansen, B., Guldberg, A., Hansen, A. W., and Riishøjgaard, L. P.: On the response of a three-dimensional general circulation model to imposed changes in the ozone distribution, Journal of Geophysical Research: Atmospheres, 102, 13 051–13 077, doi:10.1029/97JD00529, http://onlinelibrary.wiley.com/doi/10.1029/97JD00529/abstract, 1997.
- 20 Crutzen, P. J., Isaksen, I. S. A., and Reid, G. C.: Solar Proton Events: Stratospheric Sources of Nitric Oxide, Science, 189, 457–459, doi:10.1126/science.189.4201.457, 1975.
 - Funke, B., López-Puertas, M., Fischer, H., Stiller, G. P., von Clarmann, T., Wetzel, G., Carli, B., and Belotti, C.: Comment on "Origin of the January–April 2004 increase in stratospheric NO2 observed in northern polar latitudes" by Jean-Baptiste Renard et al., Geophysical Research Letters, 34, L07 813, doi:10.1029/2006GL027518, http://onlinelibrary.wiley.com/doi/10.1029/2006GL027518/abstract, 2007.
- 25 Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, Journal of Advances in Modeling
- Earth Systems, 5, 572–597, doi:10.1002/jame.20038, http://onlinelibrary.wiley.com/doi/10.1002/jame.20038/abstract, 2013.
 Graf, H.-F., Kirchner, I., and Perlwitz, J.: Changing lower stratospheric circulation: The role of ozone and greenhouse gases, Journal of Geophysical Research: Atmospheres, 103, 11251–11261, doi:10.1029/98JD00341, http://onlinelibrary.wiley.com/doi/10.1029/98JD00341/abstract, 1998.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived
 greenhouse gases: Calculations with the AER radiative transfer models, Journal of Geophysical Research: Atmospheres, 113, D13 103, doi:10.1029/2008JD009944, http://onlinelibrary.wiley.com/doi/10.1029/2008JD009944/abstract, 2008.





- Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, Journal of Advances in Modeling Earth Systems, 5, 287–315, doi:10.1029/2012MS000178, http://onlinelibrary.wiley.com/doi/10.1029/2012MS000178/abstract, 2013.
- 5 Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and von Storch, J. S.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, Journal of Advances in Modeling Earth Systems, 5, 422–446, doi:10.1002/jame.20023, http://onlinelibrary.wiley.com/doi/10.1002/jame.20023/abstract, 2013.

Karlsson, B. and Becker, E.: How Does Interhemispheric Coupling Contribute to Cool Down the Summer Polar Mesosphere?, Journal of

- Climate, 29, 8807–8821, doi:10.1175/JCLI-D-16-0231.1, http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-16-0231.1, 2016.
 Kiehl, J. T. and Boville, B. A.: The Radiative-Dynamical Response of a Stratospheric-Tropospheric General Circulation Model to Changes in Ozone, Journal of the Atmospheric Sciences, 45, 1798–1817, doi:10.1175/1520-0469(1988)045<1798:TRDROA>2.0.CO;2, http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281988%29045%3C1798%3ATRDROA%3E2.0.CO%3B2, 1988.
- Kodera, K. and Kuroda, Y.: Dynamical response to the solar cycle, Journal of Geophysical Research: Atmospheres, 107, 4749, doi:10.1029/2002JD002224, http://onlinelibrary.wiley.com/doi/10.1029/2002JD002224/abstract, 2002.
- Langematz, U., Kunze, M., Krüger, K., Labitzke, K., and Roff, G. L.: Thermal and dynamical changes of the stratosphere since 1979 and their link to ozone and CO2 changes, Journal of Geophysical Research: Atmospheres, 108, 4027, doi:10.1029/2002JD002069, http://onlinelibrary.wiley.com/doi/10.1029/2002JD002069/abstract, 2003.

Lossow, S., McLandress, C., Jonsson, A. I., and Shepherd, T. G.: Influence of the Antarctic ozone hole on the polar mesopause region as simulated by the Canadian Middle Atmosphere Model, Journal of Atmospheric and Solar-Terrestrial Physics, 74, 111–123, doi:10.1016/j.jastp.2011.10.010, http://www.sciencedirect.com/science/article/pii/S1364682611002847, 2012.

- Lubis, S. W., Omrani, N.-E., Matthes, K., and Wahl, S.: Impact of the Antarctic Ozone Hole on the Vertical Coupling of the Stratosphere–Mesosphere–Lower Thermosphere System, Journal of the Atmospheric Sciences, 73, 2509–2528, doi:10.1175/JAS-D-15-0189.1, http://journals.ametsoc.org/doi/10.1175/JAS-D-15-0189.1, 2016.
- 25 Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, Journal of Geophysical Research: Atmospheres, 102, 16663–16682, doi:10.1029/97JD00237, http://onlinelibrary.wiley.com/doi/10.1029/97JD00237/abstract, 1997.
 - Pincus, R. and Stevens, B.: Paths to accuracy for radiation parameterizations in atmospheric models, Journal of Advances in Modeling Earth Systems, 5, 225–233, doi:10.1002/jame.20027, http://onlinelibrary.wiley.com.e-bis.mpimet.mpg.de/doi/10.1002/jame.20027/abstract, 2013.
- 30
 - Randall, C. E., Harvey, V. L., Singleton, C. S., Bernath, P. F., Boone, C. D., and Kozyra, J. U.: Enhanced NOx in 2006 linked to strong upper stratospheric Arctic vortex, Geophysical Research Letters, 33, L18 811, doi:10.1029/2006GL027160, 2006.
- Randall, C. E., Harvey, V. L., Singleton, C. S., Bailey, S. M., Bernath, P. F., Codrescu, M., Nakajima, H., and Russell, J. M.: Energetic particle precipitation effects on the Southern Hemisphere stratosphere in 1992–2005, Journal of Geophysical Research: Atmospheres, 112, D08 308, doi:10.1029/2006JD007696, 2007.
- Randel, W. J. and Wu, F.: Cooling of the Arctic and Antarctic Polar Stratospheres due to Ozone Depletion, Journal of Climate, 12, 1467–1479, doi:10.1175/1520-0442(1999)012<1467:COTAAA>20.CO;2, 1999.





15

Reick, C. H., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, Journal of Advances in Modeling Earth Systems, 5, 459–482, doi:10.1002/jame.20022, http://onlinelibrary.wiley.com/doi/10.1002/jame.20022/abstract, 2013.

Rozanov, E., Callis, L., Schlesinger, M., Yang, F., Andronova, N., and Zubov, V.: Atmospheric response to NOy source due to energetic
 electron precipitation, Geophysical Research Letters, 32, doi:10.1029/2005GL023041, 2005.

Rozanov, E., Calisto, M., Egorova, T., Peter, T., and Schmutz, W.: Influence of the Precipitating Energetic Particles on Atmospheric Chemistry and Climate, Surveys in Geophysics, 33, 483–501, doi:10.1007/s10712-012-9192-0, http://link.springer.com/article/10.1007/s10712-012-9192-0, 2012.

Schmidt, H., Brasseur, G. P., Charron, M., Manzini, E., Giorgetta, M. A., Diehl, T., Fomichev, V. I., Kinnison, D., Marsh, D., and Walters,

10 S.: The HAMMONIA Chemistry Climate Model: Sensitivity of the Mesopause Region to the 11-Year Solar Cycle and CO₂ Doubling, Journal of Climate, 19, 3903–3931, doi:10.1175/JCLI3829.1, http://journals.ametsoc.org/doi/abs/10.1175/JCLI3829.1, 2006.

Schmidt, H., Rast, S., Bunzel, F., Esch, M., Giorgetta, M., Kinne, S., Krismer, T., Stenchikov, G., Timmreck, C., Tomassini, L., and Walz, M.: Response of the middle atmosphere to anthropogenic and natural forcings in the CMIP5 simulations with the Max Planck Institute Earth system model, Journal of Advances in Modeling Earth Systems, 5, 98–116, doi:10.1002/jame.20014, http://onlinelibrary.wiley.com/doi/10.1002/jame.20014/abstract, 2013.

- Schoeberl, M. R. and Strobel, D. F.: The Response of the Zonally Averaged Circulation to Stratospheric Ozone Reductions, Journal of the Atmospheric Sciences, 35, 1751–1757, doi:10.1175/1520-0469(1978)035<1751:TROTZA>2.0.CO;2, http://journals.ametsoc.org/doi/abs/10.1175/1520-0469%281978%29035%3C1751%3ATROTZA%3E2.0.CO%3B2, 1978.
- Seppälä, A., Randall, C. E., Clilverd, M. A., Rozanov, E., and Rodger, C. J.: Geomagnetic activity and polar sur face air temperature variability, Journal of Geophysical Research: Space Physics, 114, doi:10.1029/2008JA014029, http://onlinelibrary.wiley.com/doi/10.1029/2008JA014029/abstract, 2009.

Shine, K. P.: On the modelled thermal response of the Antarctic stratosphere to a depletion of ozone, Geophysical Research Letters, 13, 1331–1334, doi:10.1029/GL013i012p01331, 1986.

Sinnhuber, M., Funke, B., von Clarmann, T., Lopez-Puertas, M., Stiller, G. P., and Seppälä, A.: Variability of NOx in the polar middle

25 atmosphere from October 2003 to March 2004: vertical transport vs. local production by energetic particles, Atmos. Chem. Phys., 14, 7681–7692, doi:10.5194/acp-14-7681-2014, 2014.

Solomon, S., Rusch, D. W., Gérard, J. C., Reid, G. C., and Crutzen, P. J.: The effect of particle precipitation events on the neutral and ion chemistry of the middle atmosphere: II. Odd hydrogen, Planetary and Space Science, 29, 885–893, doi:10.1016/0032-0633(81)90078-7, 1981.

30 Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, Journal of Advances in Modeling Earth Systems, 5, 146–172, doi:10.1002/jame.20015, http://onlinelibrary.wiley.com/doi/10.1002/jame.20015/abstract, 2013.

Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, Bulletin of the American Meteorological
 Society, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, http://journals.ametsoc.org/doi/abs/10.1175/BAMS-D-11-00094.1, 2012.