

Author Response:

We thank the referees and editor for their time and effort in reviewing this manuscript. We have made the changes as requested. In the time between submission of the revised manuscript and the current review, we found an error in the way we treated the particle size dependence of molecular transport to the particle surface (the beta term in Eq. 1). The impact of error is small and does not affect the conclusions of our work. It amounts to ~10% reduction of the growth rate for the largest particle diameter examined, and very little impact on the size-dependent molecular composition. Nonetheless, we are compelled to disclose the problem and rectify it in the manuscript. For this reason, we made additional changes to the text in the manuscript and supporting information, and we updated all figures even though the change was often imperceptible. As an aid to the editor/reviewers, we have appended redline documents to this file that highlight the changes that were made. Please contact us if you have questions or require additional information.

Murray Johnston
Michael Apsokardu

Nanoparticle Growth by Particle Phase Chemistry

Michael J. Apsokardu and Murray V. Johnston

Department of Chemistry and Biochemistry, University of Delaware, Newark, DE, 19716, USA

Correspondence to: Murray V. Johnston (mvj@udel.edu)

5 Abstract

The ability of particle phase chemistry to alter the molecular composition and enhance the growth rate of nanoparticles in the 2-100 nm diameter range is investigated through the use of a growth model. The molecular components included are sulfuric acid, ammonia, water, a non-volatile organic compound, and a semi-volatile organic compound. Molecular composition and growth rate are compared for particles that grow by partitioning alone vs. those that grow by a combination of partitioning and an accretion reaction in the particle phase between two organic molecules. Particle phase chemistry causes a change in molecular composition that is particle diameter dependent, and when the reaction involves semi-volatile molecules, the particles grow faster than by partitioning alone. These effects are most pronounced for particles larger than about 20 nm in diameter. The modeling results provide a fundamental basis for understanding recent experimental measurements of the molecular composition of secondary organic aerosol showing that accretion reaction product formation increases linearly with increasing aerosol volume-to-surface area. They also allow initial estimates of the reaction rate constants for these systems. For secondary aerosol produced by either OH oxidation of the cyclic dimethylsiloxane (D₃) or ozonolysis of β-pinene, oligomerization rate constants on the order of 10⁻³ to 10⁻¹ M⁻¹s⁻¹ are needed to explain the experimental results. These values are consistent with previously measured rate constants for reactions of hydroperoxides and/or peroxyacids in the condensed phase.

1 Introduction

Atmospheric aerosols influence Earth's energy balance either directly by scattering incoming solar radiation or indirectly through cloud formation (Charlson et al., 1992; Kerminen et al., 2012; Lohmann and Feichter, 2004). A significant fraction of airborne particles originate from gas-to-particle conversion during new particle formation (NPF) (Bzdek and Johnston, 2010). Volatile compounds emitted into the atmosphere from biogenic and anthropogenic sources are oxidized to produce semi-volatile and non-volatile products. Particle formation begins when these products, in combination with other gas phase species, come together to form clusters on the order of 1-2 nm in diameter that are able to spontaneously grow to much larger sizes (Bzdek and Johnston, 2010; Dusek et al., 2006; Kulmala et al., 2013; Zhang et al., 2012). Depending on chemical composition, once the particles have grown to a size of 50-100 nm in diameter, they are able to serve as cloud condensation nuclei (CCN). The probability that a newly formed nanoparticle will grow to a CCN relevant size depends on its growth rate relative to the loss rate from e.g. coagulation or scavenging by pre-existing aerosols (Kuang et al., 2010). Uncertainties in predicting the

35 conditions that favor CCN formation make it challenging to accurately predict future impacts of radiative forcing (Carslaw et al., 2013).

The three main chemical species that contribute to ambient nanoparticle growth are sulfuric acid, a neutralizing base typically ammonia, and organic matter. The growth rate due to sulfuric acid along with neutralizing base is accurately predicted by experimental measurements of gas phase mixing ratio and particle phase composition using a condensational growth model (Bzdek et al., 2013; Pennington et al., 2013; Smith et al., 2008; 40 Stolzenburg et al., 2005), though sulfuric acid represents only a minor fraction of the total growth rate of ambient particles (Kuang et al., 2010, 2012; Weber et al., 1996; Wehner et al., 2005). Nanoparticle composition and growth rate are dominated by organic matter (Bzdek et al., 2011, 2012, 2013, 2014a, 2014b; Pennington et al., 2013; Riipinen et al., 2012; Smith et al., 2008), and though significant molecular insight has been gained (Bianchi et al., 2016; Ehn et al., 2014; Kulmala et al., 2013; Riccobono et al., 2014), current growth models for organic matter 45 appear to be incomplete (Hallquist et al., 2009; Tröstl et al., 2016).

Gas phase oxidation of volatile organic compounds occurs through a complex set of reaction pathways in both the gas and particle phases to yield particle phase products that often number in the hundreds or thousands based on accurate mass measurements (Bateman et al., 2008; Mentel et al., 2015; Reinhardt et al., 2007; Tu et al., 2016). Absorptive partitioning (Barsanti et al., 2017; Pankow, 1994) of gas phase products to the particle phase 50 forms secondary organic aerosol (SOA), which includes both non-volatile (NVOC) and semi-volatile (SVOC) organic compounds (Kroll and Seinfeld, 2008; Riipinen et al., 2012). NVOC have a negligible evaporation rate after partitioning to the particle phase, and therefore cause particles to grow at a rate given by the condensation rate. SVOC have a significant evaporation rate from the particle phase, and therefore grow particles at a slower rate than the condensation rate. Recent advances in measurement technology have greatly improved our molecular 55 understanding of NVOC (Ehn et al., 2012, 2014; Jokinen et al., 2015). More broadly, the complex distribution of molecular products and their associated wide range of volatilities can be represented by a volatility basis set (VBS) distribution where products are binned according to their gas phase saturation concentrations (Donahue et al., 2006, 2011, 2012).

Molecular analysis of SOA has shown the presence of compounds that were produced by a reaction within 60 the particle phase. Some examples include oligomers in biogenic SOA formed by accretion reactions (Barsanti and Pankow, 2004; Kalberer et al., 2004; Tolocka et al., 2004a), imine related species formed by the reaction of dicarbonyls with ammonia or amines (Galloway et al., 2014; De Haan et al., 2011; Lee et al., 2013; Stangl and Johnston, 2017), and organosulfates (Riva et al., 2016; Surratt et al., 2007; Wong et al., 2015; Xu et al., 2015). Reactions such as these increase the aerosol yield by forming additional SOA beyond what would be expected from 65 partitioning alone, if they form non-volatile products from semi-volatile reactants in the particle phase (Lopez-Hilfiker et al., 2016; Shiraiwa et al., 2014). Experimental measurements have shown that oligomers can constitute up to about 50% of the mass of SOA produced from biogenic precursors in laboratory reactors, though it is not clear how much of the oligomeric matter is produced from semi-volatile vs. non-volatile precursors (Hall IV and Johnston, 2011).

70 The role of particle phase oligomerization in SOA formation has been the focus of several modeling
studies. Owing to the high molecular weight and corresponding low volatility of oligomer products (Shiraiwa et al.,
2014), early work assumed an irreversible process (Vesterinen et al., 2007), which proved effective for predicting
the yields of freshly formed aerosol in chamber experiments and estimating the magnitude of the oligomerization
rate constant needed for the process to impact yields. More recent models have included reversibility (Roldin et al.,
75 2014; Trump and Donahue, 2014), which is needed to reproduce perturbations of freshly formed SOA such as
changes induced by isothermal dilution, thermal degradation and/or aging. Regional air quality models show that
oligomerization has the potential to significantly increase the SOA mass concentration (Aksoyoglu et al., 2011;
Lemaire et al., 2016), and accurately representing this chemistry in these models is perhaps the greatest uncertainty
for predicting SOA formation (Shrivastava et al., 2016).

80 The influence of particle size, or more precisely the relative roles of particle volume and surface area, on
oligomer formation in SOA has received relatively little attention though these effects are implicit in all of the
models. Particle size has been discussed primarily in the context of accumulation mode particles greater than 100
nm in diameter (Roldin et al., 2014; Shiraiwa et al., 2013; Vesterinen et al., 2007). The Vesterinen study did report
aerosol yield functions from simulations starting with 20 nm diameter seed particles, and the results suggested that
85 particle phase chemistry could enhance growth rates under atmospherically relevant conditions.

In the work described here, we explore computationally the impact of particle phase chemistry on both the
growth rate and composition of nanoparticles (defined here as particles smaller than 100 nm in diameter). This work
was inspired by recent composition studies of laboratory-generated secondary aerosol spanning several tens of
nanometers in diameter, which showed systematic changes in composition with increasing particle size (Tu and
90 Johnston, 2017; Wu and Johnston, 2017). In the first set of calculations, chemical composition and growth rate are
determined for particles between 2 and 100 nm assuming that growth occurs by partitioning alone. These results are
compared to a second set of calculations where semi-volatile molecules in the particle phase are allowed to undergo
an accretion reaction to produce a non-volatile product. Together, the two sets of calculations give insight into the
reaction conditions and particle size range where particle phase chemistry is most likely to alter nanoparticle
95 composition and accelerate the growth rate. The modeling approach is similar to that used in previous studies,
though the reaction conditions used here more closely resemble those of an “open” laboratory reactor where aerosol
flows into and out of the reactor (i.e. flow reactor), as opposed to a “closed” batch reactor which was the main focus
of previous modeling work. The results are discussed in the context of recent size-resolved molecular composition
measurements (performed with flow reactors) and the potential atmospheric impact of particle phase chemistry.

100

2 Model Description

The growth model used in this work includes sulfuric acid, ammonia, and ~~organic matter~~ since these are the
major components found in ambient nanoparticles during NPF (Bzdek et al., 2011, 2014a; Pennington et al., 2013;
Smith et al., 2008; Stolzenburg et al., 2005). Water is also included as predicted by the Extended Aerosol Inorganics

Owner 12/28/2017 11:55 AM

Deleted: st

Owner 12/28/2017 11:55 AM

Deleted: di

Owner 12/28/2017 11:55 AM

Deleted: SOA

Model (E-AIM) (Wexler and Clegg, 2002), and then corrected to account for the effects of particle surface curvature (Kreidenweis et al., 2005; Yli-Juuti et al., 2013). For simplicity, the gas phase precursors to SOA formation are represented by two specific organic compounds, one non-volatile (NVOC) and the other semi-volatile (SVOC). Particle phase chemistry involves a generic accretion reaction between two SVOC (or NVOC) molecules to give a dimer. Table 1 gives relevant molecular parameters and gas phase mixing ratios. The gas phase mixing ratios are assumed to be constant over time (steady state) as might be achieved in a flow reactor, and the values chosen are typical of what might be observed during NPF in a boreal forest (Vestenius et al., 2014). Molecular volatility is expressed in terms of the saturation concentration, C^* , which is the equilibrium vapor pressure in units of $\mu\text{g}/\text{m}^3$ (Donahue et al., 2006). Diffusion coefficients are obtained from the procedure described in Supporting Information (Poling et al., 2001).

Calculations begin with a 2 nm diameter particle consisting of sulfuric acid and ammonia in a 2:1 ratio of base to acid. From there, gas phase species partition to the particle phase based on their volatilities and gas phase mixing ratios, causing the particle to grow. The growth calculation is iterative. Starting from an initial particle of a given composition and volume ($V_{p,n}$), the increase in particle volume (and corresponding change in composition) is determined over a short time period for each molecular species separately, and the increases for the individual components are summed to give the total volume of the particle at the end of the time period ($V_{p,n+1}$) along with its composition. The end point for the first time period is used as the start point for the second time period, and the process repeats. A schematic of calculation of work flow is given in Supporting Information Figure S1 along with additional aspects of the approach.

2.1 Partitioning

The extent to which a compound formed in the gas phase partitions to the particle phase is determined by its saturation ratio (S_d), which is the ratio of the gas phase mixing ratio to the saturation mixing ratio. The subscript “ d ” refers to the particle diameter and acknowledges that the saturation ratio depends in part on the radius of curvature of the particle surface. Accordingly, the first step in the calculation is to determine the Kelvin effect modified vapor pressure ($KEMP_d$) for all species based on the initial particle diameter. For the conditions used in this study, sulfuric acid and NVOC have $S_d \gg 1$ for all particle diameters investigated, and as discussed previously, grow particles at their condensation rates. Ammonia, water, and SVOC have $S_d \ll 1$ and grow particles at rates slower than their condensation rates.

The collision rate for species i determines the number of condensing molecules that can be taken into the particle during time period dt , thereby incrementing the particle volume by dV_i :

$$(1) \quad \frac{dV_i}{dt} = \frac{c_i}{4} \gamma \pi d^2 C_{i,g} \beta_d V_{M,i}$$

where c_i is the mean thermal velocity, γ is the uptake coefficient, d is the particle diameter, $C_{i,g}$ is the gas phase mixing ratio of species i , β_d is the Loyalka correction factor for mass transport to a spherical particle with diameter d , and $V_{M,i}$ is the molar volume of species i . The equation for β_d is given in Supporting Information. Values of dV_i

Owner 12/28/2017 11:55 AM

Deleted: $D^2 C_{i,g} \beta_D$

Owner 12/28/2017 11:55 AM

Deleted: Fuchs-Sutugin

for condensational growth by sulfuric acid and NVOC are calculated from Eq. 1, with the inherent assumption that the surface accommodation coefficient is 1.

Equation 1 can also be used to calculate the uptake of semi-volatile species provided that the total amount taken into the particle does not exceed the equilibrium end point. Ammonia uptake is determined from the number of sulfuric acid molecules that have condensed. For a given time period dt , once the number of ammonia molecules that have been taken into the particle equals twice the number of sulfuric acid molecules that were taken during the same time period, no further ammonia uptake occurs. While dissolution of ammonia into water is possible, the amount is negligible in comparison to ammonia uptake associated with condensation of sulfuric acid (Clegg et al., 1998; Wexler and Clegg, 2002). Under the conditions used in this study, the condensation rate of ammonia is about 25 times greater than that of sulfuric acid, so the model assumes that stoichiometric uptake of ammonia is instantaneous with sulfuric acid. Water uptake is determined from E-AIM based on the combined amounts of sulfuric acid, ammonia, and NVOC that have been added during time period dt . Since the gas phase mixing ratio of water is very high, equilibrium is assumed to be achieved instantaneously during the time period.

Individual volume increments dV_i for sulfuric acid, NVOC, ammonia, and water are summed to give the total volume increment. Because the volume of particle phase has increased, SVOC is no longer in equilibrium between the gas and particle phases, and a net migration of SVOC from the gas phase to the particle phase must occur. SVOC molecules are taken into the particle at a rate described by Eq. 1 to re-establish equilibrium. The equilibrium point (Pankow, 1994) expressed as the volume ratio of SVOC in the particle phase ($V_i/V_{p,n}$) is:

$$(2) \quad \frac{V_i}{V_{p,n}} = \left(\frac{C_{i,g}}{KEMP_d} \right) \left(\frac{V_{M,i}}{V_{M,p}} \right) \frac{1}{\zeta_i}$$

where i in this case refers to SVOC, ζ_i is the activity coefficient for SVOC in the particle phase (assumed to be 1 in this study), and $V_{M,i}$ and $V_{M,p}$ are the respective the molar volumes of SVOC and the particle phase. The incremental increase of SVOC over time period dt is determined by evaluating Eq. 2 before and after the increases due to sulfuric acid, NVOC, ammonia, and water. For the conditions used in this study, the condensation rate of gas phase SVOC is generally greater than the uptake rate needed to maintain equilibrium when particle growth is restricted to partitioning. The ratio of the two rates is dependent on the volume to surface area ratio of the particle, ranging from several orders of magnitude for a 2 nm diameter particle to a factor of 200 for a 100 nm diameter particle. Because the condensation rate is generally much greater than the uptake rate needed for partitioning, no evaporation rate of particle phase SVOC was included in the current study. The one situation where condensation is not sufficient for partitioning occurs when particle phase chemistry is **included** and will be discussed later.

2.2 Particle Phase Chemistry

Particle phase chemistry is modeled as an irreversible accretion reaction where two monomers come together to produce a non-volatile DIMER. While all combinations of SVOC and NVOC molecules were considered, most calculations involved the reaction between two SVOC molecules:

Owner 12/28/2017 11:55 AM

Deleted: s

Owner 12/28/2017 11:55 AM

Deleted: , and the particle diameter is small enough that gas phase diffusion does not limit the condensation rate.

Owner 12/28/2017 11:55 AM

Deleted: considered

$$(3) \quad \frac{d[DIMER]}{dt} = - \frac{d[SVOC]}{dt} = k_{DIMER}[SVOC]^2$$

where k_{DIMER} is the second order rate constant, and $[SVOC]$ is the particle phase concentration established by partitioning between the gas and particle phases. Since the reaction consumes particle phase SVOC molecules, additional SVOC must be taken from the gas phase into the particle phase to re-establish equilibrium. The rate at which additional SVOC molecules are taken into the particle is determined by the rate of particle phase reaction. When particle phase reaction is included in the growth calculation, the change in particle volume from $V_{p,n}$ to $V_{p,n+1}$ with respect to SVOC is the sum of the volume increase from unreacted SVOC molecules needed to re-establish partitioning equilibrium, and the volume increase from additional SVOC uptake to form dimers. SVOC continues to partition to the particle phase at each iteration, resulting in a continuous supply of reactant molecules for dimerization.

Dimerization rate constants in the range of 10^{-3} to $10^{-1} \text{ M}^{-1} \text{ s}^{-1}$ were studied, with most calculations at $10^{-2} \text{ M}^{-1} \text{ s}^{-1}$, which is the rate constant reported for dimerization of glyoxal in a bulk aqueous solution (Ervens and Volkamer, 2010). Ziemann and Atkinson (Ziemann and Atkinson, 2012) have reviewed thermodynamic and kinetic data for several types of reactions relevant to biogenic SOA. The reaction of a hydroperoxide with a carbonyl to give a peroxyhemiacetal, and the reaction of a peroxyacid with a carbonyl to form an acyl peroxyhemiacetal, both have reported rate constants in the 10^{-4} to $10^{-2} \text{ M}^{-1} \text{ s}^{-1}$ range depending on reaction conditions, and are relevant to the modeling results presented here. Reactions such as aldol condensation of carbonyls and ester formation from an acid and alcohol are much slower and unlikely to be atmospherically relevant based on both kinetics (Casale et al., 2007) and thermodynamics (DePalma et al., 2013). The effects of dimer decomposition (reaction reversibility), particle phase diffusion coefficient, and phase separation are not considered in this work, though we note that all would have the effect of reducing the impact of particle phase chemistry on composition and growth rate.

3 Results and Discussion

3.1 Particle Growth by Partitioning

The first set of calculations includes particle growth by partitioning, but no particle phase reaction. Figure 1a shows the diameter growth rate and Figure 1b the dry mass fraction for each chemical species as a function of particle diameter. For sulfuric acid and NVOC, where partitioning to the particle phase occurs at the collision limit, the growth rates for these species are independent of particle diameter. A constant growth rate for non-volatile species has been noted previously (Weber et al., 1996) and can be understood by rewriting Eq. 1 in terms of the diameter growth rate:

$$(4) \quad \frac{dd}{dt} = \frac{c_i}{2} \gamma C_{i,g} \beta_d V_{M,i}$$

Since γ is independent of d for a surface-limited process such as condensation (Smith et al., 2003; Tolocka et al., 2004a), dd/dt also becomes independent of d as a first approximation, provided that the precursor gas phase mixing

Owner 12/28/2017 11:55 AM

Deleted: Initially, the diameter growth rates for all species increase quickly with increasing particle size.

Owner 12/28/2017 11:55 AM

Deleted: increase is most pronounced between 2 and 10 nm, and this size dependence arises directly from the Fuchs-Sutugin term that limits mass transport to small diameter particles. Above about 10 nm

Owner 12/28/2017 11:55 AM

Deleted: become

Owner 12/28/2017 11:55 AM

Deleted: Above about 10 nm, β_D becomes independent of D .

ratio ($C_{i,g}$) does not change. In practice, dd/dt decreases slightly with increasing particle diameter owing to the effect of molecular diffusion on transport of vapor molecules to the particle surface (β_d becomes less than 1). Ammonia uptake is driven by sulfuric acid uptake, and therefore follows the same particle diameter dependence.

230 Water and SVOC show greater increases in their growth rates with increasing particle diameter than sulfuric acid and NVOC because they have $S_{d,e} \ll 1$. In principle, semi-volatile species are subject to two particle size-dependent effects: β_d in Eq. 1 and $KEMP_d$ in Eq. 2. For the conditions studied here, the β_d term has a negligible effect on the uptake rates of water and SVOC, since uptake is determined by the equilibrium endpoint rather than the condensation rate. Instead, the difference between the semi-volatile species and non-volatile species in Figure 1a are driven by the particle size dependence of $KEMP_d$, which alters the equilibrium point for absorptive partitioning.

235 In Figure 1b, the dry particle mass fractions of sulfuric acid, NVOC, and ammonia show very little change with increasing d since their relative growth rates are independent of d . Small changes just above 2 nm are due to the choice of starting composition of the 2 nm diameter particle. In contrast, the SVOC mass fraction increases quickly with increasing particle size owing to the dependence of $KEMP_d$ on d . Figure 2 shows the mass fraction ratio of SVOC to NVOC as a function of particle size. The ratio increases quickly at the smallest particle sizes and
240 then more slowly thereafter. This plot is consistent with experimental measurements of molecular composition across a similar range of particle diameters, which show that lower volatility species are preferentially detected in smaller particles (Winkler et al., 2012; Zhao et al., 2013).

3.2 Particle Growth by a Combination of Partitioning and Particle Phase Chemistry

245 When particle phase reaction is included in the growth calculation, both the growth rate and chemical composition change. Figures 3a and b show diameter growth rates and dry mass fractions, respectively, when particle phase reaction is included. Comparing Figure 3a to Figure 1a shows that particle phase chemistry causes the diameter growth rate to continue increasing with increasing particle size above 10 nm. This diameter dependence is different from the growth rate due to partitioning alone where dd/dt becomes independent of particle size above 10 nm. The size dependence of particle growth at larger particle sizes can be understood by expressing the uptake coefficient in terms of particle diameter:
250

$$(5) \quad \gamma = k'd$$

In Eq. 5, the uptake coefficient is proportional to d , which is characteristic of a volume-limited process (Saul et al., 2006; Tolocka et al., 2004b). Inserting Eq. 5 into Eq. 4, shows that dd/dt also increases linearly with d . The number of SVOC molecules that react to form DIMER is proportional to the total volume of the particle, and therefore the
255 SVOC uptake rate needed to maintain partitioning equilibrium is also volume (and hence diameter) dependent. Note that the SVOC and DIMER diameter growth rates are both proportional to d in Figure 3a. Growth by condensation of NVOC and sulfuric acid remain surface-limited, and therefore, these diameter growth rates are unaffected by dimer formation.

Owner 12/28/2017 11:55 AM

Deleted: <

260 Figure 3b shows the dry mass fractions as a function of particle diameter when particle phase chemistry is
included. Below about 10 nm, the plots in Figure 3b are identical to those in Figure 1b where partitioning alone is
considered. Particle phase chemistry has minimal impact on the growth and composition of small particles for two
reasons. First, the particle volume to surface area ratio is very small, which favors surface-limited processes
(condensational growth by NVOC and sulfuric acid) over volume-limited processes (accretion reaction). Second,
265 the dependence of $KEMP_d$ on d causes the equilibrium concentration of SVOC in the particle phase to be very low
for small particles, which decreases the reaction rate (Eq. 3). As a result, particle phase chemistry has little impact
on the diameter growth rate or molecular composition below about 10 nm. Above 10 nm, DIMER mass starts to
accumulate in the particle, causing the mass fractions of NVOC, sulfuric acid, and ammonia decrease.

Figure 4a shows the mass fraction ratio of SVOC to NVOC as a function of particle diameter. Below 10
270 nm, where particle phase chemistry has minimal impact, the ratio in Figure 4a for particle phase reaction is
equivalent to that in Figure 2 for partitioning alone. The plots in Figures 2 and 4a diverge above about 10 nm
because the DIMER mass fraction increases, causing the NVOC mass fraction to decrease.

Figure 4b shows the mass fraction ratio of DIMER to NVOC. Below about 10 nm, hardly any DIMER is
produced. Above 10 nm, the DIMER to NVOC ratio increases approximately linearly with increasing d as expected
275 for a volume-limited process relative to a surface-limited process. Figures 3 and 4 show, for the conditions studied,
that particle size dependencies of the growth rate and composition above 10 nm are driven mainly by particle phase
chemistry rather than partitioning alone. Figure 4b illustrates how the impact of an accretion reaction might be
observed experimentally through organic molecular composition measurements. If a systematic increase in the
concentration of non-volatile dimers and higher order oligomers is observed with increasing particle diameter, then
280 particle phase chemistry is likely to have contributed to the formation of these molecules.

3.3 Factors that Influence Growth by Particle Phase Chemistry

Additional calculations were performed to investigate the roles of SVOC gas phase mixing ratio and
particle phase reaction rate constant on growth rate and composition. In Figure S2, diameter growth rates are shown
for SVOC gas phase mixing ratios between 0.35 to 1.4 pptv and compared to the partitioning calculation in Figure
285 1a (the “1.4 - NR” plot for no reaction in Figure S2). Above about 10 nm, the growth rates increase linearly with
increasing d , and the slope of the increase scales by approximately $[SVOC]^2$ as expected by Eq. 3. The actual
dependence is slightly less than $[SVOC]^2$ owing to the change in particle density that is associated with the changing
DIMER mass fraction.

Figure 5a shows the effect of reaction rate constant on diameter growth rate. The rate constants investigated
290 range over two orders of magnitude from 10^{-3} to $10^{-1} \text{ M}^{-1}\text{s}^{-1}$ while keeping the SVOC mixing ratio constant at 1.4
pptv. “NR” in Figure 5a represents no reaction and is equivalent to the plot in Figure 1a for growth by partitioning
alone. For a reaction rate constant of $10^{-3} \text{ M}^{-1}\text{s}^{-1}$, particle growth rate is only slightly larger than growth by
partitioning only. In this case, DIMER production is slow, so there is only minor enhancement of the overall growth
rate. Increasing the rate constant one order of magnitude to $10^{-2} \text{ M}^{-1}\text{s}^{-1}$, causes a substantial enhancement of the

295 growth rate above 10 nm. For a rate constant of $10^{-1} \text{ M}^{-1}\text{s}^{-1}$, a complex plot is observed. Volume-limited particle
growth occurs up to ~40 nm where the growth rate increases approximately linearly with d . Above ~40 nm, the
reaction rate becomes so fast that particle growth is limited by the condensation rate of SVOC, a surface-limited
process, as reflected by the particle diameter dependence. Since the calculation methodology used in our study did
not include an evaporation rate for SVOC, the transition from volume-limited to surface-limited kinetics is abrupt in
300 Figure 5a. In practice, one would expect a more gradual transition from one to the other. The lack of a particle size
dependence on SOA growth in the limit of a fast reaction rate has also been suggested in a modeling study of SOA
produced by α -pinene ozonolysis (Gatzsche et al., 2017). It is also consistent with the work of Vesterinen et al.
(Vesterinen et al., 2007) who showed a linear increase of particle diameter with time when the reaction rate constant
was sufficiently large.

305 Accretion chemistry is not necessarily restricted to SVOC molecules. If two NVOC molecules react in the
particle phase to form DIMER, then the diameter growth rate *will not* change with increasing particle diameter since
NVOC uptake remains unaffected, though the molecular composition *will* change with increasing diameter. A
calculation demonstrating this principle is shown in Figure S3 (growth rate vs. particle diameter) and Figure S4
(DIMER/ NVOC mass ratio vs. particle diameter). If one SVOC molecule reacts with one NVOC molecule to form
310 DIMER, both the growth rate and molecular composition change with increasing particle diameter. A calculation
demonstrating this principle is shown in Figure 5b, where growth rate vs. particle diameter are compared for SVOC-
SVOC, NVOC-NVOC and SVOC-NVOC dimer formation reactions. The SVOC-NVOC reaction enhances the
particle growth rate at a much smaller particle diameter than the SVOC-SVOC reaction. This enhancement arises
from $KEMP_d$ of SVOC, which favors formation of the SVOC-NVOC dimer over the SVOC-SVOC dimer at small
315 particle sizes. Because of the greater potential of the SVOC-SVOC reaction to transform semi-volatile matter into
non-volatile matter, its contribution to growth rate eventually overcomes that of the SVOC-NVOC reaction as the
particle size increases.

Taken together, Figures S2-S4 show that the observation of a change in DIMER to monomer ratio with
increasing particle diameter does not necessarily indicate an enhancement of the particle growth rate by accretion
320 chemistry. On the other hand, Figure 5b shows that the lack of a particle size-dependent change in the growth rate
does not necessarily mean that particle growth rate is unaffected by accretion chemistry. Nonetheless, these results
illustrate how experimental measurements of particle size-dependent changes in molecular composition and growth
rate can supplement traditional measures, such as the concentration dependence of the aerosol yield and
perturbations due to e.g. isothermal dilution, to better constrain SOA formation models with regard to particle phase
325 chemistry.

3.4 Comparison to Recent Experimental Measurements and Atmospheric Implications

For the modeling conditions studied here, particle phase chemistry influences both the molecular
composition and growth rate of sub-100 nm diameter particles under atmospherically relevant conditions, provided
that the rate constant is above about $10^{-3} \text{ M}^{-1}\text{s}^{-1}$ for semi-volatile reactants having a saturation concentration on the

Owner 12/28/2017 11:55 AM

Deleted: and

Owner 12/28/2017 11:55 AM

Deleted: growth rate and molecular composition become independent of D

Owner 12/28/2017 11:55 AM

Deleted: /

Owner 12/28/2017 11:55 AM

Deleted: is

335 order of $1 \mu\text{g}/\text{m}^3$ and gas phase mixing ratios on the low pptv level. Based on the condensed phase concentration
dependence of reactants on the reaction rate, a rate constant on the order of $10^{-1} \text{M}^{-1}\text{s}^{-1}$ would be required for semi-
volatile reactants having saturation concentrations on the order of $10 \mu\text{g}/\text{m}^3$. These rate constants (Ziemann and
Atkinson, 2012) and saturation concentrations (Trump and Donahue, 2014) are in the range of those expected for
e.g. monoterpene accretion chemistry. The overall conclusion that accretion chemistry may contribute to
340 nanoparticle growth and/or composition under atmospherically relevant conditions is similar to that reached by
(Vesterinen et al., 2007). The current work provides an additional, fundamental basis for interpreting recent
experimental measurements of molecular composition as a function of particle size in the case of monodisperse
aerosols (Tu and Johnston, 2017), or more generally, volume-to-surface area ratio for polydisperse aerosols (Wu and
Johnston, 2017). In each study, the relative concentration of accretion products increased approximately linearly
345 with particle size (or volume-to-surface area ratio) as predicted by Figure 4b. Below, we estimate the magnitudes of
the reaction rate constants needed to reproduce the experimental results.

In the Wu study (Wu and Johnston, 2017), decamethylcyclopentasiloxane (D_5) reacted with OH in the gas
phase to produce secondary aerosol having three types of products: ring-opened species, oxidized monomers
containing OH and/or CH_2OH functionalities in place of CH_3 , and dimers of D_5 and/or its monomer oxidation
350 products. The dimer concentration in the particle phase (as indicated by the signal intensities of corresponding ions
produced by electrospray ionization) increased linearly with increasing volume-to-surface area ratio of the aerosol
relative to the (non-volatile) ring-opened products. This dependence suggested that ring-opened products were
formed in the gas phase and subsequently condensed onto the particles, while dimers were produced directly in the
particle phase from monomers. Monomer gas phase mixing ratios in these experiments were on the order of a few
355 ppbv, which is similar in magnitude to the levels measured in a university lecture hall (Tang et al., 2015). Using
these values in conjunction with the basic model described above, we estimate the dimerization rate constant had to
be in the 10^{-3} to $10^{-1} \text{M}^{-1}\text{s}^{-1}$ range to achieve the level of dimerization observed, given that most of the monomers
expected to participate in the reaction had estimated saturation concentrations in the 10^2 to $10^3 \mu\text{g}/\text{m}^3$ range. Smaller
rate constants would have been unable to produce a sufficient amount of dimers, while higher rate constants would
360 not have shown a volume-to-surface area ratio dependence of the dimer concentration since the reaction rate would
have been limited by transport of reacting monomer to the particle surface.

In the Tu study (Tu and Johnston, 2017), β -pinene ozonolysis produced SOA having systematic changes in
molecular composition as a function of particle size. The relative concentrations (as indicated by the signal
intensities of ions produced by electrospray ionization) of higher order oligomers, i.e. trimers and tetramers,
365 increased linearly with increasing particle size, similar to Figure 4b. The total oligomer signal intensity was
comparable to our previous study of α -pinene SOA formed under similar reaction conditions, where the oligomer
content was experimentally determined to be about 50% of the total SOA mass (Hall IV and Johnston, 2011). For
the Tu study, we estimate the NVOC mixing ratio to be approximately 2 ppbv based on the yield estimate of Ehn et
al. (Ehn et al., 2014) for β -pinene ozonolysis. We estimate the mixing ratios for the $10 \mu\text{g}/\text{m}^3$ and $100 \mu\text{g}/\text{m}^3$
370 saturation concentration bins to be approximately 30 ppbv and 40 ppbv, respectively, based on the volatility basis set

(VBS) parameterization of Donohue and coworkers for α -pinene ozonolysis (Donahue et al., 2012; Trump and Donahue, 2014). For these conditions, condensational growth and/or partitioning of molecular species alone are able to explain the overall growth of particles to about 80-100 nm in diameter during the ~20 s residence time of the flow reactor, though the modeling results are extremely sensitive to the mixing ratios and VBS parameterization used. However, condensational growth and partitioning cannot explain the high abundance of accretion products that were measured. We estimate that the reaction rate constants needed to form dimers and higher order oligomers in sufficient amounts had to be at least on the order of $10^{-3} \text{ M}^{-1}\text{s}^{-1}$. (The required rate constant is dependent on SVOC volatility, with a larger rate constant needed for saturation concentrations above $10 \mu\text{g}/\text{m}^3$.) Together, the modeling work performed here and the experimental measurements in the Wu and Tu studies all suggest that reaction rate constants on the order of 10^{-3} to $10^{-1} \text{ M}^{-1}\text{s}^{-1}$ are needed for accretion chemistry to be relevant to SOA formation and growth.

While systematic changes in molecular composition with increasing particle size and/or aerosol volume-to-surface area ratio have been established experimentally, an enhancement of the particle growth rate in the sub-100 nm diameter range is less clear. Recent measurements by Kourtchev et al. (Kourtchev et al., 2016) of accretion oligomers in ambient aerosol from a boreal forest suggest that oligomer formation increases with increasing SOA mass concentration, an observation that is consistent with the molecular composition results reported here. More importantly, the authors noted that aerosols enriched with oligomers were strongly correlated with higher CCN activity, and they suggested that this correlation could indicate that oligomers may speed up particle growth. The modeling results presented here suggest that particle phase reactions are indeed capable of enhancing growth rates in a size range relevant to CCN activity.

The work presented here suggests that experimental measurements of molecular composition as a function of particle size and/or aerosol volume-to-surface area ratio can supplement traditional measurements (concentration-dependent aerosol mass yields, perturbations caused by isothermal dilution, etc.) for constraining aerosol formation models. Future modeling of size-dependent molecular composition should include physico-chemical parameters such as diffusion, phase separation, and reaction reversibility (Liu et al., 2014; Mai et al., 2015; Riipinen et al., 2012; Shiraiwa et al., 2012; Song et al., 2015; Trump and Donahue, 2014; Zaveri et al., 2014) that may influence the contribution of particle phase chemistry to nanoparticle growth.

Supporting Information

Four figures (S1-S4) plus additional description of the particle growth model.

Acknowledgements

This research was supported by the United States National Science Foundation under grant numbers CHE-1408455 and AGS-1649719. The authors thank Pauli Paasonen for helpful discussions on molecular transport dynamics.

Owner 12/28/2017 11:55 AM

Deleted: molecular composition

Owner 12/28/2017 11:55 AM

Deleted: ,

Owner 12/28/2017 11:55 AM

Deleted: , etc.

References

- Aksoyoglu, S., Keller, J., Barmadimos, I., Oderbolz, D., Lanz, V. A., Prévôt, A. S. H. and Baltensperger, U.:
410 Aerosol modelling in Europe with a focus on Switzerland during summer and winter episodes, *Atmos. Chem. Phys.*,
11(14), 7355–7373, doi:10.5194/acp-11-7355-2011, 2011.
- Barsanti, K. C. and Pankow, J. F.: Thermodynamics of the formation of atmospheric organic particulate matter by
accretion reactions-Part 1: aldehydes and ketones, *Atmos. Environ.*, 38, 4371–4382,
doi:10.1016/j.atmosenv.2006.03.013, 2004.
- 415 Barsanti, K. C., Kroll, J. H. and Thornton, J. A.: The Formation of Low-Volatility Organic Compounds in the
Atmosphere: Recent Advancements and Insights, *J. Phys. Chem. Lett.*, DOI: 10.1021/acs.jpcclett.6b02969,
doi:10.1021/acs.jpcclett.6b02969, 2017.
- Bateman, A. P., Walser, M. L., Desyaterik, Y., Laskin, J., Laskin, A., Nizkorodov, S. A. and Bateman, A. P.: The
Effect of Solvent on the Analysis of Secondary Organic Aerosol Using Electrospray Ionization Mass Spectrometry,
420 *Environ. Sci. Technol.*, 42(19), 7341–7346, doi:10.1021/es801226w, 2008.
- Bianchi, F., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki,
N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Manninen, H. E., Rondo, L.,
Williamson, C., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J. and Baltensperger, U.: New particle
formation in the free troposphere: A question of chemistry and timing, *Science*, 352(6289), 1109–1112,
425 doi:10.1126/science.aad5456, 2016.
- Bzdek, B. R. and Johnston, M. V.: New Particle Formation and Growth in the Troposphere, *Anal. Chem.*, 82(19),
7871–7878, doi:10.1021/ac100856j, 2010.
- Bzdek, B. R., Zordan, C. A., Luther, G. W. and Johnston, M. V.: Nanoparticle Chemical Composition During New
Particle Formation, *Aerosol Sci. Technol.*, 45(8), 1041–1048, doi:10.1080/02786826.2011.580392, 2011.
- 430 Bzdek, B. R., Zordan, C. A., Pennington, M. R., Luther III, G. W. and Johnston, M. V.: Quantitative Assessment of
the Sulfuric Acid Contribution to New Particle Growth, *Environ. Sci. Technol.*, 46(8), 4365–4373,
doi:10.1021/es204556c, 2012.
- Bzdek, B. R., Horan, A. J., Pennington, M. R., DePalma, J. W., Zhao, J., Jen, C. N., Hanson, D. R., Smith, J. N.,
McMurry, P. H. and Johnston, M. V.: Quantitative and time-resolved nanoparticle composition measurements during
435 new particle formation, *Faraday Discuss.*, 165, 25–43, doi:10.1039/c3fd00039g, 2013.
- Bzdek, B. R., Lawler, M. J., Horan, A. J., Pennington, M. R., DePalma, J. W., Zhao, J., Smith, J. N. and Johnston,
M. V.: Molecular constraints on particle growth during new particle formation, *Geophys. Res. Lett.*, 6045–6054,
doi:10.1002/2014GL060160, 2014a.
- Bzdek, B. R., Horan, A. J., Pennington, M. R., Janecek, N. J., Baek, J., Stanier, C. O. and Johnston, M. V.: Silicon

- 440 is a frequent component of atmospheric nanoparticles, *Environ. Sci. Technol.*, 48(19), 11137–11145, doi:10.1021/es5026933, 2014b.
- Carslaw, K. S., Lee, L. a, Reddington, C. L., Mann, G. W. and Pringle, K. J.: The magnitude and sources of uncertainty in global aerosol, *Faraday Discuss.*, 165, 495–512, doi:10.1039/c3fd00043e, 2013.
- Casale, M. T., Richman, A. R., Elrod, M. J., Garland, R. M., Beaver, M. R. and Tolbert, M. A.: Kinetics of acid-catalyzed aldol condensation reactions of aliphatic aldehydes, *Atmos. Environ.*, 41(29), 6212–6224, doi:10.1016/j.atmosenv.2007.04.002, 2007.
- 445 Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Hansen, J. E. and Hofmann, D. J.: Climate Forcing by Anthropogenic Aerosols, *Science*, 255, 423–430, 1992.
- Clegg, S. L., Brimblecombe, P. and Wexler, A. S.: Thermodynamic Model of the System H^+ - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O at 298 . 15 K, *J. Phys. Chem.*, 5639(3), 2155–2171, 1998.
- 450 DePalma, J. W., Horan, A. J., Hall IV, W. A. and Johnston, M. V.: Thermodynamics of oligomer formation: implications for secondary organic aerosol formation and reactivity, *Phys. Chem. Chem. Phys.*, 15(18), 6935, doi:10.1039/c3cp44586k, 2013.
- Donahue, N. M., Robinson, A. L., Stanier, C. O. and Pandis, S. N.: Coupled Partitioning, Dilution, and Chemical Aging of Semivolatile Organics, *Environ. Sci. Technol.*, 40(8), 2635–2643, doi:10.1021/es052297c, 2006.
- 455 Donahue, N. M., Epstein, S. A., Pandis, S. N. and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics, *Atmos. Chem. Phys.*, 11(7), 3303–3318, doi:10.5194/acp-11-3303-2011, 2011.
- Donahue, N. M., Kroll, J. H., Pandis, S. N. and Robinson, A. L.: A two-dimensional volatility basis set-Part 2: Diagnostics of organic-aerosol evolution, *Atmos. Chem. Phys.*, 12(2), 615–634, doi:10.5194/acp-12-615-2012, 2012.
- 460 Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S. and Andreae, M. O.: Size Matters More Than Chemistry Aerosol Particles, *Science*, 312(5778), 1375–1378, doi:10.1126/science.1125261, 2006.
- 465 Ehn, M., Kleist, E., Junninen, H., Petäjä, T., Lönn, G., Schobesberger, S., Dal Maso, M., Trimborn, A., Kulmala, M., Worsnop, D. R., Wahner, A., Wildt, J. and Mentel, T. F.: Gas phase formation of extremely oxidized pinene reaction products in chamber and ambient air, *Atmos. Chem. Phys.*, 12(11), 5113–5127, doi:10.5194/acp-12-5113-2012, 2012.
- 470 Ehn, M., Thornton, J. A., Kleist, E., Sipila, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I. H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurten, T., Nielsen, L. B., Jorgensen, S., Kjaergaard, H. G., Canagaratna, M., Dal

- Maso, M., Berndt, T., Petaja, T., Wahner, A., Kerminen, V. M., Kulmala, M., Worsnop, D. R., Wildt, J. and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506(7489), 476–479, doi:10.1038/Nature13032, 2014.
- 475 Ervens, B. and Volkamer, R.: Glyoxal processing by aerosol multiphase chemistry: Towards a kinetic modeling framework of secondary organic aerosol formation in aqueous particles, *Atmos. Chem. Phys.*, 10(17), 8219–8244, doi:10.5194/acp-10-8219-2010, 2010.
- Galloway, M. M., Powelson, M. H., Sedehi, N., Wood, S. E., Millage, K. D., Kononenko, J. A., Rynaski, A. D. and De Haan, D. O.: Secondary organic aerosol formation during evaporation of droplets containing atmospheric aldehydes, amines, and ammonium sulfate, *Environ. Sci. Technol.*, 48(24), 14417–14425, doi:10.1021/es5044479, 2014.
- 480 Gatzsche, K., Iinuma, Y., Tilgner, A., Mutzel, A., Berndt, T. and Wolke, R.: Modeling studies of SOA formation from α -pinene ozonolysis, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2017-275, 2017.
- De Haan, D. O., Hawkins, L. N., Kononenko, J. A., Turley, J. J. and Jimenez, J. L.: Formation of Nitrogen-Containing Oligomers by Methylglyoxal and Amines in Simulated Evaporating Cloud Droplets, *Environ. Sci. Technol.*, 45(3), 984–991, doi:10.1021/es102933x, 2011.
- 485 Hall IV, W. A. and Johnston, M. V: Oligomer Content of α -Pinene Secondary Organic Aerosol, *Aerosol Sci. Technol.*, 45, 37–45, doi:10.1080/02786826.2010.517580, 2011.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M. and Dommen, J.: and Physics 490 The formation, properties and impact of secondary organic aerosol: current and emerging issues, (November 2008), 5155–5236, 2009.
- Jokinen, T., Berndt, T., Makkonen, R., Kerminen, V.-M., Junninen, H., Paasonen, P., Stratmann, F., Herrmann, H., Guenther, A. B., Worsnop, D. R., Kulmala, M., Ehn, M. and Sipilä, M.: Production of extremely low volatile organic compounds from biogenic emissions: Measured yields and atmospheric implications, *Proc. Natl. Acad. Sci. U. S. A.*, 112(23), 7123–7128, doi:10.1073/pnas.1423977112, 2015.
- 495 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prevot, a S. H., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R. and Baltensperger, U.: Identification of polymers as major components of atmospheric organic aerosols., *Science*, 303(5664), 1659–1662, doi:10.1126/science.1092185, 2004.
- Kerminen, V. M., Paramonov, M., Anttila, T., Riipinen, I., Fountoukis, C., Korhonen, H., Asmi, E., Laakso, L., Lihavainen, H., Swietlicki, E., Svenningsson, B., Asmi, A., Pandis, S. N., Kulmala, M. and Petäjä, T.: Cloud condensation nuclei production associated with atmospheric nucleation: A synthesis based on existing literature and new results, *Atmos. Chem. Phys.*, 12(24), 12037–12059, doi:10.5194/acp-12-12037-2012, 2012.
- 500 Kourtchev, I., Giorio, C., Manninen, A., Wilson, E., Mahon, B., Aalto, J., Kajos, M., Venables, D., Ruuskanen, T., Levula, J., Lopenon, M., Connors, S., Harris, N., Zhao, D., Kiendler-Scharr, A., Mentel, T., Rudich, Y., Hallquist,

- 505 M., Doussin, J.-F., Maenhaut, W., Bäck, J., Petäjä, T., Wenger, J., Kulmala, M. and Kalberer, M.: Enhanced Volatile Organic Compounds emissions and organic aerosol mass increase the oligomer content of atmospheric aerosols, *Sci. Rep.*, 6, 35038, doi:10.1038/srep35038, 2016.
- Kreidenweis, S. M., Koehler, K., DeMott, P., Prenni, a. J., Carrico, C. and Ervens, B.: Water activity and activation diameters from hygroscopicity data - Part I: Theory and application to inorganic salts, *Atmos. Chem. Phys. Discuss.*, 5(1), 287–323, doi:10.5194/acpd-5-287-2005, 2005.
- 510 Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42(16), 3593–3624, doi:10.1016/j.atmosenv.2008.01.003, 2008.
- Kuang, C., Riipinen, I., Sihto, S. L., Kulmala, M., McCormick, A. V. and McMurry, P. H.: An improved criterion for new particle formation in diverse atmospheric environments, *Atmos. Chem. Phys.*, 10(17), 8469–8480, doi:10.5194/acp-10-8469-2010, 2010.
- 515 Kuang, C., Chen, M., Zhao, J., Smith, J., McMurry, P. H. and Wang, J.: Size and time-resolved growth rate measurements of 1 to 5 nm freshly formed atmospheric nuclei, *Atmos. Chem. Phys.*, 12(7), 3573–3589, doi:10.5194/acp-12-3573-2012, 2012.
- Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M. V., Smith, J. N., Ehn, M., Mentel, T. F., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M. and Worsnop, D. R.: Direct observations of atmospheric aerosol nucleation, *Science*, 339(6122), 943–6, doi:10.1126/science.1227385, 2013.
- 520 Lee, A. K. Y., Zhao, R., Li, R., Liggio, J., Li, S. M. and Abbatt, J. P. D.: Formation of light absorbing organo-nitrogen species from evaporation of droplets containing glyoxal and ammonium sulfate, *Environ. Sci. Technol.*, 47(22), 12819–12826, doi:10.1021/es402687w, 2013.
- Lemaire, V., Coll, I., Couvidat, F., Mouchel-Vallon, C., Seigneur, C. and Siour, G.: Oligomer formation in the troposphere: From experimental knowledge to 3-D modeling, *Geosci. Model Dev.*, 9(4), 1361–1382, doi:10.5194/gmd-9-1361-2016, 2016.
- 530 Liu, A. T., Zaveri, R. A. and Seinfeld, J. H.: Analytical solution for transient partitioning and reaction of a condensing vapor species in a droplet, *Atmos. Environ.*, 89, 651–654, doi:10.1016/j.atmosenv.2014.02.065, 2014.
- Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys. Discuss.*, 4(6), 7561–7614, doi:10.5194/acpd-4-7561-2004, 2004.
- 535 Lopez-Hilfiker, F. D., Mohr, C., D'Ambro, E. L., Lutz, A., Riedel, T. P., Gaston, C. J., Iyer, S., Zhang, Z., Gold, A., Surratt, J. D., Lee, B. H., Kurten, T., Hu, W. W., Jimenez, J., Hallquist, M. and Thornton, J. A.: Molecular Composition and Volatility of Organic Aerosol in the Southeastern U.S.: Implications for IEPOX Derived SOA,

- Environ. Sci. Technol., 50(5), 2200–2209, doi:10.1021/acs.est.5b04769, 2016.
- 540 Mai, H., Shiraiwa, M., Flagan, R. C. and Seinfeld, J. H.: Under What Conditions Can Equilibrium Gas-Particle Partitioning Be Expected to Hold in the Atmosphere?, Environ. Sci. Technol., 49(19), 11485–11491, doi:10.1021/acs.est.5b02587, 2015.
- 545 Mentel, T. F., Springer, M., Ehn, M., Kleist, E., Pullinen, I., Kurtén, T., Rissanen, M., Wahner, A. and Wildt, J.: Formation of highly oxidized multifunctional compounds: autoxidation of peroxy radicals formed in the ozonolysis of alkenes – deduced from structure–product relationships, Atmos. Chem. Phys., 15(12), 6745–6765, doi:10.5194/acp-15-6745-2015, 2015.
- Pankow, J. F.: An Absorption Model of Gas / Particle Partitioning of Organic Compounds in the Atmosphere, Atmos. Environ., 28(2), 185–188, doi:10.1016/1352-2310(94)90093-0, 1994.
- 550 Pennington, M. R., Bzdek, B. R., Depalma, J. W., Smith, J. N., Kortelainen, A. M., Hildebrandt Ruiz, L., Petäjä, T., Kulmala, M., Worsnop, D. R. and Johnston, M. V.: Identification and quantification of particle growth channels during new particle formation, Atmos. Chem. Phys., 13(20), 10215–10225, doi:10.5194/acp-13-10215-2013, 2013.
- Poling, B. E., Prausnitz, J. M. and O’Connell, J. P.: The properties of gases and liquids, 5th Edn., McGraw-Hill, New York, USA., 2001.
- 555 Reinhardt, A., Emmenegger, C., Gerrits, B., Panse, C., Dommen, J., Baltensperger, U., Zenobi, R. and Kalberer, M.: Ultrahigh mass resolution and accurate mass measurements as a tool to characterize oligomers in secondary organic aerosols, Anal. Chem., 79(11), 4074–4082, doi:10.1021/ac062425v, 2007.
- 560 Riccobono, F., Schobesberger, S., Scott, C. E., Dommen, J., Ortega, I. K., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hansel, A., Junninen, H., Kajos, M., Keskinen, H., Kupc, A., Kurten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petaja, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipilä, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tome, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R., Baltensperger, U., Kürten, A., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Nieminen, T., Onnela, A., Petäjä, T., Praplan, A. P., Santos, F. D., Schallhart, S., Seinfeld, J. H., Sipilä, M., Spracklen, D. V., Stozhkov, Y., Stratmann, F., Tomé, A., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Wimmer, D., Carslaw, K. S., Curtius, J., Donahue, N. M., Kirkby, J., Kulmala, M., Worsnop, D. R. and Baltensperger, U.: Oxidation Products of Biogenic Emissions Contribute to Nucleation of Atmospheric Particles, Science, 344(6185), 717–721, doi:10.1126/science.1243527, 2014.
- 570 Riipinen, I., Yli-Juuti, T., Pierce, J. R., Petäjä, T., Worsnop, D. R., Kulmala, M. and Donahue, N. M.: The contribution of organics to atmospheric nanoparticle growth, Nat. Geosci., 5(7), 453–458, doi:10.1038/ngeo1499, 2012.

- Riva, M., Budisulistiorini, S. H., Chen, Y., Zhang, Z., D'Ambro, E. L., Zhang, X., Gold, A., Turpin, B. J., Thornton, J. A., Canagaratna, M. R. and Surratt, J. D.: Chemical Characterization of Secondary Organic Aerosol from Oxidation of Isoprene Hydroxyhydroperoxides, *Environ. Sci. Technol.*, 50(18), 9889–9899, doi:10.1021/acs.est.6b02511, 2016.
- 575
- Roldin, P., Eriksson, A. C., Nordin, E. Z., Hermansson, E., Mogensen, D., Rusanen, A., Boy, M., Swietlicki, E., Svenningsson, B., Zelenyuk, A. and Pagels, J.: Modelling non-equilibrium secondary organic aerosol formation and evaporation with the aerosol dynamics, gas- and particle-phase chemistry kinetic multilayer model ADCHAM, *Atmos. Chem. Phys.*, 14(15), 7953–7993, doi:10.5194/acp-14-7953-2014, 2014.
- 580
- Saul, T. D., Tolocka, M. P. and Johnston, M. V.: Reactive uptake of nitric acid onto sodium chloride aerosols across a wide range of relative humidities, *J. Phys. Chem. A*, 110(24), 7614–7620, doi:10.1021/jp060639a, 2006.
- Shiraiwa, M., Pfrang, C., Koop, T. and Pöschl, U.: Kinetic multi-layer model of gas-particle interactions in aerosols and clouds (KM-GAP): Linking condensation, evaporation and chemical reactions of organics, oxidants and water, *Atmos. Chem. Phys.*, 12(5), 2777–2794, doi:10.5194/acp-12-2777-2012, 2012.
- 585
- Shiraiwa, M., Yee, L. D., Schilling, K. A., Loza, C. L., Craven, J. S., Zuend, A., Ziemann, P. J. and Seinfeld, J. H.: Size distribution dynamics reveal particle-phase chemistry in organic aerosol formation., *Proc. Natl. Acad. Sci. U. S. A.*, 110(29), 11746–50, doi:10.1073/pnas.1307501110, 2013.
- Shiraiwa, M., Berkemeier, T., Schilling-Fahnestock, K. A., Seinfeld, J. H. and Pöschl, U.: Molecular corridors and kinetic regimes in the multiphase chemical evolution of secondary organic aerosol, *Atmos. Chem. Phys.*, 14(16), 8323–8341, doi:10.5194/acp-14-8323-2014, 2014.
- 590
- Shrivastava, M. K., Zhao, C., Easter, R. C., Qian, Y., Zelenyuk, A., Fast, J. D., Liu, Y., Zhang, Q. and Guenther, A.: Journal of Advances in Modeling Earth Systems, *J. Adv. Model. Earths Syst.*, 8, 499–519, doi:10.1002/2015MS000554.Received, 2016.
- Smith, G. D., Woods III, E., Baer, T. and Miller, R. E.: Aerosol Uptake Described by Numerically Solving the Diffusion-Reaction Equations in the Particle, *J. Phys. Chem. A*, 107, 9582–9587, doi:10.1021/jp021843a, 2003.
- 595
- Smith, J. N., Dunn, M. J., VanReken, T. M., Iida, K., Stolzenburg, M. R., McMurry, P. H. and Huey, L. G.: Chemical composition of atmospheric nanoparticles formed from nucleation in Tecamac, Mexico: Evidence for an important role for organic species in nanoparticle growth, *Geophys. Res. Lett.*, 35(4), 2–6, doi:10.1029/2007GL032523, 2008.
- 600
- Song, M., Liu, P. F., Hanna, S. J., Li, Y. J., Martin, S. T. and Bertram, A. K.: Relative humidity-dependent viscosities of isoprene-derived secondary organic material and atmospheric implications for isoprene-dominant forests, *Atmos. Chem. Phys.*, 15(9), 5145–5159, doi:10.5194/acp-15-5145-2015, 2015.
- Stangl, C. M. and Johnston, M. V.: Aqueous Reaction of Dicarboxyls with Ammonia as a Potential Source of Organic Nitrogen in Airborne Nanoparticles, *J. Phys. Chem. A*, doi:10.1021/acs.jpca.7b02464, 2017.

- 605 Stolzenburg, M. R., McMurry, P. H., Sakurai, H., Smith, J. N., Mauldin, R. L., Eisele, F. L. and Clement, C. F.: Growth rates of freshly nucleated atmospheric particles in Atlanta, *J. Geophys. Res. Atmos.*, 110(22), 1–10, doi:10.1029/2005JD005935, 2005.
- Surratt, J. D., Lewandowski, M., Offenberg, J. H., Kleindienst, T. E., Edney, E. O., Seinfeld, J. H. and Surratt, J. D.: Effect of Acidity on Secondary Organic Aerosol Formation from Isoprene Effect of Acidity on Secondary Organic
610 Aerosol Formation from Isoprene, *Environ. Sci. Technol.*, 41(15), 5363–5369, doi:10.1021/es0704176, 2007.
- Tang, X., Misztal, P. K. and Goldstein, A. H.: Emitted from Engineering Students in a Classroom, *Environ. Sci. Technol. Lett.*, 2, 303–307, doi:10.1021/acs.estlett.5b00256, 2015.
- Tolocka, M. P., Jang, M., Ginter, J. M., Cox, J., Kamens, R. M., Johnston, M. V and Tolocka, M. P.: Formation of Oligomers in Secondary Organic Aerosol, , 38(5), 1428–1434, doi:10.1021/es035030r, 2004a.
- 615 Tolocka, M. P., Saul, T. D. and Johnston, M. V: Reactive uptake of nitric acid into aqueous sodium chloride particles using real-time single-particle mass spectrometry , *J. Phys. Chem. A*, 108, 2659–2665, doi:10.1021/jp036612y, 2004b.
- Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer,
620 A.-K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Möhler, O., Nieminen, T., Onnela, A., Petäjä, T., Piel, F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipilä, M., Smith, J. N., Steiner, G., Tomè, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Carslaw,
625 K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop, D. R., Donahue, N. M. and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533(7604), 527–531, doi:10.1038/nature18271, 2016.
- Trump, E. R. and Donahue, N. M.: Oligomer formation within secondary organic aerosols: equilibrium and dynamic considerations, *Atmos. Chem. Phys.*, 14, 3691–3701, doi:10.5194/acp-14-3691-2014, 2014.
- 630 Tu, P. and Johnston, M. V.: Particle Size Dependence of Biogenic Secondary Organic Aerosol Molecular Composition, *Atmos. Chem. Phys. Discuss.*, DOI:10.5194/acp-2017-53, doi:10.5194/acp-2017-53, 2017.
- Tu, P., Hall, W. A. and Johnston, M. V.: Characterization of Highly Oxidized Molecules in Fresh and Aged Biogenic Secondary Organic Aerosol, *Anal. Chem.*, 88(8), 4495–4501, doi:10.1021/acs.analchem.6b00378, 2016.
- 635 Vestenius, M., Hellén, H., Levula, J., Kuronen, P., Helminen, K. J., Nieminen, T., Kulmala, M. and Hakola, H.: Acidic reaction products of monoterpenes and sesquiterpenes in atmospheric fine particles in a boreal forest, *Atmos. Chem. Phys.*, 14(15), 7883–7893, doi:10.5194/acp-14-7883-2014, 2014.
- Vesterinen, M., Lehtinen, K. E. J., Kulmala, M. and Laaksonen, A.: Effect of particle phase oligomer formation on

- aerosol growth, *Atmos. Environ.*, 41(8), 1768–1776, doi:10.1016/j.atmosenv.2006.10.024, 2007.
- 640 Weber, R. J., Marti, J. J. and McMurry, P. H.: Measured Atmospheric New Particle Formation Rates: Implications for Nucleation Mechanisms, *Chem. Eng. Comm.*, 151, 53–64, 1996.
- Wehner, B., Petäjä, T., Boy, M., Engler, C., Birmili, W., Tuch, T., Wiedensohler, A. and Kulmala, M.: The contribution of sulfuric acid and non-volatile compounds on the growth of freshly formed atmospheric aerosols, *Geophys. Res. Lett.*, 32(2), 3–6, doi:10.1029/2005GL023827, 2005.
- 645 Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions H⁺, NH₄⁺, Na⁺, SO₄²⁻, NO₃⁻, Cl⁻, Br⁻, and H₂O, *J. Geophys. Res.*, 107, 1–14, doi:10.1029/2001JD000451, 2002.
- Winkler, P. M., Ortega, J., Karl, T., Cappellin, L., Friedli, H. R., Barsanti, K., McMurry, P. H. and Smith, J. N.: Identification of the biogenic compounds responsible for size-dependent nanoparticle growth, *Geophys. Res. Lett.*, 39(20), 1–6, doi:10.1029/2012GL053253, 2012.
- 650 Wong, J. P. S., Lee, A. K. Y. and Abbatt, J. P. D.: Impacts of Sulfate Seed Acidity and Water Content on Isoprene Secondary Organic Aerosol Formation, *Environ. Sci. Technol.*, 49(22), 13215–13221, doi:10.1021/acs.est.5b02686, 2015.
- Wu, Y. and Johnston, M. V.: Aerosol Formation from OH Oxidation of the Volatile Cyclic Methyl Siloxane (cVMS) Decamethylcyclopentasiloxane, *Environ. Sci. Technol.*, 51, 4445–4451, doi:10.1021/acs.est.7b00655, 2017.
- 655 Xu, L., Guo, H., Boyd, C. M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-vanwertz, G., Kreisberg, N. M., Olson, K., Koss, A., Goldstein, A. H., Susanne, V., Gouw, J. De, Baumann, K., Lee, S., Nenes, A., Weber, R. J. and Ng, N. L.: Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States, *Proc. Natl. Acad. Sci.*, 112(32), 37–42, doi:10.1073/pnas.1512277112, 2015.
- Yli-Juuti, T., Barsanti, K., Hildebrandt Ruiz, L., Kieloaho, A. J., Makkonen, U., Petäjä, T., Ruuskanen, T., Kulmala, M. and Riipinen, I.: Model for acid-base chemistry in nanoparticle growth (MABNAG), *Atmos. Chem. Phys.*, 13(24), 12507–12524, doi:10.5194/acp-13-12507-2013, 2013.
- 660 Zaveri, R. A., Easter, R. C., Shilling, J. E. and Seinfeld, J. H.: Modeling kinetic partitioning of secondary organic aerosol and size distribution dynamics: Representing effects of volatility, phase state, and particle-phase reaction, *Atmos. Chem. Phys.*, 14(10), 5153–5181, doi:10.5194/acp-14-5153-2014, 2014.
- Zhang, R., Khalizov, A., Wang, L., Hu, M. and Xu, W.: Nucleation and growth of nanoparticles in the atmosphere, *Chem. Rev.*, 112, 1957–2011, doi:10.1021/cr2001756, 2012.
- 665 Zhao, J., Ortega, J., Chen, M., McMurry, P. H. and Smith, J. N.: Dependence of particle nucleation and growth on high-molecular-weight gas-phase products during ozonolysis of α -pinene, *Atmos. Chem. Phys.*, 13(15), 7631–7644, doi:10.5194/acp-13-7631-2013, 2013.
- Ziemann, P. J. and Atkinson, R.: Kinetics, products, and mechanisms of secondary organic aerosol formation,


Tables and Figures

675 **Table 1:** Mixing ratios and relevant physical properties of chemical species included in particle growth calculations.
(T = 282 °C, RH = 50%)

| | Mixing Ratio, C_{ig} | | Saturation Concentration | Molar Mass | Density | Gas Phase Diffusion Coefficient |
|----------------------|------------------------|---------|---------------------------------------|---------------------|--------------------|---------------------------------|
| | ng m^{-3} | pptv | $\log C^*$, ($\mu\text{g m}^{-3}$) | g mol^{-1} | g cm^{-3} | $\text{cm}^2 \text{s}^{-1}$ |
| Sulfuric Acid | 0.5 | 0.12 | - | 98 | 1.8 | 0.1 |
| Ammonia | 1 | 1.25 | - | 17 | 0.7 | 0.2 |
| SVOC | 3-11 | 0.4-1.4 | 10^0 | 200 | 1.2 | 0.05 |
| NVOC | 3 | 0.4 | 10^{-4} | 200 | 1.2 | 0.05 |
| DIMER | - | - | - | 400 | 1.2 | - |

Owner 12/28/2017 11:55 AM
 Deleted: [Section Break \(Next Page\)](#)
 *
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt

681




682

683 **Figure 1:** Particle diameter dependence of a) growth rates and b) dry mass fractions of chemical species under the
684 conditions where the gas phase mixing ratios are constant and growth occurs by partitioning alone.

685

686

Unknown
Formatted: Font:(Default) Times New Roman, 10 pt




687


688 **Figure 2:** Mass fraction ratio of SVOC / NVOC vs. particle diameter under conditions where growth occurs by
 689 partitioning alone.

690

691



Deleted:
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt




693


694 **Figure 3:** Particle diameter dependence of a) growth rates and b) dry mass fractions of chemical species under the
 695 conditions where both partitioning and particle phase chemistry (SVOC-SVOC dimer formation) are included.

696

697



Deleted:
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt



699


700 **Figure 4:** Mass ratio vs. particle diameter for a) SVOC to NVOC and b) DIMER to NVOC, under conditions where
 701 both partitioning and particle phase chemistry (SVOC-SVOC dimer formation) are included.

702

703

Owner 12/28/2017 11:55 AM

Deleted:
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt



705

706 **Figure 5:** Growth rate vs. particle diameter: a) Comparison of growth rates with different DIMER formation rate
 707 constants for the SVOC-SVOC reaction. b) Comparison of growth rates for SVOC-SVOC, NVOC-NVOC and
 708 SVOC-NVOC reactions having a DIMER formation rate constant of $10^{-2} M^{-1}s^{-1}$. In these plots, the black line is
 709 equivalent to the particle diameter dependence in Figure 1a.

Owner 12/28/2017 11:55 AM

a) Rate Constant ($M^{-1}s^{-1}$)
Growth Rate (nm/h)
Diameter (nm)

b) Growth Rate (nm/h)
Diameter (nm)
— NVOC-NVO

Deleted:
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt
 Unknown
 Formatted: Font:(Default) Times New Roman, 10 pt
 Owner 12/28/2017 11:55 AM
 Deleted: for
 Owner 12/28/2017 11:55 AM
 Deleted: in

1 Supporting Information: Nanoparticle Growth by Particle 2 Phase Chemistry


3 Michael J. Apsokardu and Murray V. Johnston

4 Department of Chemistry and Biochemistry, University of Delaware, Newark, DE, 19716, USA

5 Correspondence to: mvj@udel.edu

6 Four figures (S1-S4) and associated discussion.

7 Figure S1




8 Figure S1 shows a schematic of the iterative calculations, starting with volume $V_{p,n}$, the total volume of the particle at the start of time period n . Color shaded regions delineate particle growth by condensation (red), partitioning (green), and particle phase chemistry (yellow). Calculations begin with the Kelvin effect modified vapor pressure ($KEMP_d$):

9
10
11
12
13 (S1) $KEMP_d = P_0 e^{\left[\frac{2\sigma V_{M,p}}{RT} \right]}$

14 where P_0 is the saturation vapor pressure over a flat surface, σ is the surface tension, $V_{M,p}$ is the average molar volume of the particle, d is the particle diameter, R is the universal gas constant, and T is the temperature. The subscript d , shown here with $KEMP_d$, and for other variables hereafter, represents the particle diameter and denotes that the variable is particle size dependent. The saturation ratio (S_d) is given by:

Murray Johnston 12/28/2017 1:10 PM
Deleted: Three



Deleted:

Murray Johnston 12/28/2017 1:10 PM
Deleted:) given as:

Murray Johnston 12/28/2017 1:10 PM
Deleted: $e^{\left[\frac{2\sigma V_{M,p}}{RT} \right]}$,

Murray Johnston 12/28/2017 1:10 PM
Deleted: Subscript D

Murray Johnston 12/28/2017 1:10 PM
Deleted: the size dependence of

Murray Johnston 12/28/2017 1:10 PM
Deleted: term. $KEMP_D$ defines

Murray Johnston 12/28/2017 1:10 PM
Deleted:), which determines whether uptake will occur at a rate either equal to or less than the condensation rate. The saturation ratio is

29 (S2) $S_d = C_{lg}/KEMP_d$

30 For compounds having $S_d < 1$, uptake occurs at a slower rate than the condensation rate, while for compounds
 31 having $S_d \gg 1$, uptake occurs at the condensation rate.

32 For the molecular species considered in this study, those growing the particle at the condensation rate (red
 33 shaded region of Figure S1) are sulfuric acid and non-volatile organic compound (NVOC). Equation 1 in the main
 34 text gives the uptake rate, which assumes that every collision results in uptake. The Joyalka mass transfer correction
 35 factor β_d is given by:

36 (S3)
$$\beta_d = \frac{\sqrt{\pi} Kn (1+1.333)}{1 + Kn (1.333 + 1.333\sqrt{\pi} Kn + \delta)}$$

37 where α is the mass accommodation coefficient (assumed to be 1), δ is the mass transfer jump coefficient equal to
 38 1.0161, and Kn is the Knudsen number:

39 (S4)
$$Kn = \frac{2\lambda}{d_p + d_i}$$

40 where λ is the mean free path and d is diameter. Subscripts p and i indicate particle and gas molecule respectively.
 41 The mean free path is defined as:

42 (S5)
$$\lambda = \frac{4(D_p + D_i)}{\sqrt{\pi}(c_p^2 + c_i^2)^{1/2}}$$

43 where D is the gas phase diffusion coefficient and c is the mean thermal speed. The gas phase diffusion coefficient
 44 of a particle is defined as

45 (S6)
$$D_p = \frac{k_B T C_c}{3\pi\mu_{air}d_p}$$

46 where C_c is the Cunningham slip correction factor and μ_{air} is the dynamic viscosity of air ($\mu_{air} = 1.76 \times 10^{-5} \text{ kg m}^{-1} \text{ s}^{-1}$
 47 at 282 K). The gas phase diffusion coefficient of a vapor molecule is estimated from:

48 (S7)
$$D_i = \frac{0.001 T^{1.75}}{P (\Sigma_{air}^{1/3} + \Sigma_i^{1/3})^2} \sqrt{\frac{1}{MW_{air}} + \frac{1}{MW_i}}$$

49 where P is atmospheric pressure, Σ is the atomic diffusion volume, and MW is the molecular weight. Eq. S7 is to
 50 calculate the gas phase diffusion coefficients for sulfuric acid and ammonia. For Σ_{air} and MW_{air} , the values 19.7 and
 51 28.97 g mol⁻¹ are used respectively. A list of atomic diffusion volumes can be found in Poling et al 2001. The
 52 diffusion coefficients for SVOC and NVOC vapor molecules were assumed to be 0.05 cm² s⁻¹. This value is
 53 consistent with Eq. S7 for typical molecular structures of monomer products of monoterpene ozonolysis.

54 Seinfeld and Pandis (Seinfeld, John H. and Pandis, Spyros N. "Atmospheric Chemistry and Physics: From
 55 Air Pollution To Climate Change", John Wiley and Sons, New York, 2006, ISBN-13: 978-0471720188) give a
 56 detailed discussion of various calculation procedures for the mass transfer coefficient β_d in Equation 1, including the

Murray Johnston 12/28/2017 1:10 PM
 Deleted: .

Murray Johnston 12/28/2017 1:10 PM
 Deleted: g

Murray Johnston 12/28/2017 1:10 PM
 Deleted: en

Murray Johnston 12/28/2017 1:10 PM
 Deleted: Fuchs-Sutugin

Murray Johnston 12/28/2017 1:10 PM
 Deleted: $= \frac{1+Kn}{1+(4/3\alpha)+0.337(Kn)+(4/3\alpha)(Kn)^2}$

Murray Johnston 12/28/2017 1:10 PM
 Deleted:)

Murray Johnston 12/28/2017 1:10 PM
 Deleted: $\frac{2\lambda}{D}$

Murray Johnston 12/28/2017 1:10 PM
 Deleted: particle

Murray Johnston 12/28/2017 1:10 PM
 Deleted: $\frac{k_B T}{\sqrt{2P\pi} ((D+D_i)/2)^2}$

Murray Johnston 12/28/2017 1:10 PM
 Deleted: k_B is the Boltzmann constant, T is

Murray Johnston 12/28/2017 1:10 PM
 Deleted: temperature, P

Murray Johnston 12/28/2017 1:10 PM
 Deleted: and D_i is the diameter of a gas molecule i .


70 [Loyalka approach used in our work. As a check of our procedure, we also used the Fuchs-Sutugin approach to](#)
71 [calculate \$\beta_d\$ and obtained equivalent results within the precision of the respective theories.](#)

72 Semi-volatile organic compounds (SVOC) cause particle growth at a rate that is slower than the
73 condensation rate ([green shaded region in Figure S1](#)). Based on the gas phase mixing ratio and particle properties, a
74 corresponding equilibrium particle phase concentration is calculated (Eq. 2 of the main text). The mass of such
75 species added to the particle is based on $V_{p,n}$, so by $V_{p,n+1}$, the species is no longer in equilibrium and must be re-
76 calculated. Partitioning of water is dependent on the mixing ratio and the $KEMP_d$ to determine the equilibrium mole
77 fraction x_i . For simplicity, the activity coefficient (ζ) is assumed to be 1.

78 Particle phase chemistry occurs by an accretion reaction ([yellow shaded region of Figure S1](#)). Reactions are
79 modeled by the second order decay of SVOC (or in the case of Figures S2 and S3, NVOC) to produce DIMER
80 products. Depletion of SVOC is dependent on the concentration of SVOC existing in the particle at $V_{p,n}$. When
81 equilibrium is re-calculated for SVOC at $V_{p,n+1}$, the mass added to the particle must account for both depletion by
82 reaction and dilution due to particle growth. When the volume changes for all individual species have been
83 calculated, they are summed to give the new particle volume, $V_{p,n+1}$. After volume $V_{p,n+1}$ is achieved, calculations are
84 iteratively repeated.

85

86 **Figure S2**



87
88 Figure S2 shows growth rate vs particle diameter for the comparison of the growth rate due to particle
89 phase reaction at different gas phase mixing ratios of SVOC. “1.4 - NR” indicates no particle phase reaction occurs
90 and growth is only due to condensation [and partitioning](#).

91

92


Murray Johnston 12/28/2017 1:10 PM

Deleted: yellow

Murray Johnston 12/28/2017 1:10 PM


Deleted: red

Murray Johnston 12/28/2017 1:10 PM



Deleted:

96 **Figure S3**




97

98 Figures S3a and b show the size dependent evolution of particle growth rate by a) partitioning alone and b)
99 with dimer formation from NVOC ($k_H = 10^{-3} \text{ M}^{-1}\text{s}^{-1}$). Dimer formation from NVOC does not enhance the growth
100 rate (growth still proceeds at the NVOC collision rate), but it does change the composition.



101

102 **Figure S4**



103

104 Figure S4 shows the mass fraction ratio of DIMER to NVOC, which **monotonically** increases with
105 increasing particle diameter.



Murray Johnston 12/28/2017 1:10 PM
Deleted: systematically