

Overview and Preliminary Results of the Surface Ocean Aerosol Production (SOAP) campaign

5 Law Cliff S.^{1,2}, Smith Murray J.¹, Harvey Mike J.¹, Bell Thomas G.^{3,4}, Cravigan Luke T.^{5,6}, Elliott Fiona C.¹,
Lawson Sarah J.⁷, Lizotte Martine⁸, Marriner Andrew¹, McGregor John¹, Ristovski Zoran⁶, Safi Karl A⁹,
Saltzman Eric S.⁴, Vaattovaara Petri¹⁰, Walker Carolyn F.¹

- 10 ¹ National Institute of Water and Atmospheric Research, Wellington, New Zealand
² Department of Chemistry, University of Otago, Dunedin, New Zealand
³ Plymouth Marine Laboratory, Prospect Place, The Hoe, Plymouth, UK
⁴ Department of Earth System Science, University of California, Irvine, CA, USA
⁵ Climate Science Centre, Commonwealth Scientific and Industrial Research Organisation, Aspendale,
15 Australia
⁶ International Laboratory for Air Quality and Health, Queensland University of Technology,
Brisbane, Australia
⁷ Commonwealth Scientific and Industrial Research Organisation, Oceans and Atmosphere Flagship,
Aspendale, Australia
20 ⁸ Université Laval, Department of Biology (Québec-Océan), Québec City, Québec,
Canada.
⁹ National Institute of Water and Atmospheric Research, Hamilton, New Zealand
¹⁰ University of Eastern Finland, Kuopio, Finland

25 Correspondence to Cliff Law cliff.law@niwa.co.nz

Abstract. Establishing the relationship between marine boundary layer (MBL) aerosols and surface
30 water biogeochemistry is required to understand aerosol and cloud production processes over the
remote ocean, and represent them more accurately in Earth System Models and global climate
projections. This was addressed by the SOAP (Surface Ocean Aerosol Production) campaign, which
examined air-sea interaction over biologically-productive frontal waters east of New Zealand. This
overview details the objectives, regional context, sampling strategy, and provisional findings of a pilot
35 study, PreSOAP, in austral summer 2011, and the following SOAP voyage in late austral summer 2012.
Both voyages characterised surface water and MBL composition in three phytoplankton blooms of
differing species composition and biogeochemistry, with significant regional correlation observed
between chlorophyll-*a* and DMS_{sw}. Surface seawater dimethylsulfide (DMS_{sw}) and associated air-sea

DMS flux showed spatial variation during the SOAP voyage, with maxima of 25 nmol L⁻¹ and 100
40 $\mu\text{mol m}^{-2} \text{d}^{-1}$, respectively, recorded in a dinoflagellate bloom. Inclusion of SOAP data in a regional
DMS_{sw} compilation indicates that the current climatological mean is an underestimate for this region
of the South-west Pacific. Estimation of the DMS gas transfer velocity (k_{DMS}) by independent techniques
of eddy covariance and gradient flux showed good agreement, although both exhibited periodic
deviations from model estimates. Flux anomalies were related to surface warming and sea surface
45 microlayer enrichment, and also reflected the heterogeneous distribution of DMS_{sw} and the associated
flux footprint. Other aerosol precursors measured included the halides and various volatile organic
carbon compounds, with first measurements of the short-lived gases glyoxal and methylglyoxal in
pristine Southern Ocean marine air indicating an unidentified local source. The application of a real-time
clean-sector, contaminant markers, and a common aerosol inlet facilitated multi-sensor measurement
50 of uncontaminated air. Aerosol characterisation identified variable Aitken mode, and consistent sub-
micron sized accumulation and coarse modes. Sub-micron aerosol mass was dominated by secondary
particles containing ammonium sulfate/bisulfate under light winds, with an increase in sea-salt under
higher wind-speeds. MBL measurements and chamber experiments identified a significant organic
component in primary and secondary aerosols. Comparison of SOAP aerosol number and size
55 distributions reveals an underprediction in GLOMAP-mode aerosol number in clean marine air masses,
suggesting a missing marine aerosol source in the model. The SOAP data will be further examined for
evidence of nucleation events, and also to identify relationships between MBL composition and surface
ocean biogeochemistry that may provide potential proxies for aerosol precursors and production.

60 1. Introduction

It is recognised that the surface ocean alters the properties of the lower atmosphere, and so atmospheric albedo and climate (McCoy et al., 2015; Seinfeld et al., 2016), via the direct and indirect effects of aerosols (O'Dowd and de Leeuw, 2007). Aerosols are precursors of clouds, which play a major role in the scattering and absorption of incident solar radiation (Carslaw et al., 2013), but the concentration, number and chemical properties of aerosols that act as cloud-condensation nuclei (CCN) can also influence cloud droplet size and number, and consequently precipitation and cloud albedo (Twomey, 1977). Indeed, cloud formation and properties are sensitive to relatively minor changes in aerosol concentration, particularly in remote regions (Carslaw et al., 2013). This is particularly the case in the Southern Ocean, where natural aerosol sources dominate and where CCN concentrations can range from tens per cm^3 in winter to hundreds per cm^3 in summer (Andreae and Rosenfeld, 2008), leading to seasonally variant trends in cloud albedo. However, the relationship between clouds and aerosols derived from natural sources is poorly understood, and represents a major uncertainty in the representation of low-level marine clouds and feedbacks in climate models (Wang et al., 2013; Stephens, 2005). Current models underestimate cloud over the Southern Ocean, particularly south of 55°S, resulting in excess surface shortwave radiation and a warm bias (Trenberth and Fasullo, 2010; Kay et al., 2016). This discrepancy is potentially attributable to a variety of factors, chief among which is the limited understanding of aerosol-cloud interaction and cloud water phase, compounded by a lack of regional observations and data to advance satellite retrievals and climate model simulations.

Breaking waves and associated bubble formation are a major source of Primary Marine Aerosol (PMA), supplying most the aerosol mass in the marine boundary layer (MBL) over the remote ocean (Andreae and Rosenfeld, 2008), and particularly in regions that experience high winds and breaking waves (de Leeuw et al., 2014). This is reflected in PMA contributing only ~10–20% of CCN number concentrations over the remote Pacific Ocean (Blot et al. 2013; Clarke et al. 2013), but up to 55% over the Southern Ocean (McCoy et al. 2015). Although PMA is generally regarded as primarily composed of sea-salt, recent reassessments suggest it is highly enriched in organic matter relative to bulk seawater. Organic material may in fact dominate submicron aerosol mass (Facchini et al., 2008; O'Dowd et al., 2004), with the Primary Organic Aerosol (POA) being of biogenic origin and including bacteria, carbohydrates, polymers and gels (Facchini et al., 2008; Russell et al., 2010). Although the contribution of POA to the MBL is uncertain, it may be significant over biologically active oceanic regions, as suggested by correlations between organic aerosol content and surface chlorophyll-*a* (Chl-*a*) (O'Dowd et al., 2004). There is also similarity in the composition of aerosol and surface ocean organics, and organically enriched sub-micron particles have been produced experimentally using surface seawater conditions

(Quinn and Bates, 2011). Indeed, the degree of organic enrichment may influence both the type and size of aerosols, as well as properties such as aerosol light scattering and water uptake (Vaishya et al., 2012).

It is well-established that biologically productive regions are characterised by elevated concentrations and emissions of a range of compounds that may influence aerosol production, composition and properties (Meskhidze and Nenes, 2010; Gantt and Meskhidze 2013; de Leeuw et al., 2014). However, the oceanic influence on atmospheric composition is not only attributable to PMAs but also to secondary marine aerosols (SMAs), that are produced during gas-phase reactions of volatile organic compounds (VOCs). Although SMAs have less impact upon aerosol mass they potentially have a large influence on aerosol number (Meskhidze et al., 2011). The biogeochemical origin of SMAs is reflected in their seasonality, with Aitken and accumulation mode aerosol number concentrations dominated by secondary particles in summertime (Clarke et al. 2013; Cravigan et al. 2015). Research into SMAs has primarily focussed on dimethylsulfide (DMS), the primary natural marine source of volatile sulfur, in response to early hypotheses related to its potential role in climate feedback processes (Charlson et al., 1987). The CLAW hypothesis linked the production of the DMS precursor, dimethylsulfoniopropionate (DMSP), by phytoplankton and subsequent DMS emission and oxidation to sulfate aerosol, to CCN formation and changes in cloud cover. Although well-studied, this hypothesis remains unproven and there is a lack of consensus, with a recent review identifying uncertainties regarding the role of DMS in aerosol production in the MBL (Quinn and Bates, 2011). However, there is evidence that DMS may play a role in cloud formation over larger spatial and temporal scales, via entrainment from the free troposphere (Carslaw et al., 2010).

The fundamental tenet of the CLAW hypothesis, of feedback between surface ocean biogeochemistry and climate, may be applicable via a broader spectrum of precursor species. Recent research has shown increasing complexity of potential aerosol source pathways, involving a variety of chemical species, processes and interactions (Vaattovaara et al., 2006). In addition to DMS, a variety of other gaseous aerosol precursors that originate from phytoplankton, bacterial and photochemical sources at the sea surface may undergo physical and chemical transformation to produce new particles in the MBL (Ciuraru et al., 2015). These SMA precursors include volatile organic species, such as carboxylic acids, isoprene, monoterpenes, halocarbons, iodine oxides and iodine (Vaattovaara et al., 2006; Sellegri et al., 2005). A biological source of these SMAs has been inferred from the spatial and temporal correlation between phytoplankton blooms and cloud microphysics (Meskhidze et al., 2009; Meskhidze and Nenes, 2010; Lana et al. 2012). The presence and concentration of SMA precursors in the MBL may be dependent

125 upon plankton abundance and community composition, and consequently their influence on aerosol formation will show spatial and seasonal variability (O'Dowd et al., 2004).

New particle formation may be suppressed by the interaction of aerosol precursors and SMAs with pre-existing aerosol, for example, by absorption of ammonia and gaseous sulfuric acid by coarse mode sea-salt aerosol (SSA; Cainey and Harvey, 2002). Conversely, existing particles may grow via condensation
130 which enhances their CCN capacity (Clarke et al., 2013). It has also been proposed that organic acids combine with sulfuric acid to create the critical nucleus required for aerosol formation (Zhang, 2010; Almeida *et al.* 2013). However, nucleation events over the open ocean remain elusive (O'Dowd et al., 2010; Chang et al., 2011; Willis et al., 2016), making it difficult to elucidate the primary pathways and reactants, and consequently they are currently regarded as of low significance to marine aerosol
135 formation. Following nucleation, the aerosol distribution is modified by aerosol-aerosol interaction, heterogenous reactions and removal processes, including coagulation and condensation, resulting in the longest-lived aerosol component being in the accumulation mode (0.06-0.4 μ m). With such a wide variety of potential precursors and inorganic/organic interactions affecting nucleation and CCN activation, the modelling of aerosols and their indirect influence on cloud radiative properties over the
140 remote ocean presents a major challenge (Seinfeld et al., 2016).

The production and transfer of aerosol precursors from the ocean surface is also dependent upon physical factors. Exchange across the air-sea interface is primarily controlled by near-surface turbulence, which is dependent on wind and waves. For practical purposes, this is represented by a kinetic factor, the transfer velocity k which is generated with wind-speed parameterisations (Nightingale et al., 2000;
145 Ho et al., 2006). Although wind-speed provides a reasonable broad-scale proxy for kinetic transfer, other factors such as fetch, wave development, wind-wave direction and surfactants also influence k , and so generate variation in gas exchange and deviation from k -wind-speed relationships. For example, most k -wind-speed parameterisations do not explicitly capture the solubility effects associated with bubbles (Blomquist et al. 2006), although the COAREG gas transfer model incorporates this factor into a
150 physically-based flux algorithm (Fairall et al., 2003; Fairall et al. 2011). Biogeochemical gradients near or at the ocean surface are also not considered, despite their potential to alter the air-sea exchange of gases, PMAs and SMAs (Facchini et al., 2008; Calleja et al., 2013).

Previous related research campaigns have examined the biogeochemical and physical factors influencing oceanic DMS and CO₂ fluxes, as summarised in Suppl. Table 1, but few have linked this to the physical
155 controls of air-sea exchange, and variation in aerosol and trace gas composition of the MBL. Similarly, other campaigns with an atmospheric focus, such as MAP (Decesari et al., 2011), have carried out

detailed studies of aerosol chemistry, but not interpreted this with regard to surface ocean biogeochemistry. To address this, the Surface Ocean Aerosol Production (SOAP) campaign was initiated, with the primary aim of characterising the variation in aerosol composition and concomitant marine sources, processes and pathways in the South-west Pacific. SOAP utilised a multi-disciplinary framework, encompassing surface ocean biology and biogeochemistry, transport and air-sea exchange with characterisation of aerosol number and composition, to establish controls on aerosols and gas exchange. The campaign consisted of two voyages - a pilot study, PreSOAP, which carried out a regional survey and established sampling strategies - and the following SOAP voyage, in biologically productive frontal waters along the Chatham Rise, east of New Zealand (see Figure 1). Building upon the approaches used in previous studies, the SOAP campaign targeted three phytoplankton blooms of differing plankton community composition, to determine their respective influences on biogeochemistry, gas exchange and MBL composition. The following paper details the regional context, sampling strategy, environmental conditions and some preliminary results for the SOAP campaign.

170

2. Regional context

The South-west Pacific has many features in common with the Southern Ocean, as it is characterised by low anthropogenic and terrestrial aerosol loading, long ocean fetch and high wind-speed, making it an optimal location for examining the marine contribution to aerosol production. One of the more biologically productive regions lies east of New Zealand, where the Sub-Tropical Front (STF) extends as a tongue of elevated phytoplankton production (Murphy et al., 2001), along 43.0-43.5°S over the Chatham Rise (see Figure 1a). This arises from the confluence of warmer saline subtropical waters that are relatively deplete in macronutrients, with fresher cooler subantarctic waters containing elevated macronutrients but depleted in iron (see Figure 1b; Boyd et al., 1999). Mixing across the front alleviates nutrient stress which, combined with a relatively stable water column, promotes primary production (Chiswell et al., 2013). Ocean colour climatologies show a monthly mean Chl-*a* of 0.6 mg m⁻³, reaching ~ 1 mg m⁻³ over the Chatham Rise in spring (Murphy et al., 2001), and the region is characterised by elevated marine particle export, secondary production and fish stocks (Nodder et al., 2007; Bradford-Grieve et al., 1999). In spring the phytoplankton community composition varies with water mass, with diatoms dominating the STF, cryptophytes, prasinophytes and dinoflagellates more prevalent in subtropical waters, and photosynthetic nanoflagellates dominating subantarctic waters (Chang and Gall, 1998; Delizo et al., 2007). The STF also supports spatially-extensive coccolithophore blooms (Sadeghi et al., 2012), and is situated on the northern edge of the “Great Calcite Belt” (Balch et al., 2011), a

185

latitudinal band of elevated backscatter attributed to coccolithophore liths. Surface mixed layer
190 nutrients vary spatially in response to mixing of the water masses and seasonally due to phytoplankton
uptake, with the evolution of nutrient stoichiometry and grazing determining the succession and
duration of different phytoplankton blooms (Chang and Gall, 1998; Delizo et al., 2007). The STF is
characterised by significant gradients in $p\text{CO}_2$ associated with phytoplankton blooms, with current global
climatologies indicating the region east of New Zealand as a significant carbon sink ($>1\text{mol C m}^{-2}\text{ yr}^{-1}$,
195 Landschuetzer et al., 2014).

The waters south of New Zealand are characterised by high wind-speeds which drive the
disproportionate contribution of this region to global ocean CO_2 uptake. Here, wind, waves and currents
develop unhindered by land, and strong persistent westerlies act over long fetch to generate large swells
that propagate north-east influencing the wave-climate off New Zealand. While this wave energy is
200 attenuated closer to land in the eastern Chatham Rise, the average wave energy is still 75% of values
south of New Zealand where annual mean wave heights exceed 4m. Subantarctic waters south of the
Chatham Rise region provided a prime location for a dual tracer release experiment (SAGE; Harvey et
al., 2011), aimed at constraining k at high wind-speeds. Comparison of the SAGE k -wind-speed
parameterisation with those generated in other regions, and using different techniques, showed
205 generally good agreement (Ho et al., 2006); this may be interpreted as indicating that regional influences
on exchange may be less important, supporting the application of a universal wind-speed
parameterisation. Nevertheless, other factors, such as wave age, duration and height do influence gas
exchange in this region (Smith et al., 2011; Young et al., 2012). The elevated winds also influence the
transfer of aerosols and precursors, as reflected by a zonal band of elevated sea spray aerosol mass and
210 water-insoluble organic matter over the Chatham Rise region (Vignati et al., 2010).

Both models and measurements indicate that DMS is a significant contributor to total non-sea salt
sulfate (nssSO_4) in the Southern Hemisphere (Gondwe et al., 2003; Korhonen et al., 2008). However, a
paucity of observational data in the Southern Ocean has hindered development of global climatologies
for surface seawater DMS (DMS_{sw}), with the region south-east of New Zealand represented by only a
215 few data points in a recent DMS climatology (Lana et al., 2011). Despite this shortcoming, this
climatology provides a realistic representation of atmospheric DMS and total sulfate when applied in
aerosol-climate Global Climate Models, particularly over the Southern Ocean (Mahajan et al., 2015).
Seasonal variability in atmospheric DMS is apparent at stations in New Zealand and south of 44°S (Blake
et al., 1999), with concentrations of 100-200 pptv, and maximal values associated with the transport of
220 DMS from waters to the south in summer (Harvey et al., 1993; de Bruyn et al., 2002; Wylie and de Mora,
1996). Corresponding seasonality in nssSO_4 was observed, with a maximum ($0.8\text{-}1.5\text{ }\mu\text{g m}^{-3}$) in early

austral summer at the start of the year, decreasing in late summer to 0.1-0.4 $\mu\text{g m}^{-3}$ through autumn and winter (see Figure. 2; Sievering et al., 2004; Allen et al., 1997). For comparison, coarse SSA dominates the aerosol mass at Baring Head, with concentrations of 6-10 $\mu\text{g m}^{-3}$ (Jaeglé et al., 2011; Spada et al., 2015). Similar seasonal cycles of DMS and nssSO₄ were recorded at Cape Grim (Ayers, 1991), and the observed diurnal inverse correlation between sulfur dioxide and DMS at Baring Head was applied to estimate yield and potential contribution to aerosols (de Bruyn et al., 2002). Consistent seasonal trends between activated particles and cloud droplet number concentration were also apparent, with a summer maximum over the Southern Hemisphere (Boers et al., 1996; 1998), related to phytoplankton production (Thomas et al., 2010). Overall, the temporal trends in aerosol precursors and pathways do not follow that of wind-speed and other physical drivers, but instead reflect biological processes inferring control by surface ocean biogeochemistry (Korhonen et al., 2008).

3. Research Programme and Strategy

3.1 PreSOAP

A pilot study, PreSOAP, was carried out to test technical approaches and confirm the regional source of biogenic aerosols in the Chatham Rise region on the New Zealand research vessel, *Tangaroa*, on 1-12/2/2011 (DoY32-42). The strategy of bloom location using satellite imagery and subsequent mapping of surface properties proved successful, with three blooms of differing DMS_{sw} and pCO₂ signatures located and monitored each for 3-4 days. The first bloom was initially dominated by dinoflagellates with an increase in diatom biomass after 3 days, while the second and third blooms were primarily dominated by coccolithophores and dinoflagellates, respectively. This variability in species composition resulted in significant spatial and temporal variability in DMS concentrations in the MBL (DMS_a) and DMS_{sw}. DMS_a concentration varied over two orders of magnitude, reaching 1000 ppt on DoY 36 (see Figure 3b), similar in range to that recorded at the Baring Head station near Wellington (Harvey et al., 1993; de Bruyn et al., 2002). There was no significant correlation between DMS in the two phases, with DMS_a showing a stronger relationship with wind-speed (see Figure 3). Surface Chl-*a* concentrations reached 2 mg m⁻³, but there was no significant relationship between DMS_{sw} and Chl-*a*, with the DMS_{sw} maximum of ~10 nmol L⁻¹ during the first bloom coinciding with Chl-*a* of ~1 mg m⁻³ (Figure 3d). The observed temporal and spatial variability in DMS_a and DMS_{sw} during PreSOAP highlighted the technical challenge of establishing relationships between surface ocean biogeochemistry and atmospheric composition. Provisional method development was also carried out for measurement of DMS and other parameters in near-surface waters and the sea surface microlayer (SSM).

Surface DMS_{sw} and pCO₂ were mapped, and DMS_a and CO₂ MBL concentrations and fluxes measured
continuously by sensors and collectors mounted on the bow of the vessel. Testing of the eddy covariance
(EC) flux technique identified an issue with water vapour interference that dominated the CO₂ signal
recorded by an open-path InfraRed Gas Analyser (IRGA). Preliminary studies also identified that residual
ship motion dominated over turbulence for the real-time switching of Relaxed Eddy Accumulation
measurement of flux under high swell conditions. The logistical challenges of flux measurement at
distance from the vessel were also assessed by deployment of a free-floating catamaran supporting a
mounted gradient flux sampling system (Smith et al., to be submitted). A temperature microstructure
profiler was also deployed to record near-surface temperature and turbulence structure (Stevens et al.,
2005), although this was limited to short sampling periods, highlighting the need for a mounted
thermistor array on a spar buoy for longer measurement coverage.

The utility of a baseline sector for sampling MBL composition, using relative wind direction and speed,
was also tested during PreSOAP. Measurements showed a tendency for higher condensation nuclei
concentration in the “non-baseline” sector, confirming the utility of this approach (Harvey et al., to be
submitted). A common aerosol inlet provided clean air from a height of 17.5m above sea level to
instruments and sensors in a container laboratory on deck. Particle size distribution and concentration,
including ultrafine nuclei concentrations, were continuously monitored using a scanning mobility
particle sizer (SMPS) and optical particle counters (OPC’s), with bulk ion chemistry samples collected
using a high-volume sampler. The composition of primary marine aerosols was also examined using a
0.45m³ bubble chamber, in which sea spray was formed via the bursting of bubbles produced by passing
clean compressed air through sintered glass (Mallet et al., 2016).

3.2 The SOAP voyage

The SOAP voyage employed the strategy successfully piloted on PreSOAP, of identifying phytoplankton
blooms in NASA MODIS Aqua and Terra satellite ocean colour images, with subsequent bloom location
and mapping using a suite of underway sensors (Chl-a, β_{660} backscatter, pCO₂, DMS_{sw}). The blooms were
discrete and coherent areas of elevated ocean colour, that were provisionally characterised by a
concentration of 1 mg/m³ Chl-a or higher. For each bloom, a nominal centre was identified, based upon
maximum DMS_{sw} and Chl-a concentrations, and marked by deployment of a Spar Buoy. Repeat
activities at the bloom centre included characterisation of the surface mixed layer by vertical profiling,
collection of SSM samples at distance from the main vessel, and gradient flux on a catamaran. Overnight
mapping was carried out to determine changes in bloom magnitude and position. Sampling also took

place at stations on the periphery and outside the blooms, as defined by distance from the bloom centre and a clear demarcation in surface biogeochemical variables. The SOAP voyage was nominally divided into three different bloom periods (see Figure 4), with an initial dinoflagellate bloom (B1) located 12 hours into the SOAP voyage that exhibited elevated Chl-*a* and DMS_{sw}, and pCO₂ drawdown, a
290 coccolithophore bloom (B2) with initially moderate signals that weakened, and a final bloom (B3) of mixed community composition. Following a storm the surface water column structure and biogeochemistry were significantly different, and so this bloom was subdivided into B3a and B3b.

3.2.1 Environmental conditions during the SOAP voyage

Back-trajectory analysis of particle density was calculated for each bloom using the Lagrangian
295 Numerical Atmospheric-dispersion Modelling Environment (NAME) for the lower atmosphere (see Figure 5). The meteorological situation evolved over the SOAP voyage from a high-pressure system with light winds during B1, to stronger winds during B2 and B3. The main weather features included a depression crossing the central South Island on DoY 54-55 during B2, and a second depression from the east from DoY 58 onwards. During B3 a vigorous front advanced up the east coast of the South Island
300 on DoY 61 with strong SW winds of 20 m s⁻¹, followed by a depression crossing the lower North Island on DoY 63 that maintained a fresh southerly airflow for the remainder of the voyage. Air and water temperatures during B1 were generally similar indicating near-neutral stability, whereas B2 experienced a period of warm, moist air and reversal in direction of turbulent heat fluxes, followed by a short period when air temperatures were 2-3°C higher on DoY 56-58 (See Figure 6). Waves were dominated by swell
305 from the south-southwest, with significant wave height mirroring trends in wind-speed, reaching a 5m maximum during the localised storm on DoY 61 (see Figure 6). Wave parameters obtained from NOAA WaveWatch III analyses indicated that wave height was 23% lower during B1 and B3, and 13% lower at B2, relative to wave height south of New Zealand at 50°S.

Table 1 summarises the hydrographic and biogeochemical characteristics in the surface mixed layer of
310 the three phytoplankton bloom regions. B1 was a large dinoflagellate bloom with high surface DMS_{sw} (maximum ~30 nmol L⁻¹; mean 16.8 nmol L⁻¹; Bell et al., 2015), and Chl-*a* (maximum 3.4 mg m⁻³), and significant CO₂ undersaturation with a mean surface pCO₂ of 320 ppmv (see Table 1). B1 was located, south of the Mernoo bank, a deep channel between the western end of Chatham Rise and the east coast of the South Island. This region has been previously identified as a prime location for phytoplankton
315 blooms, due to eddy-driven mixing and flow reversals arising from current and topographic interaction, which enhance iron and nutrient supply (Boyd et al., 2004). During B1, winds remained light (see Table 1) with a calm sea state, and the spar buoy drifted north-east primarily under the action of surface

currents. Solar irradiance was high and a shallow surface mixed layer developed (see Figure 6), with a significant near-surface temperature gradient (Walker et al., 2016). Mean nitrate and phosphate concentrations (5.3 and 0.4 $\mu\text{mol L}^{-1}$, respectively) were sufficient for phytoplankton growth, whereas silicate was low (see Table 1), and close to growth-limiting concentrations (Boyd et al., 1999). Although dinoflagellates dominated, coccolithophores biomass was higher at some stations, and nanoeukaryote abundance was generally low. B1 was occupied for 5-6 days, during which broader regional excursions with overnight mapping identified a bloom of high Chl-*a* but relatively low DMS_{sw} to the south-west.

The vessel re-located to a coccolithophore bloom, B2, evident at the eastern end of the Chatham Rise in MODIS true colour satellite images (see Figure 4b). Upon arrival on DoY 52 B2 showed an initial mean DMS_{sw} of 9 nmol L⁻¹ and elevated Chl-*a*, and was characterised by a relatively warmer, shallower, saltier surface mixed layer of lower nitrate concentration (compared to B1, see Table 1), typical of subtropical water. This appeared to provide optimal conditions for coccolithophores as surface water backscatter (β_{660}) was initially elevated by high lith abundances, with coccolithophores accounting for up to 40% of phytoplankton carbon. However, intrusion of warm, moist air associated with north-westerly winds, coincided with a reversal in the direction of turbulent heat fluxes, and was followed by a southwest wind shift strengthening to 17 m s⁻¹ by DoY 56 (see Figure 6). This resulted in deepening and cooling of the surface mixed layer with a corresponding increase in nutrient concentrations which, combined with a decrease in solar irradiance, resulted in a decline in Chl-*a* and DMS_{sw} (Bell et al., 2015).

Following the 5-day occupation of B2, the vessel returned to the south of Mernoo bank to assess a bloom that had developed near the original site of B1. Surface biogeochemical signals were initially weak in B3a, with a mixed community of coccolithophores and dinoflagellates and low DMS_{sw} (2.2 nmol L⁻¹) and Chl-*a* (mean 0.39 mg m⁻³). However, an intense front advanced up the South Island and resulted in strong SW winds that exceeded 20 ms⁻¹ (see Figure 6), after which mixed layer depth and associated nutrients increased. Consequently, stations before and after the storm were physically and biogeochemically disparate. B3a stations exhibited similar sea surface temperature to B1, but with a deeper surface mixed layer and a Chl-*a* half that of B1, whereas B3b stations were significantly cooler (at 13°C) and deeper (41m) than B1 (see Figure 7), with higher silicate concentration due the enhanced vertical mixing. Subsequent stabilisation of the surface mixed layer by light winds combined with elevated nutrients stimulated Chl-*a*, diatom and coccolithophore abundance in the final B3b stations (see Figures 6 and 7).

4. SOAP work programmes and observations

350 A number of parameters were measured (see Table 2) in three interlinked work programmes during the SOAP voyage, as indicated in Figure 8 and detailed below.

4.1. The distribution and composition of aerosols, precursors and trace gases in the MBL

Aerosol number concentration, size distribution, composition, water uptake and CCN concentration were measured semi-continuously during SOAP to address the overall paucity of aerosol observations, and the apparent rarity of nucleation events, over the remote ocean. These were characterised by a suite of instruments covering a particle size range of 0.01 to 10 μm (see Figure 9 and Table 2), which enabled determination of the size-dependent contribution of PMA and nssSO_4 to aerosol and CCN concentrations. Aerosol characterisation identified variable Aiken and consistent sub-micron sized accumulation and coarse modes, with the sub-micron aerosol mass dominated by secondary aerosol with ammonium sulfate/bisulfate under light winds, and with an increase in sea-salt proportion as local winds increased. Ongoing data analysis is examining whether significant nucleation events occurred.

The operational mode for underway aerosol measurement was to slowly steam at 1–2 kts into the prevailing wind, across an area of high biological productivity or significant air-sea gas gradient, generally between noon and 2:00PM when solar irradiance was maximal. The common aerosol inlet developed during PreSOAP allowed uncontaminated air from above the bridge to be sampled when the wind was on the bow, so minimizing interference from ship stack emissions. Contamination events were screened out using a real-time clean-sector sampling “baseline” flag and switch (Harvey et al., to be submitted), enabling clean collection of integrated samples. Although the vessel exhaust was the primary contaminant, other potential sources included the workboat and recirculation of polluted air around the ship, and longer range terrestrial influences were also assessed. Measurements of black carbon using an aethalometer, and CO_2 by high precision Cavity Ring-Down Laser Spectroscopy (CRDS) provided two independent variables for detecting contamination events, and some VOCs, measured by PTR-MS (see Table 2), were also used as indicators of diesel combustion. The vessel was orientated into the wind as often as possible, which resulted in a high frequency (~75%) of baseline sector conditions during the SOAP voyage. Clean marine air periods were defined post-voyage using the baseline wind sector ($225 - 135^\circ$ relative to bow and wind speed greater than 3 m s^{-1}), black carbon concentrations (less than 50 ngm^{-3}), and back trajectories indicating minimal terrestrial impact (periods when the minimum number of hours over land in 72-hour back trajectory is zero), with periods of workboat operations also removed. An ensemble of Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPPLIT) model back trajectories (Draxler and Rolph, 2013) was run for each hour of the voyage, and NAME back trajectories calculated for every three hours (Figure 5, Jones et al., 2007). Figure 10 shows

particle number and CCN concentrations, compared to the number of hours the 72-hour back trajectory spent over land calculated from HYSPLIT trajectories. Particle concentrations were generally higher during periods of terrestrial influence (see DoY 52 and 60, Figure 10), with average particle number concentrations of $1122 \pm 1482 \text{ cm}^{-3}$, double that observed for clean marine air. Ion beam analysis also revealed the presence of silicate and aluminium on ambient submicron filter samples suggesting a terrestrial source, supporting the back-trajectory modelling of continental outflow.

During the initial occupation of B1 under light winds, the particulate matter (PM₁₀) total ion mass was $9.5 \text{ } \mu\text{g m}^{-3}$ compared to subsequent samples under higher winds in the range $20\text{--}50 \text{ } \mu\text{g m}^{-3}$. The dominant components of the inorganic mass fraction were sea-salt ions and nssSO₄, although a measurable organic fraction was also present (see below). The NaCl mass in light winds during B1 was $6.6 \text{ } \mu\text{g m}^{-3}$ with >95% of > $3 \mu\text{m}$ diameter, relative to $32.5 \text{ } \mu\text{g m}^{-3}$ under stronger winds during B3b. Although 72% was > $3 \mu\text{m}$, the largest difference in mass occurred in the 1.5 to $3 \mu\text{m}$ size range. In contrast, the mass of nssSO₄ was predominantly sub-micron sized; B1 exhibited the largest nssSO₄ mass at $2.0 \text{ } \mu\text{g m}^{-3}$ with 85% in sizes < $1 \mu\text{m}$, whereas in B3b, the nssSO₄ mass was much lower at $0.6 \text{ } \mu\text{g m}^{-3}$ with 76% in < $1 \mu\text{m}$ sizes. These results confirm the influence of both physical and biogeochemical processes on aerosol composition.

Voyage particle number concentrations during clean marine periods averaged $534 \pm 338 \text{ cm}^{-3}$, with CCN concentrations of $178 \pm 87 \text{ cm}^{-3}$ ($\pm 1 \text{ sd}$) at 0.5% supersaturation, and an average particle fraction activated into CCN of 0.4 ± 0.2 . Bloom average particle number concentrations ranged from a minimum of $385 \pm 96 \text{ cm}^{-3}$ in B3b to a maximum $830 \pm 255 \text{ cm}^{-3}$ at the start of B2 (Figure 10). B1 displayed the highest CCN activation ratio, of 0.5 ± 0.2 , potentially the combination of low wind speeds, large biogeochemical signals and SMA fluxes. Comparison of the inorganic ion mass determined from high-volume sampler filters between the different blooms does not support the conclusion that the B1 activation ratio was higher simply because particles were larger. As the median particle diameters during clean marine periods were consistent between the three blooms, this suggests that particle composition, secondary organics or coagulation may have impacted CCN activation at B1. These findings are supported by preliminary results from an application of the ACCESS-UKCA model (Woodhouse, pers comm.), which simulated the additional impact of emissions of marine secondary organic carbon under the conditions determined during SOAP. In contrast, the average CCN activation ratio for B3a was 0.13 ± 0.06 . Nucleation mode particles (10 nm and 15 nm), were measured by ultra-fine organic tandem differential mobility analyser (UFO-TDMA, Vaattovaara et al., 2005), and Aitken mode particles (50 nm), by UFO-TDMA and a volatility and hygroscopicity tandem differential mobility analyser (VH-TDMA, Johnson et al. 2004a; Villani et al., 2008). This analysis typically identified a significant (up to 50% volume

fraction) secondary organic component during sunny conditions in bloom regions, particularly during B1. The TDMA results provided further evidence for secondary organic aerosol processing of the dominant secondary nssSO₄ mode during B1. Deliquescence measurements (VH-TDMA) indicate that the Aitken mode population is largely comprised of neutralised nssSO₄ i.e. ammonium sulfate. Small and sporadic contributions to the Aitken mode from a non-hygroscopic (number fraction up to 0.4) and a highly hygroscopic component (number fraction up to 0.3) were observed in addition to the secondary nssSO₄ mode (number fraction of 0.6 - 1). The water uptake and volatility of the sporadic highly hygroscopic mode indicates this may be composed of PMA.

The *in-situ* aerosol size, number and composition measurements in the MBL were complemented by *in vitro* chamber measurements of nascent SSA, to determine the PMA organic volume fraction and water uptake properties. Nascent SSA filter samples were analysed using Fourier Transform InfraRed spectroscopy (FTIR) for organic functional groups (Russell et al. 2011), and ion beam analysis for inorganic concentrations (Cohen et al. 2004). Measurements of the hygroscopic growth factor and the volatile fraction up to 450°C for 50-150 nm particles using the VH-TDMA were compared with those of reference inorganic samples (e.g. sea salt, ammonium sulfate) to determine their organic volume fractions (Modini et al. 2010). Complementing the VH-TDMA, the UFO-TDMA provided further information on the organic content of particles of 50nm and down to 10 nm. The bubble chamber observations indicate that the PMA contained a substantial primary organic fraction. VH-TDMA results indicate that the Aitken mode PMA was primarily non-volatile (78-93%), with an average organic volume fraction of 51% (ranging from 39 to 68%), and the UFO-TDMA results show an OVF ranging from 35-45%. These results are consistent with observations in the North Pacific and Atlantic, for which an Aitken mode volatile fraction of the order of 15% and OVF of 0.4-0.8 have been observed (Quinn et al. 2014). FTIR analysis indicated that the POA aerosol in the chamber experiments was largely composed of hydroxyl functional groups, with minor contributions from alkanes, amines and carboxylic acid groups, consistent with previous PMO observations (Russell et al. 2011).

Although DMS was a primary focus of measurements during SOAP, a wide variety of other VOCs that potentially contribute to secondary organic aerosol formation were also measured. Halogens and halogen oxides were measured using Multi Axis Differential Optical Absorption Spectroscopy (Max-DOAS) and Electron Capture Detector-Gas Chromatography (ECD-GC). Iodine has been identified as a potentially important precursor of nucleation in coastal regions (Sellegrì et al., 2005), and SOAP provided an opportunity to relate the presence of halogen oxides to phytoplankton biomass and composition in the surface ocean, and nucleation events in the MBL. A High Sensitivity Photon Transfer Reaction Mass Spectrometer (PTR-MS) measured continuously in H₃O⁺ mode in the range of m/z 21- m/z 155

throughout the voyage (Lawson et al., to be submitted). Aldehydes, ketones and dicarbonyls were measured with 2,4-dinitrophenylhydrazine (2,4-DNPH) cartridges and high performance liquid chromatography (HPLC; Lawson et al., 2015), and a range of VOCs were sampled using adsorbent tubes and later analysed via Thermal Desorption-Gas Chromatography - Flame Ionisation Detection - Mass Spectrometry Detection (TD-GC-FID/MS). These measurements identified a positive relationship between DMS (m/z 63), acetone (m/z 59) and methanethiol (m/z 49), indicating common biological drivers (Lawson et al., to be submitted).

The first in situ measurements of aqueous phase SMA precursors dicarbonyls, glyoxal and methylglyoxal were obtained over the remote Southern Ocean during SOAP (Lawson et al., 2015). Parallel measurements of known dicarbonyl precursors, measured by PTR-MS, were used to calculate the expected yields of glyoxal and methyl glyoxal, which accounted for < 30% of observed mixing ratios indicating an unidentified source of dicarbonyls (Lawson et al., 2015). This was corroborated by inclusion of SOAP glyoxal measurements obtained by Max-DOAS measurement in a global database, which concluded that the missing glyoxal source was an order of magnitude greater than identified sources (Mahajan et al, 2014). Surface mixing ratios of glyoxal converted to vertical columns, were significantly lower than average vertical column densities (VCDs) from satellite retrievals, possibly reflecting the difficulty of retrieving low glyoxal VCDs over the ocean, or incorrect assumptions about the vertical distribution of glyoxal in the atmosphere (Lawson et al., 2015).

4.2. Rates and controls of volatile and precursor emissions at the air-sea interface

DMS measurements were made using three different instruments during SOAP (see Table 2); an Atmospheric Pressure Ionisation-Chemical Ionisation Mass Spectrometer (API-CIMS) continuously monitored DMS in both phases (Bell et al., 2015), a PTR-MS monitored DMSa (Lawson et al., to be submitted), and discrete water measurements were made using a Sulfur Chemiluminescence Detector Gas Chromatograph (SCD-GC; Walker et al., 2016). Intercomparison of sulfur measurements is not easily or routinely performed (Bell et al., 2012), particularly at sea. Seawater DMS measurements (CIMS and SCDGC) compared well during SOAP (Walker et al., 2016) and the SCD-GC technique also compared well with traditional gas chromatography (with flame photometric detector) in an international intercalibration exercise (Swan et al., 2014). Intercomparison of the PTR-MS and SCD during SOAP, involved analysis of two air samples and two diuted DMS gas standards with a concentration range of 158 – 354 ppt. The instruments showed very good agreement, with a mean difference of 5% and maximum 10%.

Although the majority of DMS flux estimates to date have been derived by applying an independently
480 determined transfer velocity (k) to the measured DMS gradient at the ocean surface (ΔDMS), there has
been a recent increase in direct micrometeorological measurements of DMS flux. Measurements at 10-
30-minute resolution show considerable variability in flux, which may reflect methodological artefacts
or inherent variability in the distribution of DMS. SOAP provided a platform for comparing eddy
covariance (EC) flux measurements of DMS using API-CIMS (Bell et al., 2015), with a gradient flux
485 technique using a drogued catamaran within one kilometre of the vessel (Smith et al., to be submitted).
The gradient flux technique is less direct than EC but provides an alternative reference on a platform
that is relatively free of shipboard air-flow distortion. The EC system sampled from an intake on the
ships bow, with flux instruments mounted on the foremast 12.6m above sea level, and the air pumped
to a containerised laboratory on the foredeck. Additional meteorological measurements were obtained
490 from a weather station above the bridge. Both sites are subject to airflow distortion which is azimuthally
dependent (Popinet et al., 2004). The catamaran sampling framework, which consisted of four air
intakes distributed vertically on a 5.3m mast, sampled closer to the water surface where gas gradients
are largest. Flux measurements were augmented by continuous near-surface measurement of physical
parameters using a range of sensors attached to a Spar Buoy, with stratification determined by
495 temperature sensors at 0.5 m intervals (Walker et al., 2016), turbulence determined by a Vector acoustic
Doppler Velocimeter at 0.6 m depth. This permitted comparison of k_{DMS} estimates with near-surface
upper-ocean turbulence at a distance from the vessel (Smith et al., to be submitted). Wave-breaking
whitecap coverage was monitored using a Campbell Scientific 5-megapixel camera (cc5mpx) located on
the starboard side of the vessel (Scanlon and Ward, 2016). This provided an indicator of bubble
500 entrainment, which contributes to the differential transfer rate of DMS and CO_2 due to their different
solubilities (Blomquist et al., 2006; Bell et al., 2017).

Although SOAP primarily focussed on DMS fluxes, EC measurements of CO_2 flux were an important
adjunct measurement for providing insight into gas exchange mechanisms and controls, and improving
gas transfer algorithms for gases of differing solubilities. Four Licor infrared gas analyzers were used for
505 eddy covariance flux measurements of CO_2 during SOAP, following the initial trials on PreSOAP.
Comparison of EC measurements with wet and dry incoming gas streams, and an empirically-based post-
processing correction, indicated that only gas stream drying produced robust CO_2 flux and k_{CO_2} estimates
(Landwehr et al., 2014). A detailed examination of ship motion and airflow distortion effects resulted in
a significant reduction of the scatter in the CO_2 eddy covariance data (Landwehr et al., submitted). The
510 EC-derived k_{CO_2} estimates provided a better correlation with a linear fit to the EC friction velocity than
with the 10-metre neutral wind speed ($u_{10\text{N}}$), and showed good agreement with dual tracer-derived

estimates from the SAGE experiment conducted in this region in March-April 2004 (Ho et al, 2006). Measurement of DMS and CO₂ fluxes also provided further constraint of k parameterisations based upon wind-speed, and the opportunity to assess the influence of bubbles on gas exchange at high wind speeds. DMS fluxes derived by EC and gradient flux techniques showed good agreement (Bell et al., 2015; Smith et al., to be submitted), and confirmed previous observations that gas transfer is a linear function of wind-speed at low to intermediate winds (Blomquist et al., 2006; Yang et al., 2011). However, despite winds reaching 20 m s⁻¹ during the latter part of SOAP insufficient data was obtained to draw conclusions regarding the reported deviation of k_{DMS} under high winds (Bell et al., 2015). However, SOAP provided a novel estimate of the size of the EC flux footprint, and the temporal-spatial mismatch between DMS_{sw} and shipboard measured fluxes, highlighting the importance of considering skew in flux estimates arising from non-linear distribution of DMS_{sw} (Bell et al., 2015).

A further objective of SOAP was comparison of measured DMS fluxes with calculated estimates from the COAREG model (Fairall et al. 2011) based on ΔDMS , to assess potential discrepancies with modelled fluxes (Marandino et al., 2008; Walker et al., 2016). Potential factors examined here included air and water stability, and the influence of the SSM. Despite the agreement between DMS flux estimates by the two micrometeorological techniques, there was significant departure from COAREG predictions (Fairall et al., 2011) on occasions, suggesting the influence of unidentified processes (Smith et al., to be submitted). One potential example was the suppressed DMS flux during a period of atmospheric stability and reversed heat flux during B2. Concurrent EC flux measurement for DMS and CO₂ also provided an opportunity to assess other influences on k . The DMS flux data indicate that the k_{DMS} -wind speed relationship was relatively insensitive to surface biogeochemistry or wave action during SOAP (Bell et al., 2015). In addition, SOAP data was used to parameterise whitecap coverage against wind-speed, and identify that maturing waves may obscure and lead to underestimate of the variability of breaking waves (Scanlon and Ward, 2016).

4.3. Surface ocean biogeochemical influences on aerosols and volatiles

Surface mapping of DMS_{sw} and pCO₂, using API-CIMS and IRGA, respectively (Bell et al., 2015) were critical to the SOAP voyage strategy and the aims of the two workpackages discussed above. These measurements also provided insight into the covariance of DMS sources and CO₂ sinks in surface waters, and established the importance of this region to global budgets. The New Zealand Coastal province (NEWZ), which includes the frontal region (STF) studied during SOAP, is characterised in the global DMS climatology by year-round low DMS concentrations with a maximum <2 nmol L⁻¹ (Lana et al., 2011). This infers that this region has some of the lowest global DMS_{sw} concentrations, in marked contrast to the adjacent South Subtropical Convergence (SSTC) province, which occupies the remainder of the 35-50°S

latitude band and accommodates the STF, which is characterised by a mid-summer maximum of 10 nmol L⁻¹ DMS. This discrepancy between the two regions likely reflects the low number of DMS observations for the NEWZ province in the climatology (n=6; Lana et al., 2011). Previous DMS_{sw} measurements in subantarctic waters south of the Chatham Rise, and east of Tasmania in the SSTC biome (Archer et al., 2011; Griffiths et al., 1999), are consistent with this climatological estimate, whereas larger unpublished surveys have recorded elevated surface DMS_{sw} during austral spring (October 2000), with a mean DMS_{sw} of 4.5 (+/- 6.8) nmol L⁻¹ on the Chatham Rise (Harvey et al., pers. comm). Combining these measurements with data from the SOAP campaign (mean DMS_{sw} = 6.6 nmol L⁻¹) gives a weighted-mean DMS_{sw} of 5.3 nmol L⁻¹ (n=5300, see Table 3), confirming that DMS_{sw} in the NEWZ province is currently underestimated, and is in fact more typical of the SSTC province. Although the PreSOAP and SOAP sampling strategy of focussing on phytoplankton blooms may introduce bias towards higher DMS_{sw}, the BOX voyage, which had broad spatial coverage of subtropical and subAntarctic waters between 39.5-47°S, gave a similar mean DMS_{sw} to the weighted mean for all voyages. The elevated DMS_{sw} was reflected in the EC flux measurements during SOAP, which recorded maximum and mean fluxes of 100 and 16.3 μmol S m⁻²d⁻¹, respectively, (Bell et al., 2015), which exceed the climatological mean of >10 μmol S m⁻² d⁻¹ for the SSTC region (Lana et al., 2011). In addition, the high MBL DMS concentrations of 1000 ppt recorded during SOAP exceed DMS_a at coastal stations on the New Zealand North Island in summer (Harvey et al., 1993; de Bruyn et al., 2002; Wylie and de Mora, 1996). Although seasonally constrained, the SOAP measurements provide evidence that regional DMS emissions are significant in the South West Pacific. The large dataset of regional concentrations and flux will allow further refinement of global climatologies, such as the Global Surface Water DMS Database and the Surface Ocean CO₂ Atlas (SOCAT).

The spatial variability of DMS_{sw} was related to surface ocean biogeochemistry and bloom type by measurement of a suite of ancillary parameters in underway mode, including temperature and salinity, Chl-*a*, chromophoric dissolved organic matter (CDOM), β₆₆₀ backscatter, dissolved oxygen and pCO₂ (see Tables 1 and 2). The vertical variability of DMS_{sw}, and the dissolved and particulate pools of its precursor DMSP, were quantified in the surface mixed layer at stations within each bloom, and related to plankton biomass and community composition, nutrient and organic composition and physical drivers (see Suppl. Table 2). Process studies of DMSP cycling included deckboard incubations examining the bacterially-mediated pathways of DMSP cleavage and demethylation in relation to different bloom dynamics (Lizotte et al., submitted). DMSP concentrations were relatively high, reaching a maximum of 160 nmol L⁻¹, and showed significant correlation with phytoplankton biomass during SOAP. However, the yield of DMS from bacterial conversion of dissolved DMSP was variable with no spatial trend, although a

correlation with leucine incorporation indicates that DMSP was an important carbon source for bacteria. Overall, gross DMS production by bacteria in deckboard incubations of near-surface water was relatively low, at $< 6 \text{ nmol L}^{-1} \text{ d}^{-1}$, inferring that phytoplankton-mediated conversion of DMSP was likely a significant near-surface source of DMS (Lizotte et al., submitted).

The SSM is a potentially important interface controlling MBL and aerosol composition, as it is the interface across which material exchanges between atmosphere and ocean. Physical and biogeochemical processes within this thin layer have the potential to alter transfer via factors, such as the concentration of organic material and enhanced biological and photochemical processing. Near-surface CO_2 gradients have been observed (Calleja et al., 2005), and several studies report DMS enrichment in the SSM (see summary in Walker et al., 2016). If DMS consumption or production in the SSM is significant this represents a potential source of discrepancy in comparison of measured fluxes with that calculated by the COAREG model (see above). The biogeochemistry of the SSM and the upper 1.6m surface water were characterised at 10 stations during SOAP at distance from the research vessel, to determine the spatial variability in composition within, and between, different phytoplankton blooms (Walker et al., 2016). Near-surface DMS gradients were generally negligible, except during B1 where low wind-speed, near-surface stratification and high dinoflagellate abundance may have combined to enhance DMS in the SSM relative to subsurface waters. The observed DMS enrichment factors in the SSM during B1, ranging from 1.4 to 5.3, are some of the highest reported to date. The anomaly between measured DMS fluxes and COAREG estimated was also greatest during B1, inferring that DMS emissions, and associated k-wind-speed parameterisations, may be sensitive to DMS production in the SSM under certain conditions. However, the observations also raise questions as to how such significant DMS enrichment is maintained in the SSM, as high DMS production would be required to ameliorate loss processes (Walker et al., 2016).

5. Conclusions

The SOAP voyage has identified new questions in important areas of SOLAS-related research, including the influence of the SSM on DMS emissions, implications for secondary aerosol formation, and unidentified sources of organic aerosol precursors, all of which are potentially influenced by photochemistry in the surface ocean and MBL (Lawson et al., 2015). It has also addressed confounding technical challenges including small-scale heterogeneity in surface waters, clean air baseline sampling, and discrepancies between existing techniques and models. An overarching aim of the SOAP campaign was to assess potential relationships between surface water biogeochemistry and corresponding or

610 related species in the MBL, to identify the factors influencing aerosol precursors, and their potential as analogues. Chl-*a* is an indicator of phytoplankton biomass that is readily retrievable by satellite, and consequently has been investigated as a potential proxy for DMS_{sw} (Lana et al., 2011). The SOAP voyage provided a platform to validate this observation, particularly as it took place in the 40-60°S latitude band which exhibits the most significant regional correlation between Chl-*a* and DMS_{sw} (Vallina et al., 2006).
615 Overall there was a weak, but significant, correlation ($r = 0.12$, $p < 0.005$) between Chl-*a* and DMS_{sw} in the underway surface data during SOAP, but also significant variability in the slope and the sign of this relationship between the different blooms. Correlations were also apparent between Chl-*a* and DMSP (Lizotte et al., submitted), and Chl-*a* and DMS_a, but there was no relationship between Chl-*a* and DMS flux, as expected, due to the short timescales and flux footprint identified by Bell et al., 2015.
620 Correlations have been reported previously for Chl-*a* with CCN (Meskhidze and Nenes, 2006), and aerosol organic enrichment (Gantt et al., 2011), although other assessments have shown variable results (Russell et al., 2010; Rinaldi et al., 2013). The measurement of PMA and SMA composition and number coincident with multi-species characterisation of MBL and surface water composition during SOAP has provided a broad database with which to assess and develop these relationships for potential
625 application in remote sensing and Earth System Models. The first step towards this is the inclusion of SOAP aerosol and tropospheric data in the global ACCESS-UKCA model (Woodhouse et al., 2015), containing the GLOMAP-mode aerosol scheme (Mann et al., 2010, 2012), which shows very good agreement with observed distributions of condensation nuclei (Woodhouse et al, pers. comm.)

630 **6. Data availability**

The underway DMS_{sw} can be downloaded at <http://saga.pmel.noaa.gov/dms/select.php>. The remaining data is available by request email to cliff.law@niwa.co.nz

7. Supplement link (to be provided by Copernicus)

635

8. Author contribution

The SOAP campaign was led and coordinated by CSL, MJS & MJH. All authors developed analytical methods and instruments used during the SOAP campaign, and also analysed and interpreted data. CSL, MJS, MJH, TGB, LTC, FCE, SJL, ML, AM, JM, KAS, PV and CFW made measurements during the SOAP
640 voyage. CSL led manuscript preparation, with contributions from MJS, MJH, TGB, LTC, SJL, ML, RZ, PV and CFW.

The authors declare that they have no conflict of interest

645 9. Acknowledgements

We acknowledge the invaluable assistance of the Captain, officers and crew of the R/V *Tangaroa*. We thank Kim Currie, Steve George, Matt Walkington, Mark Gall, Marc Mallet, Greg Olsen, Gus Olivares and Nick Talbot for collection and analysis of samples; Christa Marandino, Warren DeBruyn and Cyril McCormick for assistance with API-CIMS measurements; Sebastian Landwehr, Scott Miller and Brian Ward for CO₂ flux measurements; Paul Johnson, Karin Kreher and Neil Harris for halocarbon and VOC measurements; Jason Ward for assistance with CCN measurement; Gordon Brailsford for Picarro set up; Paul Selleck and Min Cheng for analysis of VOC samples, Sarah Connors for the back-trajectory suite using the UK Met Office NAME model, and Matt Woodhouse for analysis of SOAP data using the global ACCESS-UKCA model. We also acknowledge Maurice Levasseur and Melita Keywood for their comments, and their support of the microbial DMSP cycling and VOC measurements, respectively.

10. Funding

This research was supported by funding from NIWA's Climate and Atmosphere Research Programme 3 – Role of the oceans (2015/16 SCI), and a Postdoctoral Fellowship (CO1X0911) for CW from the New Zealand Ministry for Business, Innovation and Employment (MBIE). Flux measurements were supported by a NSF Atmospheric Chemistry Program (grant nos. 08568, 0851472, 0851407 and 1143709), and VOC and CCN measurements supported by CSIRO's Capability Development Fund. PV participation was supported by EU COST action 735, the Academy of Finland through the Centre of Excellence, and via a Finnish Academy visiting grant no. 136841. SL participation was supported by Science Foundation Ireland as part of the US–Ireland R&D Partnership Programme under grant number 08/US/I1455, and a SFI Short-term Travel Fellowship under grant 09/US/I1758-STTF-11. We thank the Natural Sciences and Engineering Research Council of Canada (NSERC) for supporting the microbial DMSP cycling research.

670 11. Tables

	Time/Location				Meteorological			Hydrodynamic				Biogeochemical				
Bloom	Start NZST (DoY UTC)	End NZST (DoY UTC)	Bloom Centre Lat.	Bloom Centre Long.	Atmos Press. mb	Irrad W m ⁻²	U ₁₀ Range m s ⁻¹	Hs m	MLD m *	SST °C #	Sal. #	Nitrate/ Phosphate/ Silicate μmol L ⁻¹ *	Chl- <i>a</i> mg/m ³ *	pCO ₂ ppmv #	DMSsw nmolL ⁻¹ #	Dominant Phytoplankton *
B1	14/02/12 2:00 (44.6)	19/02/12 12:00 (50.0)	-44.34 -44.61	174.2 174.78	1019.1 ± 2.9	232 <1061	6.6 (5- 7.6)	2.0	14.5 ±1	14.5 ±0.4	34.48	5.3 ± 0.9 0.43 ± 0.2 0.35 ± 0.1	0.84 ±0.2 <3.4	320 ±24	16.8 ±1.5	Dinoflagellate
B2	21/02/12 16:15 (52.2)	26/02/12 12:00 (57.0)	-43.55 -43.71	180.16 180.32	1011.5 ± 5.3	196 <1079	10.4 (6.9- 12.4)	2.9	24.0 ±9	15.8 ±0.2	34.6	1.7 ± 1.0 0.27 ± 0.07 0.41 ± 0.33	0.67 ±0.3 <1.0	339 ±9	9.1 ±2.9	Coccolithophore Dinoflagellate
B3a	27/02/12 10:00 (57.9)	1/03/12 04:00 (60.67)	-44.11 -44.61	174.47 174.88	1010.0 ± 8.2	242 <1212	10.3 (8.1- 12.1)	2.6	28.6 ±1.7	14.4 ±0.2	34.32	3.7 ± 1 0.34 ± 0.06 0.3 ± 0.16	0.44 ± 0.17 <0.92	333 ±14	5.37 ±1.5	Mixed
B3b	02/03/12 06:00 (61.7)	5/03/12 17:00 (65.2)	-44.19 -44.78	174.3 174.93	1008.6 ± 9.4	182 <1016	12.6 (8.5- 14.9)	3.6	41.1 ±6	13.2 ±0.4	34.32	4.2 ± 1.1 0.39 ± 0.1 0.48 ± 0.05	0.59 ±0.2 <1.1	340 ±8	3.1 ±1.2	Mixed

Table 1. Summary of surface water characteristics during each bloom period. All values are Mean ±1 standard deviation, except where maximum value also shown by <. * indicates value derived from 2-10m depth on all stations during bloom occupation; # indicates continuous measurement in surface waters (nominal 6m depth). Abbreviations: Lat: Latitude; Long.: Longitude; Atmos. Press.: Atmospheric Pressure; Irrad.: Irradiance; U₁₀: Wind speed adjusted to 10m height (uncorrected for vessel flow distortion); Hs: Significant wave Height; MLD: Mixed Layer Depth; SST: Sea Surface Temperature; Sal>: Surface salinity; Chl-*a*: Chlorophyll-*a*.

Measurement	Mode	Instrument
WP1 Atmospheric		
Organic nuclei production	C*	UltraFine Organic-Tandem Differential Mobility Analyser (UFO-TDMA)
Aerosol water uptake and volatility	C*	Volatility Humidity Differential Mobility Analyser (VH-TDMA)
Nucleation, Aitken mode size spectra	C*	Scanning Mobility Particle Sizer (SMPS)
Condensation nuclei counts	C*	Condensation Particle Counter (CPC)
Accumulation mode aerosol number	C	PCASP
Cloud condensation nuclei concentration	C*	CCN spectrometer
Aerosol filter chemistry – Major ions	C	Hi-Vol, cascade, Ion Chromatograph
Black carbon	C*	Aetholometer
PM ₁ aerosol filters	C	Organic functional groups by FTIR and inorganic composition by IBA
Column aerosol	D	Sun photometer (Microtops II)
Nascent sea spray composition via bubble burst of sea-water samples	D	Chamber experiments
DMS	C	APIMS
CO ₂ and methane	C	Picarro CRDS
Halocarbons, Iodine and halogen oxides	C	μ-Dirac ECD-GC and Multi-Axis Differential Optical Absorption Spectroscopy (Max-DOAS)
VOCs (Acetone, DMS, Acetonitrile, Methanol, Methanethiol, Isoprene, Monoterpenes, Acetaldehyde	C	Proton Transfer Reaction Mass Spectrometer (PTR-MS)
VOCs C ₅ to C ₁₅	D	Pre-concentration and TD-GC-FID/MS
Aldehydes, ketones (including dicarbonyls), C ₂ to C ₈	D	Derivatisation and HPLC
WP2 Physics		
DMS flux	C	API-CIMS (chemical ionization mass spectrometry (mini-CIMS))
CO ₂ EC flux	C	Licors, Sonic anemometer motion sensors
DMS gradient flux	D	Catamaran, SCD-GC
Near surface TandS	D	CTD
Near-surface stratification	C	Spar buoy – temperature array, microcats
Near-surface turbulence	C	Vector, FastCat
Sea state	C	NOAA Wavewatch III
Whitecap coverage	D	Camera
Meteorological conditions	C	AWS
Bulk fluxes	C	Eppeley radiometers, raingauge; Eppeley Precision Spectral Pyranometer (PSP);
MBL Height and stability	D	Radiosonde deployment
WP3 Ocean biogeochemistry		
Chlorophyll- <i>a</i>	C, D, W	Ecotriplet
Back-scatter	C	Ecotriplet
B660 backscatter	C	Ecotriplet
pCO ₂	C	IRGA
pH	C, D, W	Spectrophotometer
DIC	D	
Nutrients	D, W	Colorimetric Autoanalyser
DOC	D, W	HTCO
CDOM	D, W	Spectrophotometer
POC/PON/PC/PN/13C/15N	D	Mass Spectrometer
Fatty Acids and Alkanes	D, W	
Dissolved DMS	C,	Meso-CIMS
Dissolved DMS	D, W	SCD/FPD
DMSP	D, W	SCD

DMSP processes	D, W	
Pigments	D	HPLC
Microbial community abundance	D, W	Flow Cytometry
Phytoplankton identification and counts	D, W	Optical Microscopy
Microzooplankton	D, W	Optical Microscopy

Table 2. Parameters sampled during the SOAP voyage. Key: C - Continuous, D - Discrete, W – Workboat, *indicates instrument sampling on common aerosol inlet.

Voyage	Date	Latitude	Longitude	Mean DMS (nmol L ⁻¹)	Std Dev	N	Method	Reference
BOX	October 2000	39.5-47°S	170-179°E	4.55	6.8	482	FPD-GC	this paper
	November 2005	49-50°S	175°E	1.75	-	2	FPD-GC	Kiene et al., 2007
SAGE	April 2006	41-46.6°S	172.5-78.5E	1.06	0.9	6	PFPD	Archer et al., 2011
PreSOAP	February 2011	42.5-44°S	174E-178°W	2.2	2.0	736	MIMS	this paper
SOAP	March-April 2012	41.7-46.5°S	172E-179°W	6.36	4.4	4132	API-CIMS	Bell et al., 2015
SOAP	March-April 2012	41.7-46.5°S	172E-179°W	11.5	9.2	22	SCD	Walker et al., 2016
S.W. Pacific	Weighted Mean	39.5-50°S	170-179°W	5.6		5380		
NEWZ		35-55S	170E-170W	0.05-2.0		6	Climatology	Lana et al., 2011
SSTC		35-50S	170W-170E	0.05-10			Climatology	Lana et al., 2011

Table 3. DMS data for the S.W. Pacific region east of New Zealand.

12. Figures

Figure 1. a) An ocean colour image on 10/2/11 during the PreSOAP voyage, showing phytoplankton blooms on the western Chatham Rise region along 44°S (data courtesy of NASA) b) The SOAP voyage track in the Chatham Rise region, overlain by Sea Surface Temperature (°C), with the study region (box) indicated in the inset bathymetric map of New Zealand.

Figure 2. nssSO₄ concentrations at New Zealand coastal atmospheric monitoring sites

Figure 3. Continuous measurement during PreSOAP of a) wind speed (m s^{-1}), b) atmospheric DMS (ppt), c) surface water DMS (nmol L^{-1}), and d) surface chlorophyll-*a* (mg m^{-3} ; quenched data removed).

Figure 4. 8-day composite images (at 4 km resolution) during the SOAP voyage for a) 10-17 Feb 2012 (DoY 41-48), b) 18-25 Feb 2012 (DoY 49-56), and c) 26th Feb-4th March (DoY 57-64), showing bloom locations (red dots), and daily true colour images for d) 16 Feb 2012 (DoY 47), e) 18 Feb 2012 (DoY 49) and f) 3 March (DoY 65) (MODIS Aqua data courtesy of NASA).

Figure 5. a)-c) Synoptic meteorology summary for each bloom period during the SOAP voyage. Surface pressure and wind plots are derived from the NZ local area Unified Model NZLAM, and the bloom location indicated by a red dot. d)-f) Back-trajectory analyses for each bloom period during the SOAP voyage. This was calculated using the Lagrangian Numerical Atmospheric-dispersion Modelling Environment (NAME) for the lower atmosphere (0-100m) as time integrated particle density (g s m^{-3}). Each plot shows the back-trajectory of 8 “releases”, i.e. one every three hours over 24 hours for the actual ship position.

Figure 6. Meteorological and hydrodynamic variables during the SOAP voyage, including a) Wind speed (WS, ms^{-1}); b) Direction (Dir., °); Wind (blue) and wave (cyan); c) Temperature (Temp., °C); Air (black) and surface water (green); d) Irradiance (Irrad., Wm^{-2}) and e) Significant wave height (Hs, m). The Bloom occupation periods are indicated by the red horizontal bars in the upper panel.

Figure 7. Surface water properties (2-10m) during the SOAP voyage: Temperature (Temp, °C), Mixed Layer Depth (ML Depth, m), Chlorophyll-*a* (Chl-*a*, mg m^{-3}), and nitrate concentration ($\mu\text{mol L}^{-1}$), with the occupation period for each bloom indicated by the vertical shaded bars.

Fig. 8. SOAP parameters and integrated work programmes

Figure 9. Aerosol characterisation during SOAP indicating size spectral (red) and total counts (black). Ambient RH measurement was used for RH correction of the PCASP, Hi Vol and SMPS, and diffusion driers (Silica Gel) were used on the inlet of the UFO-TDMA and VH-TDMA.

Figure 10. a) Marine boundary layer CN concentrations (top, CPC3772 in blue, CPC3010 in red), b) CCN concentrations (middle) and c) number of hours 72 hour back-trajectory was over land (bottom, 27-member ensemble average).

13. References

- Allen, A. G., Dick, A. L., and Davison, B. M.: Sources of atmospheric methanesulphonate, non-sea-salt sulphate, nitrate and related species over the temperate South Pacific, *Atmospheric Environment*, 31, 191-205, doi:10.1016/1352-2310(96)00194-x, 1997.
- Almeida, J., Schobesberger, S., Kürten, A., Ortega, I. K., Kupiainen-Maatta, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppa, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Trostl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H. and Kirkby, J.: Molecular understanding of sulphuric acid-amine particle nucleation in the atmosphere, *Nature*, 502(7471), 359–363, doi:10.1038/nature12663, 2013.
- Andreae, M. O., and Rosenfeld, D.: Aerosol–cloud–precipitation interactions. Part 1. The nature and sources of cloud-active aerosols, *Earth-Science Reviews*, 89, 13-41, doi:10.1016/j.earscirev.2008.03.001, 2008.
- Archer, S. D., Safi, K., Hall, A., Cummings, D. G., and Harvey, M.: Grazing suppression of dimethylsulphoniopropionate (DMSP) accumulation in iron-fertilised, sub-Antarctic waters, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 58, 839-850, doi:10.1016/j.dsr2.2010.10.022, 2011.
- Ayers, G. P., Ivey, J. P., and Gillett, R. W.: Coherence between seasonal cycles of dimethylsulfide, methanesulfonate, and sulfate in Marine Air, *Nature*, 349, 404-406, 1991.
- Balch, W. M., Drapeau, D. T., Bowler, B. C., Lyczkowski, E., Booth, E. S., and Alley, D.: The contribution of coccolithophores to the optical and inorganic carbon budgets during the Southern Ocean Gas Exchange Experiment: New evidence in support of the "Great

- Calcite Belt" hypothesis, *Journal of Geophysical Research-Oceans*, 116, doi:10.1029/2011jc006941, 2011.
- Bell, T. G., De Bruyn, W., Marandino, C. A., Miller, S. D., Law, C. S., Smith, M. J., and Saltzman, E. S.: Dimethylsulfide gas transfer coefficients from algal blooms in the Southern Ocean, *Atmos. Chem. Phys.*, 15, 1783-1794, doi:10.5194/acp-15-1783-2015, 2015.
- Bell, T. G., Landwehr, S., Miller, S. D., de Bruyn, W. J., Callaghan, A. H., Scanlon, B., Ward, B., Yang, M., and Saltzman, E. S.: Estimation of bubble-mediated air-sea gas exchange from concurrent DMS and CO₂ transfer velocities at intermediate-high wind speeds, *Atmos. Chem. Phys.*, 17, 9019-9033, 10.5194/acp-17-9019-2017, 2017.
- Bell, T. G., Malin, G., Lee, G. A., Stefels, J., Archer, S., Steinke, M., and Matrai, P.: Global oceanic DMS data inter-comparability, *Biogeochem*, 110, 147-161, 10.1007/s10533-011-9662-3, 2012.
- Blake, N. J., Blake, D. R., Wingenter, O. W., Sive, B. C., Kang, C. H., Thornton, D. C., Bandy, A. R., Atlas, E., Flocke, F., Harris, J. M., and Rowland, F. S.: Aircraft measurements of the latitudinal, vertical, and seasonal variations of NMHCs, methyl nitrate, methyl halides, and DMS during the First Aerosol Characterization Experiment (ACE 1), *Journal of Geophysical Research-Atmospheres*, 104, 21803-21817, doi:10.1029/1999jd900238, 1999.
- Blomquist, B.W., Fairall, C.W., Huebert, B.J., Kieber, D.J., Westby, G.R.: DMS sea-air transfer velocity: Direct measurements by eddy covariance and parameterization based on the NOAA/COARE gas transfer model. *Geophysical Research Letters*, 33(7), 2006.
- Blot, R., Clarke, A. D., Freitag, S., Kapustin, V., Howell, S. G., Jensen, J. B., Shank, L. M., McNaughton, C. S. and Brekhovskikh, V.: Ultrafine sea spray aerosol over the southeastern Pacific: open-ocean contributions to marine boundary layer CCN, *Atmos. Chem. Phys.*, 13(14), 7263-7278, doi:10.5194/acp-13-7263-2013, 2013.
- Boers, R., Jensen, J. B., Krummel, P. B., and Gerber, H.: Microphysical and radiative structure of wintertime marine stratocumulus clouds over the Southern Ocean, *Q. Jl R. met. Soc.*, 122, 1307-1339, 1996.
- Boers, R., Jensen, J. B., and Krummel, P. B.: Microphysical and shortwave radiative structure of stratocumulus clouds over the Southern Ocean: summer results and seasonal differences, *Q. Jl R. met. Soc.*, 124, 151-168, 1998.
- Boyd, P., LaRoche, J., Gall, M., Frew, R., and McKay, R. M. L.: Role of iron, light, and silicate in controlling algal biomass in subantarctic waters SE of New Zealand, *Journal of Geophysical Research-Oceans*, 104, 13395-13408, doi:10.1029/1999jc900009, 1999.
- Boyd, P. W., McTainsh, G., Sherlock, V., Richardson, K., Nichol, S., Ellwood, M., and Frew, R.: Episodic enhancement of phytoplankton stocks in New Zealand subantarctic waters: Contribution of atmospheric and oceanic iron supply, *Global Biogeochemical Cycles*, 18, C8 GB1029, doi:10.1029/2002GB002020 2004.
- Bradford-Grieve, J. M., Boyd, P. W., Chang, F. H., Chiswell, S., Hadfield, M., Hall, J. A., James, M. R., Nodder, S. D., and Shushkina, E. A.: Pelagic ecosystem structure and functioning in the Subtropical Front region east of New Zealand in austral winter and spring 1993, *Journal of Plankton Research*, 21, 405-428, doi:10.1093/plankt/21.3.405, 1999.
- Cainey, J., and Harvey, M.: Dimethylsulfide, a limited contributor to new particle formation in the clean marine boundary layer (DOI 10.1029/2001GL014439), *Geophysical Research Letters*, 29, 32, doi:DOI: 10.1029/2001GL014439, 2002.

- Calleja, M. L., Duarte, C. M., Navarro, N., and Agustí, S.: Control of air-sea CO₂ disequilibria in the subtropical NE Atlantic by planktonic metabolism under the ocean skin, *Geophysical Research Letters*, 32, L08606, doi:10.1029/2004GL022120, 2005.
- Calleja, M. L., Duarte, C. M., Alvarez, M., Vaquer-Sunyer, R., Agustí, S., and Herndl, G. J.: Prevalence of strong vertical CO₂ and O₂ variability in the top meters of the ocean, *Global Biogeochemical Cycles*, 27, 941-949, doi:10.1002/gbc.20081, 2013.
- Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth system, *Atmospheric Chemistry and Physics*, 10, 1701-1737, doi:10.5194/acp-10-1701-2010, 2010.
- Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., and Pierce, J. R.: Large contribution of natural aerosols to uncertainty in indirect forcing, *Nature*, 503, 66-71, 2013.
- Chang, F. H., and Gall, M. Phytoplankton assemblages and photosynthetic pigments during winter and spring in the Subtropical Convergence region near New Zealand, *New Zealand Journal of Marine and Freshwater Research*, 32, 515-530, 1998.
- Chang, F. H., Williams, M. J. M., Schwarz, J. N., Hall, J. A., Maas, E. W., and Stewart, R.: Spatial variation of phytoplankton assemblages and biomass in the New Zealand sector of the Southern Ocean during the late austral summer 2008, *Polar Biology*, 36, 391-408, doi:10.1007/s00300-012-1270-8, 2013.
- Chang, R.Y.W., Sjostedt, S.J., Pierce, J.R., Papakyriakou, T.N., Scarratt, M.G., Michaud, S., Levasseur, M., Leaitch, W.R. and Abbatt, J.P. Relating atmospheric and oceanic DMS levels to particle nucleation events in the Canadian Arctic. *Journal of Geophysical Research: Atmospheres*, 116(D17), 2011.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulfur, cloud albedo and climate, *Nature*, 326, 655-661, doi:10.1038/326655A0 1987.
- Chiswell, S. M., Bradford-Grieve, J., Hadfield, M. G., and Kennan, S. C.: Climatology of surface chlorophyll a, autumn-winter and spring blooms in the southwest Pacific Ocean, *Journal of Geophysical Research-Oceans*, 118, 1003-1018, doi:10.1002/jgrc.20088, 2013.
- Ciuraru, R., Fine, L., Pinxteren, M. V., D'Anna, B., Herrmann, H., and George, C.: Unravelling new processes at interfaces: Photochemical isoprene production at the sea surface. *Environmental Science and Technology*, 49(22), 13199-13205, 2015.
- Clarke, A. D., Freitag, S., Simpson, R. M. C., Hudson, J. G., Howell, S. G., Brekhovskikh, V. L., Campos, T., Kapustin, V. N. and Zhou, J.: Free troposphere as the dominant source of CCN in the Equatorial Pacific boundary layer: long-range transport and teleconnections, *Atmospheric Chemistry and Physics Discussions*, 13(1), 1279–1326, doi:10.5194/acpd-13-1279-2013, 2013.
- Cohen, D. D., Stelcer, E., Hawas, O. and Garton, D.: IBA methods for characterisation of fine particulate atmospheric pollution: a local, regional and global research problem, *The 13th International Conference on Particle Induced X-ray Emission (PIXE 2013)*, 219–220 IS -, 145–152, 2004.
- Cravigan, L. T., Ristovski, Z., Modini, R. L., Keywood, M. D. and Gras, J. L.: Observation of sea-salt fraction in sub-100 nm diameter particles at Cape Grim, *J. Geophys. Res. Atmos.*, 2014JD022601, doi:10.1002/2014JD022601, 2015.

- de Bruyn, W. J., Harvey, M., Caine, J. M., and Saltzman, E. S.: DMS and SO₂ at Baring Head, New Zealand: Implications for the yield of SO₂ from DMS, *Journal of Atmospheric Chemistry*, 41, 189-209, doi:10.1023/a:1014252106572, 2002.
- de Leeuw, G., Guieu, C., Arneth, A., Bellouin, N., Bopp, L., Boyd, P. W., van der Gon, H. A. C. D., Desboeufs, K. V., Dulac, F., Facchini, M. C., Gantt, B., Langmann, B., Mahowald, N. M., Maranón, E., O'Dowd, C., Olgun, N., Pulido-Villena, E., Rinaldi, M., Stephanou, E. G., and Wagener, T.: Ocean–Atmosphere Interactions of Particles, P.S. Liss and M.T. Johnson (Eds.), *Ocean-Atmosphere Interactions of Gases and Particles*, Springer Earth System Sciences, 171-246, doi:10.1007/978-3-642-25643-1_4,, 2014.
- Decesari, S., Finessi, E., Rinaldi, M., Paglione, M., Fuzzi, S., Stephanou, E.G., Tziaras, T., Spyros, A., Ceburnis, D., O'Dowd, C. and Dall'Osto, M.: Primary and secondary marine organic aerosols over the North Atlantic Ocean during the MAP experiment. *Journal of Geophysical Research: Atmospheres*, 116(D22), 2011.
- Delizo, L., Smith, W. O., and Hall, J.: Taxonomic composition and growth rates of phytoplankton assemblages at the Subtropical Convergence east of New Zealand, *Journal of Plankton Research*, 29, 655-670, doi:10.1093/plankt/fbm047, 2007.
- Draxler, R. R. and Rolph, G. D., Eds.: HYSPLIT(HYbrid Single-PARTicle Lagrangian Intergrated Trajectory) Model, NOAA Air Resources Laboratory, College Park, MD. [online] Available from: <http://www.arl.noaa.gov/HYSPLIT.php> (Accessed 1 September 2013), 2013.
- Facchini, M. C., Rinaldi, M., Decesari, S., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Ceburnis, D., Flanagan, R., Nilsson, E. D., de Leeuw, G., Martino, M., Woeltjen, J., and O'Dowd, C. D.: Primary submicron marine aerosol dominated by insoluble organic colloids and aggregates, *Geophysical Research Letters*, 35, L17814, doi:10.1029/2008GL034210, 2008.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk parameterization of air-sea fluxes: Updates and verification for the COARE algorithm, *J Climate*, 16, 571-591, 2003.
- Fairall, C. W., Yang, M. X., Bariteau, L., Edson, J. B., Helmig, D., McGillis, W., Pezoa, S., Hare, J. E., Huebert, B., and Blomquist, B.: Implementation of the Coupled Ocean-Atmosphere Response Experiment flux algorithm with CO₂, dimethyl sulfide, and O₃, *Journal of Geophysical Research-Oceans*, 116, doi:10.1029/2010jc006884, 2011.
- Gantt, B., and Meskhidze, N.: The physical and chemical characteristics of marine primary organic aerosol: a review, *Atmos. Chem. Phys.*, 13, 3979-3996, doi:10.5194/acp-13-3979-2013, 2013.
- Gantt, B., Meskhidze, N., Facchini, M. C., Rinaldi, M., Ceburnis, D., and O'Dowd, C. D.: Wind speed dependent size-resolved parameterization for the organic mass fraction of sea spray aerosol, *Atmospheric Chemistry and Physics*, 11, 8777-8790, 2011.
- Gondwe, M., Krol, M., Gieskes, W., Klaassen, W., De Baar, H.: The contribution of ocean-leaving DMS to the global atmospheric burdens of DMS, MSA, SO₂, and NSS SO₄=, *Global Biogeochemical Cycles*, 17(2), 2003.
- Griffiths, F. B., Bates, T. S., Quinn, P. K., Clementson, L. A., Parslow, J. S.: Oceanographic context of the First Aerosol Characterization Experiment (ACE 1): A physical, chemical, and biological overview, *Journal of Geophysical Research-Atmospheres*, 104, 21649-21671, doi:10.1029/1999jd900386, 1999.
- Harvey, M. J., Wylie, D. J., Martin, R. J., Clarkson, T. S., and de Mora, S. J.: Dimethylsulphide measurements at Baring Head, New Zealand, in: *Dimethylsulphide*, Oceans,

- Atmosphere and Climate, edited by: Restelli, G., and Angeletti, G., Kluwer Academic, Dordrecht, 143-151, 1993.
- Harvey, M. J., Law, C. S., Smith, M. J., Hall, J. A., Abraham, E. R., Stevens, C. L., Hadfield, M. G., Ho, D. T., Ward, B., Archer, S. D., Caine, J. M., Currie, K. I., Devries, D., Ellwood, M. J., Hill, P., Jones, G. B., Katz, D., Kuparinen, J., Macaskill, B., Main, W., Marriner, A., McGregor, J., McNeil, C., Minnett, P. J., Nodder, S. D., Peloquin, J., Pickmere, S., Pinkerton, M. H., Safi, K. A., Thompson, R., Walkington, M., Wright, S. W., and Ziolkowski, L. A.: The SOLAS air-sea gas exchange experiment (SAGE) 2004, Deep Sea Research Part II: Topical Studies in Oceanography, 58, 753-763, doi:10.1016/j.dsr2.2010.10.015, 2011.
- Harvey M.J., McGregor J.A., Lawson S. J., Cravigan L., Talbot N., Olivares G., Connors S., Woodhouse M.: Aerosol and tracer measurement at sea – defining the baseline. Atmospheric Chemistry and Physics, to be *submitted*
- Ho, D. T., Law, C. S., Smith, M. J., Schlosser, P., Harvey, M., and Hill, P.: Measurements of air-sea gas exchange at high wind speeds in the Southern Ocean: Implications for global parameterizations. doi:10.1029/2006GL026817, Geophys. Res. Lett., 33, L16611, 2006.
- Jaeglé, L., Quinn, P. K., Bates, T. S., Alexander, B., and Lin, J. T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, Atmos. Chem. Phys., 11, 3137-3157, doi:10.5194/acp-11-3137-2011, 2011.
- Johnson, G., Ristovski, Z., and Morawska, L.: Application of the VH-TDMA technique to coastal ambient aerosols, Geophys. Res. Lett., 31, doi:10.1029/2004GL020126, 2004a.
- Jones, A. R., Thomson, D. J., Hort, M., and Devenish, B.: The UK Met Office's next-generation atmospheric dispersion model, NAME III, in: Proceedings of the 27th NATO/CCMS International Technical Meeting on Air Pollution Modelling and its Application, edited by: Borrego, C., and Norman, A.-L., Springer, pp 580 - 589, 2007.
- Kay, J. E., Wall, C., Yettella, V., Medeiros, B., Hannay, C., Caldwell, P., and Bitz, C.: Global Climate Impacts of Fixing the Southern Ocean Shortwave Radiation Bias in the Community Earth System Model (CESM), Journal of Climate, 29, 4617-4636, doi:doi:10.1175/JCLI-D-15-0358.1, 2016.
- Kiene, R. P., Kieber, D. J., Slezak, D., Toole, D. A., del Valle, D. A., Bisgrove, J., Brinkley, J., and Rellinger, A.: Distribution and cycling of dimethylsulfide, dimethylsulfoniopropionate, and dimethylsulfoxide during spring and early summer in the Southern Ocean south of New Zealand, Aquatic Sciences, 69, 305-319, doi:10.1007/s00027-007-0892-3, 2007.
- Korhonen, H., Carslaw, K. S., Spracklen, D. V., Mann, G. W., and Woodhouse, M. T.: Influence of oceanic dimethyl sulfide emissions on cloud condensation nuclei concentrations and seasonality over the remote Southern Hemisphere oceans: A global model study, Journal of Geophysical Research-Atmospheres, 113, doi:10.1029/2007jd009718, 2008.
- Lana, A., Bell, T. G., Simo, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S. 2011. An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, Global Biogeochemical Cycles, 25, doi:10.1029/2010gb003850, 2011.
- Lana, A., Simo, R., Vallina, S. M., and Dachs, J.: Potential for a biogenic influence on cloud microphysics over the ocean: a correlation study with satellite-derived data, Atmospheric Chemistry and Physics, 12, 7977-7993, doi:10.5194/acp-12-7977-2012, 2012.

- Landschuetzer, P., Gruber, N., Bakker, D. C. E., and Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochemical Cycles*, 28, 927-949, doi:10.1002/2014gb004853, 2014.
- Landwehr, S., Miller, S. D., Smith, M. J., Saltzman, E. S., and Ward, B.: Analysis of the PKT correction for direct CO₂ flux measurements over the ocean, *Atmospheric Chemistry and Physics*, 14, 3361-3372, 2014.
- Landwehr, S., Miller, S. D., Smith, M. J., Saltzman, E. S., Bell, T. G., and Ward, B. Using Eddy Covariance to Measure the Dependence of Air-Sea CO₂ Exchange Rate on Friction Velocity, *Atmospheric Chemistry and Physics* *submitted*
- Lawson, S. J., Selleck, P. W., Galbally, I. E., Keywood, M. D., Harvey, M. J., Lerot, C., Helmig, D., and Ristovski, Z.: Seasonal in situ observations of glyoxal and methylglyoxal over the temperate oceans of the Southern Hemisphere, , 15, 223-240, doi:10.5194/acp-15-223-2015, 2015.
- Lawson S.J., Selleck P.W., Cheng M. and M. Harvey: VOCs in marine air over the biologically productive Chatham Rise. *Atmospheric Chemistry and Physics*, to be submitted
- Lizotte, M., Levasseur, M., Law, C. S., Walker, C. F., Safi, K. A., Marriner, A., and Kiene, R. P.: Dimethylsulfoniopropionate (DMSP) and dimethylsulfide (DMS) cycling across contrasting biological hotspots of the New Zealand Subtropical Front, *Ocean Sci. Discuss.*, 2017, 1-49, doi:10.5194/os-2017-32, 2017
- Mahajan, A. S., Fadnavis, S., Thomas, M. A., Pozzoli, L., Gupta, S., Royer, S.-J., Saiz-Lopez, A., and R., S.: Quantifying the impacts of an updated global dimethyl sulfide climatology on cloud microphysics and aerosol radiative forcing, *J. Geophys. Res. Atmos.*, 120, 2524-2536, doi:doi:10.1002/2014JD022687, 2015.
- Mahajan, A. S., et al.: Glyoxal observations in the global marine boundary layer, *J. Geophys. Res. Atmos.*, 119, 6160–6169, doi:10.1002/2013JD021388, 2014
- Mallet, M., L. Cravigan, B. Miljevic, P. Vaattovaara, E. Deschaseaux, H. Swan, G. Jones, *and* Z. Ristovski (2016), Sea spray aerosol in the Great Barrier Reef and the presence of nonvolatile organics, *J. Geophys. Res. Atmos.*, 121, 7088–7099, doi:10.1002/2016JD024966.
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J. and Johnson, C. E. Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model Geoscientific Model Development, 3, 519-551, 2012.
- Mann, G. W., Carslaw, K. S., Ridley, D. A., Spracklen, D. V., Pringle, K. J., Merikanto, J., Korhonen, S., Schwarz, J. P., Lee, L. A., Manktelow, P. T., Woodhouse, M. T., Schmidt, A., Breider, T. J., Emmerson, K. M., Chipperfield, M. P. and Pickering, S. J. Intercomparison of modal and sectional aerosol microphysics representations within the same 3-D global chemical transport model *Atmospheric Chemistry and Physics*, 12, 4449-4476, 2012.
- Marandino, C. A., De Bruyn, W. J., Miller, S. D., and Saltzman, E. S.: DMS air/sea flux and gas transfer coefficients from the North Atlantic summertime coccolithophore bloom, *Geophysical Research Letters*, 35, doi:10.1029/2008gl036370, 2008.
- McCoy, D. T., Burrows, S. M., Wood, R., Grosvenor, D. P., Elliott, S. M., Ma, P.-L., Rasch, P. J., and Hartmann, D. L.: Natural aerosols explain seasonal and spatial patterns of Southern Ocean cloud albedo, *Science Advances*, 1, doi:10.1126/sciadv.1500157, 2015.


- Meskhidze, N., Xu, J., Zhang, Y., Gantt, B., Ghan, S., Nenes, A., Liu, X., Easter, R., and Zaveri, R.: Effect of marine biogenic organic aerosols on cloud properties: Modeling study, *Geochimica Et Cosmochimica Acta*, 73, A874-A874, 2009.
- Meskhidze, N., and Nenes, A.: Phytoplankton and cloudiness in the Southern Ocean. *Science*, 314(5804), 1419-1423, 2006.
- Meskhidze, N., and Nenes, A.: Effects of Ocean Ecosystem on Marine Aerosol-Cloud Interaction, *Advances in Meteorology*, 2010, doi:10.1155/2010/239808, 2010.
- Meskhidze, N., Xu, J., Gantt, B., Zhang, Y., Nenes, A., Ghan, S. J., Liu, X., Easter, R., and Zaveri, R.: Global distribution and climate forcing of marine organic aerosol: 1. Model improvements and evaluation, *Atmospheric Chemistry and Physics*, 11, 11689-11705, doi:10.5194/acp-11-11689-2011, 2011.
- Modini, R. L., Harris, B. and Ristovski, Z.: The organic fraction of bubble-generated, accumulation mode Sea Spray Aerosol (SSA), *Atmos. Chem. Phys.*, 10(6), 2867–2877, 2010.
- Murphy, R. J., Pinkerton, M. H., Richardson, K. M., Bradford-Grieve, J. M., and Boyd, P. W.: Phytoplankton distributions around New Zealand derived from SeaWiFS remotely-sensed ocean colour data, *New Zealand Journal of Marine and Freshwater Research*, 35, 343-362, 2001.
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat M.I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochemical Cycles*, 14, 373-387, 2000.
- Nodder, S. D., Duineveld, G. C. A., Pilditch, C. A., Sutton, P. J., Probert, P. K., Lavaleye, M. S. S., Witbaard, R., Chang, F. H., Hall, J. A., and Richardson, K. M.: Focusing of phytodetritus deposition beneath a deep-ocean front, Chatham Rise, New Zealand, *Limnology and Oceanography*, 52, 299-314, 2007.
- O'Dowd, C., Monahan, C., and Dall'Osto, M.: On the occurrence of open ocean particle production and growth events, *Geophysical Research Letters*, 37, doi:10.1029/2010gl044679, 2010.
- O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and Putaud, J.-P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676-680, 2004.
- O'Dowd, C. D., and de Leeuw, G.: Marine aerosol production: a review of the current knowledge, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 365, 1753-1774, 2007.
- Popinet, S., Smith, M., and Stevens, C.: Experimental and Numerical Study of the Turbulence Characteristics of Airflow around a Research Vessel, *Journal of Atmospheric and Oceanic Technology*, 21, 1575-1589, doi:10.1175/1520-0426(2004)021<1575:EANSOT>2.0.CO;2 2004.
- Quinn, P. K., and Bates, T. S.: The case against climate regulation via oceanic phytoplankton sulphur emissions, *Nature*, 480, 51-56, 2011.
- Quinn, P.K., Bates, T.S., Schulz, K.S., Coffman, D.J., Frossard, A.A., Russell, L.M., Keene, W.C., Kieber, D.J.: Contribution of sea surface carbon pool to organic matter enrichment in sea spray aerosol. *Nature Geoscience*, 7(3), 228-232, 2014.
- Rinaldi, M., Fuzzi, S., Decesari, S., Marullo, S., Santoleri, R., Provenzale, A., von Hardenberg, J., Ceburnis, D., Vaishya, A., O'Dowd, C. D., and Facchini, M. C.: Is chlorophyll-a the best

- surrogate for organic matter enrichment in submicron primary marine aerosol?, *Journal of Geophysical Research-Atmospheres*, 118, 4964-4973, doi:10.1002/jgrd.50417, 2013.
- Russell, L. M., Bahadur, R. and Ziemann, P. J.: Identifying organic aerosol sources by comparing functional group composition in chamber and atmospheric particles, *Proceedings of the National Academy of Sciences*, 2011.
- Russell, L. M., Hawkins, L. N., Frossard, A. A., Quinn, P. K., and Bates, T. S.: Carbohydrate-like composition of submicron atmospheric particles and their production from ocean bubble bursting, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6652-6657, doi:10.1073/pnas.0908905107, 2010.
- Sadeghi, A., Dinter, T., Vountas, M., Taylor, B. B., Altenburg-Soppa, M., Peeken, I., and Bracher, A.: Improvement to the PhytoDOAS method for identification of coccolithophores using hyper-spectral satellite data, *Ocean Science*, 8, 1055-1070, doi:10.5194/os-8-1055-2012, 2012.
- Scanlon, B., and Ward, B.: The influence of environmental parameters on active and maturing oceanic whitecaps, *Journal of Geophysical Research: Oceans*, 121, doi:10.1002/2015JC011230, 2016.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, *Proceedings of the National Academy of Sciences*, 113, 5781-5790, doi:10.1073/pnas.1514043113, 2016.
- Sellegri, K., Loon, Y. J., Jennings, S. G., O'Dowd, C. D., Pirjola, L., Cautenet, S., Chen, H. W., and Hoffmann, T.: Quantification of coastal new ultra-fine particles formation from in situ and chamber measurements during the BIOFLUX campaign, *Environmental Chemistry*, 2, 260-270, doi:10.1071/en05074, 2005.
- Sievering, H., Caine, J., Harvey, M., McGregor, J., Nichol, S., and Quinn, P.: Aerosol non-sea-salt sulfate in the remote marine boundary layer under clear-sky and normal cloudiness conditions: Ocean-derived biogenic alkalinity enhances sea-salt sulfate production by ozone oxidation, *Journal of Geophysical Research-Atmospheres*, 109, doi:10.1029/2003jd004315, 2004.
- Smith, M. J., Ho, D. T., Law, C. S., McGregor, J., Popinet, S., and Schlosser, P.: Uncertainties in gas exchange parameterization during the SAGE dual-tracer experiment, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 58, 869-881, doi:10.1016/j.dsr2.2010.10.025, 2011.
- Smith M.J., Walker C.F., Bell T.G., Harvey M.J., Law C.S., Saltzman E.S. Air-sea gradient fluxes of DMS during the Surface Ocean Aerosol Production (SOAP) experiment. *Ocean Sciences*, *submitted*
- Spada, M., Jorba, O., Pérez García-Pando, C., Janjic, Z., and Baldasano, J. M.: On the evaluation of global sea-salt aerosol models at coastal/orographic sites, *Atmospheric Environment*, 101, 41-48, doi:http://dx.doi.org/10.1016/j.atmosenv.2014.11.019, 2015.
- Stephens, G. L.: Cloud feedbacks in the climate system: A critical review, *Journal of climate* 18, 237-273, 2005.
- Stevens, C. L., Hill, P., Smith, M. J., and Popinet, S.: SUCA: An engine for repetitive autonomous profiling near the ocean surface, *Limnol Oceanogr-Meth*, 3, 300-307, 2005.

- Swan, H. B., Armishaw, P., Lavetz, R., Alamgir, M., Davies, S. R., Bell, T. G., and Jones, G. B.: An interlaboratory comparison for the quantification of aqueous dimethylsulfide, *Limnology and Oceanography-Methods*, 12, 784-794, doi:10.4319/lom.2014.12.784, 2014.
- Thomas, M. A., Suntharalingam, P., Pozzoli, L., Rast, S., Devasthale, A., Kloster, S., Feichter, J., and Lenton, T. M.: Quantification of DMS aerosol-cloud-climate interactions using the ECHAM5-HAMMOZ model in a current climate scenario, *Atmospheric Chemistry and Physics*, 10, 7425-7438, 2010.
- Trenberth, K. E., and Fasullo, J. T.: Simulation of present-day and twenty-first-century energy budgets of the southern oceans, *J. Climate*, 23, 440-454, 2010.
- Twomey, S.: The influence of pollution on the shortwave albedo of clouds. *Journal of the atmospheric sciences*, 34(7), 1149-1152, 1977.
- Vaattovaara, P., Räsänen, M., Kühn, T., Joutsensaari, J., and Laaksonen, A.: A method for detecting presence of organic fraction in nucleation mode sized particles, *Atmospheric Chemistry and Physics*, 5, 3277-3287, 2005.
- Vaattovaara, P., Huttunen, P. E., Yoon, Y. J., Joutsensaari, J., Lehtinen, K. E. J., O'Dowd, C. D., and Laaksonen, A.: The composition of nucleation and Aitken modes particles during coastal nucleation events: evidence for marine secondary organic contribution, *Atmos. Chem. Phys.*, 6, 4601-4616, doi:10.5194/acp-6-4601-2006, 2006.
- Vaishya, A., Jennings, S. G., and O'Dowd, C.: Wind-driven influences on aerosol light scattering in north-east Atlantic air, *Geophysical Research Letters*, 39, 2012.
- Vallina, S.M., Simó, R., Gassó, S.: What controls CCN seasonality in the Southern Ocean? A statistical analysis based on satellite-derived chlorophyll and CCN and model-estimated OH radical and rainfall. *Global Biogeochemical Cycles*, 20(1), 2006
- Vignati, E., Facchini, M.C., Rinaldi, M., Scannell, C., Ceburnis, D., Sciare, J., Kanakidou, M., Myriokefalitakis, S., Dentener, F., O'Dowd, C.D.: Global scale emission and distribution of sea-spray aerosol: Sea-salt and organic enrichment. *Atmospheric Environment*, 44(5), 670-677, 2010.
- Villani, P., Picard, D., Michaud, V., Laj, P., and Wiedensohler, A.: Design and Validation of a Volatility Hygroscopic Tandem Differential Mobility Analyzer (VH-TDMA) to Characterize the Relationships Between the Thermal and Hygroscopic Properties of Atmospheric Aerosol Particles, *Aerosol Science and Technology*, 42, 729-741, doi:10.1080/02786820802255668, 2008.
- Walker, C. F., Harvey, M. J., Smith, M. J., Bell, T., Saltzman, E., Marriner, A. S., McGregor, J. A., and Law, C. S.: Assessing the potential for DMS enrichment at the sea-surface and its influence on sea-to-air flux *Ocean Sciences*, 12, 1033-1048, doi:doi:10.5194/os-12-1033-2016., 2016.
- Wang, H., Easter, R. C., Rasch, P. J., Wang, M., Liu, X., Ghan, S. J., Y., Q., JH., Y., P.L., M., and Vinoj, V.: Sensitivity of remote aerosol distributions to representation of cloud-aerosol interactions in a global climate model, *Geoscientific Model Development*, 6.3, 765-782, 2013.
- Willis, M.D., Burkart, J., Thomas, J.L., Köllner, F., Schneider, J., Bozem, H., Hoor, P.M., Aliabadi, A.A., Schulz, H., Herber, A.B. and Leaitch, W.R.: Growth of nucleation mode particles in the summertime Arctic: a case study. *Atmospheric Chemistry and Physics*, 16(12), 7663-7679, 2016.

- Woodhouse, M. T., Luhar, A. K., Stevens, L., Galbally, I., Thatcher, M., Uhe, P., Wolff, H., Noonan, J., and Molloy, S. Australian reactive-gas emissions in a global chemistry-climate model and initial results *Air Quality and Climate Change*, 49, 31-38, 2015.
- Wylie, D. J., and de Mora, S. J.: Atmospheric dimethylsulfide and sulfur species in aerosol and rainwater at coastal site in New Zealand, *Journal of Geophysical Research*, 101, 21041-21049, 1996.
- Yang, M., Blomquist, B. W., Fairall, C. W., Archer, S. D., and Huebert, B. J.: Air-sea exchange of dimethylsulfide in the Southern Ocean: Measurements from SO GasEx compared to temperate and tropical regions, *J. Geophys. Research*, 116,, doi:10.1029/2010JC006526, 2011.
- Young, I. R., Vinoth, J., Zieger, S., and Babanin, A. V.: Investigation of trends in extreme value wave height and wind speed, *J Geophys Res-Oceans*, 117, 2012.
- Zhang, R.: Getting to the Critical Nucleus of Aerosol Formation, *Science*, 328, 1366-1367, doi:10.1126/science.1189732, 2010.

a)



b)




Figure 1. a) An ocean colour image on 10/2/11 during the PreSOAP voyage, showing phytoplankton blooms on the western Chatham Rise region along 44°S (data courtesy of NASA) b) The SOAP voyage track in the Chatham Rise region, overlain by Sea Surface Temperature (°C), with the study region (box) indicated in the inset bathymetric map of New Zealand.




Figure 2. nssSO₄ concentrations at New Zealand coastal atmospheric monitoring sites




Figure 3. Continuous measurement during PreSOAP of a) windspeed (m s^{-1}), b) atmospheric DMS (ppt), c) surface water DMS (nmol l^{-1}), and d) surface chlorophyll-*a* (mg m^{-3} ; quenched data removed).




Figure 4. 8-day composite images (at 4 km resolution) during the SOAP voyage for a) 10-17 Feb 2012 (DoY 41-48), b) 18-25 Feb 2012 (DoY 49-56), and c) 26th Feb-4th March (DoY 57-64), showing bloom locations (red dots), and daily true colour images for d) 16 Feb 2012 (DoY 47), e) 18 Feb 2012 (DoY 49) and f) 3rd March (DoY 65) (MODIS Aqua data courtesy of NASA).




Figure 5. a)-c) Synoptic meteorology summary for each bloom period during the SOAP voyage. Surface pressure and wind plots are derived from the NZ local area Unified Model NZLAM, and the bloom location indicated by a red dot. d)-f) Back-trajectory analyses for each bloom period during the SOAP voyage. This was calculated using the Lagrangian Numerical Atmospheric-dispersion Modelling Environment (NAME)

for the lower atmosphere (0-100m) as time integrated particle density (g s m^{-3}). Each plot shows the back-trajectory of 8 “releases”, i.e. one every three hours over 24 hours for the actual ship position




Figure 6. Meteorological and hydrodynamic variables during the SOAP voyage, including a) Wind speed (WS, ms^{-1}); b) Direction (Dir., $^{\circ}$); Wind (blue) and wave (cyan); c) Temperature (Temp., $^{\circ}\text{C}$); Air (black) and surface water (green); d) Irradiance (Irrad., Wm^{-2}) and e) Significant wave height (Hs, m). The Bloom occupation periods are indicated by the red horizontal bars in the upper panel.




Figure 7. Surface water properties (2-10m) during the SOAP voyage: Temperature (Temp, °C), Mixed Layer Depth (ML Depth, m), Chlorophyll-a (Chl-a, mg m⁻³), and nitrate concentration (μmol l⁻¹), with the occupation period for each bloom indicated by the vertical shaded bars.




Fig. 8. SOAP parameters and integrated work programmes




Figure 9. Aerosol characterisation during SOAP indicating size spectral (red) & total counts (black). Ambient RH measurement was used for RH correction of the PCASP, Hi Vol and SMPS, and diffusion driers (Silica Gel) were used on the inlet of the UFO-TDMA and VH-TDMA.




Figure 10. a) Marine boundary layer CN concentrations (top, CPC3772 in blue, CPC3010 in red), b) CCN concentrations (middle) and c) number of hours 72 hour back-trajectory was over land (bottom, 27-member ensemble average).

