

1 **Supplementary Information for**

2 **A water vapor modulated aerosol impact on ice crystal size**

3

4 **Bin Zhao<sup>1</sup>, Kuo-Nan Liou<sup>1</sup>, Yu Gu<sup>1</sup>, Jonathan H. Jiang<sup>2</sup>, Qinbin Li<sup>1</sup>, Rong Fu<sup>1</sup>, Lei**  
5 **Huang<sup>1,2</sup>, Xiaohong Liu<sup>3</sup>, Xiangjun Shi<sup>3</sup>, Hui Su<sup>2</sup>, and Cenlin He<sup>1</sup>**

6 [1] Joint Institute for Regional Earth System Science and Engineering and Department of  
7 Atmospheric and Oceanic Sciences, University of California, Los Angeles, California 90095,  
8 USA.

9 [2] Jet propulsion Laboratory, California Institute of Technology, Pasadena, California 91109,  
10 USA.

11 [3] Department of Atmospheric Science, University of Wyoming, Laramie, Wyoming 82071,  
12 USA.


13

14 Correspondence to: Bin Zhao (zhaob1206@ucla.edu)


15

16


1 **Figures and Tables**



2  
3 Figure S1. The spatial domain of this study: 15°-55° N, 70°-135° E.



1 Figure S2. Changes in  $R_{ci}$  as a function of AOD for different ranges of (a) relative humidity  
 2 (RH) averaged between 100 hPa and 440 hPa, (b) the convective available potential energy  
 3 (CAPE), (c) the middle cloud layer temperature ( $T_{mid}$ ), (d) the vertical velocity at 500 hPa  
 4 (VV500), (e) the vertical velocity at 300 hPa (VV300), (f) the U-components of wind speed at  
 5 200 hPa (U200), (g) the U-components of wind speed at 1000 hPa (U1000), (h) the V-  
 6 components of wind speed at 200 hPa (V200), (i) the V-components of wind speed at 1000  
 7 hPa (V1000), (j) and the vertical wind shear (VWSH) at potential vorticity surface of  $2 \times 10^{-6}$   
 8  $\text{deg K m}^2 \text{kg}^{-1} \text{s}^{-1}$ . The meteorological parameters are divided into 3 ranges containing similar  
 9 numbers of data points, and the curves for the medium meteorological range are not shown.  
 10 The definition of error bars is the same as in Fig. 1 in the main text.



1 Figure S3. Changes in AOD as a function of meteorological parameters: (a) RH averaged  
 2 between 100 hPa and 440 hPa, (b) CAPE, (c)  $T_{\text{mid}}$ , (d) VV500, (e) VV300, (f) U200, (g)  
 3 U1000, (h) V200, (i) V1000, and (j) VWSH at the potential vorticity surface of  $2 \times 10^{-6}$  deg K  
 4  $\text{m}^2 \text{kg}^{-1} \text{s}^{-1}$ . The definition of error bars is the same as in Fig. 1 in the main text. Note that the  
 5 error bars in some panels are very small and hence not visible.  
 6

1 Table S1. Datasets used in this study.

Satellite/ Sensor	Product	Variable	Horizontal resolution
Aqua/MODIS	MYD04 (Level 2, Collection 6)	Column AOD	10 km × 10 km
	MYD06 (Level 2, Collection 6)	Cloud effective radius, cloud phase (determined by the “cloud optical property” algorithm), primary cloud retrieval outcome	1 km × 1 km
CALIPSO/ CALIOP	05kmMLay (Level 2, Version 4.10)	Aerosol/cloud layer number, layer base temperature, middle layer temperature, layer top/base height, layer aerosol/cloud optical depth, feature classification flags, CAD score, extinction QC	5 km along-track
	05kmAPro (Level 2, Version 4.10)	Vertically resolved pressure, relative humidity, and temperature	5 km along-track
--	NCEP ds083.2	Vertically resolved vertical velocity and wind speed; CAPE, wind shear	1° × 1°

2

3 Table S2. Correlation coefficients between various meteorological parameters.

	RH	CAPE	U200	T <sub>mid</sub>
RH	--	0.514	-0.535	-0.352
CAPE	0.514	--	-0.623	-0.390
U200	-0.535	-0.623	--	0.502
T <sub>mid</sub>	-0.352	-0.390	0.502	--

4 Note:  $p < 0.01$  for all cases. RH, relative humidity averaged between 100 hPa and 440 hPa; CAPE,  
5 convective available potential energy; U200, U-components of wind speed at 200 hPa; T<sub>mid</sub>, middle cloud  
6 layer temperature.

7