

acp-2017-568: Modelling carbonaceous aerosol from residential solid fuel burning with different assumptions for emissions

Ots, R. et al.

We thank the reviewer for their further review of our paper and are happy to provide more information in our responses here and to the manuscript. The most substantial clarifying additions can be found in Section 2.1 – Model description (including emissions description) and Section 2.3 – Comparison with measurements. The reviewer’s comments are presented in full below, in blue font.

Response to further review comments

The authors have provided lengthy replies, but I am afraid I still have two main issues with this paper:

i) I do not believe the authors have adequately addressed my comments about volatility. As explained below, they ignore the possibly very significant effect of evaporation of SFOA emissions. I would like to know what the UK POA emissions are, and to what extent they contain condensables. This is a reasonable request I think for a paper that focuses on such emissions from one country.

The UK POA emissions do not include condensables; we apologise for not making this explicit in our responses to the previous version in which we had argued that model simulations with (semi)volatile SFOA emissions were not required for our model-measurement comparison. We have now added text to Section 2.1 Model Description as follows.

First, we have changed the sentence that originally read “*It is likely that domestic PM_{2.5} emissions reported to CEIP vary by country according to whether condensables are included in the PM_{2.5} emissions but the specific information is not known*” to now read “*The UK emissions inventory for domestic PM_{2.5} does not include condensables, but this information is not known for the emissions reported to CEIP by other countries.*”

Secondly, we have now added the following paragraph to this section:

“The reason this study does not present model simulations with the volatile treatment of SFOA is that for the AMS-PMF data, primary (SF)OA and oxygenated (secondary) OA are separated. Therefore, the direct comparisons with SFOA measurements here do not include the semivolatile components as those would only become condensed after atmospheric ageing but then they would be measured as oxygenated OA, not SFOA. The volatile components and secondary OA precursors are not needed to test the main hypothesis of this paper – that the spatial distribution of wood and coal burning emissions should not be assumed to be zero in smoke control areas. Using primary components to demonstrate this is sufficient. This is not to say that SFOA emissions do not include precursors for SOA. The inclusion of semivolatile SOA precursors from SFOA is of course necessary to close the gap between total measured OA and total modelled OA. Indeed, the work by Xu et al. (2016) acknowledge that oxygenated OA likely contains secondary and/or aged SFOA.”

ii) The authors are responding to the referees, but often without any change in the paper (most replies to my comments were to me, not readers). The usual procedure is to change the paper

to clarify the points raised by the referees, since one cannot expect readers to churn through the whole ACPD discussion.

We had made additions to the paper where we thought these necessary but are happy to include more information in the manuscript (detailed below) whilst remaining conscious not to overly lengthen a paper that is already quite long. We have also referenced relevant previous studies as sources of further information for readers.

Our responses to the further specific comments made in this second review are given below. First, however, we indicate where we have added further material to the paper relating to comments made at the first review.

In Section 2.1, we have now directly referred to the published descriptions of the modelling framework for ASOA and BSOA by inserting the following sentence “*Descriptions of the modelling framework for ASOA and BSOA are given in Simpson et al. (2012) and Ots et al. (2016).*” before the sentence beginning “*In all the experiments presented here...*”

In Section 3.2, where text originally read “*This temporal misclassification of emissions for the UK is expected to have a detrimental effect on all of the model hourly evaluation statistics, except NMB*” we have now expanded this to read “*Whilst reducing boundary layer height is likely also to contribute to greater measured concentrations in late evening, the temporal misclassification of emissions for the UK is expected to have a detrimental effect on all of the model hourly evaluation statistics, except NMB.*”

Regarding the question in the previous review about convolved factors, we have now included the following text to Section 2.3 after this sentence that is currently in the paper: Limitations and uncertainties of these measurement datasets have been discussed in Ots et al. (2016a, b).

“In addition, the original analysis of the annual c-ToF-AMS (compact) dataset in Young et al. (2015a) identified that the SFOA and OOA factors were convolved. Section 4.4 of Young et al. (2015a) describes how these two convolved factors were dealt with. In brief, both factors had a strong and similar diurnal cycle, so the effect of being convolved was reduced by using daily averaged concentrations. In the current work presented here, these daily average concentrations are used for the annual comparisons, thus the issue of the convolved factors should not be significantly influencing the overall observations. In contrast, concentration data from the HR-ToF-AMS (high-resolution) instrument are used for the hourly comparisons presented in the current work. The PMF factors resulting from the HR-AMS data were not found to be convolved, which is likely due to a combination of the fact that the measurements were high resolution and the HR-ToF-AMS was deployed only during the intensive observation periods, rather than the full year like the c-ToF-AMS. Consequently, hourly comparisons presented in this work use the HR-ToF-AMS data.”

In more depth on the first point, the authors comment that:

a) ..we have not ‘ignored’ the issue of volatility because it is not important but because it is not needed in this examination of spatial patterns and magnitudes of primary SFOA PM2.5 emissions against primary SFOA measurements.

b) In this study we included the best-available county-by-county domestic sector PM2.5 emissions as reported to CEIP. This includes instances of countries that do report condensables as part of their PM emissions, e.g. Belgium (who report these as part of the total

PM emissions, not separately), and countries that don't. Without considerable more coordination of measurement and reporting it not possible to make any evidence-based adjustments to the CEIP emissions.

While the statement in (a) about spatial patterns may be true, the one about magnitudes certainly is not. It would only be true if the emissions included in PM_{2.5} (especially in the UK) were indeed non-condensable. This may be true for some countries, and is not for others, but this paper does not demonstrate the UK situation in any way. The statement in (b) does not help, and the comments about European emissions are misleading I think. This study is not really about European emissions; the focus is very much on the UK and indeed London. With such a focus I would have expected a proper explanation of what the United Kingdom does with regard to reporting condensables or not in its PM_{2.5} inventory. Did the authors speak to the emission inventory experts in the UK to clarify this problem?

We respond to all these comments together. (i) Yes we did speak to a UK national emissions expert and confirmed that the UK PM_{2.5} emissions do not include condensables. As indicated above, we have now added this fact explicitly to the paper. In addition, one of the authors of this paper, Jeroen Kuenen, is an emissions expert. We prefer not to refer to unpublished work too much which is why we used Belgium as an example of a country that does include condensables in their PM emissions reported to CEIP, and which is published.

(ii) As stated on the first page of our responses, we have now added a paragraph to 'Section 2.1 Methods Description' to explain why this study does not consider condensable emissions, the paragraph starting: *"The reason this study does not present model simulations with the volatile treatment of SFOA...."*

(iii) We have also now added the following extra text about emissions to Section 2.1 Model Description:

"Furthermore, the various sampling methods used to derive emission factors (which are applied by each country reporting emissions to CEIP) vary greatly (Denier van der Gon et al., 2015). The two main types are filter measurements (capturing only solid particles), and dilution tunnel measurements (capturing solid particles and condensable organics). The difference between the two methods can be large - up to 5-fold for wood-burning (Denier van der Gon et al., 2015), which is similar to what was shown by May et al. (2013) that up to 80% of the mass of POA from biomass burning may evaporate when diluted from plume to ambient conditions. Based on partitioning calculations for a range of OA (from 0.01 to 10⁵ ug m⁻³) in Donahue et al. (2006), only the 10 ug m⁻³ volatility bin exhibits a substantial portions in both gaseous and solid form. The organic components in all other bins are either mostly solid or mostly gaseous. May et al. (2013) also shows that only 10% of biomass burning emissions (Table 2 in May et al. (2013)) is in the 10 ug m⁻³ bin. Therefore, the potential overestimation arising from not letting these solids in the UK emissions inventory evaporate is no more than 10% (as the lower volatility materials is almost completely solid, and the higher volatility material is almost completely gaseous)."

(iv) In the previous revision we had also added text to make the point that without considerable more coordination of measurement and reporting (of domestic PM_{2.5} emissions) it is not possible to make any evidence-based adjustments to the CEIP emissions.

It is not clear that the authors have understood the implications of condensable/semi-volatile

POA concepts, since they only discuss (mainly in the reply to referees) aging. Evaporation is very likely to be the main issue with this work, not aging. According to May et al 2013, between 50-80% of the mass of POA from biomass burning may evaporate when diluted from plume to ambient conditions. The AMS instruments measure ambient air, so any study that ignores this will necessarily give too big a contribution from SFOA.

The measured amounts of SFOA against which we compare our model simulations are taken directly from the AMS-PMF speciation of OA components into HOA, SFOA, OA, COA. The AMS instrument measures only the condensed (aerosol particle) material in the ambient air. In the *Base* run, the model underestimates the measured (condensed) SFOA concentrations by 71% on average. In our simulation experiments we show various ways (by increasing the total amount as well as redistributing) to bring the model values of SFOA closer to what the AMS quantifies for this component. The non-volatile treatment of SFOA in the model does not make the AMS see more SFOA than there really is. If a proportion of the emissions going into the model are in reality being lost through evaporation then there would be greater discrepancy between model and measurement to account for.

In summary: the evaporation of semivolatile SFOA does not affect how much condensed SFOA the AMS measures. Our work is exploring ways of closing a model underestimation in condensed SFOA, not an overestimation.

Finally, I also saw the other referees suggestion to remove the Marleybone road plot from Fig.13, and agree with this. Having stated that 'the modelled concentrations are not fairly comparable due to differences in spatial scale (major road vs 5 km model resolution)', there is no valid argument to compare I think. The authors suggest that one needs to illustrate the range of concentrations in a megacity, but plenty of other studies do that. And major roads have little to do with SFOA modelling.

Response: We have now removed Marylebone Road from Figure 13 and deleted the text in this section that referred to that particular dataset.

Modelling carbonaceous aerosol from residential solid fuel burning with different assumptions for emissions

Riinu Ots^{1,2,*}, Mathew R. Heal¹, Dominique E. Young^{3,**}, Leah R. Williams⁴, James D. Allan^{3,5}, Eiko Nemitz², Chiara Di Marco², Anais Detournay², Lu Xu^{6,***}, Nga L. Ng^{6,7}, Hugh Coe³, Scott C. Herndon⁴, Ian A. Mackenzie⁸, David C. Green⁹, Jeroen J. P. Kuenen¹⁰, Stefan Reis^{2,11}, and Massimo Vieno²

¹School of Chemistry, University of Edinburgh, Edinburgh, UK

²Natural Environment Research Council, Centre for Ecology & Hydrology, Penicuik, UK

³School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester, UK

⁴Aerodyne Research, Inc., Billerica, MA, USA

⁵National Centre for Atmospheric Science, University of Manchester, Manchester, UK

⁶School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA, USA

⁷School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

⁸School of GeoSciences, University of Edinburgh, Edinburgh, UK

⁹MRC PHE Centre for Environment and Health, King's College London, London, UK

¹⁰TNO, Department of Climate, Air and Sustainability, Utrecht, The Netherlands

¹¹University of Exeter Medical School, European Centre for Environment and Health, Knowledge Spa, Truro, UK

* now at: Clinical Surgery, University of Edinburgh, Edinburgh, UK

** now at: Air Quality Research Center, University of California, Davis, CA, USA

*** now at: Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA.

Correspondence to: M. Heal (M.Heal@ed.ac.uk) and R. Ots (R.Ots@ed.ac.uk)

Abstract. Evidence is accumulating that emissions of primary particulate matter (PM) from residential wood and coal combustion in the UK may be underestimated and/or spatially misclassified. In this study, different assumptions for the spatial distribution and total emission of PM from solid fuel (wood and coal) burning in the UK were tested using an atmospheric chemical transport model. Modelled concentrations of the PM components were compared with measurements from aerosol mass spectrometers at four sites in central and Greater London (ClearfLo campaign, 2012), as well as with measurements from the UK black carbon network.

The two main alternative emission scenarios modelled were *Base4x* and *combRedist*. For *Base4x*, officially reported PM_{2.5} from the residential and other non-industrial combustion source sector were increased by a factor of 4. For the *combRedist* experiment, half of the baseline emissions from this same source were redistributed by residential population density to simulate the effect of allocating some emissions to the smoke control areas (that are assumed in the national inventory to have no emissions from this source). The *Base4x* scenario yielded better daily and hourly correlations with measurements than the *combRedist* scenario for year-long comparisons of the solid fuel organic aerosol (SFOA) component at the two London sites. However, the latter scenario better captured mean measured concentrations across all four sites. A third experiment, *Redist* - all emissions redistributed linearly to population density, is also presented as an indicator of the maximum concentrations an assumption like this could yield.

The modelled elemental carbon (EC) concentrations derived from the *combRedist* experiments also compared well with seasonal-average concentrations of black carbon observed across the network of UK sites. Together, the two model scenario simulations of SFOA and EC suggest both that residential solid-fuel emissions may be higher than inventory estimates and that the spatial distribution of residential solid-fuel burning emissions, particularly in smoke control areas, needs re-evaluation.

5 The model results also suggest the assumed temporal profiles for residential emissions may require review to place greater emphasis on evening (including ‘discretionary’) solid-fuel burning.

1 Introduction

Globally, wood and coal burning from residential heating and from cooking activities is a major source of both indoor and outdoor PM_{2.5} (particulate matter with diameter < 2.5 μm) air pollution (WHO, 2015). As developed countries commit to

10 renewable energy targets, the use of wood and biomass in residential heating is likely to increase (replacing some of the natural gas based heating systems; (WHO, 2015)). Within Europe, residential wood burning is estimated to be the single largest anthropogenic primary source of organic carbon (OC), contributing ~60% of total OC emissions from European countries (Denier van der Gon et al., 2015). In some countries, both wood and coal are burned in residential stoves and other small combustion plants. A number of particle source apportionment studies, particularly those based on positive matrix factorisation

15 of aerosol mass spectrometry measurements (AMS-PMF studies), have thus attributed organic aerosol (OA) from this source as solid fuel OA (SFOA), which is a combination of the commonly known biomass burning OA (BBOA) factor plus coal burning OA (Allan et al., 2010; Young et al., 2015a; Xu et al., 2016). In addition to heating, some residential solid fuel burning in European urban areas can also be attributed to recreation (i.e. fireplaces for ambience; Fuller et al. (2013)).

Since the Great London Smog of 1952 (Bell et al., 2004), several legislative interventions have substantially reduced the use

20 of solid fuels for residential heating in the UK by subsidising infrastructure and availability of oil and natural gas, as well as the implementation of smoke control areas (Fuller et al., 2013). For example, almost all of London is now a smoke control area where solid fuel burning is prohibited unless undertaken in approved wood burners (Fuller et al., 2013). This control is applied only to appliances with a chimney; incidental sources such as bonfires or barbecues are permitted.

There is evidence, however, that the smoke control legislation is no longer actively enforced. Several recent studies have re-

25 ported substantial local contributions of emissions from solid fuel burning to particle concentrations in London coinciding with days of low temperature (Fuller et al., 2014; Crilley et al., 2015). This is relevant as, currently, the UK’s National Atmospheric Emissions Inventory (NAEI) assumes zero residential emissions of non-approved solid fuel burning in smoke control areas (i.e. that there is full legal compliance). Furthermore, the NAEI only includes estimates of emissions from officially-sold solid fuels (NAEI, 2013), but there is reason to believe that much fuel wood is not obtained through commercial outlets and falls outside

30 the economic administration and therefore is not included in official statistics (Denier van der Gon et al., 2015). For example, a recent UK Wood Use Survey concluded that the official national consumption of domestic wood fuel is underestimated by a factor of 3 (Waters, 2016). Furthermore, the survey showed that 31% of wood fuel is sourced from the informal “grey” market of wood (i.e. own garden, other landowners’ gardens, waste wood, etc.).

The results of the UK survey are in line with recent evaluations of wood burning emissions in Belgium, which also concluded that their official inventory has underestimated the amount of wood burned in residential settings by more than a factor of 3 (Lefebvre et al., 2016). Belgium has since included these increased emissions estimates (adding estimates for non-commercial wood sources to officially reported sales) in the amount reported to the Centre on Emission Inventories and Projections (CEIP) of the European Monitoring and Evaluation Programme (EMEP). The authors of the Belgian study urge countries that currently do not make estimates of non-officially traded or sourced wood to follow the same practise and start reporting these.

Model-measurement comparisons of a range of gaseous and particulate pollutants (Ots et al., 2016a) show that modelled concentrations of SFOA at the North Kensington site in London are substantially underestimated (normalised mean bias, NMB, of -71%) compared to an annual dataset of PMF apportionment of AMS measurements collected during the 2012 Clean Air for London campaign (ClearfLo; Bohnenstengel et al. (2014); Young et al. (2015a); Xu et al. (2016)). The aim of this work is to use these measurements of SFOA as the basis for an atmospheric chemical transport model exploration of potential closure of this discrepancy.

2 Methods

2.1 Model description

EMEP4UK is a regional application of the EMEP MSC-W (European Monitoring and Evaluation Programme Meteorological Synthesizing Centre-West) chemical transport model (Vieno et al., 2009, 2010, 2016; Ots et al., 2016a, b). The EMEP MSC-W model is a 3-D Eulerian model that has been used for both scientific studies and to support policy making in Europe. A detailed description of the EMEP MSC-W model, including references to evaluation and application studies is available in Simpson et al. (2012), Schulz et al. (2013), and at www.emep.int. The model configuration used here was based on version v4.5. The model has 21 vertical layers, extending from the surface to 100 hPa. The lowest vertical layer is ~ 40 m in height, and the horizontal resolution over an inner domain covering the British Isles is $5 \text{ km} \times 5 \text{ km}$. The model uses one-way nesting from an extended European domain (simulated with $50 \text{ km} \times 50 \text{ km}$ horizontal resolution). The model was driven by output from the Weather Research and Forecasting (WRF) Open Source model (www.wrf-model.org, version 3.1.1) including data assimilation of 6-hourly meteorological reanalysis from the US National Center for Environmental Prediction (NCEP)/National Center for Atmospheric Research (NCAR) Global Forecast System (GFS) at 1° ($\sim 100 \text{ km}$) resolution (NCEP, 2000).

Gridded anthropogenic emissions were obtained from NAEI (NAEI, 2013) for the UK, and from CEIP (CEIP, 2015) for the rest of Europe. All emissions are apportioned across a standard set of emission source sectors, following the sector structure defined in the Selected Nomenclature for Air Pollutant (SNAP, Table 1; EEA (2013)). Daily emissions from open burning (including wild fires and agricultural burning) were taken from the Fire INventory from NCAR version 1.0 (FINNv1, Wiedinmyer et al. (2011)). Primary PM emissions reported as $\text{PM}_{2.5}$ and PM_{10} in NAEI and CEIP were speciated into elemental carbon (EC), hydrocarbon-like OA (HOA) from fossil fuel combustion, OA from domestic combustion (SFOA = BBOA + coal OA) and remaining primary PM for each source sector using splits developed by Kuenen et al. (2014), as shown in Fig. 1. [Descriptions of the modelling framework for ASOA and BSOA are given in Simpson et al. \(2012\) and Ots et al. \(2016a\).](#) In all

the experiments presented here, SFOA is assumed to be non-volatile and it does not undergo atmospheric ageing. ~~This is not because it is assumed that there is no aging of SFOA emissions but because the model simulations compare against AMS-PMF determinations of primary SFOA concentrations. It is likely that domestic~~ The UK emissions inventory for domestic PM2.5 emissions reported to CEIP vary by country according to whether condensables are included in the PM2.5 emissions but the specific information is not known. 5 does not include condensables, but this information is not known for the emissions reported to CEIP by other countries.

The reason this study does not present model simulations with the volatile treatment of SFOA is that for the AMS-PMF data, primary (SF)OA and oxygenated (secondary) OA are separated. Therefore, the direct comparisons with SFOA measurements here do not include the semivolatile components as those would only become condensed after atmospheric ageing but then they would be measured as oxygenated OA, not SFOA. The volatile components and secondary OA precursors are not needed to test the main hypothesis of this paper – that the spatial distribution of wood and coal burning emissions should not be assumed to be zero in smoke control areas. Using primary components to demonstrate this is sufficient. This is not to say that SFOA emissions do not include precursors for SOA. The inclusion of semivolatile SOA precursors from SFOA is of course necessary to close the gap between total measured OA and total modelled OA. Indeed, the work by Xu et al. (2016) acknowledge that oxygenated OA likely contains secondary and/or aged SFOA.

Furthermore, the various sampling methods used to derive emission factors (which are applied by each country reporting emissions to CEIP) vary greatly (Denier van der Gon et al., 2015). The two main types are filter measurements (capturing only solid particles), and dilution tunnel measurements (capturing solid particles and condensable organics). The difference between the two methods can be large - up to 5-fold for woodburning (Denier van der Gon et al., 2015), which is similar to what was shown by May et al. (2013) that up to 80% of the mass of POA from biomass burning may evaporate when diluted from plume to ambient conditions. Based on partitioning calculations for a range of OA (from 0.01 to 10^5 ug m^{-3}) in Donahue et al. (2006), only the 10 ug m^{-3} volatility bin exhibits a substantial portions in both gaseous and solid form. The organic components in all other bins are either mostly solid or mostly gaseous. May et al. (2013) also shows that only 10% of biomass burning emissions (Table 2 in May et al. (2013)) is in the 10 ug m^{-3} bin. Therefore, the potential overestimation arising from not letting these solids in the UK emissions inventory evaporate is no more than 10% (as the lower volatility materials is almost completely solid, and the higher volatility material is almost completely gaseous).

The performance of this version of the EMEP4UK model simulating a standard suite of gas-phase components and secondary inorganic aerosol PM components is reported in Ots et al. (2016a) comparing with a full year of measurements in London in 2012. In brief, Ots et al. (2016a) reported an NMB of -1% and $r = 0.79$ for ozone, an NMB of -32% and $r = 0.78$ for NO_x , an NMB of $+6\%$ and $r = 0.73$ for SO_4^{2-} , an NMB of -12% and $r = 0.65$ for NH_4^+ , and an NMB of -23% and $r = 0.57$ for NO_3^- .

2.2 Model experiments

In this study, four different cases were considered. The *Base* case model experiment uses the same emission inventory dataset as Ots et al. (2016a) (i.e. as reported by the NAEI using the splits in Fig. 1), but with a small adjustment in the daily variation

Table 1. SNAP source sectors as specified in the emissions input to the model (CEIP, 2015).

SNAP1	Combustion in energy and transformation industries
SNAP2	Residential and non-industrial combustion
SNAP3	Combustion in manufacturing industry
SNAP4	Production processes
SNAP5	Extraction and distribution of fossil fuels
SNAP6	Solvent and other product use
SNAP7	Road transport
SNAP8	Other mobile sources and machinery
SNAP9	Waste treatment and disposal
SNAP10	Agriculture




Figure 1. Annual UK PM_{2.5} emissions by SNAP sector as specified in the NAEI (for year 2012), with each sector split into primary OA (HOA or SFOA), EC, and remaining PM following Kuenen et al. (2014). Source: Ots et al. (2016a).

in emissions due to temperature, called degree-day factors (Simpson et al., 2012). Recent studies in London have shown that local contributions of SFOA coincide the days of low temperature (Fuller et al., 2014; Crilley et al., 2015). Therefore, degree-day factors were included to modulate the daily variation in emissions from the SNAP2 sector according to ambient temperature (i.e. increasing the emissions during colder days). SNAP2 includes PM_{2.5} emissions from both residential and small (non-industrial) commercial combustion, but the residential part dominates with annual emissions for 2012 of 19 Gg from residential, 1 Gg from commercial and 5 Gg from stationary military combustion (NAEI, 2013). The domination of the residential emissions means that no large additional uncertainty is introduced by applying the degree-day factors to the whole of SNAP2.

A degree-day is defined as $H_{dd,j} = \max(18^\circ\text{C} - T_j^{24h}, 1)$, where j is the day number and T_j^{24h} is the daily averaged temperature in $^\circ\text{C}$. These degree-days are divided by the annual mean ($\overline{H_{dd}}$) to obtain degree-day factors (Simpson et al., 2012). For this work, degree-day factors pre-calculated by the EMEP MSC-W Centre based on European Centre for Medium-Range Weather Forecasts (ECMWF) 2012 meteorological simulations for the 50 km \times 50 km domain (Schulz et al., 2013) were

Table 2. Summary of the four experiments for PM_{2.5} emissions from SNAP2. In all experiments, PM_{2.5} is split into three components as follows: 48% is SFOA, 26% EC, and 26% other/mineral PM (as in Fig. 1).

Experiment	SNAP2 PM _{2.5} emission	Spatial distribution
<i>Base</i>	25 Gg	NAEI
<i>Base4x</i>	100 Gg	NAEI
<i>Redist</i>	25 Gg	Population density
<i>combRedist</i>	25 Gg	NAEI + Population density

disaggregated using the simple area-weighting method assuming homogeneity for degree-day factors within the 50 km to 5 km conversion.

In the second model experiment (*Base4x*), emissions from SNAP2 were increased by a factor of 4 (based on the NMB of -71% at the London North Kensington site). SNAP2 is the only sector with SFOA emissions in the set-up used (Fig. 1). The spatial distribution was unchanged.

For the third experiment (*Redist*), the annual reported PM_{2.5} emissions from SNAP2 were re-gridded linearly by residential population density (census data from Reis et al. (2016)) in order to test the assumption made by the NAEI that only smokeless fuels are used in smoke control areas. The total emission was unchanged.

Finally, the fourth experiment, *combRedist*, was a hybrid of the *Base* and *Redist* experiments, where, for each grid cell, half the emissions of *Base* and half of *Redist* (of the SNAP2 sector) were added together, i.e. $combRedist_{emis} = 0.5 \times Base_{emis} + 0.5 \times Redist_{emis}$.

The experiments *Base*, *Redist*, and *combRedist* therefore use the same total emission; the only difference is in the spatial distribution of the emissions. All four experiments use the same temporal variation for SNAP2, including degree-day factors. The emissions used for the four experiments are summarised in Table 2. In all of these experiments, SFOA is assumed to be non-volatile (see Sect. 2.1 for more information on this). The emission maps of these four experiments are shown in Fig. 2, and the resulting modelled annual average surface concentrations of SFOA are shown in Fig. 3.

2.3 Comparison with measurements

Modelled OA_{2.5} (OA with diameter $< 2.5 \mu\text{m}$) is compared with non-refractory submicron (NR-PM₁) OA measured by Aerodyne AMS instruments at the London North Kensington and London Marylebone Road sites in central London (urban background and roadside, respectively), and at the Harwell and Detling rural sites located to the west and east of London, respectively (Bohnenstengel et al., 2014; Young et al., 2015a, b; Xu et al., 2016). The locations of the AMS measurement sites used in this study are shown in Fig. 4. The discrepancy introduced by the different size fractions of modelled and measured concentrations is small as at an urban background site, $\sim 90\%$ of organic carbon in PM_{2.5} is in the submicron fraction (Harrison and Yin, 2008).




Figure 2. Total SFOA emissions (defined as 48% of $PM_{2.5}$ from SNAP2) for the year 2012 in the inner nesting domain for the four scenarios of this study: (a) *Base*: as in the NAEI, (b) *Base4x*: Base increased by a factor of 4 over the whole of UK, (c) *Redist*: UK emissions redistributed to residential population density (national total same as Base), (d) *combRedist*: half of the total emission redistributed to residential population density, half as reported by NAEI (a combination of (a) and (c)). The national total is the same for (a), (c), and (d). All UK emissions are aggregated to the $5 \text{ km} \times 5 \text{ km}$ grid (20 Mg per grid square is 0.8 Mg km^{-2}) from an initial resolution of $1 \text{ km} \times 1 \text{ km}$ reported by the NAEI. The emission resolution is $50 \text{ km} \times 50 \text{ km}$ for other countries (as in CEIP). Note the non-linear scale.

Different types of AMS instruments were deployed in the ClearLo campaign. At the London North Kensington site a compact time-of-flight AMS (cToF-AMS) was deployed for a full calendar year (January 2012 – January 2013), and an additional high-resolution time-of-flight AMS (HR-ToF-AMS, DeCarlo et al. (2006)) was deployed for the winter Intensive Observation Period (IOP). The measurements at Marylebone Road were taken with a Q-AMS (Quadrupole AMS; Jayne et al. (2000)).

5 HR-ToF-AMS instruments were deployed at Detling and Harwell during the winter IOP (not the whole year). PMF analysis was applied to each dataset to apportion measured OA into different components (Ulbrich et al., 2009). A detailed description of the derivation and optimization of the factors retrieved from the AMS data at Detling can be found in Xu et al. (2016), at London North Kensington in Young et al. (2015a) and Young et al. (2015b), at London Marylebone Road in Detournay et al. (2017) (all of these analyses were performed with the PMF2 solver), and at Harwell in Di Marco et al. (2017) (using

10 the ME-2 solver). The limitations and uncertainties of these measurement datasets have been discussed in Ots et al. (2016a,




Figure 3. Annual average modelled surface SFOA concentrations for the year 2012 using the emission scenarios shown in Fig. 2: (a) *Base*: as in the NAEI, (b) *Base4x*: *Base* increased by a factor of 4 over the whole of UK, (c) *Redist*: UK emissions redistributed by residential population density (national total same as *Base*), (d) *combRedist*: half of the total emission redistributed by residential population density, half as reported by NAEI (a combination of (a) and (c)). The national total is the same for (a), (c), and (d).

b) . In addition, the original analysis of the annual c-ToF-AMS (compact) dataset in Young et al. (2015a) identified that the SFOA and OOA factors were convolved. Section 4.4 of Young et al. (2015a) describes how these two convolved factors were dealt with. In brief, both factors had a strong and similar diurnal cycle, so the effect of being convolved was reduced by using daily averaged concentrations. In the current work presented here, these daily average concentrations are used for the annual comparisons, thus the issue of the convolved factors should not be significantly influencing the overall observations. In contrast, concentration data from the HR-ToF-AMS (high-resolution) instrument are used for the hourly comparisons presented in the current work. The PMF factors resulting from the HR-AMS data were not found to be convolved, which is likely due to a combination of the fact that the measurements were high resolution and the HR-ToF-AMS was deployed only during the intensive observation periods, rather than the full year like the c-ToF-AMS. Consequently, hourly comparisons presented in this work use the HR-ToF-AMS data.

The following numerical metrics were used for model evaluation: FAC2 (Factor of 2) - the proportion of modelled concentrations that are within a factor of 2 of the measured concentrations; NMB - normalised mean bias; NMGE - normalised mean



Figure 4. Locations of the ClearfLo measurement sites used in this work. London North Kensington is an urban background site, London Marylebone Road is a roadside site, Harwell and Detling are rural background sites (Source: Ots et al. (2016a)). Underlying map from © OpenStreetMap contributors.

gross error, which is defined as:

$$NMGE = \frac{\frac{1}{n} \sum_{i=1}^n |M_i - O_i|}{\bar{O}}, \quad (1)$$

where M_i is the i th modelled value, O_i is the corresponding measured value, \bar{O} is the mean measured value, and n is the total number of observations; r - correlation coefficient; and COE - coefficient of efficiency, which is defined as:

$$COE = 1.0 - \frac{\sum_{i=1}^n |M_i - O_i|}{\sum_{i=1}^n |\bar{O} - O_i|}. \quad (2)$$

A COE of 1 indicates perfect agreement between model and measurements. Although the COE does not have a lower bound, a zero or negative COE implies that the model cannot explain any of the variation in the observations (Legates and McCabe, 2013).

3 Results and Discussion

Figure 3 shows the annual mean modelled SFOA surface concentrations for the year 2012. In the model, SFOA is emitted as 48% of $PM_{2.5}$ from the SNAP2 source sector, and it is advected as a non-volatile and chemically inert species (but it is included in the total OA budget for the absorptive partitioning of secondary organic aerosol species). The gradients of SFOA surface concentrations visible over the North Sea and the English Channel are indicators of European transport. Over the UK, SFOA concentrations follow the pattern of the prescribed local emissions, with the spatial distributions of the experiments with and without redistributed national emissions being substantially different from each other. The *Base* and *Base4x* scenarios (spatially gridded as reported by the NAEI) assigned most emissions to Northern England, Wales and Northern Ireland, leaving

major cities such as London, Manchester, Leeds, Birmingham, and Glasgow almost unnoticeable in the concentration fields (these, and many other urban locations, are designated as smoke control areas). In contrast, the *Redist* experiment highlights all of these urban areas, because the SFOA emissions were redistributed linearly by residential population density. The *combRedist* experiment shows these residential hot-spots while also retaining some of the spatial pattern from the officially reported distribution.

In our experiments, we did not modify European emissions, we used exactly what has been reported to the CEIP. While there is reason to believe European emissions of SFOA are also underreported, we do not believe this to have a major influence on the surface concentrations of SFOA over the UK as even our Base4x experiment (Fig. 3b) only indicates very modest regional transport of our SFOA to Europe.

3.1 Daily evaluation - London Marylebone Road and North Kensington annual datasets

Time-series of measured and modelled daily-average SFOA concentrations for the London Marylebone Road and North Kensington sites are shown in Fig. 5 (*Base*, and *Base4x*) and Fig. 6 (*Base*, and *combRedist*). These figures also highlight the date of the annual bonfire celebrations around Guy Fawkes Night (5th of November) to draw attention to an increase in SFOA emissions that can not be simulated with the model, given that the temporal variation is prescribed using a regular approach (i.e. hour-of-day, day-of-week, month-of-year) which does not include information about specific days and events. (The small difference between the *Base* experiment time-series in these figures and a similar comparison in Ots et al. (2016a) is due to the use of degree-day factors.) For convenience, the evaluation statistics presented on these time-series (as well as for the *Redist* experiment) are given in Table 3.

These two sites are ~ 4.5km apart, in adjacent model grid cells, and represent different kinds of urban areas. North Kensington is comparatively more residential whereas Marylebone Road is near central London and therefore exhibits very high numbers of people during the workday. More discussion on this, including detailed maps can be found in Ots et al. (2016b)

Table 3. Evaluation statistics for modelled vs measured daily-average concentrations of SFOA for the ClearfLo year-long datasets (year 2012).

Site	Experiment	NMB	NMGE	r	COE
Marylebone Road	<i>Base</i>	-59%	64%	0.60	-0.23
	<i>Base4x</i>	14%	54%	0.67	-0.02
	<i>Redist</i>	57%	79%	0.57	-0.51
	<i>combRedist</i>	-1%	51%	0.59	0.03
North Kensington	<i>Base</i>	-69%	70%	0.73	0.03
	<i>Base4x</i>	-18%	42%	0.78	0.42
	<i>Redist</i>	33%	57%	0.71	0.21
	<i>combRedist</i>	-18%	43%	0.73	0.41




Figure 5. Time-series of measured and modelled (*Base* and *Base4x* experiments) daily-average SFOA concentrations at the (a) Marylebone Road, and (b) North Kensington measurement sites, year 2012. The vertical line marks the date of ‘bonfire night’, November 5th. The annual average measured SFOA concentrations at these sites were $0.9 \mu\text{g m}^{-3}$ at Marylebone Road, and $1.0 \mu\text{g m}^{-3}$ at North Kensington.




Figure 6. Time-series of measured and modelled (*Base* and *combRedist* experiments) daily-average SFOA concentrations at the (a) Marylebone Road, and (b) North Kensington measurement sites, year 2012. The vertical line marks the date of ‘bonfire night’, November 5th. The annual average measured SFOA concentrations at these sites were $0.9 \mu\text{g m}^{-3}$ at Marylebone Road, and $1.0 \mu\text{g m}^{-3}$ at North Kensington.

The experiments result in better daily-average SFOA model-measurements agreement than the *Base* case at the two sites. The only exception is the *Redist* simulation at Marylebone Road, where the underestimation of the *Base* case is replaced with an equivalent overestimation (NMB of -59% and $+57\%$, *Base* and *Redist*, respectively, Table 3). There is a small decrease in the r -value, and an increase of the NMGE, which is caused by the modelled values of *Redist* being greater than those of the *Base* experiment. For the London North Kensington site, the *Redist* experiment is an improvement compared to the *Base* run, although the concentrations are also overestimated (NMB = $+33\%$, *Redist*). This is expected, as areas with high population densities would include large apartment buildings which are unlikely to have individual fireplaces. Therefore a completely linear redistribution of residential emissions is not correct, but this experiment gives an indication of the maximum effect that population density could have on SFOA concentrations.

Both the *Base4x* and *combRedist* experiments have better predictive abilities for the AMS-PMF measured concentrations of SFOA at the two sites in London than the *Base* case emissions simulation. The NMGE at the Marylebone Road site is reduced to 54% (*Base4x*) or 51% (*combRedist*), compared to 64% for the *Base* case simulation. At the North Kensington site, NMGE is reduced to 42% (*Base4x*) or 43% (*combRedist*), compared to 70% for the *Base* case simulation. The *Base4x* results in improvements in the r -value: 0.67 (*Base*: 0.60) and 0.78 (*Base*: 0.73) at Marylebone and North Kensington, respectively, whereas with the *combRedist* emissions, the r -values of daily-average concentrations remain the same as *Base*: 0.59 (0.60) and 0.73 (0.73), at Marylebone Road and North Kensington, respectively. The improvement in COE values is similar for both experiments (Table 3) at both sites. Both experiments decreased the NMB at North Kensington to -18% (*Base4x* and *combRedist*) from -69% (*Base*), whereas at Marylebone Road, the *Base4x* reaches an overestimation of NMB = $+14\%$ while the *combRedist* matches the measured mean SFOA: NMB = -1% (NMB of *Base* at the Marylebone Road measurement site: -59%). It should be noted, however, that there are several days in November and December where both the *Base4x* and *combRedist* experiments overestimate SFOA concentrations compared to measurements.

In summary, in comparison with daily-average measurements of SFOA concentrations at two sites in London the *Base4x* and the *combRedist* experiments resulted in similar improvements in NMGE and COE, the *Base4x* experiment had better r -values, and the *combRedist* experiment better matched the annual mean concentrations of SFOA at the two sites. Nevertheless, it should be noted that AMS-PMF apportionment measurements are also subject to uncertainty which limits the expected correlation with the model. For example, Ots et al. 2016a presented scatter plots of daily averaged concentrations of the different OA components derived from measurements with the two different AMS instruments at the North Kensington site during the winter IOP (the cToF-AMS versus the HR-ToF-AMS). While these comparisons showed good correlations between the two measurements (0.88 to 0.95 for the primary OA components, 0.77 for secondary OA), on some days the absolute measured concentrations of specific components do differ, sometimes by more than a factor of two. For further discussion on this see Ots et al. (2016a, b).

The following sections evaluate these experiments with respect to hourly-average measurements taken with the High Resolution (HR-ToF-AMS) instruments during the ClearfLo winter IOP, which included two rural background sites - Harwell and Detling - as well as the London North Kensington site.

3.2 Hourly averaged diurnal profiles of SFOA concentrations, winter IOP

Hourly averaged diurnal profiles of measured and modelled SFOA concentrations at the ClearfLo winter IOP sites are shown in Fig. 7. The profiles of measured concentrations at the rural background sites (Detling and Harwell), and the urban background site (North Kensington) all show a pronounced maximum in the evening (after 18:00), and a much smaller peak at around 9:00. In the model, the hourly emission factors applied to the SNAP2 sector have similar magnitudes for morning and evening thus underestimating the higher evening concentrations seen in the measurements (in this work, the same diurnal emission profile was used for all countries). This Whilst reducing boundary layer height is likely also to contribute to greater measured concentrations in late evening, the temporal misclassification of emissions for the UK is expected to have a detrimental effect on all of the model hourly evaluation statistics, except NMB.




Figure 7. Hourly-averaged diurnal profiles of modelled and measured SFOA concentrations at the (a) Detling, (b) Harwell, and (c) North Kensington measurement sites, winter IOP 2012.

3.3 High SFOA episode: 13-Jan–18-Jan, 2012

Time-series of hourly-average measured and modelled (*Base*, *Base4x*, and *combRedist* experiments) SFOA concentrations at the ClearfLo winter IOP sites are shown in Fig. 8. Note the very high concentrations measured at Harwell from 13 to 18 Jan. North Kensington and Detling also exhibited elevated concentrations (although not as high as Harwell) on these days, especially on 17 Jan. During this episode, both the *Base4x* and *combRedist* experiments simulate similar concentrations for the North Kensington site, whereas for Harwell and Detling, the *Base4x* gives higher concentrations than *combRedist* and is therefore slightly closer to the very high measured concentrations. Note that *Base4x* only increased the UK emissions, not European ones. Hence the increase in *Base4x* concentrations compared with *Base* is not fourfold.

Daily-average maps of modelled SFOA surface concentrations during these days are shown in Fig. 9 (*Base4x*), and Fig. 10 (*combRedist*). Time-series of daily-average concentrations during the winter IOP are shown in Fig. 11. At Detling, the measurements commenced the morning of 16-Jan (Fig. 8a) but since we used a data capture threshold of 75%, 16-Jan and 17-Jan

did not include sufficient hourly data points to present measured daily averages for this site. Ots et al. (2016a) demonstrated that a rural background site can on occasion exhibit substantially higher concentrations than central London due to atmospheric import of polluted air masses from Europe creating a strong spatial gradient. The daily-average concentration maps, however, do not indicate European gradients over southern England during 13 to 18 Jan. Nevertheless, southern England did experience a sustained high-pressure weather system during these days, including noticeably lower temperatures for 14–18 Jan than average (Crilley et al. (2015): Fig. 2). Sustained high pressure usually leads to a very stable atmosphere with descending air masses. Therefore, these high concentrations could have been caused by meteorological build-up, and it is possible the model set-up underestimated the strength of this effect. Furthermore, 14–15 Jan was a weekend whereby people are more likely to spend time home and therefore potentially use their fireplaces more than on weekdays.

10 The exceptional concentrations measured especially at Harwell (and to a lesser extent at Detling) could have been caused by (i) missing local sources in the area, (ii) over-reporting of the concentrations by AMS measurements or by the PMF analysis




Figure 8. Time-series of measured and modelled hourly average SFOA concentrations at the (a) Detling, (b) Harwell, and (c) London North Kensington measurement sites, winter IOP 2012. Note the different scales on the y-axis.

applied to apportion measured OA into its components, (iii) meteorological build-up, or (iv) a combination of these. However, the specific origin of the large discrepancy between model and measurements at the Harwell site during these four days remains unknown.

3.4 Hourly evaluation statistics during the rest of the ClearfLo winter IOP, 2012

5 Table 4 presents the hourly evaluation statistics at the Detling, Harwell, and London North Kensington sites during the winter IOP (as in Fig. 8) but excluding the period of largely unexplained high SFOA concentrations episode between 13 Jan–18 Jan. These r -values (0.35–0.53; range of hourly r -values for all three sites) are lower than the daily-average r -values in comparison with the annual datasets shown in Figs. 5 and 6 (0.59–0.78). This is expected as the diurnal emissions profile used for all European countries assigns equal amounts of residential combustion emissions to the morning and evening which does not
 10 match the measured diurnal SFOA profiles at these three sites (as was shown in Fig. 7).

For Detling, the *Base* case run underestimates the mean measured SFOA concentration by $-27%$ (NMB), the *Base4x* experiment results in an overestimation of $+49%$, and the *combRedist* yields a close match with $-3%$. For Harwell, the *Base* scenario has a NMB of $-36%$, which becomes an overestimation of $+64%$ with the *Base4x* experiment. The *combRedist* emis-




Figure 9. Daily-average modelled (*Base4x* experiment) SFOA surface concentrations during the episode of high SFOA concentrations at the beginning of the winter IOP, year 2012. The black crosses mark the measurement site locations, left: Harwell, right: London North Kensington.




Figure 10. Same as Fig. 9, but for the *combRedist* experiment.




Figure 11. Daily-average measured and modelled SFOA concentrations at the (a) Detling, (b) Harwell, and (c) North Kensington measurement sites, year 2012. On panel (b), the concentration measured at 16-Jan is given as a text label.

sions have a minor effect on modelled concentrations at Harwell (NMB of -31% ; compared with -36% for the *Base* case; the other evaluation statistics of *combRedist* are also close to those of the *Base* case). At North Kensington, hourly comparison shows similar results for the two experiments as was also seen in the daily evaluation. Both emissions cases capture the mean concentration well, but the *Base4x* experiment yields a better r -value than *combRedist* experiment .

Table 4. Evaluation statistics for modelled vs measured hourly-average concentrations of SFOA during the ClearfLo winter IOP measurement sites as in Fig. 8, but excluding the period of 13 Jan–18 Jan.

Site	Experiment	NMB	NMGE	r	COE
Detling	<i>Base</i>	-27%	66%	0.38	0.19
	<i>Base4x</i>	48%	92%	0.45	-0.14
	<i>combRedist</i>	-3%	68%	0.41	0.16
Harwell	<i>Base</i>	-36%	69%	0.43	0.18
	<i>Base4x</i>	64%	105%	0.42	-0.24
	<i>combRedist</i>	-31%	68%	0.44	0.19
North Kensington	<i>Base</i>	-64%	66%	0.53	0.06
	<i>Base4x</i>	-1%	56%	0.53	0.20
	<i>combRedist</i>	12%	73%	0.35	-0.04

5 3.5 Comparison of modelled EC with measured EC and BC

Locations for the UK black carbon (BC) measurement network existing in 2012 are shown in Fig. 12. All of the measurement sites use an Aethalometer to measure optically-absorbing aerosol on a filter tape. Optical absorption is converted to an effective black carbon (eBC, referred to as just BC in this work) concentration using a mass-specific absorption cross-section (MAC). Three sites (Harwell, London North Kensington, and London Marylebone Road) also have daily-average filter measurements that are analysed for elemental carbon (EC) with a thermal optical technique in which the split between EC and OC can vary depending on instrumental and analysis parameters. Comparisons of the daily measurements of EC-R (measured EC corrected using reflectance), EC-T (corrected using transmittance), and daily-averaged BC are presented in Appendix A, along with a discussion of the sources of method biases. Substantial discrepancies in reported concentrations exist (at times, more than a factor of 2). Furthermore, the sign and the magnitude of discrepancies differ by season and by measurement site. Consequently, detailed (i.e. hourly or daily) model-measurement evaluation is not justified, and only seasonally-averaged concentrations are presented in this section.

Seasonal boxplots of daily-average concentrations of measured BC, EC-T, EC-R, and modelled concentrations of EC for Harwell ,~~North Kensington , and Marylebone Road~~ and North Kensington are shown in Fig. 13. In addition to emissions from domestic heating, these concentrations also include all other sources of EC, mainly traffic. Marylebone Road also measures




Figure 12. BC-network measurement site locations in 2012. Site names are abbreviated as follows: GLA - Glasgow Centre, STAB - Strabane, BEL - Belfast Centre, LISB - Lisburn Dunmurry, BIR - Birmingham Tyburn, NORW - Norwich Lakenfields, CAR - Cardiff, HAR - Harwell, NKEN - London North Kensington, MARY - London Marylebone Road. Underlying map from © OpenStreetMap contributors.

all these components, but since it is a roadside site, the traffic-influenced concentrations measured there are much higher than the $5 \text{ km} \times 5 \text{ km} \times 40 \text{ m}$ grid cell average. Therefore the measured and modelled concentrations of EC are not comparable for that site.

Within each panel, the different datasets were made to be of equal size, i.e. days with missing measurements, or with measurements below the limit of detection for EC-R or EC-T values, were also removed from measured BC as well as from the modelled time series.

~~Although the pollutant levels measured close to the traffic source at the Marylebone Road roadside site and the modelled concentrations are not fairly comparable due to the differences in spatial scale (major road vs 5 km model resolution), they are included to illustrate the range of concentrations in a megacity.~~

For Harwell, modelled EC concentrations from the *Base* and *combRedist* experiments are similar, but concentrations from the *Base4x* experiment are higher. This is consistent with the comparison of modelled and measured SFOA concentrations in the *Base4x* experiment which overestimates wood and coal burning contributions at the two rural sites near London (Harwell and Detling, excluding 13–18-Jan). Fig. 13 shows that the *Base* and *combRedist* modelled EC concentrations are close to measured EC-R during all seasons except autumn for which overestimations of several pollutants for many different days and periods are discussed in Ots et al. (2016a). Generally, measured BC concentrations are higher than measured or modelled

EC, but during winter, measured BC, measured EC-R and modelled EC (*Base* or *combRedist* experiments) are a close match, whereas measured EC-T is substantially lower.




Figure 13. Seasonal boxplots of daily-average concentrations of measured BC, EC-T, EC-R, and modelled EC concentrations at Harwell, [and London North Kensington, and London Marylebone Road, the three](#) measurement sites that measure both BC and EC, year 2012. Middle line: median, boxes: 25th and 75th quartiles (i.e. the interquartile range - IQR), whiskers extend to $1.5 \times \text{IQR}$, and all daily values beyond the whiskers are plotted individually.

At North Kensington, both the *Base4x* and *combRedist* experiments result in similarly higher concentrations than the *Base* run, and these experiments match the measurements of EC-R well. ~~At Marylebone Road, modelled concentrations are~~
 5 ~~substantially lower than are measured. This is expected as Marylebone Road is a roadside site with heavy traffic flows and the concentrations captured at the measurement site are therefore much higher than a $5 \text{ km} \times 5 \text{ km} \times 40 \text{ m}$ model grid cell average.~~

Seasonal boxplots of daily-average concentrations of measured BC and modelled EC at all the other BC network measurement sites are presented in Fig. 14. For these urban sites, both the *combRedist* case and the *Base4x* case yield a higher modelled
 10 concentration than the *Base* run and bring the modelled EC into better agreement with measured BC. There are no sites for which the *combRedist* yields a lower modelled concentration than the *Base* run (i.e. for these, urban, sites, the redistribution does not make the comparison worse). Therefore, based on BC measurements at these sites in different parts of the UK (which are on average higher than measurements of EC-T or EC-R), the experiments conducted for the investigation of spatial distribution of residential wood and coal burning do not result in unrealistic EC concentrations. The three sites in Northern Ireland
 15 (BEL, LISB, and STAB in Fig. 14) exhibit a stronger seasonal cycle in the measurements than the other sites (i.e. relatively greater increase for autumn and winter), indicating stronger traditions of residential solid fuel burning in this part of the UK.




Figure 14. Seasonal boxplots of daily-average concentrations of measured BC and modelled EC concentrations at the BC-network measurement sites, year 2012. Site name abbreviations are given in Fig. 12. Each box is the interquartile range - IQR, whiskers extend to $1.5 \times \text{IQR}$, and all daily values beyond the whiskers are plotted individually.

4 Conclusions

In this study, different assumptions for the spatial distribution and total emitted amount of PM emissions from solid fuel burning in the UK were tested with the EMEP4UK atmospheric chemical transport model. These model experiments were conducted to investigate the large model underestimations of SFOA concentrations compared with aerosol source apportionment measurements which arise when using the officially-reported PM emissions inventory (a NMB of -71% at the London North Kensington urban background site, for example). The two main scenarios considered were *Base4x*, and *combRedist*. For *Base4x*, officially reported $PM_{2.5}$ from the SNAP2 emission source sector (residential and other non-industrial combustion) were increased by a factor of 4. For the *combRedist* experiment, half of the emissions from SNAP2 were redistributed linearly by residential population density to extend emissions into smoke control areas. The emission total for the *combRedist* experiment was the same as that officially reported (i.e. equal to the *Base* scenario, and 4 times less than in the *Base4x* scenario). A third experiment, *Redist* - all emissions redistributed linearly to population density, is also presented as an indicator of the maximum concentrations an assumption like this could yield. This is not however completely realistic as the most densely populated areas (of large apartment buildings) are unlikely to have many individual fireplaces.

Comparison of model output with AMS-PMF measurements of SFOA concentrations at an urban background and roadside site in central London (a smoke control area), shows that *Base4x* yielded better daily and hourly correlations than the *combRedist*. Therefore, for certain air masses, the spatial distribution reported by the national emissions inventory appears reasonable. However, the *Base4x* overestimated SFOA concentrations at the rural sites, whereas the *combRedist* better captured mean measured concentrations across the range of site types and locations. The *combRedist* was intentionally simplistic (exactly 50% of the national total was spatially redistributed), so a better agreement might be, for example, *Base2x* + 30% redistributed to population density (or another combination of *Base* emissions and redistribution to include emissions for smoke control areas). The results also suggest that refinement of the prescribed temporal profiles for residential emissions may also be required as the measurements indicated higher levels of SFOA concentrations in the evening than in the morning, whereas the emissions profiles used here emitted relatively more during the morning. It is acknowledged that the experiments undertaken here investigated only potential discrepancies in the national atmospheric emissions inventory, and not other potential sources of model-measurement discrepancy such as shortcomings in model quantification of dry and wet deposition, both of which also depend on accurate model description of the meteorology.

Modelled concentrations of elemental carbon were compared with different measurements: EC-R, EC-T, and black carbon (BC). Substantial discrepancies were noted between the different measurements of this component of PM, so detailed comparison with the model was not presented. However, based on seasonal-average concentrations at the BC network sites over the UK, it was shown that the concentrations derived from the *combRedist* experiments improved the comparisons of modelled vs observed concentrations. Therefore, some redistribution of SNAP2 emissions into smoke control areas, as also suggested by the SFOA comparisons, appears justified. Overall, our results suggest that simulations of SFOA can be improved by adjusting the spatial distribution of the national emissions inventory.

5 Code availability

The EMEP model is OpenSource and can be downloaded from www.emep.int. The WRF model is OpenSource and can be downloaded from <https://www.mmm.ucar.edu/weather-research-and-forecasting-model>.

6 Data availability

- 5 Processed measurement data used in this study are available through the ClearfLo project archive at the British Atmospheric Data Centre (<http://badc.nerc.ac.uk/browse/badc/clearflo>). The model data (input, scripts, relevant output) are archived at the University of Edinburgh and are available on request.

Appendix A: Measured EC-R, EC-T, BC

A1 Overview

- 10 Because of its diverse origins and chemical processing, airborne carbonaceous particulate matter exists in a continuum of different forms – from pure graphitic-like elemental carbon (EC) at one end through to an array of highly chemically-functionalised organic compounds at the other (Gelencsér, 2004). This raises the issue of at what level of chemical oxidation/functionalisation should EC no longer be categorised as EC but as organic carbon (OC). In practice, EC, and an alternative measure for the ‘sooty carbon’ content of particulate matter – black carbon (BC) – are both measurement defined, rather than chemically defined (Pet-
15 zold et al., 2013; Lack et al., 2014). EC is defined by thermal methods (heating up a PM sample and burning off the more volatile organic components such that what is only burnt off at highest temperatures is called EC; Chow et al. (2007)), and BC is defined by optical methods (measuring how opaque the material is to transmission of visible or near infrared wavelengths to determine effective black carbon, or eBC; Bond and Bergstrom (2006)). So by definition EC and eBC are not measuring the same thing, and both methods require choices in the quantification process. For example, the EC method requires choice on
20 the temperature programme and pyrolysis correction approach used to assign a distinction between EC and OC (Chow et al., 2004; Cavalli et al., 2010), while the eBC method requires a ‘shadowing’ correction (Virkkula et al., 2007) and imposition of a mass-specific absorption coefficient (MAC) to convert optical absorbance to mass concentration (Andreae and Gelencsér, 2006; Quincey et al., 2009). These issues have been the subject of much discussion in the literature, some of which is cited above, with development of standard protocols and terminology. In general, however, measurements of EC and eBC on the
25 same samples are very highly correlated, and to a first approximation EC and eBC data values can both be used in comparison of model output of EC against measurements. For the remainder of this Appendix, eBC is referred to as BC.

A2 Measurements of EC-T and EC-R

The UK Particle Numbers and Concentration Network collects daily samples of PM₁₀ onto binder-free pure quartz filters using a Partisol 2025 sampler at three measurement stations: London Marylebone Road (kerbside), London North-Kensington

(urban background) and Harwell (rural background). These filters are analysed at the UK National Physics Laboratory (on a Sunset Laboratory Carbon Aerosol Analysis Lab instrument). The protocol used to quantify total carbon (TC) of the sample is a variation of the NIOSH protocol known as Quartz (Quincey et al., 2009). During the heating of the sample, some organic matter will be converted to elemental carbon by pyrolysis. This conversion is monitored by continuously measuring the transmittance (T) and reflectance (R) of the sample. The T or R signal is used to apportion TC into OC and EC by taking account of carbonaceous material that was OC, but become pyrolysed into EC during the heating process. However, the quantification of the pyrolysed OC differs whether T or R is used, adding uncertainty to the measurements.

Figure A1 illustrates the uncertainty of the EC split from TC using transmittance (EC-T) and EC split from TC using reflectance (EC-R). For Harwell and North Kensington, EC-R is higher than EC-T (on average over the full dataset) which agrees with the findings of Chow et al. (2004). However, it can be noted that for both Harwell and North Kensington, in the lower range ($< \sim 0.4 \mu\text{g m}^{-3}$ for Harwell, $< \sim 0.6 \mu\text{g m}^{-3}$ for North Kensington), EC-T is higher than EC-R, whereas above these values, EC-R is higher than EC-T.

Figures A2, A3, and A4 split the points in Fig. A1 into seasons for Harwell, North Kensington, and Marylebone Road, respectively. In spring and winter, EC-T is lower than EC-R (NMB from -7% to -35%), whereas during summer EC-T is higher than EC-R (NMB from $+19\%$ to $+31\%$). In autumn, average measured concentrations of EC-R and EC-T are similar (NMB from -2% to $+10\%$), but substantial scatter can be seen for some days as measured concentrations of EC-T and EC-R can differ from each other by more than a factor of 2. These consistent findings at all three sites suggest that the different methods for quantifying of EC with the same instrument are dependent on season.

A3 Measurements of BC

Measurements of BC use an Aethalometer (Magee AE22) and involve the collection of $\text{PM}_{2.5}$ onto a quartz tape, for which the absorption $\alpha [\text{m}^{-1}]$ is measured by single-pass attenuation of 880 nm light, corrected for scattering. The absorption is converted to BC concentration using a mass-specific absorption cross section (MAC). The measurements are reported with an hourly timestep. The assumption of a single value for the MAC can lead to uncertainties due to (Heal and Quincey, 2012):




Figure A1. Scatterplot of measured daily-average EC split from TC using transmittance (EC-T) and EC split from TC using reflectance (EC-R), year 2012. Data below the detection limit ($< 0.1 \mu\text{g m}^{-3}$) have been removed, leading to the lower number of data points for Harwell.




Figure A2. Scatterplot of measured daily-average EC split from TC using transmittance (EC-T) and EC split from TC using reflectance (EC-R) at the Harwell site split by seasons, year 2012.




Figure A3. Scatterplot of measured daily-average EC split from TC using reflectance (EC-R) against EC split from TC using transmittance (EC-T) at the North Kensington site split by seasons, year 2012.

- Atmospheric oxidation changing the MAC of the sample (making it appear less dark).
- Variations in the size distribution of the particles leading to variations in the MAC.
- Variations in the external and internal mixing with other particles in the sample leading to variations in the optical properties of the sample.




Figure A4. Scatterplot of measured daily-average EC split from TC using reflectance (EC-R) against EC split from TC using transmittance (EC-T) at the Marylebone Road site split by seasons, year 2012.




Figure A5. Time-series of daily-average measured EC-R and BC concentrations at the (a) Harwell, (b) North Kensington, and (c) Marylebone Road measurement sites, year 2012.

A4 Measured BC vs measured EC

Figure A5 shows daily-average time-series of measured EC-R and BC for the three measurement sites that have both sets of measurements. For Jan–March and Oct–Dec, measurements of BC and EC-R at the Harwell and North Kensington sites are a close match to each other, whereas from April to September, BC is consistently higher than EC-R. At Marylebone Road,

winter and early spring months have better agreement than the summer, but BC is overestimated compared to EC-R throughout the year.

A5 Appendix summary

There are several inherent and methodological uncertainties in quantifying the refractory part of carbonaceous aerosol as EC or BC. Overall, the mean absolute values of the different measurements relate to each other as follows: BC > EC-R > EC-T, but the magnitudes of the differences, and in some cases also the order, vary for seasons and for individual sites.

Author contributions. R.O. designed, executed and analysed the model experiments. R.O. and M.R.H. wrote the manuscript. L.X., L.R.W., S.C.H., and N.L.N. collected and processed the AMS measurements at the Detling Winter IOP site, D.E.Y., J.D.A, H.C, and D.C.G. at the London North Kensington site, and E.M., C.M, and A.D. at the London Marylebone Road site and the Harwell site. M.V manages the EMEP4UK project. S.R. processed the population density data. I.A.M. processed the Base case emissions inventory. All authors contributed to the final version of the manuscript.

Competing interests. The authors declare that they have no conflicts of interest.

Disclaimer. None.

Acknowledgements. The authors acknowledge the UK Department for Environment, Food and Rural Affairs (Defra) and the Devolved Administrations for funding aspects of the development of the EMEP4UK model (AQ0727), for partial support for the aerosol measurements, as well as access to the AURN data, which were obtained from uk-air.defra.gov.uk and are subject to Crown 2014 copyright, Defra, licensed under the Open Government Licence (OGL). Partial support for the EMEP4UK modelling from the European Commission FP7 ECLAIRE project is gratefully acknowledged. This work was supported in part by the UK Natural Environment Research Council (NERC) ClearLo project [grant ref. NE/H003169/1]. R.O. was supported by a PhD studentship (University of Edinburgh and NERC-CEH contract 587/NEC03805). D.E.Y. was supported by a NERC PhD studentship [ref. NE/I528142/1]. L.X., L.R.W., S.C.H., and N.L.N. acknowledge support from US Department of Energy (grant no. DE-SC000602).

NCAR command language (NCL) was used to produce the maps (NCAR, 2015), and R, `openair` and `ggplot2` for the analysis and all other plots (R Core Team, 2016; Carslaw and Ropkins, 2012; Wickham, 2009).

References

- Allan, J. D., Williams, P. I., Morgan, W. T., Martin, C. L., Flynn, M. J., Lee, J., Nemitz, E., Phillips, G. J., Gallagher, M. W., and Coe, H.: Contributions from transport, solid fuel burning and cooking to primary organic aerosols in two UK cities, *Atmos. Chem. Phys.*, 10, 647–668, doi:10.5194/acp-10-647-2010, 2010.
- 5 Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, doi:10.5194/acp-6-3131-2006, 2006.
- Bell, M. L., Davis, D. L., and Fletcher, T.: A retrospective assessment of mortality from the London smog episode of 1952: the role of influenza and pollution., *Environmental Health Perspectives*, 112, 6–8, <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1241789/>, 2004.
- Bergström, R.: Carbonaceous Aerosol in Europe Out of the Woods and into the Blue?, Ph.D. thesis, University of Gothenburgh, 2015.
- 10 Bohnenstengel, S. I., Belcher, S. E., Aiken, A., Allan, J. D., Allen, G., Bacak, A., Bannan, T. J., Barlow, J. F., Beddows, D. C. S., Bloss, W. J., Booth, A. M., Chemel, C., Coceal, O., Di Marco, C. F., Dubey, M. K., Faloon, K. H., Fleming, Z. L., Furger, M., Gietl, J. K., Graves, R. R., Green, D. C., Grimmond, C. S. B., Halios, C. H., Hamilton, J. F., Harrison, R. M., Heal, M. R., Heard, D. E., Helfter, C., Herndon, S. C., Holmes, R. E., Hopkins, J. R., Jones, A. M., Kelly, F. J., Kotthaus, S., Langford, B., Lee, J. D., Leigh, R. J., Lewis, A. C., Lidster, R. T., Lopez-Hilfiker, F. D., McQuaid, J. B., Mohr, C., Monks, P. S., Nemitz, E., Ng, N. L., Percival, C. J., Prévôt, A. S. H., Ricketts, H.
- 15 M. A., Sokhi, R., Stone, D., Thornton, J. A., Tremper, A. H., Valach, A. C., Visser, S., Whalley, L. K., Williams, L. R., Xu, L., Young, D. E., and Zotter, P.: Meteorology, Air Quality, and Health in London: The ClearLo Project, *Bulletin of the American Meteorological Society*, 96, 779–804, doi:10.1175/BAMS-D-12-00245.1, 2014.
- Bond, T. C. and Bergstrom, R. W.: Light Absorption by Carbonaceous Particles: An Investigative Review, *Aerosol Science and Technology*, 40, 27–67, doi:10.1080/02786820500421521, 2006.
- 20 Carslaw, D. C. and Ropkins, K.: openair — An R package for air quality data analysis, *Environmental Modelling & Software*, 27–28, 52–61, doi:10.1016/j.envsoft.2011.09.008, 2012.
- Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3, 79–89, doi:10.5194/amt-3-79-2010, 2010.
- CEIP: WebDab EMEP database: Emissions as used in EMEP models, http://www.ceip.at/ms/ceip_home1/ceip_home/webdab_emepdatabase/, emis. data downloaded on 15.07.2015, 2015.
- 25 Chow, J. C., Watson, J. G., Chen, L. A., Arnott, W. P., Moosmüller, H., and Fung, K.: Equivalence of Elemental Carbon by Thermal/Optical Reflectance and Transmittance with Different Temperature Protocols, *Environmental Science & Technology*, 38, 4414–4422, doi:10.1021/es034936u, 2004.
- Chow, J. C., Yu, J. Z., Watson, J. G., Ho, S. S. H., Bohannon, T. L., Hays, M. D., and Fung, K. K.: The application of thermal methods for determining chemical composition of carbonaceous aerosols: A review, *Journal of Environmental Science and Health, Part A*, 42, 1521–1541, doi:10.1080/10934520701513365, 2007.
- 30 Crilley, L. R., Bloss, W. J., Yin, J., Beddows, D. C. S., Harrison, R. M., Allan, J. D., Young, D. E., Flynn, M., Williams, P., Zotter, P., Prevot, A. S. H., Heal, M. R., Barlow, J. F., Halios, C. H., Lee, J. D., Szidat, S., and Mohr, C.: Sources and contributions of wood smoke during winter in London: assessing local and regional influences, *Atmos. Chem. Phys.*, 15, 3149–3171, doi:10.5194/acp-15-3149-2015, 2015.
- 35 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-Deployable, High-Resolution, Time-of-Flight Aerosol Mass Spectrometer, *Analytical Chemistry*, 78, 8281–8289, doi:10.1021/ac061249n, 2006.

- Denier van der Gon, H. A. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D., and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe – revised estimates and an evaluation, *Atmos. Chem. Phys.*, 15, 6503–6519, doi:10.5194/acp-15-6503-2015, 2015.
- Detournay, A., Nemitz, E., Di Marco, C. F., and et al.: A 1-year study of submicron aerosol composition at a kerbside location in London, In Preparation, 2017.
- Di Marco, C. F., Nemitz, E., and et al.: Chemical characterisation of the aerosol at a background site in southern England with emphasis on the organic fraction and nitrogen compounds, In Preparation, 2017.
- Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled Partitioning, Dilution, and Chemical Aging of Semivolatile Organics, *Environmental Science & Technology*, 40, 2635–2643, doi:10.1021/es052297c, PMID: 16683603, 2006.
- 10 EEA: EMEP/EEA air pollutant emission inventory guidebook - 2013, <http://www.eea.europa.eu/publications/emep-eea-guidebook-2013>, 2013.
- Fuller, G. W., Sciare, J., Lutz, M., Moukhtar, S., and Wagener, S.: New Directions: Time to tackle urban wood burning?, *Atmospheric Environment*, 68, 295–296, doi:10.1016/j.atmosenv.2012.11.045, 2013.
- Fuller, G. W., Tremper, A. H., Baker, T. D., Yttri, K. E., and Butterfield, D.: Contribution of wood burning to PM10 in London, *Atmospheric Environment*, 87, 87–94, doi:10.1016/j.atmosenv.2013.12.037, 2014.
- 15 Gelencsér, A.: Carbonaceous Aerosol, Springer Netherlands, doi:10.1007/978-1-4020-2887-8, 2004.
- Harrison, R. M. and Yin, J.: Sources and processes affecting carbonaceous aerosol in central England, *Atmospheric Environment*, 42, 1413–1423, doi:10.1016/j.atmosenv.2007.11.004, 2008.
- Heal, M. R. and Quincey, P.: The relationship between black carbon concentration and black smoke: A more general approach, *Atmospheric Environment*, 54, 538–544, doi:10.1016/j.atmosenv.2012.02.067, 2012.
- 20 Jayne, J. T., Leard, D. C., Zhang, X., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development of an Aerosol Mass Spectrometer for Size and Composition Analysis of Submicron Particles, *Aerosol Science and Technology*, 33, 49–70, doi:10.1080/027868200410840, 2000.
- Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and Denier van der Gon, H. A. C.: TNO-MACC_II emission inventory; a multi-year (2003–2009) consistent high-resolution European emission inventory for air quality modelling, *Atmos. Chem. Phys.*, 14, 10963–10976, doi:10.5194/acp-14-10963-2014, 2014.
- 25 Lack, D. A., Moosmüller, H., McMeeking, G. R., Chakrabarty, R. K., and Baumgardner, D.: Characterizing elemental, equivalent black, and refractory black carbon aerosol particles: a review of techniques, their limitations and uncertainties, *Analytical and Bioanalytical Chemistry*, 406, 99–122, doi:10.1007/s00216-013-7402-3, 2014.
- 30 Lefebvre, W., Fierens, F., Vanpoucke, C., Renders, N., Jaspers, K., Vercauteren, J., Deutsch, F., and Janssen, S.: The Effect of Wood Burning on Particulate Matter Concentrations in Flanders, Belgium, in: *Air Pollution Modeling and its Application XXIV*, edited by Steyn, D. G. and Chaumerliac, N., Springer Proceedings in Complexity, pp. 459–464, Springer International Publishing, doi:10.1007/978-3-319-24478-5_73, 2016.
- Legates, D. R. and McCabe, G. J.: A refined index of model performance: a rejoinder, *International Journal of Climatology*, 33, 1053–1056, doi:10.1002/joc.3487, 2013.
- 35 May, A. A., Levin, E. J. T., Hennigan, C. J., Riipinen, I., Lee, T., Collett, J. L., Jimenez, J. L., Kreidenweis, S. M., and Robinson, A. L.: Gas-particle partitioning of primary organic aerosol emissions: 3. Biomass burning, *Journal of Geophysical Research: Atmospheres*, 118, 11,327–11,338, doi:10.1002/jgrd.50828, 2013JD020286, 2013.

- NAEI: UK Emission Mapping Methodology 2009, http://naei.defra.gov.uk/reports/reports?report_id=785, last accessed: 19-07-2017, 2013.
- NCAR: The NCAR Command Language (Version 6.3.0) [Software], Boulder, Colorado: UCAR/NCAR/CISL/TDD, doi:10.5065/D6WD3XH5, 2015.
- NCEP: NCEP FNL Operational Model Global Tropospheric Analyses, continuing from July 1999, Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, Boulder CO, doi:10.5065/D6M043C6, 2000.
- 5 Ots, R., Young, D. E., Vieno, M., Xu, L., Dunmore, R. E., Allan, J. D., Coe, H., Williams, L. R., Herndon, S. C., Ng, N. L., Hamilton, J. F., Bergström, R., Di Marco, C., Nemitz, E., Mackenzie, I. A., Kuenen, J. J. P., Green, D. C., Reis, S., and Heal, M. R.: Simulating secondary organic aerosol from missing diesel-related intermediate-volatility organic compound emissions during the Clean Air for London (ClearLo) campaign, *Atmos. Chem. Phys.*, 16, 6453–6473, doi:10.5194/acp-16-6453-2016, 2016a.
- 10 Ots, R., Vieno, M., Allan, J. D., Reis, S., Nemitz, E., Young, D. E., Coe, H., Di Marco, C., Detournay, A., Mackenzie, I. A., Green, D. C., and Heal, M. R.: Model simulations of cooking organic aerosol (COA) over the UK using estimates of emissions based on measurements at two sites in London, *Atmospheric Chemistry and Physics*, 16, 13 773–13 789, doi:10.5194/acp-16-13773-2016, 2016b.
- Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S.-M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, X.-Y.: Recommendations for reporting "black carbon" measurements, *Atmos. Chem. Phys.*, 13, 8365–8379, doi:10.5194/acp-13-8365-2013, 2013.
- 15 Quincey, P., Butterfield, D., Green, D., Coyle, M., and Cape, J. N.: An evaluation of measurement methods for organic, elemental and black carbon in ambient air monitoring sites, *Atmospheric Environment*, 43, 5085–5091, doi:10.1016/j.atmosenv.2009.06.041, 2009.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org/>, 2016.
- 20 Reis, S., Steinle, S., Carnell, E. J., Leaver, D., Vieno, M., Beck, R., and Dragosits, U.: UK gridded population based on Census 2011 and Land Cover Map 2007, NERC Environmental Information Data Centre, doi:10.5285/61f10c74-8c2c-4637-a274-5fa9b2e5ce44, 2016.
- Schulz, M., Gauss, M., Benedictow, A., Jonson, J. E., Tsyro, S., Nyíri, A., Simpson, D., Steensen, B. M., Klein, H., Valdebenito, A., Wind, P., Kirkevåg, A., Griesfeller, J., Bartnicki, J., Olivie, D., Grini, A., Iversen, T., Seland, ., Semeena, S. V., Fagerli, H., Aas, W., Hjellbrekke, A., Mareckova, K., Wankmuller, R., Schneider, P., Solberg, S., Svendby, T., Liu, L., Posch, M., Vieno, M., Reis, S., Kryza, M., Werner, M., and Walaszek, K.: Transboundary Acidification, Eutrophication and Ground Level Ozone in Europe in 2011, Tech. rep., Norwegian Meteorological Institute, 2013.
- 25 Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard, C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, S. V., Tsyro, S., Tuovinen, J. P., Valdebenito, A., and Wind, P.: The EMEP MSC-W chemical transport model - technical description, *Atmos. Chem. Phys.*, 12, 7825–7865, doi:10.5194/acp-12-7825-2012, 2012.
- 30 Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009.
- Vieno, M., Dore, A. J., Wind, P., Marco, C. D., Nemitz, E., Phillips, G., Tarrasón, L., and Sutton, M. A.: Application of the EMEP Unified Model to the UK with a Horizontal Resolution of 5 x 5 km², in: *Atmospheric Ammonia*, edited by Sutton, M. A., Reis, S., and Baker, S. M. H., pp. 367–372, Springer Netherlands, doi:10.1007/978-1-4020-9121-6_21, 2009.
- 35 Vieno, M., Dore, A. J., Stevenson, D. S., Doherty, R., Heal, M. R., Reis, S., Hallsworth, S., Tarrason, L., Wind, P., Fowler, D., Simpson, D., and Sutton, M. A.: Modelling surface ozone during the 2003 heat-wave in the UK, *Atmos. Chem. and Phys.*, 10, 7963–7978, doi:10.5194/acp-10-7963-2010, 2010.

- Vieno, M., Heal, M. R., Williams, M. L., Carnell, E. J., Nemitz, E., Stedman, J. R., and Reis, S.: The sensitivities of emissions reductions for the mitigation of UK PM_{2.5}, *Atmos. Chem. Phys.*, 16, 265–276, doi:10.5194/acp-16-265-2016, 2016.
- Virkkula, A., Mäkelä, T., Hillamo, R., Yli-Tuomi, T., Hirsikko, A., Hämeri, K., and Koponen, I. K.: A Simple Procedure for Correcting Loading Effects of Aethalometer Data, *Journal of the Air & Waste Management Association*, 57, 1214–1222, doi:10.3155/1047-3289.57.10.1214, 2007.
- 5 Waters, L.: Summary results of the domestic wood use survey, Tech. rep., Renewable Energy Statistics, <https://www.gov.uk/government/publications/energy-trends-march-2016-special-feature-article-summary-results-of-the-domestic-wood-use-survey>, last accessed: 19-07-2017, 2016.
- WHO: Residential heating with wood and coal: health impacts and policy options in Europe and North America, Tech. rep., World Health Organisation, Copenhagen, Denmark, <http://www.euro.who.int/en/publications/abstracts/residential-heating-with-wood-and-coal-health-impacts-and-policy-options-in-europe-and-north-america>, 2015.
- 10 Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer-Verlag New York, <http://ggplot2.org>, 2009.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, doi:10.5194/gmd-4-625-2011, 2011.
- 15 Xu, L., Williams, L. R., Young, D. E., Allan, J. D., Coe, H., Massoli, P., Fortner, E., Chhabra, P., Herndon, S., Brooks, W. A., Jayne, J. T., Worsnop, D. R., Aiken, A. C., Liu, S., Gorkowski, K., Dubey, M. K., Fleming, Z. L., Visser, S., Prévôt, A. S. H., and Ng, N. L.: Wintertime aerosol chemical composition, volatility, and spatial variability in the greater London area, *Atmos. Chem. Phys.*, 16, 1139–1160, doi:10.5194/acp-16-1139-2016, <http://www.atmos-chem-phys.net/16/1139/2016/>, 2016.
- 20 Young, D. E., Allan, J. D., Williams, P. I., Green, D. C., Flynn, M. J., Harrison, R. M., Yin, J., Gallagher, M. W., and Coe, H.: Investigating the annual behaviour of submicron secondary inorganic and organic aerosols in London, *Atmos. Chem. Phys.*, 15, 6351–6366, doi:10.5194/acp-15-6351-2015, 2015a.
- Young, D. E., Allan, J. D., Williams, P. I., Green, D. C., Harrison, R. M., Yin, J., Flynn, M. J., Gallagher, M. W., and Coe, H.: Investigating a two-component model of solid fuel organic aerosol in London: processes, PM₁ contributions, and seasonality, *Atmos. Chem. Phys.*, 15, 2429–2443, doi:10.5194/acp-15-2429-2015, 2015b.
- 25