

# First long-term and near real-time measurement of trace elements in China's urban atmosphere: temporal variability, source apportionment, and precipitation effect

Yunhua Chang<sup>1, 2</sup>, Kan Huang<sup>3</sup>, Mingjie Xie<sup>4</sup>, Congrui Deng<sup>3</sup>, Zhong Zou<sup>5</sup>, Shoudong Liu<sup>1, 2</sup>, and Yanlin Zhang<sup>1, 2 \*</sup>

<sup>1</sup>Yale-NUIST Center on Atmospheric Environment, International Joint Laboratory on Climate and Environment Change (ILCEC), Nanjing University of Information Science & Technology, Nanjing 210044, China

<sup>2</sup>Key Laboratory of Meteorological Disaster, Ministry of Education (KLME)/ Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters (CIC-FEMD), Nanjing University of Information Science & Technology, Nanjing 210044, China

<sup>3</sup>Center for Atmospheric Chemistry Study, Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention (LAP<sup>3</sup>), Department of Environmental Science and Engineering, Fudan University, Shanghai 200433, China

<sup>4</sup>School of Environmental Science and Engineering, Nanjing University of Information Science & Technology, Nanjing 210044, China

<sup>5</sup>Pudong New Area Environmental Monitoring Station, Shanghai 200135, China

*Correspondence to:* Yanlin Zhang (dryanlinzhang@outlook.com or zhangyanlin@nuist.edu.cn)

**Abstract:** Atmospheric trace elements, especially metal species, are an emerging environmental and health concern with insufficient understanding of their levels and sources in Shanghai, the most important industrial megacity in China. Here we continuously performed a one-year (from March 2016 to February 2017) and hourly-resolved measurement of eighteen elements in fine particles (PM<sub>2.5</sub>) at Shanghai urban center with a Xact multi-metals monitor and several collocated instruments. Mass concentrations (mean $\pm$ 1 $\sigma$ ; ng m<sup>-3</sup>) determined by Xact ranged from detection limits (nominally 0.1 to 20 ng m<sup>-3</sup>) to 15  $\mu$ g m<sup>-3</sup>. Elements related oxidized species comprised

an appreciable fraction of PM<sub>2.5</sub> during all seasons, accounting for 8.3% on average. As  
30 a comparison, atmospheric elements concentration level in Shanghai was comparable  
with that in other industrialized cities in East Asia but one or two orders of magnitude  
higher than at sites in North America and Europe. Positive matrix factorization (PMF)  
was applied to identify and apportion the sources of elements in PM<sub>2.5</sub> mass. Five  
different factors were resolved (notable elements and relative contribution in brackets):  
35 traffic-related (Ca, Fe, Ba, Si; 46%), shipping (V, Ni; 6%), nonferrous metal melting  
(Ag, Cd, Au; 15%), coal combustion (As, Se, Hg, Pb; 18%), and ferrous metal smelting  
(Cr, Mn, Zn; 15%). Contributed by exhaust and non-exhaust vehicle emissions, traffic-  
related factor shows strong bimodal diurnal profile with average concentration over two  
times higher during rush hour than during nighttime. Shipping factor was firmly  
40 identified because V and Ni, two recognized tracers of shipping emissions, almost  
exclusively transported from East China Sea and their ratio (around 0.31) fallen within  
the variation range of V/Ni in particles emitted from heavy oil combustion. Interestingly,  
nearly half of K were derived from coal combustion with high mineral affinity  
(elements associated with aluminosilicates, carbonates and other minerals in coal ash).  
45 The contributions of (non)ferrous metal smelting to trace elements are consistent with  
a newly-developed emission inventory. Although the precipitation scavenging effect on  
the mass concentration of trace elements varied among different species and sources,  
precipitation could effectively lower traffic- and coal combustion-related trace elements.  
Therefore, water spray to simulate natural types of precipitation could be one of the  
50 abatement strategies to facilitate the reduction of ambient PM<sub>2.5</sub> trace elements in urban  
atmosphere. Collectively, our findings in this study provide baseline levels and sources  
of trace elements with high detail, which are needed for developing effective control  
strategies to reduce the high risk of acute exposure to atmospheric trace elements in  
China's megacities.

## 55 1 Introduction

It is well known that personal exposure to atmospheric aerosols have detrimental

consequences and aggravating effects on human health such as respiratory, cardiovascular, and allergic disorders (Pope III et al., 2002; Pope III et al., 2009; Shah et al., 2013; West et al., 2016; Burnett et al., 2014). Among the chemical components relevant for aerosol health effects, airborne heavy metals (a very imprecise term without authoritative definition (John, 2002), loosely refers to elements with atomic density greater than  $4.5 \text{ g cm}^{-3}$  (Streit, 1991)) are of particular concern as they typically feature with unique properties of bioavailability and bioaccumulation (Morman and Plumlee, 2013; Tchounwou et al., 2012; Fergusson, 1990; Kastury et al., 2017), representing 7 of the 30 hazardous air pollutants identified by the US Environmental Protection Agency (EPA) in terms of posing the greatest potential health threat in urban areas (see [www.epa.gov/urban-air-toxics/urban-air-toxic-pollutants](http://www.epa.gov/urban-air-toxics/urban-air-toxic-pollutants)). Depending on aerosol composition, extent and time of exposure, previous studies have confirmed that most elements components of fine particles ( $\text{PM}_{2.5}$ ; particulate matter with aerodynamic diameter equal to or less than  $2.5 \mu\text{m}$ ) exerted a multitude of significant diseases from pulmonary inflammation, to increased heart rate variability, to decreased immune response (Fergusson, 1990; Morman and Plumlee, 2013; Leung et al., 2008; Hu et al., 2012; Pardo et al., 2015; Kim et al., 2016).

Guidelines for atmospheric concentration limits of many trace elements are provided by the World Health Organization (WHO) (WHO, 2005). In urban atmospheres, ambient trace elements typically represent a small fraction of  $\text{PM}_{2.5}$  on a mass basis, while elements species like Cd, As, Co, Cr, Ni, Pb and Se are considered as human carcinogens even in trace amounts (Iyengar and Woittiez, 1988; Wang et al., 2006; Olujimi et al., 2015). It has been shown that Cu, Cr, Fe and V have several oxidation states that can participate in many atmospheric redox reactions (Litter, 1999; Brandt and van Eldik, 1995; Seigneur and Constantinou, 1995; Rubasinghege et al., 2010a), which can catalyze the generation of reactive oxygenated species (ROS) that have been associated with direct molecular damage and with the induction of biochemical synthesis pathways (Charrier and Anastasio, 2012; Strak et al., 2012; Rubasinghege et al., 2010b; Saffari et al., 2014; Verma et al., 2010; Jomova and Valko, 2011).

Additionally, lighter elements such as Si, Al and Ca are the most abundant crustal elements next to oxygen, which can typically constitute up to 50% of elemental species in remote continental aerosols (Usher et al., 2003; Ridley et al., 2016). These species are usually associated with the impacts of aerosols on respiratory diseases and climate (Usher et al., 2003; Tang et al., 2017).

Health effects of airborne elements species are not only seen from chronic exposure, but also from short-term acute concentration spikes in ambient air (Kloog et al., 2013; Strickland et al., 2016; Huang et al., 2012). In addition, atmospheric emissions, transport, and exposure of trace elements to human receptors may depend upon rapidly evolving meteorological conditions and facility operations (Tchounwou et al., 2012; Holden et al., 2016). Typical ambient trace elements sampling devices collect 12 to 24-hr integrated average samples, which are then sent off to be lab analyzed in a time-consuming and labor-intensive way. As a consequence, daily integrated samples inevitably ignore environmental shifts with rapid temporality, and thereby hinder the efforts to obtain accurate source apportionment results such as short-term elements pollution spikes related to local emission sources. In fact, during a short-term trace elements exposure event, 12 or 24-hr averaged sample concentrations for elements species like Pb and As may be one order of magnitude lower than the 4-hr or 15-min average concentration from the same day (Cooper et al., 2010). Current source apportionment studies are mainly performed by statistical multivariate analysis such as receptor models (e.g., Positive Matrix Factorization, PMF), which could greatly benefit from high inter-sample variability in the source contributions through increasing the sampling time resolution. In this regard, continuous monitoring of ambient elements species on a real-time scale is essential for studies on trace elements sources and their health impacts.

Currently, there are only a few devices available for the field sampling of ambient aerosols with sub-hourly or hourly resolution, i.e., the Streaker sampler, the DRUM (Davis Rotating-drum Unit for Monitoring) sampler, and the SEAS (Semi-continuous

Elements in Aerosol Sampler) (Visser et al., 2015b; Visser et al., 2015a; Bukowiecki et  
115 al., 2005; Chen et al., 2016). Mass loadings of trace elements collected by these  
samplers can be analyzed with highly sensitive accelerator-based analytical techniques,  
in particular particle-induced X-ray emission (PIXE) or synchrotron radiation X-ray  
fluorescence (SR-XRF) (Richard et al., 2010; Bukowiecki et al., 2005; Maenhaut, 2015;  
Traversi et al., 2014). More recently, Aerosol Time-Of-Flight Mass Spectrometry  
120 (ATOFMS) (Murphy et al., 1998; Gross et al., 2000; DeCarlo et al., 2006), National  
Institute for Standards and Technology (NIST)-traceable reference aerosol generating  
method (QAG) (Yanca et al., 2006), distance-based detection in a multi-layered device  
(Cate et al., 2015), environmental magnetic properties coupled with support vector  
machine (Li et al., 2017), and Xact<sup>TM</sup> 625 automated multi-metals analyzer (Fang et al.,  
125 2015; Jeong et al., 2016; Phillips-Smith et al., 2017; Cooper et al., 2010) have been  
developed for more precise, accurate, and frequent measurement of ambient elements  
species. The Xact method is based on nondestructive XRF analysis of aerosol deposits  
on reactive filter tapes, which has been validated by US Environmental Technology  
Verification testing and several other field campaigns (Fang et al., 2015; Phillips-Smith  
130 et al., 2017; Jeong et al., 2016; Yanca et al., 2006; Cooper et al., 2010; Park et al., 2014;  
Furger et al., 2017).

Located at the heart of the Yangtze River Delta (YRD), Shanghai is home to nearly 25  
million people as of 2015, making it the largest megacity in China (Chang et al., 2016).  
Shanghai city is one of the main industrial centers of China, playing a vital role in the  
135 nation's heavy industries, including but not limited to, steel making, petrochemical  
engineering, thermal power generation, auto manufacture, aircraft production, and  
modern shipbuilding (Normile, 2008; Chang et al., 2016; Huang et al., 2011). Shanghai  
is China's most important gateway for foreign trade, which has the world's busiest port,  
handling over 37 million standard containers in 2016 (see  
140 [www.simic.net.cn/news\\_show.php?lan=en&id=192101](http://www.simic.net.cn/news_show.php?lan=en&id=192101)). As a consequence, Shanghai  
is potentially subject to substantial quantities of trace elements emissions (Duan and  
Tan, 2013; Tian et al., 2015). Ambient concentrations of trace elements, especially Pb

and Hg, in the Shanghai atmosphere have been sporadically reported during the past two decades (Shu et al., 2001; Lu et al., 2008; Wang et al., 2013a; Zheng et al., 2004; Huang et al., 2013; Wang et al., 2016). Of current interest are V and Ni, which are often indicative of heavy oil combustion from ocean-going vessels (Fan et al., 2016; Liu et al., 2017). However, previous work rarely presented a full spectrum of elements species in ambient aerosols. Furthermore, recent attribution of hospital emergency-room visits in China to PM<sub>2.5</sub> constituents failed to take short-term variations of trace elements into account (Qiao et al., 2014), which could inevitably underestimate the toxicity of aerosols and potentially misestimate the largest influence of aerosol components on human health effects (Honda et al., 2017).

In this study, the first of its kind, we conducted a long-term and near real-time measurement of atmospheric trace elements in PM<sub>2.5</sub> with a Xact multi-metals analyzer in Shanghai, China, from March 2016 to February 2017. The primary target of the present study is to elucidate the levels and sources of atmospheric trace elements in a complex urban environment, which can be used to support future health studies. Meanwhile, the potential effect of precipitation scavenging on the mass concentration of trace elements will be investigated to examine if water spray could be proposed as an effective approach to curb severe trace elements pollution in China's urban atmosphere.


## **2 Methods**

### **2.1 Field measurements**

#### **2.1.1 Site description**

Figure 1a shows the map of eastern China with provincial borders and land cover, in which Shanghai city (provincial level) sits in the middle portion of China's eastern coast and its metropolitan area (indicated as the densely-populated area in Fig. 1b) concentrated on the south edge of the mouth of the Yangtze River. The municipality borders the provinces of Jiangsu and Zhejiang to the north, south and west, and is bounded to the east by the East China Sea (Fig. 1a). Shanghai has a humid subtropical

climate and experiences four distinct seasons. Winters are chilly and damp, with northwesterly winds from Siberia which can cause nighttime temperatures to drop below freezing. Air pollution in Shanghai is low compared to other cities in northern China, such as Beijing, but still substantial by world standards, especially in winter (Han et al., 2015; Chang et al., 2017).



**Figure 1.** Land use map indicating the location of Shanghai (a; black box), as well as the population density (b) and the major point sources (c) around the sampling site (PEMC). The emissions of V (d) and Ni (e) from shipping in the YRD and the East China Sea within 400 km of the coastline were estimated based on an automatic identification system model (adopted from Fan et al. (2016)).

Field measurements were performed at the rooftop (~18 m above ground level) of Pudong Environmental Monitoring Center (PEMC; 121.5447°E, 31.2331°N; ~7 m above sea level) in Pudong New Area of southwestern Shanghai, a region with dense population (Fig. 1b). Pudong New Area is described as the "showpiece" of modern China due to its height-obsessed skyline and export-oriented economy. For PEMC, there were no metal-related sources (except for road traffic) or high-rise buildings nearby to obstruct observations, so the air mass could flow smoothly. More broadly, as

indicted in Fig. 1c, PEMC is surrounded by a multitude of emissions sources such as coal-fired power plants (CFPP) in all directions and iron and steel smelting in the northwest. Furthermore, a high level of ship exhaust emissions in 2010 such as V (Fig. 1d) and Ni (Fig. 1e) in the YRD and the East China Sea within 400 km of China's coastline was recently quantified based on an automatic identification system model (Fan et al., 2016). Therefore, PEMC can be regarded as an ideal urban receptor site of diverse emission sources. More information regarding the sampling site has been given elsewhere (Chang et al., 2017; Chang et al., 2016).

### 2.1.2 Hourly elemental species measurements

From March 1<sup>st</sup> 2016 to February 28<sup>th</sup> 2017, hourly ambient mass concentrations of eighteen elements (Si, Fe, K, Ca, Zn, Mn, Pb, Ba, V, Cu, Cd, As, Ni, Cr, Ag, Se, Hg, and Au) in PM<sub>2.5</sub> were determined by a Xact multi-metals monitor (Model Xact™ 625, Cooper Environmental Services LLT, OR, USA) (Phillips-Smith et al., 2017; Jeong et al., 2016; Fang et al., 2015; Yanca et al., 2006). Specifically, the Xact sampled the air on a reel-to-reel Teflon filter tape through a PM<sub>2.5</sub> cyclone inlet (Model VSCC-A, BGI Inc., MA, USA) at a flow rate of 16.7 L min<sup>-1</sup>. The resulting PM<sub>2.5</sub> deposit on the tape was automatically advanced into the analysis area for nondestructive energy-dispersive X-ray fluorescence analysis to determine the mass of selected elemental species as the next sampling was being initiated on a fresh tape spot. Sampling and analysis were performed continuously and simultaneously, except during advancement of the tape (~20 sec) and during daily automated quality assurance checks. For every event of sample analysis, the Xact included a measurement of pure Pd as an internal standard to automatically adjust the detector energy gain. The XRF response was calibrated using thin film standards for each elements of interest. These standards were provided by the manufacturer of Xact, produced by depositing vapor phase elements on blank Nuclepore (Micromatter Co., Arlington, WA, USA). The Nuclepore filter of known area was weighed before and after the vapor deposition process to determine the concentration (μg cm<sup>-2</sup>) of each element. In this study, excellent agreement between the



measured and standard masses for each element was observed, indicating a deviation of < 5%. The 1-hr time resolution minimum detection limits (in ng m<sup>-3</sup>) were: Si (17.80), K (1.17), Ca (0.30), V (0.12), Cr (0.12), Mn (0.14), Fe (0.17), Ni (0.10), Cu (0.27), Zn (0.23), As (0.11), Se (0.14), Ag (1.90), Cd (2.50), Au (0.23), Ba (0.39), Hg (0.12), and Pb (0.13).

As a reference method to validate the Xact on-line measurements, daily PM<sub>2.5</sub> samples were also collected at the PEMC site using a four-channel aerosol sampler (Tianhong, Wuhan, China) on 47 mm cellulose acetate and glass fiber filters at a flow rate of 16.7 L min<sup>-1</sup>. The sampler was operated once a week with a 24-hr sampling time (starting from 10:00 am). In total 48 filter samples (26 cellulose acetate filter samples and 22 glass fiber filter samples) were collected, in which 8 paired samples were simultaneously collected by cellulose acetate and glass fiber filters. In the laboratory, elemental analysis procedures strictly followed the latest national standard method “Ambient air and stationary source emission-Determination of metals in ambient particulate matter-Inductively coupled plasma/mass spectrometer (ICP-MS)” (HJ 657-2013) issued by the Chinese Ministry of Environmental Protection. A total of 24 elements (Al, Fe, Mn, Mg, Mo, Ti, Sc, Na, Ba, Sr, Sb, Ca, Co, Ni, Cu, Ge, Pb, P, K, Zn, Cd, V, S, and As) were measured using Inductively coupled plasma-mass spectrometer (ICP-MS; Agilent, CA, USA). The results of the 8 paired samples were first compared. Significant correlations were observed for K, Cr, Mn, Fe, Ni, Cu, As, Cd, Ba, Zn, and Pb (Table S1), and these species were used to validate the performance of Xact. In Table S1, the slope values of Cr (1.9) and Ba (2.6) were higher than those of other species, this can be explained by higher background values of Cr and Ba collected by the cellulose acetate filters.

### 2.1.3 Auxiliary measurements, quality assurance and quality control

Meteorological data, including ambient temperature (*T*), relative humidity (RH), wind direction (WD) and wind speed (WS), were provided by Shanghai Meteorological Bureau at Century Park station (located approximately 2 km away from PEMC). The

routine procedures, including the daily zero/standard calibration, span and range check,  
 245 station environmental control, and staff certification, followed the Technical Guideline  
 of Automatic Stations of Ambient Air Quality in Shanghai based on the national  
 specification HJ/T193–2005. This was modified from the technical guidance  
 established by the USEPA. QA/QC for the Xact measurements was implemented  
 throughout the campaign. The internal Pd, Cr, Pb, and Cd upscale values were recorded  
 250 after the instrument's daily programmed test, and the PM<sub>10</sub> and PM<sub>2.5</sub> cyclones were  
 cleaned weekly.

## 2.2 Data analysis

### 2.2.1 Positive Matrix Factorization (PMF) analysis for source apportionment

The Positive Matrix Factorization or PMF is an effective source apportionment method  
 255 to identify and quantify possible emission sources of measurements using the bilinear  
 factor model (Paatero and Tapper, 1994)

$$x_{ij} = \sum_{k=1}^p g_{ik} f_{kj} + e_{ij}, \quad (1)$$

where  $x_{ij}$  is the  $j^{\text{th}}$  species concentration measured in the  $i^{\text{th}}$  sample,  $g_{ik}$  is the contribution  
 of the  $k^{\text{th}}$  source to the  $i^{\text{th}}$  sample (factor time series) and  $f_{kj}$  is the concentration of the  
 260  $j^{\text{th}}$  species in the  $k^{\text{th}}$  source (factor profiles). The part of the data remaining unexplained  
 by the model is represented by the residual matrix  $e_{ij}$ . The entries of  $g_{ik}$  and  $f_{kj}$  (required  
 to be non-negative) are fit using a least-squares algorithm that iteratively minimizes the  
 objective function  $Q$ :

$$Q = \sum_{i=1}^n \sum_{j=1}^m \left( \frac{e_{ij}}{\sigma_{ij}} \right)^2. \quad (2)$$

265 where  $\sigma_{ij}$  are the measurement uncertainties.

In this work, the US Environmental Protection Agency (EPA) PMF version 5.0 was

applied to attribute PM<sub>2.5</sub> trace elements to specific factors/sources. One-year hourly-resolved measurements ( $n = 8784$ ) of eighteen elements in PM<sub>2.5</sub> fraction were obtained and included for PMF analysis. The measurements ( $n = 1265$ ) with all elements missed were excluded. An estimated fractional uncertainty of 10% was used to derive the uncertainty data set (Kim et al., 2005; Kim and Hopke, 2007), which did not impact the interpretability of the PMF results. The missing values of individual elements were replaced by their geometric mean of the remaining observations, and their accompanying uncertainties were set to four times the geometric mean. The measurements below detection limit (BDL) were set to half the detection limit, with uncertainties set at five-sixths the detection limit (Polissar et al. 1998). The EPA PMF 5.0 has three uncertainty estimation methods, including bootstrapping (BS), displacement (DISP), and bootstrapping enhanced with DISP (BS-DISP) (Norris et al., 2014; Brown et al., 2015; Paatero et al., 2014; Wang et al., 2017). BS-DISP analysis is time consuming due to the huge data set ( $7519 \times 18$ ), and only BS and DISP analysis were conducted individually. Details of the uncertainty analysis are described in the supporting information (Text S1). In this study, PMF solutions using 3-10 factors were considered, and the final factor number is determined based on the interpretability as well the uncertainty analysis with BS and DISP methods.

### **2.2.2 Conditional probability function and bivariate polar plot for tracing source regions**

The determination of the geographical origins of trace elements in Shanghai requires the use of diagnostic tools such as the conditional probability function (CPF) and bivariate polar plot (BPP), which are very useful in terms of quickly gaining an idea of source impacts from various wind directions and have already been successfully applied to various atmospheric pollutants and pollution sources (Chang et al., 2017; Carslaw and Ropkins, 2012). In this study, the CPF and BPP were performed on the one-year data set for the major trace elements with similar source. The two methods have been implemented in the R “openair” package and are freely available at [www.openair-](http://www.openair-)

295 project.org (Carslaw and Ropkins, 2012).


The CPF is defined as  $CPF = m_{\theta}/n_{\theta}$ , where  $m_{\theta}$  is the number of samples in the wind sector  $\theta$  with mass concentrations greater than a predetermined threshold criterion, and  $n_{\theta}$  is the total number of samples in the same wind sector. CPF analysis is capable of showing which wind directions are dominated by high concentrations and with which probability. In this study, the 90<sup>th</sup> percentile of a given element species was set as threshold, and 24 wind sectors were used ( $\Delta\theta = 15^{\circ}$ ). Calm wind ( $< 1 \text{ m s}^{-1}$ ) periods were excluded from this analysis due to the isotropic behavior of the wind vane under calm winds.

The BPP demonstrates how the concentration of a targeted species varies synergistically with wind direction and wind speed in polar coordinates, which thus is essentially a non-parametric wind regression model to alternatively display pollution roses but include some additional enhancements. These enhancements include: plots are shown as a continuous surface and surfaces are calculated through modelling using smoothing techniques. These plots are not entirely new as others have considered the joint wind speed-direction dependence of concentrations (see for example Liu et al. (2015)). However, plotting the data in polar coordinates and for the purposes of source identification is new. The BPP has described in more detail in Carslaw et al. (2006) and the construction of BPP has been presented in our previous work (Chang et al., 2017).

### 3 Results and discussion

#### 315 3.1 Mass concentrations

The temporal patterns and summary statistics of hourly elemental species concentrations determined by the Xact at PEMC during March 2016-February 2017 are presented in Fig. 2. The mass concentrations of the 18 elements measured in Shanghai were sorted from high to low in Fig. 3. The one-year data set presented in the current study, to the best of our knowledge, represents the longest on-line continuous measurement series of atmospheric trace elements.




**Figure 2.** General statistical summaries of 18 trace elements measured in Shanghai.

The plots in the left panel show the time series data, where blue shows the presence of data and red shows missing data. The mean daily values are also shown in pale yellow scaled to cover the range in the data from zero to the maximum daily value. As such,

the daily values are indicative of an overall trend rather than conveying quantitative information. For each elemental species (at hourly resolution), the overall summary statistics are given. The panels on the right show the distribution of each elemental species using a histogram plot.

Taking the study period as a whole, ambient average mass concentrations of elemental species varied between detection limit (ranging from 0.05 to 20 ng m<sup>-3</sup>) and nearly 15 µg m<sup>-3</sup>, with Si as the most abundant element (mean ± 1σ; 638.7 ± 1004.5 ng m<sup>-3</sup>), followed by Fe (406 ± 385 ng m<sup>-3</sup>), K (389 ± 326 ng m<sup>-3</sup>), Ca (192 ± 383) ng m<sup>-3</sup>, Zn (120 ± 131 ng m<sup>-3</sup>), Mn (32 ± 39 ng m<sup>-3</sup>), Pb (27 ± 26 ng m<sup>-3</sup>), Ba (24 ± 25 ng m<sup>-3</sup>), V (13 ± 15 ng m<sup>-3</sup>), Cu (12 ± 11 ng m<sup>-3</sup>), Cd (10 ± 4 ng m<sup>-3</sup>), As (7 ± 7 ng m<sup>-3</sup>), Ni (6 ± 5 ng m<sup>-3</sup>), Cr (5 ± 6 ng m<sup>-3</sup>), Ag (4 ± 2.6 ng m<sup>-3</sup>), Se (2.6 ± 2.9 ng m<sup>-3</sup>), Hg (2.2 ± 1.7 ng m<sup>-3</sup>), and Au (2.2 ± 3.4 ng m<sup>-3</sup>). According to the ambient air quality standards of China (GB 3095-2012), EU (DIRECTIVE 2004/107/EC) and WHO, the atmospheric concentration limits for Cd, Hg, As, Cr (VI), Mn, V, and Ni are 5, 50 (1000 for WHO), 6 (6.6 for WHO), 0.025, 150 (WHO), 1000 (WHO), and 20 (25 for WHO) ng m<sup>-3</sup>, respectively. Therefore, airborne metal pollution in Shanghai is generally low by the current limit ceilings. Nevertheless, information regarding the specific metal compounds or chemical forms is rarely available given that most analytical techniques only record data on total metal content. In the absence of this type of information, it is generally assumed that many of the elements of anthropogenic origin (especially from combustion sources) are present in the atmosphere as oxides. Here we reconstructed the average mass concentrations of metal and crustal oxides as 5.2, 5.0, 2.8, and 3.1 µg m<sup>-3</sup> in spring, summer, fall, and winter, respectively, while the annual average concentration as 3.9 µg m<sup>-3</sup>, which accounting for 8.3% of total PM<sub>2.5</sub> mass (47 µg m<sup>-3</sup>) in 2016. Detailed calculation of the reconstructed mass has been fully described elsewhere (Dabek-Zlotorzynska et al., 2011).



355 **Figure 3.** Average mass concentrations of 18 trace elements measured in Shanghai as sorted from high to low (log10 scaling). The dark red line indicates one standard deviation.

**Table 1.** Overview of long-term and high-time resolution measurements of ambient trace elements concentrations (ng m<sup>-3</sup>) in fine particles.

Species	Shanghai, CN <sup>a</sup>	Gwangju, KP <sup>b</sup>	London, UK <sup>c</sup>	London, UK <sup>d</sup>	Barcelona, ES <sup>e</sup>	Wood Buffalo, CA <sup>f</sup>	Toronto, CA <sup>g</sup>
Ag	3.9	/	/	/	/	/	/
As	6.6	9.6	/	/	/	/	/
Au	2.2	/	/	/	/	/	/
Ba	24.2	52.0	10.3	3.7	/	/	1.9
Ca	191.5	122	78.7	50.1	130.0	54.0	54.0
Cd	9.6	/	/	/	/	/	/
Cr	4.5	/	2.3	0.8	8.0	0.04	0.24
Cu	12.0	15.5	12.8	4.9	8.0	2.04	3.1
Fe	406.2	293.0	350.3	118.9	131.0	60.0	76.8
Hg	2.2	/	/	/	/	/	/
K	388.6	732.0	27.2	23.7	82.0	31.0	27.1
Mn	31.7	24.0	4.8	2.5	6.0	1.12	1.8
Ni	6.0	3.8	0.5	0.2	3.0	0.08	0.21
Pb	27.2	49.0	2.3	1.8	12.0	/	2.4
Se	2.6	4.3	/	/	/	/	0.3
Si	638.7	/	/	/	/	143	/
V	13.4	4.6	1.3	0.6	8.0	0.21	0.11
Zn	120.3	103.0	8.9	5.3	25.0	0.88	11.3

360 **Note:** a, this study; b, Park et al., 2014; c, PM<sub>0.3-2.5</sub>, Marylebone Road (Visser et al.,

2015b); d, PM<sub>0.3-2.5</sub>, North Kensington (Visser et al., 2015b); e, Road site (Dall'Osto et al., 2013); f, Phillips-Smith et al., 2017; g, Sofowote et al., 2015. We noticed that although a huge data set of hourly resolved trace metals had been reported in Jeong et al. (2016) and Visser et al. (2015a), but that no detailed information regarding the specific mass concentrations of trace elements were given.


The toxicological effect of hazardous elements species is more evident and well known in soils and aquatic ecosystems, while few (if any) studies on the geochemical cycle of trace metals have considered the fast dynamics of trace elements in the atmosphere. Using a diversity of chemical, physical, and optical techniques, elevated atmospheric concentrations of various element species have been observed globally; however, a tiny minority of them were performed with high time resolution. As a comparison, we compiled previous work related to the near real-time measurements of trace elements concentrations in Table 1. The concentrations of most trace elements in Shanghai were commonly an order or two orders of magnitude higher than those measured in Europe and North America, and generally ranged in the same level as industrialized city like Kwangju in South Korea. Exceptionally, the concentrations of V and Ni in Shanghai were up to three times higher than that of Kwangju City. This is expected since Shanghai has the world's busiest container port, and V and Ni were substantially and almost exclusively emitted from heavy oil combustion in ship engines of ocean-going vessels (see more discussion in Section 3.2 and 3.3).

In contrast to traditional trace elements measurements, the on-line XRF used in the current study enables measurement of elements species concentrations with 1 hr resolution, which are useful both for source discrimination and in determining the processes contributing to elevated trace elements levels through investigation of their seasonal, weekly, weekday-weekend, and diurnal cycles (Fig. S2-S8; see discussion below).

### 3.2 Source analysis



In the PMF analysis, three to ten factor solutions were initially examined, from which possible solutions (i.e., four to six factor solutions) were chosen based on the change of  $Q/Q_{\text{exp}}$ , the achievement of the constant and global minimum of  $Q$ , the displacement of factor elements, and the interpretation of physically meaningful factors (Text S1; see discussion below). The most reliable solution was explained by five factors. The chemical profiles and average contributions of the five factors are presented in Fig. 4 with the time-series evolution of these factors included in the Supplement (Fig. S9). On the one hand, we will use various mathematical and physical criteria to constrain different solutions of source apportionment. On the other hand, we will take CPF and BPP as diagnostic tools for quickly gaining the idea of potential source regions, which in turn will contribute to further analysis of source apportionment. Ultimately, the five factors were assigned to different sources, i.e., traffic-related, shipping, nonferrous metal smelting, coal combustion, and ferrous metal smelting.




**Figure 4.** PMF-resolved source profiles (concentration and % of species apportioned

to factor) and average contributions (in the parentheses) of individual sources to measured total PM<sub>2.5</sub> elements in Shanghai. The notable species for each factor/source are marked in red.

### 3.2.1 Traffic-related

Factor 1 was characterized by a large mass fraction of Ca, Fe, Ba, and Si, which explained 89.5%, 50.3%, 76.5%, and 62.6% of the variation, respectively. This mixed factor is similar to that reported by Amato et al. (2009, 2013), Bukowiecki et al. (2010), Harrison et al. (2012), and Visser et al. (2015b). In urban atmosphere, Fe can be released from engine oil or catalyst equipped gasoline vehicles (Chen et al., 2007). Besides, Fe is linked to non-exhaust emissions such as brake wear because it is the support material for brake pads, and the agents present in brake linings typically consist of Ba, Mn and Cu (Lough et al., 2005; Hjortenkrans et al., 2007; Dall'Osto et al., 2016). Therefore, Fe and Ba can be regarded as chemical tracers for traffic-related source (exhaust and non-exhaust) (Thorpe et al., 2008; Lin et al., 2015). Ca and Si are known as two of the most abundant elements in the upper continental crust, and their atmospheric origins typically attributed to wind-blown dust. Located on the eastern coast of China, Shanghai rarely receives long-range transport of crustal matter from aeolian dust and the Gobi Desert in northwestern China (Huang et al., 2013). Sampling in the urban area of Shanghai, airborne Ca and Si should be dominated by anthropogenic activities like road fugitive dust or urban construction works. In Fig. S10, significant correlations are observed for Ca, Si, Fe, and Ba, suggesting that the measured Ca and Si during our study period were more likely derived from road fugitive dust. Therefore, factor 1 can be assigned to traffic-related source and it was the largest source in Shanghai, accounting for 46.0% (683.9 ng m<sup>-3</sup>) of the total measured elemental mass in PM<sub>2.5</sub>.




**Figure 5.** Diurnal variation of PMF-derived elemental concentration for factor 1. The red line, bottom boundary, and upper boundary represent the mean, 1<sup>st</sup> quartile, and 3<sup>rd</sup> quartile of the concentration value, respectively.

Hourly measurements over one-year periods provide a unique opportunity to examine the diurnal profile of factor 1. As reported in Fig. 5, the concentration of trace elements contributed by factor 1 shows a marked bimodal diurnal cycle, with average values at rush hours are over two times higher than at nighttime. Such variation pattern agrees well with the diurnal variation of traffic flow in Shanghai (Chang et al., 2016a), further confirming that factor 1 can be interpreted as traffic-related emissions.

### 3.2.2 Shipping

In Fig. 4, V (100%) and Ni (74%) comes almost exclusively from factor 2, while factor 2 contributes to less than 10% of any other elemental species. V is typically emitted from oil and petrochemical refining and combustion, and natural gas extraction and processing (Duce and Hoffman, 1976; Hope, 1994; Shafer et al., 2012). From CPF and BPP analysis (Fig. S11), higher concentrations of both V and Ni were observed when winds originated from east, northeast, and southeast directions. The most dominant directions were east and southeast, suggesting the influence from coastal port cluster or petroleum refinery industry located east/southeast of Shanghai (Fig. 1). Gathering evidence revealed that the ratio of V/Ni can serve as a robust indicator of shipping emissions (Tao et al., 2013; Celo et al., 2015; Liu et al., 2017; Viana et al., 2009). Recent study in Shanghai port suggested that the ratio of V/Ni in aerosols emitted from heavy oil combustion of ocean-going ship engines was 3.4 on average (Zhao et al., 2013).

Here measured in urban area, the average ratio of V/Ni in our study was 3.1 with slightly seasonal changes (Fig. 6), indicating V- and Ni-containing aerosols from shipping emissions subject to minor atmospheric transformation. In short, factor 2 likely corresponds to shipping emissions (instead of petrochemical refining), which is consistent with the results of many previous source apportionment works (e.g., Liu et al., 2017; Zhao et al., 2013; Cesari et al., 20014; Healy et al., 2010).




**Figure 6.** Linear correlation analysis between V (x axis) and Ni (y axis) in Shanghai during four seasons.

Although shipping emissions only contribute to 5.9% of trace elements in Shanghai urban center, its share can be expected to greatly increase in harbor district (Zhao et al., 2013). The good news is that since 1 January 2016, the sea areas of Shanghai and its neighboring ports were designed as shipping emission control area, requiring use of lower sulfur fuels in place of heavy fuel oil in main engines of ships (Zhen et al., 2018). Therefore, it is critically important to assess the impacts of fuel changes on air quality in Shanghai in the future through continuous measurements of trace elements.

### 3.2.3 Nonferrous metal smelting

The predominant elements found in factor 3 were Au (100%), Cd (65%), and Ag (63%) with 37% of Hg. These four heavy metals are important associated elements in Cu, Pb, and Zn ores. In fact, Cu, Pb, and Zn smelting represent the three most common form of nonferrous metal smelting in China (Tian et al., 2015). Because of high temperatures during roasting, sintering and smelting process for the extraction of Cu, Pb, and Zn

from ores, metals like Au, Cd, Ag, and Hg in nonferrous metal ores will inevitably be vaporized and released into the flue gas (Pacyna and Pacyna, 2001; Wu et al., 2012). Therefore, factor 3 was interpreted as nonferrous metal smelting emissions and the contribution of this factor was 14.9% ( $221.6 \text{ ng m}^{-3}$ ) to total measured elemental mass in  $\text{PM}_{2.5}$ .




**Figure 7.** Percentile rose plot of Au (a) and Hg (b) concentrations in Shanghai between March 2016 and February 2017. The percentile intervals are shaded and shown by wind direction.

To further pinpoint the specific subsector of nonferrous metal smelting, here we calculate percentile concentration levels of Au and Hg, and plot them by wind direction in Fig. 7 (and Ag, Cd in Fig. S12). It clearly shows Au and Hg largely share the same source region that different from Ag and Cd, indicating that Au and Hg were emitted from a similar subsector of nonferrous metal smelting. In Shanghai, Zn smelting is the most important contributor of Hg emissions from the nonferrous metal smelting sector. Therefore, element Au resolved in factor 3 during our study period can be expected to be originated from Zn smelting.

### 3.2.4 Coal combustion

The most abundant elements found in factor 4 were As, Se, Pb, Hg (explaining 56% to 95% of the variation) with some contributions of Cu (30%), Zn (29%) and unexpected large amount of K (53%). As, Se, Pb, Hg, and Cu are typical marker elements for coal combustion. In China, 73% of As, 62% of Se, 56% of Pb, and 47% of Hg were found to be emitted from coal combustion. Coal consumption in southern China (including

Shanghai) is mainly driven by industrial boilers and power plant, while in norther China, coal-based heating is also a major sector of coal consumption (Tian et al., 2015). Seasonally, the average mass concentration of coal combustion-related PM<sub>2.5</sub> trace elements during winter (407 ng m<sup>-3</sup>) was much higher than that during spring (296 ng m<sup>-3</sup>), summer (148 ng m<sup>-3</sup>), and fall (210 ng m<sup>-3</sup>) (Fig. 8). This seasonal pattern was not observed for other sources (not shown). Shanghai has a humid subtropical climate and experiences four distinct seasons. Winters are chilly and damp, with northwesterly winds from norther China can transport air pollutants (including trace elements) caused by coal-based heating to Shanghai atmosphere (Huang et al., 2013; Chang et al., 2017). As the largest city-scale coal consumer in China, coal combustion contributed to 275.4 ng m<sup>-3</sup> or 18.5% of PM<sub>2.5</sub> trace elements during our study period.




**Figure 8.** Seasonal variation of elements concentrations contributed by coal combustion in Shanghai. The error bar indicates one standard derivation.

Traditionally, K in particles was considered to be originated from biomass burning along with some contribution of fugitive dust (Zhang et al., 2010; Hueglin et al., 2005; Fang et al., 2015). Here we show that over half of element K in urban Shanghai were derived from coal combustion. The reason for this discrepancy maybe that in most previous studies, K in particles was pretreated using deionized water to extract (Wang et al., 2013b). In fact, K has high mineral affinity (elements associated with aluminosilicates, carbonates and other minerals in coal ash), and in some extreme cases, only about 1% of K in fly ash from coal combustion can be extracted by water (Querol

et al., 1996). For example, particles collected from coal combustion by Wang et al. (2013b) were extracted with deionized water, then atomized and measured by an ATOFMS. The ATOFMS mass spectrum contained relatively low K peak. The observation by Wang et al. (2013) was not consistent with that of Suess et al. (2002), in which they detected larger K peaks in ATOFMS spectra for coal combustion particles in an in situ measurement (i.e. freshly emitted particles were directly introduced into ATOFMS and measured).

### 3.2.5 Ferrous metal smelting

Factor 5 was distinguished by high levels of Cr, Mn, and Zn representing 100%, 56%, and 52% of the explained variation, respectively. These elements are typically emitted from ferrous metal smelting. For example, the steel production industry represents the dominant contributor to Zn emissions, accounting for about 60% in China (Tian et al., 2015). Driven by rapid modernization of its infrastructure and manufacturing industries, China produced more than 49% of world steel production in 2017 (around 830 million tons), and 6 of 10 of the largest steel producers are in China (data retrieved from <https://www.worldsteel.org>). Headquartered in Shanghai (20 km northwest of the sampling site), the Baosteel is the fifth-largest steel producer in the world measured by crude steel output, with an annual output of around 35 million tons. Meanwhile, there are several factories of ferrous metal processing located in western Shanghai (Fig. 1). Since element Cr is reported to be transported over distances by air flow (Perry et al., 1999), the presence of ferrous metal smelting activities in the west/northwest of the sampling site is inferred to be associated with this factor based on the results of CPF and BBP in Fig. 9. Overall, ferrous metal smelting contributed 218.9 ng m<sup>-3</sup> or 14.7% of PM<sub>2.5</sub> trace elements in Shanghai.



**Figure 9.** Conditional probability function analysis (left) and bivariate polar plots

(right) of seasonal concentrations (in  $\text{ng m}^{-3}$ ) of Cr in Shanghai between March 2016 and February 2017. The center of each plot (centered at the sampling site) represents a  
545 wind speed of zero, which increases radially outward. The concentration is shown by  
the color scale.

### 3.3 Precipitation effect


Theoretically, precipitation could enhance the wet scavenging of airborne pollutants and reduce their ability to suspend because the increased moisture might capture the  
550 particles on the road surface (Kuhns et al., 2003; Karanasiou et al., 2011). Water spray (from sprinkler on road or atop tall building) to simulate natural types of precipitation has been proposed as an important abatement strategy to facilitate the reduction of ambient PM concentrations (including trace elements) in urban China (Liu et al., 2014; Yu, 2014). However, several field measurements revealed that water spray activities did  
555 not influence PM mass levels (e.g., Karanasiou et al., 2012, 2014). Taking advantage of our simultaneous and hourly record of precipitation amount (up to 36.8 mm) and elements concentration, here we evaluate the effects of precipitation on the mitigation of  $\text{PM}_{2.5}$  trace elements. The precipitation (all were in the form of rainfall) distributed during the full year of measurements is shown in Figure S13. The mass concentrations  
560 of trace elements six hours before and after precipitation events were compared from the perspective of individual species and sources. The precipitation event in this study is defined as (1) there are at least six consecutive hours with hourly rainfall amount higher than 1 mm; (2) the consecutive no-rainy time in a precipitation event should less than six hours; (3) the total no-rainy time should less than 1/3 of the entire time of a precipitation event; (4) if the rainfall amount of a specific hour is less than 0.1 mm, and there are at least three no-rainy hours before and after the rainy hour, then this hour should be treated as no-rainy hour. Consequently, 12 precipitation events during our study period were identified with the duration time and accumulated rainfall ranging from 7 to 55 hours, and 26.4 to 217.5 mm, respectively (Table S2).

#### 3.3.1 Change of mass concentration by species

The average mass concentration of each elemental species before, during, and after



every precipitation event is presented in Fig. S13. If precipitation effectively scavenge and remove aerosol, then the mass concentrations of trace elements during a precipitation event should lower than that before and after this precipitation event. However, there is no uniform variation pattern in Fig. S14, indicating that precipitation may not be the predominant factor to influence ambient elements mass in some cases. For example, most elemental species had a relatively higher mass concentration during the 12<sup>th</sup> precipitation event (lasted from 09:00 25 December to 22:00 26 December; Fig. S14). This can be explained by the much less anthropogenic activities during the periods prior to (3:00 to 8:00) and after (23:00 to 3:00 the next day) the 12<sup>th</sup> precipitation event.



**Figure 10.** (a) The variation of the overall mass concentration of each elemental species before, during, and after the total 12 precipitation events; (b) scatter plot of the relationship between the rainfall amount of each precipitation event and the decrement rate of Zn concentration from the period before precipitation to the period after precipitation.


For each elemental species, the variation of mass concentration before, during, and after every precipitation event were aggregated and reported in Fig. 10a. Before precipitation

events, the mass concentrations of all species except Cr and Au were higher than that during precipitation events (notably V, Zn, Fe), suggesting that water spray could generally help to reduce PM<sub>2.5</sub> trace elements load in the atmosphere. After precipitation events, there were six species (notably V and Au) with their mass concentrations lower than that during precipitation events, indicating potential long-lasting aftereffect of precipitation scavenging. Among all elemental species, the mass concentrations of Zn and Se fluctuate as the most ideal V-shape, which properly reflect the cycle of precipitation. However, as shown in Fig. 10b, a linear relationship cannot be observed between the decrement rate of Zn concentration and rainfall amount of each precipitation event. Although we failed to pinpoint the exact value in this study, our results imply that there is a threshold of precipitation amount to lower ambient PM<sub>2.5</sub> trace elements mass.

### **3.3.1 Change of mass concentration by sources**

The variation of the overall mass concentration of trace elements contributed by each source and their relative contributions before, during, and after the total 12 precipitation events is shown in Fig. 11a to 10e, and Fig. 11f, respectively. The mass concentration of traffic-related trace elements experienced the sharpest decrease during the transition of no-rainy hours to rainy hours (159%), and a moderate rebound after precipitation (35%). Fang et al. (2015) found that mobile source emissions generated through mechanical processes (re-entrained road dust, tire and brake wear) and processing by secondary sulfate were major contributors to water-soluble metals. In our study, traffic-related source mainly includes road dust and brake wear, which can not only be easily removed through precipitation but also can hardly be blown up from wet road surface after raining. In comparison, the mass contribution of coal combustion source was also wet removed rapidly first (139%) due to its tracer elements like As, Se, Pb, and Hg have larger water-soluble fraction. However, after precipitation, the contribution of coal combustion source dramatically increased over two times (Fig. 11d and 11f). This can be explained that different from traffic-related source, coal combustion-related trace elements are generally emitted through elevated chimneys in the sectors of industrial


broilers and power plants. The mass concentrations of trace elements contributed by nonferrous and ferrous metal smelting during the three periods kept quite flat (Fig. 11c and 11d), suggesting that precipitation has little effect on ambient trace elements emitted from metal smelting activities. Nevertheless, given that traffic-related and coal combustion are the dominant contributors to ambient PM<sub>2.5</sub> trace elements, our results validate that water spray could be an effective approach to help curb the severe atmospheric metal pollution in many Chinese cities.



**Figure 11.** The variation of the overall mass concentration of trace elements contributed by traffic-related (a), shipping (b), nonferrous metal smelting (c), coal combustion (d), and ferrous metal smelting (e), and their relative contributions (f) before, during, and after the total 12 precipitation events.

In Fig. 11b and 11f, the contribution of shipping emissions to ambient trace elements (mainly V and Ni) during the three periods reduced continuously. Mostly transported from Eastern China sea, V and Ni almost exclusively originated from the east of the sampling site. In other words, the contribution of shipping emissions to urban atmosphere is supposed to be very sensitive to wind speed and wind direction in Shanghai. The wind roses for the three periods are presented in Fig. 12. It shows that before precipitation events, the average wind speed ( $\pm 1\sigma$ ) was the lowest ( $2.3 \pm 1.3 \text{ m s}^{-1}$ ), and easterly winds prevail in most times. These factors are favorable to the transportation of shipping emissions from Eastern China sea and then accumulated in Shanghai urban atmosphere. In contrast to the period before precipitation events, the average wind speed after precipitation events was the highest ( $3.1 \pm 2.1 \text{ m s}^{-1}$ ) with

northwesterly and northerly winds from mainland China can dilute shipping-related trace elements to the lowest levels (Fig. 11b). In brief, the mass concentration of shipping-related trace elements in Shanghai urban atmosphere is more likely to be influenced by winds instead of precipitation.



**Figure 12.** Wind roses plots for the periods before, during, and after the twelve precipitation events. The average wind speed ( $\pm 1\sigma$ ) for each period is shown in bracket.

#### 4. Conclusion and outlook

This paper presents the results from a year-long, near real-time measurement study of 18 trace elements (Si, Fe, K, Ca, Zn, Mn, Pb, Ba, V, Cu, Cd, As, Ni, Cr, Ag, Se, Hg, and Au) in  $PM_{2.5}$  using a Xact multi-metal monitor, conducted at an urban site in Shanghai from March 2016 to February 2017. The scientific significance of this work can be reflected by the general findings as follows:

-The Xact multi-metals monitor was demonstrated as a valuable and practical tool for ambient monitoring of atmospheric trace elements by comparing online monitoring results with ICP analyses of offline filter samples.

-The metal concentrations in Shanghai are one or two orders of magnitude higher than in north America and Europe, highlighting the need to allocate more scientific, technical, and legal resources on controlling metal emissions in China.

-The total of metal related species comprised approximately 8.3% of the  $PM_{2.5}$  mass, which should not be ignored in China's recent epidemiologic study of attributing hospital emergency-room visits to  $PM_{2.5}$  chemical constituents.

-The full coverage of trace elemental species (18) measurement and the high temporal frequency (hourly) in the work provided unprecedented details regarding the temporal evolution of metal pollution and its potential sources in Shanghai.

-Five sources, i.e., traffic-related, shipping, nonferrous metal melting, coal combustion, and ferrous metal smelting were identified by PMF analysis, which contributed to 46%, 6%, 15%, 18%, and 15% of ambient PM<sub>2.5</sub> trace elements, respectively.

-The dominant contributors of trace elements (traffic-related and coal combustion) can be effectively removed through precipitation from the atmosphere, suggesting that water spray can be used to curb PM<sub>2.5</sub> trace elements in urban atmosphere.

A greater value and more interesting topic to the scientific community would be to fully assess the role of PM<sub>2.5</sub> chemical constituents (including metal species) and sources of emission to human health. Looking towards the future, three major steps will be taken toward thoroughly addressing these questions. Firstly, characterizing the chemical and isotopic (including metal species) signatures of emission sources will be intensively undertaken through field sampling as well as for laboratory simulations (see example of Geagea et al. (2007)). Secondly, the Xact multi-metals monitor, Sunset OC/EC analyzer (Chang et al., 2017), and MARGA (Monitoring of AeRosols and Gases) platform will be collocated across a rural-urban-background transect to simultaneously measure hourly metal species, carbonaceous aerosols, and inorganic aerosol components in PM<sub>2.5</sub>. Lastly, integrating all available information regarding PM<sub>2.5</sub> chemical species and isotopes into a receptor model or atmospheric chemical transport model will be carried forward to create more specific and confident source apportionment results.

### **Competing interests**

The authors declare that they have no competing interests.

### **Data availability**

Data are available from the corresponding authors on request.

## Acknowledgements

This study was supported by the National Key Research and Development Program of China (2017YFC0212700), National Natural Science Foundation of China (Grant nos. 41705100, 91644103, 41603104, 41429501, and 91644105), Provincial Natural Science Foundation of Jiangsu (BK20170946), University Science Research Project of Jiangsu Province (17KJB170011), and the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD). Many thanks go to Tony Dore at the Centre for Ecology and Hydrology, Edinburgh, United Kingdom for his linguistic corrections.

## Reference

- Amato, F., Pandolfi, M., Escrig, A., Querol, X., Alastuey, A., Pey, J., Perez, N., and Hopke, P. K.: Quantifying road dust resuspension in urban environment by Multilinear Engine: A comparison with PMF2, *Atmos. Environ.*, 43, 2770-2780, doi: 10.1016/j.atmosenv.2009.02.039, 2009.
- Amato, F., Pandolfi, M., Moreno, T., Furger, M., Pey, J., Alastuey, A., Bukowiecki, N., Prevot, A. S. H., Baltensperger, U., and Querol, X.: Sources and variability of inhalable road dust particles in three European cities, *Atmos. Environ.*, 45, 6777-6787, doi: 10.1016/j.atmosenv.2011.06.003, 2011.
- Amato, F., Schaap, M., Denier van der Gon, H. A. C., Pandolfi, M., Alastuey, A., Keuken, M., and Querol, X.: Short-term variability of mineral dust, metals and carbon emission from road dust resuspension, *Atmos. Environ.*, 74, 134-140, doi: 10.1016/j.atmosenv.2013.03.037, 2013.
- Brandt, C. and van Eldik, R.: Transition metal-catalyzed oxidation of sulfur (IV) oxides. Atmospheric-relevant processes and mechanisms, *Chem. Rev.*, 95, 119-190, doi: 10.1021/cr00033a006, 1995.
- Bukowiecki, N., Hill, M., Gehrig, R., Zwicky, C. N., Lienemann, P., Hegedüs, F., Falkenberg, G., Weingartner, E., and Baltensperger, U.: Trace metals in ambient air: Hourly size-segregated mass concentrations determined by synchrotron-XRF, *Environ. Sci. Technol.*, 39, 5754-5762, doi: 10.1021/es048089m, 2005.
- Bukowiecki, N., Lienemann, P., Hill, M., Furger, M., Richard, A., Amato, F., Prévôt, A. S. H., Baltensperger, U., Buchmann, B., and Gehrig, R.: PM<sub>10</sub> emission factors for non-exhaust particles generated by road traffic in an urban street canyon and along a freeway in Switzerland, *Atmos. Environ.*, 44, 2330-2340, doi: 10.1016/j.atmosenv.2010.03.039, 2010.
- Burnett, R. T., Pope, C. A., III, Ezzati, M., Olives, C., Lim, S. S., Mehta, S., Shin, H. H., Singh, G., Hubbell, B., Brauer, M., Anderson, H. R., Smith, K. R., Balmes, J. R., Bruce, N. G., Kan, H., Laden, F., Pruss-Ustun, A., Turner, M. C., Gapstur, S. M., Diver, W. R., and Cohen, A.: An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure,

- 730 Environ. Health. Perspect., 122, 397-403, doi: 10.1289/ehp.1307049, 2014.
- Carslaw, D. C., Beevers, S. D., Ropkins, K., and Bell, M. C.: Detecting and quantifying aircraft and other on-airport contributions to ambient nitrogen oxides in the vicinity of a large international airport, *Atmos. Environ.*, 40, 5424-5434, doi: 10.1016/j.atmosenv.2006.04.062, 2006.
- 735 Carslaw, D. C. and Ropkins, K.: Openair - An R package for air quality data analysis, *Environ. Modell. Softw.*, 27-28, 52-61, doi: 10.1016/j.envsoft.2011.09.008, 2012.
- Cate, D. M., Noblitt, S. D., Volckens, J., and Henry, C. S.: Multiplexed paper analytical device for quantification of metals using distance-based detection, *Lab Chip*, 15, 2808-2818, doi: 10.1039/c5lc00364d, 2015.
- 740 Celo, V., Dabek-Zlotorzynska, E., and McCurdy, M.: Chemical characterization of exhaust emissions from selected Canadian marine vessels: the case of trace metals and lanthanoids, *Environ. Sci. Technol.*, 49, 5220-5226, doi: 10.1021/acs.est.5b00127, 2015.
- Cesari, D., Genga, A., Ielpo, P., Siciliano, M., Mascolo, G., Grasso, F. M., and Contini, D.: Source apportionment of PM<sub>2.5</sub> in the harbour-industrial area of Brindisi (Italy): Identification and estimation of the contribution of in-port ship emissions, *Sci. Total Environ.*, 497-498, 392-400, doi: 10.1016/j.scitotenv.2014.08.007, 2014.
- 745 Chang, Y., Zou, Z., Deng, C., Huang, K., Collett, J. L., Lin, J., and Zhuang, G.: The importance of vehicle emissions as a source of atmospheric ammonia in the megacity of Shanghai, *Atmos. Chem. Phys.*, 16, 3577-3594, doi: 10.5194/acp-16-3577-2016, 2016.
- 750 Chang, Y., Deng, C., Cao, F., Cao, C., Zou, Z., Liu, S., Lee, X., Li, J., Zhang, G., and Zhang, Y.: Assessment of carbonaceous aerosols in Shanghai, China: Long-term evolution, seasonal variations and meteorological effects, *Atmos. Chem. Phys. Discuss.*, 2017, 1-46, doi: 10.5194/acp-2017-50, 2017.
- 755 Charrier, J. G. and Anastasio, C.: On dithiothreitol (DTT) as a measure of oxidative potential for ambient particles: evidence for the importance of soluble transition metals, *Atmos. Chem. Phys.*, 12, 9321-9333, doi: 10.5194/acp-12-9321-2012, 2012.
- 760 Chen, B., Stein, A. F., Castell, N., Gonzalez-Castanedo, Y., Sanchez de la Campa, A. M., and de la Rosa, J. D.: Modeling and evaluation of urban pollution events of atmospheric heavy metals from a large Cu-smelter, *Sci. Total Environ.*, 539, 17-25, doi: 10.1016/j.scitotenv.2015.08.117, 2016.
- 765 Chen, L. W. A., Watson, J. G., Chow, J. C., and Magliano, K. L.: Quantifying PM<sub>2.5</sub> source contributions for the San Joaquin Valley with multivariate receptor models, *Environ. Sci. Technol.*, 41, 2818-2826, doi: 10.1021/es0525105, 2007.
- Cooper, J. A., Petterson, K., Geiger, A., Siemers, A., and Rupprecht, B.: Guide for developing a multi-metals, fence-line monitoring plan for fugitive emissions using X-ray based monitors, Cooper Environmental Services, Portland, Oregon, 1-42, 2010.
- 770 Dabek-Zlotorzynska, E., Dann, T. F., Kalyani Martinelango, P., Celo, V., Brook, J. R., Mathieu, D., Ding, L., and Austin, C. C.: Canadian National Air Pollution Surveillance (NAPS) PM<sub>2.5</sub> speciation program: Methodology and PM<sub>2.5</sub> chemical

composition for the years 2003–2008, *Atmos. Environ.*, 45, 673-686, doi: 10.1016/j.atmosenv.2010.10.024, 2011.

775 Dall'Osto, M., Beddows, D. C. S., Harrison, R. M., and Onat, B.: Fine iron aerosols are internally mixed with nitrate in the urban European atmosphere, *Environ. Sci. Technol.*, 50, 4212-4220, doi: 10.1021/acs.est.6b01127, 2016.

780 Dall'Osto, M., Querol, X., Amato, F., Karanasiou, A., Lucarelli, F., Nava, S., Calzolari, G., and Chiari, M.: Hourly elemental concentrations in PM<sub>2.5</sub> aerosols sampled simultaneously at urban background and road site during SAPUSS - diurnal variations and PMF receptor modelling, *Atmos. Chem. Phys.*, 13, 4375-4392, doi: 10.5194/acp-13-4375-2013, 2013.

785 DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281-8289, doi: 10.1021/ac061249n, 2006.

790 Duan, J. and Tan, J.: Atmospheric heavy metals and arsenic in China: Situation, sources and control policies, *Atmos. Environ.*, 74, 93-101, doi: 10.1016/j.atmosenv.2013.03.031, 2013.

Duce, R. A., and Hoffman, G. L.: Atmospheric vanadium transport to the ocean, *Atmos. Environ.*, 10, 989-996, doi: 10.1016/0004-6981(76)90207-9, 1976.

795 Fan, Q., Zhang, Y., Ma, W., Ma, H., Feng, J., Yu, Q., Yang, X., Ng, S. K. W., Fu, Q., and Chen, L.: Spatial and seasonal dynamics of ship emissions over the Yangtze River Delta and East China Sea and their potential environmental influence, *Environ. Sci. Technol.*, 50, 1322-1329, doi: 10.1021/acs.est.5b03965, 2016.

800 Fang, T., Guo, H., Verma, V., Peltier, R. E., and Weber, R. J.: PM<sub>2.5</sub> water-soluble elements in the southeastern United States: Automated analytical method development, spatiotemporal distributions, source apportionment, and implications for health studies, *Atmos. Chem. Phys.*, 15, 11667-11682, doi: 10.5194/acp-15-11667-2015, 2015.

Fergusson, J. E.: *The heavy elements: Chemistry, environmental impact and health effects*, Pergamon Press, Oxford, UK, pp 614, 1990.

805 Furger, M., Minguillón, M. C., Yadav, V., Slowik, J. G., Hüglin, C., Fröhlich, R., Petterson, K., Baltensperger, U., and Prévôt, A. S. H.: Elemental composition of ambient aerosols measured with high temporal resolution using an online XRF spectrometer, *Atmos. Meas. Tech.*, 10, 2061-2076, doi: 10.5194/amt-10-2061-2017, 2017.

810 Geagea, M. L., Stille, P., Millet, M., and Perrone, T.: REE characteristics and Pb, Sr and Nd isotopic compositions of steel plant emissions, *Sci. Total Environ.*, 373, 404-419, doi: 10.1016/j.scitotenv.2006.11.011, 2007.

Gross, D. S., Gälli, M. E., Silva, P. J., and Prather, K. A.: Relative sensitivity factors for alkali metal and ammonium cations in single-particle aerosol time-of-flight mass spectra, *Anal. Chem.*, 72, 416-422, doi: 10.1021/ac990434g, 2000.

815 Han, T., Qiao, L., Zhou, M., Qu, Y., Du, J., Liu, X., Lou, S., Chen, C., Wang, H., Zhang, F., Yu, Q., and Wu, Q.: Chemical and optical properties of aerosols and their interrelationship in winter in the megacity Shanghai of China, *J. Environ. Sci.*



- (China), 27, 59-69, doi: 10.1016/j.jes.2014.04.018, 2015.
- 820 Harrison, R. M., Jones, A. M., Gietl, J., Yin, J., and Green, D. C.: Estimation of the contributions of brake dust, tire wear, and resuspension to nonexhaust traffic particles derived from atmospheric measurements, *Environ. Sci. Technol.*, 46, 6523-6529, doi: 10.1021/es300894r, 2012.
- 825 Healy, R. M., Hellebust, S., Kourtshev, I., Allan, A., O'Connor, I. P., Bell, J. M., Healy, D. A., Sodeau, J. R., and Wenger, J. C.: Source apportionment of PM<sub>2.5</sub> in Cork Harbour, Ireland using a combination of single particle mass spectrometry and quantitative semi-continuous measurements, *Atmos. Chem. Phys.*, 10, 9593-9613, doi: 10.5194/acp-10-9593-2010, 2010.
- 830 Holden, P. A., Gardea-Torresdey, J. L., Klaessig, F., Turco, R. F., Mortimer, M., Hund-Rinke, K., Cohen Hubal, E. A., Avery, D., Barceló, D., Behra, R., Cohen, Y., Deydier-Stephan, L., Ferguson, P. L., Fernandes, T. F., Herr Harthorn, B., Henderson, W. M., Hoke, R. A., Hristozov, D., Johnston, J. M., Kane, A. B., Kapustka, L., Keller, A. A., Lenihan, H. S., Lovell, W., Murphy, C. J., Nisbet, R. M., Petersen, E. J., Salinas, E. R., Scheringer, M., Sharma, M., Speed, D. E., Sultan, Y., Westerhoff, P., White, J. C., Wiesner, M. R., Wong, E. M., Xing, B., Steele
- 835 Horan, M., Godwin, H. A., and Nel, A. E.: Considerations of environmentally relevant test conditions for improved evaluation of ecological hazards of engineered nanomaterials, *Environ. Sci. Technol.*, 50, 6124-6145, doi: 10.1021/acs.est.6b00608, 2016.
- 840 Honda, T., Eliot, M. N., Eaton, C. B., Whitsel, E., Stewart, J. D., Mu, L., Suh, H., Szpiro, A., Kaufman, J. D., Vedal, S., and Wellenius, G. A.: Long-term exposure to residential ambient fine and coarse particulate matter and incident hypertension in post-menopausal women, *Environ. Int.*, 105, 79-85, doi: 10.1016/j.envint.2017.05.009, 2017.
- 845 Hueglin, C., Gehrig, R., Baltensperger, U., Gysel, M., Monn, C., and Vonmont, H.: Chemical characterisation of PM<sub>2.5</sub>, PM<sub>10</sub> and coarse particles at urban, near-city and rural sites in Switzerland, *Atmos. Environ.*, 39, 637-651, doi: 10.1016/j.atmosenv.2004.10.027, 2005.
- 850 Hu, X., Zhang, Y., Ding, Z., Wang, T., Lian, H., Sun, Y., and Wu, J.: Bioaccessibility and health risk of arsenic and heavy metals (Cd, Co, Cr, Cu, Ni, Pb, Zn and Mn) in TSP and PM<sub>2.5</sub> in Nanjing, China, *Atmos. Environ.*, 57, 146-152, doi: 10.1016/j.atmosenv.2012.04.056, 2012.
- 855 Huang, C., Chen, C. H., Li, L., Cheng, Z., Wang, H. L., Huang, H. Y., Streets, D. G., Wang, Y. J., Zhang, G. F., and Chen, Y. R.: Emission inventory of anthropogenic air pollutants and VOC species in the Yangtze River Delta region, China, *Atmos. Chem. Phys.*, 11, 4105-4120, DOI 10.5194/acp-11-4105-2011, 2011.
- Huang, K., Zhuang, G., Lin, Y., Wang, Q., Fu, J. S., Fu, Q., Liu, T., and Deng, C.: How to improve the air quality over megacities in China: Pollution characterization and source analysis in Shanghai before, during, and after the 2010 World Expo, *Atmos. Chem. Phys.*, 13, 5927-5942, doi: 10.5194/acp-13-5927-2013, 2013.
- 860 Huang, W., Cao, J., Tao, Y., Dai, L., Lu, S.-E., Hou, B., Wang, Z., and Zhu, T.: Seasonal variation of chemical species associated with short-term mortality effects of PM<sub>2.5</sub>

in Xi'an, a central city in China, *Am. J. Epidemiol.*, 175, 556-566, doi: 10.1093/aje/kwr342, 2012.

Hope, B. K.: A global biogeochemical budget for vanadium, *Sci. Total Environ.*, 141, 1-10, doi: 10.1016/0048-9697(94)90012-4, 1994.

Iyengar, V. and Woittiez, J.: Trace elements in human clinical specimens: evaluation of literature data to identify reference values, *Clin. Chem.*, 34, 474-481, 1988.

Jeong, C. H., Wang, J. M., and Evans, G. J.: Source apportionment of urban particulate matter using hourly resolved trace metals, organics, and inorganic aerosol components, *Atmos. Chem. Phys. Discuss.*, 1-32, doi: 10.5194/acp-2016-189, 2016.

John, D. H.: "Heavy metals"-a meaningless term?, *Pure Appl. Chem.*, 74, 793-807, doi: 10.1351/pac200274050793, 2002.

Jomova, K. and Valko, M.: Advances in metal-induced oxidative stress and human disease, *Toxicol.*, 283, 65-87, doi: 10.1016/j.tox.2011.03.001, 2011.

Karanasiou, A., Moreno, T., Amato, F., Lumberras, J., Narros, A., Borge, R., Tobías, A., Boldo, E., Linares, C., Pey, J., Reche, C., Alastuey, A., and Querol, X.: Road dust contribution to PM levels – Evaluation of the effectiveness of street washing activities by means of Positive Matrix Factorization, *Atmos. Environ.*, 45, 2193-2201, doi: 10.1016/j.atmosenv.2011.01.067, 2011.

Karanasiou, A., Moreno, T., Amato, F., Tobías, A., Boldo, E., Linares, C., Lumberras, J., Borge, R., Alastuey, A., and Querol, X.: Variation of PM<sub>2.5</sub> concentrations in relation to street washing activities, *Atmos. Environ.*, 54, 465-469, doi: 10.1016/j.atmosenv.2012.02.006, 2012.

Karanasiou, A.: Road dust emission sources and assessment of street washing effect, *Aerosol Air Qual. Res.*, doi: 10.4209/aaqr.2013.03.0074, 2014.

Kastury, F., Smith, E., and Juhasz, A. L.: A critical review of approaches and limitations of inhalation bioavailability and bioaccessibility of metal(loid)s from ambient particulate matter or dust, *Sci. Total Environ.*, 574, 1054-1074, doi: 10.1016/j.scitotenv.2016.09.056, 2017.

Khalaf, F., Literathy, P., and Anderlini, V.: Vanadium as a tracer of oil pollution in the sediments of Kuwait, *sediment/freshwater interaction*, Dordrecht, 1982, 147-154.

Kim, K. H., Kabir, E., and Jahan, S. A.: A review on the distribution of Hg in the environment and its human health impacts, *J. Hazard. Mater.*, 306, 376-385, doi: 10.1016/j.jhazmat.2015.11.031, 2016.

Kloog, I., Ridgway, B., Koutrakis, P., Coull, B. A., and Schwartz, J. D.: Long- and short-term exposure to PM<sub>2.5</sub> and mortality: Using novel exposure models, *Epidemiol.*, 24, 555-561, doi: 10.1097/EDE.0b013e318294beaa, 2013.

Kuhns, H., Etyemezian, V., Green, M., Hendrickson, K., McGown, M., Barton, K., and Pitchford, M.: Vehicle-based road dust emission measurement-Part II: Effect of precipitation, wintertime road sanding, and street sweepers on inferred PM<sub>10</sub> emission potentials from paved and unpaved roads, *Atmos. Environ.*, 37, 4573-4582, doi: 10.1016/S1352-2310(03)00529-6, 2003.

Leung, A. O. W., Duzgoren-Aydin, N. S., Cheung, K. C., and Wong, M. H.: Heavy metals concentrations of surface dust from e-waste recycling and its human health

- implications in Southeast China, *Environ. Sci. Technol.*, 42, 2674-2680, doi: 10.1021/es071873x, 2008.
- Li, H., Wang, J., Wang, Q., Tian, C., Qian, X., and Leng, X.: Magnetic properties as a proxy for predicting fine-particle-bound heavy metals in a support vector machine approach, *Environ. Sci. Technol.*, *Environ. Sci. Technol.*, 51, 6927-6935, doi: 10.1021/acs.est.7b00729, 2017.
- Lin, Y. C., Tsai, C. J., Wu, Y. C., Zhang, R., Chi, K. H., Huang, Y. T., Lin, S. H., and Hsu, S. C.: Characteristics of trace metals in traffic-derived particles in Hsuehshan Tunnel, Taiwan: size distribution, potential source, and fingerprinting metal ratio, *Atmos. Chem. Phys.*, 15, 4117-4130, doi: 10.5194/acp-15-4117-2015, 2015.
- Litter, M. I.: Heterogeneous photocatalysis: Transition metal ions in photocatalytic systems, *Appl. Catal. B: Environ.*, 23, 89-114, doi: 10.1016/S0926-3373(99)00069-7, 1999.
- Liu, Z., Hu, B., Wang, L., Wu, F., Gao, W., and Wang, Y.: Seasonal and diurnal variation in particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>) at an urban site of Beijing: Analyses from a 9-year study, *Environ. Sci. Pollut. Res. Int.*, 22, 627-642, doi: 10.1007/s11356-014-3347-0, 2015.
- Liu, Z., Lu, X., Feng, J., Fan, Q., Zhang, Y., and Yang, X.: Influence of ship emissions on urban air quality: A comprehensive study using highly time-resolved online measurements and numerical simulation in Shanghai, *Environ. Sci. Technol.*, 51, 202-211, doi: 10.1021/acs.est.6b03834, 2017.
- Liu, S., Triantis, K., and Zhang, L.: The design of an urban roadside automatic sprinkling system: mitigation of PM<sub>2.5-10</sub> in ambient air in megacities, *Chinese J. Engineering*, 12, doi: 10.1155/2014/618109, 2014.
- Lu, S., Yao, Z., Chen, X., Wu, M., Sheng, G., Fu, J., and Paul, D.: The relationship between physicochemical characterization and the potential toxicity of fine particulates (PM<sub>2.5</sub>) in Shanghai atmosphere, *Atmos. Environ.*, 42, 7205-7214, doi: 10.1016/j.atmosenv.2008.07.030, 2008.
- Lough, G. C., Schauer, J. J., Park, J. S., Shafer, M. M., DeMinter, J. T., and Weinstein, J. P.: Emissions of metals associated with motor vehicle roadways, *Environ. Sci. Technol.*, 39, 826-836, doi: 10.1021/es048715f, 2005.
- Maenhaut, W.: Present role of PIXE in atmospheric aerosol research, *Nucl. Instrum. Meth. B*, 363, 86-91, doi: 10.1016/j.nimb.2015.07.043, 2015.
- Morman, S. A. and Plumlee, G. S.: The role of airborne mineral dusts in human disease, *Aeolian Res.*, 9, 203-212, doi: 10.1016/j.aeolia.2012.12.001, 2013.
- Murphy, D. M., Thomson, D. S., and Mahoney, M. J.: In situ measurements of organics, meteoritic material, mercury, and other elements in aerosols at 5 to 19 kilometers, *Science*, 282, 1664-1669, doi: 10.1126/science.282.5394.1664, 1998.
- Normile, D.: China's living laboratory in urbanization, *Science*, 319, 740-743, doi: 10.1126/science.319.5864.740, 2008.
- Olujimi, O. O., Oputu, O., Fatoki, O., Opatoyinbo, O. E., Aroyewun, O. A., and Baruani, J.: Heavy metals speciation and human health risk assessment at an illegal gold mining site in Igun, Osun State, Nigeria, *J. Heal. Pollut.*, 5, 19-32, doi: 10.5696/i2156-9614-5-8.19, 2015.

- 950 Pacyna, J. M. and Pacyna, E. G.: An assessment of global and regional emissions of trace metals to the atmosphere from anthropogenic sources worldwide, *Environ. Rev.*, 9, 269-298, doi: 10.1139/a01-012, 2001.
- Pardo, M., Shafer, M. M., Rudich, A., Schauer, J. J., and Rudich, Y.: Single exposure to near roadway particulate matter leads to confined inflammatory and defense responses: Possible role of metals, *Environ. Sci. Technol.*, 49, 8777-8785, doi: 955 10.1021/acs.est.5b01449, 2015.
- Park, S. S., Cho, S. Y., Jo, M. R., Gong, B. J., Park, J. S., and Lee, S. J.: Field evaluation of a near-real time elemental monitor and identification of element sources observed at an air monitoring supersite in Korea, *Atmos. Pollut. Res.*, 5, 119-128, 960 doi: 10.5094/apr.2014.015, 2014.
- Perry, K. D., Cahill, T. A., Schnell, R. C., and Harris, J. M.: Long-range transport of anthropogenic aerosols to the National Oceanic and Atmospheric Administration baseline station at Mauna Loa Observatory, Hawaii, *J. Geophys. Res.*, 104, 18521-18533, doi: 10.1029/1998JD100083, 1999.
- 965 Phillips-Smith, C., Jeong, C. H., Healy, R. M., Dabek-Zlotorzynska, E., Celo, V., Brook, J. R., and Evans, G.: Sources of particulate matter in the Athabasca oil sands region: Investigation through a comparison of trace element measurement methodologies, *Atmos. Chem. Phys. Discuss.*, 2017, 1-34, doi: 10.5194/acp-2016-966, 2017.
- Pope III, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and 970 Thurston, G. D.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *J. Am. Med. Assoc.*, 1132-1141, 2002.
- Pope III, C. A., Ezzati, M., and Dockery, D. W.: Fine-particulate air pollution and life expectancy in the United States, *New Engl. J. Med.*, 360, 376-386, doi: 10.1056/NEJMsa0805646, 2009.
- 975 Qiao, L., Cai, J., Wang, H., Wang, W., Zhou, M., Lou, S., Chen, R., Dai, H., Chen, C., and Kan, H.: PM<sub>2.5</sub> constituents and hospital emergency-room visits in Shanghai, China, *Environ. Sci. Technol.*, 48, 10406-10414, doi: 10.1021/es501305k, 2014.
- Querol, X., Juan, R., Lopez-Soler, A., Fernandez-Turiel, J., and Ruiz, C. R.: Mobility of trace elements from coal and combustion wastes, *Fuel*, 75, 821-838, doi: 980 10.1016/0016-2361(96)00027-0, 1996.
- Richard, A., Bukowiecki, N., Lienemann, P., Furger, M., Fierz, M., Minguillón, M. C., Weideli, B., Figi, R., Flechsig, U., Appel, K., Prévôt, A. S. H., and Baltensperger, U.: Quantitative sampling and analysis of trace elements in atmospheric aerosols: impactor characterization and Synchrotron-XRF mass calibration, *Atmos. Meas. 985 Tech.*, 3, 1473-1485, doi: 10.5194/amt-3-1473-2010, 2010.
- Ridley, D. A., Heald, C. L., Kok, J. F., and Zhao, C.: An observationally constrained estimate of global dust aerosol optical depth, *Atmos. Chem. Phys.*, 16, 15097-15117, doi: 10.5194/acp-16-15097-2016, 2016.
- Rubasinghege, G., Elzey, S., Baltrusaitis, J., Jayaweera, P. M., and Grassian, V. H.: 990 Reactions on atmospheric dust particles: Surface photochemistry and size-dependent nanoscale redox chemistry, *J. Phys. Chem. Lett.*, 1, 1729-1737, doi: 10.1021/jz100371d, 2010a.
- Rubasinghege, G., Lentz, R. W., Scherer, M. M., and Grassian, V. H.: Simulated

- atmospheric processing of iron oxyhydroxide minerals at low pH: Roles of particle size and acid anion in iron dissolution, *P. Natl. Acad. Sci.*, 107, 6628-6633, doi: 10.1073/pnas.0910809107, 2010b.
- 995 Saffari, A., Daher, N., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Global perspective on the oxidative potential of airborne particulate matter: A synthesis of research findings, *Environ. Sci. Technol.*, 48, 7576-7583, doi: 10.1021/es500937x, 2014.
- 1000 Seigneur, C. and Constantinou, E.: Chemical kinetic mechanism for atmospheric chromium, *Environ. Sci. Technol.*, 29, 222-231, doi: 10.1021/es00001a029, 1995.
- Shafer, M. M., Toner, B. M., Overdier, J. T., Schauer, J. J., Fakra, S. C., Hu, S., Herner, J. D., and Ayala, A.: Chemical speciation of vanadium in particulate matter emitted from diesel vehicles and urban atmospheric aerosols, *Environ. Sci. Technol.*, 46, 189-195, doi: 10.1021/es200463c, 2012.
- 1005 Shah, A. S. V., Langrish, J. P., Nair, H., McAllister, D. A., Hunter, A. L., Donaldson, K., Newby, D. E., and Mills, N. L.: Global association of air pollution and heart failure: A systematic review and meta-analysis, *Lancet*, 382, 1039-1048, doi: 10.1016/s0140-6736(13)60898-3, 2013.
- 1010 Shu, J., Dearing, J. A., Morse, A. P., Yu, L., and Yuan, N.: Determining the sources of atmospheric particles in Shanghai, China, from magnetic and geochemical properties, *Atmos. Environ.*, 35, 2615-2625, doi: 10.1016/S1352-2310(00)00454-4, 2001.
- Sofowote, U. M., Su, Y., Dabek-Zlotorzynska, E., Rastogi, A. K., Brook, J., and Hopke, P. K.: Sources and temporal variations of constrained PMF factors obtained from multiple-year receptor modeling of ambient PM<sub>2.5</sub> data from five speciation sites in Ontario, Canada, *Atmos. Environ.*, 108, 140-150, doi: 10.1016/j.atmosenv.2015.02.055, 2015.
- 1015 Strak, M., Janssen, N. A., Godri, K. J., Gosens, I., Mudway, I. S., Cassee, F. R., Lebre, E., Kelly, F. J., Harrison, R. M., Brunekreef, B., Steenhof, M., and Hoek, G.: Respiratory health effects of airborne particulate matter: The role of particle size, composition, and oxidative potential-the RAPTES project, *Environ. Health. Perspect.*, 120, 1183-1189, doi: 10.1289/ehp.1104389, 2012.
- 1020 Streit, B.: *Lexikon der Okotoxikologie*, Wiley-VCH, Weinheim, Germany, 1991.
- 1025 Strickland, M. J., Hao, H., Hu, X., Chang, H. H., Darrow, L. A., and Liu, Y.: Pediatric emergency visits and short-term changes in PM<sub>2.5</sub> concentrations in the U.S. State of Georgia, *Environ. Health. Perspect.*, 124, 690-696, doi: 10.1289/ehp.1509856, 2016.
- Suess, D. T.: Single particle mass spectrometry combustion source characterization and atmospheric apportionment of vehicular, coal and biofuel exhaust emissions, PhD, Chemistry, University of California, Riverside, CA, USA, 2002.
- 1030 Tang, M., Huang, X., Lu, K., Ge, M., Li, Y., Cheng, P., Zhu, T., Ding, A., Zhang, Y., Gligorovski, S., Song, W., Ding, X., Bi, X., and Wang, X.: Heterogeneous reactions of mineral dust aerosol: Implications for tropospheric oxidation capacity, *Atmos. Chem. Phys. Discuss.*, 2017, 1-124, doi: 10.5194/acp-2017-458, 2017.
- 1035 Tao, L., Fairley, D., Kleeman, M. J., and Harley, R. A.: Effects of switching to lower sulfur marine fuel oil on air quality in the San Francisco Bay area, *Environ. Sci.*

- Technol., 47, doi: 10171-10178, 10.1021/es401049x, 2013.
- 1040 Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., and Sutton, D. J.: Heavy metals toxicity and the environment. In *Molecular, Clinical and Environmental Toxicology*, 133-164, ISBN: 978-3-7643-8337-4, 2012.
- 1045 Tian, H. Z., Zhu, C. Y., Gao, J. J., Cheng, K., Hao, J. M., Wang, K., Hua, S. B., Wang, Y., and Zhou, J. R.: Quantitative assessment of atmospheric emissions of toxic heavy metals from anthropogenic sources in China: Historical trend, spatial distribution, uncertainties, and control policies, *Atmos. Chem. Phys.*, 15, 10127-10147, doi: 10.5194/acp-15-10127-2015, 2015.
- 1050 Traversi, R., Becagli, S., Calzolari, G., Chiari, M., Giannoni, M., Lucarelli, F., Nava, S., Rugi, F., Severi, M., and Udisti, R.: A comparison between PIXE and ICP-AES measurements of metals in aerosol particulate collected in urban and marine sites in Italy, *Nucl. Instrum. Meth. B*, 318, 130-134, doi: 10.1016/j.nimb.2013.05.102, 2014.
- Usher, C. R., Michel, A. E., and Grassian, V. H.: Reactions on mineral dust, *Chem. Rev.*, 103, 4883-4940, doi: 10.1021/cr020657y, 2003.
- 1055 Verma, V., Shafer, M. M., Schauer, J. J., and Sioutas, C.: Contribution of transition metals in the reactive oxygen species activity of PM emissions from retrofitted heavy-duty vehicles, *Atmos. Environ.*, 44, 5165-5173, doi: 10.1016/j.atmosenv.2010.08.052, 2010.
- 1060 Viana, M., Amato, F., Alastuey, A., Querol, X., Moreno, T., García Dos Santos, S., Hecce, M. D., and Fernández-Patier, R.: Chemical tracers of particulate emissions from commercial shipping, *Environ. Sci. Technol.*, 43, 7472-7477, 10.1021/es901558t, 2009.
- 1065 Visser, S., Slowik, J. G., Furger, M., Zotter, P., Bukowiecki, N., Canonaco, F., Flechsig, U., Appel, K., Green, D. C., Tremper, A. H., Young, D. E., Williams, P. I., Allan, J. D., Coe, H., Williams, L. R., Mohr, C., Xu, L., Ng, N. L., Nemitz, E., Barlow, J. F., Halios, C. H., Fleming, Z. L., Baltensperger, U., and Prévôt, A. S. H.: Advanced source apportionment of size-resolved trace elements at multiple sites in London during winter, *Atmos. Chem. Phys.*, 15, 11291-11309, doi: 10.5194/acp-15-11291-2015, 2015a.
- 1070 Visser, S., Slowik, J. G., Furger, M., Zotter, P., Bukowiecki, N., Dressler, R., Flechsig, U., Appel, K., Green, D. C., Tremper, A. H., Young, D. E., Williams, P. I., Allan, J. D., Herndon, S. C., Williams, L. R., Mohr, C., Xu, L., Ng, N. L., Detournay, A., Barlow, J. F., Halios, C. H., Fleming, Z. L., Baltensperger, U., and Prévôt, A. S. H.: Kerb and urban increment of highly time-resolved trace elements in PM<sub>10</sub>, PM<sub>2.5</sub> and PM<sub>10</sub> winter aerosol in London during ClearfLo 2012, *Atmos. Chem. Phys.*, 15, 2367-2386, doi: 10.5194/acp-15-2367-2015, 2015b.
- 1075 Wang, F., Chen, Y., Meng, X., Fu, J., and Wang, B.: The contribution of anthropogenic sources to the aerosols over East China Sea, *Atmos. Environ.*, 127, 22-33, doi: 10.1016/j.atmosenv.2015.12.002, 2016.
- 1080 Wang, J., Hu, Z., Chen, Y., Chen, Z., and Xu, S.: Contamination characteristics and possible sources of PM<sub>10</sub> and PM<sub>2.5</sub> in different functional areas of Shanghai, China, *Atmos. Environ.*, 68, 221-229, doi: 10.1016/j.atmosenv.2012.10.070,

2013a.

Wang, X., Bi, X., Sheng, G., and Fu, J.: Hospital indoor PM<sub>10</sub>/PM<sub>2.5</sub> and associated trace elements in Guangzhou, China, *Sci. Total Environ.*, 366, 124-135, doi: 10.1016/j.scitotenv.2005.09.004, 2006.

Wang, X., Williams, B. J., Wang, X., Tang, Y., Huang, Y., Kong, L., Yang, X., and Biswas, P.: Characterization of organic aerosol produced during pulverized coal combustion in a drop tube furnace, *Atmos. Chem. Phys.*, 13, 10919-10932, doi: 10.5194/acp-13-10919-2013, 2013b.

West, J. J., Cohen, A., Dentener, F., Brunekreef, B., Zhu, T., Armstrong, B., Bell, M. L., Brauer, M., Carmichael, G., Costa, D. L., Dockery, D. W., Kleeman, M., Krzyzanowski, M., Kunzli, N., Liousse, C., Lung, S. C., Martin, R. V., Poschl, U., Pope, C. A., 3rd, Roberts, J. M., Russell, A. G., and Wiedinmyer, C.: "What we breathe impacts our health: Improving understanding of the link between air pollution and health", *Environ. Sci. Technol.*, 50, 4895-4904, doi: 10.1021/acs.est.5b03827, 2016.

WHO (World Health Organization) Air quality guidelines - global update 2005, available online at [http://www.who.int/phe/health\\_topics/outdoorair/outdoorair\\_aqg/en/](http://www.who.int/phe/health_topics/outdoorair/outdoorair_aqg/en/), 2005. (last accessible: 7/7/2017)

Wu, Q. R., Wang, S. X., Zhang, L., Song, J. X., Yang, H., and Meng, Y.: Update of mercury emissions from China's primary zinc, lead and copper smelters, 2000-2010, *Atmos. Chem. Phys.*, 12, 11153-11163, doi: 10.5194/acp-12-11153-2012, 2012.

Yanca, C. A., Barth, D. C., Petterson, K. A., Nakanishi, M. P., Cooper, J. A., Johnsen, B. E., Lambert, R. H., and Bivins, D. G.: Validation of three new methods for determination of metal emissions using a modified Environmental Protection Agency Method 301, *J. Air Waste Manage. Assoc.*, 56, 1733-1742, doi: 10.1080/10473289.2006.10464578, 2006.

Yu, S.: Water spray geoengineering to clean air pollution for mitigating haze in China's cities, *Environ. Chem. Lett.*, 12, 109-116, doi: 10.1007/s10311-013-0444-0, 2014.

Zhang, X., Hecobian, A., Zheng, M., Frank, N. H., and Weber, R. J.: Biomass burning impact on PM<sub>2.5</sub> over the southeastern US during 2007: integrating chemically speciated FRM filter measurements, MODIS fire counts and PMF analysis, *Atmos. Chem. Phys.*, 10, 6839-6853, doi: 10.5194/acp-10-6839-2010, 2010.

Zhao, J., Lewinski, N., and Riediker, M.: Physico-chemical characterization and oxidative reactivity evaluation of aged brake wear particles, *Aerosol Sci. Technol.*, 49, 65-74, doi: 10.1080/02786826.2014.998363, 2015.

Zhao, M., Zhang, Y., Ma, W., Fu, Q., Yang, X., Li, C., Zhou, B., Yu, Q., and Chen, L.: Characteristics and ship traffic source identification of air pollutants in China's largest port, *Atmos. Environ.*, 64, 277-286, doi: 10.1016/j.atmosenv.2012.10.007, 2013.

Zhen, L., Li, M., Hu, Z., Lv, W., and Zhao, X.: The effects of emission control area regulations on cruise shipping, *Transport. Res. D-Tr. E.*, 62, 47-63, doi: 10.1016/j.trd.2018.02.005, 2018.

Zheng, J., Tan, M., Shibata, Y., Tanaka, A., Li, Y., Zhang, G., Zhang, Y., and Shan, Z.: Characteristics of lead isotope ratios and elemental concentrations in PM<sub>10</sub> fraction of airborne particulate matter in Shanghai after the phase-out of leaded gasoline, *Atmos. Environ.*, 38, 1191-1200, doi: 10.1016/j.atmosenv.2003.11.004, 2004.