

Coupling of organic and inorganic aerosol systems and the effect on gas-particle partitioning in the southeastern United States

Havala O. T. Pye¹, Andreas Zuend², Juliane L. Fry³, Gabriel Isaacman-VanWertz^{4,5}, Shannon L. Capps⁶, K. Wyatt Appel¹, Hosein Foroutan¹, Lu Xu⁷, Nga L. Ng^{8,9}, and Allen H. Goldstein^{5,10}

¹National Exposure Research Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, USA

²Department of Atmospheric and Oceanic Sciences, McGill University, Montreal, Québec, CAN

³Department of Chemistry, Reed College, Portland, Oregon, USA

⁴Department of Civil and Environmental Engineering, Virginia Polytechnic Institute and State University, Blacksburg, Virginia, USA

⁵Department of Environmental Science, Policy, and Management, University of California, Berkeley, California, USA

⁶Civil, Architectural, and Environmental Engineering, Drexel University, Philadelphia, Pennsylvania, USA

⁷Department of Environmental Science and Engineering, California Institute of Technology, Pasadena, California, USA

⁸School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA, USA

⁹School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA

¹⁰Department of Civil and Environmental Engineering, University of California, Berkeley, California, USA

Correspondence to: Havala O. T. Pye (pye.havala@epa.gov)

Abstract. Several models were used to describe the partitioning of ammonia, water, and organic compounds between the gas and particle phase for conditions in the southeastern United States during summer 2013. Existing equilibrium models and frameworks were found to be sufficient although additional improvements in terms of estimating pure-species vapor pressures are needed. Thermodynamic model predictions were consistent, to first order, with a molar ratio of ammonium to sulfate of approximately 1.6 to 1.8 (Ratio of ammonium to $2 \times$ sulfate, $R_{N/2S} \approx 0.8$ to 0.9) with approximately 70% of total ammonia and ammonium (NH_x) in the particle. Southeastern Aerosol Research and Characterization (SEARCH) network gas and aerosol and Southern Oxidant and Aerosol Study (SOAS) Monitor for Aerosols and Gases in Air (MARGA) aerosol measurements were consistent with these conditions. CMAQv5.2 regional chemical transport model predictions did not reflect these conditions due to a factor of three overestimate of the nonvolatile cations. In addition, gas-phase ammonia was overestimated in the CMAQ model leading to an even lower fraction of total ammonia in the particle. Chemical Speciation Network (CSN) and Aerosol Mass Spectrometer (AMS) measurements indicated less ammonium per sulfate than SEARCH and MARGA measurements and were inconsistent with thermodynamic model predictions. Organic compounds were predicted to be present to some extent in the same phase as inorganic constituents, modifying their activity and resulting in a decrease in $[\text{H}^+]_{\text{air}}$ (H^+ in $\mu\text{g m}^{-3}$ air), increase in ammonia partitioning to the gas phase, and increase in pH compared to complete organic vs. inorganic liquid-liquid phase separation. In addition, accounting for non-ideal mixing modified the pH such that a fully interactive inorganic-organic system had a pH roughly 0.7 units higher than predicted by traditional methods (pH= 1.5 vs 0.7). Particle-phase interactions of organic and inorganic compounds were found to increase partitioning towards the particle phase (vs. gas phase) for highly

oxygenated ($O:C \geq 0.6$) compounds including several isoprene-derived tracers as well as levoglucosan, but decrease particle-phase partitioning for low O:C, monoterpene-derived species.

1 Introduction

Ambient particles consist of organic and inorganic compounds. The organic compounds present in the gas and particle phase are diverse and numerous (Goldstein and Galbally, 2007), ranging from relatively unoxidized, long-chain alkanes in fresh emissions to small, highly soluble compounds formed through multiple generations of atmospheric chemistry. Major inorganic constituents include water, sulfate, ammonium, and nitrate with additional contributions from species such as calcium, potassium, magnesium, sodium, and chloride (Reff et al., 2009). The extent to which organic and inorganic components of particulate matter interact within a particle depends on the mixing state (e.g. internal vs external) of the aerosol population as well as degree of phase separation (or number of phases) within the particle. Internally mixed populations, as typically assumed in chemical transport models such as the Community Multiscale Air Quality (CMAQ) model, may exhibit one fairly homogeneous liquid phase state or be heterogeneous in composition. Heterogeneous configurations occur as a result of phase separation and may include a liquid and solid phase or multiple liquid phases. A common heterogeneous configuration under conditions of liquid-liquid or solid-liquid phase separation is that of a core-shell morphology; alternatively, partially engulfed morphologies have been predicted by theory and observed in laboratory experiments (Kwamena et al., 2010; Song et al., 2013; Reid et al., 2011).

Currently, the CMAQ model, as well as other chemical transport models, considers accumulation mode aerosol to form a heterogeneous internal mixture in which organic and inorganic constituents partition between the gas and aerosol phase independently of each other. Pye et al. (2017) examined how assumptions about phase separation of internally mixed particles affect organic aerosol concentrations in the southeast United States as predicted by the CMAQ model. When organic compounds were allowed to mix with the aqueous inorganic phase under conditions of high relative humidity and high degree of oxygenation (You et al., 2013; Bertram et al., 2011; Song et al., 2012), the concentration of organic aerosol was predicted to increase significantly (Pye et al., 2017). While the effects of phase separation on organic compounds are potentially large, they are highly dependent on an accurate parameterization of activity coefficients and a reliable prediction of the composition of individual particle phases (Zuend and Seinfeld, 2012).

Recent work highlights potential discrepancies between current gas-particle partitioning models, which assume equilibrium is attained on short timescales, and observations for both inorganic and organic compounds. Silvern et al. (2017) found that models predict higher ratios of particulate ammonium to sulfate than observed in the eastern U.S. and proposed that organic compounds in an organic-rich phase at the particle surface may reduce ammonia partitioning to the particle via a kinetic inhibition. In addition, organic compound vapor pressure estimation method predictions can vary by orders of magnitude (Topping et al., 2016; O'Meara et al., 2014; Pankow and Asher, 2008) and have often been adjusted downward to improve model predictions (Chan et al., 2009; Johnson et al., 2006). Furthermore, isoprene-epoxydiol-derived organic aerosol partitions to the particle phase to a greater degree than structure-based vapor pressures would suggest (Isaacman-VanWertz et al., 2016;

Lopez-Hilfiker et al., 2016; Hu et al., 2016). Since $PM_{2.5}$ (particulate matter concentration from particles of diameters less than $2.5 \mu m$) is regulated via the National Ambient Air Quality Standards (NAAQS) in the U.S., while similar ambient standards are not set for the gas-phase counterparts (NH_3 and organic compound vapors), errors in partitioning will affect model performance with implications for metrics used in regulatory applications. The model sensitivity of $PM_{2.5}$ to emission changes can also be
5 too high or too low if compounds are erroneously partitioned.

In this work, gas-particle partitioning of ammonia and several isoprene-, monoterpene-, and biomass burning-derived organic compounds was examined using common air quality modeling treatments and advanced approaches. Results address the degree to which techniques accounting for organic-inorganic interactions, deviations in ideality, and phase separation reproduce observations. Models were evaluated for their ability to predict ammonia versus ammonium as well as gas-particle partitioning
10 of organic compounds. In addition, the effects of organic compounds on aerosol pH were examined.

2 Methods

2.1 Model Approaches

Several box-model approaches as well as CMAQ regional chemical transport model calculations were used to represent the partitioning of compounds between the gas and particle phases. CMAQ version 5.2-gamma was run over the continental U.S.
15 at 12 km by 12 km horizontal resolution for 1 June – 15 July 2013, coinciding with the Southern Oxidant and Aerosol Study (SOAS) field campaign and the Centreville, Alabama, U.S. field site. WRF v3.8 meteorology with lightning assimilated into the convection scheme (Heath et al., 2016) was processed for use with the CMAQ model (Otte and Pleim, 2010). Emissions were based on the 2011 National Emission Inventory version 2 (ek). Emissions influenced by model meteorology (biogenic compounds, mobile sector) or monitored (electrical generation units) were year 2013 specific. Windblown dust emissions
20 followed the scheme of Foroutan et al. (2017). Ammonia emissions and deposition from croplands were parameterized as a bidirectional exchange (Pleim et al., 2013). CMAQ used ISORROPIA v2.1 (Fountoukis and Nenes, 2007; Nenes et al., 1998) with gas and aerosol composition and environmental conditions (temperature, relative humidity) as input to predict the Aitken and Accumulation mode ammonium, nitrate, and chloride mass concentrations. CMAQ predicted PM_1 and $PM_{2.5}$ were computed based on the fraction of the Aitken and Accumulation modes less than 1 or 2.5 microns in diameter as appropriate
25 (Nolte et al., 2015).

Consistent with the CMAQ regional model, partitioning of ammonia between the gas and particle phases was also predicted using ISORROPIA as a box model driven with observed aerosol (reverse mode) or gas and aerosol (forward mode) concentrations of ammonia, ammonium, nitrate, nitric acid, calcium, potassium, magnesium, sodium, and chloride. Output from the ISORROPIA box-model was either gas-phase ammonia in equilibrium with the observed aerosol ammonium (reverse mode),
30 or ammonia vs. ammonium based on total gas and aerosol conditions (forward mode). ISORROPIA does not consider the effects of organic compounds on aerosol pH or explicitly treat liquid-liquid phase separation.

Algorithms that allowed for inorganic-organic interactions were applied using a thermodynamic equilibrium gas-particle partitioning model (Zuend et al., 2010; Zuend and Seinfeld, 2012) based on the Aerosol Inorganic-Organic Mixtures Func-

tional groups Activity Coefficients (AIOMFAC) model (Zuend et al., 2008, 2011). AIOMFAC provided an estimate of activity coefficients for aerosol systems of specified functional group composition, which was used in two modes: (i) predefined complete liquid-liquid phase separation (CLLPS) in which the organic compounds did not mix with the inorganic salts and (ii) equilibrium (EQLB) in which the Gibbs energy of the system was minimized and up to two liquid phases of any composition were allowed to form in the particle as predicted by a modified liquid-liquid phase separation algorithm based on the method by Zuend and Seinfeld (2013). For purposes of AIOMFAC calculations, observed calcium, potassium, and magnesium concentrations were converted to charge-equivalent sodium amounts since the former's interactions with the bisulfate ion in solution are not treated by the model.

Several quantities, including pH and molar ratios, were calculated to evaluate the inorganic aerosol system. Solution acidity can be expressed in different ways, the most common one being the pH value. However, many definitions of pH exist, with several definitions only applicable to highly dilute aqueous solutions. Thermodynamics-based pH definitions vary with the choice of composition scale (molality, molarity, or mole fraction basis) and the solvent into which H^+ is assumed to dissolve, which may be strictly water associated with inorganic constituents as in ISORROPIA II, or include the diluting effect of water associated with organic compounds (Guo et al., 2015), organic compounds themselves (Zuend et al., 2008), or other aerosol constituents (Budisulistiorini et al., 2017). Furthermore activity coefficients of H^+ may not be unity as is frequently assumed. In this work, pH was defined following the thermodynamic definition on a molality basis, as recommended by IUPAC (<http://goldbook.iupac.org/html/P/P04524.html>) and computed by the AIOMFAC model. By expressing the molality of H^+ in terms of concentration per volume of air, the following results:

$$pH = -\log_{10}(\gamma_{H^+} [H^+]_{\text{air}}/[S]) \quad (1)$$

where γ_{H^+} is the molality based activity coefficient for H^+ in the liquid phase, $[H^+]_{\text{air}}$ is the concentration of the hydronium ion in the liquid phase in moles per volume of air and $[S]$ is the solvent mass in that liquid phase (kg per volume of air), i.e. $[H^+]_{\text{air}}/[S]$ is the molality of H^+ . The solvent included water associated with inorganic compounds (W_i), water associated with organic compounds (W_o), and organic compounds (C_{org}) as appropriate based on the predicted phase composition. ISORROPIA pH calculations assumed $[S] = [W_i]$ and an activity coefficient of unity thus following previous methods (Guo et al., 2017a). The molar ratio of ammonium to 2 × sulfate was defined as:

$$R_{N/2S} = \frac{n_{\text{NH}_4^+}}{2 \times n_{\text{SO}_4^{2-}}}, \quad (2)$$

and the electric charge normalized molar ratio of cations to anions that participate in ISORROPIA was:

$$R_{+/-} = \frac{n_{\text{NH}_4^+} + n_{\text{Na}^+} + 2 \times n_{\text{Ca}^{2+}} + n_{\text{K}^+} + 2 \times n_{\text{Mg}^{2+}}}{2 \times n_{\text{SO}_4^{2-}} + n_{\text{NO}_3^-} + n_{\text{Cl}^-}}. \quad (3)$$

Since ambient measurements and CMAQ model output do not distinguish bisulfate from sulfate, the sulfate in these ratios represented total sulfate ($\text{SO}_4^{2-} + \text{HSO}_4^-$).

To employ the AIOMFAC-based equilibrium models, organic aerosol positive matrix factorization (PMF) analysis results of ambient data (next section) were converted to molecular structures of known functional group composition as surrogates for a

range of organic compound classes in ambient particles as described in Tables S1-S3 thus providing a complete characterization of the organic aerosol partitioning medium. Several isoprene-derived (2-methyltetrols, C₅-alkene triols, 2-methylglyceric acid) and monoterpene-derived (pinic acid, pinonic acid, hydroxyglutaric acid) compounds as well as levoglucosan, a semivolatile indicator of biomass burning, were explicitly represented in box-model calculations. Pure species' vapor pressures (sub-cooled liquid) were obtained via the EVAPORATION model (Compernelle et al., 2011). The temperature dependence was parameterized by using the same Antoine-like function that is also employed by the EVAPORATION model. A sensitivity calculation (referred to as Adj Psat) reduced EVAPORATION-based vapor pressures by a factor of 4.2, thus maintaining the compound to compound variability predicted by EVAPORATION but correcting for potential overestimates in pure compound vapor pressures. The magnitude of the adjustment was based on the effective saturation concentration obtained via regression needed to reproduce observations in a traditional absorptive partitioning framework (Equation S1). This adjustment factor is similar in magnitude to the difference between SIMPOL (Pankow and Asher, 2008) and EVAPORATION (Compernelle et al., 2011) predicted vapor pressures for several species, but not all (see Table S4). The effective saturation concentration, C*, of a species, i, was defined as (Zuend and Seinfeld, 2012):

$$C_i^* = \frac{C_i^g \sum C_k^{PM}}{C_i^{PM}} \quad (4)$$

where C_i^g is the mass-based gas-phase concentration of species i , C_i^{PM} is the mass-based liquid-phase concentration of species i , and C_k^{PM} is the total mass-based concentration of the liquid phase where the summation index k includes organic species, inorganic species, and water. See Equation S2 for C_i^* in terms of the mole-fraction based activity coefficient.

2.2 Ambient Data

Regional model predictions of inorganic aerosol were evaluated against the Chemical Speciation Network (CSN) and Southeastern Aerosol Research and Characterization (SEARCH) network observations (at different ground sites). The Interagency Monitoring of Protected Visual Environments (IMPROVE) network (Solomon et al., 2014) also measures some chemical speciation of PM_{2.5} throughout the U.S., but does not include ammonium. CSN determines anions and cations via ion chromatography of extracts from nylon filters (Solomon et al., 2014). Solomon et al. (2014) estimate the precision of CSN measured ammonium is 11% and sulfate is 7% (for co-located samples during 2012) but the actual measurement uncertainty is likely higher (and not quantified). The SEARCH network operates at fewer sites and exclusively in the Southeast U.S. It uses a multi-channel approach employing nylon, teflon, and citric acid-coated cellulose filters to measure speciated 24-hour average PM_{2.5} (Edgerton et al., 2005). SEARCH reports the precision of measured sulfate and ammonium in PM_{2.5} is 2-3% (Edgerton et al., 2005). The SEARCH 24-hour filter measurements are also used to adjust the co-located continuous measurements (Edgerton et al., 2006).

In addition to the network data, ambient data from SOAS at the Centreville, AL (CTR; 87.25° W, 32.90° N) site from June and July, 2013 were used as input to the box models and for model evaluation. The High Resolution Time of Flight Aerosol Mass Spectrometer (HR-ToF-AMS, hereafter AMS) operated by the Georgia Institute of Technology was the primary source of SOAS PM₁ organic mass, ammonium, and sulfate (Xu et al., 2015a). This AMS dataset was consistent with the other

AMS instrument operating at CTR as well as AMS measurements aboard an aircraft over the southeastern U.S. (Table S5-S6). When AMS data was used as input in PM₁ box modeling, inorganic nitrate was set to zero as nitrate measured by the AMS contained significant contributions from organic nitrogen-containing compounds (Xu et al., 2015b). Thus, AMS calculations assumed the inorganic aerosol was composed only of ammonium, sulfate, bisulfate, and the hydronium ion (referred to in subsequent sections as the A⁺ system). The assignment of measured ammonium and sulfate to specific salts (ammonium sulfate vs ammonium bisulfate) for use as input electrolyte components to AIOMFAC was determined by mass balance. Inorganic PM_{2.5}, including ammonium, sulfate, nitrate, calcium, potassium, magnesium, sodium, and chloride, was measured at CTR by a Monitor for Aerosols and Gases in Air (MARGA) (Allen et al., 2015). Less than 5% of the PM_{2.5} MARGA data used in this work had elevated nitrate (>0.8 μg m⁻³) due to supermicron crustal material and sea salt episodes (Allen et al., 2015). The data from MARGA was used in two ways for model calculations with AIOMFAC. (1) All the measured ion concentrations were considered, but the molar amounts of the cations Ca²⁺, K⁺, Mg²⁺, and Na⁺ were mapped to a charge-equivalent amount of Na⁺ (see Section 2.1). (2) Only the measured concentrations of ammonium and sulfate ions were considered and mapped to the electrolyte components ammonium sulfate, ammonium bisulfate, and sulfuric acid for AIOMFAC model input purposes. For ammonium-sulfate only conditions (option 2) in which the moles NH₄⁺ ≥ 2×SO₄²⁻, a small amount (1 × 10⁻⁴ μg m⁻³) of ammonium bisulfate was added to the AIOMFAC input for MARGA calculations in order to trigger a potential partial association of sulfate and H⁺ ions to bisulfate following the equilibrium constant of that reaction. Such conditions did not occur with AMS data. Hourly gas-phase ammonia was obtained from the CTR SEARCH network site via a corrected Thermo Scientific citric acid-impregnated denuder. Relative humidity (RH) and temperature were obtained from the routine SEARCH network measurements at the SOAS site.

The entire organic aerosol composition was characterized in terms of functional groups for use with AIOMFAC. The semi-volatile thermal desorption aerosol gas chromatograph (SV-TAG) with in situ derivatization (Isaacman-VanWertz et al., 2016; Isaacman et al., 2014) provided measured gas- and aerosol-phase concentrations of 2-methyltetrols, C₅-alkene triols, 2-methylglyceric acid, pinic acid, pinonic acid, hydroxyglutaric acid, and levoglucosan. More oxidized-oxygenated organic aerosol (MO-OOA), biomass burning organic aerosol (BBOA), Isoprene-OA, and less oxidized-oxygenated organic aerosol (LO-OOA) PMF factors from the AMS were represented with specific functional groups and associated surrogate chemical structures (Table S1). As previous work indicates a fraction of measured 2-methyltetrols may be decomposition products of low-volatility accretion products (Isaacman-VanWertz et al., 2016; Lopez-Hilfiker et al., 2016), 50% (as a rough estimate) of measured 2-methyltetrols (in the particle phases) were assumed to be dimer decomposition products when EVAPORATION-based vapor pressures were used (see Table S4). In the sensitivity calculation (Adj Psat), 2-methyltetrols were assumed to be present only in the monomer form as including dimers increased the model bias.

The overlap in the input data sets resulted in 180 hours of measurement coverage. Additional measurements of ammonium, sulfate, and ammonia (not used in this work) are summarized in Tables S5-S7 for reference.

3 Results and Discussion

3.1 Regional ammonium-sulfate conditions

Figure 1 shows the molar ratios of ammonium to $2 \times$ total sulfate and cations to anions over the eastern U.S. for 1 June - 15 July 2013 based on observations from the CSN network and predicted by CMAQv5.2. CMAQ predicted a mean $R_{N/2S}$ of 0.73 over the U.S. compared to the observed mean of 0.67. The model showed higher values (near 1) over the central U.S. and lower values over the southeast U.S. The magnitude of the CMAQ predicted $R_{N/2S}$ over the southeast U.S. (mean of 0.6) was only slightly higher than that from the CSN network (mean of 0.4). CMAQ-predicted sulfate was relatively unbiased in the southeastern U.S. (normalized mean bias of 5% compared to CSN), but ammonium was high by a factor of 1.5 (Table S10). Despite only a moderate bias in $R_{N/2S}$, significant discrepancies existed between the model and observations for the ratio of cations to anions. In CMAQ, the ratio of cations to anions was approximately one indicating that ammonia tended to be pulled into the particle in an amount necessary to neutralize sulfate not already associated with nonvolatile cations. Molar ratios are not robust indicators of aerosol pH (Hennigan et al., 2015) as a result of the role of relative humidity and associated liquid water content as well as buffering by bisulfate (Guo et al., 2015). However, chemical transport model biases in ion ratios should result in biases in acidity and gas-particle partitioning of volatile acids and bases (e.g. NH_3) considering other factors (such as RH) held constant.

Also included in Figure 1(a-b) are observations of $R_{N/2S}$ based on the SEARCH network (triangles) which are much higher (>0.8) than the CSN values (<0.6) in the southeast U.S. While there could be spatial heterogeneity in the southeast U.S., differences so large are unlikely and not present in CMAQ, thus indicating potential problems in one set of measurements. Nylon filters (used by CSN for inorganic ions) can collect 4-5% of gas-phase sulfur dioxide (Benner et al., 1991; Hansen et al., 1986), leading to a small but positive sulfate mass concentration artifact. In addition, nylon filters tend to measure lower ammonium concentrations than other filter types (Solomon et al., 2000; Yu et al., 2006). These ammonium artifacts are not restricted to ammonium nitrate since more than twice as much NH_4^+ was lost compared to nitrate on nylon filters from Great Smoky Mountains National Park, TN, U.S. (Yu et al., 2006). 6-14% of total NH_4^+ can volatilize in federal reference method (FRM) collection, and the SEARCH network best estimates of $\text{PM}_{2.5}$ result in higher ammonium on an absolute basis and as a fractional contribution to $\text{PM}_{2.5}$ compared to the FRM equivalent mass (Edgerton et al., 2005). Consider that a 10% underestimate in ammonium PM and 10% overestimate in sulfate, for example, will lead to almost a 20% underestimate in $R_{N/2S}$.

An overabundance of cations in the CMAQ model (Figure S1, Table S10) means that ammonium was displaced from the particle and $R_{N/2S}$ was biased low for the southeast U.S. An evaluation of the individual cations and anions (Figure S1, Table S10) indicated CMAQ over predicted the non-volatile ISORROPIA cations (Na^+ , Ca^{2+} , K^+ , Mg^{2+}) by factors of 2 to 6 individually and by a charge equivalent factor of 3 overall in the Southeast. A factor of 3 overestimate in nonvolatile cations indicates ammonium predicted by CMAQ was low by about 26%. Appel et al. (2013) have previously shown that even when anthropogenic fugitive dust and windblown dust emissions are removed from the CMAQ model, crustal elements are still typically overestimated compared to observations. Coal combustion, for example, is a major source of trace metals in the U.S. (Reff

et al., 2009). Trace metal emissions were overestimated (and/or physical mixing was underestimated) since CMAQ overestimated their measured concentration, which included soluble and insoluble contributions (Solomon et al., 2014). A sensitivity simulation in which all Aitken and accumulation mode Na^+ , Ca^{2+} , K^+ , and Mg^{2+} were removed from the partitioning thermodynamics resulted in a mean predicted $R_{\text{N}/2\text{S}}$ of 0.96 for the southeast U.S. Since ISORROPIA should only consider the cations associated with sulfate, nitrate, and chloride, but CMAQ includes cations that are part of insoluble metal oxides (Reff et al., 2009), additional error was incurred in CMAQ by allowing all of the calcium, potassium, magnesium, and sodium present in aerosol to participate in ISORROPIA calculations. Thus, the apparent consistency in ammonium to sulfate ratios between CSN and CMAQ, should not be used to confirm the reasonableness of either. The ratio of cations to anions indicates discrepancies between CSN and CMAQ, specifically, that the CMAQ model tends to achieve charge balance as defined by $R_{+/-}=1$ while observations indicate otherwise.

3.2 Ammonia gas-particle partitioning during SOAS

Consistent with CMAQ predictions over the greater southeastern U.S. region, CMAQ predicted an average ratio of ammonium to $2\times$ sulfate ($R_{\text{N}/2\text{S}}$) of 0.64-0.61 (for PM_{10} and $\text{PM}_{2.5}$ respectively) and 24-28% of total ammonia in the particle as ammonium (b and g in Figure 2) at CTR. CMAQ also predicted that the cation to anion charge ratio, $R_{+/-}$, was near one during SOAS. Thus, CMAQ predictions for SOAS CTR site were representative of the southeastern United States for further investigating CMAQ model issues related to inorganic molar ratios and ammonia partitioning.

As shown in Figure 2, the CMAQ predicted $R_{\text{N}/2\text{S}}$ (b) was similar to the Georgia Tech AMS derived value (a). It was also similar to the regional SEAC⁴RS AMS-derived values (Silvern et al., 2017) (Table S6) which averaged near 0.6. ISORROPIA predictions using AMS measured ammonium and sulfate as input (c), thus exactly reproducing observed $R_{\text{N}/2\text{S}}$, showed much higher partitioning of ammonia to the particle phase (mean $\text{NH}_x F_p$ of 0.8) than indicated by AMS aerosol data combined with SEARCH ammonia. Using total ammonia and ammonium as model input resulted in a similar fraction of NH_x in the particle as using only aerosol composition as input, but the $R_{\text{N}/2\text{S}}$ value significantly increased to around 0.8 (d). The AIOMFAC-based equilibrium model run with aerosol-only inputs (e) was qualitatively consistent with ISORROPIA (c) in terms of the fraction of NH_x in the particle. Since no box model simulation of AMS data in this work was able to reproduce both the $\text{NH}_x F_p$ and $R_{\text{N}/2\text{S}}$, these tests indicated that AMS measurements at SOAS CTR were inconsistent with ISORROPIA and AIOMFAC thermodynamic calculations, as found in previous model evaluation (Silvern et al., 2017).

The $R_{\text{N}/2\text{S}}$ determined from the MARGA instrument for $\text{PM}_{2.5}$ (f) was significantly higher than that derived from the AMS measurements and closer to the values based on SEARCH measurements (Table S5, Figure 1). The AMS tended to measure much less ammonium than the MARGA, and as a result, the fraction of ammonia partitioned to the particle using SEARCH NH_3 and MARGA aerosol measurements was higher than would be estimated using AMS data. The CMAQ model calculations showed a small but similar trend as observations for PM_{10} to $\text{PM}_{2.5}$ in terms of ammonia gas-particle partitioning (since $\text{PM}_{2.5} \geq \text{PM}_{10}$ and $F_p = \text{PM} / (\text{PM} + \text{gas})$) but did not show significantly increased $R_{\text{N}/2\text{S}}$ with increased particle size. Note that in full CMAQ model calculations, the predicted nonvolatile cation concentrations were so high that they erroneously affected the partitioning of ammonium (Table S10, Figure S1). Removing nonvolatile cations from CMAQ (h) allowed more ammonium

into the particle and led to increased $R_{N/2S}$, but $NH_x F_p$ was still low indicating overestimates in gas-phase ammonia in the CMAQ model are not primarily due to the displacement of ammonium by nonvolatile cations.

ISORROPIA $PM_{2.5}$ calculations using both gas and aerosol inputs were run with (j in Figure 2) and without (k) aerosol calcium, potassium, magnesium, sodium, nitrate, and chloride and the results were qualitatively the same in terms of mean fraction of ammonia partitioned to the particle and ratio of NH_4^+ to sulfate in the particle. Thus, the difference between AMS and MARGA was primarily driven by the difference in ammonium and sulfate measured by the AMS versus MARGA instrument, not the availability of nonvolatile cations. Comparing the change in mean $NH_x F_p$ with (m) and without (l) organic compound interactions indicates that organic compounds have a larger effect on ammonia gas-particle partitioning than the inclusion (j) or lack (k) of calcium, potassium, magnesium, sodium, nitrate, and chloride. Overall, ISORROPIA and AIOMFAC were qualitatively consistent with MARGA measurements of $R_{N/2S}$, but not with AMS measurements.

The inconsistency in AMS data and box models indicated in Figure 2, but ability of models to simulate MARGA data, indicates the AMS data alone may not be suitable for equilibrium thermodynamic modeling. Contributing factors could include: missing ammonium residing in particles larger than PM_1 but smaller than $PM_{2.5}$, potential missing nonvolatile cations, uncertainty in AMS-measured concentrations of sulfate and ammonium, organosulfate contributions to sulfate, or other issues. Guo (2017) indicates differences in ammonium to sulfate ratios for PM_1 measured during the first half of SOAS versus $PM_{2.5}$ measured during the second half of SOAS using the same instrument (the Particle into Liquid Sampler, PILS, (Guo et al., 2017b)) suggesting the role of particle size on ammonium to sulfate ratios. Furthermore, thermodynamic predictions of ammonia were degraded in the work of Guo et al. (2015) when the PILS inlet switched from $PM_{2.5}$ to PM_1 . Size alone does not explain the difference in AMS (PM_1) vs $PM_{2.5}$ data as AMS sulfate can be similar to (MARGA) or exceed by 20% (PILS, Guo et al. (2015)) collocated $PM_{2.5}$ sulfate. Future work that characterizes ammonium and sulfate in $PM_{2.5}$ - PM_1 would be helpful for understanding differences in AMS versus other datasets as well as to facilitate connections between AMS data and regulatory metrics including the U.S. NAAQS for $PM_{2.5}$. Guo et al. (2017b) suggest when AMS (or PILS) data are used together with nonvolatile cations, thermodynamic models can predict ammonia partitioning accurately. However, the levels of nonvolatile cations would need to be larger than current measurements indicate (Guo et al., 2017b). Furthermore, MARGA simulations in this work (Figure 2 j,k) indicated little sensitivity of $R_{N/2S}$ or $NH_x F_p$ to inclusion of measured calcium, potassium, magnesium, sodium, nitrate, and chloride.

The differences in the AMS and MARGA datasets in terms of $R_{N/2S}$ are larger than can be explained by known measurement precision. However, uncertainty for AMS measured ammonium (34%) and sulfate (36%) are large (Bahreini et al., 2009). A contributor to this uncertainty is the AMS collection efficiency (CE), and AMS instruments are known to have a higher collection efficiency for acidic (H_2SO_4 -enriched) vs $(NH_4)_2SO_4$ -enriched aerosol (Middlebrook et al., 2012). Furthermore organosulfates (Budisulistiorini et al., 2015; Hettiyadura et al., 2017) can be measured in the AMS as sulfate. However, organosulfates have been estimated to account for only 5% of AMS-measured sulfate during SOAS (Hu et al., 2017).

3.3 Phase composition

Figure 3 shows the average concentration of aerosol components predicted in the electrolyte-rich (α) and organic-rich (β) aerosol phases as well as under conditions in which only one liquid phase was predicted (single phase) based on AIOMFAC equilibrium calculations (EQLB) of the aqueous ammonium-sodium-sulfate-nitrate-chloride (A) and organic surrogates system. In all cases, water was predicted to be a major contributor to the phase accounting for 60%, 35%, and 90% of the mass in the average α , β , and single phases respectively. In addition, inorganic ions were present in all phases including the organic-rich phase. This means that the effects of inorganic species on organic compounds were not limited to times when one single liquid phase was predicted. Higher concentrations of organic species were generally associated with an increase in the predicted frequency of phase separation. However, LO-OOA, the least oxygenated (Table S2) and least water-soluble secondary organic aerosol PMF factor (Xu et al., 2017), was not more or less abundant when phase separated vs. single phase conditions were predicted.

The mean $R_{N/2S}$ varied slightly by phase with the α phase having a value of 0.8 and the phases with a greater proportion of organic compounds (β and single) having a value of 0.9. The β phase, with its higher concentration of organic species, showed a lower average $[H^+]_{\text{air}}$ (0.1 nmol m^{-3}) compared to the α phase (1.5 nmol m^{-3}), while the activity-based pH values were predicted to be similar in both phases, typically within 0.2 pH units (as expected from equilibrium thermodynamics). The ammonium-sulfate only (in terms of inorganic ion representation) system was predicted to have the same frequency of phase separation and trend in $[H^+]_{\text{air}}$, but less difference in the $R_{N/2S}$ between the phases.

Phase separation into electrolyte-rich and organic-rich phases was predicted to occur 70% of the time. The frequency of phase separation predicted for SOAS conditions was higher than the frequency predicted in previous CMAQ work (Pye et al., 2017) that calculated separation relative humidities based on average O:C ratios using the empirical parameterization of You et al. (2013) for a particular inorganic salt type. Both the previous CMAQ calculations (Pye et al., 2017) and this work predicted the same diurnal variation with a greater frequency of phase separation during the day driven by lower relative humidities (Figure S7).

3.4 Effects of organic compounds on acidity

Acidity (pH) is an important aerosol property as it promotes dissolution of metals (Fang et al., 2017), increases nutrient availability (Stockdale et al., 2016), and catalyzes particle-phase reactions (Eddingsaas et al., 2010). Current recommended methods for estimating aerosol pH include thermodynamic models and ammonia-ammonium partitioning (Hennigan et al., 2015) as direct measurements are difficult to make (Rindelaub et al., 2016). AIOMFAC predicted a median molal pH of 1.4 (ammonium-sulfate system) to 1.5 (ammonium-sodium-sulfate-nitrate-chloride system) for SOAS conditions (Table 1). AIOMFAC occasionally showed high pH (pH = 7, Figure 4) which occurred when an excess of cations compared to anions were observed, leading to the absence of H^+ and bisulfate in the input compositions used with the model. Similar behavior has occurred with ISORROPIA and the AIM thermodynamic models using aerosol-only inputs (Hennigan et al., 2015) and likely resulted from measurement uncertainty and a resulting high-bias in the measured amounts of cations compared to charge-

equivalent anions. The ISORROPIA predicted pH for the subset of conditions used here (pH = 0.7 to 1.1) was similar to those previously reported for SOAS (pH = 0.9) and Southeast Nexus (SENEX) aircraft campaign (pH = 1.1) using other datasets as summarized by Guo et al. (2017a).

Regardless of whether only ammonium-sulfate or ammonium-sodium-sulfate-nitrate-chloride systems were treated, AIOMFAC predicted an increase in the concentration of gas-phase ammonia (decrease in $\text{NH}_x F_p$, Figure 2 m compared to l or o compared to n) along with a decrease in acidity when organic compounds were considered in the calculation of partitioning (EQLB vs CLLPS, Table 1, Figure 4). The presence of organic compounds in the same phase as H^+ and other ions (EQLB case) shifted free H^+ towards increased association with sulfate to form bisulfate as AIOMFAC predicts bisulfate to be more miscible with organic compounds than H^+ and other small cations. Interactions with organic compounds resulted in a 34-36% decrease in median $[\text{H}^+]_{\text{air}}$ and a 0.1 unit increase (11-12% increase) in median pH.

If the pH for forced complete phase separation conditions was recalculated using AIOMFAC CLLPS predicted $[\text{H}^+]_{\text{air}}$ and assuming an activity coefficient of one (traditional method), the resulting pH has a median of 0.7 (Figure 4b), the same value obtained by ISORROPIA using only aerosol inputs and an activity coefficient of unity. Thus, traditional methods resulted in an artificially low pH. Taking into account activity coefficients other than unity, phase separation, and the diluting effect of organic compounds and their associated water (EQLB) resulted in a pH 0.7 pH units higher than traditional methods. This increase is substantial given that the pH scale is logarithmic; a 0.7 pH unit higher value is equivalent to a five times lower molal H^+ activity in solution. The activity coefficient value was a major driver of this difference with a secondary role for solvent abundance and change in $[\text{H}^+]_{\text{air}}$.

3.5 Partitioning of organic compounds under ambient conditions

For organic compounds with $\text{O:C} \geq 0.6$ (C_5 -alkene triols, levoglucosan, 2-methyltetrols, hydroxyglutaric acid, 2-methylglyceric acid), the particle-phase fraction, F_p , was predicted to increase when the electrolyte-rich and organic-rich phases were allowed to equilibrate (EQLB compared to CLLPS, Figures 5-6) as a result of an increase in the abundance of the partitioning medium. For compounds with lower O:C (specifically pinic and pinonic acid) F_p decreased as a result of unfavorable liquid-phase interactions. The increase in F_p for most species generally resulted in a decrease in the mean bias and mean error of F_p compared to observations (Figure 5b). With the pure-species adjusted vapor pressure (Adj Psat sensitivity), the mean bias in F_p for all organic species was less than 0.2 and emphasized that information about the pure species vapor pressure is important for accurate gas-particle partitioning calculations. The influence of inorganic constituents on organic compound partitioning was not limited to the times when one single phase was present. In the case of hydroxyglutaric acid (Figure 6g), predictions of F_p were found to be most sensitive to assumptions regarding condensed phase mixing during the day when phase separation was most common (coinciding with a lower average RH during midday and afternoon hours, as expected). This occurred because the organic-rich phase still contained a significant amount of inorganic ions (Figure 3) which modified the partitioning medium and impacted the predicted activity of the organic species.

The change in F_p between CLLPS and EQLB calculations was consistent with the change in effective saturation concentrations (Figure 5c). The effective C^* (equation 4) under equilibrium (EQLB) conditions compared to CLLPS (EQLB

C*/CLLPS C*) was a strong function of the compound O:C ratio (Pearson's $r^2=0.79$) with higher O:C species having lower EQLB C*/CLLPS C* ratios. The mean activity coefficient value was predicted to either stay the same (2-methylglyceric acid) or increase (all other explicit organic species) in EQLB compared to CLLPS. Thus, the driving factor for increased partitioning to the particle phase (indicated by increased F_p and decreased C*) for species with O:C>0.6 under EQLB compared to CLLPS was the ability of the increased partitioning medium size to overcome the increased activity coefficients. The increased partitioning medium gained by interacting with the inorganic species and their water lowered the mole fraction of the organic species in the particle, thus leading to lower predicted particle-phase activity and gas-phase concentrations via modified Raoult's law. In some cases, like for 2-methyltetrols, the species exhibited negative deviations from ideality ($\gamma < 1$) in both CLLPS and EQLB, but the activity coefficient still increased from CLLPS to EQLB (Table S9). For pinic and pinonic acid, the deviation ($\gamma > 1$) was positive in CLLPS and its activity coefficient even larger in magnitude in EQLB such that the larger partitioning medium did not overcome the deviation in ideality resulting in the species being more abundant in the gas phase in EQLB compared to CLLPS. Interestingly, levoglucosan was the only species predicted to have an activity coefficient near 1 for the organic-rich (β) phase in EQLB calculations (Table S9). Due to the effect on the size of the partitioning medium resulting from additional species (specifically water and inorganic salts) in the β phase during EQLB, the effective C* for levoglucosan was predicted to be 35% of its pure species value ($1.4 \mu\text{g m}^{-3}$, Table S8).

Predicted unfavorable interactions (limited miscibility within both the organic-rich and inorganic-rich liquid phases) resulted in pinonic acid (Figure 6f) being partitioned to the gas phase to a much greater degree than the measurements indicated. Model performance was consistent with previous work in which multiple measurement techniques showed slightly higher F_p than model predictions (Thompson et al., 2017). Formation of a second organic-rich phase (a third liquid phase) containing lower O:C compounds, which was not allowed in the AIOMFAC calculations, could improve pinonic acid partitioning predictions. The lack of a resolved hydrocarbon-like organic aerosol (HOA) component (Xu et al., 2015a) and representation of its associated functional groups in the model may have also contributed to an unfavorable environment for low O:C compounds.

Overall, the treatment of liquid phase mixing vs separation did not improve the mean bias in 2-methyltetrol predicted F_p . It also did not significantly change the mean error. The average fraction of 2-methyltetrols in the particles was represented fairly well by assuming half of the measured 2-methyltetrols are actually decomposition products of a fairly nonvolatile ($C^*=10^{-6} \mu\text{g m}^{-3}$) dimer compound (dark grey square, Figure 5a,b). However, this assumption did not perform equally well at all times of day. Figure 6a indicates that the 50% dimer assumption leads to an underestimate in 2-methyltetrol F_p during the day and overestimate at night. Modeling 2-methyltetrols as entirely monomers with a pure species C* of $8 \mu\text{g m}^{-3}$ at 298.15K (factor of 4.2 reduction in EVAPORATION predicted vapor pressure) reproduced the daytime 2-methyltetrol partitioning well, but overestimated partitioning to the particle at night. Even with the reduced P^{sat} (in the Adj Psat sensitivity), 2-methyltetrol monomers remained slightly more volatile than predicted by SIMPOL ($C^*=5 \mu\text{g m}^{-3}$) at reference conditions. The average effective 2-methyltetrol C* (accounting for the effects of temperature and partitioning medium) in the case of CLLPS was $6 \mu\text{g m}^{-3}$ while in the equilibrium calculations (EQLB) it was reduced further to $3.7 \mu\text{g m}^{-3}$ (Table S8). Thus, 2-methyltetrols behaved like compounds with an effective mean saturation concentration roughly half of the pure species value due to the influence of temperature and presence of other species in the particle.

4 Conclusions

In this work, conditions over the eastern United States were examined with a focus on gas-particle partitioning during the Southern Oxidant and Aerosol Study (SOAS). Different measurement techniques indicated fairly different ratios of ammonium to $2 \times$ total sulfate with the AMS instruments having the lowest values followed by CSN. The MARGA instrument (Allen et al., 2015) and SEARCH network indicated the highest ratios of ammonium to $2 \times$ sulfate of slightly less than 1 (mean of 0.8 to 0.9). The lack of agreement of AMS and CSN data with thermodynamic models, but the agreement between MARGA observations and models, indicates a potential bias in CSN measurements and that AMS data alone may not be suitable for thermodynamic modeling. The diversity in observational datasets can explain why some work has concluded thermodynamic models fail (Silvern et al., 2017) while others indicate models are adequate (Weber et al., 2016). This work finds thermodynamic equilibrium models (both ISORROPIA and AIOMFAC) are consistent with high ammonium to $2 \times$ sulfate ratios in conjunction with about 70 to 80% of ammonia in the particle. Lower ammonium to sulfate ratios imply much higher fractions of total ammonia in the particle as thermodynamic equilibrium assumptions (and models) generally do not allow a large excess of gas-phase ammonia under highly acidic conditions. While consideration of inorganics mixing in liquid phases with organic compounds may increase pH significantly compared to estimates from traditional models like ISORROPIA, that effect is likely not the cause of current inorganic aerosol model evaluation issues.

By performing ISORROPIA and AIOMFAC box modeling, this work demonstrates that our current thermodynamic understanding of ammonium and sulfate aerosol is consistent with (MARGA) observations in the southeastern U.S. atmosphere. Since models like CMAQ use the same thermodynamic basis, specifically ISORROPIA, these results build confidence that regional models can capture the thermodynamics of the ambient atmosphere. However, our results also demonstrate that for the partitioning of ammonia and ammonium to be correct, errors in emissions of nonvolatile cations, on the order of a factor of 3, must be resolved as well.

AIOMFAC-based predictions of gas-particle partitioning of organic compounds were sensitive to pure species vapor pressure estimates and predictions generally had a lower mean bias when EVAPORATION-based vapor pressures were adjusted downward by a factor of 4.2 and close to values estimated by SIMPOL for 2-methyltetrols, pinic acid, and hydroxyglutaric acid. AIOMFAC predicted organic compounds interact with significant amounts of water and inorganic ions. 2-methyltetrol predictions had roughly the same error in particle fraction (F_p) assuming 50% of measured particulate 2-methyltetrols were decomposition products or if their vapor pressure was adjusted downward by a factor of 4.2 (to $P^{\text{sat}}=1.4 \times 10^{-4}$ Pa at 298.15 K).

Code and data availability. CMAQ model code is available at <https://github.com/USEPA/CMAQ> and v5.2-gamma was used in this work.

ISORROPIA is available from <http://isorro피아.eas.gatech.edu/>.

AIOMFAC can be run online (<http://www.aiomfac.caltech.edu/>) or via contact with A. Zuend.

SOAS data is available at <https://esrl.noaa.gov/csd/groups/csd7/measurements/2013senex/>.

CSN data is available at <https://www.epa.gov/outdoor-air-quality-data>.

Model output associated with the final article will be available from the EPA Environmental Dataset Gateway at <https://edg.epa.gov/> if accepted.

Competing interests. The authors declare no competing interests.

Disclaimer. The US EPA through its Office of Research and Development supported the research described here. It has been subjected to Agency administrative review and approved for publication but may not necessarily reflect official Agency policy.

Acknowledgements. We thank J. Jimenez, G. Ruggeri, S. Takahama, and S. Lee for providing additional datasets that are summarized in the supporting information. We thank the CSN and SEARCH networks for providing long-term measurements. We thank two reviewers at EPA. We thank Paul Solomon for useful discussion. We thank CSRA for preparing emissions and meteorology input for CMAQ simulations. AZ was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), grant RGPIN/04315-2014. JLF acknowledges support from EPA-STAR RD-83539901. GIVW was supported by the NSF Graduate Research Fellowship (DGE 1106400). F_p of organic compounds collected by SV-TAG at SOAS was supported by grants to UC Berkeley including NSF Atmospheric Chemistry Program 1250569 and Department of Energy SBIR grant DE-SC0004698. LX and NLN acknowledge support from National Science Foundation (NSF) Grant 1242258 and US Environmental Protection Agency (EPA) STAR Grant RD-83540301.

References

- Allen, H. M., Draper, D. C., Ayres, B. R., Ault, A., Bondy, A., Takahama, S., Modini, R. L., Baumann, K., Edgerton, E., Knote, C., Laskin, A., Wang, B., and Fry, J. L.: Influence of crustal dust and sea spray supermicron particle concentrations and acidity on inorganic NO_3^- aerosol during the 2013 Southern Oxidant and Aerosol Study, *Atmos. Chem. Phys.*, 15, 10669–10685, doi:10.5194/acp-15-10669-2015, 5 2015.
- Appel, K. W., Pouliot, G. A., Simon, H., Sarwar, G., Pye, H. O. T., Napelenok, S. L., Akhtar, F., and Roselle, S. J.: Evaluation of dust and trace metal estimates from the Community Multiscale Air Quality (CMAQ) model version 5.0, *Geosci. Model Dev.*, 6, 883–899, doi:10.5194/gmd-6-883-2013, 2013.
- Bahreini, R., Ervens, B., Middlebrook, A. M., Warneke, C., de Gouw, J. A., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., Neuman, J. A., Ryerson, T. B., Stark, H., Atlas, E., Brioude, J., Fried, A., Holloway, J. S., Peischl, J., Richter, D., Walega, J., Weibring, P., Wollny, A. G., and Fehsenfeld, F. C.: Organic aerosol formation in urban and industrial plumes near Houston and Dallas, Texas, *J. Geophys. Res.*, 114, n/a–n/a, doi:10.1029/2008JD011493, <http://dx.doi.org/10.1029/2008JD011493>, d00F16, 2009.
- Benner, C. L., Eatough, D. J., Eatough, N. L., and Bhardwaja, P.: Comparison of annular denuder and filter pack collection of $\text{HNO}_3(\text{g})$, $\text{HNO}_2(\text{g})$, $\text{SO}_2(\text{g})$, and particulate-phase nitrate, nitrite and sulfate in the south-west desert, *Atmos. Environ.*, 25, 1537–1545, 15 doi:10.1016/0960-1686(91)90013-W, 1991.
- Bertram, A. K., Martin, S. T., Hanna, S. J., Smith, M. L., Bodsworth, A., Chen, Q., Kuwata, M., Liu, A., You, Y., and Zorn, S. R.: Predicting the relative humidities of liquid-liquid phase separation, efflorescence, and deliquescence of mixed particles of ammonium sulfate, organic material, and water using the organic-to-sulfate mass ratio of the particle and the oxygen-to-carbon elemental ratio of the organic component, *Atmos. Chem. Phys.*, 11, 10995–11006, doi:10.5194/acp-11-10995-2011, 2011.
- 20 Budisulistiorini, S. H., Li, X., Bairai, S. T., Renfro, J., Liu, Y., Liu, Y. J., McKinney, K. A., Martin, S. T., McNeill, V. F., Pye, H. O. T., Nenes, A., Neff, M. E., Stone, E. A., Mueller, S., Knote, C., Shaw, S. L., Zhang, Z., Gold, A., and Surratt, J. D.: Examining the effects of anthropogenic emissions on isoprene-derived secondary organic aerosol formation during the 2013 Southern Oxidant and Aerosol Study (SOAS) at the Look Rock, Tennessee ground site, *Atmos. Chem. Phys.*, 15, 8871–8888, doi:10.5194/acp-15-8871-2015, 2015.
- Budisulistiorini, S. H., Nenes, A., Carlton, A. G., Surratt, J. D., McNeill, V. F., and Pye, H. O. T.: Simulating aqueous-phase isoprene-epoxydiol (IEPOX) secondary organic aerosol production during the 2013 Southern Oxidant and Aerosol Study (SOAS), *Environ. Sci. Technol.*, 51, 5026–5034, doi:10.1021/acs.est.6b05750, 2017.
- Chan, M. N., Chan, A. W. H., Chhabra, P. S., Surratt, J. D., and Seinfeld, J. H.: Modeling of secondary organic aerosol yields from laboratory chamber data, *Atmos. Chem. Phys.*, 9, 5669–5680, doi:10.5194/acp-9-5669-2009, 2009.
- Compernelle, S., Ceulemans, K., and Müller, J. F.: EVAPORATION: a new vapour pressure estimation method for organic molecules including non-additivity and intramolecular interactions, *Atmos. Chem. Phys.*, 11, 9431–9450, doi:10.5194/acp-11-9431-2011, 2011.
- Eddingsaas, N. C., VanderVelde, D. G., and Wennberg, P. O.: Kinetics and products of the acid-catalyzed ring-opening of atmospherically relevant butyl epoxy alcohols, *J. of Phys. Chem. A*, 114, 8106–8113, doi:10.1021/jp103907c, 2010.
- Edgerton, E. S., Hartsell, B. E., Saylor, R. D., Jansen, J. J., Hansen, D. A., and Hidy, G. M.: The Southeastern Aerosol Research and Characterization Study: Part II. Filter-based measurements of fine and coarse particulate matter mass and composition, *J. Air Waste* 35 *Manag. Assoc.*, 55, 1527–1542, 2005.

- Edgerton, E. S., Hartsell, B. E., Saylor, R. D., Jansen, J. J., Hansen, D. A., and Hidy, G. M.: The Southeastern Aerosol Research and Characterization Study, part 3: Continuous measurements of fine particulate matter mass and composition, *J. Air Waste Manag. Assoc.*, 56, 1325–1341, 2006.
- Fang, T., Guo, H., Zeng, L., Verma, V., Nenes, A., and Weber, R. J.: Highly acidic ambient particles, soluble metals, and oxidative potential: a link between sulfate and aerosol toxicity, *Environ. Sci. Technol.*, 51, 2611–2620, doi:10.1021/acs.est.6b06151, 2017.
- Foroutan, H., Young, J., Napelenok, S., Ran, L., Appel, K. W., Gilliam, R. C., and Pleim, J. E.: Development and evaluation of a physics-based windblown dust emission scheme implemented in the CMAQ modeling system, *J. Adv. Model Earth Sy.*, 9, 585–608, doi:10.1002/2016MS000823, 2017.
- Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K^+ - Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O aerosols, *Atmos. Chem. and Phys.*, 7, 4639–4659, 2007.
- Goldstein, A. H. and Galbally, I. E.: Known and unexplored organic constituents in the Earth’s atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, doi:10.1021/es072476p, 2007.
- Guo, H.: Interactive comment on “The underappreciated role of nonvolatile cations on aerosol ammonium-sulfate molar ratios” by Hongyu Guo et al., *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2017-737-SC3, 2017.
- 15 Guo, H., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Hite Jr, J. R., Carlton, A. G., Lee, S. H., Bergin, M. H., Ng, N. L., Nenes, A., and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmos. Chem. Phys.*, 15, 5211–5228, doi:10.5194/acp-15-5211-2015, 2015.
- Guo, H., Liu, J., Froyd, K. D., Roberts, J. M., Veres, P. R., Hayes, P. L., Jimenez, J. L., Nenes, A., and Weber, R. J.: Fine particle pH and gas–particle phase partitioning of inorganic species in Pasadena, California, during the 2010 CalNex campaign, *Atmos. Chem. Phys.*, 17, 5703–5719, doi:10.5194/acp-17-5703-2017, 2017a.
- 20 Guo, H., Nenes, A., and Weber, R. J.: The underappreciated role of nonvolatile cations on aerosol ammonium-sulfate molar ratios, *Atmos. Chem. Phys. Discuss.*, 2017, 1–19, doi:10.5194/acp-2017-737, 2017b.
- Hansen, L. D., White, V. F., and Eatough, D. J.: Determination of gas-phase dimethyl sulfate and monomethyl sulfate, *Environ. Sci. Technol.*, 20, 872–878, doi:10.1021/es00151a004, 1986.
- 25 Heath, N. K., Pleim, J. E., Gilliam, R. C., and Kang, D.: A simple lightning assimilation technique for improving retrospective WRF simulations, *J. Adv. Model Earth Sys.*, 8, 1806–1824, doi:10.1002/2016MS000735, 2016.
- Hennigan, C. J., Izumi, J., Sullivan, A. P., Weber, R. J., and Nenes, A.: A critical evaluation of proxy methods used to estimate the acidity of atmospheric particles, *Atmos. Chem. Phys.*, 15, 2775–2790, doi:10.5194/acp-15-2775-2015, 2015.
- Hettiyadura, A. P. S., Jayarathne, T., Baumann, K., Goldstein, A. H., de Gouw, J. A., Koss, A., Keutsch, F. N., Skog, K., and Stone, E. A.: Qualitative and quantitative analysis of atmospheric organosulfates in Centreville, Alabama, *Atmos. Chem. and Phys.*, 17, 1343–1359, doi:10.5194/acp-17-1343-2017, 2017.
- 30 Hu, W., Palm, B. B., Day, D. A., Campuzano-Jost, P., Krechmer, J. E., Peng, Z., de Sá, S. S., Martin, S. T., Alexander, M. L., Baumann, K., Hacker, L., Kiendler-Scharr, A., Koss, A. R., de Gouw, J. A., Goldstein, A. H., Seco, R., Sjostedt, S. J., Park, J. H., Guenther, A. B., Kim, S., Canonaco, F., Prévôt, A. S. H., Brune, W. H., and Jimenez, J. L.: Volatility and lifetime against OH heterogeneous reaction of ambient isoprene-epoxydiols-derived secondary organic aerosol (IEPOX-SOA), *Atmos. Chem. Phys.*, 16, 11 563–11 580, doi:10.5194/acp-16-11563-2016, 2016.
- 35

- Hu, W., Campuzano-Jost, P., Day, D. A., Croteau, P., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., and Jimenez, J. L.: Evaluation of the new capture vaporizer for aerosol mass spectrometers (AMS) through field studies of inorganic species, *Aerosol Sci. Tech.*, 51, 735–754, doi:10.1080/02786826.2017.1296104, 2017.
- 5 Isaacman, G., Kreisberg, N. M., Yee, L. D., Worton, D. R., Chan, A. W. H., Moss, J. A., Hering, S. V., and Goldstein, A. H.: Online derivatization for hourly measurements of gas- and particle-phase semi-volatile oxygenated organic compounds by thermal desorption aerosol gas chromatography (SV-TAG), *Atmos. Meas. Tech.*, 7, 4417–4429, doi:10.5194/amt-7-4417-2014, 2014.
- Isaacman-VanWertz, G., Yee, L. D., Kreisberg, N. M., Wernis, R., Moss, J. A., Hering, S. V., de Sá, S. S., Martin, S. T., Alexander, M. L., Palm, B. B., Hu, W., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Riva, M., Surratt, J. D., Viegas, J., Manzi, A., Edgerton, E., Baumann, K., Souza, R., Artaxo, P., and Goldstein, A. H.: Ambient gas-particle partitioning of tracers for biogenic oxidation, *Environ. Sci. Technol.*, 10 50, 9952–9962, doi:10.1021/acs.est.6b01674, 2016.
- Johnson, D., Utembe, S. R., Jenkin, M. E., Derwent, R. G., Hayman, G. D., Alfarra, M. R., Coe, H., and McFiggans, G.: Simulating regional scale secondary organic aerosol formation during the TORCH 2003 campaign in the southern UK, *Atmos. Chem. Phys.*, 6, 403–418, doi:10.5194/acp-6-403-2006, 2006.
- Kwamena, N. O. A., Buajareern, J., and Reid, J. P.: Equilibrium morphology of mixed organic/inorganic/aqueous aerosol droplets: investigating the effect of relative humidity and surfactants, *J. Phys. Chem. A*, 114, 5787–5795, doi:10.1021/jp1003648, 2010.
- 15 Lopez-Hilfiker, F. D., Mohr, C., D'Ambro, E. L., Lutz, A., Riedel, T. P., Gaston, C. J., Iyer, S., Zhang, Z., Gold, A., Surratt, J. D., Lee, B. H., Kurten, T., Hu, W. W., Jimenez, J., Hallquist, M., and Thornton, J. A.: Molecular composition and volatility of organic aerosol in the southeastern U.S.: implications for IEPOX derived SOA, *Environ. Sci. Technol.*, 50, 2200–2209, doi:10.1021/acs.est.5b04769, 2016.
- Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of composition-dependent collection efficiencies for the Aerodyne Aerosol Mass Spectrometer using field data, *Aerosol Sci. Technol.*, 46, 258–271, doi:10.1080/02786826.2011.620041, 20 2012.
- Nenes, A., Pandis, S. N., and Pilinis, C.: ISORROPIA: A new thermodynamic equilibrium model for multiphase multicomponent inorganic aerosols, *Aquat. Geochem.*, 4, 123–152, doi:10.1023/A:1009604003981, 1998.
- Nolte, C. G., Appel, K. W., Kelly, J. T., Bhave, P. V., Fahey, K. M., Collett Jr, J. L., Zhang, L., and Young, J. O.: Evaluation of the Community 25 Multiscale Air Quality (CMAQ) model v5.0 against size-resolved measurements of inorganic particle composition across sites in North America, *Geosci. Model Dev.*, 8, 2877–2892, doi:10.5194/gmd-8-2877-2015, 2015.
- O'Meara, S., Booth, A. M., Barley, M. H., Topping, D., and McFiggans, G.: An assessment of vapour pressure estimation methods, *Phys. Chem. Chem. Phys.*, 16, 19453–19469, doi:10.1039/C4CP00857J, 2014.
- Otte, T. L. and Pleim, J. E.: The Meteorology-Chemistry Interface Processor (MCIP) for the CMAQ modeling system: Updates through 30 MCIPv3.4.1, *Geosci. Model Dev.*, 3, 243–256, 2010.
- Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting vapor pressures and enthalpies of vaporization of multifunctional organic compounds, *Atmos. Chem. Phys.*, 8, 2773–2796, doi:10.5194/acp-8-2773-2008, 2008.
- Pleim, J. E., Bash, J. O., Walker, J. T., and Cooter, E. J.: Development and evaluation of an ammonia bidirectional flux parameterization for air quality models, *J. Geophys. Res.*, 118, 3794–3806, doi:10.1002/jgrd.50262, 2013.
- 35 Pye, H. O. T., Murphy, B. N., Xu, L., Ng, N. L., Carlton, A. G., Guo, H., Weber, R., Vasilakos, P., Appel, K. W., Budisulistiorini, S. H., Surratt, J. D., Nenes, A., Hu, W., Jimenez, J. L., Isaacman-VanWertz, G., Misztal, P. K., and Goldstein, A. H.: On the implications of aerosol liquid water and phase separation for organic aerosol mass, *Atmos. Chem. Phys.*, 17, 343–369, doi:10.5194/acp-17-343-2017, 2017.

- Reff, A., Bhave, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D., and Houyoux, M.: Emissions inventory of PM_{2.5} trace elements across the United States, *Environ. Sci. Technol.*, 43, 5790–5796, doi:10.1021/es802930x, 2009.
- Reid, J. P., Dennis-Smith, B. J., Kwamena, N.-O. A., Miles, R. E. H., Hanford, K. L., and Homer, C. J.: The morphology of aerosol particles consisting of hydrophobic and hydrophilic phases: hydrocarbons, alcohols and fatty acids as the hydrophobic component, *Phys. Chem. Chem. Phys.*, 13, 15 559–15 572, doi:10.1039/C1CP21510H, 2011.
- 5 Rindelaub, J. D., Craig, R. L., Nandy, L., Bondy, A. L., Dutcher, C. S., Shepson, P. B., and Ault, A. P.: Direct measurement of pH in individual particles via Raman microspectroscopy and variation in acidity with relative humidity, *J. of Phys Chem. A*, 120, 911–917, doi:10.1021/acs.jpca.5b12699, 2016.
- Silvern, R. F., Jacob, D. J., Kim, P. S., Marais, E. A., Turner, J. R., Campuzano-Jost, P., and Jimenez, J. L.: Inconsistency of ammonium–sulfate aerosol ratios with thermodynamic models in the eastern US: a possible role of organic aerosol, *Atmos. Chem. Phys.*, 17, 5107–5118, doi:10.5194/acp-17-5107-2017, 2017.
- 10 Solomon, P. A., Mitchell, W., Tolocka, M., Norris, G., Gemmill, D., Wiener, R., Vanderpool, R., Murdoch, R., Natarajan, S., and Hardison, E.: Evaluation of PM_{2.5} chemical speciation samplers for use in the EPA National PM_{2.5} Chemical Speciation Network, Report EPA-454/R-01-005, United States Environmental Protection Agency, 2000.
- 15 Solomon, P. A., Crumpler, D., Flanagan, J. B., Jayanty, R. K. M., Rickman, E. E., and McDade, C. E.: U.S. National PM_{2.5} Chemical Speciation Monitoring Networks—CSN and IMPROVE: Description of networks, *J. Air Waste Manag. Assoc.*, 64, 1410–1438, doi:10.1080/10962247.2014.956904, 2014.
- Song, M., Marcolli, C., Krieger, U. K., Zuend, A., and Peter, T.: Liquid-liquid phase separation in aerosol particles: Dependence on O:C, organic functionalities, and compositional complexity, *Geophys. Res. Lett.*, 39, L19 801, doi:10.1029/2012GL052807, 2012.
- 20 Song, M., Marcolli, C., Krieger, U. K., Lienhard, D. M., and Peter, T.: Morphologies of mixed organic/inorganic/aqueous aerosol droplets, *Faraday Discuss.*, 165, 289–316, doi:10.1039/C3FD00049D, 2013.
- Stockdale, A., Krom, M. D., Mortimer, R. J. G., Benning, L. G., Carslaw, K. S., Herbert, R. J., Shi, Z., Myriokefalitakis, S., Kanakidou, M., and Nenes, A.: Understanding the nature of atmospheric acid processing of mineral dusts in supplying bioavailable phosphorus to the oceans, *P. Natl. Acad. Sci. USA*, 113, 14 639–14 644, doi:10.1073/pnas.1608136113, 2016.
- 25 Thompson, S. L., Yatavelli, R. L. N., Stark, H., Kimmel, J. R., Krechmer, J. E., Day, D. A., Hu, W., Isaacman-VanWertz, G., Yee, L., Goldstein, A. H., Khan, M. A. H., Holzinger, R., Kreisberg, N., Lopez-Hilfiker, F. D., Mohr, C., Thornton, J. A., Jayne, J. T., Canagaratna, M., Worsnop, D. R., and Jimenez, J. L.: Field intercomparison of the gas/particle partitioning of oxygenated organics during the Southern Oxidant and Aerosol Study (SOAS) in 2013, *Aerosol Sci. Technol.*, 51, 30–56, doi:10.1080/02786826.2016.1254719, 2017.
- Topping, D., Barley, M., Bane, M. K., Higham, N., Aumont, B., Dingle, N., and McFiggans, G.: UManSysProp v1.0: an online and open-source facility for molecular property prediction and atmospheric aerosol calculations, *Geosci. Model Dev.*, 9, 899–914, doi:10.5194/gmd-9-899-2016, 2016.
- 30 Weber, R. J., Guo, H., Russell, A. G., and Nenes, A.: High aerosol acidity despite declining atmospheric sulfate concentrations over the past 15 years, *Nature Geosci.*, 9, 282–285, doi:10.1038/ngeo2665 <http://www.nature.com/ngeo/journal/v9/n4/abs/ngeo2665.html#supplementary-information>, 2016.
- 35 Xu, L., Guo, H., Boyd, C. M., Klein, M., Bougiatioti, A., Cerully, K. M., Hite, J. R., Isaacman-VanWertz, G., Kreisberg, N. M., Knote, C., Olson, K., Koss, A., Goldstein, A. H., Hering, S. V., de Gouw, J., Baumann, K., Lee, S.-H., Nenes, A., Weber, R. J., and Ng, N. L.: Effects of anthropogenic emissions on aerosol formation from isoprene and monoterpenes in the southeastern United States, *P. Natl. Acad. Sci. USA*, 112, 37–42, doi:10.1073/pnas.1417609112, 2015a.

- Xu, L., Suresh, S., Guo, H., Weber, R. J., and Ng, N. L.: Aerosol characterization over the southeastern United States using high-resolution aerosol mass spectrometry: spatial and seasonal variation of aerosol composition and sources with a focus on organic nitrates, *Atmos. Chem. Phys.*, 15, 7307–7336, doi:10.5194/acp-15-7307-2015, 2015b.
- Xu, L., Guo, H., Weber, R. J., and Ng, N. L.: Chemical characterization of water-soluble organic aerosol in contrasting rural and urban environments in the southeastern United States, *Environ. Sci. Technol.*, 51, 78–88, doi:10.1021/acs.est.6b05002, 2017.
- 5 You, Y., Renbaum-Wolff, L., and Bertram, A. K.: Liquid-liquid phase separation in particles containing organics mixed with ammonium sulfate, ammonium bisulfate, ammonium nitrate or sodium chloride, *Atmos. Chem. Phys.*, 13, 11 723–11 734, doi:10.5194/acp-13-11723-2013, 2013.
- Yu, X.-Y., Lee, T., Ayres, B., Kreidenweis, S. M., Malm, W., and Collett Jr, J. L.: Loss of fine particle ammonium from denuded nylon filters, *Atmos. Environ.*, 40, 4797–4807, doi:https://doi.org/10.1016/j.atmosenv.2006.03.061, 2006.
- Zuend, A. and Seinfeld, J. H.: Modeling the gas-particle partitioning of secondary organic aerosol: the importance of liquid-liquid phase separation, *Atmos. Chem. Phys.*, 12, 3857–3882, doi:10.5194/acp-12-3857-2012, 2012.
- Zuend, A. and Seinfeld, J. H.: A practical method for the calculation of liquid-liquid equilibria in multicomponent organic-water-electrolyte systems using physicochemical constraints, *Fluid Phase Equilib.*, 337, 201–213, doi:10.1016/j.fluid.2012.09.034, 2013.
- 15 Zuend, A., Marcolli, C., Luo, B. P., and Peter, T.: A thermodynamic model of mixed organic-inorganic aerosols to predict activity coefficients, *Atmos. Chem. Phys.*, 8, 4559–4593, doi:10.5194/acp-8-4559-2008, 2008.
- Zuend, A., Marcolli, C., Peter, T., and Seinfeld, J. H.: Computation of liquid-liquid equilibria and phase stabilities: implications for RH-dependent gas/particle partitioning of organic-inorganic aerosols, *Atmos. Chem. Phys.*, 10, 7795–7820, doi:10.5194/acp-10-7795-2010, 2010.
- 20 Zuend, A., Marcolli, C., Booth, A. M., Lienhard, D. M., Soonsin, V., Krieger, U. K., Topping, D. O., McFiggans, G., Peter, T., and Seinfeld, J. H.: New and extended parameterization of the thermodynamic model AIOMFAC: calculation of activity coefficients for organic-inorganic mixtures containing carboxyl, hydroxyl, carbonyl, ether, ester, alkenyl, alkyl, and aromatic functional groups, *Atmos. Chem. and Phys.*, 11, 9155–9206, 2011.




Figure 1. Molar ratio of aerosol ammonium to $2\times$ sulfate ($R_{N/2S}$) (a-b) and cations to anions ($R_{+/-}$) (c-d) over the eastern US for June 1- July 15, 2013 based on observations and predicted by the CMAQ model.




Figure 2. Gas-particle partitioning of ammonia ($\text{NH}_x \text{F}_p = \text{ammonium}/(\text{ammonia}+\text{ammonium})$), mean $R_{\text{NI}/2\text{S}}$ (red \times), and mean $R_{+/-}$ (blue \circ) for PM_{10} measured by the Georgia Tech AMS (Xu et al., 2015a) and $\text{PM}_{2.5}$ measured by a MARGA (Allen et al., 2015) as well as predicted by a CMAQ regional chemical transport model calculation and box models for SOAS conditions at CTR. F_p boxplots indicate the maximum, 75th percentile, median, 25th percentile, and minimum. Short dashes within the boxplots indicate the mean F_p . Box model inputs were either the aerosol (A) or aerosol and gas concentrations (A+G). Box models were run with either the ammonium-sulfate system (A') or including all cations and anions (A). AIOMFAC calculations assumed complete liquid-liquid phase separation between the organic-rich and electrolyte-rich phases (CLLPS) or employed a full equilibrium calculation with organic compounds in which phase separation was calculated based on composition (EQLB). Observed gas-phase ammonia concentrations are from the SEARCH network at CTR. Boxplots are labeled by a letter for easier reference in the text. Shading of the boxplot interquartile range distinguishes different models (CMAQ, ISORROPIA, and AIOMFAC). The horizontal lines correspond to mean observed $\text{NH}_x \text{F}_p$ (black) and $R_{\text{NI}/2\text{S}}$ (red). A simulation is consistent with observations if it reproduces both $\text{NH}_x \text{F}_p$ and $R_{\text{NI}/2\text{S}}$.




Figure 3. Average composition of the α (electrolyte-rich), β (organic-rich), and single phase in terms of (a) mass (organic and inorganic components) and (b) moles (ions only) predicted by AIOMFAC using $\text{PM}_{2.5}$ aerosol composition observed during SOAS. Species are stacked in the same order as indicated by the legend.

Table 1. $[\text{H}^+]_{\text{air}}$ and pH predicted for $\text{PM}_{2.5}$ at SOAS CTR (median \pm one standard deviation) under conditions of complete liquid-liquid phase separation between the organic-rich and electrolyte-rich phases (CLLPS) or in a full equilibrium calculation in which phase separation was calculated based on composition (EQLB).

Model	CLLPS	EQLB
$[\text{H}^+]_{\text{air}}$ in nmol m^{-3} air		
AIOMFAC (A')	1.9 ± 1.9	1.3 ± 1.6
AIOMFAC (A)	1.8 ± 2.1	1.1 ± 1.8
ISORROPIA (A)	2.0 ± 2.8	NA
ISORROPIA (A+G)	0.5 ± 1.5	NA
$pH = -\log_{10}(\gamma_{\text{H}^+}[\text{H}^+]_{\text{air}}/[S])$		
AIOMFAC (A')	1.3 ± 1.2	1.4 ± 1.2
AIOMFAC (A)	1.3 ± 2.1	1.5 ± 2.0
ISORROPIA (A)	0.7 ± 2.5	NA
ISORROPIA (A+G)	1.1 ± 0.7	NA




Figure 4. (a) [H⁺]_{air} and (b) pH predicted for PM_{2.5} using AIOMFAC. Dashed lines in panel (a) indicate a factor of two difference from the 1:1 line. Dashed lines in (b) represent a ± 0.5 shift in pH while dotted lines represent a ± 1 shift in pH. Series marked in open circles (○) are summarized in Table 1. All calculations used the ammonium-sodium-sulfate-nitrate-chloride and organic compounds system.




Figure 5. Observed and AIOMFAC-based predictions of equilibrium partitioning of organic compounds in the presence of MARGA-measured $PM_{2.5}$ inorganics. In panel (b), mean bias = $\frac{1}{n} \sum_{i=1}^n (M_i - O_i)$ and mean error = $\frac{1}{n} \sum_{i=1}^n |M_i - O_i|$ where M_i is the model prediction and O_i is the observation of F_p . The ratio of mean saturation concentration under EQLB compared to CLLPS conditions (c) uses predictions from the adjusted vapor pressure calculations (Adj Psat). Modeled particulate 2-methyltetrols are 50% dimers except with Adj Psat.




Figure 6. Fraction of each explicit organic species in the particle as a function of hour of day between 1 June and 15 July 2013 at CTR. 2-methyltetrols were modeled as 50% dimers in the particle for CLLPS and EQLB. When the pure species vapor pressure was adjusted, 2-methyltetrols were assumed to be entirely monomers. Fit is based on traditional absorptive partitioning to an organic compounds-only phase (Equation S1).