

Supplement information

Open burning of rice, corn and wheat straw: primary emissions, photochemical aging, and secondary organic aerosol formation

Zheng Fang^{1,3}, Wei Deng^{1,3}, Yanli Zhang^{1,2}, Xiang Ding¹, Mingjin Tang¹, Tengyu Liu¹, Qihou Hu¹, Ming Zhu^{1,3}, Zhaoyi Wang^{1,3}, Weiqiang Yang^{1,3}, Zhonghui Huang^{1,3}, Wei Song^{1,2}, Xinhui Bi¹, Jianmin Chen⁴, Yele Sun⁵, Christian George⁶, Xinming Wang^{1,2,*}

¹State Key Laboratory of Organic Geochemistry and Guangdong Key Laboratory of Environment Protection and Resources Utilization, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

²Center for Excellence in Regional Atmospheric Environment, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention, Department of Environmental Science & Engineering, Fudan University, Shanghai 200433, China

⁵Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

⁶Institut de Recherches sur la Catalyse et l'Environnement de Lyon (IRCELYON), CNRS, UMR5256, Villeurbanne F-69626, France

Correspondence to: X. Wang (wangxm@gig.ac.cn)

23 **Table S1.** Emission factors for the non-methane hydrocarbons (NMHCs) species from the straw burning.

No. Species	EF (g kg ⁻¹)			No. Species	EF (g kg ⁻¹)		
	Rice	Corn	Wheat		Rice	Corn	Wheat
1 ethene	1.316±0.492	0.540±0.484	0.777±0.666	35 methyl-cyclopentane	0.003±0.004	0.001±0.001	0.001±0.001
2 acetylene	0.958±0.328	0.149±0.108	0.293±0.294	36 2,4-dimethylpentane	0.001±0.002	ND ^a	ND
3 propene	0.315±0.323	0.342±0.295	0.332±0.272	37 cyclohexane	0.020±0.042	0.001±0.001	ND
4 1-butene	0.045±0.018	0.027±0.029	0.044±0.071	38 2-methyl-hexane	0.002±0.004	ND	0.001±0.001
5 1,3-butadiene	0.030±0.038	0.116±0.110	0.104±0.080	39 2,3-dimethyl-pentane	0.001±0.004	ND	ND
6 trans-2-butene	0.016±0.005	0.035±0.031	0.040±0.038	40 3-methyl-hexane	0.006±0.009	0.001±0.002	ND
7 cis-2-butene	0.013±0.004	0.025±0.022	0.030±0.028	41 2,2,4-trimethyl-pentane	0.002±0.003	ND	ND
8 3-methyl-1-butene	0.007±0.003	0.009±0.009	0.008±0.006	42 n-heptane	0.037±0.058	0.003±0.004	0.003±0.002
9 1-pentene	0.017±0.011	0.029±0.031	0.021±0.025	43 methyl-cyclohexane	0.003±0.006	ND	ND
10 2-methyl-1-butene	0.012±0.008	0.004±0.002	0.013±0.011	44 2,3,4-trimethyl-pentane	0.000±0.001	ND	ND
11 isoprene	0.096±0.101	0.026±0.035	0.060±0.045	45 2-methyl-heptane	0.011±0.025	0.001±0.001	ND
12 trans-2-pentene	0.010±0.011	0.012±0.010	0.016±0.013	46 3-methyl-heptane	0.001±0.001	ND	ND
13 cis-2-pentene	0.010±0.005	0.007±0.006	0.007±0.006	47 n-octane	0.005±0.005	0.002±0.003	0.002±0.002
14 2-methyl-2-butene	0.018±0.021	0.013±0.012	0.015±0.012	48 n-nonane	0.015±0.024	0.002±0.003	0.002±0.001
15 cyclopentene	0.005±0.007	0.008±0.007	0.008±0.007	49 n-decane	0.005±0.006	0.004±0.005	ND
16 4-methyl-1-pentene	0.003±0.004	0.002±0.003	0.001±0.001	50 n-undecane	ND	0.007±0.012	ND
17 1-hexene	0.007±0.004	0.016±0.025	ND	51 benzene	0.567±0.172	0.163±0.124	0.249±0.200
18 trans-2-hexene	ND	0.003±0.005	ND	52 toluene	0.206±0.148	0.142±0.186	0.134±0.099
19 cis-2-hexene	0.001±0.003	0.001±0.001	0.001±0.001	53 ethyl-benzene	0.027±0.017	0.016±0.024	0.012±0.009
20 3-hexene	0.002±0.003	0.002±0.002	0.001±0.002	54 m/p-xylene	0.045±0.043	0.020±0.029	0.030±0.023
21 a-pinene	0.008±0.022	ND	ND	55 styrene	0.095±0.033	0.035±0.031	0.029±0.021
22 b-pinene	0.011±0.019	ND	ND	56 o-xylene	0.016±0.014	0.008±0.011	0.008±0.006
23 ethane	0.492±0.603	0.452±0.357	0.586±0.615	57 isopropylbenzene	0.003±0.004	0.001±0.001	0.001±0.001
24 propane	0.171±0.196	0.127±0.127	0.103±0.155	58 n-propylbenzene	0.003±0.003	0.003±0.004	0.002±0.001
25 isobutane	0.052±0.043	0.011±0.011	0.004±0.007	59 m-ethyltoluene	0.005±0.003	0.003±0.003	0.005±0.004
26 n-butane	0.062±0.060	0.034±0.030	0.078±0.129	60 p-ethyltoluene	0.005±0.004	0.002±0.003	0.003±0.003
27 isopentane	0.036±0.051	0.015±0.016	0.031±0.035	61 1,3,5-trimethyl-benzene	0.003±0.004	0.001±0.002	0.002±0.003
28 n-pentane	0.016±0.008	0.011±0.014	0.006±0.009	62 o-ethyltoluene	0.003±0.002	0.002±0.003	0.002±0.002
29 2,2-dimethyl-butane	0.004±0.007	ND	ND	63 1,2,4-trimethylbenzene	0.006±0.006	0.004±0.006	0.006±0.006
30 cyclopentane	ND	0.001±0.001	0.002±0.004	64 1,2,3-trimethylbenzene	0.039±0.086	0.007±0.010	0.005±0.005
31 2,3-dimethylbutane	ND	ND	0.001±0.001	65 m-diethylbenzene	0.003±0.005	0.001±0.001	0.001±0.001
32 2-methylpentane	0.008±0.011	0.002±0.002	0.001±0.001	66 p-diethylbenzene	0.004±0.007	0.002±0.002	0.001±0.001
33 3-methylpentane	0.006±0.007	0.008±0.014	ND	67 o-diethylbenzene	0.002±0.004	0.001±0.001	ND
34 n-hexane	0.120±0.175	0.001±0.002	0.001±0.001				

(^a ND=not detected)

25 **Table S2.** Secondary organic aerosol yields reported in literature and used in our work for
 26 different precursors.

compound	yields reported in literature			yield used in our work		
	yield reference	seed ^a	yield range ^b	applied yield ^c	lower bound	upper bound
acrolein	Chhabra et al., 2011	yes	0.022	0.026	0.022	0.035
	Chan et al., 2010	yes	0.022-0.035			
furan	Gómez Alvarez et al., 2009	no	0.019-0.072	0.019	0.019	0.072
crotonaldehyde or methacrolein ^d	Chhabra et al., 2011	yes	0.02	0.02	0.019	0.194
	Chan et al., 2010	yes	0.019-0.194			
	Chhabra et al., 2011	yes	0.02			
	Chan et al., 2010	yes	0.019-0.044			
benzene	Ng et al., 2007	yes	0.28-0.37	0.33	0.28	0.37
	Nakao et al., 2011	no	0.19-0.28			
	Borras and Tortajada-Genaro, 2012	no	0.016-0.097			
2/3-methylfuran	Gómez Alvarez et al., 2009	no	0.055-0.085	0.085	0.055	0.085
	Chan et al., 2010	yes	0.008-0.391			

27
 28
 29

30 **Table S2.** Secondary organic aerosol yields reported in literature and used in our work for
 31 different precursors (continued).

compound	literature data			application of yield data		
	yield reference	seed ^a	yield range ^b	applied yield ^c	lower bound	upper bound
toluene	Chhabra et al., 2011	yes	0.11-0.36	0.26	0.08	0.66
	Ng et al., 2007	yes	0.08-0.31			
	Nakao et al., 2011	no	0.17-0.23			
	Hildebrandt et al., 2009	yes	0.26-0.66			
phenol	Yee et al., 2013	yes	0.24-0.54	0.38	0.13	0.54
	Nakao et al., 2011	no	0.38-0.45			
	Borras and Tortajada-Genaro, 2012	no	0.13-0.18			
	Chhabra et al., 2011	yes	0.34-0.38			
2,4-/2,5-dimethylfuran ^e				0.38	0.13	0.54
styrene ^f				0.22	0.04	0.4
benzaldehyde ^g				0.38	0.27	0.49
m-xylene	Chhabra et al., 2011	yes	0.06-0.4	0.22	0.04	0.4
	Ng et al., 2007	yes	0.04-0.38			
	Nakao et al., 2011	no	0.1-0.32			
o/m-cresol	Nakao et al., 2011	no	0.27-0.49	0.38	0.27	0.49
catechol or benzenediol	Nakao et al., 2011	no	0.39	0.53	0.39	0.53
	Borras and Tortajada-Genaro, 2012	no	0.45-0.53			
dimethylphenol	Nakao et al., 2011	no	0.13-0.9	0.52	0.13	0.9
guaiacol	Yee et al., 2013	yes	0.35-0.5	0.4	0.35	0.5
	Chhabra et al., 2011	yes	0.36-0.39			

32 **Table S2.** Secondary organic aerosol yields reported in literature and used in our work for
 33 different precursors (continued).

compound	literature data			application of yield data		
	yield reference	seed ^a	yield range ^b	applied yield ^c	lower bound	upper bound
naphthalene	Chhabra et al., 2011	yes	0.36-0.39			
	Chhabra et al., 2011	yes	0.33-0.67	0.36	0.11	0.74
	Chan et al., 2009	yes	0.2-0.74			
	Shakya and Griffin, 2010	no	0.11-0.12			
1/2-methylnaphthalene	Chan et al., 2009	yes	0.19-0.71	0.45	0.19	0.71
	Shakya and Griffin, 2010	no	0.04-0.15			
acenaphthalene	Shakya and Griffin, 2010	no	0.03-0.04	0.03	0.03	0.04
acenaphthene	Shakya and Griffin, 2010	no	0.04-0.05	0.05	0.04	0.05
1,2-dimethylnaphthalene	Chan et al., 2009	yes	0.31	0.31	0.31	0.31

34 (a Yields obtained without seed aerosol are not taken into account if yields obtained with seed aerosol are available;

35 b the yields at the maximum and minimum NO_x/NMOGs ratios are determined to be the boundary values; c if the

36 NO_x/NMOGs ratio in this study (1.2±0.9) are within the range in the corresponding literature, the average value

37 of boundary values is used, or the yield at NO_x/NMOGs ratio closer to this study is used; besides, yields obtained

38 from different studies were averaged; d averaged SOA yields were applied for isomers since they cannot be


39 resolved by PTR-TOF-MS; e because of lack of available reported value, 2,4-/2,5-dimethylfuran is assumed to

40 have the same SOA yield as phenol; f styrene is assumed to have the same SOA yield as m-xylene; g benzaldehyde



41 is assumed to have the same SOA yield as o/m-cresol)

42

43



44
 45 **Figure S1** Pearson coefficients of correlations between the modified combustion efficiency
 46 (MCE) and individual NMHC concentrations. The number order of NMHC species is the same
 47 as Table S1.



48

49 **Figure S2** Correlations of modified combustion efficacy (MCE) with emission factors (EFs)

50 for (a) particulate matter and (b) primary organic carbon.

51 **Reference**

- 52 Borrás, E., and Tortajada-Genaro, L. A.: Secondary organic aerosol formation from the photo-
53 oxidation of benzene, *Atmos. Environ.*, 47, 154-163, doi:10.1016/j.atmosenv.2011.11.020,
54 2012.
- 55 Chan, A. W. H., Kautzman, K. E., Chhabra, P. S., Surratt, J. D., Chan, M. N., Crouse, J. D.,
56 Kürten, A., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol
57 formation from photooxidation of naphthalene and alkylnaphthalenes: implications for
58 oxidation of intermediate volatility organic compounds (IVOCs), *Atmos. Chem. Phys.*, 9,
59 3049-3060, doi:10.5194/acp-9-3049-2009, 2009.
- 60 Chan, A. W. H., Chan, M. N., Surratt, J. D., and Chhabra, P. S.: Role of aldehyde chemistry
61 and NO_x concentrations in secondary organic aerosol formation, *Atmos. Chem. Phys.*, 10,
62 7169-7188, doi:10.5194/acp-10-7169-2010, 2010.
- 63 Chhabra, P. S., Ng, N. L., Canagaratna, M. R., Corrigan, A. L., Russell, L. M., Worsnop, D. R.,
64 Flagan, R. C., and Seinfeld, J. H.: Elemental composition and oxidation of chamber
65 organic aerosol, *Atmos. Chem. Phys.*, 11, 8827-8845, doi:10.5194/acp-11-8827-2011,
66 2011.
- 67 Christian, T. J., Yokelson, R. J., Cardenas, B., Molina, L. T., Engling, G., and Hsu, S. C.: Trace
68 gas and particle emissions from domestic and industrial biofuel use and garbage burning
69 in central Mexico, *Atmos. Chem. Phys.*, 10, 565-584, doi:10.5194/acp-10-565-2010, 2010.
- 70 Gómez Alvarez, E., Borrás, E., Viidanoja, J., and Hjorth, J.: Unsaturated dicarbonyl products
71 from the OH-initiated photo-oxidation of furan, 2-methylfuran and 3-methylfuran, *Atmos.*
72 *Environ.*, 43, 1603-1612, doi:10.1016/j.atmosenv.2008.12.019, 2009.
- 73 Hildebrandt, L., Donahue, N. M., and Pandis, S. N.: High formation of secondary organic
74 aerosol from the photo-oxidation of toluene, *Atmos. Chem. Phys.*, 9, 2973-2986,
75 doi:10.5194/acp-9-2973-2009, 2009.
- 76 Kim Oanh, N. T., Tipayarom, A., Bich, T. L., Tipayarom, D., Simpson, C. D., Hardie, D., and
77 Sally Liu, L. J.: Characterization of gaseous and semi-volatile organic compounds emitted
78 from field burning of rice straw, *Atmos. Environ.*, 119, 182-191,
79 doi:10.1016/j.atmosenv.2015.08.005, 2015.
- 80 Li, C., Hu, Y., Zhang, F., Chen, J., Ma, Z., Ye, X., Yang, X., Wang, L., Tang, X., Zhang, R., Mu,
81 M., Wang, G., Kan, H., Wang, X., and Mellouki, A.: Multi-pollutant emissions from the
82 burning of major agricultural residues in China and the related health-economic effects,
83 *Atmos. Chem. Phys.*, 17, 4957-4988, doi:10.5194/acp-17-4957-2017, 2017.
- 84 Li, X. G., Wang, S. X., Duan, L., Hao, J., Li, C., Chen, Y. S., and Yang, L.: Particulate and trace
85 gas emissions from open burning of wheat straw and corn stover in China, *Environ. Sci.*
86 *Technol.*, 41, 6052-6058, doi:10.1021/es0705137, 2007.
- 87 Nakao, S., Clark, C., Tang, P., Sato, K., and Iii, D. C.: Secondary organic aerosol formation
88 from phenolic compounds in the absence of NO_x, *Atmos. Chem. Phys.*, 11, 2025-2055,
89 doi:10.5194/acp-11-10649-2011, 2011.
- 90 Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.:
91 Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem.*
92 *Phys.*, 7, 3909-3922, doi:10.5194/acp-7-3909-2007, 2007.
- 93 Ni, H. Y., Han, Y. M., Cao, J. J., Chen, L. W. A., Tian, J., Wang, X. L., Chow, J. C., Watson, J.

- 94 G., Wang, Q. Y., Wang, P., Li, H., and Huang, R. J.: Emission characteristics of
95 carbonaceous particles and trace gases from open burning of crop residues in China,
96 *Atmos. Environ.*, 123, 399-406, doi:10.1016/j.atmosenv.2015.05.007, 2015.
- 97 Shakya, K. M., and Griffin, R. J.: Secondary Organic Aerosol from Photooxidation of
98 Polycyclic Aromatic Hydrocarbons, *Environ. Sci. Technol.*, 44, 8134-8139,
99 doi:10.1021/es1019417, 2010.
- 100 Yee, L. D., Kautzman, K. E., Loza, C. L., Schilling, K. A., Coggon, M. M., Chhabra, P. S.,
101 Chan, M. N., Chan, A. W. H., Hersey, S. P., Crounse, J. D., Wennberg, P. O., Flagan, R.
102 C., and Seinfeld, J. H.: Secondary organic aerosol formation from biomass burning
103 intermediates: phenol and methoxyphenols, *Atmos. Chem. Phys.*, 13, 8019-8043,
104 doi:10.5194/acp-13-8019-2013, 2013.
- 105 Yokelson, R. J., Burling, I. R., Urbanski, S. P., Atlas, E. L., Adachi, K., Buseck, P. R.,
106 Wiedinmyer, C., Akagi, S. K., Toohey, D. W., and Wold, C. E.: Trace gas and particle
107 emissions from open biomass burning in Mexico, *Atmos Chem Phys*, 11, 6787-6808,
108 doi:10.5194/acp-11-6787-2011, 2011.