

Contributions of natural and anthropogenic sources to ambient ammonia in the Athabasca Oil Sands and north-western Canada

Cynthia Whaley¹, Paul A. Makar¹, Mark W. Shephard¹, Leiming Zhang¹, Junhua Zhang¹, Qiong Zheng¹, Ayodeji Akingunola¹, Gregory R. Wentworth^{2,3}, Jennifer G. Murphy², Shailesh K. Kharol¹, and Karen E. Cady-Pereira⁴

¹Air Quality Research Division, Environment and Climate Change Canada, 4905 Dufferin Street, Toronto, Ontario, Canada

²Dept of Chemistry, University of Toronto, 80 St George Street, Toronto, Ontario, Canada

³Environmental Monitoring and Science Division, Alberta Environment and Parks, 9888 Jasper Ave NW, Edmonton, Alberta, Canada

⁴Atmospheric and Environmental Research (AER), Lexington, Massachusetts, USA

Correspondence to: Cynthia Whaley (cynthia.whaley@canada.ca)

Abstract. Atmospheric ammonia (NH₃) is a short-lived pollutant that plays an important role in aerosol chemistry and nitrogen deposition. Dominant NH₃ emissions are from agriculture and forest fires, both of which are increasing globally. ~~The Alberta Oil Sands region has~~ Even remote regions with relatively low ambient NH₃ concentrations ~~because of its remote location~~, such as northern Alberta and Saskatchewan in northern Canada ~~; however, a may be of interest because of industrial Oil Sands emissions, and a sensitive ecological system.~~ A previous attempt to model NH₃ in the region showed a substantial negative bias compared to satellite profiles and aircraft observations. Known missing sources of NH₃ in the model were re-emission of NH₃ from plants and soils (bidirectional flux), and forest fire emissions, but the relative impact of these sources on NH₃ concentrations ~~and column totals~~ was unknown. Here we have used a research version of the high-resolution air quality forecasting model, GEM-MACH, to quantify the relative impacts of ~~natural~~ semi-natural (bidirectional flux of NH₃ and forest fire emissions) and direct anthropogenic (Oil Sands operations, combustion of fossil fuels, and agriculture) sources on ammonia concentrations, both at the surface and aloft, with a focus on the Athabasca Oil Sands region ~~;~~ during a measurement-intensive campaign in the summer of 2013. The addition of fires and bidirectional flux to GEM-MACH has improved the model bias, slope and correlation coefficients relative to ground, aircraft, and satellite NH₃ measurements significantly.

By running the ~~GEM-MACH~~ GEM-MACH-Bidi model in three configurations and calculating their differences, we find that averaged over Alberta and Saskatchewan during this time period ~~;~~ an average of 23.1% of surface NH₃ came from direct anthropogenic sources, 56.6% (or 1.24 ppbv) from bidirectional flux (re-emission from plants and soils), and 20.3% (or 0.42 ppbv) from forest fires. In the NH₃ total column, an average of 19.5% came from direct anthropogenic sources, 50.0%

from bidirectional flux, and 30.5% from forest fires. The addition of bidirectional flux and fire emissions caused the overall average net ~~flux of NH_3~~ deposition of NH_x across the domain to be positive ~~(upward). It also increased the NH_4^+ wet deposition by nearly a factor of three during the period simulated~~ increased by 24.5%. Note that forest fires are very episodic and their contributions will vary significantly for different time periods and regions.

This study is the first use of the bidirectional flux scheme in GEM-MACH, which could be generalized for other volatile or semi-volatiles species. It is also the first time CrIS satellite observations of NH_3 have been used for model evaluation, and the first use of fire emissions in GEM-MACH at 2.5-km resolution.

1 Introduction

Ammonia (NH_3) is a short-lived pollutant that is receiving global attention because of its increasing concentrations. Emissions of NH_3 – which are in large part from agricultural fertilizer, livestock (Behera et al., 2013; Environment and Climate Change Canada, 2016), and biomass burning (Olivier et al., 1998; Krupa, 2003) – have not been regulated to the same extent as other nitrogen species. NH_3 is the only aerosol precursor whose global emissions are projected to rise throughout the next century (Moss et al., 2010; Lamarque et al., 2010; Ciais et al., 2013).

NH_3 has an atmospheric lifetime of hours to a day (Seinfeld and Pandis, 1998; Aneja et al., 2001).

It is a base that reacts in the atmosphere with sulphuric acid (H_2SO_4) and nitric acid (HNO_3) to form crystalline sulphate, nitrate salts (e.g., $(\text{NH}_4)_2\text{SO}_4$, NH_4HSO_4 , NH_4NO_3) and aqueous ions (SO_4^{2-} , HSO_4^- , NO_3^-), (Nenes et al., 1998; Makar et al., 2003) which are significant components of fine particulate matter ($\text{PM}_{2.5}$) (e.g., Jimenez et al., 2009; [Environment Canada, 2001](#)), thus causing health (Pope III et al., 2002; Lee et al., 2015) and climate impacts (IPCC, 2013). A large portion of NH_3 is readily deposited in the first 4-5 km from its source, but when in fine particulate form (as NH_4^+), its lifetime is days to several weeks (Galperin and Sofiev, 1998; Park et al., 2004; Behera et al., 2013; Paulot et al., 2014) and can be transported hundreds of kilometers (Krupa, 2003; Galloway et al., 2008; Makar et al., 2009). Deposition of NH_3 and these aerosols can lead to nitrogen eutrophication and soil acidification (Fangmeier et al., 1994; Sutton et al., 1998; Dragosits et al., 2002; Carfrae et al., 2004). ~~Thus,~~ NH_3 is listed as a Criteria Air Contaminant (Environment and Climate Change Canada, 2017) in order to help address air quality issues such as smog and acid rain.

Modelling ~~provides information on NH_3 concentrations where there are no measurements, and~~ can be used to better understand NH_3 processes. Recent NH_3 models have focused on improving bidirectional flux processes and impacts of livestock. Measurements of NH_3 bidirectional flux include those in Farquhar et al. (1980); Sutton et al. (1993, 1995); Asman et al. (1998); Nemitz et al. (2001), with indirect support for bidirectional flux also in Ellis et al. (2011). Thus, these studies were the motivation for the recent design of parameterizations to describe this important process (Wu et al., 2009;

Wichink Kruit et al., 2010; Massad et al., 2010; Zhang et al., 2010; Zhu et al., 2015; Fu et al., 2015; Hansen et al., 2017). Additionally, satellite observations are providing valuable insight on ammonia concentrations and emissions both on regional and global scales (Beer et al., 2008; Clarisse et al., 2009; Shephard et al., 2011; Shephard and Cady-Pereira, 2015; Van Damme et al., 2014; Zhu et al., 2013).

The Athabasca Oil Sands region (AOSR), located in the north-eastern part of the province of Alberta, Canada, is a large source of air (Gordon et al., 2015; Liggio et al., 2016; Li et al., 2017), ecosystem (Kelly et al., 2009; Kirk et al., 2014; Hsu and Clair, 2015) and greenhouse gas (Charpentier et al., 2009) ~~pollution~~ pollution to air (Gordon et al., 2015; Liggio et al., 2016; Li et al., 2017) and ecosystems (Kelly et al., 2009; Kirk et al., 2014; Hsu and Clair, 2015), as well as a source of greenhouse gases (Charpentier et al., 2009) due to mining and processing by the oil industry. ~~However~~ While NH_3 concentrations surrounding the AOSR in northern Alberta and Saskatchewan remain relatively low (e.g., averaging 0.63 ± 0.57 ppbv at the surface, around 0.6-1.2 ppbv background (this study) ~~and averaging 1.2 \pm 0.2 ppbv aloft (Shephard et al., 2015) compared to agricultural areas in the south of the provinces and Shephard et al., 2015) – due to low population and lack of agriculture, the northern Alberta and Saskatchewan ecosystems are sensitive to nitrogen deposition (Clair and Percy, 2015; Wieder et al., 2016a, b; Vitt, 2016; M, and the modelled background NH_3 must be correct in order to understand the relative impacts of the oil sands operations. It is important to understand if the AOSR facilities are causing critical levels of NH_3 , and if not, if any other kinds of sources (e.g., Makar et al., 2009) ~~However, a fires, re-emissions~~ are. A monitoring study from 2005 to 2008 found NH_3 concentrations near Fort McMurray and Fort McKay (population centers in the vicinity of the oil sands facilities) to be highly variable in space and time with a range of 1.1 to 8.8 ppbv (Bytnerowicz et al., 2010) ~~– (where the upper end corresponds to NH_3 levels found in agricultural regions of Canada and the U.S.), with NH_3 concentrations 1.5-3 \times higher than HNO_3 concentrations (Bytnerowicz et al., 2010). Hsu and Clair (2016) also found NH_3 concentrations in the AOSR to be much higher than HNO_3 , NO_3^- , and NH_4^+ concentrations (by 5, 23, and 1.8 \times , respectively). Thus, NH_3 may contribute the largest fraction of deposited nitrogen in the AOSR compared to other nitrogen species like NO_2 and HNO_3^- (Bytnerowicz et al., 2010; Hsu and Clair, 2015; Kharol et al., 2017). Estimates of deposition of nitrogen and sulphur compounds in the AOSR are described in Makar et al. (2017) in this issue, however they did not include NH_3 bidirectional flux or forest fires in their model simulations.~~~~

In a previous study by Shephard et al. (2015) it was found that the GEM-MACH air quality forecasting model (Moran et al., 2010, 2013; Makar et al., 2015a, b; Gong et al., 2015), using a domain covering the Canadian provinces of Alberta and Saskatchewan, at 2.5-km resolution, under-predicted summertime tropospheric ammonia concentrations by 0.4-0.6 ppbv (which is 36-100 % depending on altitude - see Fig. 16 in Shephard et al., 2015) in the AOSR when compared to Tropospheric Emission Spectrometer (TES) satellite measurements and aircraft measurements. ~~NH₃~~ Having too much modelled NH_3 deposition is a cause that was ruled out when Makar et al. (2017) showed

95 that GEM-MACH actually underestimates NH_x deposition. Underestimating anthropogenic and
agricultural emissions was also ruled out as a cause because the GEM-MACH model performs
well in southern Canada and the U.S when compared to the U.S. Ambient Ammonia Monitoring
Network (AMoN). NH_3 sources known to be missing from the GEM-MACH model were forest fire
emissions and re-emission of deposited NH_3 from soils and plants (the latter referred to as bidirec-
100 tional flux, hereafter). ~~These~~, which would have the greatest impact in background areas, such as
northern Alberta and Saskatchewan. Therefore, these two sources were added to an updated version
of GEM-MACH and model simulations were repeated for a 2013 summer period (12 August to 7
September 2013) during which an intensive measurement campaign occurred. We utilize ground,
aircraft and satellite measurements of NH_3 and related species to evaluate the model and to quantify
105 the impacts of the different sources on atmospheric NH_3 and its deposition.

Section 2 provides the model description. Section 3 provides a brief description of ammonia mea-
surements during the campaign. Section 4 presents the evaluation of three model scenarios against
three different types of measurements (surface, aircraft, and satellite), and Section 5 presents our
quantitative assessment on the impacts of different sources of NH_3 to ambient concentrations and
110 NH_x deposition in the region. Our conclusions appear in Section 6.

2 GEM-MACH model description

GEM-MACH (Global Environment Multiscale-Modelling Air quality and CHemistry) is an on-line
chemical transport model, which is embedded in GEM, Environment and Climate Change Canada
(ECCC)'s numerical weather prediction model (Moran et al., 2010). This means that the chemical
115 processes of the model (gas-phase chemistry, plume rise emissions distribution, vertical diffusion
and surface fluxes of tracers, and a particle chemistry package including particle microphysics, cloud
processes, and inorganic heterogeneous chemistry) are imbedded within the meteorological model's
physics package, this component in turn is imbedded within the meteorological model's dynamics
package, which also handles chemical tracer advection. A detailed description of the process repre-
120 sentation of GEM-MACH, and an evaluation of its performance for pollutants such as ozone and par-
ticulate matter (PM) appears in Moran et al. (2013); Makar et al. (2015a, b); Gong et al. (2015); Akingunola et al. (2017)
Moran et al. (2013); Makar et al. (2015a, b); Gong et al. (2015).

GEM-MACH is used operationally to issue twice-daily, 48-hour public forecasts of criteria air pol-
lutants (ozone, nitrogen oxides, PM), as well as the the Air Quality Health Index [https://ec.gc.ca/cas-aqhi/].
125 Any improvements to NH_3 in the model may result in better AQHI predictions, since NH_3 is a ma-
jor precursor of $\text{PM}_{2.5}$, as mentioned in the introduction. We start with a similar, research version of
GEM-MACHv2 to make the bidirectional flux modifications. The key differences between this and
older versions are the use of a more recent meteorological package (GEMv4.8), the capability to nest

in the vertical dimension as well as the horizontal dimension, and improvements to the treatment of
130 fluxes, vertical diffusion, and advection(~~Akingunola et al. (2017), this issue~~).

GEM-MACH can be run for many different spatial domains, at various spatial resolutions, and in
2-bin or 12-bin aerosol size distribution modes. For this study we run the model in the 2-bin mode
(for computational efficiency), using a nested set of domains. The outer domain at 10-km resolution
~~covering covers~~ North America, and the inner domain at 2.5-km resolution ~~covering covers~~ the
135 provinces of Alberta and Saskatchewan. The latter is referred to as the 2.5-km Oil Sands domain.
~~And this~~ This set up, along with the emissions described in the next section is hereafter called our
“base” simulation.

2.1 Emissions

The emissions of 25 species (SO₂, SO₄ (gas), sulphate, nitrate, NH₄⁺, NO, NO₂, NH₃, CO, nitrous
140 acid, benzene, propane, higher alkanes, higher alkenes, ethene, toluene, aromatics, formaldehyde,
aldehydes, methyl ethyl ketone, creosol, isoprene, crustal material, elemental carbon, and primary
carbon) used in GEM-MACH (base case) come from Canadian and U.S. emissions inventories: 2011
National Emissions Inventory (NEI) version 1 for U.S. emissions, and the Air Pollutant Emission
Inventory (APEI) 2013 for Canadian emissions (2010 for onroad and offroad emissions). Emissions
145 were processed with SMOKE (Sparse Matrix Operator Kernel Emissions, <https://www.cmascenter.org/smoke/>)
to convert the inventories into model-ready gridded hourly emissions files for modeling, separated
into major point emissions (typically industrial emissions from stacks, emitted into the model layers
that correspond to the stack height, at the reported temperature and velocity in the inventory’s
stack parameters), and area emissions (emissions from spread-out sources, such as transportation
150 and agriculture, emitted into the first model layer). For more details about these emissions, see
Moran et al. (2015) and (~~Zhang et al.,~~ Zhang et al. (2017, this issue).

The emissions data for NH₃ from oil sands sources are reported to the Canadian National Pollu-
tant Release Inventory (NPRI) on a “total annual emissions per facility” basis. NH₃ emissions are
generally more uncertain than SO₂ and NO_x emissions because NH₃ emissions are not measured to
155 the same extent as those two. The oil sands represent only 1% of total Alberta NH₃ emissions, at
approximately 1438 tonnes in 2013. For comparison, about 18× more NO_x and 57× more SO₂ was
emitted from the oil sands facilities that year (<http://www.ec.gc.ca/inrp-npri/donnees-data/index.cfm?lang=En>).
However, we found an issue with NH₃ in this inventory that impacted our model evaluation in the
region, which we describe below. The emissions data for NH₃ from oil sands sources are reported to
160 the Canadian National Pollutant Release Inventory (NPRI) on a “total annual emissions per facility”
basis. NH₃ emissions are generally more uncertain than SO₂ and NO_x emissions because NH₃
emissions are not measured to the same extent as those two. The oil sands represent only 1% of total
Alberta NH₃ emissions, at approximately 1438 tonnes in 2013. For comparison, about 18× more
NO_x and 57× more SO₂ was emitted from the oil sands facilities that year (<http://www.ec.gc.ca/inrp-npri/donnees-data/index.cfm?lang>

165 However, we found an issue with NH₃ in this inventory that impacted our model evaluation in the region, which we describe below.

If stack parameters (e.g., stack height and diameter, volume flow rates, temperatures, etc.) are included as part of that data, then the emissions are allocated to large stacks in our configuration of the SMOKE emissions processing system. In the absence of this information, SMOKE will assign
170 default stack parameters based on its source category code. For the Syncrude Canada Ltd. - Mildred Lake Plant Site, NPRI ID 2274 ([a facility in the AOSR](#)), the default stack parameters were: 18.90 m for the stack height (which is within the first model layer), 0.24 m for the stack diameter, 320.0 K for the exhaust temperature, and 0.58 m/s for the exhaust velocity. However, when these defaults were applied in initial model simulations, they were found to result in erroneous short term
175 plume events with simulated surface NH₃ levels up to 2 orders of magnitude higher than ground observations ([Wentworth et al., 2017](#)), and modelled concentrations aloft too low compared to aircraft measurements (see Section 3). Conversely, for species such as SO₂, for which stack parameters were reported, the model was able to correctly place the SO₂ enhancements in space and time, relative to observations. When the stack parameters of the main stack for this facility were used for NH₃
180 emissions as well (stack height=183 m, stack diameter=7.9 m, exit temperature=513 K, exit velocity=23.9 m/s, from the NPRI website), the simulation of surface NH₃ was greatly improved. All subsequent simulations reported here make use of this correction.

2.2 Ammonia bidirectional flux parameterization

~~The bidirectional flux~~ NH₃ can be both deposited from the atmosphere to the ground, and re-emitted
185 from soils and plants back to the atmosphere. The two taken together are called bidirectional flux, since the flux of NH₃ can go both up and down. The source of NH₃ available for re-emissions are from the accumulated NH₃ in the soil and stomatal water, which can arise from increased deposition from anthropogenic sources, as well as from organic nitrogen decomposition (Booth et al., 2005), N₂-fixation (Vile et al., 2014), and natural microbial action (McCalley and Sparks, 2008).

The bidirectional flux scheme of Zhang et al. (2010) was applied within the GEM-MACHv2 model, replacing the original deposition velocity for NH₃ only (deposition velocity of other gas species follows a scheme based on a multiple resistance approach and a single-layer “big leaf” approach (Wesely, 1989; Zhang et al., 2002; Robichaud and Lin, 1991; Robichaud, 1994)). The bidirectional flux scheme is described in detail in Zhang et al. (2010), but we summarize it here.

195 Bidirectional exchange occurs between air-soil and air-stomata interfaces. The bidirectional flux (F_t) equation is:

$$F_t = -\frac{C_a - C_c}{R_a + R_b} \quad (1)$$

where R_a and R_b are the aerodynamic and quasi-laminar resistances, respectively. C_a is the NH_3 concentration in the air, and C_c is the canopy compensation point concentration, given by Eq. (2).

$$C_c = \frac{\frac{C_a}{R_a + R_b} + \frac{C_{st}}{R_{st}} + \frac{C_g}{R_{ac} + R_g}}{(R_a + R_b)^{-1} + (R_{st})^{-1} + (R_{ac} + R_g)^{-1} + (R_{cut})^{-1}} \quad (2)$$

where C_{st} and C_g are the stomatal and ground compensation points, and R_i are the resistances in s/m of the ground/soil (R_g), stomata (R_{st}), cuticle (R_{cut}), and in-canopy aerodynamic (R_{ac}). All resistance formulas can be found in Zhang et al. (2003).

Stomata (st) and ground (g) compensation points are both calculated using Eq. (3):

$$C_{st,g} = \frac{A}{T_{st,g}} \exp\left(\frac{-B}{T_{st,g}}\right) \exp\left(\frac{-B}{T_{st,g}}\right) \Gamma_{st,g} \quad (3)$$

A and B are constants derived from the equilibria constants for $\text{NH}_3(\text{g})$ in leaves' stomatal cavities to NH_4^+ and OH^- in the water contained in the apoplast within the leaf and in the soil where $\text{NH}_3(\text{g})$ in the soil pore air space is in equilibrium with the NH_4^+ and OH^- dissolved in soil water (Pleim et al., 2013). $A=161500 \text{ mol K/L}$ (Nemitz et al., 2000), or $2.7457 \times 10^{15} \text{ ugK/m}^3$ (Pleim et al., 2013) for NH_3 for both stomata and soil. $B=10380$ (Nemitz et al., 2000). $\Gamma_{st,g}$ is the emission potential of the stomata and ground, respectively and, in theory, is equal to the NH_4^+ concentration over the H^+ concentration in the apoplast water of the canopy leaves or soil water:

$$\Gamma_{st,g} = \frac{[\text{NH}_4^+]_{st,g}}{[\text{H}^+]_{st,g}} \quad (4)$$

However, since there are no modeled NH_4^+ and H^+ apoplast water concentrations to use, we use $\Gamma_{st,g}$ ~~in from~~ Wen et al. (2014), which are based on long-term empirical averages. Wen et al. (2014) gives a range of values for emission potentials for 26 land use categories (LUCs), and we use the low-end of the values in our model with the following exceptions: We further lower the Γ_g for agriculture LUCs to 800, and increase Γ_{st} of boreal forest LUCs to 3000, all of which were necessary in order to achieve realistic NH_3 concentrations ~~(e.g., compared to reported AMoN values)~~, while staying consistent with Γ findings from the literature.

This version of the model, which we call GEM-MACH-Bidi (or just "bidi" hereafter) was quite sensitive to the selection of emission potentials, which are themselves highly uncertain (Wen et al., 2014). GEM-MACH-Bidi uses the exact same emissions as in the base case, described in the previous section. However, when the sign of F_t in Eq. (1) becomes positive (that is, when $C_a < C_c$), the bidirectional flux acts effectively as an additional source of NH_3 gas, releasing stored NH_3 until and unless the ambient concentration rises to the compensation point concentration. When the flux is negative, net deposition of NH_3 occurs.

It is important to note that $C_{st,g}$ values are exponentially dependent on temperature (Fig. 1 shows an example of this relationship for the dominant LUCs in the northern part of the domain), and the higher the compensation point is, the greater the likelihood there will be upward flux. The lower $C_{st,g}$

is, the more likely there will be deposition. Since our simulation occurred in August and September 2013, when the average temperature in the AOSR was about 18°C (agriculture.alberta.ca/acis/alberta-weather-data-viewer.jsp), we expect to have more NH₃ re-emission than at other times of the year. During the rest of the year (e.g., the preceeding winter and spring), the compensation point would be much lower, greatly increasing the likelihood to have net deposition, even in northern Alberta/Saskatchewan where ambient NH₃ concentrations are low. Other meteorological factors affect the magnitude of bidirectional via the resistance terms. For example canopy compensation points have been observed to decrease with decreasing wind velocity, and increased precipitation (Flechard and Fowler, 1998; Fowler et al., 1998; Biswas et al., 2005; Zhang et al., 2010). In other words, we expect more re-emission during higher winds and drier conditions.

Other chemical transport models, such as GEOS-Chem and CMAQ use a similar method as Zhang et al. (2010), however, instead of the constant average soil emission potentials used here, they utilize a CMAQ-agroecosystem coupled simulation to calculate a soil pool from which to estimate Γ_g (Bash et al., 2013; Pleim et al., 2013; Zhu et al., 2015). In this case, the emission potential will vary and can go to zero if the NH₄⁺ in the pool is depleted. However, it was shown in Wen et al. (2014) that their $\Gamma_{st,g}$ worked well during the same time of year as this investigation (August and September). This time of year was also shown in Zhu et al. (2015) to not have a large effect on emissions from the NH₄⁺ pool. ~~Our investigation also has a short time scale of about a month, thus it is expected that the the soil pool would not be depleted over this time period~~ Additionally, Wentworth et al. (2014) calculated the approximate relative abundances of NH_x in the boundary layer versus NH₄⁺ in the soil pool to assess whether surface-to-air fluxes were sustainable. They found that soil NH₄⁺ were much greater than boundary layer NH_x (by over two orders of magnitude), further supporting the assumption made here. In addition, the turnover time for soil NH₄⁺ is on the order of one day, hence it is unlikely that NH₃ bi-directional fluxes would significantly deplete/enhance soil NH₄⁺ pools. Finally, given that GEM-MACH is used for real-time air quality forecasts at Environment and Climate Change Canada, it is not desirable for our bidirectional flux scheme to have to rely in advance on another model's output. Therefore, we use this simplified version, and assess whether its results provide ~~a good enough improvement to an~~ improvement (smaller biases and better correlations to measurements) to simulated NH₃ for less cost in run time.

2.3 Addition of forest fire emissions

Our third model scenario (called "fire+bidi" hereafter) uses the GEM-MACH-Bidi model, and the exact same area emissions and anthropogenic major point emissions as the base and bidi scenarios. However, in addition, we add hourly North American forest fire emissions for all species to the major point emissions. The forest fire emissions system for GEM-MACH (called "Fire-work") is described in detail in Pavlovic et al. (2016). Briefly, to calculate the fire emissions for input to FireWork, biomass burning areas are first identified in near real time by the Canadian Wildland Fire Information System (CWFIS), which is operated by the Canadian Forest Service

(<http://cwfis.cfs.nrcan.gc.ca/home>). CWFIS uses fire hotspots detected by NASA's Moderate Resolution Imaging Spectroradiometer (MODIS) and NOAA's Advanced Very High Resolution Radiometer (NOAA/AVHRR) and Visible Infrared Imaging Radiometer Suite (VIIRS) imagery as inputs. Daily
270 total emissions per hotspot are then estimated by the Fire Emission Production Simulator (FEPS) module of the BlueSky Modeling Framework (Larkin et al., 2009). SMOKE was then used to prepare model-ready hourly emissions of several species (including NH_3) in a point-source format for model input.

In Environment and Climate Change Canada's operational forest fire forecasts, these emissions are
275 used at 10-km resolution for the domain encompassing North America, with forest fires being treated as point sources with specific plume rise (Pavlovic et al., 2016). We have added 2013 forest fire emissions which were originally created for the 2013 Firework forecasts to the anthropogenic point source emissions used in the base case simulation, and modified the GEM-MACH model to be able to accommodate the changing number of major point sources each day (as the fires are parametrized
280 as major points, and their number changes daily). Fire plume rise is an ongoing area of investigation (e.g., Heilman et al., 2014; Paugam et al., 2016); smoldering emissions tend to be emitted directly at the surface, whereas flaming emissions can inject plumes to the upper troposphere. Here, we have set all fire emissions to be distributed evenly throughout the boundary layer, which is a simplification, but one that averages out smoldering and flaming plume heights. Different parameterizations of
285 fire plume rise are currently under development in GEM-MACH. The Fireworks fire emissions are described in detail in Zhang et al., (2017, this issue), and this study represents the first time they have been used at a 2.5-km horizontal resolution.

2.4 Model setup for three scenarios

The base, bidi, and fire+bidi models were all run with the following input files: Analysis files, which
290 are the products of meteorological data assimilation and provide optimized initial conditions for the 12 UTC hour of each day, were obtained from ECCO archives (Buehner et al., 2013, 2015; Caron et al., 2015), and the numerical weather prediction regional GEM model was run regionally at 10-km and the high resolution GEM model was run at 2.5-km resolution to produce meteorological files to drive the model simulation. The base, bidi, and fire+bidi scenarios were run from 1 August to
295 7 September, 2013, where the first 11 days were "spin up" in order to allow chemical concentrations to stabilize, and are not used in our evaluation. This is a sufficient amount of spinup time, given that the atmospheric lifetime of NH_3 is typically up to 1 day (Seinfeld and Pandis, 1998; Aneja et al., 2001), and given that it is close to the transport time of air crossing the larger North American domain. The time period ~~of~~ from 12 August to 7 September was chosen to coincide with the intensive
300 measurement campaign described in Section 3.

The model was run in a nested setup, whereby the North American domain was run at 10-km resolution using "climatological" chemical initial and boundary conditions from a 1-year MOZART

simulation for all pollutants (Giordano et al., 2015). The nested Oil Sands region (which covers most of Alberta and Saskatchewan) was run at 2.5-km horizontal resolution, using the initial and boundary conditions from the 10-km North American model run. Figure 2 shows the two model domains.

The model simulations for the pilot and nested domains were not run as a continuous multiday forecast, but rather following to the operational air quality forecast process, where the meteorological values are updated regularly with new analyses, in order to prevent chaotic drift of the model meteorology from observations. Consequently, our simulation setup comprises simulations on the North American domain in 30-hour cycles starting at 12 UTC, and the Oil Sands domain in 24-hour cycles starting at 18 UTC (the 6 hour lag being required to allow meteorological spinup of the lower resolution model). The next cycle uses the chemical concentrations from the end of the last cycle as initial conditions for the next 24-30 hours. This system of staggered meteorological driving forecasts with a continuous concentration record continues until the full time period completes.

We run GEM-MACH in the 2-bin particle mode, which means that particles fall in either fine mode (diameter 0-2.5 μm) or coarse mode (diameter 2.5-10 μm), for computational efficiency (although sub-binning is used in some particle microphysics processes in order to ensure an accurate representation of particle microphysics (Moran et al., 2010)), and in order to follow the setup used for the operational 10-km resolution GEM-MACH forecast.

3 Measurements

Our three model simulations (base, bidi, and fire+bidi) are evaluated with surface, aircraft, and Cross-track Infrared Sounder (CrIS) satellite measurements. We briefly describe each of these observations below.

3.1 AMS13 ground measurements

An extensive suite of instrumentation was deployed at monitoring site AMS13 (57.1492°N, 111.6422°W, 270 m.a.s.l., Fig. 3) from 7 August 2013 until 12 September 2013. Mining operations and bitumen upgrading facilities are 5 km to the south and north of the site. It is surrounded by boreal forest, with dominant winds from the west, averaging 1.9 m/s throughout the year. The average temperatures in the region for August are highs in the low 20°C, and lows around 10°C, which is warm enough to make upward NH_3 flux more likely (recall Fig. 1), but temperatures drop rapidly at the end of August, into September, where the September highs average around 15°C, lows around 5°C. The skies are the clearest during August, with at least partly clear skies 50% of the time. That said, the warm season (May through September) is the wetter season (20% chance daily average), with more precipitation than during the cold season (7% chance daily average), but year round precipitation, as well as relative humidity are both relatively low in the AOSR. During the cold season (November through February), the average temperatures range from -21°C to -5°C, when the forest and soils are

more likely to be a deposition sink for NH_3 . During November to April, it is also much cloudier, with February having cloudy conditions 77% of the time. (All weather data cited here are from the annual report at Fort McMurray, found here: <https://weatherspark.com/y/2795/Average-Weather-in-Fort-McMurray-Canada-Year-Round>).

340

NH_3 , fine particulate ammonium and nitrate, and other species were measured by the Ambient Ion Monitor-Ion Chromatograph (AIM-IC), via an inlet 4.55 m off the ground (Wentworth et al., 2017). The uncertainty of these measurements is $\pm 15\%$. These measurements appear in this issue (Wentworth et al., 2017), and are described in more detail therein Markovic et al. (2012).

345 Data gaps sometimes appeared in the surface NH_3 time series for the following reasons: instrument zero (Aug 14/15 and 17/18), instrument maintenance (Aug 19) and a power outage (Aug 27/28).

3.2 Aircraft measurements

During the Oil Sands Monitoring Intensive campaign, there were a total of 22 flights spanning 13 August to 7 September 2013. These measurements are described in detail in Shephard et al. (2015); 350 Gordon et al. (2015); Liggio et al. (2016); Li et al. (2017), and are summarized here. Aircraft NH_3 measurements were conducted with a dual quantum cascade laser (QCL) trace gas monitor (Aerodyne Inc., Billerica, MA, USA; McManus et al., 2008), collecting data every 1 s. Outside air was sampled through a heated Teflon inlet tube shared with a high-resolution time-of-flight chemical 355 ionization mass spectrometer (HR-ToF-CIMS); the flow rate through the QCL was 10.8 L min^{-1} . The 1σ uncertainty for each measurement is estimated to be $\pm 0.3 \text{ ppbv}$ ($\pm 35\%$) (Shephard et al., 2015).

Particulate NH_4^+ and NO_3^- ($0 < 1 \mu\text{m}$ in diameter) were measured by the Aerodyne high-resolution time-of-flight aerosol mass spectrometer (HR-ToF-AMS) instrument on board the same flights, 360 which collected data every 10 s. The ambient air was drawn through a forward facing, shrouded isokinetic particle inlet from which the HR-ToF-AMS sub-sampled. The total residence time in the inlet and associated tubing was approximately 1 second. The error on these measurements is $\pm 9\%$. (Liggio et al., 2016)

Figure 3 shows a sample flight path from the campaign from 13 August 2013 – one of the thirteen 365 flights with valid NH_3 measurements. The others took place on 15-17, 19 (two this day), 22-24, 26, 28 August, and 5-6 September 2013. NH_3 data on the other nine flights were invalidated due to instrument issues (those on 14, 20-21, 29, 31 August, and 2-4 September 2013), but were successful for the NH_4^+ and NO_3^- measurements.

3.3 CrIS satellite measurements

370 CrIS was launched in late October 2011 on board the Suomi NPP platform. CrIS follows a sun-synchronous orbit with a daytime overpass time at 13:30 (ascending) and a night time equator over-

pass at 1:30 (descending), local time. The instrument scans along a 2200 km swath using a 3 x 3 array of circular pixels with a diameter of 14 km at nadir for each pixel. The CrIS Fast Physical Retrieval (CFPR) described by Shephard and Cady-Pereira (2015) is used to perform satellite profile retrievals of ammonia volume mixing ratio (VMR) given the infrared emission spectrum from the atmosphere. This retrieval uses an optimal estimation approach (Rogers, 2000) that provides the satellite vertical sensitivity (averaging kernels) and an estimate of the total errors (error covariance matrix).

We take the CrIS retrieved profile and match it up with the closest model profile in both distance and time. ~~Compute~~, compute the distance between the CrIS pixel and model field for each time step, and then select the time step that best matches the satellite overpass time. Since the model time steps are every hour with a 10-km spatial resolution they are always matched up to better than half an hour, and within 5 km. ~~Figure 10 shows a map with four boxed regions indicating where model-measurement pairs were sampled for this study. These and the corresponding satellite fields will be discussed later in the model evaluation section below.~~

4 Model evaluation

An older version of GEM-MACH (v1.5.1) has been compared to TES satellite and aircraft measurements of ammonia over the AOSR (Shephard et al., 2015). Simulations with that version of the model were shown to be biased low, by about -0.5 ppbv, throughout the lower-tropospheric vertical profile. This represented a substantial deficit in the model predicted sources of NH_3 , prompting the current work. We now compare our three GEM-MACH simulations (base, bidi, and fire+bidi) against surface point measurements at the measurement site near an oil sands facility (AMS13), aircraft measurements over the broader AOSR, and satellite measurements over the Alberta and Saskatchewan area. We will discuss which simulation agrees best with measurements and where there may still be room for additional model improvement.

4.1 At the AMS13 ground site

Figure 4 shows the timeseries of the daily average (for clarity) concentrations of NH_3 and ~~its reaction products~~, fine-particulate NH_4^+ ~~and~~, NO_3^- , ~~as well as its aerosol sink and~~ SO_4^{2-} at the AMS13 Oil Sands ground site for the observations and three model simulations. ~~Note that the y-scale is logarithmic to better show the differences between the three model scenarios and the measurements – however, this means that when concentrations drop to zero in the model, the line becomes disjointed. This is mainly evident in the NO_3^- time series (Fig. 4c). Figure 4~~ The hourly data were also studied, but not shown in the time series.

We first note that the NH_3 concentrations seen in the measured time series are relatively low with mean, median, and maximum of 0.6 ppbv, 0.426 ppbv, and 2.98 ppbv, respectively in the hourly data,

which are lower than the 1-8 ppbv range in Bytnerowicz et al. (2010), and the 2.7 ppbv summertime mean given in Hsu and Clair (2016). However, this may be due to the different time periods and locations measured. Our mean measured values at the AMS13 site are similar to the concentrations found at U.S. AMoN background sites (<http://nadp.sws.uiuc.edu/amon/>).

410 Figure 4a shows that the base model (green) background concentrations of NH_3 are ~~biased-very~~ low (nearly 0 ppbv when there is no plume influence) compared to the measurements (orange). Only during the spike on September 3-4th does the base model exceed the measured values, probably indicating a local plume event fumigating to a lesser extent in the observations than was assumed in the model. The NH_3 concentrations of the base case are biased low compared ~~to~~ the surface
415 measurements by a median of -0.35 ppbv (Fig. 5a) over the time period of the campaign – comparable to the bias observed in satellite observations in Shephard et al. (2015). ~~The model-measurement differences are shown in Fig. 5a. In Fig. In Figure~~ 4, the bidi model (blue line) and fire+bidi model (red line) show a significant improvement to the NH_3 concentrations ~~during the background times,~~ compared to the base ~~case~~-model (green line). Unfortunately, during some time periods, these two
420 versions of the model overestimate NH_3 : During August 13th, the model adds a significant level of NH_3 due to fire emissions, however the surface *in situ* observations show no evidence of fire impact. During other time periods (e.g., 30 August to 3 September, and 4-7 September), the bidi model appears to have put too much NH_3 into the system. Therefore, the bidi model bias (Fig. 5a) is now 0.30 ppbv too high (median), and the fire+bidi bias is 0.32 ppbv high (median) over the time period of
425 the campaign, resulting in an overall improvement of only 0.03 ppbv in the model bias.

~~When the influence of local plumes (defined as the base model exceeding 0.5 ppbv) is removed from the time series, the median biases are as follows: -0.36 ppbv for the base model, +0.26 ppbv for the bidi model, and +0.28 ppbv for the fire model – corresponding to an overall improvement of 0.08 ppbv in model bias for background concentrations.~~

430 While the bias improvement is small, the bidi and fire+bidi both have greatly improved correlation coefficients (from $R=0.2$ ~~0.1~~ to 0.4) and slopes much closer to 1 (from ~~0.035 to 0.614~~ ~~0.1~~ to 0.7), showing that those added sources are important to improve model results (Fig. 6a). ~~The correlation coefficients and slopes were similarly improved when the plume influence was removed, and only background-level concentrations were examined. Additionally, the diurnal cycle (not shown) was~~
435 ~~improved in the bidi simulation, with both it and the measurements shaped like a sine curve having a minimum at 3:00-4:00am local time, and a maximum at noon local time, although the amplitude of the cycle was underestimated. Whereas, the base model diurnal cycle was flat from midnight to noon local time, and spikey from noon to midnight.~~

While Fig. 4a to 6a show that the addition of bidirectional flux significantly improves the model
440 correlation coefficient, slope, and bias, there is still room for improvement. [Paired t-test results indicate that the fire+bidi and measurements are still significantly different \(see Table 2 for comparison statistics of all three simulations\).](#) While inherent limitations from model resolution and uncertain-

ties may be responsible for the remaining bias, it is likely that (a) the emission potentials for the land use categories (LUCs) in the region may be causing too much re-emission of NH_3 , and need refinement, and (b) the fire emissions of NH_3 are not properly distributed in the vertical, placing too much NH_3 near the surface and/or the fire emission factors for NH_3 are too high.

Refinement needed for the emission potentials and LUCs may be a significant cause of the bidi and fire+bidi model biases. Rooney et al. (2012) have shown that about 64% of the AOSR are wetlands (fens, bogs and marshes), which ~~should~~ should be mapped to the swamp LUC. However, our model currently assigns the AOSR landscape to evergreen needleleaf trees, deciduous broadleaf trees, inland lake, mixed shrubs, and mixed forests (and none of the region to swamp). This would lead to an overestimation of re-emission given that bogs are fairly acidic and our swamp emission potential is lower than the aforementioned LUCs. Other evidence for these two explanations will be presented below in Section 4.3.

The time series, model-vs-measured correlations, and model biases of NH_4^+ , NO_3^- , and SO_4^{2-} are also shown in Fig. 4 to 6 (b, c, and d, respectively). For NH_4^+ and SO_4^{2-} there is very little change despite the increase in NH_3 that the bidirectional flux yields. The bias is very small for all three model scenarios, and the correlation coefficients are all relatively poor. So while there is an improvement to modelled NH_3 with bidirectional flux, there is a neutral affect on fine particulate NH_4^+ . This may be because the charge of NH_4^+ in the particles is already enough in the base model to balance the charge of $2 \times \text{SO}_4^{2-} + \text{NO}_3^-$ in the aerosols, thus, causing any additional NH_3 (from bidi and fires) to remain in the gas phase. Or it could be due to additional wet scavenging of the additional ammonium, which will be discussed in Section 5.2. The change in NH_3 concentrations has no effect on SO_4^{2-} since particulate SO_4^{2-} is not sensitive to the amount of $\text{NH}_3/\text{NH}_4^+$ available, and is dominated by anthropogenic and fire emissions. For NO_3^- , the base model bias was quite small at $0.01 \mu\text{m}/\text{m}^3$, however the addition of bidi and fire+bidi further reduced that bias to 0.0011 and $0.0004 \mu\text{m}/\text{m}^3$, respectively, which is a significant improvement. The correlation coefficient for NO_3^- also improved from about 0.1 to 0.3 (Fig. 6c).

4.2 Along the OS campaign flight paths

There were 13 flights during the OS campaign that had valid (above detection limit, and no instrument error) NH_3 measurements, and 22 flights that had valid NH_4^+ ($0-1 \mu\text{m}$ diameter) measurements. The flight path of the first flight, which occurred on 13 August 2013 is shown in Fig. 3; chosen as an example because this flight sampled mainly background NH_3 concentrations (rather than facility plumes).

Figure 7 shows the NH_3 concentrations along this flight path over time. Here the hourly model output is interpolated to the same time frequency as the measurements. The model also has spatial resolution limits when comparing to the aircraft. However, we clearly see that for this flight, the bidirectional flux has increased NH_3 concentrations, bringing them closer to the measured val-

ues ~~-(median biases for this flight are -1.38, 0.68, and 0.69 ppbv in the base, bidi, and fire+bidi~~
480 ~~simulations). There is little change when fires are added (Fig 7d vs c) because this flight did not pass~~
~~through a fire plume.~~

Figure 8 shows the ~~corresponding~~-model-measurement ~~difference~~-differences and the model vs
measurement scatter plots for the combined set of all flight paths for hourly-average concentra-
tions of NH_3 and NH_4^+ . For NH_3 the median base model bias is -0.75 ppbv, comparable to the
485 bias observed in Shephard et al., 2015, with the bidi model bias improving to -0.24 ppbv, and
the fire+bidi bias to -0.23 ppbv. Also the best correlation coefficient and slope is achieved by the
fire+bidi scenario. The use of the bidirectional flux has thus reduced the model bias relative to the
aircraft observations by a factor of three. The fire+bidi simulation has the best statistics compared to
measurements, as summarized in Table 2.

490 Again, the NH_4^+ results show little change despite the increase in NH_3 concentrations. The small
bias from the base case gets insignificantly smaller, and the slope and correlation coefficients are all
negligibly changed.

4.3 In the vertical profiles across the region

The CrIS satellite has many observations over North America during the 2013 Oil Sands campaign.

495 We have evaluated the model with these observations in a number of ways:

1. ~~Daytime-overall-average-in-this-area~~ All daytime data from Aug 12 - September 7th, 2013;
model-measurement comparisons over a large region encompassing Alberta and Saskatchewan (~~blue~~
~~box in Fig. 10~~, latitude range: 48-60 °N, longitude range: 100-122 °W), which contains agri-
cultural areas, a number of cities, the northern boreal forest, and the Oil Sands facilities.
- 500 2. Case studies where we attempt to isolate fire emissions (~~magenta box in Fig. 10~~), and non-fire
conditions (~~cyan and black boxes in Fig. 10~~) to evaluate both new components (fires and bidi)
of the model.

The latitude and longitude ranges of our model-measurement pairs are given in Table 1. The satellite
passes over these regions at approximately 1pm and 1am local time.

505 ~~In the large box spanning northern Alberta and Saskatchewan, there~~ There were over 60 000
model-measurement pairs between the model and the CrIS satellite over the model domain during
August 12th to September 7th, 2013. Figure ~~10 shows the surface NH_3 concentrations over that~~
~~region on three sample days (3 Sept, 1 Sept and 12 Aug, 2013), as well as sample Aqua-MODIS true~~
~~colour composite map for that day. Figure 9 presents statistics~~ 9top panel presents model biases for
510 the entire dataset in a box and whiskers plot of the vertical NH_3 profiles at five vertical levels, ~~along~~
~~with the model-satellite bias for the base model, bidi model, and fire+bidi model. The~~. The left-most
panel shows the NH_3 concentrations measured by CrIS, and the right-most panel shows the diagonal
elements of the CrIS averaging kernels, illustrating the sensitivity of the satellite measurements to

each vertical level. The NH_3 concentrations over Alberta and Saskatchewan measured by CrIS are very similar to those found by TES in the Shephard et al. (2015) study for the AOSR region.

In Fig. 9 we see that the middle panels show the model biases from the three simulations. The fire+bidirectional model has the smallest bias in the highest three layers, but the bidirectional model has the smallest bias in the two lowest layers, whereas the fire+bidirectional model increases NH_3 concentrations further (though still a smaller absolute bias compared to the base case, Fig. 9top). This could be due to an overestimate of the bidirectional flux re-emissions or of the fire emissions, or to an underestimate of the altitude of the fire emissions, or a combination of all three factors. In order to distinguish between these possibilities, two case studies were examined further below. The statistics from the model-CrIS comparison can be found in Table 2. That summary shows that the fire+bidirectional simulation performs better than the base and the bidirectional simulations.

The spatial distribution of modelled NH_3 can also be evaluated with CrIS measurements, as shown in the bottom panel of Figure 9. These are maps of the average surface NH_3 concentrations from the base model, the fire+bidirectional model, and the CrIS satellite. The fire+bidirectional model over-predicts the effect of fires in the middle of northern Saskatchewan, but appears to be missing fires in north-western Manitoba. Other than fire influence, the spatial distribution in the fire+bidirectional model is the same as that of the base model, but with significant increases in overall concentrations. And the spatial distribution of the model simulations is different from the spatial distribution that CrIS measures. For example, the model predicts much higher NH_3 concentrations near the city of Edmonton than CrIS shows. That said, the addition of bidirectional flux has greatly improved the NH_3 concentrations in the northern part of the province, where there was almost none in the base model.

We selected three sample days (3 Sept, 1 Sept and 12 Aug, 2013) that we use for the case studies, and surface NH_3 concentrations over that region as well as sample Aqua MODIS true colour composite maps for those days are shown (Fig. 10). The four boxed regions on those maps indicate where model-measurement pairs were sampled for this study. The cyan and black boxes in Fig. 10a and b are the regions where we sample clear-sky, no-fire conditions on 3 and 1 September 2013, respectively. The magenta box in Fig. 10c is the region where we isolated our fire case study on 12 August 2013. The blue box is the region we discussed above, which we analysed for the full time period simulated (12 Aug - 7 Sep 2013, Fig. 9top).

4.3.1 Case study 1: clear-sky days with little fire influence - evaluating bidirectional

In order to evaluate the bidirectional flux component separately from the fire component, we selected September 1st (southern, agricultural region - black box in Fig. 10b), and 3rd (northern, boreal forest and AOSR region - cyan box in Fig. 10a), where the MODIS map (EOSDIS NASA World view map, worldview.earthdata.nasa.gov) shows very little hot spots from fires, and that the conditions were relatively cloud and smoke free (which yield the most CrIS observations). See Table 1 for the latitude and longitude ranges. Figure 10 also shows the surface NH_3 concentrations as observed by

CrIS on each of those days. Figure 11a shows that in the north, the bidi model improves the bias from -0.84 ppbv to -0.07 ppbv in the lowest vertical level, and smaller, but still significant, improvements to the bias at the other levels. The fire+bidi model has a nearly identical impact as the bidi model, which is expected in a fire-free zone. Therefore, the GEM-MACH-Bidi model performs very well in northern Alberta and Saskatchewan where there is mainly boreal forest, and background-level NH_3 concentrations. This also implies that the LUC assignment discussed in Section 4.1 may only apply to a small region around the AOSR, and not to the overall large region we've defined here.

In the southern region (Fig. 11b), the addition of bidirectional flux moves the bias from near-zero to +1.02 ppbv in the lowest level. In this case, the base model with no bidirectional flux appears to be the most accurate model in areas dominated by agricultural sources. There are two possible explanations: a) agricultural emissions are too high in the base model, and the addition of the bidirectional flux leads to an overestimation of the NH_3 amounts, or b) re-emissions from bidirectional flux from crops are not significant. The literature (Bash et al., 2010; Massad et al., 2010; Zhang et al., 2010; Zhu et al., 2015) indicate that crops do indeed re-emit NH_3 , therefore, (a) is the more likely explanation. The agriculture NH_3 emission inventory we used was created by the NAESI (National Agriculture Environmental Standards Initiative) project (Bittman et al., 2008; Ayres et al., 2009; Makar et al., 2009) have about 30-200% uncertainty associated with them (Bouwman et al., 1997; Asman et al., 1998). Therefore, with improved national NH_3 emission inventories, the GEM-MACH-Bidi should improve model results across the domain.

4.3.2 Case study 2: a clear day with significant fire influence - evaluating fires

In order to evaluate the fire component separately from the bidirectional flux, we selected August 12th (a northern region with little-to-no agricultural contributions) where the MODIS map shows numerous hot spots from fires and smokey conditions (Fig. 10c, magenta box). The base and bidi models underestimate NH_3 concentrations (Fig. 11c) by -6.22 and -5.84 ppbv, respectively (in the lowest vertical layer), but ~~that~~ the fire+bidi model overestimates NH_3 by +4.06 ppbv. The fire+bidi version of the model still has the lowest bias of the three simulations, however, either (a) the fire+bidi model does not distribute the fire emissions properly in the vertical, (b) the fire emissions of NH_3 are too high, and/or (c) the model is not properly representing NO_2 and SO_2 in the fire, and so the conversion of NH_3 to NH_4^+ is underestimated. ~~Both~~ It is potentially a combination of all three explanations, and we further elaborate below.

- For explanation (a), both fire plume rise and fire emission factors are on-going areas of study. ~~It is likely a combination of all three explanations; in~~ In the model the fire emissions are distributed evenly ~~through-out~~ throughout the boundary layer (the first 3-4 layers ~~from~~ in Fig. 11c), however, Shinzuka et al. (2011) suggest that sometimes the fire plumes are ~~in a Gaussian distribution located in a distributed normally in a~~ thin layer aloft ~~(which is an option for GEM-MACH that is currently under development)~~. However, should that be the case, ~~the~~

~~bias would move for the real-life fires in this case study, the model bias would be~~ negative at at least one of the levels in Fig. 11c, which it ~~does not (unless the fire plume is actually above 4 km, however, it was found that the plume heights for the Fort McMurray fires of 2016 reached only up to 3-3.5 km, shown by the lidar on CALIPSO satellite)~~is not. Figure 11c shows that the positive bias extends throughout the first three vertical layers, and in the top two vertical layers, the bias does not move further negative (as would happen in the fire plume were actually at those altitudes in real life).

- ~~(a) Our bias would be very high at low levels if the real fire plumes were actually above 4 km (above the altitudes we studied), however, it was found that the plume heights for most of the Fort McMurray fires of 2016 reached only up to 3-4 km altitude range based on the NASA Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) and Multi-angle Imaging SpectroRadiometer (MISR) satellite observations. Therefore, this is not likely the case.~~
- Explanation (b) seems the most likely, as the uncertainty on emission factors for NH₃ from wildfires is very large (e.g., 50-100% depending on the fuel type Urbanski, 2014), and could easily be overestimated. ~~Similarly, the NO_x~~
- ~~The NO_x~~ and SO₂ ~~emission factors have fire emission factors (c) have smaller~~ uncertainties of 10-40% (Urbanski, 2014).

Unfortunately, there were no flights that captured the fine structure of the fire plumes during the 2013 monitoring intensive campaign that can be used to further corroborate the vertical distribution of the fire plumes. There will however be flight observations of fires during the planned 2018 AOSR measurement campaign. Therefore, the model may be further improved with reduced NH₃ emission factors for fires, and/or improved vertical distribution of fire plumes.

5 Impacts of bidirectional flux and forest fires on NH₃ concentrations

5.1 Effect on ambient concentrations

Given that the overall fire+bidi model agrees best with measurements in the greater Alberta/Saskatchewan region (discussed throughout Section 4) and contains all known missing sources of NH₃, we can use the model to answer one of our key questions: What percent contributions to total ambient NH₃ concentrations came from bidirectional flux and from forest fires during the study time period? We do so by subtracting the bidi model output from the fire+bidi model output to get the forest fire component, and subtracting the base model output from the bidi model output to get the bidi component. The absolute differences are calculated as follows:

$$bidicomponent = NH_3^{bidi} - NH_3^{base} \quad (5)$$

$$620 \quad firecomponent = NH_3^{fire+bidirectional} - NH_3^{bidirectional}, \quad (6)$$

which tell us how many ppbv of NH_3 on average comes from re-emissions of NH_3 (upward component of bidirectional flux), and from fire emissions.

The percent differences are calculated as follows:

$$bidipercents = \frac{NH_3^{bidirectional} - NH_3^{base}}{NH_3^{fire+bidirectional}} \times 100\% \quad (7)$$

$$625 \quad firepercent = \frac{NH_3^{fire+bidirectional} - NH_3^{bidirectional}}{NH_3^{fire+bidirectional}} \times 100\%, \quad (8)$$

which tell us what percent of ~~total~~ total NH_3 concentrations on average comes from re-emissions of NH_3 (upward component of bidirectional flux), and from fire emissions, assuming the NH_3 from our fire+bidirectional simulation is the true total NH_3 .

630 ~~Over the 2.5-km model domain (averaged over~~ We perform this calculation on the averaged model output (12 August to 7 September 2013) , we do this calculation (See Fig. 12) over the 2.5-km model domain, and get an average of 20.3% (or 0.42 ppbv) ~~of~~ and a median of 10.4% for ambient surface NH_3 concentrations ~~comes that come~~ from forest fires —though the median amount is only 10.4% for fires. (Fig. 12). The mean and median are so different because fires are sporadic,
635 large contributions to NH_3 concentrations, and the mean value is more sensitive to the big outliers. We get an average of 56.6% (or 1.24 ppbv) from bidirectional flux (56.3%~~median~~). ~~The median), and the~~ remaining 23.1%~~(average)~~ , average (33.3 % (, median) comes from direct emissions from anthropogenic sources (agriculture, fossil fuel combustion, oil sands industry, etc). These numbers are summarized in Table 3. The increase in NH_3 concentrations due to the bidirectional flux scheme is of the same order of magnitude as that found in the Zhu et al. (2015) study using the GEOS-Chem model, during the month of July, in the United States (where they found 1 ppbv increase in surface concentrations due to bidirectional flux). It is also similar to values found in Europe in the Wichink Kruit et al. (2010) study.

Over the model domain, the minimum bidi influence on surface NH_3 is just north of Edmonton,
645 where only 1% of NH_3 comes from bidi. Similarly, two AOSR facilities north of Fort McMurray stand out as having small bidi influence (12-40 %, surrounded by values in the 90s% - Fig. 12,d). Also, any remote region with fire emissions will have a small percentage contribution from bidirectional flux during the fires, as they are in northern Saskatchewan (Fig. 12,d). This is expected given that the average concentrations in cities and near large sources are very close to, or exceed the compensation point. The absolute maximum in the bidi component map is 4.5 ppbv in the lower right corner (an agricultural region with high NH_3 emissions), and the minimum is 0 ppbv (Fig. 12,b). This means that nowhere in the domain, did the bidirectional flux formula result in ~~more~~ more net deposition than the base model calculated via the Welesley/Robichaud/Zhang scheme. The maximum fire

contribution to surface NH_3 is 27.9 ppbv where large fires occurred in northern Saskatchewan (Fig.

655 12,c).

5.2 Effect on Deposition

Similar to our analysis from the previous section, we can use the model to determine how bidirectional flux and fires impact daily NH_3 ~~dry deposition~~ deposition (which equals the dry deposition of NH_3 + the wet deposition of NH_4^+). Figure 13 shows the average daily net ~~dry~~ deposition (or net flux) of NH_3 ~~from the base, bidi, and fire+bidi models~~. Negative (or blue) indicates net deposition (downward flux), and positive (or red), net emission (upward flux). The base model (Fig. 13a) had no re-emission (upward flux) option, thus NH_3 was always net dry deposited in that scenario, ~~and was 9.85×10^{-6} moles/m²/day on average~~. The bidi (Fig. 13b) and fire+bidi (Fig. 13c) maps show that ~~over much most~~ of the Alberta and Saskatchewan area, ~~there is net emission of NH_3 , and net deposition only occurs where concentrations are highest~~ (has net deposition (e.g., near the cities, agriculture, and forest fires)). ~~Average flux has changed to~~, but that some regions (with low atmospheric NH_3 concentrations) have net emission of NH_x . The dry NH_3 flux is net positive over the domain, ~~with averages of $+2.44 \times 10^{-5}$ moles/m²/day and $+2.10 \times 10^{-5}$ moles/m²/day~~ for however, when the increase in wet NH_4^+ is accounted for, the net flux of NH_x is still negative (downward). This is very similar to what Wichink Kruit et al. (2010) found in a 2007 study in Europe; a reduction in dry NH_3 , compensated by an increase wet NH_4^+ deposition.

Note that the bidi and fire+bidi cases, respectively. Note that this result is partly due to our assumption of an infinite soil pool of NH_4^+ . Following the in our bidirectional flux scheme has not caused an overwhelming upward flux of NH_x . In fact, the average results across the domain actually have more deposition in the fire+bidi scenario than in our base scenario. Table 4 shows the mean and median net NH_x flux for each scenario (presented as deposition, so negative signs removed). That said, following the soil pool approach (Pleim et al., 2013; Zhu et al., 2015), the soil pool of NH_4^+ would may eventually get depleted, thus the average net flux may not be as largely positive as we calculated in this study. However, that is very unlikely for the following reasons: (1) Deposition of NH_x throughout the year continually replenishes the soil pool – especially when temperatures are cooler in winter, spring, and fall, since the compensation point is exponentially dependent on temperature. (2) The short time frame of this study would not be long enough to deplete the soil pool. For example, Zhu et al. (2015) needed to spin up their model for three months in order to get the NH_4 soil pool stable.

685 In the AOSR near Fort McMurray, we can compare our NH_3 dry deposition results to those calculated in Hsu and Clair (2015). Their values range from 0.7 to 1.25 kg-N/ha/year (or 1.13 to 2.01×10^{-5} ~~mol~~moles/m²/day), and ours are $10 \times$ ~~times~~ lower at around 0.13 kg-N/ha/year (or 2.12×10^{-6} ~~mol~~moles/m²/day) near Fort McMurray, and do not vary much among our three model scenarios. Our deposition underestimate may be partially due to the fact that our modelled ambient NH_3

690 concentrations are also low compared to those measured in Hsu and Clair (2016) near Fort McMurray. They measured an average of 1.55 ± 0.6 ppbv ($1.9 \mu\text{g}/\text{m}^3$) at Fort McMurray, whereas our fire+bidi model has an average of 1.01 ppbv there (0.73 ppbv in bidi, and 0.39 in base). There may also be differences in that our model has more of the NH_x deposition coming down as NH_4^+ , rather than as NH_3 .

695 ~~Figure ?? (a) shows the difference in deposition between the bidi and base cases — essentially the contribution of bidi to the total flux. The bidi model has increased flux in the positive direction everywhere by an average of +3.43~~ Our fire+bidi NH_x deposition values (Table 4 are well in line with reported NH_3 deposition in Kharol et al. (2017), who report satellite-derived NH_3 deposition of about $2.1\text{--}7.0 \times 10^{-5} \mu\text{moles}/\text{m}^2$ across the domain. ~~Figure ?? (b) shows the /day in Alberta), and~~ are at the low end of NH_3 deposition values reported within Behera et al. (2013).

700 The difference in deposition between the fire+bidi and bidi cases — which is the contribution of fires to the total flux. The fires have decreased flux (in other words NH_x flux — showed that the fires increased downward flux, or deposition) /deposition over large swaths of the domain (e.g., difference between Figure 13c and b). The fires contributed an average of $-3.471.954 \times 10^{-6} \mu\text{moles}/\text{m}^2$ of NH_x deposition across the domain.

705 ~~The wet deposition of NH_4^+ was also modelled and analysed in a similar way (Fig. ??). Unlike NH_3 , since there is no re-emissions of NH_4^+ in the model, all flux is negative. While the particulate NH_4^+ concentrations did not change much in our three simulations (see Sections 4.1 and 4.2), the wet deposition of NH_4^+ increased significantly going from the base to bidi to fire+bidi models (Fig. ?? from a to b to c).~~ This is in contrast to what Zhu et al. (2015) found, which was little change to NH_4^+ due to bidirectional flux. However that could be due other parameters, such as the meteorological conditions, scavenging parameters, and/or gas-particle partitioning of NH_x . It would seem that in GEM-MACH-Bidi, the increased NH_3 concentrations were scavenged by precipitation. The average NH_4^+ deposition from the three simulations was: $-2.04 \times 10^{-5} \mu\text{moles}/\text{m}^2$ for base, $-4.25 \times 10^{-5} \mu\text{moles}/\text{m}^2$ for bidi, and $-5.86 \times 10^{-5} \mu\text{moles}/\text{m}^2$ for fire+bidi. That is nearly a had a nearly three-fold increase in the NH_4^+ deposition due to the increased NH_3 concentrations that the fire+bidi simulation yields. ~~Note that in the soil pool approach of Pleim et al. (2013); Zhu et al. (2015), this deposited NH_4^+ would contribute towards the soil pool, which could be re-emitted as NH_3 . In our case, once deposited, the NH_4^+ is not re-emitted~~ wet deposition for our fire+bidi simulation is $5.86 \times 10^{-5} \text{ moles}/\text{m}^2/\text{day}$, which is in between values reported in the United States in Stensland et al. (2000) (where they found an average of $1.9 \times 10^{-5} \text{ moles}/\text{m}^2/\text{day}$ over the country), and in Japan in Murano et al. (1998) (where they found an average of $10.3 \times 10^{-5} \text{ moles}/\text{m}^2/\text{day}$ over the country).

720 In the three scenarios, the average daily relative ratio of dry/wet deposition was: 0.43 for base, -0.77 for bidi, and -0.51 for fire+bidi (the negative value for the bidi and fire+bidi cases are because of the average upward direction of NH_3). Since all average ratios are less than 1, this means that most

of the removal process is from wet deposition, rather than dry deposition (even for the base case that had no re-emission of NH_3). Therefore, increased monitoring of wet deposition in the region would be useful. These results may also be useful for AEP terrestrial/aquatic scientists interested in nitrogen eutrophication. ~~The negative value for the bidi and fire+bidi cases are because of the average upward direction of NH_3 .~~ Maps of these ratios can be found in the supplemental material.

6 Conclusions

The GEM-MACHv2 air quality forecasting model was altered to include both the Zhang et al. (2010) bidirectional flux scheme for NH_3 and forest fire emissions of all species. This new “fire+bidi” model greatly improves the simulated NH_3 in the modelled Oil Sands domain at 2.5-km resolution when compared to independent *in situ* measurements at the ground (at the AMS13 oil sands monitoring site) and aloft (aircraft measurements), as well as at 10-km resolution when compared to ~~remote sensing cutting-edge satellite measurements~~ from the CrIS instrument. ~~We have also shown that for further improvements in the Alberta/Saskatchewan region, the NH_3 emission factors for fires, and the NH_3 emissions from agriculture likely need to be reduced.~~ Almost all comparison statistics are best with our fire+bidi simulation. This suggests that the fire+bidi model shows promise for improving NH_3 model predictions elsewhere and during other time periods. However, more work is required to validate the model in other regions of the continent (e.g., with the Wood Buffalo Environmental Association (WBEA) and the U.S. Ammonia Monitoring Network (AMoN) surface networks, and further CrIS satellite measurements), and for different time periods (e.g., springtime fertilizer season, cooler conditions, etc.). We ~~ve have~~ also shown that for further improvements in the Alberta/Saskatchewan region, the NH_3 emission factors for fires, and the NH_3 emissions from agriculture likely need to be reduced.

Despite the significant increase in NH_3 concentrations with these additional sources, the impact on its byproduct, NH_4^+ , was miniscule - as was the change to SO_4^{2-} concentrations ($0.02 \mu\text{g}/\text{m}^3$ for each). The model bias for those species was not significantly changed in either direction. This is probably because of the extra NH_4^+ wet scavenging by precipitation, and the NH_3 concentrations were already high enough (before adding the extra sources) to charge balance the SO_4^{2-} and NO_3^- in the aerosols. Thus, any additional NH_3 would remain in the gas phase. That said, the model bias for NO_3^- at the AMS13 ground station was essentially removed with the fire+bidi model.

By running the base, bidi, and fire+bidi model scenarios, and taking the fire+bidi results as “true”, we were able to calculate their differences and determine the average contributions from each source. We found that, on average, during the 12 August to 7 September 2013 time period in the Alberta/Saskatchewan model domain, 23.1% of surface NH_3 comes from ~~anthropogenic sources~~direct anthropogenic emissions, 56.6% of surface NH_3 comes from bidirectional flux (re-emission from soils and plants), and 20.3% of NH_3 comes from forest fires. Possible sources of error that remain

in the bidi and fire+bidi simulations are the agricultural and fire emissions of NH_3 , as well as the emission potentials for different land-use categories. The fraction of NH_3 from fires is highly variable depending on the time periods and spatial domain analysed: on average from 12 August to 7
765 September 2013, the largest impact was in northern Saskatchewan. We expect the re-emission source to be near the highest at this time of year because of the high temperatures, and this source should be much lower during the cold season, when deposition is expected to dominate the bidirectional flux process.

The bidirectional flux process has decreased NH_{3-x} deposition on average ~~,resulting in across the~~
770 ~~domain, with some areas having~~ a net emission of NH_3 ~~across the domain. This increase.~~ However, ~~that upward flux is~~ due to the ~~bidirectional flux, coupled with the increase driven by fires, enhances wet deposition of NH_4^+ by a factor of three~~ low atmospheric concentrations and high temperatures, and does not exceed the amount of NH_x deposition that occurs during the cooler winter and spring times. When fires are also taken into account, the net NH_x deposition is greater, on average across
775 ~~the domain, compared to the base model.~~

Acknowledgements. The project was supported by Environment and Climate Change Canada's Oil Sands Monitoring program (OSM), and the Climate Change and Air Quality Program (CCAP). We would also like to acknowledge the University of Wisconsin-Madison Space Science and Engineering Center Atmosphere SIPS team sponsored under NASA contract NNG15HZ38C for providing us with the CrIS level 1 and 2 input data,
780 in particular Keven Hrpcek and Liam Gumley.

Data availability: The CrIS-FRP- NH_3 science data products used in this study can be made available on request (M. W. Shephard, Environment and Climate Change Canada, Toronto, Ontario, Canada). Similarly, the AMS13 observations can be made available on request from Greg Wentworth (AEP), and the model output or code from Cynthia Whaley (ECCC). The aircraft observations are on the ECCC data portal (<http://donnees.ec.gc.ca/data/air/monitor/ambient-air-quality>).

785 References

- Akingunola, A., Makar, P. A., Zhang, J., Darlington, A., Li, S. M., Moran, M. D., and Zheng, Q.: Impact of continuous monitoring emissions stack parameters on model simulations using 2.5 km resolution GEM-MACH, *Atmos. Chem. Phys. Disc.*, 2017.
- Aneja, V., Bunton, B., Walker, J., and Malik, B.: Measurement and analysis of atmospheric ammonia emissions from anaerobic lagoons, *Atmos. Environ.*, 35, 1949 – 1958, doi:https://doi.org/10.1016/S1352-2310(00)00547-1, 2001.
- Asman, W. A. H., Sutton, M. A., and Schjorring, J. K.: Ammonia: emission, atmospheric transport and deposition, *New Phytologist*, 139, 27–48, doi:10.1046/j.1469-8137.1998.00180.x, 1998.
- Ayres, J., Bittman, S., Girdhar, S., Sheppard, S., Niemi, D., Ratte, D., and Smith, P.: Chapter 5: Sources of Ammonia Emissions, in: *The 2008 Canadian Atmospheric Assessment of Agricultural Ammonia, Environment and Climate Change Canada*, Gatineau, QC, Canada, 2009.
- Bash, J. O., Walker, J. T., Katul, G. G., Jones, M. R., Nemitz, E., and Robarge, W. P.: Estimation of In-Canopy Ammonia sources and sinks in a fertilized Zea mays field, *Environ. Sci. Tech.*, 44, 1683–1689, doi:10.1021/es9037269, 2010.
- 800 Bash, J. O., Cooter, E. J., Dennis, R. L., Walker, J. T., and Pleim, J. E.: Evaluation of a regional air-quality model with bidirectional NH₃ exchange coupled to an agroecosystem model, *Biogeosciences*, 10, 1635–1645, doi:10.5194/bg-10-1635-2013, 2013.
- Beer, R., Shephard, M. W., Kulawik, S. S., Clough, S. A., Eldering, A., Bowman, K. W., Sander, S. P., Fisher, B. M., Payne, V. H., Luo, M., Osterman, G. B., and Worden, J. R.: First satellite observations of lower tropospheric ammonia and methanol, *Geophys. Res. Lett.*, 35, doi:10.1029/2008GL033642, 109801, 2008.
- 805 Behera, S. N., Sharma, M., Aneja, V. P., and Balasubramanian, R.: Ammonia in the atmosphere: a review on emission sources, atmospheric chemistry and deposition on terrestrial bodies, *Environ. Sci. Pollut. Res. Int.*, 20, 8092–8131, doi:10.1007/s11356-013-2051-9, 2013.
- Biswas, H., Catterjee, A., Mukhopadhyaya, S. K., De, T. K., Sen, S., and Jana, T. K.: Estimation of ammonia exchange at the land ocean boundary condition of Sundarban mangrove, northeast coast of Bay of Bengal, India, *Atmos. Environ.*, 39, 4489–4499, doi:10.1016/j.atmosenv.2005.03.041, 2005.
- 810 Bittman, S., Ayres, J., S. Sheppard, S., and Girdhar, S.: Chapter 4: Emission Inventory Development, in: *The 2008 Canadian Atmospheric Assessment of Agricultural Ammonia, Environment and Climate Change Canada*, Gatineau, QC, Canada, 2008.
- 815 Booth, M. S., Stark, J. M., and Rastetter, E.: Controls on nitrogen cycling in terrestrial ecosystems: a synthetic analysis of literature data, *Ecol. Monogr.*, 75, 139–157, 2005.
- Bouwman, A., Lee, D. S., Asman, W. A. H., Dentener, F. J., van der Hoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Glob. Biogeochem. Cycles*, 11, 561–587, doi:10.1029/97GB02266, 1997.
- 820 Buehner, M., Morneau, J., and Charette, C.: Four-dimensional ensemble–variational data assimilation for global deterministic weather prediction, *Nonlinear Processes Geophys.*, 20, 669–682, doi:10.5194/npg-20-669-2013, 2013.
- Buehner, M., McTaggart-Cowan, R., Beaulne, A., Charette, C., Garand, L., Heilliette, S., Lapalme, E., Laroche, S. S. R. M., Morneau, J., and Zadra, A.: Implementation of Deterministic Weather Forecasting Systems based

- 825 on Ensemble-Variational Data Assimilation at Environment Canada. Part I: The Global System, *Mon. Wea. Rev.*, 143, 2532–2559, doi:10.1175/MWR-D-14-00354.1, 2015.
- Bytnerowicz, A., Fraczek, W., Schilling, S., and Alexander, D.: Spatial and temporal distribution of ambient nitric acid and ammonia in the Athabasca Oil Sands Region, Alberta, *J. Limnol.*, 69, 11–21, doi:10.3274/JL10-69-S1-03, 2010.
- 830 Carfrae, J. A., Sheppard, L. J., Raven, J., Stein, W., Leith, I. D., Theobald, A., and Crossley, A.: Early effects of atmospheric ammonia deposition on *Calluna vulgaris* (L.) hull growing on an ombrotrophic peat bog, *Water Air Soil Pollut. Focus*, 4, 229–239, doi:10.1007/s11267-004-3033-1, 2004.
- Caron, J.-F., Milewski, T., Buehner, M., Fillion, L., Reszka, M., Macpherson, S., and St-James, J.: Implementation of deterministic weather forecasting systems based on ensemble–variational data assimilation at Environment Canada. Part II: The regional system, *Mon. Wea. Rev.*, 143, 2560–2580, doi:10.1175/MWR-D-14-00353.1, 2015.
- 835 Charpentier, A. D., Bergerson, J. A., and McLean, H. L.: Understanding the Canadian oil sands industry’s greenhouse gas emissions, *Environ. Res. Lett.*, 4, 1–11, doi:10.1088/1748-9326/4/1/014005, 2009.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles, in: *Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., p. 465–570, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 840 2013.
- Clair, T. A. and Percy, K. E.: Assessing forest health in the Athabasca Oil Sands Region, Wbea technical report, Wood Buffalo Environmental Agency, Alberta, 2015.
- Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nature Geosci.*, 2, doi:10.1038/ngeo551, 109801, 2009.
- 850 Dragosits, U., Theobald, M., Place, C., Lord, E., Webb, J., Hill, J., ApSimon, H., and Sutton, M.: Ammonia emission, deposition and impact assessment at the field scale: a case study of sub-grid spatial variability, *Environmental Pollution*, 117, 147 – 158, doi:https://doi.org/10.1016/S0269-7491(01)00147-6, 2002.
- Ellis, R., Murphy, J. G., Markovic, M. Z., VandenBoer, T. C., Makar, P. A., Brooks, J., and Mihele, C.: The influence of gas-particle partitioning and surface-atmosphere exchange on ammonia during BAQS-Met, *Atmos. Chem. Phys.*, 11, 133–145, doi:10.5194/acp-11-133-2011, 2011.
- 855 Environment and Climate Change Canada: Canadian Environmental Sustainability Indicators: Air Pollutant Emissions, Technical report, Environment and Climate Change Canada, ECCC Public Inquiries Centre, 200 Sacre-Coeur boul. Gatineau, QC, K1A 0H3, <http://www.ec.gc.ca/indicateurs-indicators/default.asp?lang=en&n=E79F4C12-1>, 2016.
- 860 Environment and Climate Change Canada: Criteria air contaminants, Tech. rep., Environment and Climate Change Canada and Canadian Council of Ministers of the Environment, <https://www.ec.gc.ca/air/default.asp?lang=En&n=7C43740B-1>, last referenced 9 June 2017, 2017.
- Environment Canada: Precursor contributions to ambient fine particulate matter in Canada, Report, Air Quality Research Division, Environment and Climate Change Canada, Toronto, Ontario, Canada, 2001.

- 865 Fangmeier, A., Hadwiger-Fangmeier, A., der Eerden, L. V., and Jäger, H.-J.: Effects of atmospheric ammonia on vegetation—A review, *Environmental Pollution*, 86, 43 – 82, doi:[http://dx.doi.org/10.1016/0269-7491\(94\)90008-6](http://dx.doi.org/10.1016/0269-7491(94)90008-6), 1994.
- Farquhar, G. D., Firth, P. M., Wetselaar, R., and Weir, B.: On the Gaseous Exchange of Ammonia between Leaves and the Environment Determination of the Ammonia Compensation Point, *Plant Physiol.*, 66, 710–714, doi:10.1104/pp.66.4.710, 1980.
- 870 Flechard, C. R. and Fowler, D.: Atmospheric ammonia at a moorland site. II: Long term surface atmosphere micrometeorological flux measurements, *Q. J. Roy. Meteor. Soc.*, 124, 759–791, doi:10.1002/qj.49712454706, 1998.
- Fowler, D., Flechard, C. R., Sutton, M. A., and Storeton-West, R. L.: Long term measurements of the land atmosphere exchange of ammonia over moorland, *Atmos. Environ.*, 32, 453–459, doi:10.1016/S1352-2310(97)00044-7, 1998.
- 875 Fu, X., Wang, S. X., Rau, L. M., Pleim, J. E., Cooter, E., Bash, J. O., Benson, V., and Hao, J. M.: Estimating NH₃ emissions from agricultural fertilizer application in China using the bi-directional CMAQ model coupled to an agro-ecosystem model, *Atmos. Chem. Phys.*, 15, 6637–6649, doi:10.5194/acp-15-6637-2015, 2015.
- 880 Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z. C., Freney, J. R., Martinelli, L. A., Seitzinger, S. P., and Sutton, M. A.: Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions, *Science*, 320, 889–892, doi:10.1126/Science.1136674, 2008.
- Galperin, M. and Sofiev, M.: The long-range transport of ammonia and ammonium in the Northern Hemisphere, *Atmos. Environ.*, 32, 373 – 380, doi:10.1016/S1352-2310(97)00045-9, 1998.
- 885 Giordano, L., Brunner, D., Flemming, J., Hogrefe, C., Im, U., Bianconi, R., Badia, A., Balzarini, A., Baró, R., Chemel, C., Curci, G., Forkel, R., Jiménez-Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Jorba, O., Knote, C., Kuenen, J., Makar, P., Manders-Groot, A., Neal, L., Pérez, J., Pirovano, G., Pouliot, G., José, R. S., Savage, N., Schröder, W., Sokhi, R., Syrakov, D., Torian, A., Tuccella, P., Werhahn, J., Wolke, R., Yahya, K., Zabkar, R., Zhang, Y., and Galmarini, S.: Assessment of the MACC reanalysis and its influence as chemical boundary conditions for regional air quality modeling in AQMEII-2, *Atmos. Environ.*, 115, 371–388, doi:10.1016/j.atmosenv.2015.02.034, 2015.
- 890 Gong, W., Makar, P. A., Zhang, J., Milbrandt, J., Gravel, S., Hayden, K. L., Macdonald, A. M., and Leaitch, W. R.: Modelling aerosol cloud meteorology interaction: A case study with a fully coupled air quality model GEM-MACH, *Atmos. Environ.*, 115, 695–715, doi:10.1016/j.atmosenv.2015.05.062, 2015.
- 895 Gordon, M., Li, S.-M., Staebler, R., Darlington, A., Hayden, K., O'Brien, J., and Wolde, M.: Determining air pollutant emission rates based on mass balance using airborne measurement data over the Alberta oil sands operations, *Atmos. Meas. Tech.*, 8, 3745–3765, doi:10.5194/amt-8-3745-2015, 2015.
- Hansen, K., Personne, E., Skjoth, C. A., Loubet, B., Ibrom, A., Jensen, R., Sorensen, L. L., and Boegh, E.: Investigating sources of measured forest-atmosphere ammonia fluxes using two-layer bi-directional modelling, *Agriculture and Forest Meteorology*, 237–238, 20–94, doi:10.1016/j.agrformet.2017.02.008, 2017.
- 900 Heilman, W. E., Liu, Y., Urbanski, S., Kovalev, V., and Mickler, R.: Wildland fire emissions, carbon, and climate: Plume rise, atmospheric transport, and chemistry processes, *Forest Ecology and Management*, 317, 70 – 79, doi:<http://dx.doi.org/10.1016/j.foreco.2013.02.001>, wildland fire emissions, carbon, and climate: Science overview and knowledge needs, 2014.

- 905 Hsu, Y.-M. and Clair, T. A.: Measurement of fine particulate matter water-soluble inorganic species and precursor gases in the Alberta Oil Sands Region using an improved semicontinuous monitor, *J. Air Waste Manage. Assoc.*, 65, 423–435, doi:10.1080/10962247.2014.1001088, 2015.
- Hsu, Y.-M. and Clair, T. A.: Measurement of fine particulate matter water-soluble inorganic species and precursor gases in the Alberta Oil Sands Region using an improved semicontinuous monitor, *J. Air Waste Manage. Assoc.*, 568, 285–295, doi:10.1016/j.scitotenv.2016.05.205, 2016.
- 910 IPCC: Climate Change 2013: The Physical Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, in: *IPCC Fifth Assessment Report*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 915 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., and et al: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.
- Kelly, E. N., Short, J. W., Schindler, D. W., Hodson, P. V., Ma, M., Kwan, A. K., and Fortin, B. L.: Oil sands development contributes polycyclic aromatic compounds to the Athabasca River and its tributaries, *Proc. Natl. Acad. Sci.*, 106, 22 346–22 351, doi:10.1073/pnas.0912050106, 2009.
- 920 Kharol, S. K., Shephard, M. W., McLinden, C. A., Zhang, L., Sioris, C. E., O’Brien, J. M., Vet, R., Cady-Pereira, K. E., Hare, E., Siemons, J., and Krotkov, N. A.: Dry deposition of reactive nitrogen from satellite observations of ammonia and nitrogen dioxide over North America, *Geophys. Res. Lett.*, doi:10.1002/2017GL075832, 2017.
- 925 Kirk, J. L., Muir, D. C. G., Gleason, A., Wang, X., Lawson, G., Frank, R. A., Lehnher, I., and Wrona, F.: Atmospheric Deposition of Mercury and Methylmercury to Landscapes and Waterbodies of the Athabasca Oil Sands Region, *Environ. Sci. Technol.*, 48, 7374–7383, doi:10.1021/es500986r, 2014.
- Krupa, S.: Effects of atmospheric ammonia (NH₃) on terrestrial vegetation: a review, *Environmental Pollution*, 124, 179 – 221, doi:10.1016/S0269-7491(02)00434-7, 2003.
- 930 Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., Mc-Connell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- 935 Larkin, N. K., O’Neill, S. M., Solomon, R., Raffuse, S., Strand, T., Sullivan, D., Krull, C., Rorig, M., Peterson, J., and Ferguson, S. A.: The BlueSky smoke modeling framework, *Int. J. Wildland Fire*, 18, 906–920, doi:10.1071/WF07086, 2009.
- Lee, C. J., Martin, R. V., Henze, D. K., Brauer, M., Cohen, A., and van Donkelaar, A.: Response of Global particulate matter related mortality to changes in local precursor emissions, *Environ. Sci. and Tech.*, 49, 4335–4344, doi:10.1021/acs.est.5b00873, 2015.
- 940 Li, S.-M., Leithead, A., Moussa, S. G., Liggio, J., Moran, M. D., Wang, D., Hayden, K., Darlington, A., Gordon, M., Staebler, R., Makar, P. A., Stroud, C. A., McLaren, R., Liu, P. S. K., O’Brien, J., Mittermeier, R. L., Zhang, J., Marson, G., Cober, S. G., Wolde, M., and Wentzell, J. J. B.: Differences between measured and

reported volatile organic compound emissions from oil sands facilities in Alberta, Canada, *Proc. Nat. Acad. Sci.*, doi:10.1073/pnas.1617862114, 2017.

945 Liggio, J., Li, S.-M., Hayden, K., Taha, Y. M., Stroud, C., Darlington, A., Drollette, B. D., Gordon, M., Lee, P., Liu, P., Leithead, A., Moussa, S. G., Wang, D., O'Brien, J., Mittermeier, R. L., Brook, J. R., Lu, G., Staebler, R. M., Han, Y., Tokarek, T. W., Osthoff, H. D., Makar, P. A., Zhang, J., L. Plata, D., and Genthner, D. R.: Oil sands operations as a large source of secondary organic aerosols, *Nature*, 534, 91–94, doi:10.1038/nature17646, 2016.

950 Makar, P., Bouchet, V. S., and Nenes, A.: Inorganic chemistry calculations using HETV - a vectorized solver for the SO_4^{2-} - NO_3^- - NH_4^+ system based on the ISORROPIA Algorithms, *Atmos. Environ.*, 37, 2279–2294, doi:10.5194/acp-9-7183-2009, 2003.

Makar, P., Moran, M., Zheng, Q., Cousineau, S., Sassi, M., Duhamel, A., Besner, M., Davignon, D., Crevier, L.-P., and Bouchet, V. S.: Modelling the impacts of ammonia emissions reductions on North American air quality, *Atmos. Chem. Phys.*, 9, 7183–7212, doi:10.1016/S1352-2310(03)00074-8, 2009.

955 Makar, P., Gong, W., Hogrefe, C., Zhang, Y., Curci, G., Zabkar, R. ., Milbrandt, J., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jiménez-Guerrero, P., Langer, M., Moran, M., Pabla, B., Pérez, J., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., Zhang, J., and Galmarini, S.: Feedbacks between air pollution and weather, part 2: Effects on chemistry, *Atmos. Environ.*, 115, 499–526, doi:10.1016/j.atmosenv.2014.10.021, 2015a.

960 Makar, P. A., Gong, W., Milbrandt, J., Hogrefe, C., Zhang, Y., Curci, G., Zabkar, R. ., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jiménez-Guerrero, P., Langer, M., Moran, M., Pabla, B., Pérez, J., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., Zhang, J., and Galmarini, S.: Feedbacks between air pollution and weather, part 1: Effects on weather, *Atmos. Environ.*, 115, 442–469, doi:10.1016/j.atmosenv.2014.12.003, 2015b.

965 Makar, P. A., Akingunola, A., Aherne, J., Cole, A. S., Aklilu, Y., Zhang, J., Wong, I., Hayden, K., Li, S. M., Kirk, J., Scott, K., Moran, M. D., Robichaud, A., Cathcart, H., Baratzehah, P., Pabla, B., Cheung, P., Zheng, Q., and Jeffries, D. S.: Estimates of exceedances of critical loads for acidifying deposition in Alberta and Saskatchewan, *Atmos. Chem. Phys. Disc.*, submitted to ACPD Oil Sands special issue, 2017.

970 Markovic, M. Z., VandenBoer, T. C., and Murphy, J. G.: Characterization and optimization of an online system for the simultaneous measurement of atmospheric water-soluble constituents in the gas and particle phases, *J. Env. Mon.*, 14, 1872–1884, doi:1, 2012.

Massad, R.-S., Nemitz, E., and Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere, *Atmos. Chem. Phys.*, 10, 10 359–10 386, doi:10.5194/acp-10-10359-2010, 2010.

975 McCalley, C. K. and Sparks, J. P.: Controls over nitric oxide and ammonia emissions from Mojave Desert soils, *Oecologia*, 156, 871–881, doi:10.1007/s00442-008-1031-0, 2008.

Moran, M., Menard, S., Gravel, S., Pavlovic, R., and Anselmo, D.: RAQDPS Versions 1.5.0 and 1.5.1: Upgrades to the CMC Operational Regional Air Quality Deterministic Prediction System Released in October 2012 and February 2013, Technical report, Canadian Meteorological Centre, Canadian Meteorological Centre, Dorval, Quebec, 2013.

980

- Moran, M., Zheng, Q., Zhang, J., and Pavlovic, R.: RAQDPS Version 013: Upgrades to the CMC Operational Regional Air Quality Deterministic Prediction System Released in June 2015, Technical report, Canadian Meteorological Centre, Canadian Meteorological Centre, Dorval, Quebec, 2015.
- Moran, M. D., Ménard, S., Talbot, D., Huang, P., Makar, P. A., Gong, W., Landry, H., Gravel, S., Gong, S., Crevier, L.-P., Kallaur, A., and Sassi, M.: Particulate-matter forecasting with GEM-MACH15, a new Canadian air-quality forecast model, in: Air pollution modelling and its application XX, edited by Steyn, D. G. and Rao, S. T., pp. 289–292, Springer, Dordrecht, 2010.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, doi:10.1038/nature08823, 2010.
- Murano, K., Mukai, H., Hatakeyama, S., Oishi, O., Utsunomiya, A., and Shimohara, T.: Wet deposition of ammonium and atmospheric distribution of ammonia and particulate ammonium in Japan, *Environmental Pollution*, 102, 321 – 326, doi:10.1016/S0269-7491(98)80050-X, 1998.
- Nemitz, E., Sutton, M. A., Schjoerring, J. K., Husted, S., and Wyers, G. P.: Resistance modelling of ammonia exchange over oilseed rape, *Agr. For. Meteorol.*, 105, 405–425, doi:10.1016/S0168-1923(00)00206-9, 2000.
- Nemitz, E., Milford, C., and Sutton, M. A.: A two-layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia, *Q. J. Roy. Meteorol. Soc.*, 127, 815–833, doi:10.1002/qj.49712757306, 2001.
- Nenes, A., Pilinis, C., and Pandis, S.: ISORROPIA: A New Thermodynamic Model for Multiphase Multicomponent Inorganic Aerosols, *Aquat. Geochem.*, 4, 123–152, doi:10.1023/A:1009604003981, 1998.
- Olivier, J., Bouwman, A., der Hoek, K. V., and Berdowski, J.: Global air emission inventories for anthropogenic sources of NO_x, NH₃ and N₂O in 1990, *Environmental Pollution*, 102, 135 – 148, doi:http://dx.doi.org/10.1016/S0269-7491(98)80026-2, 1998.
- Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *J. Geophys. Res.*, 109, doi:10.1029/2003JD004473, d15204, 2004.
- Paugam, R., Wooster, M., Freitas, S., and Val Martin, M.: A review of approaches to estimate wildfire plume injection height within large-scale atmospheric chemical transport models, *Atmos. Chem. Phys.*, 16, 907–925, doi:10.5194/acp-16-907-2016, 2016.
- Paulot, F., Jacob, D. J., Pinder, R. W., Bash, J. O., Travis, K., and Henze, D. K.: Ammonia emissions in the United States, European Union, and China derived by high-resolution inversion of ammonium wet deposition data: Interpretation with a new agricultural emissions inventory (MASAGE NH₃), *J. Geophys. Res.*, 119, 4343–4364, doi:10.1002/2013JD021130, 2014.
- Pavlovic, R., Chen, J., Anderson, K., Moran, M., Beaulieu, P.-A., Davignon, D., and Cousineau, S.: The Fire-Work air quality forecast system with near-real-time biomass burning emissions: Recent developments and evaluation of performance for the 2015 North American wildfire season, *J. Air Waste Manage. Assoc.*, 66, 819–841, doi:10.1080/10962247.2016.1158214, 2016.

- Pleim, J. E., Bash, J. O., Walker, J. T., and Cooter, E. J.: Development and evaluation of an ammonia bidirectional flux parameterization for air quality models, *J. Geophys. Res.*, 118, 3794–3806, doi:10.1002/jgrd.50262, 2013.
- Pope III, C., Burnett, R., Thun, M., and et al: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *JAMA*, 287, 1132–1141, doi:10.1001/jama.287.9.1132, 2002.
- Robichaud, A.: Statistical links between meteorological factors and tropospheric ozone levels at the Duchesnay forest site, *Climat*, 12, 31–57, 1994.
- Robichaud, A. and Lin, C. A.: The linear steady response of a stratified baroclinic atmosphere to elevated diabatic forcing, *Atmosphere - Ocean*, 29, 619–635, doi:10.1080/07055900.1991.9649421, 1991.
- 1030 Rogers, C. D.: *Inverse methods for atmospheric Sounding: Theory and Practice*, World Sci., Hackensack, NJ, 2000.
- Rooney, R. C., Bayley, S. E., and Schindler, D. W.: Oil sands mining and reclamation cause massive loss of peatland and stored carbon, *Proc. Natl. Acad. Sci.*, 109, 4933–4937, doi:10.1073/pnas.1117693108, 2012.
- Seinfeld, J. H. and Pandis, S. N., eds.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley-Intersci., New York, 1326 pp., 1998.
- 1035 Shephard, M. W. and Cady-Pereira, K. E.: Cross-track Infrared Sounder (CrIS) satellite observations of tropospheric ammonia, *Atmos. Meas. Techn.*, 8, 1323–1336, doi:10.5194/amt-8-1323-2015, 2015.
- Shephard, M. W., Cady-Pereira, K. E., Luo, M., Henze, D. K., Pinder, R. W., Walker, J. T., Rinsland, C. P., Bash, J. O., Zhu, L., Payne, V. H., and Clarisse, L.: TES ammonia retrieval strategy and global observations of the spatial and seasonal variability of ammonia, *Atmos. Chem. Phys.*, 11, 10743–10763, doi:10.5194/acp-11-10743-2011, 2011.
- 1040 Shephard, M. W., McLinden, C. A., Cady-Pereira, K. E., Luo, M., Moussa, S. G., Leithead, A., Liggio, J., Staebler, R. M., Akingunola, A., Makar, P., Lehr, P., Zhang, J., Henze, D. K., Millet, D. B., Bash, J. O., Zhu, L., Wells, K. C., Capps, S. L., Chaliyakunnel, S., Gordon, M., Hayden, K., Brook, J. R., Wolde, M., and Li, S.-M.: Tropospheric Emission Spectrometer (TES) satellite observations of ammonia, methanol, formic acid, and carbon monoxide over the Canadian oil sands: validation and model evaluation, *Atmos. Meas. Tech.*, 8, 5189–5211, doi:10.5194/amt-8-5189-2015, 2015.
- 1045 Shinozuka, Y., Redemann, J., Livingston, J. M., Russell, P. B., Clarke, A. D., Howell, S. G., Freitag, S., O'Neill, N. T., Reid, E. A., Johnson, R., Ramachandran, S., McNaughton, C. S., Kapustin, V. N., Brekhovskikh, V., Holben, B. N., and McArthur, L. J. B.: Airborne observation of aerosol optical depth during ARC-TAS: vertical profiles, inter-comparison and fine-mode fraction, *Atmos. Chem. Phys.*, 11, 3673–3688, doi:10.5194/acp-11-3673-2011, 2011.
- Stensland, G. J., Bowersox, V. C., Larson, B., and Claybrooke, R. D.: Comparison of Ammonium in USA Wet Deposition to Ammonia emission estimates, technical report, Illinois State Water Survey, 2204 Griffith Drive, Champaign, IL, 61820, <https://www3.epa.gov/ttn/chief/conference/ei10/ammonia/stensland.pdf>, 2000.
- 1055 Sutton, M., Milford, C., Dragosits, U., Place, C., Singles, R., Smith, R., Pitcairn, C., Fowler, D., Hill, J., ApSimon, H., Ross, C., Hill, R., Jarvis, S., Pain, B., Phillips, V., Harrison, R., Moss, D., Webb, J., Espenhahn, S., Lee, D., Hornung, M., Ulyett, J., Bull, K., Emmett, B., Lowe, J., and Wyers, G.: Dispersion, deposition and impacts of atmospheric ammonia: quantifying local budgets and spatial variability, *Environmental Pollution*, 102, 349 – 361, doi:http://dx.doi.org/10.1016/S0269-7491(98)80054-7, 1998.
- 1060

- Sutton, M. A., Fowler, D., and Moncrieff, J. B.: The Exchange of Atmospheric Ammonia with Vegetated Surfaces .1. Unfertilized Vegetation, *Q. J. Roy. Meteorol. Soc.*, 119, 1023–1045, doi:10.1002/qj.49711951309, 1993.
- Sutton, M. A., Schjorring, J. K., and Wyers, G. P.: Plant Atmosphere Exchange of Ammonia, *Philos. T. Roy. Soc. A.*, 351, 261–276, <http://www.jstor.org/stable/54415>, 1995.
- 1065 Urbanski, S.: Wildland fire emissions, carbon, and climate: Emission factors, *Forest Ecology and Management*, 317, 51–60, doi:10.1016/j.foreco.2013.05.045, 2014.
- Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., and Coheur, P. F.: Global distributions, time series and error characterization of atmospheric ammonia (NH₃) from IASI satellite observations, *Atmos. Chem. Phys.*, 14, 2905–2922, doi:10.5194/acp-14-2905-2014, 2014.
- 1070 Vile, M. A., Kelman Wieder, R., Živković, T., Scott, K. D., Vitt, D. H., Hartsock, J. A., Iosue, C. L., Quinn, J. C., Petix, M., Fillingim, H. M., Popma, J. M. A., Dynarski, K. A., Jackman, T. R., Albright, C. M., and Wyckoff, D. D.: N₂-fixation by methanotrophs sustains carbon and nitrogen accumulation in pristine peatlands, *Biogeochemistry*, 121, 317–328, doi:10.1007/s10533-014-0019-6, 2014.
- 1075 Vitt, D. H.: Nitrogen addition experiments in boreal ecosystems: understanding the fate of atmospheric deposited nitrogen in order to determine nitrogen critical loads, Cema report, Wood Buffalo Environmental Agency, Alberta, 2016.
- Wen, D., Zhang, L., Lin, J. C., Vet, R., and Moran, M. D.: An evaluation of ambient ammonia concentrations over southern Ontario simulated with different dry deposition schemes within STILT-Chem v0.8, *Geosci. Model Dev.*, 7, 1037–1050, doi:10.5194/gmd-7-1037-2014, 2014.
- 1080 Wentworth, G. R., Murphy, J. G., Gregoire, P. K., Cheyne, C. A. L., Tevlin, A. G., and Hems, R.: Soil–atmosphere exchange of ammonia in a non-fertilized grassland: measured emission potentials and inferred fluxes, *Biogeosciences*, 11, 5675–5686, doi:10.5194/bg-11-5675-2014, 2014.
- 1085 Wentworth, G. R., Murphy, J. G., Willis, M. D., Lee, M. K. Y., Abbat, J. P. D., Li, S.-M., Brook, J. R., Makar, P., Akingunola, A., and Whaley, C.: Ammonia and particle acidity in the Athabasca Oil Sands Region, tbd, tbd, tbd, doi:tbd, 2017.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.*, 23, 1293–1304, doi:10.1016/0004-6981(89)90153-4, 1989.
- 1090 Wichink Kruit, R., van Pul, W., Sauter, F., van den Broek, M., Nemitz, E., Sutton, M., Krol, M., and Holtslag, A.: Modeling the surface–atmosphere exchange of ammonia, *Atmos. Environ.*, 44, 945 – 957, doi:10.1016/j.atmosenv.2009.11.049, 2010.
- Wieder, R. K., Vile, M. A., Albright, C. M., Scott, K. D., Vitt, D. H., Quinn, J. C., and Burke-Scoll, M.: Effects of altered atmospheric nutrient deposition from Alberta oil sands development on *Sphagnum fuscum* growth and C, N and S accumulation in peat, *Biogeochemistry*, 129, 1–19, doi:10.1007/s10533-016-0216-6, 2016a.
- 1095 Wieder, R. K., Vile, M. A., Scott, K. D., Albright, C. M., McMillen, K. J., Vitt, D. H., and Fenn, M. E.: Differential effects of high atmospheric N and S deposition on bog plant/lichen tissue and porewater chemistry across the Athabasca Oil Sands Region, *Environ. Sci. Technol.*, 50, 12 630–12 640, doi:10.1021/acs.est.6b03109, 2016b.




Figure 1. Compensation point (C_a) relationship to temperature: C_a for evergreen needleleaf LUC shown as example.

- 1100 Wu, Y., Walker, J., Schwede, D., Peters-Lidard, C., Dennis, R., and Robarge, W.: A new model of bi-directional ammonia exchange between the atmosphere and biosphere: Ammonia stomatal compensation point, *Agr. Forest Meteorol.*, 149, 263 – 280, doi:10.1016/j.agrformet.2008.08.012, 2009.
- Zhang, J., Moran, M. D., Zheng, Q., Makar, P. A., Baratzadeh, P., Marsen, G., Liu, P., and Li, S.-M.: Emissions preparation and analysis for multiscale air quality modelling over the Athabasca oil sands region of Alberta, Canada, *Atmos. Chem. Phys. Disc.*, submitted to ACPD Oil Sands special issue, 2017.
- 1105 Zhang, L., Moran, M., Makar, P., Brook, J., and Gong, S.: Gaseous Dry Deposition in AURAMS A Unified Regional Air-quality Modelling System, *Atmos. Environ.*, 36, 537–560, doi:10.1016/S1352-2310(01)00447-2, 2002.
- Zhang, L., Brook, J. R., and Vet, R.: A revised parametrization for gaseous dry deposition in air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, doi:10.5194/acp-3-2067-2003, 2003.
- 1110 Zhang, L., Wright, L. P., and Asman, W. A. H.: Bi-directional air-surface exchange of atmospheric ammonia: A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models, *J. Geophys. Res.*, 115, D20 310, doi:10.1029/2009JD013589, 2010.
- Zhu, L., Henze, D. K., Cady-Pereira, K. E., Shephard, M. W., Luo, M., Pinder, R. W., Bash, J. O., and Jeong, G.-R.: Constraining U.S. ammonia emissions using TES remote sensing observations and the GEOS-Chem adjoint model, *J. Geophys. Res.*, 118, 3355–3368, doi:10.1002/jgrd.50166, 2013.
- 1115 Zhu, L., Henze, D., Bash, J., Jeong, G.-R., Cady-Pereira, K., Shephard, M., Luo, M., Paulot, F., and Capps, S.: Global evaluation of ammonia bidirectional exchange and livestock diurnal variation schemes, *Atmos. Chem. Phys.*, 15, 12 823–12 843, doi:10.5194/acp-15-12823-2015, 2015.
- 1120 ~~Maps modelled daily average (a) bidi and (b) fire components to dry deposition. In all maps, red/positive represents upward flux, and blue/negative represents downward flux. These are daily amounts, averaged over 12 August to 7 September, 2013.~~




Figure 2. Map of 10-km resolution continental piloting model domain (green), and 2.5-km resolution nested model domain (purple).




Figure 3. Flight path on 13 August 2013, where elevation (in meters) is denoted by the colour scale, and the AMS13 site is indicated by a black circle.

Maps of the modelled average NH_4^+ wet deposition for (a) base (b) bidi, and (c) fire+bidi models. Negative represents downward flux. These are daily amounts, averaged over 12 August to 7 September, 2013.

(Top panels) Images of the Alberta/Saskatchewan region with clouds and fire hotspots from MODIS. (Bottom panels) Map of CrIS-measured surface NH_3 concentrations, with coloured boxes showing the regions where model and satellite measurements were sampled. These three examples are for (a) northern-bidi case study (cyan), (b) southern-bidi case study (black), and (c) fire case study (magenta), discussed in Section 4.3), and the blue box is the region of our overall comparison.




Figure 4. Surface daily average (a) NH_3 , (b) fine particulate NH_4^+ , (c) NO_3^- , and (d) SO_4^{2-} concentrations at the AMS13 ground site in the AOSR. Measurements in orange, base model in green, bidirectional flux model in blue, and fire+bidi model in red. **Y-scale is logarithmic.**

Table 1. Latitude and longitude ranges that the model was evaluated over with the CrIS satellite measurements

domain	date (in 2013)	lat range (°)	lon range (°)
AB/SK large domain	12 Aug to 7 Sept	48 to 60 N	-122.0 to -100.0 W
northern, no-fire case study	3 Sept	55 to 60 N	-120.0 to -110.0 W
southern, no-fire case study	1 Sept	49 to 53.5 N	-117.0 to -106.0 W
northern, fire case study	12 Aug	56.5 to 60 N	-110.0 to -104.4 W




Figure 5. ~~Model-measurement~~ Hourly model-measurement bias in surface (a) NH_3 , (b) NH_4^+ , (c) NO_3^- and (d) SO_4^{2-} concentrations at the AMS13 ground site in the AOSR.




Figure 6. ~~Modelled-Hourly modelled~~ vs measured surface (a) NH₃, (b) NH₄⁺, (c) NO₃⁻ and (d) SO₄²⁻ concentrations at the AMS13 ground site in the AOSR. Base model in grey, bidirectional flux model in blue, and fire+bidirectional model in red.




Figure 7. NH₃ concentrations aloft (colour scale) over the OS region during the 13 August 2013 flight. (a) measurements, (b) base model, (c) fire+bidi model, and (d) bidi model.




Figure 8. Hourly averages along all flight paths over the OS region during the summer 2013 campaign: Model-measurement bias in (a) NH_3 and (b) NH_4^+ . Modelled vs measured (c) NH_3 and (d) NH_4^+ concentrations aloft. Base model in grey, bidirectional flux model in blue, and fire+bidi model in red.





Figure 9. Top panel: (a) NH_3 vertical profile as measured by CrIS satellite, difference between measurement and (b) base model, (c) bidi model, and (d) fire+bidi model, and (e) averaging kernel of CrIS satellite for NH_3 retrieval. Bottom panel: average (12 Aug - 7 Sep 2013) surface NH_3 concentrations given by the (a) base model, (b) fire+bidi model, and (c) CrIS satellite.




Figure 10. (Top panels) Images of the Alberta/Saskatchewan region with clouds and fire hotspots from MODIS. (Bottom panels) Map of CrIS-measured surface NH_3 concentrations, with coloured boxes showing the regions where model and satellite measurements were sampled. These three examples are for (a) northern bidi case study (cyan), (b) southern bidi case study (black), and (c) fire case study (magenta), discussed in Section 4.3), and the blue box is the region of our overall comparison.




Figure 11. As in Fig. 9, but for our (a) northern “bidi-only” case study (3 Sept 2013), (b) southern “bidi-only” case study (1 Sept 2013), and (c) northern “fire-only” case study (12 Aug 2013). Regions are shown in Figure 10a (cyan), 10b (black), and 10c (magenta) boxes, respectively).




Figure 12. Maps of the modelled average (a) surface NH_3 concentrations (b) absolute bidirectional flux contribution, (c) absolute fire contribution, (d) percent bidirectional flux contribution, and (e) percent fire contribution to surface NH_3 . These are averaged averages over 12 August to 7 September, 2013.




Figure 13. Maps of the modelled average NH_3 ~~dry~~ deposition for (a) base (b) bidi, and (c) fire+bidi models. In all maps, red/positive represents upward flux, and blue/negative represents downward flux. These are daily amounts, averaged over 12 August to 7 September, 2013.

Table 2. Model-measurement NH₃ comparison statistics from 12 August to 7 September 2013: R=correlation coefficient; slope is of the line-of-best fit between model vs. measurement; p and t are from a paired t-test between model and measurement data pairs (p>0.05 and |t|<1 means that the model is statistically indistinguishable from measurements); the median model bias; RMSE=root-mean-square error; and FE=fractional error of the models. CrIS (troposphere) results are for the entire model domain at all tropospheric levels shown in Figure 9(top), and CrIS (surface) results are for the lowest retrieval level (both are during mid-day satellite overpass times); aircraft results are from the 12 flight paths over the oil sands facilities, hourly averages during the daytime; and AMS13 results are from hourly data (day and night) at the one ground station.

	<u>R</u>	<u>slope</u>	<u>p</u>	<u>t</u>	<u>bias (ppbv)</u>	<u>RMSE (ppbv)</u>	<u>FE</u>
<u>CrIS (troposphere)</u>							
<u>base</u>	<u>0.248</u>	<u>0.076</u>	<u><2E-16</u>	<u>-247.5</u>	<u>-0.430</u>	<u>2.02</u>	<u>-5.3E-6</u>
<u>bidi</u>	<u>0.302</u>	<u>0.205</u>	<u><2E-16</u>	<u>-77.4</u>	<u>-0.176</u>	<u>1.93</u>	<u>-1.2E-6</u>
<u>fire+bidi</u>	<u>0.338</u>	<u>0.425</u>	<u><2E-16</u>	<u>36.2</u>	<u>-0.126</u>	<u>2.45</u>	<u>5.9E-7</u>
<u>CrIS (surface)</u>							
<u>base</u>	<u>0.272</u>	<u>0.118</u>	<u><2E-16</u>	<u>-19.0</u>	<u>-1.11</u>	<u>5.72</u>	<u>-1.6E-3</u>
<u>bidi</u>	<u>0.289</u>	<u>0.162</u>	<u><2E-16</u>	<u>-12.8</u>	<u>-0.66</u>	<u>5.32</u>	<u>-8.9E-4</u>
<u>fire+bidi</u>	<u>0.566</u>	<u>1.195</u>	<u>1.4E-06</u>	<u>4.9</u>	<u>-0.19</u>	<u>8.67</u>	<u>3.7E-4</u>
<u>aircraft (hourly)</u>							
<u>base</u>	<u>0.368</u>	<u>0.114</u>	<u>8.5E-14</u>	<u>-10.3</u>	<u>-0.751</u>	<u>1.14</u>	<u>-2.5E-3</u>
<u>bidi</u>	<u>0.549</u>	<u>0.503</u>	<u>0.0026</u>	<u>-3.2</u>	<u>-0.244</u>	<u>0.69</u>	<u>-5.0E-4</u>
<u>fire+bidi</u>	<u>0.560</u>	<u>0.519</u>	<u>0.0052</u>	<u>-2.9</u>	<u>-0.233</u>	<u>0.68</u>	<u>-4.5E-4</u>
<u>AMS13 (hourly)</u>							
<u>base</u>	<u>0.103</u>	<u>0.116</u>	<u><2E-16</u>	<u>-12.4</u>	<u>-0.35</u>	<u>0.92</u>	<u>-1.6E-3</u>
<u>bidi</u>	<u>0.413</u>	<u>0.652</u>	<u><2E-16</u>	<u>12.1</u>	<u>-0.30</u>	<u>0.95</u>	<u>8.0E-4</u>
<u>fire+bidi</u>	<u>0.403</u>	<u>0.691</u>	<u><2E-16</u>	<u>13.1</u>	<u>0.32</u>	<u>1.04</u>	<u>9.0E-4</u>

Table 3. Average source contributions to ambient NH₃ concentrations over the AB/SK model domain during 12 Aug to 7 Sep 2013.

source	median (ppbv)	median (%)	average (ppbv)	average (%)
total surface NH ₃	1.60	100	2.53	100
from fires to surface	0.25	10.4	0.42	20.3
from bidi to surface	0.97	56.3	1.24	56.6
from anthro to surface	0.38	33.3	0.87	23.1
total column NH ₃	18.8	100	25.6	100
from fires to total column	6.1	27.7	8.1	30.5
from bidi to total column	8.8	48.1	11.15	50.0
from anthro to total column	3.9	24.2	6.35	19.5

Table 4. Average NH_3 deposition (downward flux) over the AB/SK model domain during 12 Aug to 7 Sep 2013.

<u>Net Flux (moles/m²/day)</u>	<u>base</u>	<u>bidi</u>	<u>fire+bidi</u>
<u>mean</u>	<u>3.025×10^{-5}</u>	<u>1.811×10^{-5}</u>	<u>3.765×10^{-5}</u>
<u>median</u>	<u>2.061×10^{-5}</u>	<u>1.299×10^{-5}</u>	<u>2.843×10^{-5}</u>