

# ***Interactive comment on “Ozone trends over the United States at different times of day” by Yingying Yan et al.***

**Yingying Yan et al.**

yanyy09@pku.edu.cn

Received and published: 23 November 2017

Referee #2

The authors have conducted a very interesting study to investigate the ozone trends at different times of a day. This work includes many important details and could have important implications for future ozone air quality control. So I recommend this paper could be published after they have addressed some issues as listed below.

We thank the reviewer for comments, which have been incorporated to the revised manuscript.

Major comments.

Printer-friendly version

Discussion paper



When trying to quantify the influence of meteorology on U.S. ozone trends during 1990–2014, the authors focused on two climate patterns—AMO and ONI. A new index named AMONI is also constructed. Even they have found strong correlations between the annual mean ozone with these climate indices, but the underlying physical mechanism is unclear to the readers. It may be too reckless to say US ozone is dependent on AMONI. Here are my concerns.

There are a lot of evidence in the literature on the relevance of AMO and ONI (ENSO) to the US ozone. We discussed some evidence in the original manuscript, and we have further improved the discussion in the revised Sect. 2.2:

“We relate the interannual variability of ozone to two major climate indices relevant to the US air quality (Sutton et al., 2007; Lin et al., 2015): the Atlantic Multi-decadal Oscillation (AMO; <https://www.esrl.noaa.gov/psd/data/timeseries/AMO/>) index and the Oceanic Niño index (ONI; [http://origin.cpc.ncep.noaa.gov/products/analysis\\_monitoring/ensostuff/ONI\\_v5.php](http://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php)).

De-trended annual AMO index time series over 1990–2014 is calculated from the unsmoothed Kaplan sea surface temperature (SST) dataset of the National Oceanic and Atmospheric Administration (NOAA) Earth System Research Laboratory (<http://www.esrl.noaa.gov/psd/data/correlation/amon.us.data>). Observational and model studies have shown that the recent multi-decadal fluctuations in Atlantic SST were associated with large-scale climate anomalies (Enfield et al., 2001; Johannessen et al., 2004; Hu et al., 2011; Oglesby et al., 2012) important for the ozone chemistry. The warm conditions in the North Atlantic Ocean (positive AMO) have been associated with increased air temperatures at northern latitudes (by more than 3°C locally especially in winter) (Johannessen et al., 2004) and reduced summertime rainfall and increased droughts over much of the US (Enfield et al., 2001; McCabe et al., 2004). Delworth and Mann (2000) showed evidence of AMO-related variations in sea level pressure (SLP), and hence atmospheric circulation, in the North Atlantic region. Such AMO-associated meteorological changes can alter the distribution of tropospheric con-

Printer-friendly version

Discussion paper



stituents (Olsen et al., 2016), including ozone (Shen et al., 2015; Lin et al., 2014). For example, warmer temperature under a positive AMO phase tends to enhance the ozone production over the US (Lin et al., 2014); less precipitation means less cloudy and higher radiation for photochemistry (Kunkel et al., 2008); and the change in circulation affects the ozone transport (Fu et al., 2015).

De-trended annual ONI index time series over 1990–2014 is calculated from the NOAA Climate Prediction Center dataset ([http://www.cpc.noaa.gov/products/analysis\\_monitoring/ensostuff/ensoyears.shtml](http://www.cpc.noaa.gov/products/analysis_monitoring/ensostuff/ensoyears.shtml)). The ONI index refers to ERSST.v4 SST anomalies in the Niño 3.4 region, as an indicator of the El Niño–Southern Oscillation (ENSO). These SST anomalies are associated with global climate variability, including changes in the temperature and precipitation patterns over the United States (Ropelewski and Halpert, 1987; Yu et al., 2012; Yu and Zou, 2013; Liang et al., 2015), through alterations in atmospheric circulations and teleconnection patterns (Bjerknes, 1969; Enfield, 1989). The Niño 3.4 index (referred as ONI here) has been widely used to examine the impact of ENSO on surface ozone over the United States (e.g., Lin et al., 2015; Xu et al., 2017). Xu et al. (2017) showed that over 1993–2013, the monthly ozone decreases (increases) during El Niño (La Niña) years, with the amplitude varying from 0.4 ppb (for the US average) up to 1.8 ppb (for the southeastern US) per standard deviation of the Niño 3.4 index. The La Niña years (strongly negative ONI) tend to be associated with a more meandering jet over the central western US in favor of stratosphere–troposphere exchange (STE) of ozone but not transport from Asia (Lin et al., 2015).

In order to indicate the climate variability that influence the whole US, we combined the de-trended and normalized AMO and ONI indices to obtain a third index, named AMONI:  $AMONI = \frac{AMO_{detrended} - \overline{AMO_{detrended}}}{\sigma_{AMO_{detrended}}} + \frac{ONI_{detrended} - \overline{ONI_{detrended}}}{\sigma_{ONI_{detrended}}}$ . The normalization could adjust the AMO and ONI values measured on different scales to a common scale and keep the individual characteristics of the original AMO and ONI indices. The negative sign for

Printer-friendly version

Discussion paper



ONI in the formula accounts for the negative correlation between de-trended ozone and ONI anomalies (see Sect. 4). Thus a positive AMO and a negative ONI, both of which are closely related to higher ozone mixing ratios over the US, contribute to a positive AMONI index.

A recent work by Shen et al. (2017) developed two metrics,  $\text{MAM-}\Delta\text{SST}$  and  $\text{MAM-}\Delta\text{SLP}$ , to study June-July-August (JJA) MDA8 ozone variability across much of the eastern US. They found that  $\text{MAM-}\Delta\text{SST}$  is highly correlated to the summer ozone ( $R = \sim 0.7$ ), which level of correlation is comparable to our results for daytime and daily mean ozone (see Sect. 4)."

1. According to a new study (Shen et al., 2017), the U.S. summertime ozone is associated with a tripole SST pattern in the Atlantic and a dipole pattern in the northeastern Pacific. This raises the question that why the authors choose to use AMO and ONI. Is this the best choice?

See above for the choice of AMO and ONI.

The correlation between AMONI and ozone (0.71 for daytime mean ozone and 0.62 for daily mean ozone) is comparable to the correlation ( $\sim 0.7$ ) found by Shen et al. (2017).

We have added a discussion of Shen et al. (2017) in the revised Sect. 2.2.:

"A recent work by Shen et al. (2017) developed two metrics,  $\text{MAM-}\Delta\text{SST}$  and  $\text{MAM-}\Delta\text{SLP}$ , to study June-July-August (JJA) MDA8 ozone variability across much of the eastern US. They found that  $\text{MAM-}\Delta\text{SST}$  is highly correlated to the summer ozone ( $R = \sim 0.7$ ), which level of correlation is comparable to our results for daytime and daily mean ozone (see Sect. 4)."

2. The mechanism related to ENSO as proposed by Xu et al. [2017] only applies to the spring or fall season (e.g. the subtropical jet wind only explains the springtime ozone decreases in a small region in the southwest US). Since annual mean ozone is used in this study, the authors can't simply use the mechanism in Xu et al. [2017] to support

[Printer-friendly version](#)[Discussion paper](#)

their conclusion here.

See above for why we chose the ONI index.

Xu et al. (2017) showed that over 1993–2013, the monthly ozone decreases (increases) during El Niño (La Niña) years, with the amplitude varying from 0.4 ppb (for the US average) up to 1.8 ppb (for the southeastern US) per standard deviation of the Niño 3.4 index (our ONI index). Also, Fig. 1b in Xu et al. (2017) shows that the correlation coefficient between monthly ozone anomalies and the Niño 3.4 index are negative across the whole US. Our calculated correlation between US annual mean ozone anomalies and the ONI index are also negative (Figure 6c).

Xu et al. (2017) showed that the largest ENSO influences occur over two southern US regions during fall and over two western US regions during winter to spring. They showed that ENSO affects surface ozone via chemical processes during the warm seasons in the southern regions, where favorable meteorological conditions occur, but via dynamic transport during the cold seasons in western regions, where the ENSO-induced circulation variations are large. Our seasonal and regional AMONI-ozone correlation results (Table 4), with higher correlations during summer and fall over the eastern US regions and during winter and spring over the US western regions, are also consistent with Xu et al. (2017).

We have added more discussion on the seasonal and regional features of AMONI-ozone correlations in the revised Sect. 4.

3. The influence of AMO on U.S. weather varies by season and region. See Sutton et al. [2007] for more details. Also, AMO not only influence the temperature but also the subsidence, precipitation, drought and surface wind. Throughout this paper, the authors only mention the impact on temperature, which is not enough.

We have added discussions of AMO-related and ozone-relevant climate impacts, including temperature, precipitation, droughts and atmospheric circulation, in the revised

## Sect. 2.2.

We have also updated the analysis to explain the AMONI-ozone correlations in the revised Sect. 4. In particular, we show that “A positive AMONI anomaly is associated with increased temperature over the east (Fig. 8(c)), which enhances biogenic emissions, changes chemical reaction rates, and changes atmospheric circulation that overall lead to increased ozone formation/buildup (Jacob and Winner, 2009; Shen et al., 2015; Fu et al., 2015; Xu et al., 2017).”.

4. Does annual mean ozone strongly depend on temperature? It seems the authors mainly mention the influence of temperature changes when they explain the relationship of AMONI and annual mean ozone.

We have revised the third paragraph in Sect. 4:

“Figure 7(a, b) further shows the correlation between de-trended AMONI and de-trended daily and daytime mean ozone in individual grid cells. The AMONI-ozone correlation is positive and statistically significant over most of the eastern US, and it reaches 0.82 over parts of the southeast. A positive AMONI anomaly is associated with increased temperature over the east (Fig. 8(c)), which enhances biogenic emissions, changes chemical reaction rates, and changes atmospheric circulation that overall lead to increased ozone formation/buildup (Jacob and Winner, 2009; Shen et al., 2015; Fu et al., 2015; Xu et al., 2017). AMONI also correlate positively to ozone over the high-altitude west. This is because a negative ONI anomaly (La Niña-like) means a decrease in lower-tropospheric transport of ozone-poor air from the Eastern Pacific (Xu et al., 2017) and a more meandering subtropical jet and strengthened ozone transport from the stratosphere that compensates for weakened transport from Asia (Lin et al., 2015). AMONI correlates negatively to ozone over southern California, likely reflecting reduced temperature there associated with a positive AMONI (negative ONI) anomaly (Fig. 8(b, c)).”

Minor comments.

P1 L32. Should mention that his mechanism only applies to the surface ozone.

We have changed this line to “Chemically, surface ozone is produced in the sunshining daytime and destroyed mainly by nitrogen oxides (NO<sub>x</sub>) at night”.

P1 L35. We usually use “MDA8” instead of “DM8A”.

We have changed “DM8A” to “MDA8” throughout the text.

P2 L1-7. The summary here is too general. Try to give more details or you can just cite two review papers (Jacob and Winner, 2009; Fiore et al., 2015).

We have simplified this sentence to “Previous observational and modeling studies have revealed important impacts of varying climate conditions and anthropogenic precursor emissions on the near-surface daytime, MDA8 or daily mean ozone over the United States (US) (Jacob and Winner, 2009; Fiore et al., 2015).”

P2 L19-24. Move this paragraph to the last paragraph of the introduction part.

Moved and updated.

P2 L25-34. These studies choose to focus on one region or one season due to potential occurrence of ozone episodes. So why should we care about the low ozone part, e.g. the ozone in the wintertime?

The low-level ozone still affects health (Bell et al., 2006; Peng et al., 2013; Yang et al., 2012), as reviewed in the first paragraph of the introduction. In addition, analysis of their trends and underlying driving factors helps to understand the regional and seasonal ozone changes in the context of reductions in anthropogenic emissions and warming in the climate.

P3 L8-9. What is the meaning of “21.3-28.5% of data are missing”? It is unclear to the readers.

We have changed this sentence to “The fraction of hours in any year with missing data

[Printer-friendly version](#)[Discussion paper](#)

ranges from 21.3% to 28.5%.”

P4 L13-15. Please make it clear why you choose these two climate indices?

Please see our response to major comment 1.

P4 L19-22. The summary of the influence of AMO on U.S. weather is too general.

Updated. Please see our response to major comment 1.

P4 L16-23. What definition of AMO is used here? What SST product (ERSST V4 or HadISST?) is used to calculate the AMO index? Does Fu 2015 mention that AMO can enhance the temperature anomaly and ozone production in US? They just speculate that AMO could change the ozone transport between the north and south US. Also to my knowledge, the influence of AMO on US weather varies in different seasons. Please refer to Sutton et al. [2007].

Updated. Please see our response to major comment 1.

Fu et al. (2015) does not mention that AMO could enhance the temperature anomaly and ozone production in US. And we did not imply this. We have modified the text for clarification.

P4 L24-30. What SST product is used here? Why the authors choose to use Nino 3.4 rather than other indices like Nino 1+2, Nino3 or Nino 4?

Please see our response to major comment 1.

P4 L31-36. Please specify the detrending method here. Is there a reason that you want to subtract the influence of ONI from AMO?

The detrended AMO and ONI indices were downloaded from the National Oceanic and Atmospheric Administration (NOAA) website. The method used for these two indices are linear detrending (Enfield et al., 2001).

The negative sign for ONI in the formula accounts for the negative correlation between



de-trended ozone and ONI anomalies (see more discussion of AMONI in Sect. 2.2).

P8 L33. Change “Relation” to “Relationship”.

Modified.

P8 L35-37. Why do you use linear detrending given the fact that the domestic NO<sub>x</sub> emissions are not linearly decreased?

Here we would like to show the relationship between interannual ozone variability and interannual climate variability. Linear de-trending allows to keep all signals of interannual fluctuations other than linear trend.

P9 L13-16. The evidence shown here is not sufficient to support that positive AMONI leads to enhanced temperatures. This may be just a coincidence. None of the three references listed here can support the conclusion here.

We did not intend to imply the causality between AMONI and temperature. Rather, we regarded AMONI as an indicator of large-scale climate variability that is also associated with temperature. We put in the revised Sect. 4 that “A positive AMONI anomaly is associated with increased temperature over the east (Fig. 8(c)), which enhances biogenic emissions, changes chemical reaction rates, and changes atmospheric circulation that overall lead to increased ozone formation/buildup (Jacob and Winner, 2009; Shen et al., 2015; Fu et al., 2015; Xu et al., 2017).”

P9 L17. Can the mechanism proposed in Xu et al. [2017] be able to explain the relationship? Xu’s mechanism only applies to the springtime ozone decreases in a small region in the southwest US, while the authors here are discussing the annual mean ozone.

Please see our response to major comment 2.

P9 L26-29. It is good to try AO, PDO and NAO. But there are also a lot of other climate indices. This doesn’t mean AMONI is the best. See the metrics used in Shen et al.

[2017].

Please see our response to major comment 1. Further, we have discussed Shen et al. (2017) in the revised Sect. 2.2.

P10 L16-24. The reason is that a heatwave swept across much of the US in 2012. The authors may need to mention this here.

Updated. Thanks for the suggestion.

P11 L9. Do you use the emissions in 2004 here?

Yes, we fixed global anthropogenic emissions at the 2004 levels here.

P12 L19-27. I appreciate the author's efforts in quantifying the contribution of anthropogenic emissions and meteorological variability. But are the influences of these two processes independent? Why  $R$  is used here? Is it better to use  $R^2$ ?

We expect the two mechanisms to be largely independent here, as typically assumed. We have redefined the contribution of anthropogenic emission as  $C_{anth} = (R_{anth}^2) / ((R_{anth}^2 + R_{clim}^2))$ , which has led to stronger contrast between the contributions of emissions and climate. The revised Fig. 11 shows that the emission contribution dominates in the nighttime hours (relatively constant at about 70%), with a reduction in the morning hours, an increase in the late afternoon hours, and a minimum value (at 10%) around 15:00.

Please also note the supplement to this comment:

<https://www.atmos-chem-phys-discuss.net/acp-2017-659/acp-2017-659-AC2-supplement.pdf>

Interactive comment on Atmos. Chem. Phys. Discuss., <https://doi.org/10.5194/acp-2017-659>, 2017.