

Dear Editor, dear referees,

I thank the first referee for the acceptance of the manuscript and the second referee for their very conscientious reading of the whole paper pointing out some final inconsistencies introduced during the first phase of the reviewing process. The first 4 main remarks of the second referee are answered in the first part of this response, whereas the minor points are then clarified. The new version of the manuscript including track changes is added after the answers, so that the reviewer can rapidly catch the modifications.

1) **Neglecting lifting processes other than thermally induced, convective transport.**

- The authors do agree that, since the ABL-TopoIndex is presently only verified with aerosol measurements, it is mostly representative of the aerosol layer. Consequently we changed the title to “*The topography contribution to the influence of the aerosol layer at high altitude stations*”. The authors however still hope that the ABL-TopoIndex will be correlated with gaseous species measurements in future studies, allowing to enlarge its validity as representative of the ABL influence.
- Moreover, considering the referee’s completely correct comment about the lifting mechanism with precipitation, the authors modified the discussion by adding more detailed explanations in § 4.1 and 4.3:

1. P. 16-17: “*Zellweger et al. (2003) concluded that, in contrast to NO<sub>y</sub>, the major process for upward transport of aerosol is the thermally induced vertical transport, confirming that the aerosol parameters used in this study should be in most cases good tracers for ABL influence. However, one has to remain conscious that many lifting processes co-occur with precipitation involving hence aerosol washout. The present validation of the ABL-TopoIndex by aerosol measurements at high altitude stations is consequently valid only for thermally induced processes without precipitation. ABL events involving air masses with low aerosol concentrations –for example synoptic lifting or foehn- were not directly analyzed.*”
2. P. 20: “*However, the aerosol parameters and, particularly, the absorption coefficient cannot be considered as unique tracers of the ABL, particularly in case of lifting processes with precipitation. Analysis of other ABL marker (gaseous species, radon, wind turbulences, etc.) can provide information on additional transport mechanisms which would allow for refinement of this topographic analysis by adding further parameters*”

2) **Emissions**

- Our first answer was somewhat imprecise on this subject. The authors do agree that the amplitude of the diurnal cycle will be influenced by the aerosol concentration emitted into the ABL and consequently by the emission sources in the case of a high altitude site situated part of the time in the FT. In case of a large influence of the residual layer during the night, the dependence of the diurnal cycle amplitude on the emission sources will be largely decreased or even suppressed. The timing of the diurnal cycle will however not be affected by local pollutant sources.

- The reason why ZEP, SUM and NCOS were considered as outliers is not at all due to the low aerosol concentration or the low emissions near the station. As explained in the paper (see 3.5 first §) the topographical environments of SUM and NCOS (high plains that cannot be considered as a complex topography) lead to very high ABL-TopoIndex that are considered as outliers. The exclusion of ZEP is due to its very bad representation in GTOPO30: ZEP is only 306 higher than its grid cell altitude, but this corresponds to 65% of its altitude and also explains its very high ABL-TopoIndex and its outlier status.
- The referee completely identifies the problem of the pollutant sources when he says that the source inventory should include a factor depending on the altitude and on the horizontal distance. These altitude and horizontal distance factors would however also depend on the time of the year and/or the surface temperature in order to take into account the seasonal variability of the ABL height. For example in the alps, wood heating leads to very high aerosol concentration during winter when the ABL influence to high altitude sites are so low that they can be neglected. This is a great subject for a future study. The authors have consequently developed the part of the discussion about the emission problem on p. 16 :

*“Because there are variable pollution levels in the vicinity of the stations, a single absolute value of a pollutant cannot be used to evaluate the ABL-TopoIndex (or ABL influence in general) when considering multiple high altitude stations. An inventory of the proximate pollution sources bounded to a 3D thermodynamic model adapted to complex topographies would be required before using absolute pollutant concentrations as indicators of ABL influence at high altitude sites. Another possibility would be to weight the pollution source inventories by factors depending on their vertical and horizontal distances to the high altitude stations, as well as on a seasonal parameter representing the potential ABL height. A further use of DBinv to restrict the area of potential pollution sources could also be envisaged since this parameter describes the domain from which pollutants can reach the high altitude station by convection and without crossing topographical barriers. This delicate issue can however be avoided by instead considering dynamical parameters such as the various temporal cycles.”*

### 3) **ABL events:**

- **Skewness:** The aerosol parameters used here have a distribution that is neither normal nor lognormal but a Johnson distribution, even if it can be considered as lognormal for some applications (but not all). The skewness is also defined for lognormally distributed data. The skewness of the data and of the logarithm of the data were calculated for the JFJ station for which data of different instruments measuring the same quantity are available. For all aerosol parameters and all instruments, the skewness of the logarithm of each parameter is always negative, leading to the conclusion that the low concentration events are more frequent than the high concentration events. The data skewness (without taking the logarithm) clearly depends on the instrument and on the wavelength. For example, the absorption coefficient is measured at the JFJ by 3 different instruments, two Aethalometers (AE31 and AE33) measuring at 7 wavelengths and the MAAP at 637 nm. The absorption coefficient skewness at similar wavelengths (between 630 and 660 nm) varies from -7.9 to 22.8 among the three instruments. For the Aethalometers, the skewness varies between 5.1 to 6.7

(AE31) and from 9.8 to -7.9 (AE33) as a function of the wavelength. In conclusion, the skewness results depend mostly on the resolution (and artifact) of the instruments and cannot be used to make any assessments about the frequency of pollution events.

- 5 - **The diurnal cycles** are not averaged over a whole year (see §2.4 and supplement 1), but for each month. This allows us to find the minimal and maximal amplitude of the diurnal cycles over the entire year. These minima and maxima of the amplitude of the diurnal cycles are used in Fig. 9, as well as the amplitude of the seasonal cycle. The seasonal variation of the energy available for convection is consequently taken into account and is not the reason for the difficulties in statistically extracting the cycles from the time series. This was not so clearly described in the previous version of the results section so that the following sentence was added to p. 15:

10 *“The diurnal cycles were calculated for the 12 months, so that Dmin and Dmax correspond to the lowest and the greatest monthly amplitudes, respectively.”*

- 15 - The authors do agree that the difficulties in evaluating the diurnal cycle are not fully described in the discussion so that they add further comments in § 4.1:

20 *“However, the statistical determination of the diurnal and seasonal cycle amplitudes suffer from several difficulties: 1) the low aerosol concentration at high altitude often results in measurements near the detection limit leading to high uncertainties, 2) the high hourly autocorrelation of the data require a pre-whitening procedure (see supplement) in order to be able to detect the diurnal and seasonal cycle, 3) meteorological conditions (e.g., cloud coverage, precipitation or seasonal fluctuations) modify the theoretical cycles and lead to a broadening of the time of the extrema. These difficulties make obtaining clear statistical cycles another reason contributing to the observed low correlations.”*

25 4) **Topographic parameters and their selection:**

- 30 - **DBinv:** The referee is still not convinced that DBinv is a necessary parameter of the ABL-TopoIndex. The authors allow themselves to argue once again in favor of DBinv:
- a. One of the referee’s arguments is that “there is no reason what so-ever why a hydrological divide should also hinder atmospheric flow” and the authors do agree with this sentence. The main idea to introduce the DBinv is to determine from where air masses can reach the high altitude station only due to convection, that is without crossing topographical barriers. In that sense, DBinv is a kind of “reservoir” not for air convection (that does not exist, as explicitly mentioned by the referee), but for pollutants that could reach the high altitude sites during long fair-weather periods without precipitation. Let’s take the unrealistic example of a summit encircled by a continuous ridge at higher altitude than this isolated summit. The potential influence of ground emitted pollutants will be lower than if the ridge is not continuous but is lowered at several places by valleys. DBinv will take this difference into account much better than Hypso% or hypsoD50.
  - b. As explained in the discussion (§ 4.3), the present study uses only aerosol measurements to validate the ABL-TopoIndex. The authors however expect that the ABL-TopoIndex will be further validated with gaseous species or radon measurements as well as used with other
- 40

5 studies involving, for example, new particle formation. DBinv presents an only marginal correlation with the aerosol parameters and cycles, but could be an important parameter in other contexts, so that the authors still wish to include DBinv as a ABL-TopoIndex parameter. Moreover, the use of DBinv does not decrease the significance of the correlation between the ABL-TopoIndex and the aerosol parameters. In summary, DBinv has a potential for explaining other variables and does not disturb the present analysis.

- 10 c. The very accurate remarks of the referee concerning the use of source inventories clearly assess the difficulties of using these data in a meaningful way. The authors think that DBinv could be used to define the domain whose sources will contribute to the pollutant concentrations at the high altitude sites. This problem could be solved either by using NWP models to calculate a probability density for a region to affect high altitude sites with the disadvantage of using a lot of computer time to obtain a wind climatology for each site and the not always accurate topography. One could also consider only all the sources in the DBinv or, as suggested by the referee, use DBinv as a probability weight for sources to influence the high altitude site, taking into account that transport through advection is not considered.

15 In order to replace the misleading/incorrect description of CBinv as “a reservoir for air convection” and to better explain this concept, its description was modified and it is now referred in the paper as a “drainage basin for thermally lifted pollutants”:

20 *“Parameter 5 – DBinv: Since the air masses have to be thermally lifted from the valleys and plains towards the summit to influence the station measurements, the size of the drainage basin (DBinv) for thermally lifted pollutants can be calculated with standard hydrology tools using an inverse topography, where the altitude Z is changed to  $-Z$  allowing the summit to become a hole. It represents the region from which pollutants such as aerosols can be thermally lifted without crossing any topographical barrier. Figures 4d and 5d are examples of the DBinv calculation for BEO and PYR. The DBinv is related to criterion 4. The ABL influence should increase with increasing size of the convection drainage basin.”*

25 The authors hope that these modifications associated with the developed arguments will convince the referee as well as the editor to accept that DBinv should be kept in the determination of the ABL-TopoIndex. The authors remain however open to further discussion on this particular point.

- 30 - **The PYR example** : The phenomena described by the referee corresponds to an advection, which is not taken into account in our paper dealing with convection. The authors do agree that the convection from low altitude places in the DBinv for BEO is not a usual phenomenon, but it still corresponds to a region from which pollution is more likely to influence the high altitude station, particularly in case of long summer periods with fair weather days when the pollutants stay in the atmosphere for a long time.
- 35 - **Latitude and altitude**: The referee suggested to eventually add the latitude and the altitude as ABL-TopoIndex parameters. One aim of this study is to show that the altitude is not the right parameter to quantify the ABL influence, so that the altitude has not been included into the ABL-TopoIndex.

However the correlation with the altitude was also calculated and results in poor correlations. The latitude is really a predictive parameter, but it does not relate to the topography of the high altitude stations. In that sense, it has to be considered as another independent parameter.

- **G8 and LocSlope:** The authors do agree with the referee that G8 and LocSlope are the two parameters having the greatest correlation since the morphology of mountainous ranges are usually similar at several scale ranges. G8 describes however the slope steepness in the direct vicinity (about 1km or less depending on the latitude) of the station whereas LocSlope characterizes the steepness in a 10 km range. The authors find that it is worth analyzing the steepness locally as well as at a regional scale.
  
- **Regression model :** First, the authors are unaware of a non-parametric regression model allowing to estimate the better proxy variables to explain, for example, the aerosol measurements. Regression models are constructed for normally distributed data or can be applied on the logarithm of the data. In our case, the aerosol parameters have a Johnson distribution that can be approximated by a lognormal distribution in some cases, but this approximation can also lead to clear errors. Second, the authors hope that the ABL-TopoIndex will be tested with other measured parameters and not only with aerosol optical properties and number concentration. For example, new particle formation, radon and gaseous compounds could be also used. In that sense, the ABL-TopoIndex should not be restricted to the parameters explaining the aerosol measurements, so that a restriction of the ABL-TopoIndex parameters to the those presenting a high correlation with the aerosol measurements is not desirable.

The type of the aerosol distribution was added to the manuscript on p. 14:

*“In order to have a robust estimate of the correlation between the aerosol measurements (following a Johnson distribution) and the topographical parameters (following a normal or a log-normal distribution depending on the parameters) the Spearman’s rank correlation was calculated.”*

#### 5) **Further minor comments:**

**P2, L2:** “drainage basin for air convection”. Once more there is no such thing and the analogy is not valid. Since I suggest dropping parameter DBinv this term and concept can also be removed from the manuscript altogether. See the discussion above.

- As developed under §4) of this document, the authors do agree that the expression “drainage basin for air convection” is not adequate and have changed it to “drainage basin for thermally lifted pollutants”. The explanation of this parameter was also modified in the manuscript (p. 7-8). While remaining open for a further discussion, the authors do consider DBinv as a potentially important parameter for further correlation studies.

**P6, L5f:** Not clear what the accuracy refers to: the altitude differences between sites and GTOPO or the specification of GTOPO? However, what you should be worried about is not the accuracy of GTOPO which refers to the average altitude in each grid cell but the representativeness of GTOPO for the whole grid cell and the resulting mismatch with your station altitudes. So the results in Table S1 are the more important numbers.

- The accuracy refers to the specification of GTOPO. The referee is right in the sense that the important point is really the representativeness of GTOPO. As already commented in the first review round, further studies should use topographical models with lower grid size in order to better represent the reality.

I am also confused about the new text inserted in the revised manuscript. It is not the same as given in the replies (see page 14 of replies), which discusses the altitude differences in much more detail. Just there it is said that the altitude difference for NCOS is 1100 m, which is not true (see Table S1). However, some of the other large altitude differences should be mentioned in the manuscript not just in the reply.

- Table S1: Thank you for pointing this out. It is the station LAN and not NCOS (also called NAM) that has the greatest negative altitude difference with the GTOPO altitude. The text was consequently modified:

*“Due to its location at 1.2 km horizontal distance from the Yala peak (5500 m), LAN’s altitude (3920 m) is 1110 m lower than its DEM grid point altitude (by 28%) and this corresponds to the largest overestimation of the station altitude.”*

- The text given on page 14 of the answer was inserted into the supplement. The authors think that it is too much detailed to be part of the manuscript, which was however complemented (in this version of the manuscript) with the following information (p. 6):

*“Due to the altitude averaging over each grid cell, there is typically an altitude difference between the true station altitude and its corresponding grid location. The mean and median of the differences between the station altitude and its representative grid point are 190 m (8.6%) and 140 m (5.8%). For stations situated near the summits, the difference can be significant: five stations have an underestimation of their altitude greater than 500 m corresponding to 15-32 % (see the supplement and particularly Table S1 for further details), despite, according to the manual, the high reported GTOPO30 accuracy (minimal accuracy of 250 m at 90% confidence level with a RMSE of 152 m).”*

**P7, L5:** Why are these two sites discussed when there are sites like WLG and MUK also showing larger hypso%? There are many more sites with large values as seen in Table S1. So the statement that most sites have hypso% smaller 5 % is not true! Actually 22 of 46 sites have values larger 5 %. Please be more precise in all your discussion! Don't let the reviewers do your work!

- The stations mentioned at this place in the text are the ones taken as examples on the plot and were not intended at all as a description of all station hypso% values. The limit of 5% was taken from the analysis on the larger domain size (750km x 750 km) and was unfortunately not changed in the revised manuscript. The manuscript was consequently modified (p. 7):

*“Figure 3a presents several normalized hypsometric curves with dots indicating the station hypsometric value. While most of the high altitude stations (65%) have hypso% values less than 10%, some stations are situated on wide inflection points of the hypsometric curve (see for example PYR and NCOS on Figure 3a). Six stations (see Table S1) have hypso% larger than 50%.”*

**P7, L14:** 6 out of 46 sites are situated under hypso50. What is the consequence? Should they even be discussed? The new sentences just before suggest that these sites cannot claim to be high altitude stations!

- As explained in the paper, a very small value of 10 was attributed to the 6 stations with negative hypsoD50. The main consequence is that these 6 stations have the same hypsoD50 even if the negative values are fall between -120 m and -872 m. In order to take into account the difference in hypsoD50 negative values, we introduce a new parametrization in the new version of the manuscript: instead of replacing the negative hypsoD50 by a constant value of 10, the value of  $1000/\text{abs}(\text{hypsoD50})$  is taken. The only consequence is to increase, to some extent, the ABL-TopoIndex of these stations that are already the stations with the highest ABL-TopoIndex values. This new parametrization induces no change in the correlation analysis because a method based on rank is used to determine that.
- If we consider the real altitude of the station instead of the GTOPO altitude, there is only 3 stations having a negative hypsoD50 (HLE: -392, LQO: -68 m and TLL:-97 m). Due to our decision to average over 9 grid cells for this parameter, it is not possible to use the real station altitude since that is not clearly defined for the adjacent grid cells.

**P7, L16f:** This is a hypothesis so far not a fact. Please rephrase.

- The sentence was changed to “*The ABL influence should decrease with increasing values of hypsoD50*”.

**P7, L23:** A “plateau” is a high plain (<https://en.wikipedia.org/wiki/Plateau>). So here the word valley or plain should be used instead.

- The word “plateau” was changed with “high plain” in the whole text, but for “the Tibetan plateau” that is a tradition expression used in geography.

**P9, L30f:** This argumentation is confusing and misleading. First of all it is not the hydrological drainage basin that is used in the ABL index but the inverse drainage basin (5d rather than 5c). Then the size of the basin is made responsible for the high ABL index. However, just above we have seen that a large inverse drainage basin does not lead to a high ABL index (inverse drainage basin for BEO is actually larger than the one for PYR). So I would say that one of the other parameters is the more important one in terms of this discussion, most likely hypso% which differs by 3 orders of magnitude between the two sites. However, the discussion of DBinv will need to be removed from the manuscript anyway.

- The authors wanted to say that PYR has a high ABL-TopoIndex not only due to the large DB but also due to its position at the foot of Mount Everest. The authors really referred to the hydrological drainage basin of Fig. 5c, since high altitude stations are frequently at the edges of several drainage basin due to their dominant position. The sentences were misleading and was changed:  
“*PYR (5079 m) is the second highest station considered here, but the station is located at the foot of Mount Everest (8848 m) at a confluence point of several valleys (Fig. 5a and b). Figure 5c shows that PYR is situated in the middle of a very large hydrological drainage basin enhancing the fact that PYR station isn't placed at a dominant position. The PYR ABL-TopoIndex is consequently quite high (3.43) and supports the observation of a large ABL influence in the Himalaya region (Bonasoni et al., 2008).*”
- In that §, no mention of DBinv are made.

**P10, L6:** This should be 50 x 50 to 1000 x 1000 km<sup>2</sup>.

- If the mentioned page and line are correct, the size given in the manuscript corresponds to the sizes on which G8 and LocSlope are computed and not to the various sizes on which the ABL-TopoIndex was tested.

5 **P13, L6:** MWO supposed to have a “low DBinv”, but a value of 127457 for DBinv at MWO means that the site has a DBinv larger than the mean, larger than the median and even larger than the 75% per-centile of all sites (see Table S1).

- The referee is right, this is a mistake. MWO has one the lowest hypso% and the greatest LocSlope and hypsoD50 of North America as well as the second greatest G8 of North America. The manuscript was modified:

10 *“MWO is consequently the North American station with the lowest ABL-TopoIndex due to very low hypso% and relatively high G8, LocSlope and hypsoD50.”*

15 **P14, L12ff:** Once more, I can only say that it is surprising that the ABL-Index does not work for such sites that are clearly not high altitude sites. How can we then trust the method to rank sites at different degrees of high altitude? One common factor at all three sites is the absence of local to region emissions! Hence, aerosols no longer are a good estimator of FT vs ABL conditions.

- The reason why the SUM and NCOS stations are discarded is that the topography at these places is not a “complex topography”. Both stations are really situated on high plains and the ABL-TopoIndex is usable only in a “mountain” context and not in a “plain” context. This is clearly explained in the manuscript (§ 3.5).
- ZEP was not included at all at the beginning of the study. ZEP is also usually not presented at all as a “high altitude” station. In Gawsis (<https://gawsis.meteoswiss.ch/GAWSIS/index.html#/search/station/stationReportDetails/459>), the ZEP station is described as “land”. It was just my own wish to test the limits of the ABL-TopoIndex by introducing some middle altitude stations for comparison purpose. I tried to clearly describe the peculiar status of these sites in the manuscript (§ 2.1 and 3.5).

25 **P15, L9ff:** One should mention that these two parameters are actually strongly correlated ( $r=0.76$ , spearman)! The highest correlation between all parameter pairs! So it is no surprise that they both show similar characteristics and actually one could be sufficient for this study.

- As described on p.2 of this document, G8 and LocSlope both correspond to the steepness of the slope but at different ranges (about 1 and 10 km, respectively). The authors think that keeping both parameters could be important for various phenomena not yet studied, such as air lifting improving the new particle formation (G8) or valley winds (LocSlope).

## The topography contribution to the influence of the **atmospheric boundaryaerosol** layer at high altitude stations

Martine Collaud Coen<sup>1</sup>, Elisabeth Andrews<sup>2,3</sup>, Diego Aliaga<sup>4</sup>, Marcos Andrade<sup>4</sup>, Hristo Angelov<sup>5</sup>, Nicolas Bukowiecki<sup>6</sup>, Marina Ealo<sup>7</sup>, Paulo Fialho<sup>8</sup>, Harald Flentje<sup>9</sup>, A. Gannet Hallar<sup>10,11</sup>, Rakesh Hooda<sup>12,13</sup>, Ivo Kalapov<sup>5</sup>, Radovan Krejci<sup>14</sup>, Neng-Huei Lin<sup>15</sup>, Angela Marinoni<sup>16</sup>, Jing Ming<sup>17,18</sup>, Nhat Anh Nguyen<sup>19</sup>, Marco Pandolfi<sup>7</sup>, Véronique Pont<sup>20</sup>, Ludwig Ries<sup>21</sup>, Sergio Rodríguez<sup>22</sup>, Gerhard Schauer<sup>23</sup>, Karine Sellegri<sup>24</sup>, Sangeeta Sharma<sup>25</sup>, Junying Sun<sup>26</sup>, Peter Tunved<sup>14</sup>, Patricio Velasquez<sup>27</sup> and Dominique Ruffieux<sup>1</sup>

10 <sup>1</sup>Federal Office of Meteorology and Climatology, MeteoSwiss, 1530 Payerne, Switzerland

<sup>2</sup>University of Colorado, CIRES, Boulder, Colorado, 80305, USA

<sup>3</sup>National Oceanic and Atmospheric Administration, Earth System Research Laboratory, Boulder, Colorado, 80305, USA

<sup>4</sup>Laboratory for Atmospheric Physics, Institute for Physics Research, Universidad Mayor de San Andres, Campus Universitario Cota Cota calle 27, Edificio FCPN piso 3, La Paz, Bolivia

15 <sup>5</sup>Institute for Nuclear Research and Nuclear Energy, 1784 Sofia, Bulgaria

<sup>6</sup>Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen PSI, Switzerland

<sup>7</sup>Institute of Environmental Assessment and Water Research, c/ Jordi-Girona 18-26, 08034 Barcelona, Spain

<sup>8</sup>Instituto de Investigação em Vulcanologia e Avaliação de Riscos - IVAR, Rua da Mãe de Deus, 9500-321 Ponta Delgada, Portugal

20 <sup>9</sup>Deutscher Wetterdienst, Met. Obs. Hohenpeissenberg, D-82383 Hohenpeissenberg, Germany

<sup>10</sup>Department of Atmospheric Science, University of Utah, Salt Lake City, UT, USA

<sup>11</sup>Storm Peak Laboratory, Desert Research Institute, Steamboat Springs, CO, USA

<sup>12</sup>Finnish Meteorological Institute, P.O. Box 503, 00101 Helsinki, Finland

<sup>13</sup>The Energy and Resources Institute, IHC, Lodhi Road, New Delhi -110003, India

25 <sup>14</sup>Department of Environmental Science and Analytical Chemistry (ACES), Atmospheric Science Unit, S 106 91 Stockholm, Sweden

<sup>15</sup>Department of Atmospheric Sciences, National Central University, Taoyuan, Taiwan

<sup>16</sup>Institute of Atmospheric Sciences and Climate, National Research Council of Italy, 40129, Bologna, Italy

<sup>17</sup>Max Planck Institute for Chemistry, Mainz 55128, Germany

30 <sup>18</sup>Guest at the State Key Laboratory of Cryospheric Sciences, Chinese Academy of Sciences, Lanzhou 730000, China

<sup>19</sup>Hydro-Meteorological and Environmental Station Network Center (HYMENET), National Hydro-Meteorological Service (NHMS), Hanoi, Vietnam

<sup>20</sup> Université Toulouse III - Laboratoire d'aérodynamique UMR 5560, 31400 Toulouse- France

<sup>21</sup> German Environment Agency, Platform Zugspitze, GAW-Global Observatory Zugspitze/Schneefernerhaus, Zugspitze 5, 82475 Zugspitze

<sup>22</sup> Izaña Atmospheric Research Centre, AEMET, Joint Research Unit to CSIC “Studies on Atmospheric Pollution”, Santa Cruz de Tenerife, Spain.

<sup>23</sup> Sonnblick Observatory, Zentralanstalt für Meteorologie und Geodynamik (ZAMG), 5020 Salzburg, Austria

<sup>24</sup> Laboratoire de Météorologie Physique, UMR6016, Université Blaise Pascal, 63170 Aubière, France

<sup>25</sup> Climate Chemistry Measurements Research, Climate Research Division, Environment and Climate Change Canada, 4905 Dufferin Street, Toronto, M3H 5T4 Canada

<sup>26</sup> Key Laboratory of Atmospheric Chemistry of CMA, Chinese Academy of Meteorological Sciences, Beijing 100081, China

<sup>27</sup> Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland

15 *Correspondence to:* Martine Collaud Coen ([martine.collaudcoen@meteoswiss.ch](mailto:martine.collaudcoen@meteoswiss.ch))

**Abstract.** High altitude stations are often emphasized as free tropospheric measuring sites but they remain influenced by atmospheric boundary layer (ABL) air masses due to convective transport processes. The local and meso-scale topographical features around the station are involved in the convective boundary layer development and in the formation of thermally induced winds leading to ABL air lifting. The station altitude alone is not a sufficient parameter to characterize the ABL influence. In this study, a topography analysis is performed allowing calculation of a newly defined index called ABL-TopoIndex. The ABL-TopoIndex is constructed in order to correlate with the ABL influence at the high altitude stations and long-term aerosol time series are used to assess its validity. Topography data from the global digital elevation model GTopo30 were used to calculate 5 parameters for ~~46-43~~ high and 3 middle altitude stations situated in five continents. The geometric mean of these 5 parameters determines a topography based index called ABL-TopoIndex which can be used to rank the high altitude stations as a function of the ABL influence. To construct the ABL-TopoIndex, we rely on the criteria that the ABL influence will be low if the station is one of the highest points in the mountainous massif, if there is a large altitude difference between the station and the valleys or ~~plateaus~~ high plains, if the slopes around the station are steep, and finally if the drainage basin for ~~air convection-~~ thermally lifted pollutants is small. All stations on volcanic islands exhibit a low ABL-TopoIndex whereas stations in the Himalaya and the Tibetan Plateau have high ABL-TopoIndex values. Spearman's rank correlation between aerosol optical properties and number concentration from 28 stations and the ABL-TopoIndex, the altitude and the latitude are used to validate this topographical approach. Statistically significant (s.s.) correlations are found between the 5 and 50 percentiles of all aerosol parameters and the ABL-TopoIndex whereas no s.s.

correlation is found with the station altitude. The diurnal cycles of aerosol parameters seem to be best explained by the station latitude although a s.s. correlation is found between the amplitude of the diurnal cycles of the absorption coefficient and the ABL-TopoIndex.

## 1. Introduction

5 Climate monitoring programs aim to measure climatically relevant parameters at remote sites and to monitor rural, arctic, coastal and mountainous environments. The majority of these programs consist of in-situ instruments probing the Atmospheric Boundary Layer (ABL). The high altitude stations provide a unique opportunity to make long-term, continuous in-situ observations of the free troposphere (FT) with high time and space resolution. It is however well-known that, even if located at high altitudes, the stations designed to measure the FT may be influenced by the transport of boundary layer air masses. Remote sensing instruments can be used to complement in-situ measurements in order to provide more information about the FT. For example, sun photometers measure aerosol optical depth of the integrated atmospheric column including the FT although they don't provide vertical information to enable separation of FT and ABL conditions. Light Detection and Ranging (LIDAR) type instruments measure the profile of various atmospheric parameters (meteorological, aerosol, gas-phase) and thus can provide information not only on the ABL but also on the FT. They can be used to detect the ABL and Residual Layer (RL) heights at high altitude stations from a convenient site at lower elevation (Haeffelin et al., 2012; Ketterer et al., 2014; Poltera et al., 2017). These instruments are however limited in the presence of fog and low clouds and they don't measure above the cloud cover. Further, the use of LIDAR to attribute the various aerosol gradients to ABL layers remains a delicate problem. Lastly, few LIDAR instruments are currently installed in regions of complex topography. Instrumented airplanes can make detailed measurements of the vertical and spatial distribution of atmospheric constituents and are used either during limited measurement campaigns or on regular civil aircraft (see for example the IAGOS CARIBIC project, <http://www.caribic-atmospheric.com/>), but, because of the limited temporal scope of most measurement campaigns, cannot provide long-term, continuous context for the measurements. Ideally, to make FT measurements, a combination of these techniques would be used, but due to limited resources that is rarely possible. Thus, it is important to evaluate the constraints of each technique. The high altitude time series from surface measurements remain the most numerous and the longest data sets to characterize the FT and its evolution during the last decades. Here we focus on identifying factors controlling the influence of ABL air on high altitude surface stations hoping to sample FT air.

The ABL is the lowest part of the atmosphere that directly interacts with the Earth's surface and is most of the time structured into several sub layers. In the case of fair-weather days, the continental ABL has a well-defined structure and diurnal cycle leading to the development of a Convective Boundary Layer (CBL), also called a mixing or mixed layer, during the day and a Stable Boundary Layer (SBL) which is capped by a Residual Layer (RL) during the night (Stull, 1988). During daytime, the aerosol concentration is maximum in the CBL and remains high in the RL. During nighttime, the surface-emitted species accumulate in the SBL. In the case of cloudy or rainy conditions as well as in the case of advective

Formatted: Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 0.63 cm

weather situations, free convection is no longer driven primarily by solar heating, but by ground thermal inertia, cold air advection and/or cloud top radiative cooling. In those cloudy cases the CBL development remains weaker than in the case of clear sky conditions. Long range or RL advection can however lead to a high aerosol concentration above the CBL during daytime, leading to high altitude aerosol layers (AL) that can be decoupled from the CBL and the SBL.

5 There are several rather complex mechanisms able to bring ABL air up to high altitude (Rotach et al., 2015; Stull, 1988; De Wekker and Kossmann, 2015). An important factor in many of these mechanisms is how the CBL develops over mountainous massifs. In their extensive review of concepts, De Wekker and Kossmann (2015) studied the CBL development over slope, valley, basin, [plateau-high plain](#) as well as over complex mountainous massifs and concluded that the CBL height behavior can be categorized into four distinct patterns describing their spatial extent as a function of the surface topography: 10 the hyper-terrain following, the terrain following, the level and the contra-terrain following. The type of CBL height behavior depends on several factors such as the atmospheric stability, synoptic wind speed and vertical and horizontal scale of the orography. Stull (1992) concluded that the CBL height tends to become more horizontal (level behavior) at the end of the day, that deeper CBLs are less terrain following than shallower ones, and that the CBL top is less level over orographic features with a large horizontal extent. Even if the CBL height remains lower than the mountainous ridges, thermally driven 15 winds develop along slopes, or in valleys or basins and these winds are able to bring ABL air masses up to mountainous ridges and summits. These phenomena were extensively modeled (Gantner et al., 2003; Zardi and Whiteman, 2012) and also measured (Gantner et al., 2003; Rotach and Zardi, 2007; Rucker et al., 2008; Venzac et al., 2008; Whiteman et al., 2009) and are part of the active mountainous effects allowing a vertical transport of polluted air masses to the FT. For example, a Continuous Aerosol Layer (CAL) is often measured above the CBL during dry, clear-sky and convective synoptic situations 20 (Poltera et al., 2017). Finally, ABL air masses can also be dynamically lifted by frontal systems, deep convections or foehn as well as be advected from mesoscale or wider regions and influence high altitude measurements by all these atmospheric processes.

The ABL influence of the mesoscale regions at high altitude sites were directly shown by airborne LIDAR measurements over the Alps and the Apennine (Nyeki et al., 2000, 2002; De Wekker et al., 2003) and more indirectly by the seasonal and 25 diurnal cycles of aerosol parameters at high altitude stations (Andrews et al., 2011). Many methods have been used to separate FT from ABL influenced measurements, including those based on time of day and time of year approach (Baltensperger et al., 1997; Gallagher et al., 2011), wind sectors (Bodhaine et al., 1980), the vertical component of the wind (García et al., 2014), wind variability (Rose et al., 2016), NO<sub>x</sub>/NO<sub>y</sub>, NO<sub>y</sub>/CO ratios or radon concentrations (Griffiths et al., 2014; Herrmann et al., 2015a, 2015b; Zellweger et al., 2003) and water vapor concentrations (Ambrose et al., 2011; Obrist et al., 2008), although none of these methods leads to an absolute screening procedure to ensure the measurement of pure FT 30 atmosphere.

The altitude range of stations which claim they sample in the FT (at least some of the time) spans from about 1000 to more than 5320 m a.s.l., but a simple analysis of the aerosol parameters (for example, the black carbon concentration) as a function of altitude suggests that higher altitude stations are not necessarily less influenced by anthropogenic pollution.

While station altitude may not be the main parameter explaining the ABL influence, topographical features around the station are nevertheless involved in the CBL development and in the formation of thermally induced winds leading to ABL air lifting (Andrews et al., 2011; Kleissl et al., 2007). In addition to topography there are other important parameters determining the ABL influence at mountainous stations such as the wind velocity and direction, soil moisture and albedo, synoptic weather conditions, pollution sources ~~and as well as~~ sea surface temperature for islands, but none of these parameters will be considered in this study, which is solely restricted to the analysis of the topographic influence.

The aim of this paper is twofold: (i) to define a topography based index called ABL-TopoIndex that can be utilized to rank the high altitude stations as a function of the ABL influence and (ii) to compare the potential ABL influence of several locations in a mountainous range in order, for example, to choose the best sampling location. ~~Several tools~~ Some concepts ~~tested or~~ used in this study are taken from the hydrology analysis field, since both air and water flow along defined, though (often) different, flow paths. The ABL air masses flow towards high altitudes and can extend into a three dimensional space, in contrast to the downward flow of water that can be restricted to a two dimensional space. However, similar to hydrological concepts, the ABL air mass reservoirs are found in the plains and valleys.

## 2. Experimental

### 2.1. Stations

Forty-three high altitude stations (Table 1 and Fig. 1) were selected based on various criteria, such as the presence of aerosol or gaseous measurements, their representativeness of the mountainous massif and/or the possibility to compare several stations from the same mountainous massif. They are representative of 5 continents and their altitudes range between 1074 (SHN) and 5352 m a.s.l (CHC). Even if clearly situated within the ABL, some stations like HPB, MSY or ZEP were added to this analysis to verify the results of the ABL-TopoIndex at lower altitude sites. Several mountainous massifs such as the Alps, the Himalaya, the Rocky Mountains and the Andes Cordillera are well represented with three to five stations. Some other stations such as BEO in the Balkan Peninsula, HAC in the Peloponnese, WLG in China, PDI in Vietnam, MKN in Kenya and the high plateau-plain of ASK in the Hoggar Mountains of southern Algeria are the only representative of their massif. The volcanic islands form a category in themselves, despite being located in different oceans and at various latitudes.

### 2.2. Topography data and analysis

The topography data were taken from the global digital elevation model GTopo30 (<https://lta.cr.usgs.gov/GTOPO30>). GTopo30 has a horizontal grid spacing of 30 arc seconds corresponding to a spatial resolution between 928 m in the East/West direction at the equator, 598 m at WHI (50° N) and 373 m at the SUM polar station (72.6° N). In the North/South direction, 30 arc seconds are almost constant with latitude and correspond to 921 m at the equator and 931 m at the poles. The geographical coordinate system WGS84 (World Geodetic System revised in 1984) from GTopo30 was projected in the Universal Transverse Mercator (UTM) conformal projection to ensure homogeneity in vertical and horizontal coordinates.

**Formatted:** Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 0.63 cm

**Formatted:** Indent: Left: 0 cm, Hanging: 0.75 cm, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.63 cm + Indent at: 1 cm

**Formatted:** Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 0.63 cm

**Formatted:** Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.63 cm + Indent at: 1 cm

**Formatted:** Indent: Left: -0.01 cm, Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 1.4 cm

Due to the altitude averaging over each grid cell, there is typically an altitude difference between the true station altitude and ~~their~~-its corresponding grid location. The mean and median of the differences between the station altitude and its representative grid point are 190 m (8.6%) and 140 m (5.8%). For stations situated near the summits, the difference can be significant: ~~five stations have an underestimation of their altitude greater than 500 m corresponding to 15-32 % (see the supplement and particularly Table S1 for further details), even if despite, according to the manual, the high reported~~ GTOPO30 accuracy ~~remains high~~ (minimal accuracy of 250 m at 90% confidence level with a RMSE of 152 m). If not specified, the given altitudes correspond to altitude above sea level (a.s.l.).

The TopoToolbox-master version of the free shareware TopoToolbox (<https://topotoolbox.wordpress.com/>), which is a set of matlab functions offering analytical GIS utilities in a non-GIS environment (Schwanghart and Scherler, 2014), was used as a principal tool for the topographic relief and flow pathways analysis in the Digital Elevation Model (DEM) analysis. The DEM were preprocessed by filling holes with a carving process prior to calculate the flow directions and the water flow paths were calculated with the single flow direction representation.

### 2.3. ABL-TopoIndex

To construct the ABL-TopoIndex, we rely on the following four criteria to indicate that the ABL influence will be low if:

- 1) the station is one of the highest points in the mountainous massif,
- 2) there is a large altitude difference between the station and the valleys, ~~plateaus~~-high plains or the average domain elevation,
- 3) the slopes around the station are steep, and
- 4) the «drainage basin» for ~~air convection~~-thermally lifted pollutants is small.

Based on these criteria, the red station on Fig. 2 will be less influenced by the ABL than the blue station, despite being situated at lower altitude. A quantitative estimation of these criteria depends clearly on the domain considered. The minimal size requirement for such a topographical analysis is that the domain should contain the whole mountainous massif. An airborne Lidar measurement of the ABL over the Alps (Nyeki et al., 2002) clearly stated that the convective boundary layer is formed over a large-scale and leads to an elevated and extended layer. Nyeki et al. (2002) also quantified this “large-scale” to extend more than 200 km from the mountainous massif. A rectangular domain size of 500 km x 500 km centered on each site was thus chosen for this analysis (see § 3.2 for a discussion of the effect of the domain size). These four criteria listed above are then represented using five parameters (Table S2 lists topographical and hydrological parameters considered but rejected for this analysis):

1. **Parameter 1 – hypso%:** A hypsometric curve is the cumulative distribution function of elevation on the considered domain. The frequency percentage of the hypsometric curve at the station altitude (hypso%) provides a representation of criterion 1 for a large spatial scale. Figure 3a presents ~~some~~-several normalized hypsometric curves with dots indicating the station hypsometric value. While most of the high altitude stations

**Formatted:** Indent: Left: -0.01 cm  
Outline numbered + Level: 2 +  
Numbering Style: 1, 2, 3, ... + Start  
1 + Alignment: Left + Aligned at: 0  
cm + Indent at: 1.4 cm

**Formatted:** Numbered + Level: 1 +  
Numbering Style: 1, 2, 3, ... + Start  
1 + Alignment: Left + Aligned at: 1  
cm + Indent at: 1.9 cm

(65%) have hypso% values less than ~~510%~~, some stations are situated on wide inflection points of the hypsometric curve (see for example PYR and NCOS on Figure 3a). Six stations (see Table S1) have hypso% larger than 50%. ~~are situated respectively at 26% and 65% on wide inflection points of the hypsometric curve.~~ BEO and FWS are found at less than 0.1% of the curve indicating they are located at one of the highest points of their respective mountainous massifs. The ABL influence should increase with increasing value of hypso%.

2. **Parameter 2 – hypsoD50:** The second parameter (hypsoD50) is the difference between the station altitude and the altitude at 50% of the hypsometric curve. The median of the hypsometric curve was chosen first because a station claiming to be a high altitude site should typically be at higher altitude than half of its geographical environment and, second, because the median is a commonly used statistical concept to determine the central value of a sample. The parameter hypsoD50 corresponds to criterion 2 for a large spatial scale. In some cases (see Fig. 3a and Table S1), the station is situated under the 50% of the hypsocurve leading to a negative hypsoD50. For these sites the hypsoD50 is set to the very small value of  $1000/abs(hypsoD50)$  to allow the geometric mean to be applied (see equation 1). The ABL influence ~~should decrease~~ with increasing values of hypsoD50.
3. **Parameter 3 – LocSlope:** The altitude difference between the station and the minima in a circular domain centered at the station is plotted as a function of the domain radius on Fig. 3b. The slope of this curve between 1 km and 10 km is then calculated (LocSlope) and corresponds to criteria 2 and 3 for a small spatial scale. The steepness of the slopes (criterion 3) around the station is only evaluated from the station toward the lowest elevations. The distance of 10 km to calculate the LocSlope was then chosen as representative of the maximal distance to the next adjacent ~~plateau-high plain~~ for almost all stations. Figure 3b shows that the change in the altitude difference as a function of domain radius can be very different from station to station. For example, there is a rapid decrease of the elevation minima with increasing distance that gradually levels off for radius greater than 7 km for JFJ and for radius greater than 4 km for MBO; there is a continuous decrease of the minima elevation for PYR and ASK up to radius larger than 30 km; and there are some steps for CHC and BEO. NCOS appears very different than the other sites plotted since the NCOS station is near the vast Nam Lake (surface area 1950 km<sup>2</sup>) situated at 4718 m. The ABL influence should increase with decreasing LocSlope.
4. **Parameter 4 – G8:** The mean gradient in elevation in the eight directions (N, NE, E, SE, S, SW, W, NW) at the station is called G8. This parameter takes into account the slopes towards lower and higher elevations over a local scale (0.5-1.1 km, which is the distance between by two grid cells, with the size of the grid depending on latitude) and corresponds to criterion 3. The ABL influence should decrease for increasing G8 gradient.
5. **Parameter 5 – DBinv:** Since the air masses have to ~~“flow”~~ be thermally lifted from the valleys and plains towards the summit to influence the station measurements, the size of the drainage basin (DBinv) for ~~convection-thermally lifted pollutants~~ can be calculated with standard hydrology tools using an inverse

topography, where the altitude  $Z$  is changed to  $-Z$  allowing the summit to become a hole. It represents the region from which pollutants such as aerosols can be thermally lifted without crossing any topographical barrier. Figures 4d and 5d are examples of the DBinv calculation for BEO and PYR. The DBinv is related to criterion 4. The ABL influence should increase with increasing size of the convection drainage basin.

5

To summarize, the ABL influence should increase with decreasing values of hypsoD50, LocSlope and G8 and with increasing values of hypso% and DBinv. Thus, to determine the ABL-TopoIndex, the geometric mean is calculated on the inverse of hypsoD50, LocSlope and G8 along with the values of hypso% and DBinv. To avoid any particularities of the station site and due to the fact that the ABL influence is a regional factor, the mean of the values at the grid cell containing the station and at the eight neighboring grid cells (recall that grid spacing is 30 arc seconds) are used to calculate the ABL-TopoIndex. The ABL-TopoIndex is then taken as the geometric mean of the five parameters:

10

$$ABL-TopoIndex = \sqrt[5]{hypso\% \times \frac{1}{hypsoD50} \times \frac{1}{LocSlope} \times \frac{1}{G8} \times DBinv} \quad (1)$$

15

The geometric mean is used here on strictly positive parameters that have widely different numeric ranges (e.g., Table 2). The geometric mean is used instead of the arithmetic mean because it effectively "normalizes" the various parameter ranges, so that no parameter dominates the weighting. Further, a given percentage change in any of the parameters will yield an identical change in the calculated geometric mean value. In that sense the variability of each parameter is also normalized, leading to similar modifications of the ABL-TopoIndex for similar parameter's variations. Because of these properties, the geometric mean is the recommended method to determine a meaningful indices from multiple parameters (Ebert and Welsch, 2004). The extrema, median and mean of the parameters constituting the ABL-TopoIndex are reported in Table 2.

20

The value of the ABL-TopoIndex has no significance in itself, so that the units are not important, but it allows ranking of the stations as a function of the ABL influence due to convection.

#### 2.4 Aerosol parameters

25

Aerosol datasets from 25 high altitude and 3 mid-altitude stations (Table 1) were available for this study, 21 of them coming from GAW (Global Atmospheric Watch) stations. The datasets comprise absorption coefficient, scattering coefficient and/or number concentration and cover time periods ranging from at least one year up to more than one decade of measurement (see supplement Table S3). Stations with time series shorter than one year were not used, since they are not representative of a complete seasonal cycle. Due to the non-normal distribution of the aerosol parameters, the 5, 50 and 95 percentiles were taken as representative of the minimal, central position and maximal concentrations.

30

No correction for standard pressure and temperature was applied in order to use the measured aerosol properties and concentration at high altitude. For consistency, the measured hourly absorption and scattering coefficients were adjusted to a wavelength of 550 nm if reported at a different wavelength using an Ångström exponent of 1. Additionally, the scattering coefficients were corrected for truncation error. Because of their measurement technique and the low aerosol concentrations

at many high altitudes, filter-based photometers regularly measure negative absorption coefficients at some of these sites. Some datasets contain up to 20-30% of negative absorption values. Depending on the data owner's policy, these negative values were either left in the dataset, set to zero (or to a minimal value) or considered as missing values. To ensure a similar treatment for all datasets, negatives values, zeros or minimal values attributed to negatives were therefore set to missing values.

The diurnal and seasonal cycles were only analyzed on datasets longer than 2 years. To be able to statistically calculate the diurnal and seasonal cycles, the autocorrelations at one hour (first lag) were first removed from the dataset by a whitening procedure (Wang and Swail, 2001). The autocorrelations at each lag time were then calculated on the whitened dataset taking into account missing data (see supplement for further explanations). Only autocorrelation values statistically significant at 95% confidence level were kept. Since the diurnal (24 h) and annual cycles (365 days) were not well defined due to variable meteorological conditions and some shorter datasets, the autocorrelation at lags 22 to 26 h and at lags 350 to 380 days were summed to obtain the strength (i.e., the cycle amplitude) of the diurnal and seasonal cycles, respectively. Noise in the aerosol measurements makes the strength of the cycle a somewhat qualitative value. The diurnal cycles were calculated for each month of the year in order to observe the seasonal change of the diurnal cycles.

### 3. Results

#### 3.1. Case studies

Mount Moussala (BEO) is the highest summit not only of Bulgaria but of the whole Balkan massif. The regional GAW station is located at the summit (2925 m). The topographic dominance of BEO can be visualized on the topography map (Fig. 4a). Figure 4a also shows the main hydrological flow paths which follow the Iskar, Martisa and Metsa rivers. Figure 4b shows the water flow accumulations, which are the accumulated flows of all cells flowing into each downslope cell in the output raster, allowing visualization of the greatest features of the Bulgaria hydrographic network. BEO is at the junction of four drainage basins corresponding to the four main rivers (Fig. 4c). Figure 4d shows that when the convection drainage basin is calculated with the inverse topography, BEO is in the center of a large convection drainage basin that covers most of the plotted domain. Even though BEO's altitude is under 3000 m, BEO's ABL-TopoIndex of 0.52 is one of the lowest due to an almost zero hypso% (0.034 %), a high hypsoD50 of 2136 m and a small DBinv of  $1.15 \cdot 10^5$  (Table 2 and S2). HAC is a very similar case to BEO since it is situated almost at the top of Mount Helmos, the third highest mountain of the Peloponnese (Greece).

PYR (5079 m) is the second highest station considered here, but the station is located at the foot of Mount Everest (8848 m) at a confluence point of several valleys (Fig. 5a and b). Figure 5c shows that PYR is situated in the middle of a very large hydrological drainage basin enhancing the fact that PYR station isn't placed at a dominant position. The PYR ABL-TopoIndex is consequently quite high (3.43) and supports the observation of a large ABL influence in the Himalaya region

**Formatted:** Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 0.63 cm

**Formatted:** Indent: Left: -0.01 cm + Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 1.4 cm

(Bonasoni et al., 2008). The daily arrival of polluted air masses from the Indo-Gangetic plain is frequently reported in PYR data analyses (Bonasoni et al., 2010, 2012; Marinoni et al., 2010).

### 3.2. Relation between ABL-TopoIndex and domain size

5 The ABL-TopoIndex depends on the size of the chosen domain (Fig. 6a) so that the various algorithms were tested to several domain sizes ranging from 50 to 1000 km<sup>2</sup>. The gradient G8 and the local slope LocSlope are calculated on small fixed horizontal scales (0.5-1 and 10 km, respectively) and are consequently constant with domain size (Fig 6e,f), although there are small fluctuations due to some distortions occurring during the projection of GTOPO30 in the UTM WGS84 coordinate system, mostly when the analyzed domain extends beyond two UTM zones (see for example BEO). The other  
10 three parameters do change with domain size which is the reason that the ABL-TopoIndex also is a function of the domain area. DBinv tends to increase with the domain size for all stations (Fig. 6b), since the low altitude area potentially contributing to ~~the ABL influence~~ increase the thermally lifted pollutants concentration increases with domain size. The hypso% decreases continuously for stations situated in a dominant position in the ~~ir whole~~ mountainous massif such as JFJ, SBO or BEO (Fig. 6c). For stations located at a lower position in the ~~ir~~ massif (see for example HPB), the hypso% first  
15 increases before decreasing once the domain contains all the highest peaks of the massif. Finally, stations situated atop a high local mountain but surrounded by higher mountains such as MUK (not shown in Fig. 6) have a continuously increasing hypso% up to very large domain sizes (10<sup>6</sup> km<sup>2</sup> for MUK). HypsoD50, the difference between the station elevation and the minimum of elevation in the domain, always increases (or at least stays constant) with domain size but changes more or less rapidly depending on the domain topography (Fig. 6d). In general, the ABL-TopoIndex usually increases with domain size  
20 (i.e., more ABL influence). The greatest increases are usually found for the stations with the highest ABL-TopoIndex at small domain sizes and are due to an increase in DBinv overcoming the decrease in hypso% and the increase in hypsoD50.

### 3.3. Relation between ABL-TopoIndex and altitude

As stated in the introduction, the development of the ABL-TopoIndex relies on the assumption that the station position in the  
25 mountain massif is a better criterion for determining the ABL influence than the station altitude alone. To compare these two parameters, the ABL-TopoIndex is reported as a function of the altitude for all grid cells in a 5km x 5km domain around some stations on Fig. 7. For the grid cells at the highest altitudes, there is a clear dependence between the ABL-TopoIndex and the altitude, with ABL-TopoIndex decreasing (less ABL influence) as altitude increases; Fig. 7 shows that the OMP and PYR regions have ~~the a very steepest~~ and ASK ~~the a very flattest~~ ABL-TopoIndex decrease with altitude. At ~~middle-lower~~  
30 altitudes for each massif, the valleys, high ~~plateaus~~ plains, various mountainous slopes and networks lead to a wide range of the altitudes corresponding to the same ABL-TopoIndex value. For example, the altitude range corresponding to an ABL-TopoIndex of 3 varies between 3000 m and 6000 m at PYR, while at OMP an ABL-TopoIndex of 2 is achieved at an altitude

**Formatted:** Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 0.63 cm

**Formatted:** Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.63 cm + Indent at: 1 cm

**Formatted:** Indent: Left: -0.01 cm + Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 1.4 cm

**Formatted:** Indent: Left: -0.01 cm + Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 1.4 cm

range between 250-350 m. At PYR and CHC, there are discrete groupings of points likely corresponding to the basins of different valleys around the site. The ABL-TopoIndex values of the stations are indicated by the square markers, allowing visualization of their relative situation in their respective mountain massifs: OMP, HAC and CHC and to some extent SBO were constructed at places with the lowest ABL-TopoIndex of their regions thus minimizing potential ABL influence. In contrast the region around PYR (and to a lesser extent ASK) shows locations with much lower ABL-TopoIndex (less ABL influence) at similar altitudes to the stations.

### 3.4. Relation between the ABL-TopoIndex and the station location

The ABL-TopoIndex values for the forty-six stations are grouped on Fig. 8 by continents and mountainous massifs or regions (see Table 1) that can correspond to various geomorphologies. The first obvious observation is that all islands have very low ABL-TopoIndex (note the logarithmic scale for the ABL-TopoIndex), whereas the stations in the Himalaya massif have the greatest ABL-TopoIndex. The values of the ABL-TopoIndex and of all its constituting parameters are given in Table S1. Further conclusions that can be derived include:

- **Islands:** the islands with sites included in this study have a small area, are delimited by the large flat ocean (though most of them are grouped in archipelagos) and their summits were formed by volcanic activities leading to steep slopes. All these factors lead to very low ABL-TopoIndex values. The Teide Observatory in Izaña, an island of the Canary Islands archipelago (TDE) and Pico Mountain Observatory (OMP) in the Portuguese Azores archipelago ranks as the monitoring stations with the lowest ABL influence. The low ABL-TopoIndex of both stations is caused by the following reasons: 1) both mountains are the only summit of the island and the highest mountain (3715 m and 2350 m, respectively) of their archipelago, 2) both islands have small surface area (2034 km<sup>2</sup> and 447 km<sup>2</sup>) and 3) both research stations are just below the mountain summits (177 m and 126 m from the summit). The effect of the proximity to the summit can be clearly seen by the difference between TDE (ABL-TopoIndex of 0.22) and IZO (ABL-TopoIndex of 0.57). These two sites, both located on the island of Tenerife, are separated by only 15 km in horizontal distance but a vertical distance of 1165 m, with TDE being the higher station. Taiwan, where LLN is located, has the greatest surface area (36193 km<sup>2</sup>) of the islands considered here and, additionally, is in close proximity to a continent (China is 130 km to the West). Both of these facts explain LLN's high ABL-TopoIndex in the island category. MLO in Hawaii is at high altitude (3397 m), but the island of Hawaii has a second summit, Mauna Kea (4205 m). Further, the MLO research station is 870 m beneath the volcano top and has a relatively low G8, explaining why it has a higher ABL-TopoIndex than most of the islands. This difference in ABL-TopoIndex between OMP and MLO is confirmed by an almost daily occurrence of buoyant upslope flow at MLO while such flow patterns are much less frequent (<20% of the time) at OMP (Kleissl et al., 2007).
- **Alps:** The European Alps consist of a broad mountainous massif with the highest summits between 4500 and 4800 m. The four high research stations (JFJ, SBO and ZUG/ZSF) are located between 2900 m and 3600 m (ZSF being only some 300 m below ZUG). HPB (985 m) was added to this study as a low elevation station in the Alps. All the

Formatted: Indent: Left: -0.01 cm  
Outline numbered + Level: 2 +  
Numbering Style: 1, 2, 3, ... + Start  
1 + Alignment: Left + Aligned at: 0  
cm + Indent at: 1.4 cm

Formatted: Bulleted + Level: 1 +  
Aligned at: 0.63 cm + Indent at: 1  
cm

three high elevation stations have low ABL-TopoIndex: JFJ (0.64), SBO (1.24) and ZUG (1.35). Their ABL-TopoIndex values are generally a little higher than those determined for the islands. As expected, the ABL influence at HPB is much stronger (ABL-TopoIndex is 5.38) due to both its lower altitude and position near the bottom of the Zugspitze massif.

5 | - **Pyrenees:** the Pyrenees are a natural border between France and Spain and peak at 3400 m. PDM is a high altitude station (2877 m) with an ABL-TopoIndex similar to the European alpine high altitude stations. MSA is located at a mid-altitude range of the massif and has a median ABL-TopoIndex, while the low altitude MSY station, added for comparison purposes, has a high ABL-TopoIndex.

10 | - **Other European stations:** BEO and HAC are situated at the highest points of their massifs and have therefore very low ABL-TopoIndex values, comparable to those of the island high elevation sites. The lower altitudes of CMN and PUY, their middle position in mountainous massifs containing several higher summits and, to a lesser extent, their proximity other massifs such as the Alps and the Pyrenees result in higher ABL-TopoIndex values for these two sites.

15 | - **Himalaya and Tibetan Plateau:** The Himalaya is the highest mountainous massif on Earth with 14 summits peaking at more than 8000 m. The altitude of the research stations between 2200 and 5100 m are therefore at relatively low elevation in comparison to the summits. This is clearly reflected in their high ABL-TopoIndex values (between 3 and 30). MUK and SZZ are both situated in the foothills of the Himalaya in India (Uttarakhand region) and in south China (Yunnan region), respectively, and both have an ABL-TopoIndex value in the 3-10 range. Although MUK is at a lower altitude than SZZ, it is located at a higher position than SZZ relative to the mean altitude of its meso-scale environment. The high ABL-TopoIndex values for HLE and NCOS are due to their position in a large valley and on the edge of a vast lake, respectively, that largely decreases all the parameters related to criteria 1, 2 and 3 (see Sect. 2.3). WLG is constructed within some ten's of meter of Mount Waliguan's summit at the northeastern part of the Tibetan plateau, so that its dominant position in its meso-scale domain leads to a middle range ABL-TopoIndex value.

25 | - **Japan :** Mount Fuji is the highest peak of Japan and the research station is located at the top of the symmetric volcano located near the coast. The second highest peak in Japan is some 500 m lower than Mount Fuji. This particular topography leads to an ABL-TopoIndex similar to the volcanic islands. The two other Japanese stations are at much lower altitudes and have mid-range ABL-TopoIndex values.

30 | - **North America:** Mount Washington Observatory is located in the Presidential Range of the White Mountains. It is the highest peak in the Northeastern United States and the most prominent mountain east of the Mississippi River. MWO is consequently the North American station with the lowest ABL-TopoIndex due to very low hypso% and relatively high G8, LocSlope and hypsoD50 and ~~low DBinv~~. Four stations (MZW, NWR, SPL and YEL) are situated in the Rocky Mountains, whose summits peak at 4400 m. The three stations higher than 3000 m have lower ABL-TopoIndex values similar to some of the European mountains, whereas YEL is situated on the large

Yellowstone ~~plateau~~-high plain at an average elevation of 2400 m resulting in a high ABL-TopoIndex (7.2) that is similar to the values for NCOS and HLE. Mount Bachelor (MBO) is located near the top of an isolated volcano from the Cascade volcanic arc that dominates the plain surrounding it, explaining its low ABL-TopoIndex. WHI is located in the Pacific Coast Mountains, the mountain range name referring to the vicinity between the high altitude massif and the ocean coast. The highest peaks in the Pacific Coast Mountains have summits between 3000 and 4000 m (WHI is at 2182 m). WHI has middle range ABL-TopoIndex (1.4) despite its low altitude due to the proximity of the ocean and to the rather narrow width of the massif (300 km). Both APP and SHN are situated at the same altitude in the Blue Ridge mountains of the Appalachian range. At the latitude of SHN, the width of the Blue Ridge mountain range is much narrower than at APP's latitude. Moreover SHN is almost on the top of the ridge whereas APP is on a ~~plateau~~high plain. SHN therefore has higher G8 and LocSlope and lower hyps% and DBinv leading to much lower ABL-TopoIndex than found for APP.

- **Andes:** CHC (5320 m), the highest station in this study, is located in the Cordillera Oriental, itself a sub-range of the Bolivian Andes massif, and is part of the mountain belt surrounding the Altiplano (literal translation high plain) with an average height of 3750 m. This position explains its mid-range ABL-TopoIndex of about 1.3 due to relatively high hypso% (1.03%) and low hypsoD50 (1311 m). PEV (4765 m), the South America station with the lowest ABL-TopoIndex, is located at the extreme northeastern extension of the South America's Andes mountain range that peaks at about 5000 m. Its high position in its mountain range is characterized by a very low hypso% (0.28 %) and the highest hypsoD50 of 4019 m. TLL is situated in the foothills of the Andes in Chile near to the Pacific ocean and has a similar ABL-TopoIndex to SZZ due similarities in topography. LQO is at higher altitude than TLL but located in the middle of the Altiplano leading to an ABL-TopoIndex larger than 20.

- **Africa:** Mount Kenya (5199 m), the second highest peak of Africa and the highest in Kenya, is an isolated volcanic massif with several peaks. MKN observatory is located some 1500 m under Mount Kenya's summit resulting in a mid-range ABL-TopoIndex of about 1. Assekrem (ASK, 2710 m) is located on a small (about 2.5 km<sup>2</sup>) high ~~plateau~~ plain in the Hoggar Mountains located in central Sahara. The highest summit in the Hoggar range peaks at 2908 m. Despite being situated on a flat area, ASK has quite low ABL-TopoIndex value because of its relatively high elevation in the Hoggar Mountains.

- **Arctic:** SUM is located high atop the Greenland ice sheet in the central Arctic. The ice sheet has a very smooth topography due to its build up by glaciation and precipitation. While SUM has a high hypso%, its hypsoD50, G8 and LocSlope are very low and its DBinv is large leading to high ABL-TopoIndex. The Zeppelin Observatory (475 m) in Svalbard is located near the top of Zeppelinfjellet (556 m) above Ny-Ålesund but cannot be considered as a high altitude site and was added to the study for comparison purpose. Its ABL-TopoIndex is consequently very high since the highest summit on the Spizbergen island is at 1717 m.

### 3.5. Correlation between aerosol parameters and the ABL-TopoIndex

While Fig. 8 shows that there are some clear patterns in the ABL-TopoIndex, it is also instructive to see how the ABL-TopoIndex relates to measurements at mountain sites. The NCOS and SUM stations have a very high ABL-TopoIndex due to their situation on a high altitude plain near a vast lake and on the smooth shape of the Greenland inland ice sheet, respectively. Since they are not situated in a complex topography, they were excluded from this analysis due to their clear outlier status. ZEP, situated at very low altitude (475 m), also has a very high ABL-TopoIndex values. It was also not included in the correlation analysis since its seasonal and diurnal cycles exhibit different features than the high altitude stations (see Sect. 4.1).- In order to have a robust estimate of the correlation between the aerosol measurements (following a Johnson distribution) and the topographical parameters (following a normal or a log-normal distribution depending on the parameters) the Spearman's rank correlation was calculated. It should be noted that the Kendall's tau correlation analysis leads to the same conclusions (see Table S4). The Spearman's rank correlation measures the strength and direction between two ranked variables without the requirement that the variables be normally distributed. Here it is also used to verify that the assumed relationships between topographical and aerosol parameters correspond to those proposed in section 2.3 (e.g., that a positive correlation with aerosol loading as a surrogate for ABL influence in case of lifting processes without precipitation is found for the ABL-TopoIndex, hypso%, DBinv and station altitude and an anti-correlation for hypsoD50, LocSlope and G8). That is the case for all used topographical parameters. The Spearman's rank correlation coefficients of the 5th, 50th and 95th percentiles of the measured aerosol parameters with the altitude (mean the altitude over the 9 grid cells, similarly to the ABL-TopoIndex calculation), the latitude, the ABL-TopoIndex as well as all the individual parameters constituting the ABL-TopoIndex are presented on Fig. 9.

The ABL-TopoIndex has statistically significant (s.s.) correlation for the all the percentiles of all aerosol parameters except for the 95 percentile of the scattering coefficient. The highest correlation and s.s. are found for the 5 percentile of the absorption and scattering coefficient, whereas the 50 percentile has the highest correlation for the number concentration. The correlation coefficient with the maxima of the aerosol parameters (95th percentile) is always lower than with the minima (5% percentile) and is s.s. at 95% of confidence level only for the absorption coefficient. The minima of the aerosol parameters - particularly of the absorption coefficient- correspond to the measurement of air masses with the lowest aerosol concentration, namely FT air masses with the lowest ABL influence and no advection of polluted air masses. In contrast, the maxima correspond to the advection or convection of air masses with high aerosol loads and can, to some extent, be caused by special events such as dust or biomass burning events. In contrast to the absorption coefficient, the particle number concentration (and, to a far lower extent, the scattering coefficient) depend not only on the ABL influence but also on the new particle formation (NPF) that can be enhanced at high altitudes (Boulon et al., 2011; Rose et al., 2015). Thus, the high correlations of the ABL-TopoIndex with the minima of the aerosol absorption coefficient as well as its lower correlation with the absorption coefficient maxima, with the number concentration minima and with the scattering coefficient suggest the ABL-TopoIndex is indeed a promising indicator for ABL influence based on station topography.

Formatted: Indent: Left: -0.01 cm  
Outline numbered + Level: 2 +  
Numbering Style: 1, 2, 3, ... + Start  
1 + Alignment: Left + Aligned at: 0  
cm + Indent at: 1.4 cm

The hypso%, a large scale parameter, has s.s correlations with all the percentages of all the aerosol optical properties, whereas LocSlope and G8, two small scale parameters, have s.s anticorrelations except for the 95 percentile of the scattering coefficient. The hypsoD50 is s.s. for the 5% and 50% of the absorption and scattering coefficients and for the 50% and 95% of the number concentration. Only the DBinv exhibits no s.s. correlation with any of the aerosol parameters.

5 There are no s.s. correlations between the station altitude and the percentiles of any of the aerosol parameters. The station elevation alone is therefore not a good predictor of the ABL influence (at least as it relates to particle concentration and aerosol optical properties). The latitude has s.s. anticorrelation with 5% and 50% of the scattering coefficient.

The correlations of the topographical parameters with the diurnal and seasonal cycles of the aerosol measurements exhibit a clearly different pattern. Fig. 10 shows the Spearman's rank correlation coefficients of the topographical parameters with the minimum (Dmin) and maximum (Dmax) of the monthly diurnal cycle strength as well as with the seasonal cycle (Season) of the aerosol parameters. The diurnal cycles were calculated for the 12 months, so that Dmin and Dmax correspond to the lowest and the greatest monthly amplitudes, respectively. Both the diurnal and the seasonal cycles were calculated as the strength of the autocorrelation function (see § 2.4 and supplement) so that the underlying parameters are de facto normalized and that the cycles of each stations can be directly compared. The greatest correlation is found between the amplitudes of the diurnal cycles and the latitude for all three aerosol parameters. This anticorrelation is particularly marked for the number concentration diurnal cycles. At low latitudes, the stronger insolation enhances the surface temperature and the thermal convection leading to stronger diurnal cycles, particularly in summer, and the convective flow is less likely to be inhibited during the winter due to longer daylight hours. Together these effects result in a greater ABL influence year round and explain the high correlations with the diurnal cycle amplitude. The high correlation between the maximal diurnal cycle and the number concentration can also be explained by the promotion of NPF by the stronger insolation at low latitude.

15 The ABL-TopoIndex is s.s. correlated with the diurnal cycle minimal and maximal strengths of the absorption coefficient. This correlation is once again principally due to the hypso% and G8, and to a lower extent, the LocSlope. The correlation with the diurnal cycle minimal amplitude occurs because the stations that remain in the FT during the whole day should not exhibit any systematic diurnal cycles. The maximal amplitude of the diurnal cycles occurs when the site is in the FT during the night (without any influence of the RL) and influenced by the ABL during the day. Once again, the greatest correlation is found for the absorption coefficient which directly depends on ABL air masses uplift, where the number concentration and scattering coefficient cycles are also influenced by gas-to-particle conversion processes such as NPF that can be enhanced at low temperature (that is in a opposite seasonal cycle than the CBL height). The only s.s. correlation with station altitude is found for the scattering coefficient seasonal cycle. Similar to the correlation with the percentiles, there is a high anticorrelation between the particle number concentration diurnal cycles and G8 suggesting that the slope steepness in the vicinity of the stations inhibited both the transport of polluted air masses and NPF. Apart from a correlation at 90% confidence level between DBinv and the absorption coefficient, the lack of further s.s. correlations with the seasonal cycles can be attributed to several factors: (i) the relatively small time period (2-5 years) covered by most of the datasets leading to difficulties in the statistical determination of a yearly periodicity due to inter-annual variability, (ii) the low aerosol

concentration at high altitude sites inducing measurements part of the time near the detection limits of the instruments (see for example the problem with the absorption coefficient at § 2.4) and (iii) the necessary whitening procedure (see supplement) increasing the dataset noise .

## 4. Discussion

5 In this section the assessments, improvements and applications of the ABL-TopoIndex are discussed. First the possible species and phenomena enabling the estimation of the ABL influence are summarized and the occurrence of diurnal and seasonal cycles as a function of the station elevation are discussed. Second, the significance of the correlations between the topographical and the aerosol parameters are further interpreted. Finally, possible additional parameters that could increase the significance and the application of the ABL-TopoIndex are mentioned, in addition to the criteria relevant for choosing  
10 future sites to sample FT air masses.

### 4.1. Using measurements to assess the ABL influence

In order to test the relevance of the ABL-TopoIndex, it is first necessary to find a parameter commonly measured at high altitude stations that can be used as an ABL tracer. Pollutants emitted at the Earth's surface and having a (typically) minimal concentration in the FT could act as potential tracers of the ABL influence. Our results showed that of the three aerosol  
15 parameters tested in this study (number concentration, absorption coefficient and scattering coefficient), absorption coefficient has the greatest correlation with the ABL-TopoIndex values. Other possible candidates for testing the ABL-TopoIndex include the aerosol mass concentration, size distribution and chemical composition, the water vapor and the trace gases concentrations (e.g., CO<sub>2</sub>, PAN, NO<sub>x</sub>, NO<sub>y</sub>, O<sub>3</sub>, SO<sub>2</sub>, isotopologue ratio of water vapor) and the radon<sup>222</sup> concentration. These parameters have been used in different studies to provide information about the seasonal and diurnal cycles (e.g.,  
20 Collaud Coen et al., 2011; Griffiths et al., 2014; Marinoni et al., 2010; McClure et al., 2016; Okamoto and Tanimoto, 2016; Pandolfi et al., 2014; Ripoll et al., 2015; Zellweger et al., 2009), the sources and transport of aerosol to the site ( e.g., Cuevas et al., 2013; García et al., 2017; Pandey Deolal et al., 2014; Ripoll et al., 2014), the local orographic flows and the effect of the synoptic- and meso-scale weather types ( e.g., Bonasoni et al., 2010; Gallagher et al., 2011; González et al., 2016; Henne et al., 2005; Kleissl et al., 2007; Tsamalis et al., 2014; Zellweger et al., 2002). All of the extensive aerosol parameters, the  
25 radon<sup>222</sup>, water vapor concentration, particulate nitrate ( $NO_3^-$ ) and organics have been shown to be correlated with ABL transport whereas CO/NO<sub>y</sub> and NO<sub>x</sub>/NO<sub>y</sub> ratios are anticorrelated (Legreid et al., 2008; Zellweger et al., 2003). Zellweger et al. (2003) concluded that, in contrast to NO<sub>y</sub>, the major process for upward transport of aerosol is the thermally induced vertical transport, confirming that the aerosol parameters used in this study should be in most cases good tracers for ABL influence. However, one has to remain conscious that many lifting processes co-occur with precipitation involving hence aerosol washout. The present validation of the ABL-TopoIndex by aerosol measurements at high altitude stations is  
30

Formatted: Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 0.63 cm

Formatted: Indent: Left: -0.01 cm Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0 cm + Indent at: 1.4 cm

consequently valid only for thermally induced processes without precipitation. ABL events involving air masses with low aerosol concentrations due to washout –for exemple synoptic lifting or foehn- were not directly analyzed.

Because there are variable pollution levels in the vicinity of the stations, a single absolute value of a pollutant cannot be used to evaluate the ABL-TopoIndex (or ABL influence in general) when considering multiple high altitude stations. An inventory of the proximate pollution sources bounded to a 3D thermodynamic model adapted to complex topographies would be required before using absolute pollutant concentrations as indicators of ABL influence at high altitude sites.

Another possibility would be to weight the pollution source inventories by factors depending on their vertical and horizontal distances to the high altitude stations, as well as on a seasonal parameter representing the potential ABL height. A further use of DBinv to restrict the area of potential pollution sources could also be envisaged since this parameter describes the domain from which pollutants can reach the high altitude station by convection without crossing topographical barriers. This ~~problem~~ delicate issue can however be avoided by instead considering dynamical parameters such as the various temporal cycles.

At most of the high altitude stations, a seasonal cycle in ABL-indicator species is de facto observed. The maximum values of the seasonal cycles are correlated with ABL transport and typically occur in summer or in the pre-monsoon season while the minimum of the seasonal cycle occurs in winter or monsoon seasons. Usually the spring leads to higher aerosol loading than the autumn probably related to higher ABL height in the spring. These seasonal cycles are explained by the stronger thermal heating of the soil which induces convection and buoyancy in summer and by the atmospheric cleaning effect of precipitation during the monsoon. It would be expected that stations continuously situated in the ABL throughout the year could exhibit different seasonal cycles than high altitude sites due to the seasonal modification of the sources and/or of the synoptic and meso-scale meteorological conditions (see for example the difference between HPB and JFJ on Fig. S2). In contrast, a station located such that it stayed continuously in the FT would have a seasonal cycle that depends only on long-range, high altitude transport climatology (e.g., long-range transport of Asian dust and pollution at MLO in spring (Collaud Coen et al., 2013), North-America ABL transport to IZO through westerlies in spring (García et al., 2017), and dust events in EU spring and autumn (Collaud Coen et al., 2004)). Since seasonally changing parameters (e.g., temperatures, cloud cover, solar radiation, wind speeds, surface albedo, precipitation) were not studied and since the length of most of the time series are too short to smooth these effects, the ABL-TopoIndex will probably not represent an overall picture of ABL influence except at seasonally invariant sites (e.g., very low latitude sites).

The typical diurnal cycle of ABL pollutants at high altitude stations that are partially influenced by the ABL consists of a minimum in the early morning (4h-6h LTC) followed by an increase of the compound with a maximum in the late afternoon (15-17h LTC) and a decrease during the night. If the ABL influence is mostly due to orographic winds, upslope/valley winds begin to flow some hours after sunrise and downslope/mountain winds initiate after the occurrence of negative vertical heat flux. Stations always situated in the FT should exhibit no systematic diurnal cycles whereas the stations always situated in the ABL often show various diurnal cycles that can be explained by the behavior of local sources, the diurnal cycle of the ABL height and/or local meteorological conditions. At high elevation and high latitude stations the diurnal cycle typically

vanishes during winter but is clearly present during summer, spring and, to a lesser extent, autumn. For stations at lower altitude that stay in the ABL (or CBL, SBL or RL) during the whole day in summer (e.g., MSA (Pandolfi et al., 2013), HPB and PUY (Hervo et al., 2014)), the diurnal cycle may also vanish during that period.

Testing the ABL-TopoIndex using pollutant diurnal cycles is further complicated by the presence of the residual layer (RL) that keeps the pollutants brought to high altitudes during the previous days at those elevated levels during the nighttime. The climatology of the RL height usually exhibits a similar seasonality as the ABL height, with a maxima in summer (or pre-monsoon) season and a minima in winter (Birmili et al., 2009, 2010; Collaud Coen et al., 2014; Wang et al., 2016). Further, the RL also has similar dependency as the ABL as a function of latitude. The RL's maximum height also depends, therefore, on the duration of the incoming radiation. The RL pollutant concentrations are much higher than nighttime FT concentrations, leading to less marked diurnal cycles in summer than in spring (Blay-Carreras et al., 2014; Collaud Coen et al., 2011; Hallar et al., 2016; Hervo et al., 2014). The impact of the RL on the aerosol concentration is probably one of the most important reasons for the low correlation between the topographical parameters and the aerosol temporal cycles.

However, the statistical determination of the diurnal and seasonal cycle amplitudes suffer from several difficulties: 1) the low aerosol concentration at high altitude often results in measurements near the detection limit leading to high uncertainties, 2) the high hourly autocorrelation of the data require a pre-whitening procedure (see supplement) in order to be able to detect the diurnal and seasonal cycle, 3) meteorological conditions (e.g., cloud coverage, precipitation or seasonal fluctuations) modify the theoretical cycles and lead to a broadening of the time of the extrema. These difficulties make obtaining clear statistical cycles another reason contributing to the observed low correlations.

Recently, the influences of the local and of the more regional or meso-scale ABL at the JFJ were separated by differentiating the Local Convective Boundary Layer (LCBL) height from the high altitude aerosol layer (Poltera et al., 2017). The LCBL was found to rarely influence the JFJ research station (never in winter, 4% of the time which corresponds to 22% of the days in summer), whereas the continuous aerosol layer has a large influence on the JFJ pollutant concentrations (21% of the time in winter and 41% of the time corresponding to 77% of the days in summer). This suggests that the mechanisms explaining the heights of the LCBL and the more horizontally extended aerosol layer have different causes and do not follow the same diurnal pattern. This phenomenon will be more pronounced at continental high altitude stations than at marine isolated island stations since the marine ABL is less prone to strong diurnal cycles.

#### **4.2 Correlation between the topography and the aerosol parameters**

The correlations between topographical and aerosol parameters presented under Sect. 3.5 can now be further discussed in light of the pollutant temporal cycles. The absorption coefficient is primarily due to the presence of black carbon emitted from combustion processes occurring mostly in the ABL and rarely near the high altitude stations; additionally, BC aerosol is not produced by any secondary processes. Among the aerosol parameters studied here, the absorption coefficient is therefore the best tracer for anthropogenic pollution and biomass burning and consequently for ABL influence. It is then expected that a better correlation will be obtained between the topography parameters increasing the ABL influence and the

absorption coefficient. The ABL-TopoIndex reflects this correspondence, particularly through the contribution of the hypso% parameter (recall that hypso% represents the relative altitude of a station in its mountain range), the LocSlope and G8. The best correlation for both ABL-TopoIndex, hypso%, Locslope and hypsoD50 are found for the 5<sup>th</sup> percentile of the absorption coefficient, since the minima of the aerosol loading is a better tracer of the lowest ABL influence, whereas the maxima is much more dependent on source intensity and special events. Similar to this result, a clear correlation was also found between the continuous aerosol layer maximum height and the absorption coefficient measured in-situ at the JFJ (Fig. 8 in (Poltera et al., 2017)). The absorption coefficient amplitudes of the diurnal cycle are also the only aerosol cycles having a s.s. correlation with the ABL-TopoIndex.

It is more difficult to directly tie scattering and number concentration to the ABL incursions. This is because the formation of new particles and their subsequent growth are well-known to be very efficient processes at high altitudes due to the high insolation and the low temperature. Moreover, the NPF is also enhanced by local thermal winds and forced convection due to favorable changes in thermodynamic conditions (Boulon et al., 2011; Rose et al., 2015). It was found at the JFJ and confirmed at other stations that new particle formation, and particularly strong nucleation events, occur mostly when the air masses were in contact with the ABL within 2 days before arriving at high altitudes (Bianchi et al., 2016). NPF and subsequent growth of the particles have a large impact on the number concentration and its temporal cycles and a smaller influence on the scattering coefficient. The parameters describing the local topography (G8 and LocSlope) have the greatest correlation with the number concentration and are probably more relevant to the local CBL transport than to the longer range continuous aerosol layer as defined in Poltera et al. (2017). The number concentration and, to a lesser extent, the absorption coefficient percentiles and diurnal cycles are anti-correlated with the local (G8: 0.5-1 km) and regional (LocSlope and hypsoD50: 10 km) slopes, suggesting there is an increase of particle number concentration when there are small altitude differences and gentle slopes around the station. This dependence on the ease of local transport can be explained by transport to the station not only of aerosol, but also of gaseous precursors for NPF and of newly formed particles at lower elevations. Globally, NPF is the reason why the greatest correlations are found with the 50 percentile of the number concentration, instead of with the 5 percentile found for the absorption and scattering coefficients. The greater correlation of local slope (G8 and LocSlope) with the number concentration rather than with the absorption coefficient can be explained both by the very scarce sources of black carbon in the near vicinity of most of the high altitude stations and by the smooth pressure decrease experienced by the precursors during their upslope transport along gentle slopes leading to more condensation processes and nucleation.

The aerosol diurnal cycles are influenced by numerous phenomena (see Sect. 4.1) leading to a non-trivial relationship with the ABL influence. The study of the diurnal cycles can bring valuable results if specific cases are analyzed and compared, the statistical approach used here leads to is less obvious results due to the noise in the data (low aerosol concentration and whitening process), to the inter-annual variability of the meteorological processes and to cloud, precipitation and long-range advection involving a large day to day variability. There are consequently few statistical correlations between topography parameters and the diurnal cycles. The clearest correlation is the influence of the insolation on the aerosol diurnal cycles

amplitudes. This dependence between the latitude and the aerosol concentration was already mentioned by Kleissl et al. (Kleissl et al., 2007) and is easily understandable, the convection and the new particle formation being directly dependent on the solar radiation intensity. The other correlations found between some topography parameters (ABL-TopoIndex, hypso%, G8 and LocSlope) and the absorption coefficient are however directly bounded to the ABL influence.

### 5 4.3 Improving and applying the ABL-TopoIndex

The choice of the 5 parameters included in the ABL-TopoIndex was initially based on several assumptions relating the topography to the ABL influence (see Sect. 2.3). Several other parameters such as the topographical wetness index, the upstream catchment area, the Efremov-Krcho classification, the hypsometric curve, integral and index as well as the topographic prominence were tested but were finally eliminated as being not relevant for various reasons (Table S2). Indeed, most of the parameters comprising the ABL-TopoIndex exhibit some correlations with aerosol parameters. The hypso%, LocSlope and G8 are the parameters explaining the greatest variance in the aerosol optical properties, with the hypsoD50 having a lower influence than the other three parameters. It also seems evident that the topographical parameters linked to the steepness and the altitude differences (G8 and LocSlope) are clear indicators for NPF. The DBinv seems to be the least explanatory parameter in terms of ABL influence and this large scale parameter should probably be bounded with a source inventory to increase its relevance for identifying boundary layer influence (see § 4.19). DBinv has however a clear influence on the statistical significance of the correlations between the ABL-TopoIndex and the aerosol cycles (not shown in the paper) and has the greatest correlation with aerosol seasonal cycles. However, the aerosol parameters and, particularly, the absorption coefficient cannot be considered as unique tracers of the ABL, particularly in case of lifting processes with precipitation. Analysis of other ABL marker (gaseous species, radon, wind turbulences, etc.) can provide information on additional transport mechanisms which would allow for refinement of this topographic analysis by adding further parameters.

East-oriented slopes are heated early during the day and have therefore a greater contribution to the thermal convection and the associated valley winds. A parameter weighting the east slope area could therefore be added to the ABL-TopoIndex. The various geomorphologies of the mountainous ranges included in this study also raise the question of whether the stations should all be combined together for analysis as was done here, or if a morphological parameter should instead be found for each massif. The mountain steepness (at a larger scale than LocSlope and G8) also determines the necessary velocity for the wind to cross the mountains and could be an additional parameter. Finally, future studies should attempt to build a direction dependent ABL-TopoIndex that also takes into account the topography of each valley up to the meso-scale range.

It is important to understand the FT versus ABL influence on historical data sets from established high altitude observatories. The ABL-TopoIndex is one tool that can help elucidate the different influences. A further improvement could include an angular dependency of the ABL-TopoIndex allowing quantifying the potential direction of the maximal ABL influence and the most influencing pollutant sources. The ABL-TopoIndex may also be useful *a priori* in locating measurements for a field campaign or identifying potential sites for long-term observatories if FT measurements are the goal, particularly when no

previous measurements exist. For example, in-situ aerosol measurements are done at IZO at an altitude of 2373 m whereas aerosol optical depth and water vapor isotopologues measurements are done at TDE at 3538 m on the same volcano. TDE has a much lower ABL-TopoIndex than IZO and consequently TDE's measurements are more likely to represent the FT. The topography around the PYR station suffers from several inconveniences (see Fig. 7 and Sect. 3.1) leading to a high ABL influence. Even if the choice of the actual site is driven by compelling and practical logistical arguments, other emplacements at similar altitudes would have ensured lower pollution impact. Finally, some stations such as BEO (2925 m), HAC (2314 m) and MWO (1916 m) are not situated at very high altitudes but present excellent locations for FT sampling. Obviously there are other issues to consider when deploying instruments as well (e.g., ease of access, power availability, presence of local pollution sources, etc.), but the ABL-TopoIndex is one factor that could be considered to maximize the potential for FT sampling.

## Conclusion

The ABL-TopoIndex is a topographical index based on the hypsometric curve, the slope of the terrain around the station and the drainage basin for ~~convection~~thermally lifted pollutants. It allows one to rank the high altitude stations as a function of their ABL influence or to optimize choice of site location for FT sampling. High altitude stations situated on volcanic islands, the highest stations in the Alps, in the Andes and in the Pyrenees have low ABL-TopoIndex values. Stations situated at or near the summit of their mountainous ranges such as BEO, HAC and MWO also have low ABL-TopoIndex values. Stations situated at altitudes between 4000 and 5500 m in the Himalaya and the Tibetan Plateau have high ABL-TopoIndex values due to their relatively low position compared to the ~~Himalayan~~summits. Statistically significant correlations between the ABL-TopoIndex and the aerosol parameters measured at high altitude sites allow validation of the methodological approach. The greatest correlations are found with the minima of the aerosol parameters that represent the minimal ABL influence or, in other words, the most likely FT air masses. The maxima of aerosol parameters are more representative of the intensity of aerosol sources and of advection of air masses with high aerosol concentrations. There are also strong anticorrelations between the local steepness of the slope and the particle number concentration, suggesting that new particle formation could be largely influenced by this topographical parameter. If high altitude stations undergo daytime ABL air influence due to convection, a pronounced diurnal cycle of aerosol parameters is usually measured. The amplitude of the diurnal cycle of the absorption coefficient is s.s. correlated with the ABL-TopoIndex and is, thus, likely to be representative of ABL influence. The strength of the diurnal cycles of the scattering coefficient and the number concentration is however mostly explained by the latitude of the station, leading to the conclusion that the sun radiation intensity and duration drive the aerosol diurnal cycle.

## Acknowledgments

The authors greatly acknowledge:

- W. Schwanghart, the programmer of the TopoToolBox for putting his codes as a freeware and for all the kind and always rapid support he offers.
- CHC: the Chacaltaya consortium and all Laboratory for Atmospheric Physics at UMSA for taking care of the station and collecting the data. Swedish participation was supported by The Swedish Foundation for International Cooperation in Research and Higher Education (STINT) and The Swedish research council FORMAS.
- CMN: the European Commission funded ACTRIS, ACTRIS-2 EU project and NEXTDATA National project funded by MIUR.
- IZO: Global Atmospheric Watch program funded by AEMET and by the project AEROATLAN (CGL2015-66299-P, funded by the Ministry of Economy and Competitiveness of Spain and the European Regional Development Fund - ERDF).
- JFJ: International Foundation High Altitude Research Stations Jungfrauoch and Gornergrat (HFSJG) and the Federal Office of Meteorology and Climatolofy, MeteoSwiss, within the Swiss program of the Global Atmosphere Watch (GAW) of the World Meteorological Organization, funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 654109 (ACTRIS2), and the Swiss State Secretariat for Education, Research and Innovation (SERI) under contract number 15.0159-1. The opinions expressed and arguments employed herein do not necessarily reflect the official views of the Swiss Government.
- LLN: by Taiwan Environmental Protection Administration.
- MSA and MSY: MINECO (Spanish Ministry of Economy and Competitiveness), the MAGRAMA (Spanish Ministry of Agriculture, Food and Environment), the Generalitat de Catalunya (AGAUR 2014 SGR33 and the DGQA) and FEDER funds under the PRISMA project (CGL2012- 39623-C02/00). This work has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 654109. Marco Pandolfi is funded by a Ramón y Cajal Fellowship (RYC-2013-14036) awarded by the Spanish Ministry of Economy and Competitiveness
- MLO, NWR and SUM: the NOAA observatory staff at each station and the NOAA Climate Program Office's Atmospheric Chemistry, Carbon Cycle and Climate (AC4) program
- MUK: Ministry of Foreign Affairs of Finland, Academy of Finland (Project number 264242) for the financial support and an esteemed collaboration of FMI and TERI.
- NCOS: Max Planck Institute for Chemistry and Chinese Academy of Sciences (Project SKLCS-ZZ-2017 and KJZD-EW-G03-03).

**Formatted:** Bulleted + Level: 1 +  
Aligned at: 0.63 cm + Indent at: 1  
cm

- PDI: Federal Office of Meteorology and Climatology MeteoSwiss through the project Capacity Building and Twinning for Climate Observing Systems (CATCOS) Phase 2, Contract no. 81025332 between the Swiss Agency for Development and Cooperation and MeteoSwiss
- PEV: Support from Swedish Research Council (Vetenskapsrådet) and Swedish International Development Cooperation Agency (SIDA)
- PDM: Pyrenean Platform for Observation of the Atmosphere P2OA (<http://p2oa.aero.obs-mip.fr>), University Paul Sabatier, Toulouse, France, and CNRS (Centre National de la Recherche Scientifique).
- OMP: United States National Oceanic and Atmospheric Administration (NOAA) grants NA16GP1658, NA86GP0325 and NA030AP430002; United States National Science Foundation (NSF) grants ATM-0215843 and INT-0110397; Portuguese Foundation for Science and Technology (FCT) grants POCTI-32649-CTA-2000; SFRH/BD/9049/2002; and Portuguese Regional Govern of Azores and especially Professor Richard Honrath (1961-2009).
- PYR: the framework of the UNEP — ABC (Atmospheric Brown Clouds) and EvK2CNR — SHARE (Stations at High Altitude for Research on the Environment) projects and the appreciated collaboration with the technical Nepalese staff.
- SBO: TU Wien, Commission of Climate and Air Quality - ÖAW, Local Government of Salzburg - Immissionsschutz, Umweltbundesamt GmbH and Aerosol d.o.o .
- SPL: Randolph Borys, Ian McCubbin, Douglas Lowenthal, Peter Atkins, and Joe Messina, the US National Science Foundation (grant AGS-0079486), the Steamboat Ski Resort and the Desert Research Institute, a permittee of the Medicine-Bow Routt National Forests
- TLL: Federal Office of Meteorology and Climatology MeteoSwiss through the project Capacity Building and Twinning for Climate Observing Systems (CATCOS) Phase 2, Contract no. 81025332 between the Swiss Agency for Development and Cooperation and MeteoSwiss, G. Wehrle from Paul Scherrer institute, the site operator Luis Valle from DGAC and to the maintenance staff from MeteoChile.
- WLG: National Scientific Foundation of China (41675129), National Key Project of MOST (2014CB441201) and the CMA Innovation Team for Haze-fog Observation and Forecast.
- ZEP: Support from Swedish Environmental Protection Agency (Naturvårdsverket), The Swedish research council FORMAS, Knut and Alice Wallenberg Foundation (KWA) and Norwegian Polar Institute (NPI)
- ZUG: H.E. Scheel for performing aerosol measurements at ZUG and W. Junkermann, Karlsruhe Institute of Technology, Garmisch-Partenkirchen, Germany, who give us access to this dataset.

Field Code Changed

Field Code Changed

## References

- Ambrose, J. L., Reidmiller, D. R. and Jaffe, D. A.: Causes of high O<sub>3</sub> in the lower free troposphere over the Pacific Northwest as observed at the Mt. Bachelor Observatory, *Atmos. Environ.*, 45(30), 5302–5315, doi:10.1016/j.atmosenv.2011.06.056, 2011.
- 5 Andrade, M., Zaratti, F., Forno, R., Gutiérrez, R., Moreno, I., Velarde, F., Ávila, F., Roca, M., Sánchez, M. F., Laj, P., Jaffrezo, J. L., Ginot, P., Sellegri, K., Ramonet, M., Laurent, O., Weinhold, K., Wiedensohler, A., Krejci, R., Bonasoni, P., Cristofanelli, P., Whiteman, D., Vimeux, F., Dommergue, A. and Magand, O.: Set to work of a new climate monitoring station in the central andes of Bolivia: the Gaw/Chacaltaya station, *Rev. Boliv. Física*, 26(26), 6–15, 2015.
- Andrews, E., Ogren, J. A., Bonasoni, P., Marinoni, A., Cuevas, E., Rodríguez, S., Sun, J. Y., Jaffe, D. A., Fischer, E. V.,  
10 Baltensperger, U., Weingartner, E., Collaud Coen, M., Sharma, S., Macdonald, A. M., Leaitch, W. R., Lin, N.-H., Laj, P., Arsov, T., Kalapov, I., Jefferson, A. and Sheridan, P.: Climatology of aerosol radiative properties in the free troposphere, *Atmos. Res.*, 102(4), 365–393, doi:10.1016/j.atmosres.2011.08.017, 2011.
- Angelov, C., Nikolova, N., Arsov, T., Kalapov, I., Tchordadjieff, A., Penev, I. and Angelov, I.: BEO Moussala: complex for environmental studies, in *Sustainable Development in Mountain Regions*, edited by G. Zhelezov, pp. 349–365, Springer  
15 International Publishing, Switzerland., 2016.
- Backman, J., Schmeisser, L., Virkkula, A., Ogren, J. A., Asmi, E., Starkweather, S., Sharma, S., Eleftheriadis, K., Uttal, T., Jefferson, A., Bergin, M. and Makshtas, A.: On Aethalometer measurement uncertainties and multiple scattering enhancement in the Arctic, *Atmos. Meas. Tech. Discuss.*, (December), 1–31, doi:10.5194/amt-2016-294, 2016.
- Baltensperger, U., Gäggeler, H. W., Jost, D. T., Lugauer, M., Schwikowski, M. and Weingartner, E.: Aerosol climatology at  
20 the high-alpine site Jungfraujoch, Switzerland, *J. Geophys. Res.*, 102(D16), 19707–19715, 1997.
- Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H. E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J. and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing,  
25 *Science* (80-. ), 352(6289), 1109–1112, doi:10.1126/science.aad5456, 2016.
- Birmili, W., Ries, L., Sohmer, R., Anastou, A., Sonntag, A., König, K. and Levin, I.: Feine und ultrafeine Aerosolpartikeln an der GAW-Station Schneefernerhaus/Zugspitze, *Gefahrstoffe-Reinhaltung der Luft*, 69, 31–35, 2009.
- Birmili, W., Göbel, T., Sonntag, A., Ries, L., Sohmer, R., Gilge, S., Levin, I. and Stohl, A.: A case of transatlantic aerosol transport detected at the Schneefernerhaus Observatory (2650 m) on the northern edge of the Alps, *Meteorol. Zeitschrift*,  
30 19(6), 591–600, doi:10.1127/0941-2948/2010/0465, 2010.
- Blay-Carreras, E., Pino, D., Vilà-Guerau De Arellano, J., Van De Boer, A., De Coster, O., Darbieu, C., Hartogensis, O., Lohou, F., Lathon, M. and Pietersen, H.: Role of the residual layer and large-scale subsidence on the development and evolution of the convective boundary layer, *Atmos. Chem. Phys.*, 14(9), 4515–4530, doi:10.5194/acp-14-4515-2014, 2014.

- Bodhaine, B. A.: Aerosol absorption measurements at Barrow, Mauna Loa and the south pole, *J. Geophys. Res. Atmos.*, 100(D5), 8967–8975, doi:10.1029/95JD00513, 1995.
- Bodhaine, B. A., Harris, J. M., Herbert, G. A. and Komhyr, W. D.: Identification of volcanic episodes in aerosol data at Mauna Loa Observatory, *J. Geophys. Res. Ocean.*, 85(C3), 1600–1604, doi:10.1029/JC085iC03p01600, 1980.
- 5 Bonasoni, P., Laj, P., Angelini, F., Arduini, J., Bonafè, U., Calzolari, F., Cristofanelli, P., Decesari, S., Facchini, M. C., Fuzzi, S., Gobbi, G. P., Maione, M., Marinoni, A., Petzold, A., Roccatò, F., Roger, J. C., Sellegri, K., Sprenger, M., Venzac, H., Verza, G. P., Villani, P. and Vuilleumoz, E.: The ABC-Pyramid Atmospheric Research Observatory in Himalaya for aerosol, ozone and halocarbon measurements, *Sci. Total Environ.*, 391(2–3), 252–261, doi:10.1016/j.scitotenv.2007.10.024, 2008.
- 10 Bonasoni, P., Laj, P., Marinoni, A., Sprenger, M., Angelini, F., Arduini, J. and Bonafè, U.: and Physics Atmospheric Brown Clouds in the Himalayas : first two years of continuous observations at the Nepal Climate Observatory-Pyramid ( 5079 m ) , , 10, 7515–7531, doi:10.5194/acp-10-7515-2010, 2010.
- Bonasoni, P., Cristofanelli, P., Marinoni, A. and Vuilleumoz, E.: Atmospheric Pollution in the Hindu Kush – Himalaya Region, *Mt. Res. Dev.*, 32(4), 468–479, 2012.
- 15 Boulon, J., Sellegri, K., Hervo, M., Picard, D., Pichon, J. M., Fréville, P. and Laj, P.: Investigation of nucleation events vertical extent: A long term study at two different altitude sites, *Atmos. Chem. Phys.*, 11(12), 5625–5639, doi:10.5194/acp-11-5625-2011, 2011.
- Bukowiecki, N., Weingartner, E., Gysel, M., Collaud Coen, M., Zieger, P., Herrmann, E., Steinbacher, M., Gäggeler, H. W. and Baltensperger, U.: A review of more than 20 years of aerosol observation at the high altitude research station
- 20 Jungfraujoch, Switzerland (3580 m asl), *Aerosol Air Qual. Res.*, doi:10.4209/aaqr.2015.05.0305, 2016.
- Collaud Coen, M., Weingartner, E., Schaub, D., Hueglin, C., Corrigan, C., Henning, S., Schwikowski, M. and Baltensperger, U.: Saharan dust events at the Jungfraujoch: detection by wavelength dependence of the single scattering albedo and first climatology analysis, *Atmos. Chem. Phys.*, 4, 1–16 [online] Available from: [www.atmos-chem-phys.org/acp/4/1/](http://www.atmos-chem-phys.org/acp/4/1/), 2004.
- Collaud Coen, M., Weingartner, E., Furger, M., Nyeki, S., Prévôt, A. S. H., Steinbacher, M. and Baltensperger, U.: Aerosol
- 25 climatology and planetary boundary influence at the Jungfraujoch analyzed by synoptic weather types, *Atmos. Chem. Phys.*, 11, 5931–5944, doi:10.5194/acp-11-5931-2011, 2011.
- Collaud Coen, M., Andrews, E., Asmi, A., Baltensperger, U., Bukowiecki, N., Day, D., Fiebig, M., Fjaeraa, A. M., Flentje, H., Hyvärinen, A., Jefferson, A., Jennings, S. G., Kouvarakis, G., Lihavainen, H., Lund Myhre, C., Malm, W. C., Mihapopoulos, N., Molnar, J. V., O'Dowd, C., Ogren, J. A., Schichtel, B. A., Sheridan, P., Virkkula, A., Weingartner, E.,
- 30 Weller, R. and Laj, P.: Aerosol decadal trends-Part 1: In-situ optical measurements at GAW and IMPROVE stations, *Atmos. Chem. Phys.*, 13, 869–894, doi:10.5194/acp-13-869-2013, 2013.
- Collaud Coen, M., Praz, C., Haeferle, A., Ruffieux, D., Kaufmann, P. and Calpini, B.: Determination and climatology of the planetary boundary layer height above the Swiss plateau by in situ and remote sensing measurements as well as by the COSMO-2 model, *Atmos. Chem. Phys.*, doi:10.5194/acp-14-13205-2014, 2014.

- Cristofanelli, P., Landi, T. C., Calzolari, F., Duchi, R., Marinoni, A., Rinaldi, M. and Bonasoni, P.: Summer atmospheric composition over the Mediterranean basin: Investigation on transport processes and pollutant export to the free troposphere by observations at the WMO/GAW Mt. Cimone global station (Italy, 2165 m a.s.l.), *Atmos. Environ.*, 141, 139–152, doi:10.1016/j.atmosenv.2016.06.048, 2016.
- 5 Cuevas, E., González, Y., Rodríguez, S., Guerra, J. C., Gómez-Peláez, A. J., Alonso-Pérez, S., Bustos, J. and Milford, C.: Assessment of atmospheric processes driving ozone variations in the subtropical North Atlantic free troposphere, *Atmos. Chem. Phys.*, 13(4), 1973–1998, doi:10.5194/acp-13-1973-2013, 2013.
- Ealo, M., Alastuey, A., Ripoll, A., Pérez, N., Minguillón, M. C., Querol, X. and Pandolfi, M.: Detection of Saharan dust and biomass burning events using near-real-time intensive aerosol optical properties in the north-western Mediterranean, *Atmos. Chem. Phys.*, 16(19), 12567–12586, doi:10.5194/acp-16-12567-2016, 2016.
- 10 Ebert, U. and Welsch, H.: Meaningful environmental indices: A social choice approach, *J. Environ. Econ. Manage.*, 47, 270–283, doi:10.1016/j.jeem.2003.09.001, 2004.
- Fialho, P., Hansen, A. D. A. and Honrath, R. E.: Absorption coefficients by aerosols in remote areas: a new approach to decouple dust and black carbon absorption coefficients using seven-wavelength Aethalometer data, *J. Aerosol Sci.*, 36(267–282), 0–28, 2004.
- 15 Flentje, H., Briel, B., Beck, C., Collaud Coen, M., Fricke, M., Cyrys, J., Gu, J., Pitz, M. and Thomas, W.: Identification and monitoring of Saharan dust: An inventory representative for south Germany since 1997, *Atmos. Environ.*, 109, 87–96, doi:10.1016/j.atmosenv.2015.02.023, 2015.
- Gallagher, J. P., Mckendry, I. G., Macdonald, A. M. and Leitch, W. R.: Seasonal and diurnal variations in aerosol concentration on whistler mountain: Boundary layer influence and synoptic-scale controls, *J. Appl. Meteorol. Climatol.*, 20 50(11), 2210–2222, doi:10.1175/JAMC-D-11-028.1, 2011.
- Gantner, L., Hornsteiner, M., Egger, J. and Hartjenstein, G.: *Gantner\_Diurnal\_Circulation\_Of\_Zugspitzplatt.pdf*, *Meteorol. Zeitschrift*, 12(2), 95–102, doi:10.1127/0941-2948/2003/0012-0095, 2003.
- García, M. I., Rodríguez, S., González, Y. and García, R. D.: Climatology of new particle formation at Izaña mountain GAW 25 observatory in the subtropical North Atlantic, *Atmos. Chem. Phys.*, 14(8), 3865–3881, doi:10.5194/acp-14-3865-2014, 2014.
- Gheusi, F., Ravetta, F., Delbarre, H., Tsamalis, C., Chevalier-Rosso, A., Leroy, C., Augustin, P., Delmas, R., Ancellet, G., Athier, G., Bouchou, P., Campistron, B., Cousin, J. M., Fourmentin, M. and Meyerfeld, Y.: Pic 2005, a field campaign to investigate low-tropospheric ozone variability in the Pyrenees, *Atmos. Res.*, 101(3), 640–665, doi:10.1016/j.atmosres.2011.04.014, 2011.
- 30 González, Y., Schneider, M., Dyroff, C., Rodríguez, S., Christner, E., García, O. E., Cuevas, E., Bustos, J. J., Ramos, R., Guirado-Fuentes, C., Barthlott, S., Wiegele, A. and Sepúlveda, E.: Detecting moisture transport pathways to the subtropical North Atlantic free troposphere using paired H<sub>2</sub>O- $\delta$ D in situ measurements, *Atmos. Chem. Phys.*, 16(7), 4251–4269, doi:10.5194/acp-16-4251-2016, 2016.
- Griffiths, A. D., Conen, F., Weingartner, E., Zimmermann, L., Chambers, S. D., Williams, A. G. and Steinbacher, M.:

- Surface-to-mountaintop transport characterised by radon observations at the Jungfraujoch, *Atmos. Chem. Phys.*, 14(23), 12763–12779, doi:10.5194/acp-14-12763-2014, 2014.
- Haeffelin, M., Angelini, F., Morille, Y., Martucci, G., Frey, S., Gobbi, G. P., Lolli, S., O'Dowd, C. D., Sauvage, L., Xueref-Rémy, I., Wastine, B. and Feist, D. G.: Evaluation of mixing-height retrievals from automatic profiling Lidars and ceilometers in view of future integrated networks in Europe, *Boundary-Layer Meteorol.*, 143(1), 49–75, doi:10.1007/s10546-011-9643-z, 2012.
- Hallar, A. G., Petersen, R., Andrews, E., Michalsky, J., McCubbin, I. B. and Ogren, J. A.: Contributions of dust and biomass burning to aerosols at a Colorado mountain-top site, *Atmos. Chem. Phys.*, 15(23), 13665–13679, doi:10.5194/acp-15-13665-2015, 2015.
- 10 Hallar, A. G., Petersen, R., McCubbin, I. B., Lowenthal, D., Lee, S., Andrews, E. and Yu, F.: Climatology of new particle formation and corresponding precursors at storm peak laboratory, *Aerosol Air Qual. Res.*, 16(3), 816–826, doi:10.4209/aaqr.2015.05.0341, 2016.
- Hamburger, T., Matisans, M., Tunved, P., Ström, J., Calderon, S., Hoffmann, P., Hochschild, G., Gross, J., Schmeissner, T., Wiedensohler, A. and Krejci, R.: Long-term in situ observations of biomass burning aerosol at a high altitude station in  
15 Venezuela -sources, impacts and interannual variability, *Atmos. Chem. Phys.*, 13(19), 9837–9853, doi:10.5194/acp-13-9837-2013, 2013.
- Henne, S., Dommen, J., Neining, B., Reimann, S., Staehelin, J. and Prévôt, A. S. H.: Influence of mountain venting in the Alps on the ozone chemistry of the lower free troposphere and the European pollution export, *J. Geophys. Res. Atmos.*, 110(22), 1–18, doi:10.1029/2005JD005936, 2005.
- 20 Herrmann, E., Weingartner, E., Henne, S., Vuilleumier, L., Bukowiecki, N., Steinbacher, M., Conen, F., Collaud Coen, M., Hammer, E., Jurányi, Z., Baltensperger, U. and Gysel, M.: Analysis of long-term aerosol size distribution data from Jungfraujoch with emphasis on free tropospheric conditions, cloud influence, and air mass transport, *Journal of Geophysical Research: Atmospheres*, 120(18), 9459–9480, <https://doi.org/10.1002/2015JD023660>, 2015.
- Hervo, M., Sellegri, K., Pichon, J. M., Roger, J. C. and Laj, P.: Long term measurements of optical properties and their  
25 hygroscopic enhancement, *Atmos. Chem. Phys. Discuss.*, 14(20), 27731–27767 [online] Available from: <http://www.atmos-chem-phys-discuss.net/14/27731/2014/>, 2014.
- Hsiao, T.-C., Chen, W.-N., Ye, W.-C., Lin, N.-H., Tsay, S.-C., Lin, T.-H., Lee, C.-T., Chuang, M.-T., Pantina, P. and Wang, S.-H.: Aerosol optical properties at the Lulin Atmospheric Background Station in Taiwan and the influences of long-range transport of air pollutants, *Atmos. Environ.*, 150, 366–378, doi:10.1016/j.atmosenv.2016.11.031, 2017.
- 30 Hyvärinen, A.-P., Lihavainen, H., Komppula, M., Sharma, V. P., Kerminen, V.-M., Panwar, T. S. and Viisanen, Y.: Continuous measurements of optical properties of atmospheric aerosols in Mukteshwar, northern India, *J. Geophys. Res.*, 114(D8), D08207, doi:10.1029/2008JD011489, 2009.
- García, M. I., Van Drooge, B. L., Rodríguez, S. and Alastuey, A.: Speciation of organic aerosols in the Saharan Air Layer and in the free troposphere westerlies, *Atmos. Chem. Phys.*, 17(14), 8939–8958, doi:10.5194/acp-17-8939-2017, 2017.

- Ketterer, C., Zieger, P., Bukowiecki, N., Collaud Coen, M., Maier, O., Ruffieux, D. and Weingartner, E.: Investigation of the planetary boundary layer in the Swiss alps using remote sensing and in situ measurements, *Boundary-Layer Meteorol.*, 151, 317–334, doi:10.1007/s10546-013-9897-8, 2014.
- Kleissl, J., Honrath, R. E., Dziobak, M. P., Tanner, D., Val Martín, M., Owen, R. C. and Helmig, D.: Occurrence of upslope flows at the Pico mountaintop observatory: A case study of orographic flows on a small, volcanic island, *J. Geophys. Res. Atmos.*, 112(10), 1–16, doi:10.1029/2006JD007565, 2007.
- Legreid, G., Folini, D., Staehelin, J., Lööv, J. B., Steinbacher, M. and Reimann, S.: Measurements of organic trace gases including oxygenated volatile organic compounds at the high alpine site Jungfraujoch (Switzerland): Seasonal variation and source allocations, *J. Geophys. Res. Atmos.*, 113(5), 1–11, doi:10.1029/2007JD008653, 2008.
- Lugauer, M., Baltensperger, U., Furger, M., Gäggeler, H. W., Jost, D. T., Schwikowski, M. and Wanner, H.: Aerosol transport to the high Alpine sites Jungfraujoch (3454 m asl) and Colle Gnifetti (4452 m asl), *Tellus, Ser. B Chem. Phys. Meteorol.*, 50(1), 76–92, doi:10.3402/tellusb.v50i1.16026, 1998.
- Marcq, S., Laj, P., Roger, J. C., Villani, P., Sellegri, K., Bonasoni, P., Marinoni, A., Cristofanelli, P., Verza, G. P. and Bergin, M.: Aerosol optical properties and radiative forcing in the high Himalaya based on measurements at the Nepal Climate Observatory-Pyramid site (5079 m a.s.l.), *Atmos. Chem. Phys.*, 10, 5859–5872, doi:10.5194/acp-10-5859-2010, 2010.
- Marinoni, A., Cristofanelli, P., Calzolari, F., Roccatò, F., Bonafè, U. and Bonasoni, P.: Continuous measurements of aerosol physical parameters at the Mt. Cimone GAW Station (2165 m asl, Italy), *Sci. Total Environ.*, 391(2), 241–251, doi:10.1016/j.scitotenv.2007.10.004, 2008.
- Marinoni, A., Cristofanelli, P., Laj, P., Duchi, R., Calzolari, F., Decesari, S., Sellegri, K. and Vuillermoz, E.: and Physics Aerosol mass and black carbon concentrations , a two year record at NCO-P ( 5079 m , Southern Himalayas ), *Atmos. Chem. Phys.*, 10, 8551–8562, doi:10.5194/acp-10-8551-2010, 2010.
- McClure, C. D., Jaffe, D. A. and Gao, H.: Carbon dioxide in the free troposphere and boundary layer at the Mt. Bachelor observatory, *Aerosol Air Qual. Res.*, 16(3), 717–728, doi:10.4209/aaqr.2015.05.0323, 2016.
- Nyeki, S., Kalberer, M., Colbeck, I., De Wekker, S., Furger, M., Gäggeler, H. W., Kossmann, M., Lugauer, M., Steyn, D., Weingartner, E., Wirth, M. and U. Baltensperger, U.: Convective Boundary Layer Evolution to 4 km asl over High-Alpine Terrain: Airborne Lidar Observations in the Alps, *Geophys. Res. Lett.*, 27(5), 689–692, 2000.
- Nyeki, S., Eleftheriadis, K., Baltensperger, U., Colbeck, I., Fiebig, M., Fix, A., Kiemle, C., Lazaridis, M. and Petzold, A.: Airborne Lidar and in-situ Aerosol Observations of an Elevated Layer , Leeward of the European Alps and Apennines, *Geophys. Res. Lett.*, 29(17), 3–6, doi:10.1029/2002GL014897, 2002.
- Obrist, D., Hallar, A. G., McCubbin, I., Stephens, B. B. and Rahn, T.: Atmospheric mercury concentrations at Storm Peak Laboratory in the Rocky Mountains: Evidence for long-range transport from Asia, boundary layer contributions, and plant mercury uptake, *Atmos. Environ.*, 42(33), 7579–7589, doi:10.1016/j.atmosenv.2008.06.051, 2008.
- Okamoto, S. and Tanimoto, H.: A review of atmospheric chemistry observations at mountain sites, *Prog. Earth Planet. Sci.*,

- 3(34), doi:10.1186/s40645-016-0109-2, 2016.
- Pandey Deolal, S., Henne, S., Ries, L., Gilge, S., Weers, U., Steinbacher, M., Staehelin, J. and Peter, T.: Analysis of elevated springtime levels of peroxyacetyl nitrate (PAN) at the high Alpine research sites Jungfraujoch and Zugspitze, *Atmos. Chem. Phys.*, 14(22), 12553–12571, doi:10.5194/acp-14-12553-2014, 2014.
- 5 Pandolfi, M., Cusack, M., Alastuey, A. and Querol, X.: Variability of aerosol optical properties in the Western Mediterranean Basin, *Atmos. Chem. Phys.*, 11(15), 8189–8203, doi:10.5194/acp-11-8189-2011, 2011.
- Pandolfi, M., Martucci, G., Querol, X., Alastuey, A., Wilsenack, F., Frey, S., O’Dowd, C. D. and Dall’Osto, M.: Continuous atmospheric boundary layer observations in the coastal urban area of Barcelona during SAPUSS, *Atmos. Chem. Phys.*, 13(9), 4983–4996, doi:10.5194/acp-13-4983-2013, 2013.
- 10 Pandolfi, M., Ripoll, A., Querol, X. and Alastuey, A.: Climatology of aerosol optical properties and black carbon mass absorption cross section at a remote high-altitude site in the western Mediterranean Basin, *Atmos. Chem. Phys.*, 14(12), 6443–6460, doi:10.5194/acp-14-6443-2014, 2014.
- Panwar, T. S., Hooda, R. K., Lihavainen, H., Hyvarinen, A. P., Sharma, V. P. and Viisanen, Y.: Atmospheric aerosols at a regional background Himalayan site - Mukteshwar, India, *Environ. Monit. Assess.*, 185(6), 4753–4764, doi:10.1007/s10661-012-2902-8, 2013.
- 15 Poltera, Y., Martucci, G., Collaud Coen, M., Hervo, M., Emmenegger, L., Henne, S., Brunner, D. and Haeefe, A.: PathfinderTURB: an automatic boundary layer algorithm. development, validation and application to study the impact on in-situ measurements at the Jungfraujoch, *Atmos. Chem. Phys.*, 17, 10051–10070, doi:10.5194/acp-2016-962, 2017.
- Ripoll, A., Pey, J., Minguillón, M. C., Pérez, N., Pandolfi, M., Querol, X. and Alastuey, A.: Three years of aerosol mass, black carbon and particle number concentrations at Montsec (southern Pyrenees, 1570 m a.s.l.), *Atmos. Chem. Phys.*, 14(8), 4279–4295, doi:10.5194/acp-14-4279-2014, 2014.
- 20 Ripoll, A., Minguillón, M. C., Pey, J., Jimenez, J. L., Day, D. A., Sosedova, Y., Canonaco, F., Prévôt, A. S. H., Querol, X. and Alastuey, A.: Long-term real-time chemical characterization of submicron aerosols at Montsec (southern Pyrenees, 1570 m a.s.l.), *Atmos. Chem. Phys.*, 15(6), 2935–2951, doi:10.5194/acp-15-2935-2015, 2015.
- 25 Rodríguez, S., Alastuey, A. and Querol, X.: A review of methods for long term in situ characterization of aerosol dust, *Aeolian Res.*, 6, 55–74, doi:10.1016/j.aeolia.2012.07.004, 2012.
- Rose, C., Sellegrì, K., Asmi, E., Hervo, M., Freney, E., Colomb, A., Junninen, H., Duplissy, J., Sipilä, M., Kontkanen, J., Lehtipalo, K. and Kulmala, M.: Major contribution of neutral clusters to new particle formation at the interface between the boundary layer and the free troposphere, *Atmos. Chem. Phys.*, 15(6), 3413–3428, 2015.
- 30 Rose, C., Sellegrì, K., Moreno, I., Velarde, F., Ramonet, M., Weinhold, K., Krejci, R., Wiedensohler, A., Ginot, P. and Laj, P.: CCN production by new particle formation in free troposphere, *Atmos. Chem. Phys.*, 17(2), 1529–1541, doi:10.5194/acp-2016-696, 2017.
- Rotach, M. W. and Zardi, D.: On the boundary-layer structure over highly complex terrain: Key findings from MAP, *Q. J. R. Meteorol. Soc.*, doi:10.1002/qj.71, 2007.

- Rotach, M. W. . A. G., Lang, M. N. ., Leukauf, D., Stiperski, I. and Wagner, J. S.: On the Vertical Exchange of Heat, Mass, and Momentum Over Complex, Mountainous Terrain, *Front. Earth Sci.*, 3, doi:10.3389/feart.2015.00076, 2015.
- Rucker, M., Banta, R. M. and Steyn, D. G.: Along-Valley Structure of Daytime Thermally Driven Flows in the Wipp Valley, *J. Appl. Meteorol. Climatol.*, 47, 733–751, doi:10.1175/2007JAMC1319.1, 2008.
- 5 Schauer, G., Kasper-Giebl, A. and Mocnik, G.: Increased PM concentrations during a combined wildfire and saharan dust event observed at high-altitude sonnblick observatory, Austria, *Aerosol Air Qual. Res.*, 16(3), 542–554, doi:10.4209/aaqr.2015.05.0337, 2016.
- Schmeissner, T., Krejci, R., Ström, J., Birmili, W., Wiedensohler, A., Hochschild, G., Gross, J., Hoffmann, P. and Calderon, S.: Analysis of number size distributions of tropical free tropospheric aerosol particles observed at Pico Espejo (4765 m a.s.l.), Venezuela, *Atmos. Chem. Phys.*, 11(7), 3319–3332, doi:10.5194/acp-11-3319-2011, 2011.
- 10 Schwanghart, W. and Scherler, D.: Short Communication: TopoToolbox 2 - MATLAB-based software for topographic analysis and modeling in Earth surface sciences, *Earth Surf. Dyn.*, 2(1), 1–7, doi:10.5194/esurf-2-1-2014, 2014.
- Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Kluwer Aca., Springer Science {&} Business Media, Dordrecht/Boston /London., 1988.
- 15 Stull, R. B.: A theory for mixed-layer-top levelness over irregular topography, in *Proceedings of the 10th AMS Symposium on Turbulence and Diffusion*, 1992.
- Tsamalis, C., Ravetta, F., Gheusi, F., Delbarre, H. and Augustin, P.: Mixing of free-tropospheric air with the lowland boundary layer during anabatic transport to a high altitude station, *Atmos. Res.*, 143, 425–437, doi:10.1016/j.atmosres.2014.03.011, 2014.
- 20 Tunved, P., Ström, J. and Krejci, R.: Arctic aerosol life cycle: Linking aerosol size distributions observed between 2000 and 2010 with air mass transport and precipitation at Zeppelin station, Ny-Ålesund, Svalbard, *Atmos. Chem. Phys.*, 13(7), 3643–3660, doi:10.5194/acp-13-3643-2013, 2013.
- Venzac, H., Sellegri, K., Villani, P., Picard, D. and Laj, P.: Seasonal variation of aerosol size distributions in the free troposphere and residual layer at the puy de Dôme station, France, *Atmos. Chem. Phys.*, 9, 1465–1478, doi:10.5194/acp-9-1465-2009, 2009.
- 25 Venzac, H., Laj, P. and Sellegri K.: High frequency new particle formation in the Himalayas, *Pnas*, 105, 15666–15671, doi:10.1073/pnas.0801355105, 2008.
- Wang, W., Gong, W., Mao, F. and Pan, Z.: An improved iterative fitting method to estimate nocturnal residual layer height, *Atmosphere*, 106, 1–11, doi:10.3390/atmos7080106, 2016.
- 30 Wang, X. L. and Swail, V. R.: Changes of Extreme Wave Heights in Northern Hemisphere Oceans and Related Atmospheric Circulation Regimes, *J. Clim.*, 14(10), 2204–2221, [https://doi.org/10.1175/1520-0442\(2001\)014<2204:COEWHI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<2204:COEWHI>2.0.CO;2), 2001.
- De Wekker, S. F. J., Steyn, D. G. and Nyeki, S.: A comparison of aerosol-layer and convective boundary-layer structure over a mountain range during staaarte '97, *Boundary-Layer Meteorol.*, 113, 249–271, 2004.

- De Wekker, S. F. J. and Kossmann, M.: Convective Boundary Layer Heights Over Mountainous Terrain—A Review of Concepts, *Front. Earth Sci.*, 3, 77, doi:10.3389/feart.2015.00077, 2015.
- Whiteman, C. D., Hoch, S. W. and Poulos, G. A.: Evening Temperature Rises on Valley Floors and Slopes : Their Causes and Their Relationship to the Thermally Driven Wind System, *J. Appl. Meteorol. Climatol.*, 48, 776–788, doi:10.1175/2008JAMC2028.1, 2009.
- 5 Zardi, D. and Whiteman, C. D.: Diurnal Mountain Wind Systems, in *Mountain weather research and forecasting*, edited by F. K. Chow, S. F. J. De Wekker, and B. Snyder, Berlin, 2012.
- Zhang, X., Ming, J., Li, Z., Wang, F. and Zhang, G.:The online measured black carbon aerosol and source orientations in the Nam Co region, Tibet, *Environ. Sci. Pollut. Res.*, doi:10.1007/s11356-017-0165-1, 2017.
- 10 Zellweger, C., Forrer, J., Hofer, P., Nyeki, S., Schwarzenbach, B., Weingartner, E., Ammann, M. and Baltensperger, U.: Partitioning of reactive nitrogen (NO<sub>y</sub>) and dependence on meteorological conditions in the lower free troposphere, *Atmos. Chem. Phys.*, 3, 779-796, doi:10.5194/acp-3-779-2003, 2003.
- Zellweger, C., Hüglin, C., J, K., M, S., Vollmer, M. and Buchmann, B.: Inter-comparison of four different carbon monoxide measurement techniques and evaluation of the long-term carbon monoxide time series of Jungfraujoch, *Atmos. Chem. Phys.*, 15 9, 3491–3503, 2009.

## Tables

Table 1: List of station names, acronyms, latitude [°], longitude [°], altitude [m], their mountain range or region and continent. If aerosol time series were used, the station name is given in bold. The references principally describe the station measurement program and, particularly, the aerosol parameters measured.

Station	Latitude	Longitude	Altitude	Massif	Continent	References
<b>HPB</b> <b>Hohenpeissenberg, Germany</b>	47.8015	11.0096	985	Alps	Europe	(Flentje et al., 2015)
<b>JFJ</b> <b>Jungfrauoch, Switzerland</b>	46.5477	7.985	3580			(Bukowiecki et al., 2016)
<b>SBO</b> <b>Sonnblick, Austria</b>	47.0539	12.951	3106			(Schauer et al., 2016)
<b>ZSF</b> <b>Schneefernhaus, Germany</b>	47.4165	10.9796	2671			(Birmili et al., 2009)
<b>ZUG</b> <b>Zugspitze, Germany</b>	47.4211	10.9859	2962			--
<b>MSA</b> <b>Montsec, Spain</b>	42.05	0.7333	1570	Pyrenees	Europe	(Ealo et al., 2016; Pandolfi et al., 2014; Ripoll et al., 2014)
<b>MSY</b> <b>Montseny, Spain</b>	41.7795	2.3579	700			(Pandolfi et al., 2011)
<b>PDM</b> <b>Pic du Midi, France</b>	42.9372	0.1411	2877			(Gheusi et al., 2011, Hulin et al., 2017)
<b>BEO</b> <b>Moussala, Bulgaria</b>	42.1792	23.5856	2925	Balkan		(Angelov et al., 2016)
<b>CMN</b> <b>Monte Cimone, Italy</b>	44.1667	10.6833	2165	Apennines		(Cristofanelli et al., 2016; Marinoni et al., 2008)
<b>HAC</b> <b>Mount Helmos, Greece</b>	37.9843	22.1963	2314	Peloponnese		

Field Code Changed

<b>PUY</b> <b>Puy de Dôme, France</b>	45.7723	2.9658	1465	Central massif		(Venzac et al., 2009)
<b>CHC</b> <b>Chacaltaya, Bolivia</b>	-16.200	-68.100	5320			(Andrade et al., 2015)
<b>LQO</b> La Quiaca Observatorio, Argentina	-22.100	-65.599	3459	Andes	South America	(Hamburger et al., 2013; Schmeissner et al., 2011)
<b>PEV</b> <b>Pico Espeje, Venezuela</b>	8.5167	-71.05	4765			
<b>TLL</b> <b>Cerro Tololo, Chile</b>	- 30.1725	-70.7992	2220			
<b>MZW</b> Mount Zirkel Wildness, USA	40.5433	-106.6844	3243			
<b>NWR</b> <b>Niwot Ridge, USA</b>	40.04	-105.54	3035	Rocky Mountains		
<b>SPL</b> <b>Steamboat, USA</b>	40.455	-106.744	3220			
<b>YEL</b> Yellowstone NP, USA	44.5654	-110.4003	2430		North America	YEL Yellowstone NP, USA
<b>APP</b> Appalachian State University, USA	36.2130	-81.6920	1076	Appalachian		
<b>SHN</b> Shenandoah National Park, USA	38.5226	-78.4358	1074			
<b>MBO</b> Mount Bachelor, USA	43.979	-121.687	2743			
<b>MWO</b> Mount Washington	44.2703	-71.3033	1916			
<b>WHI</b> <b>Whistler, Canada</b>	50.0593	-122.9576	2182			(Gallagher et al., 2011)

Field Code Changed

Formatted: Italian (Switzerland)

<b>HLE</b> Henle, India	32.7794	78.9642	4517		
<b>LAN</b> Langtang, Nepal	28.2200	85.6200	3920		
<b>MUK</b> <b>Mukteshwar, India</b>	29.4371	79.6194	2180	Himalaya	(Hyvärinen et al., 2009; Panwar et al., 2013)
<b>NCOS</b> <b>Nam Co, China</b>	30.7728	90.9621	4730		(Zhang et al., 2017)
<b>PYR</b> <b>ABC Pyramid, Nepal</b>	27.9578	86.8149	5079		(Bonasoni et al., 2010; Marcq et al., 2010; Marinoni et al., 2010)
<b>SZZ</b> Shangrimla ZhuZhang, China	27.9998	99.4266	3583		
<b>WLG</b> <b>Mount Waligan, China</b>	36.2875	100.8963	3810	Tibetan Plateau	(Andrews et al., 2011)
<b>PDI</b> <b>Pha Din, Vietnam</b>	21.5728	103.5160	1466	--	--
<b>FWS</b> Mount Fuji, Japan	35.3606	138.7273	3776		
<b>HPO</b> Mount Happa, Japan	36.6972	137.7989	1850		
<b>MTA</b> Mount Takayama, Japan	36.1461	137.4230	1420	Japan Alps	
<b>IZO</b> <b>Izaña, Spain</b>	28.309	-16.4994	2373	Atlantic	(Rodríguez et al., 2012)
<b>LLN</b> <b>Mount Lulin, Taiwan</b>	23.4686	120.8736	2862	Pacific	(Hsiao et al., 2017)
<b>MLO</b> <b>Mauna Loa, USA</b>	19.5362	-155.576	3397	Pacific	(Bodhaine, 1995)
<b>OMP (previously PICO-NARE)</b> <b>Pico Mountain, Azores, Portugal</b>	38.4704	-28.4039	2225	Atlantic	(Fialho et al., 2004)


<b>RUN</b> Ile de la Réunion, France	-21.0795	55.3831	2160	Indian	
<b>TDE</b> Izaña, Spain	28.2702	-16.6385	3538	Atlantic	
<b>ASK</b> Assekrem, Algeria	23.2667	5.6333	2710		<b>Africa</b>
<b>MKN</b> Mount Kenia	-0.0622	37.2972	3678		
<b>SUM</b> Summit, Arctic	72.58	-38.48	3238		<b>Arctic</b>
<b>ZEP</b> Zeppelin Observatory, Norway	78.9067	11.8893	475		

Table 2: Extrema, median and mean of the topographical parameters for the 46 stations studied.

5

Parameter	min	median	mean	max
ABL-TopoIndex	0.22	1.72	4.11	30.12
Hypso% [%]	0.005	4.8	16.4	79.1
HypsoD50 [m]	-872	1192	1160	4019
LocSlope (*10 <sup>-3</sup> )	1.7	86	93	259
G8 [tangent]	0.0024	0.1743	0.2053	0.4982
DBinv [km <sup>2</sup> ]	423	86426	93287	249464
Altitude [m]	475	2771	2802	5320
Latitude  [°]	0.06	37.3	36.2	78.9

Figures:



5

Figure 1: Map of the stations colored by their mountain ranges or region.




Figure 2: Schematic view of the topographical features underlying the ABL-TopoIndex.




Figure 3: a) Normalized hypsocurves for some selected high altitude stations for a 500 km x 500 km domain centered on the station. The filled and open circles correspond to the normalized station elevations within the domain and indicate the value of hypso% (e.g., PYR hypso% is 26). The vertical dashed line corresponds to 50% of the hypsometric curve b) Difference between the station altitude and the elevation minimum in a domain of radius R around the station as a function of R. The vertical dashed line indicates the part of the curve selected to calculate LocSlope.




Figure 4: a) Topography on a  $750 \times 750 \text{ km}^2$  domain around BEO (Moussala, white dot) in Bulgaria. The main hydrologic flow paths from the station grid cell are given by the cyan lines. The color scale on the left only applies to Fig. 4a. b) hydrographical network, c) hydrologic drainage basins calculated from the real topography, the different drainage basins are defined by various colors and d) “convective drainage basin” calculated from the inverse topography (DBinv).

10




Figure 5: Idem Fig. 4 for PYR (Nepal Climate Observatory - Pyramid) station in the Himalaya, Nepal.




Figure 6: a) ABL-TopoIndex, b) drainage basin for convection, c) hypsometric percentage of the station elevation, d) hypsometric percentage of the station elevation minus the 50% hypsometry, e) local slope in a circle of 10 km radius centered on the station, f) gradient in elevation as a function of the domain size for some European high altitude stations.





Figure 7: ABL-TopoIndex as a function of elevation of all grid cells of a 625 km<sup>2</sup> domain centered on the ASK, CHC, HAC, OMP, PYR and SBO stations. The squares indicate the ABL-TopoIndex values and the altitudes of the stations.



5

Figure 8: ABL-TopoIndex for all stations as a function of continents and mountainous ranges. The color scheme corresponds to that in Fig. 1.

10




Figure 9: Spearman's rank correlation coefficient characterizing the correlation between aerosol parameters (absorption coefficient, scattering coefficient, number concentration) and various topographic parameters (the ABL-TopoIndex, mean altitude over the 9 grid cells, station latitude and the 5 parameters constituting the ABL-TopoIndex (G8, DB, LocSlope, hypso% and hypsoD50)). Correlations were calculated for the 5th, 50th and 95th percentiles of the aerosol parameters. Statistically significant correlation values at 95% and 90% confidence levels are marked by large and medium symbol sizes and the positive and negative correlations are plotted with upward and downward triangles, respectively. The correlations were performed with 21, 23 and 17 stations for the absorption coefficient, the scattering coefficient and the number concentration, respectively.




Figure 10: Spearman's rank correlation coefficient characterizing the correlation between all the topographic parameters (see Fig. 9) and the minimum and the maximum of the monthly diurnal cycles, as well as the seasonal cycle of the aerosol parameters. The correlations are performed with 21, 22 and 15 stations for the absorption coefficient, the scattering coefficient and the number concentration, respectively.