

**Reply to Interactive comment on “Determination of the refractive index of insoluble organic extracts from atmospheric aerosol over the visible wavelength range using optical tweezers”  
by**

**Rosalie H. Shepherd et al.  
Anonymous Referee #2**

In this manuscript, Shepherd et al. present an optical trapping technique combined with white light spectroscopy to measure the real and imaginary parts of the refractive index for samples of insoluble material from ambient aerosol samples. They use the data to estimate the effects of organic film-coated particles in the atmosphere revealing significant changes in the top of the atmosphere albedo compared to an assumption of equal sized water droplets. This work is interesting and relevant and should be published once the following minor points are clarified or addressed:

Thank you.

1. In the introduction, references to the work of Reid and coworkers on refractive index measurements should be included.

The following references have been added:-

R.E.H. Miles, J.S. Walker, D.R. Burnham and J.P. Reid, ‘Retrieval of the Complex Refractive Index of Aerosol Droplets from Optical Tweezers Measurements’, *Physical Chemistry Chemical Physics* 14 (2012) 3037–3047.

H.-B. Lin, J.D. Eversole, A.J. Campillo, “Identification of morphology dependent resonances in stimulated Raman scattering from microdroplets”, *Optics Communications*, 77(5,6) (1990) 407-410

With the following text added to the paper:

“The use of morphological dependent resonances in Raman Spectra to determine refractive index at a fixed wavelength has been reported by Lin et al. (1990) and references therein and Miles et al. (2012).”

2. The statement of an “unparalleled level of accuracy” seems too strong given the much greater precision and accuracy achievable using cavity-enhanced Raman scattering in optical tweezers.

The statement has been removed and the text now reads:

“Application of the optical trapping technique was successfully employed to determine the refractive index of aerosol over a wide wavelength range”.

Note the Reviewer's comment on the greater precision and accuracy using the cavity enhanced Raman scattering in optical tweezers is erroneous as both techniques use the same fundamental principles of light scattering from a sphere. The Miles et al. (2012) study is undertaken with ideal samples where greater precision is possible owing to the ability to generate a spectrum with a lot of structure and allowing a precise fit. The samples presented in this work are atmospheric, very limited in amount and supplies were exhausted within a few minutes of nebulization. In our previous studies (Jones et al., 2013), we have achieved similar precisions to the cavity enhanced Raman scattering values.

The accuracy comment is surprising as the accuracy is ultimately based on the resolution of the spectrographs used which are similar here and in the work of Miles et al. (2012).

3. Was all the material extracted during sonication with chloroform and water? What about components that are insoluble in these solvents?

Two extra references (should) have been included:-

1. Folch J., Lees M., Sloane Stanley G. H. 1957. A simple method for the isolation and purification of total lipides from animal tissues. *J. Biol. Chem.* 226: 497-509.
2. Bligh, E. G. & Dyer, W. J. 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.* 37: 911-917.

And the text now reads:-

“The organic material was extracted from atmospheric aerosol based on techniques adapted from Folch et al(1957) and Bligh and Dyer (1959) and the refractive...”

Only material that was soluble in chloroform was analyzed. Extraction by chloroform is an accepted method to remove insoluble surface active compounds from complex media. The cited reference for this process, Bligh and Dyer has been cited on ~44,000 occasions.

Material that did not dissolve in either chloroform or water remained on the filter material or as solid detritus that was filtered out of the chloroform. The following text has been added:-

“The filter debris with un-extracted material was discarded.”

4. Did the authors observe any artifacts of the sonication process (due to formation of radicals) that indicate further chemistry was occurring and changing the samples? Was this controlled for (i.e. a short duration sonication versus much longer sonication)?

There was no change in chemical properties noted owing to sonication. However checks were performed and the following text has been added:-

“The sonication in the extraction process was not found to change the Langmuir isotherm of atmospheric material at the air-water interface.”

5. Why was the mass concentration of humic acid in the droplets so low? At 30% RH, surely most of the water is lost?

The mass concentration of humic acid in the nebulizing solution was kept “low” to prevent potential aerosol heating. As noted in Miles et al (2012) the local RH of a droplet and that measured by the RH probe are frequently different. However the reviewer’ query has highlighted an error which we have corrected with the following text:-

“The concentration of the trapped humic acid was determined to be  $0.016 \text{ g cm}^{-3}$ . The concentration of the aqueous humic acid droplet had increased by a factor of  $\sim 32$  upon trapping, thus demonstrating that water had evaporated from the droplet during the trapping and aerosol equilibration process.”

6. The x-axis in figure 5 is confusing – as it reads, if the shell and core volumes are equal, this parameters goes to infinity. Please clarify. The y-axis label does not match the text (should it be ARE\_TOT?). Further, in the caption, top, middle and bottom are referenced, but the figure is horizontal.

The x-axis has been fixed.

7. Are the droplets fully spherical in these experiments? What would be the signatures of asphericity and how would this impact the fitting of the data?

The following text has been added

“The liquid droplets are assumed to be perfectly spherical. Mie scattering from droplets experiencing small deformation has been shown<sup>3,4</sup> to result in resonances which shift, broaden and split as the droplet asymmetry increases.”

3. S. Arnold, D. E. Spock and L. M. Folan, Opt. Lett., 1990,15, 1111.

4 G. Schweiger, Opt. Lett., 1990, 15, 156.

The formation of an aspherical liquid droplet trapped in a gaseous environment would be somewhat surprising as the interfacial tension responsible for the spherical shape are in the 10’s of  $\text{mN m}^{-1}$  range and far exceed those exerted by the laser trap.

Also, orientation of a spherical particle in an optical trap is independent of the Mie spectrum recorded and thus a rotating spherical particle will not affect the recorded Mie spectrum. However, the orientation of an aspherical particle will give different Mie Spectra depending on orientation. Thus, an aspherical particle will give a changing Mie spectrum. During our previous work with Mie scattering of aerosol liquid droplets we have not encountered such behavior. We have experienced gross asymmetry during collisions of droplets and solid beads (unpublished) which results in complete loss of resonance behavior. Other tell-tale signs to any asphericity of the trapped particles would be an inability to fit the trapped particle to reasonable values of the Cauchy Coefficient as discussed in the new reference, Miles et al. (2012).

***Interactive comment on “Determination of the refractive index of insoluble organic extracts from atmospheric aerosol over the visible wavelength range using optical tweezers” by Rosalie H. Shepherd et al. Anonymous Referee #3***

**Anonymous Referee #3**

Review of Shepherd et al., ‘Determination of the refractive index of insoluble organic extracts from atmospheric aerosol over the visible wavelength range using optical tweezers’.

In this manuscript, the authors present measurements of the complex refractive index of insoluble ambient aerosol matter collected on filters from a range of different environments. Refractive index (RI) measurements were made using a combination of optical trapping, white light spectroscopy and UV-VIS spectroscopy. The retrieved refractive index values were then used to model the effect of core-shell particles coated with the organic matter on the aerosol radiative effect.

The reported research is original and well presented, and is suitable for publication

Thank you,

subject to the corrections/clarifications listed below being addressed:

1) Given that the paper looks at extracting the complex refractive index from optically trapped droplets using Mie theory, the authors should be aware of the following reference and include it in the introduction.

R.E.H. Miles, J.S. Walker, D.R. Burnham and J.P. Reid, ‘Retrieval of the Complex Refractive Index of Aerosol Droplets from Optical Tweezers Measurements’, *Physical Chemistry Chemical Physics* 14 (2012) 3037–3047.

The following references have been added:-

R.E.H. Miles, J.S. Walker, D.R. Burnham and J.P. Reid, ‘Retrieval of the Complex Refractive Index of Aerosol Droplets from Optical Tweezers Measurements’, *Physical Chemistry Chemical Physics* 14 (2012) 3037–3047.

H.-B. Lin, J.D. Eversole, A.J. Campillo, ‘Identification of morphology dependent resonances in stimulated Raman scattering from microdroplets’, *Optics Communications*, 77(5,6), (1990) 407-410

With the following text added to the paper:

“The use of morphological dependent resonances in Raman Spectra to determine refractive index at a fixed wavelength has been reported by Lin et al. (1990) and references therein and Miles et al. (2012).”

2) Page 3, Line 8-9.

The authors state that ‘The technique allowed the refractive index to be resolved to within 0.015 over a large wavelength range of 460 to 700 nm’. I take this sentence to mean that the uncertainty in the RI values extracted using this technique is +/- 0.015. However, in Table 1 much reduced uncertainties are given for the samples tested, which correspond instead to the standard deviation in RI values retrieved from multiple droplets. For some of the samples, no uncertainty is given at all. If the precision of the RI measurement is indeed +/- 0.015 as stated in the introduction (and again on page 10, line 22), then this is the error which should be included in Table 1.

The text in the caption for table 1 has been edited to be clearer:-

“Where more than one particle was analyzed for a single type of sample, e.g. urban spring, the average and standard deviation of the Cauchy coefficients and real refractive index were reported for the particle studied. The standard deviation does not reflect the uncertainty estimated from the fitting process (which is  $\pm 0.015$  for the real refractive index) but a spread of values obtained for the few particles studied. The range in particle sizes studied is also reported.”

The distinction is important as the purpose of table 1 is to demonstrate the number of experiments that were performed (as demonstrated by the first line of the caption “Table described the number of aerosols studied, their determined...” ) and the range of values measured. The

3) Page 6, Line 2-4.

The authors state that ‘The values of the three empirical constants and the radius of the trapped aerosol were iterated until a good comparison was achieved between the simulated and the experimentally obtained Mie spectrum’

Were the experimental and simulated spectra compared by eye or was a fitting algorithm used?


The following text has been added:-

“Typically, the radius of droplet was fixed and the values of A, B, and C varied until a good fit between measured and simulated Mie spectra was achieved by simple comparison (inspection) of peak, trough and inflection point positions. The value of the radius was then iterated through a series of radii with optimization of the values of A, B, and C as a function of wavelength repeated at each radius. Thus, a qualitative grid search was performed over parameter space. Parameter space was A varying from 1.3 to 1.7, B from 0 to 20,000 nm<sup>-2</sup> and C from 0 to  $1 \times 10^9$  nm<sup>4</sup>. The value of the radius was between 0 to 3 microns typically. The imaginary component of the refractive index was varied only after the grid search for the woodsmoke and humic acid samples shown in fig. 2.”

The authors should include a figure which shows how sensitive the simulated spectra are to small changes in the radius and RI, in particular for the more featureless spectra, so the reader can see for themselves the sensitivity of the fit.

The following text and figures have been added to the paper:-

“The sensitivity of the simulated Mie spectra to the refractive index ( $\pm 0.015$ ) and radius ( $\pm 6\text{nm}$ ) of the droplet are shown in figure 5. The simulated spectra, with the stated variations and the experimental Mie spectra for the Spring Urban aerosol extract are plotted. Figure 5 also contains a third simulated set of Mie spectra calculated by re-optimizing the values of A, B, and C in the Cauchy equation to achieve a fit between simulated and experimental Mie spectra for particles with a radius  $\pm 12\text{ nm}$  from the optimum fit to the experimental data. Figure 5 demonstrates that the quoted uncertainties in radius ( $\pm 6\text{nm}$ ) and refractive index ( $\pm 0.015$ ) are realistic.”



(New)”Figure 5: The experimental Mie spectra for the urban spring aerosol extract with the simulated best fit perturbed in three different scenarios to demonstrate the sensitivity of fitting simulated Mie spectra to experimental Mie spectra. Initially the simulated fit (red solid line) is recalculated with a refractive index increased and decreased by 0.015 ( $n\pm 0.015$ ), followed by the simulated fit (red solid line) recalculated with a radius increased and decreased by 6nm ( $r\pm 6\text{nm}$ ). The final set of simulated Mie spectra consider the simulated fit (red solid line) is recalculated with a droplet radius increased and decreased by 12nm, but the refractive index re-optimized to get the best fit to the experimental fit ( $r\pm 12\text{nm}$  and RI re-optimized). A clear demonstration that the quoted uncertainties in radius ( $\pm 6\text{nm}$ ) and refractive index ( $\pm 0.015$ ) are conservative and more than adequate.”



4) Page 6, Line 21-22.

The authors should state what the instrument detection limit of the UV-VIS spectrometer was in order to provide an upper bound on the absorbance of the atmospheric samples.

The baseline flatness of Perkin Elmer lambda 950 is quoted as  $\pm 0.0008A$  with a photometric reproducibility of  $0.00016A$  and a RMS photometric noise of  $0.0002A$ . The following text has been added

“The quoted photometric noise for the UV-Vis spectrometer was  $\sim 0.0002A$ . However, the urban and remote aerosol extracts were diluted in isopropanol to fill the UV-Vis spectrometer cuvette and thus a value three orders of magnitude larger than  $\sim 0.0002A$  may provide an upper bound for the absorbance of the samples reported below the detection limit.”

5) Figure 7

In keeping with the description in the text (Equation 4) the y-axis label should be ARE(TOT). Based on the description of the x-variable as the volume fraction of the shell (page 11, lines 2-3) the x-axis label also needs changing to  $V_{shell}/(V_{shell}+V_{core})$ .

The x-axis has been fixed.

6) There are several places in the text which seem to contradict each other as to how the Angstrom exponent was determined ie. from optical trapping or from UV-VIS spectroscopy, or from both. Please could the authors clarify the following explicitly: Were values of the Angstrom exponent extracted from the Mie spectra and from the UV-VIS spectra independently and these values then compared to show they were equivalent? The authors should include somewhere the values of the Angstrom exponent determined from each method, along with associated uncertainties.

The Angstrom exponent was determined from the measurement of the absorbance of an aerosol sample in isopropanol using a UV-Vis spectrometer. The value of Angstrom exponent was then adjusted (converted) for use with the imaginary refractive index as described in the appendix. The simulated Mie Spectra in Fig.2 were then calculated with and without an Angstrom component to demonstrate that the attenuation of the Mie Resonances (especially at the shorter wavelengths) was consistent with the measured Mie spectra (in figure2). The following text has been added:-

“The Angstrom coefficient determined for the Absorbance in isopropanol or water was adjusted for use with the imaginary refractive index Angstrom relationship (see Appendix). Simulated Mie spectra of wood smoke aerosol and humic acid aerosol were then calculated with and without an Angstrom exponent



absorption to demonstrate that the attenuation in Mie resonance intensity was consistent with absorption as shown in figure 2.”

and

“The simulated Mie spectra in Fig 2. were calculated with and without absorption described by an Angstrom exponent to demonstrate that the attenuation of Mie resonances (especially at shorter wavelengths) was consistent with measured Mie spectra.”

and

“In a UV-Vis spectrometer the absorption coefficient,  $\beta$ , can be related to the Absorbance, Abs, by,

$$\text{Abs} = -\beta l \quad (3)$$

where  $l$  is the pathlength (1 cm for the work described here) and absorbance, Abs has been corrected from base 10 to base e (Petty 2006). The absorption coefficient can be related to the imaginary refractive index,  $k(\lambda)$ , by

$$k(\lambda) = \frac{\lambda\beta}{4\pi} \quad (4)$$

as described by Petty (2006). Substitution of equation (4) into equation (3) and subsequently equation (2) demonstrates that the Ångström relationship for absorbance (equation 2) is modified to

$$\frac{k}{k_0} = \left(\frac{\lambda}{\lambda_0}\right)^{-(\alpha-1)} \quad (5)$$

for describing the imaginary refractive index (see appendix). In essence the value of Ångström exponent,  $\alpha$ , measured by the UV-Vis spectrometer is larger than the corresponding value for the imaginary refractive index. Note for the work described here  $\lambda_0=460\text{nm}$ . The values of  $k_0$  and  $\alpha$  were measured for dilute solutions of the wood smoke extract in isopropanol and humic acid in water. In the optical trap the trapped droplet of wood smoke extract in isopropanol lost all of the isopropanol solvent to evaporation as expected, leaving pure wood smoke extract. The aqueous humic acid solution lost some water to evaporation, but remained an aqueous and more concentrated, solution. As will be described below the mass density of the woodsmoke extract was measured independently. Thus, for the wood smoke droplet the values of  $k_0$  and  $\text{Abs}_0$  were corrected for the mass density of wood smoke extract in the optical trap and the attenuation of the resulting Mie spectrum will be shown to be consistent. For the aqueous humic acid solution, the value of  $k_0$  was determined by fitting the attenuation of the Mie spectrum by inspection, *i.e.* by changing the value of  $k_0$  until the intensity

attenuation of the simulated and experimental Mie spectra matched, thereby calculating the value of the mass density in the trapped humic acid droplet.”

The values and uncertainties for the Angstrom coefficient are reported in a new table 2.

Sample	Mass density / g cm <sup>-3</sup>	Absorption coefficient, $\beta$ , at $\lambda_0 = 460$ nm / cm <sup>-1</sup>	$k_0$ ( $\lambda_0 = 460$ nm)	Abso ( $\lambda_0 = 460$ nm)	$\alpha$
Pure Wood smoke extract B	1.64	3033	0.0111±0.0010	-	9.0±0.1
Wood smoke extract B in isopropanol (Fig. 3 - top pane)	6.60×10 <sup>-5</sup>	0.122	(4.47±0.40)×10 <sup>-7</sup>	0.122	9.0±0.1
Aqueous Humic acid (Fig. 3 - top pane)	7.00×10 <sup>-5</sup>	1.513	(4.214±0.38)×10 <sup>-6</sup>	0.499	4.2±0.1
Pure Humic acid	1.52	25,000	0.092±0.046	-	4.2±0.1
Optically trapped aqueous Humic acid droplet	0.016	273.2	(1.00±0.50)×10 <sup>-3</sup>	-	4.2±0.1

**Reply to Interactive comment on “Determination of the refractive index of insoluble organic extracts from atmospheric aerosol over the visible wavelength range using optical tweezers”  
by**

**Rosalie H. Shepherd et al.**

**Anonymous Referee #1**

Interesting experimental work dealing with the characterization of aerosol particles where the imaginary part of the refractive index cannot be neglected.

Thank you

Questions/comments that should be addressed:

1. To echo what the other two reviewers stated: It seems like some important references regarding the determination of the refractive index of single aerosol particles using Mie theory are missing from the introduction.

The following references have been added:-

R.E.H. Miles, J.S. Walker, D.R. Burnham and J.P. Reid, ‘Retrieval of the Complex Refractive Index of Aerosol Droplets from Optical Tweezers Measurements’, *Physical Chemistry Chemical Physics* 14 (2012) 3037–3047.

H.-B. Lin, J.D. Eversole, A.J. Campillo, “Identification of morphology dependent resonances in stimulated Raman scattering from microdroplets”, *Optics Communications*, 77(5,6) (1990) 407-410

With the following text added to the paper:

“The use of morphological dependent resonances in Raman Spectra to determine refractive index at a fixed wavelength has been reported by Lin et al. (1990) and references therein and Miles et al. (2012).”

2. Were the Mie spectra calculated by integrating over the acceptance angle of the objective? Please add a sentence or two that qualitatively describes the calculation. On page 3 it is stated "simulating the spectrum with Mie calculations." That is too vague.

Yes, the following text has been changed to read:-

“The optically trapped aerosol droplet was illuminated with white light and the elastically backscattered light was collected over a 25-degree cone angle, by an

objective lens. Further optics (described in Jones et al, 2013) focused the light onto a spectrometer (Acton, SP1500i).”

and

“The measured Mie spectrum was simulated through the application of Bohren and Huffman(1983) formalism of Mie theory integrating over 25 degree cone angle of backscattered light. In conjunction with the Cauchy Equation...”

3. Can you provide more details on how the size and Cauchy parameters were found. A simple grid search? If so, over what space? Was the imaginary part of the refractive index included in the search space?

The following text has been added:-

“Typically, the radius of droplet was fixed and the values of A, B, and C varied until a good fit between measured and simulated Mie spectra was achieved by simple comparison (inspection) of peak, trough and inflection point positions. The value of the radius was then iterated through a series of radii with optimization of the values of A, B, and C as a function of wavelength repeated at each radius. Thus, a qualitative grid search was performed over parameter space. Parameter space was A varying from 1.3 to 1.7, B from 0 to 20,000 nm<sup>-2</sup> and C from 0 to 1×10<sup>9</sup> nm<sup>4</sup>. The value of the radius will be between 0 to 3 microns typically. The imaginary component of the refractive index was varied only after the grid search for the woodsmoke and humic acid samples.”

The Imaginary part of the refractive index was described by an Ångström equation. The value of the Ångström exponent was calculated from the UV-Vis Absorbance measurement of the bulk sample and shown to be consistent with the attenuated Mie Spectra in figure 2. The issue and changes to text are further addressed in our answer to point 6, from reviewer 3.

4. In Section 4.4, the uncertainty associated with fitting less structured Mie spectra is mentioned. What is the origin of this increased uncertainty when fitting Mie spectra of absorbing particles or submicron particles studied at optical wavelengths? Presumably, Mie theory allows you to calculate the observed spectrum very accurately. Yet, when sharp peaks are absent, the uncertainty in the retrieved parameters increases substantially. A thorough answer to this question is not necessary but a slightly more detailed comment in the text would be helpful.

The following text has been added:-

“The Mie spectra in figure 1 of Summer Urban aerosol extract are structured with pronounced peak shapes that allow the facile fitting between simulated and

measured Mie Spectra. Such spectra allow a relatively small range of values of radius, Cauchy coefficients A, B and C to provide a good fit between measured and simulated Mie Spectra. The rest of the Mie spectra in Figure 1 have significantly fewer Mie resonances and their peak shapes are less pronounced. The uncertainties become larger as the spectra become less structured.”

# Determination of the refractive index of insoluble organic extracts from atmospheric aerosol over the visible wavelength range using optical tweezers

Rosalie H. Shepherd<sup>1,2</sup>, Martin D. King<sup>2</sup>, Amelia Marks<sup>2</sup>, Neil Brough<sup>3</sup> and Andrew D. Ward<sup>1</sup>

5 <sup>1</sup>Central Laser Facility, Research Complex, STFC Rutherford Appleton Laboratory, Oxford, OX11 0FA, UK

<sup>2</sup>Department of Earth Sciences, Royal Holloway University of London, Egham, Surrey, TW20 0EX, UK

<sup>3</sup>British Antarctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, UK

Correspondence to: Martin King (m.king@rhul.ac.uk)

10 Optical trapping combined with Mie spectroscopy is a new technique used to record the refractive index of insoluble organic material extracted from atmospheric aerosol samples over a wide wavelength range with sub nanometer resolutions. The refractive index of the insoluble organic extracts was shown to follow a Cauchy equation between 460 to 700 nm for organic aerosol extracts collected from urban (London) and remote (Antarctica) locations. Cauchy coefficients for the remote sample were for the  
15 Austral summer and gave the Cauchy coefficients to be  $A = 1.467$  and  $B = 1000 \text{ nm}^2$  with a real refractive index of 1.489 at a wavelength of 589 nm. Cauchy coefficients for the urban samples varied with season, with extracts collected during summer having Cauchy coefficients of  $A = 1.465 \pm 0.005$  and  $B = 4625 \pm 1200 \text{ nm}^2$  with a representative real refractive index of 1.478 at a wavelength of 589 nm, whilst samples extracted during autumn had larger Cauchy coefficients of  $A = 1.505$  and  $B = 600 \text{ nm}^2$  with a representative real refractive index of 1.522 at a wavelength of 589 nm. The refractive index of  
20 absorbing aerosol was also recorded. The technique applied in the presented study allowed the absorption Ångström exponent to be determined for wood smoke and humic acid aerosol extract. Typical values of the Cauchy coefficient for the wood smoke aerosol extract were  $A = 1.541 \pm 0.03$  and  $B = 14800 \pm 2900 \text{ nm}^2$  resulting in a real refractive index of  $1.584 \pm 0.007$  at a wavelength of 589 nm and an absorption Ångström exponent of 8.0. The measured values of refractive index compare well with  
25 previous monochromatic or very small wavelength range measurements of refractive index. In general, the real component of the refractive index increases from remote to urban to wood smoke. A one-dimensional radiative-transfer calculation of the top of the atmosphere albedo was applied to model an atmosphere containing a 3 km thick layer of aerosol comprising of pure water, insoluble organic aerosol or an aerosol consisting of an aqueous core-with an insoluble organic shell. The calculation  
30 demonstrated that the top of the atmosphere albedo increases by 0.01 to 0.04 for pure organic particles relative to water particles of the same size and the top of the atmosphere albedo increases by 0.03 for aqueous core-shell particles as volume fraction of the shell material increases by 25 percent.

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Deleted: Ångstrom

Formatted: Not Superscript/ Subscript

Deleted: Ångstrom

Deleted: 7

## 1. Introduction

Atmospheric aerosols affect the radiative balance of our planet e.g Pöschl (2005), Ramanathan (2001), Stocker (2013) and Wild (2009). Aerosols directly affect the radiative balance by absorbing or scattering incoming solar radiation e.g. Moise (2015), and indirectly through their role as cloud condensation nuclei e.g Breon (2002), Lohmann and Feichter (2005), Rosenfeld (2008) and Charlson (2001). However, current understanding of the atmospheric aerosol radiative balance and the cloud albedo effect is currently regarded as low e.g. Stocker (2013) and Fuzzi (2005).

Owing to the complexity of material in the atmosphere, atmospheric aerosols often contain a mixture of different compounds e.g. Cappa (2011), Cai (2016) and Cochran (2016). The contrasting properties of the different compounds can result in atmospheric aerosols being coated in a film of organic material e.g. Gill (1983) and Donaldson and Vaida (2006). The presence of a film may alter the physical, chemical and optical properties of the cloud droplet or aerosol particle by (a) reducing the rate of evaporation from the droplets or particles e.g. Davies (2013), Eliason (2003), Gill (1983), Kaiser (1996) and McFiggans (2005), (b) altering the transport of chemicals between gas and liquid phase e.g. Donaldson and Anderson (1999) and Donaldson and Valsaraj (2010), (c) affecting gaseous uptake e.g. Enami, (2010), (d) reducing the scavenging by larger cloud droplets e.g. Andreaea and Rosenfeld (2008) and Feingold and Chuang (2002) and (e) altering the optical properties of the droplet e.g. Donaldson and Vaida (2006) and Li (2011). To quantify the consequences of film formation on atmospheric aerosol particles and cloud droplets, further understanding of aerosol films is urgently required. However, obstacles such as accurately quantifying anthropogenic and natural aerosol emissions (Kanakidou (2005)) or understanding the hugely varying chemistry of atmospheric aerosols (Jacobson and Hansson (2000)) means the current understanding of atmospheric aerosol remains low e.g. Stocker (2013) and Flores (2014).

Furthering the current understanding of film formation on cloud droplets or aerosols has largely been addressed by replicating the film with atmospheric proxy compounds such as oleic acid e.g. King (2004, 2009) and Jones (2015), methyl oleate e.g. Pfrang (2014) and Sebastiani (2015), pinonic acid e.g. Enami and Sakamoto (2016), DPPC (Thompson et al. 2010), or nonanoic acid e.g. Tinel (2016). However organic material extracted from the environment has only featured in a limited number of studies, e.g. Jones (2017) investigated the consequences of exposing organic films extracted from atmospheric aerosol and sea-water to ozone, whilst Zhou (2014) studied the heterogeneous chemistry of material collected from the surface of the sea. Research that focus on the sea-surface layer is highly relevant to atmospheric studies owing to sea-surface material often becoming aerosols (Blanchard (1964)).

The work presented describes the application of the optical trapping technique to measure the refractive index of organic material that may form an organic film on atmospheric aerosol. The organic material was extracted from atmospheric aerosol (using the techniques of Folch and Lee, 1957 and Bligh and Dyer, 1959) and the refractive index of the aerosol measured through the application of white light scattering in conjunction with Mie spectroscopy e.g. Bohren and Huffman (1983). The values of the refractive index were subsequently used to estimate the change in the top of the atmosphere albedo from radiative transfer calculations that modelled an aqueous aerosol with a thin film of the atmospheric aerosol film material.

Deleted:



Determining the refractive index of atmospheric aerosol is paramount to understanding the light scattering and absorption properties of atmospheric aerosol, and hence its contribution to global temperatures. Considerable work has focused on determining the real and imaginary component of the refractive indices of atmospheric aerosols as demonstrated by the work carried out by Liu (2013), Wex (2009), Barkey (2007), Meskhidze (2013), Nakayama (2010) and Lang-Yona (2010) to list a few. [The use of morphological dependent resonances in Raman Spectra to determine refractive index at a fixed wavelength has been reported by Lin et al. \(1990\) and references therein and Miles et al. \(2012\).](#) The absorbing properties of atmospheric aerosols have been extensively studied e.g. Zhao (2013) studied the optical extinction properties of aerosols, Washenfelder (2013) studied aerosol extinction in the ultra-violet region and Liu (2016) investigated the absorbing properties of brown carbon. In the study presented, wavelength dependent real refractive indices were determined for common aerosol types such as organic atmospheric aerosol sourced from urban and remote locations and from wood smoke aerosol. In addition, the wavelength-dependent refractive index of the proxy atmospheric aerosol humic acid was determined.

Application of the optical trapping technique was successfully employed to determine the refractive index of aerosol over a wide wavelength range. The technique allowed the refractive index to be resolved to within 0.015 over a large wavelength range of 460 to 700 nm e.g. David (2016), Jones (2013) and Jones (2015). Previous studies have determined the refractive index of atmospheric aerosols either over a narrower or monochromatic wavelength range. Flores (2014) determined the refractive index for secondary organic aerosol over the wavelength range of 360 to 420 nm, whilst Lambe (2013), Guyon (2003) and Nakayama (2015) studied the refractive index of aerosols at individual wavelengths.

The technique also allowed the imaginary component of the refractive index and absorption Ångström exponent to be calculated for the wood smoke aerosol extracts and aqueous humic acid aerosol owing to the samples absorbing at smaller wavelengths. The absorption Ångström exponent describes the wavelength dependence of the absorption of light by aerosols e.g. Moosmüller (2011). The absorption Ångström exponent of atmospheric aerosols has been shown to be sensitive to wavelength e.g. Chakrabarty (2010) and Lewis (2008), chemical composition e.g. Ajtai (2011), Chakrabarty (2010), Flowers (2010), Park and Yu (2016), Russell (2009) and Sandradewi (2008), morphology e.g. Liu (2008) and Utry (2014) and size e.g. Utry (2014) and Gyawali (2009).

Deleted: to an unparalleled level of accuracy

Deleted: -

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

## 25 2. Materials and Methods

To optically trap aerosol extracts, aerosol was collected from the atmosphere by pulling air through a pre-combusted quartz filter using an air pump. The aerosol material was then extracted from the filter and transferred to isopropanol from which airborne aerosols could be generated by ultrasonic nebulization. Additionally, a commercial sample of humic acid in an aqueous solution was studied as an aerosol. The airborne aerosols were optically trapped and illuminated with white light, the backscattered white light was collected to generate a Mie spectrum of scattered light intensity as a function of wavelength. From the Mie spectrum the wavelength dependent refractive index and radius could be determined by replication with Mie calculations.

## 2.1. Collection of aerosol extracts

Three types of atmospheric aerosol were sampled to demonstrate the technique. Three samples were chosen to represent (a) a “dirty” sample from an urban environment, (b) a “clean” sample from a remote location and (c) samples with strong light absorbing properties. Wood smoke aerosol extract and aqueous humic acid aerosol were chosen for this purpose.

5 The urban sample was collected from the campus of Royal Holloway, University of London. The proximity of Central London (30 km), major motorways (M25, M40 and M4) and the large international airport Heathrow (8 km) mean the sample has been categorized as urban for the purposes of the study. The urban atmospheric aerosol extracts were collected over 30-day periods and combined in the extraction process to allow seasonal (spring, summer, autumn and winter) analysis. Combining the urban aerosol extracts into seasons ensured enough material was present to create airborne aerosols and to  
10 optically trap in a relatively inefficient process as described in Sect. 2.2.

The remote atmospheric aerosol extract was collected at the Halley Clean Air Sector Laboratory operated by the British Antarctic Survey (Jones (2008)). The sample was collected over the Antarctic summer of 2015 for 60 consecutive days. Antarctica is situated far from human populated areas, and therefore the sample has been called remote aerosol for the purpose of the study.

15 The wood smoke was collected over a six-hour time period from the smoke plume of a flaming log fire burning in a domestic wood burner. The firewood used in the wood burner was sourced from wild cherry trees. Two wood smoke samples were collected from two separate fires, and were labelled extract A and extract B.

To determine the contamination for each sample, corresponding analytical filter blanks were collected. To ensure all possible contamination sources were accounted for, the remote and wood smoke analytical blanks travelled to the field sites  
20 and back with the collected sample.

The urban and wood smoke aerosols were sampled by using an air pump with a flow of 30 L min<sup>-1</sup> through clean stainless steel pipelines into a PFA (perfluoroalkoxy) Savillex filter holder, whilst remote aerosols were sampled from Antarctic ambient air by using a short length of quarter inch O.D. PFA tubing at a flow rate of 20 L min<sup>-1</sup> onto a filter holder using a Staplex low volume air sampler (Model VM-4). All filter holders contained pre-combusted quartz filters (SKC Ltd.)  
25 with a diameter of 47 mm. The air was pulled through the filter for a known time period, after which the sample and filter were frozen in the dark at -18 °C until the sample could be extracted from the filter (typically days).

After sample collection, the filter holder was disassembled in a clean glove box to prevent contamination. The filter encased in the filter holder was cut into two on clean PFA blocks with a stainless steel blade: one half was for analysis, the other for reference. The filter was placed in a glass conical flask with 10 ml of chloroform (Sigma-Aldrich, 0.5 to 1 % EtOH as stabilizer) and 10 ml of ultrapure water (> 18 MΩm<sup>-1</sup>), and the mixture sonicated for 10 minutes. After sonication, the  
30 mixture was filtered through pre-combusted quartz filters to remove filter debris. The filter debris with un-extracted material was discarded. Sequentially, the filtrate was poured into a glass separating funnel and the chloroform layer drawn off. The atmospheric aerosol extract studied was soluble in chloroform. The chloroform was removed from the atmospheric aerosol

Deleted: paper

extract by evaporation under dry nitrogen. Once all the chloroform had been removed, 2 ml of isopropanol (Sigma-Aldrich, purity  $\geq 99.8\%$ ) was added to the sample and the sample was stored in the dark at  $-18\text{ }^{\circ}\text{C}$  until use. All instrumentation used in the collection and extraction of the atmospheric aerosol extract was cleaned with ultrapure water and chloroform before use.

The sonication in the extraction process was not found to change the Langmuir-Isotherm of atmospheric material at the air-water interface.

Every sample extracted from the atmosphere had a corresponding analytical blank to check for potential contamination of the filters. The analytical blanks for each sample were extracted following the same procedure above.

In addition to atmospheric aerosol extracts, aqueous humic acid aerosol (humic acid sodium salts, STBD5313V Sigma-Aldrich) was studied. the nebulised aqueous humic acid aerosol sample was prepared at a concentration of  $0.0005\text{ g cm}^{-3}$ .

## 2.2. Optical trapping of the aerosols

A vertically aligned, counter propagating optical trap was used to optically catch and levitate the aerosols. The optical trapping description is described in full by Jones (2013), however a brief description will be given here. The optical trap consisted of two laser beams that were fibre coupled from a 1064 nm continuous wave Nd:YAG laser (Laser Quantum). The laser beams passed through beam expansion optics to two vertically opposed microscope objectives (Mitutoyo M Plan Apo  $50\times\text{ NA }0.42$ ). An aluminium sample cell (volume  $38.4\text{ cm}^3$ ) was placed between the two microscope objectives and used to contain the aerosols. Borosilicate coverslip windows allowed the focused laser beams to pass into the sample cell and form the optical trap. The aerosol could be held for more than 24 hours once trapped e.g. Rkiouak (2014). The relative humidity and the temperature of the trapping environment were held at ambient conditions (30 percent relative humidity and  $20\text{ }^{\circ}\text{C}$ ).

A nebulizer (ultrasonic nebulizer, Omron) was used to create aerosols from either the atmospheric aerosol extract in isopropanol or aqueous humic acid aerosol. Inlet and exhaust ports allowed aerosol delivery into the sample cell. The solvent for the atmospheric aerosol extracts was exchanged from chloroform (removed by blowing down with nitrogen) to isopropanol prior to optically trapping as chloroform was found to be unsuitable for ultrasonic nebulisation. Isopropanol was added in the volume ratio 5:1 isopropanol to atmospheric extract, and this mixture was nebulised to deliver airborne droplets of the extract. Isopropanol evaporated from the aerosol both during transit and immediately upon capture in the optical trap.

## 2.3. Data analysis

The optically trapped aerosol droplet was illuminated with white light and the elastically backscattered light was collected over a 25 degree cone angle as a function of wavelength by an objective lens. Further optics described in Jones et al.(2013) focussed the light onto a spectrometer (Acton SP2500i). The resulting spectrum was a function of light intensity versus wavelength and will be called a Mie spectrum henceforth in the paper. The Mie spectrum covered the wavelength range 460- to 700 nm, with a resolution of 0.06 nm per pixel. The measured Mie spectrum was simulated through application of Bohren

Deleted: ) as a function of wavelength.

Deleted: ..

and Huffman(1983) formalism of Mie theory, integrating over a 25 degree cone angle of backscattered light, in conjunction with the Cauchy equation e.g. Bohren and Huffman (1983):

$$n = A + \frac{B}{\lambda^2} + \frac{C}{\lambda^4}, \quad (1)$$

to determine both size and a precise estimation of refractive index as a function of wavelength. The values in Eq. (1) represent the refractive index ( $n$ ), Cauchy empirical constants ( $A$ ,  $B$  and  $C$ ) and wavelength ( $\lambda$ ). The values of the three empirical constants and the radius of the trapped aerosol were iterated until a good comparison was achieved between the simulated and the experimentally obtained Mie spectrum. Typically, the radius of droplet was fixed and the values of A, B, and C varied until a good fit between measured and simulated Mie spectra was achieved by simple comparison (inspection) of peak, trough and inflection point positions. The value of the radius was then iterated through a series of radii with optimization of the values of A, B, and C as a function of wavelength repeated at each radius. Thus, a qualitative grid search was performed over parameter space. Parameter space was A varying from 1.3 to 1.7, B from 0 to 20,000 nm<sup>-2</sup> and C from 0 to 1×10<sup>9</sup> nm<sup>4</sup>. When simulating the Mie spectra of atmospheric aerosol extracts, small adjustment of the empirical constant, C, did not alter the simulated Mie spectrum noticeably and this value was therefore held at zero. An estimation of the uncertainty in the derived values of the refractive index and radius of the aerosol could be determined by variation of radius, A and B in turn and comparing the experimental and simulated spectra. The value of the radius was between 0 to 3 microns typically. The imaginary component of the refractive index was varied only after the grid search for the woodsmoke and humic acid samples.

Formatted: English (US)

#### 2.4. Ångström absorption coefficient

Deleted: Ångstrom

Samples including humic acid and those obtained from atmospheric sampling may have a measurable wavelength dependent absorption that can be defined by an Ångström exponent. In the context of Mie scattering, absorption is observed as a decrease in spectral intensity of the Mie spectrum. Inclusion of a wavelength dependent imaginary refractive index term in the Bohren and Huffman formalism can simulate the attenuation of intensity observed in the Mie spectra owing to absorption.

Deleted: Ångstrom

The absorption Ångström exponent was also determined by fitting an Ångström equation (Moosmüller, 2011) to the absorbance spectra of the atmospheric aerosol extract in isopropanol or humic acid in water obtained by using UV-VIS spectroscopy:

Deleted: variations

Deleted: As well as determining the Ångstrom exponent from the Mie spectrum, the

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: . . . . .

$$\frac{Abs}{Abs_0} = \left(\frac{\lambda}{\lambda_0}\right)^{-\alpha} \quad (2)$$

where  $Abs$  is the absorbance measured by a UV-VIS spectrometer (Perkin Elmer Lambda 950),  $\lambda$  is the wavelength and  $\alpha$  is the absorption Ångström exponent. It should be noted that  $Abs_0$  is the value of absorbance at the reference wavelength  $\lambda_0=460\text{nm}$ . The absorbance spectra of the bulk atmospheric aerosol extracts dissolved in isopropanol or the humic acid dissolved in water were recorded with a spectrometer covering the wavelength range 460 to 640 nm. Wood smoke aerosol extract and aqueous humic acid both demonstrated measureable absorption at low wavelengths, however the spectra from the other samples were below the instrument detection limits. The quoted photometric noise for the UV-Vis spectrometer was ~0.0002A. However, the urban and remote aerosol extracts were diluted in isopropanol to fill the UV-Vis spectrometer cuvette

Deleted: the

Deleted: ,

Deleted: Ångstrom

Formatted: Font:Not Italic

and thus a value three orders of magnitude larger than  $\sim 0.0002A$  may provide an upper bound for the absorbance of the samples reported below the detection limit.

In a UV-Vis spectrometer the absorption coefficient,  $\beta$ , can be related to the Absorbance, Abs, by

$$Abs = -\beta l \quad (3)$$

5 where  $l$  is the pathlength (1 cm for the work described here) and absorbance, Abs has been corrected from base 10 to base  $e$  (Petty 2006). The absorption coefficient can be related to the imaginary refractive index,  $k(\lambda)$ , by

$$k(\lambda) = \frac{\lambda \beta}{4\pi} \quad (4)$$

10 as described by Petty (2006). Substitution of equation (4) into equation (3) and subsequently equation (2) demonstrates that the Ångström relationship for absorbance (equation 2) is modified to

$$\frac{k}{k_0} = \left(\frac{\lambda}{\lambda_0}\right)^{-(\alpha-1)} \quad (5)$$

15 for describing the imaginary refractive index (see appendix). In essence the value of Ångström exponent,  $\alpha$ , measured by the UV-Vis spectrometer is larger than the corresponding value for the imaginary refractive index. Note for the work described here  $\lambda_0=460\text{nm}$ . The values of  $k_0$  and  $\alpha$  were measured for dilute solutions of the wood smoke extract in isopropanol and humic acid in water. In the optical trap the trapped droplet of wood smoke extract in isopropanol lost all of the isopropanol solvent to evaporation, as expected, leaving pure wood smoke extract. The aqueous humic acid solution lost some water to evaporation, but remained an aqueous and more concentrated, solution. As will be described below the mass density of the woodsmoke extract was measured independently. Thus, for the wood smoke droplet the values of  $k_0$  and  $Abs_0$  were corrected for the mass density of wood smoke extract in the optical trap and the attenuation of the resulting Mie spectrum will be shown to be consistent. For the aqueous humic acid solution, the value of  $k_0$  was determined by fitting the attenuation of the Mie spectrum by inspection, i.e. by changing the value of  $k_0$  until the intensity attenuation of the simulated and experimental Mie spectra matched, thereby calculating the value of the mass density in the trapped humic acid droplet. For the wood smoke aerosol extract, the mass density of the wood smoke aerosol extract in isopropanol was determined from a measurement on an Anton Parr densitometer, whilst the mass density of the pure material was calculated gravimetrically by evaporating isopropanol from a pre-weighed sample of the wood smoke aerosol extract. For the measurement of mass density obtained by the densitometer, a plot of the inverse of the density versus its corresponding weight fraction allowed the mass density to be determined. The Ångström coefficient determined for the Absorbance in isopropanol or water was adjusted for use with the imaginary refractive index Ångström relationship (see Appendix). Simulated Mie spectra of wood smoke aerosol and humic

Formatted: English (US)

Deleted: By substitution of Eq. (3) (Petty (2006)) into Eq. 2, it was observed that the absorption Ångstrom exponent is related to the imaginary component of the refractive index by  $\alpha-1$  (see Appendix A). ... [1]

Formatted: English (US)

Deleted: ( $\rho_1$ )

Deleted: measured

Deleted: ,  $\rho_2$ ,

acid aerosol were then calculated with and without an Ångström exponent absorption to demonstrate that the attenuation in Mie resonance intensity was consistent with absorption.

### 3. Results

Aqueous humic acid aerosol and extracts of atmospheric aerosols were optically trapped and the real component of the refractive index determined through comparison of experimentally obtained Mie spectra to simulated Mie spectra by varying the radius and the Cauchy coefficients  $A$  and  $B$ . The correct simulation of the Mie spectra requires the variation of absorption with wavelength to be described in terms of the imaginary refractive index:

$$\left(\frac{k}{k_0}\right) = \left(\frac{\lambda}{\lambda_0}\right)^{-(\alpha-1)}, \quad (4)$$

where  $k$  represents the imaginary refractive index,  $\lambda$  is the wavelength and  $\alpha$  is the absorption Ångström exponent. The Mie spectra for urban and remote atmospheric aerosol extracts are shown in Fig. 1, and the Mie spectra for the wood smoke aerosol extract and aqueous humic acid aerosol are shown in Fig. 2. Typically, small aerosols (with a radius of approximately 0.600  $\mu\text{m}$ ) were trapped when studying the atmospheric aerosol extracts and consequently only a few resonances were observed in the Mie spectra depicted in Figs. 1 and 2. Table 1 displays the Cauchy coefficients, refractive index and radius of the aerosols studied. In general the real component of the refractive index increases from remote to urban to wood smoke.

The mass density of the two pure wood smoke aerosol extracts was determined to be 1.47  $\text{g cm}^{-3}$  for extract A and 1.64  $\text{g cm}^{-3}$  for extract B. The values are similar to the values Hoffer *et al.* (2005) reported for humic like substances (HULIS) sourced from a biomass burning plume: 1.502 to 1.569  $\text{g cm}^{-3}$  and Dinar *et al.* (2008) reported 1.42-1.51  $\text{g cm}^{-3}$ . The mass density of the aqueous humic acid solution in the nebuliser was 0.0005  $\text{g cm}^{-3}$ , the mass density of the pure humic acid was reported from the supplier Sigma-Aldrich to be 1.52  $\text{g cm}^{-3}$ . The absorption spectrum and imaginary component of the refractive index over the wavelength range of 460 to 640 nm for the wood smoke and humic acid samples are shown in Fig. 3. Table 2 contains the values of the  $k_0$ ,  $\text{Abs}_0$ , and  $\alpha$  determined in the study presented here. The dependence of the Mie spectral intensity with and without contribution from the absorption are shown for the two absorbing samples in Fig. 2. Absorption attenuates the intensity of the Mie spectra and is most notable at shorter wavelengths where the mass absorption coefficient is largest.

Figure 2 demonstrates that it may be possible to determine the absorption spectra from Mie spectra recorded upon optically trapped aerosol. The solvent (isopropanol) was lost from the trapped aerosol extracts by experimental design to ensure the pure extract was studied. In contrast the aqueous humic acid droplet retained some of its water in equilibrium with the local humidity of the optical trapping cell. The trapped droplet was observed to lose water and consequently shrink in size. The simulated Mie spectra in Fig. 2, were calculated with and without absorption described by an Ångström exponent to demonstrate that the attenuation of Mie resonances (especially at shorter wavelengths) was consistent with measured Mie spectra. The mass density of humic acid in an aqueous droplet is proportional to the imaginary refractive index, allowing the mass density of the optically trapped humic acid droplet to be calculated. The concentration of the trapped humic acid was determined to be 0.016

Deleted: Ångstrom

Deleted: if

Deleted: (

Deleted: .

Deleted: aerosol

Deleted: whilst

Deleted: Ångstrom exponent

Deleted: two

Deleted: 3. The absorption Ångstrom exponent

Deleted: wood smoke aerosol extracts was calculated to be 6, whilst the imaginary component of the refractive index was

Deleted: to be 0.0085 at 440 nm. The absorption Ångstrom exponent of the aqueous humic acid solution was calculated to be 2.2 and the imaginary component of

Deleted: refractive index to be 0.0742 at 440 nm

Deleted: As shown

Formatted: English (US)

Deleted: .

Formatted: English (US)

Deleted: .

Formatted: English (US)

Deleted: imaginary refractive index

Formatted: English (US)

Deleted: varied until the simulated and

Formatted: English (US)

Deleted: matched

Deleted: concentration

Deleted: 0009

$\text{g cm}^{-3}$ . The concentration of the aqueous humic acid droplet had increased by a factor of ~32 upon trapping, thus demonstrating that water had evaporated from the droplet during the trapping and aerosol equilibration process.

Deleted: almost doubled

#### 4. Discussion

Studies reporting the refractive index of atmospheric aerosol extracts are predominantly conducted at individual wavelengths e.g. Guyon (2003), Hoffer (2005), Kim and Paulson (2013), Lambe (2013), Lang-Yona (2010), Nakayama (2013), Redemann (2000) and Stelson (1990). Yamasoe (1998) determined the real component of the refractive index of smoke aerosol extracts to be 1.53, 1.55, 1.59, and 1.58, for wavelengths of 438, 670, 870, and 1020 nm respectively, whilst Shingler (2016) conducted in situ aerosol particle measurements of wildfire, biogenic, marine and urban air masses and discovered a refractive index of 1.52 to 1.54 at a wavelength of 532 nm. Contrastingly, the refractive index in the study presented here was not calculated at a single wavelength but over a large continuous wavelength range (460 to 700 nm).

##### 4.1. Refractive index of atmospheric aerosol extracts

The variation in refractive index between the extracts investigated in the study indicates a distinctive difference between each sample from each location source. Figure 4 graphically compares the refractive index dispersion with wavelength obtained for the atmospheric aerosol extracts analysed in the study presented to selected values from literature. Large differences between the values of refractive index for remote, urban and wood smoke aerosol extracts can be easily observed in Fig. 4. Wood smoke aerosol extracts have the largest values of refractive index, followed by urban and then remote aerosol extracts. Antarctica is considered a clean environment owing to the physical remoteness of the continent and air that reaches Antarctica is considered relatively cleansed of anthropogenic particles (Wolff, 1990). However, some particles do reach Antarctica, examples of such aerosol sources include sea spray, the transport of industrial emissions (McConnell, 2014) and particulate material from biomass burning and tropical forest fires (Tomasi, 2007). Aerosols from such sources have travelled far and have likely undergone chemical ageing, and are thus likely to be very different in chemical composition than their initial composition. The review authored by Moise, Flores and Rudich (2015) demonstrates the importance of chemical reactions in the alteration of the optical properties of atmospheric aerosols during atmospheric transport.

The urban aerosol extract samples have a wide distribution of the values of refractive index with values ranging from 1.478 to 1.522 at 589 nm. From the small sample analysed in the study presented, the autumn and winter samples are at the larger end of the refractive index range, with spring and summer at the mid to low end. The wavelength dependent refractive index values determined for atmospheric aerosol extracts lie in good agreement with previous monochromatic literature results. Studies focusing on anthropogenic aerosols determined values of refractive index varying from 1.498 to 1.653 nm at 532 nm; the refractive index range found in literature encompasses the urban and wood smoke aerosol extract wavelength dependent refractive indices determined in the study e.g. Adler (2011), Yamasoe (1998), Hoffer (2005) and Shingler (2016). Similarly, studies focused on aerosols sourced from the Antarctica demonstrate refractive index values similar to the refractive index



values determined for the aerosol extract sourced from Antarctica e.g. Virkkula (2006) measured a refractive index that generally lay between 1.4 and 1.5, but also reached as low as 1.3 for Antarctic aerosol extracts (these values are an average of measurements from the wavelengths 450, 550 and 700 nm).

5 Contrastingly, Guyon (2003) studied the refractive index of biomass burning aerosols collected from the Amazon tropical forest and determined a refractive index of 1.41 at 545 nm (0.1812 lower than the wavelength dependent refractive index determined in the study presented here at the same wavelength). A likely reason for the variation is that Guyon collected samples from a station suspended 54 m above ground level (and 22 m above the forest canopy), whereas in the study presented here samples were drawn directly from the smoke plume from a burning fire and consequently samples did not have the opportunity to chemically age or to mix with other material present in the atmosphere. Refractive index values from literature 10 for secondary organic aerosols demonstrate a wide range of refractive index values, for example the work carried out by Kim (2010), Kim (2013), Lambe (2013), Lang-Yona (2010), Spindler (2007) and Yu (2008). The work indicates that a single refractive index value cannot be used to describe secondary organic aerosol, perhaps owing to the wide range of organics present in the aerosol e.g. Yu (2008).

#### 4.2. Refractive index of aqueous humic acid aerosol

15 Aqueous humic acid was studied to demonstrate the ability of using optical trapping and Mie spectroscopy to study a known absorbing aerosol. Humic acid was dissolved in water to form a solution prior to nebulisation. The measured refractive index of the aqueous aerosol droplet will have a different refractive index than a pure sample of humic acid owing to the water content of the aerosol. The refractive index of humic-like substances has been reported to be quite large, for example Hoffer (2005) reported a refractive index of 1.653 for HULIS samples. However in the study presented, the aqueous humic acid 20 droplets had a refractive index lower than any of the atmospheric aerosol extracts because the atmospheric aerosol extracts were trapped as concentrated aerosol extracts (the solvent, isopropanol, used to allow the aerosol extracts to become airborne evaporated during nebulisation and trapping).

#### 4.3. Calculation of the absorption Ångström exponent

25 Unlike the urban and remote atmospheric aerosol extracts, the wood smoke aerosol extract and aqueous humic acid aerosol had strong absorption properties as shown by UV-VIS experiments on samples dissolved in isopropanol and water respectively. The imaginary component of the refractive index and absorption Ångström exponent could then be determined for the two samples.

30 Previous studies in the field have determined values for the absorption Ångström exponents to range from 2 to 16 for carbonaceous aerosol over the visible wavelength range e.g. Hoffer (2005), Lewis (2008); Chakrabarty (2010), Flowers (2010), Moosmüller (2011), Utry (2013), Zhang (2013), He (2015), Garg (2016), Pokhrel (2016), Rathod (2016) and Shen (2017). More specifically, values in the range of 3.5 to 8.3 have been calculated for studies focusing on the absorption Ångström exponent for smoke aerosols. Lewis (2008) studied the combustion of a variety of fuels with a dual wavelength photo-acoustic

Deleted: Ångstrom

Deleted: bulk

Deleted: .

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

instrument to determine an absorption Ångström exponent of 3.5. Contrastingly, Hoffer (2005) and Park and Yu (2016) obtained a much higher absorption Ångström exponents of 6 to 7 and 7.4 to 8.3 respectively for biomass burning aerosols.

The calculated absorption Ångström exponent for the wood smoke aerosol extract correlates with the absorption Ångström exponent measured in previous studies for biomass burning aerosols. Interestingly, it has been suggested that fire type plays a role in the amount of black carbon produced. A flaming fire has been shown to produce more particles e.g. Reid (2005), and in particular produce more black carbon than smouldering fires e.g. Hoffer (2005) and Yamasoe (2000). The fire from which the wood smoke aerosol extracts were collected from was flaming and hence a high absorption Ångström exponent is more likely. It ought to be noted that the wood smoke extract was included in this work as an exploratory sample with a strong absorption and future work will explore smoke aerosol where the fuel and fire temperature are carefully controlled.

Considering the absorption Ångström exponent for the aqueous humic acid aerosol, it can be observed that the exponent correlates with studies that determined the absorption Ångström exponent for biomass fuels. For example, Schnaiter (2006) deduced an absorption Ångström exponent between 2.2 and 3.5 for aerosols produced from the combustion of propane and Schnaiter (2003) determined an absorption Ångström exponent of 1 for emissions produced from the combustion of diesel.

#### 4.4. Uncertainty in Mie spectra fitting

The collection technique applied to extract remote and urban aerosols from the atmosphere was limited by airflow and filter size and therefore sample was very limited. An air pump with a significantly larger flow rate was used, however the aerosol extracts that were collected contained contamination in their analytical blanks warranting the samples to not be used. The contamination was attributed to the quality of the filters used, demonstrating that pre-combusted filters are critical.

Owing to limited sample, only small aerosols were optically trapped causing the collected Mie spectra to have little structure. A less structured Mie spectrum reduces the accuracy of the determined wavelength dependent refractive index, radius and absorption Ångström exponent. The Mie spectra in figure 1 of Summer Urban aerosol extract are structured with pronounced peak shapes that allow the facile fitting between simulated and measured Mie Spectra. Such spectra allow a relatively small range of values of radius, Cauchy coefficients A, B and C to provide a good fit between measured and simulated Mie Spectra. The rest of the Mie spectra in Figure 1 have significantly fewer Mie resonances and their peak shapes are less pronounced. The uncertainties become larger as the spectra become less structured. Despite the limitations in Mie spectra simulation, the typical uncertainty in radius and refractive index for a Mie spectra was typically  $\pm 6$  nm and  $\pm 0.015$  respectively, whilst the uncertainty for the absorption Ångström exponent was 7 percent for the wood smoke aerosol extract and 5 percent for the aqueous humic acid aerosol.

The liquid droplets are assumed to be perfectly spherical. Mie scattering from droplets experiencing small deformation has been shown by Arnold et al.(1990) and Schweiger(1990) to result in resonances which shift, broaden and split as the droplet asymmetry increases.

The sensitivity of the simulated Mie spectra to the refractive index ( $\pm 0.015$ ) and radius ( $\pm 6$ nm) of the droplet are shown in figure 5. The simulated spectra, with the stated variations and the experimental Mie spectra for the Spring Urban

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom

Deleted: Ångstrom exponent.

Formatted: English (US)

Deleted: Ångstrom

aerosol extract are plotted. Figure 5 also contains a third simulated set of Mie spectra calculated by re-optimising the values of A, B, and C in Cauchy equation to achieve a fit between simulated and experimental Mie spectra for particles with a radius  $\pm 12$  nm from the optimum fit to the experimental data. Figure 5 demonstrates that the quoted uncertainties in radius ( $\pm 6$  nm) and refractive index ( $\pm 0.015$ ) are realistic.

5

#### 4.5. Atmospheric Implications

Atmospheric aerosol can increase the top of the atmosphere albedo by scattering incoming solar radiation and decrease the top of the atmosphere albedo by absorbing solar radiation. Using the refractive index data collected in the study presented, a radiative transfer model was applied to consider the change in top of the atmosphere albedo owing to an aerosol film forming with the same optical properties as the extracts studied in the presented study. The material extracted from the atmospheric samples described in the study may form an organic shell at the air-water interface of an aqueous aerosol e.g. Gill (1983). An atmospheric radiative-transfer model (Stamnes, 1988) was applied to study an atmospheric aerosol layer consisting of core aqueous aerosol coated in an organic shell with the optical properties of the atmospheric aerosol extract measured within the work presented here. The change in the top of the atmospheric albedo was calculated as the proportions of water and organic material were varied for different size aerosols.

The top of the atmosphere albedo was calculated for an aerosol layer with the composition of an aqueous core aerosol surrounded by a shell of either urban, wood smoke or remote atmospheric aerosol extract with the volume fraction varying from 0 to 1 (i.e. pure water to pure organic). The calculations represent only a small-scale study to simply identify potential effects on the top of the atmosphere albedo owing to the presence of pure core-shell particles in the atmosphere versus no aerosol present. The change in the top of the atmosphere albedo is reported as an aerosol relative effect, following the approach of Mishra (2015).

The atmospheric radiative-transfer model uses the DISORT code (Stamnes, 1988). The model uses values of the scattering, absorption and the asymmetry parameter of aerosols to calculate the change in solar radiation through the atmosphere. To calculate the scattering and absorption parameters for coated spheres Mie calculations were performed for the core-shell particles using BHCOAT, a code developed by Bohren and Huffman (1983), which was later modified to also include a calculation of the asymmetry parameter. Scattering, absorption and asymmetry parameter for the particle are calculated from the refractive index of the core and shell. For all aerosol particles the refractive index of the core is a wavelength dependent value for water (IAPWS, 1997) and the refractive index of the surrounding medium a wavelength independent value for air of 1.00-0.0i. The shell of the aerosol has a wavelength dependent refractive index of the urban, remote or wood smoke aerosol extracts, as displayed in Table 1. In addition, the absorption properties of the wood smoke aerosol extract were included in the calculation.

The core-shell Mie calculation was used to obtain scattering and absorption cross-sections and asymmetry parameters for particles with a radius of 100 to 10,000 nm (in 100 nm intervals from 100 to 1000 nm, and 1000 nm intervals from 1000

to 10,000 nm), and with the shell volume making up 0.01 to 0.99 of the whole particle volume. Calculations were performed over wavelengths covering 350 to 750 nm. The ground albedo was set to 0.1. Aerosol of one size was placed in three consecutive 1 km thick layers at the surface, forming a 3 km thick aerosol layer. The aerosol optical depth for each of these layers was set to 0.126, the global average set by Mao (2014), and no aerosol or cloud was placed in any subsequent layers.

The solar zenith angle was set at 60°. The albedo of the top of the atmosphere was calculated as the ratio of incoming to outgoing irradiance at 100km altitude and averaged over wavelengths from 350 to 750 nm for each particle size. Calculations were also performed for no aerosol present in the atmosphere. The aerosol radiative effect was then calculated using Eq. (5).

$$ARE_{TOT} = TOA\ albedo_{aerosol} - TOA\ albedo_{no\ aerosol} \quad (5)$$

where  $ARE_{TOT}$  stands for total aerosol radiative effect,  $TOA\ albedo_{aerosol}$  stands for the top of the atmosphere albedo with aerosol present and  $TOA\ albedo_{no\ aerosol}$  stands for the top of atmosphere albedo without aerosol present.

The results of these calculations are presented in Fig. 6. From Fig. 6, it can be observed that all aerosols have a positive effect on the total aerosol radiative effect, with the most positive effect observed for particle sizes of 600 and 800 nm for the urban and remote atmospheric aerosol extracts and 200 and 400 nm for the wood smoke aerosol extract. Note the change in top of the atmosphere albedo is most pronounced when the volume of shell increases from 0 to 25 percent of the total volume.

## 5. Conclusions

A new technique employing optical trapping techniques applied alongside Mie spectroscopy was employed to determine the real and imaginary components of the refractive index of insoluble organic material from atmospheric aerosol extracts over a wide wavelength range. The atmospheric aerosol extract was successfully trapped, demonstrating that the material forms spherical liquid droplets onto which Mie theory could be applied. From application of Mie theory, the refractive indices of the atmospheric aerosol extract were determined to vary from 1.470 for aerosol extracted from Antarctica to 1.588 for wood smoke aerosol extracts at the wavelength 589 nm, whilst seasonal refractive index dependence was observed for atmospheric aerosol extracted from an urban environment.

Additionally, owing to the efficient light absorbing nature of the wood smoke aerosol extract and aqueous humic acid extract, the absorption Ångström exponent could be determined with a high level of certainty for the extract; through applying optical trapping and Mie spectroscopy alongside UV-VIS spectroscopy it was possible to determine the real and imaginary component of the refractive index simultaneously.

The aerosol collected may exist as an aerosol in the atmosphere or may coagulate with other atmospheric aerosols to form a film. Use of a simple one dimensional radiative-transfer model to study a shell of atmospheric aerosol extracts on an aqueous aerosol indicates that the albedo of the top of the atmosphere albedo may change by 0.03 relative to a pure aqueous droplet.

Deleted: 5

Deleted: 5

Deleted: Ångström



## Competing Interests

The authors declare that they have no conflict of interest.

## Acknowledgments

The authors would like to thank the Central Laser Facility for granting access time on the optical trapping equipment at Rutherford Appleton Laboratories, Oxfordshire under the grant number ISI002. Rosalie H. Shepherd would like to thank STFC for funding the student grant STL504279\1.

## References

- Adler, G., Flores, J. M., Riziq, A. A., Borrmann, S. and Rudich, Y. (2011) 'Chemical, physical, and optical evolution of biomass burning aerosols : a case study', *Atmospheric Chemistry and Physics*, 11, pp. 1491–1503. doi: 10.5194/acp-11-1491-2011.
- Ajtai, T., Filep, A., Utry, N., Schnaiter, M., Linke, C., Bozoki, Z., Szabo, G. and Leisner, T. (2011) 'Inter-comparison of optical absorption coefficients of atmospheric aerosols determined by a multi-wavelength photoacoustic spectrometer and an Aethalometer under sub-urban wintry conditions', *Journal of Aerosol Science*, 42(12), pp. 859–866. doi: 10.1016/j.jaerosci.2011.07.008.
- Andreaea, M. O. and Rosenfeld, D. (2008) 'Aerosol–cloud–precipitation interactions. Part 1. The nature and sources of cloud-active aerosols', *Earth Science Reviews*, 89(1–2), pp. 13–41. doi: 10.1016/j.earscirev.2008.03.001.
- [Arnold, A., Spock, D.E., Folan, L.M., \(1990\) 'Electric-field-modulated light scattering near a morphological resonance of a trapped aerosol particle', \*Optics Letters\*, 15\(20\) pp: 1111-1113 doi: 10.1364/OL.15.001111](#)
- Barkey, B., Paulson, S. E. and Chung, A. (2007) 'Genetic Algorithm Inversion of Dual Polarization Polar Nephelometer Data to Determine Aerosol Refractive Index', *Aerosol Science and Technology*, 41(8), pp. 751–760. doi: 10.1080/02786820701432640.
- Blanchard, D. C. (1964) 'Material, Sea-to-Air Transport of Surface Active', *Science*, 146, pp. 396–397. doi: 10.1126/science.146.3642.396.
- [Bligh, E.G. and Dyer, W.J. \(1959\) 'A rapid method of total lipid extraction and purification' \*Canadian Journal of Biochemistry and Physiology\*, 37, pp497-509](#)
- Bohren, C. F. and Huffman, D. R. (1983) 'Absorption and Scattering of Light by Small Particles', Wiley Scientific.
- Breon, F. M., Tanfè, D. and Generoso, S. (2002) 'Aerosol Effect on Cloud Droplet Size Monitored from Satellite', *Science*, 295, pp. 834–839. doi: 10.1126/science.1066434.
- Cai, C., Miles, R. E. H., Cotterell, M. I., Marsh, A., Rovelli, G., Rickards, A. M. J., Zhang, Y. H. and Reid, J. P. (2016) 'Comparison of methods for predicting the compositional dependence of the density and refractive index of organic-aqueous

- aerosols', *Journal of Physical Chemistry A*, 120(33), pp. 6604–6617. doi: 10.1021/acs.jpca.6b05986.
- Cappa, C. D., Che, D. L., Kessler, S. H., Kroll, J. H. and Wilson, K. R. (2011) 'Variations in organic aerosol optical and hygroscopic properties upon heterogeneous OH oxidation', *Journal of Geophysical Research: Atmospheres*, 116(15), pp. 1–12. doi: 10.1029/2011JD015918.
- 5 Chakrabarty, R. K., Moosmüller, H., Chen, L. W. A., Lewis, K., Arnott, W. P., Mazzoleni, C., Dubey, M. K., Wold, C. E., Hao, W. M. and Kreidenweis, S. M. (2010) 'Brown carbon in tar balls from smoldering biomass combustion', *Atmospheric Chemistry and Physics*, 10(13), pp. 6363–6370. doi: 10.5194/acp-10-6363-2010.
- Charlson, R. J., Seinfeld, J. H., Nees, A., Kulmala, M., Laaksonen, A. and Facchini, M. C. (2001) 'Reshaping the theory of cloud formation', *Science*, pp. 2025–2026. doi: 10.1126/science.1060096.
- 10 Cochran, R. E., Laskina, O., Jayarathne, T., Laskin, A., Laskin, J., Lin, P., Sultana, C., Lee, C., Moore, K. A., Cappa, C. D., Bertram, T. H., Prather, K. A., Grassian, V. H. and Stone, E. A. (2016) 'Analysis of Organic Anionic Surfactants in Fine and Coarse Fractions of Freshly Emitted Sea Spray Aerosol', *Environmental Science and Technology*, 50(5), pp. 2477–2486. doi: 10.1021/acs.est.5b04053.
- David, G., Esat, K., Ritsch, I. and Signorell, R. (2016) 'Ultraviolet broadband light scattering for optically-trapped submicron-sized aerosol particles', *Physical chemistry chemical physics*, pp. 5477–5485. doi: 10.1039/C5CP06940H.
- 15 Davies, J. F., Miles, R. E. H., Haddrell, A. E. and Reid, J. P. (2013) 'Influence of organic films on the evaporation and condensation of water in aerosol', *Proceedings of the National Academy of Sciences of the United States of America*, 110(22), pp. 8807–12. doi: 10.1073/pnas.1305277110.
- Dinar, E., Riziq, A. A., Spindler, C., Erlick, C., Kiss, G. and Rudich, Y. (2008) 'The complex refractive index of atmospheric and model humic-like substances ( HULIS ) retrieved by a cavity ring down aerosol spectrometer (CRD-AS)', *Faraday Discussions*, 137, pp. 137, doi: 10.1039/B703111D.
- 20 Donaldson, D. J. and Anderson, D. (1999) 'Adsorption of atmospheric gases at the air-water interface. 2. C<sub>1</sub>-C<sub>4</sub> alcohols, acids, and acetone', *J. Phys. Chem. A*, 103(7), pp. 871–876. doi: 10.1021/jp983963h.
- Donaldson, D. J. and Vaida, V. (2006) 'The influence of organic films at the air-aqueous boundary on atmospheric processes', *Chemical Reviews*, 106(4), pp. 1445–1461. doi: 10.1021/cr040367c.
- 25 Donaldson, D. J. and Valsaraj, K. T. (2010) 'Adsorption and reaction of trace gas-phase organic compounds on atmospheric water film surfaces: A critical review', *Environmental Science and Technology*, 44(3), pp. 865–873. doi: 10.1021/es902720s.
- Eliason, T. L., Aloisio, S., Donaldson, D. J., Cziczo, D. J. and Vaida, V. (2003) 'Processing of unsaturated organic acid films and aerosols by ozone', *Atmospheric Environment*, 37(16), pp. 2207–2219. doi: 10.1016/S1352-2310(03)00149-3.
- 30 Enami, S., Hoffmann, M. R. and Colussi, A. J. (2010) 'Molecular control of reactive gas uptake "on Water"', *Journal of Physical Chemistry A*, 114(18), pp. 5817–5822. doi: 10.1021/jp1019729.
- Enami, S. and Sakamoto, Y. (2016) 'OH-Radical Oxidation of Surface-Active cis-Pinonic Acid at the Air-Water Interface', *Journal of Physical Chemistry A*, 120(20), pp. 3578–3587. doi: 10.1021/acs.jpca.6b01261.
- Feingold, G. and Chuang, P. Y. (2002) 'Analysis of the Influence of Film-Forming Compounds on Droplet Growth:

Deleted: .



- Implications for Cloud Microphysical Processes and Climate', *Journal of the Atmospheric Sciences*, 59(12), pp. 2006–2018. doi: 10.1175/1520-0469.
- Flores, J. M., Washenfelder, R. A., Adler, G., Lee, H. J., Segev, L., Laskin, J., Laskin, A., Nizkorodov, S. A., Brown, S. S. and Rudich, Y. (2014) 'Complex refractive indices in the near-ultraviolet spectral region of biogenic secondary organic aerosol aged with ammonia.', *Physical chemistry chemical physics*, 16(22), pp. 10629–42. doi: 10.1039/c4cp01009d.
- Flores, J. M., Zhao, D. F., Segev, L., Schlag, P., Kiendler-Scharr, A., Fuchs, H., Watne, A. K., Bluvshstein, N., Mentel, T. F., Hallquist, M. and Rudich, Y. (2014) 'Evolution of the complex refractive index in the UV spectral region in ageing secondary organic aerosol', *Atmospheric Chemistry and Physics*, 14(11), pp. 5793–5806. doi: 10.5194/acp-14-5793-2014.
- Flowers, B. A., Dubey, M. K., Mazzoleni, C., Stone, E. A., Schauer, J. J., Kim, S. W. and Yoon, S. C. (2010) 'Optical-chemical-microphysical relationships and closure studies for mixed carbonaceous aerosols observed at Jeju Island; 3-laser photoacoustic spectrometer, particle sizing, and filter analysis', *Atmospheric Chemistry and Physics*, 10(21), pp. 10387–10398. doi: 10.5194/acp-10-10387-2010.
- Folch, J., Lees, M., Sloane Stanley, G.H., (1957) 'A simple method for the isolation and purification of total lipides ffrom animal tissues' *Journal of Biological Chemistry*, 226, pp 497-509.
- Fuzzi, S., Andreae, M. O., Huebert, B. J., Kulmala, M., Bond, T. C., Boy, M., Doherty, S. J., Guenther, A., Kanakidou, M., Kawamura, K., Kerminen, V. M., Lohmann, U., Russell, L. M. and Pöschl, U. (2005) 'Critical assessment of the current state of scientific knowledge, terminology, and research needs concerning the role of organic aerosols in the atmosphere, climate, and global change', *Atmospheric Chemistry and Physics Discussions*, 5(6), pp. 11729–11780. doi: 10.5194/acpd-5-11729-2005.
- Garg, S., Chandra, B. P., Sinha, V., Sarda-Esteve, R., Gros, V. and Sinha, B. (2016) 'Limitation of the Use of the Absorption Ångström Exponent for Source Apportionment of Equivalent Black Carbon: A Case Study from the North West Indo-Gangetic Plain', *Environmental Science and Technology*, 50(2), pp. 814–824. doi: 10.1021/acs.est.5b03868.
- Gill, P. S., Graedel, T. E. and Weschler, C. J. (1983) 'Organic films on atmospheric aerosol particles, fog droplets, cloud droplets, raindrops, and snowflakes', *Reviews of Geophysics*, 21(4), p. 903. doi: 10.1029/RG021i004p00903.
- Guyon, P., Boucher, O., Graham, B., Beck, J., Mayol-Bracero, O. L., Roberts, G. C., Maenhaut, W., Artaxo, P. and Andreae, M. O. (2003) 'Refractive index of aerosol particles over the Amazon tropical forest during LBA-EUSTACH 1999', *Journal of Aerosol Science*, 34(7), pp. 883–907. doi: 10.1016/S0021-8502(03)00052-1.
- Gyawali, M., Arnott, W. P., Lewis, K. and Moosmüller, H. (2009) 'In situ aerosol optics in Reno, NV, USA during and after the summer 2008 California wildfires and the influence of aerosol coatings', *Atmospheric Chemistry and Physics Discussions*, 9, pp. 8007–8015. doi: 10.5194/acp-9-8007-2009.
- He, Q., Zhao, X., Lu, J., Zhou, G., Yang, H., Gao, W., Yu, W. and Cheng, T. (2015) 'Impacts of biomass-burning on aerosol properties of a severe haze event over Shanghai', *Particuology. Chinese Society of Particuology*, 20, pp. 52–60. doi: 10.1016/j.partic.2014.11.004.
- Hoffer, A., Gelencsér, A., Guyon, P., Kiss, G., Schmid, O., Frank, G. P., Artaxo, P. and Andreae, M. O. (2005) 'Optical

Deleted: Angstrom

- properties of humic-like substances (HULIS) in biomass-burning aerosols', *Atmospheric Chemistry and Physics Discussions*, 5(4), pp. 7341–7360. doi: 10.5194/acpd-5-7341-2005.
- Hoffer, A., Tóth, A., Nyirő-Kósa, I., Pósfai, M. and Gelencsér, A. (2016) 'Light absorption properties of laboratory-generated tar ball particles', *Atmospheric Chemistry and Physics*, 16(1), pp. 239–246. doi: 10.5194/acp-16-239-2016.
- 5 IAPWS (1997) 'Release on the Refractive Index of Ordinary Water Substance as a Function of Wavelength, Temperature and Pressure'.
- Jacobson, M. C. and Hansson, H. (2000) 'Organic atmospheric aerosols: Review and state of the science', *Reviews of Geophysics*, (38), pp. 267–294. doi: 10.1029/1998RG000045.
- Jauffred, L., Taheri, S. M. R., Schmitt, R., Linke, H. and Oddershede, L. B. (2015) 'Optical Trapping of Gold Nanoparticles in Air', *Nano Letters*, 15(7), pp. 4713–4719. doi: 10.1021/acs.nanolett.5b01562.
- 10 Jones, A. E., Wolff, E. W., Salmon, R. A., Bauguutte, S. J. B., Roscoe, H. K., Anderson, P. S., Ames, D., Clemitshaw, K. C., Fleming, Z. L., Bloss, W. J., Heard, D. E., Lee, J. D., Read, K. A., Hamer, P., Shallcross, D. E., Jackson, A. V., Walker, S. L., Lewis, A. C., Mills, G. P., Plane, J. M. C., Saiz-Lopez, A., Sturges, W. T. and Worton, D. R. (2008) 'Chemistry of the Antarctic Boundary Layer and the Interface with Snow: an overview of the CHABLIS campaign', *Atmospheric Chemistry and Physics*, 8(14), pp. 3789–3803. doi: 10.5194/acp-8-3789-2008.
- 15 Jones, S. H., King, M. D. and Ward, A. D. (2013) 'Determining the unique refractive index properties of solid polystyrene aerosol using broadband Mie scattering from optically trapped beads.', *Physical chemistry chemical physics*, 15(47), pp. 20735–41. doi: 10.1039/c3cp53498g.
- Jones, S. H., King, M. D. and Ward, A. D. (2015) 'Atmospherically relevant core-shell aerosol studied using optical trapping and Mie scattering', *Chem. Commun., Royal Society of Chemistry*, 51(23), pp. 4914–4917. doi: 10.1039/C4CC09835H.
- 20 Jones, S. H., King, M. D., Ward, A. D., Rennie, A. R., Jones, A. C. and Arnold, T. (2017) 'Are organic films from atmospheric aerosol and sea water inert to oxidation by ozone at the air-water interface', *Atmospheric Environment*, 161, pp. 274-287, doi: 10.1016/j.atmosenv.2017.04.025.
- Kaiser, T., Roll, G. and Schweiger, G. (1996) 'Investigation of coated droplets in an optical trap: Raman-scattering, elastic-light-scattering, and evaporation characteristics.', *Applied optics*, 35(30), pp. 5918–5924. doi: 10.1364/AO.35.005918.
- 25 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G. and Wilson, J. (2005) 'Organic aerosol and global climate modelling: a review', *Atmospheric Chemistry and Physics*, 5, pp. 1053–1123. doi: 10.5194/acp-5-1053-2005.
- 30 Kim, H., Barkey, B. and Paulson, S. E. (2010) 'Real refractive indices of alpha and beta pinene and toluene secondary organic aerosols generated from ozonolysis and photo-oxidation', *Journal of Geophysical Research Atmospheres*, 115(23). doi: 10.1029/2010JD014549.
- Kim, H. and Paulson, S. E. (2013) 'Real refractive indices and volatility of secondary organic aerosol generated from photooxidation and ozonolysis of limonene,  $\alpha$ -pinene and toluene', *Atmospheric Chemistry and Physics*, 13(15), pp. 7711–

7723. doi: 10.5194/acp-13-7711-2013.

King, M.D., Thompson, K.C. and Ward, A.D., (2004) 'Laser Tweezers Raman Study of Optically Trapped Aerosol Droplets of Seawater and Oleic Acid Reacting with Ozone: Implications for Cloud-Droplet Properties' *Journal of the American Chemical Society*, 126 (51), 16710-16711. doi: 10.1021/ja044717o

- 5 King, M. D., Rennie, A. R., Thompson, K. C., Fisher, F. N., Dong, C. C., Thomas, R. K., Pfrang, C. and Hughes, A. V (2009) 'Oxidation of oleic acid at the air-water interface and its potential effects on cloud critical supersaturations.', *Physical chemistry chemical physics*. 11(35), pp. 7699–7707. doi: 10.1039/b906517b.
- Lambe, A. T., Cappa, C. D., Massoli, P., Onasch, T. B., Forestieri, S. D., Martin, A. T., Cummings, M. J., Croasdale, D. R., Brune, W. H., Worsnop, D. R. and Davidovits, P. (2013) 'Relationship between oxidation level and optical properties of secondary organic aerosol', *Environmental Science and Technology*, 47(12), pp. 6349–6357. doi: 10.1021/es401043j.
- 10 Lang-Yona, N., Abo-Riziq, A., Erlick, C., Segre, E., Trainic, M. and Rudich, Y. (2010) 'Interaction of internally mixed aerosols with light.', *Physical chemistry chemical physics*. 12(1), pp. 21–31. doi: 10.1039/b913176k.
- Lang-Yona, N., Rudich, Y., Mentel, T. F., Bohne, A., Buchholz, A., Kiendler-Scharr, A., Kleist, E., Spindler, C., Tillmann, R. and Wildt, J. (2010) 'The chemical and microphysical properties of secondary organic aerosols from Holm Oak emissions',
- 15 *Atmospheric Chemistry and Physics*, 10(15), pp. 7253–7265. doi: 10.5194/acp-10-7253-2010.
- Lewis, K., Arnott, W. P., Moosmuller, H. and Wold, C. E. (2008) 'Strong spectral variation of biomass smoke light absorption and single scattering albedo observed with a novel dual-wavelength photoacoustic instrument', *Journal of Geophysical Research Atmospheres*, 113(16), pp. 1–14. doi: 10.1029/2007JD009699.
- Li, Y., Ezell, M. J. and Finlayson-Pitts, B. J. (2011) 'The impact of organic coatings on light scattering by sodium chloride particles', *Atmospheric Environment*, 45(25), pp. 4123–4132. doi: 10.1016/j.atmosenv.2011.05.031.
- 20 Lin, H.-B., Eversole, J.D., Campillo, A.J. (1990) 'Identification of morphology dependent resonances in stimulated Raman scattering from microdroplets', *Optics communications*, 77(5,6), pp407-410
- Liu, J., Lin, P., Laskin, A., Laskin, J., Kathmann, S. M., Wise, M., Caylor, R., Imholt, F., Selimovic, V. and Shilling, J. E. (2016) 'Optical properties and aging of light-absorbing secondary organic aerosol', *Atmospheric Chemistry and Physics*,
- 25 16(19), pp. 12815–12827. doi: 10.5194/acp-16-12815-2016.
- Liu, L., Mishchenko, M. I. and Patrick Arnott, W. (2008) 'A study of radiative properties of fractal soot aggregates using the superposition T-matrix method', *Journal of Quantitative Spectroscopy and Radiative Transfer*, 109(15), pp. 2656–2663. doi: 10.1016/j.jqsrt.2008.05.001.
- Liu, P., Zhang, Y. and Martin, S. T. (2013) 'Complex refractive indices of thin films of secondary organic materials by spectroscopic ellipsometry from 220 to 1200 nm', *Environmental Science and Technology*, 47(23), pp. 13594–13601. doi: 10.1021/es403411e.
- Lohmann, U. and Feichter, J. (2005) 'Global indirect aerosol effects : a review', *Atmospheric Chemistry and Physics*, 5, pp. 715–737. doi: 1680-7324/acp/2005-5-715.
- Mao, K., Ma, Y., Xia, L., Chen, W., Shen, X., He, T. and Xu, T. (2014) 'Global aerosol change in the last decade: An analysis

- based on MODIS data', *Atmospheric Environment*, 94, pp. 680–686. doi: 10.1016/j.atmosenv.2014.04.053.
- McConnell, J. R., Maselli, O. J., Sigl, M., Vallelonga, P., Neumann, T., Anschutz, H., Bales, R. C., Curran, M. A. J., Das, S. B., Edwards, R., Kipfstuhl, S., Layman, L. and Thomas, E. R. (2014) 'Antarctic-wide array of high-resolution ice core records reveals pervasive lead pollution began in 1889 and persists today.', *Scientific reports*, 4, p. 5848. doi: 10.1038/srep05848.
- 5 McFiggans, G., Artaxo, P., Baltensperger, U., Coe, H., Facchini, M. C., Feingold, G., Fuzzi, S., Gysel, M., Laaksonen, A., Lohmann, U., Mentel, T. F., Murphy, D. M., O'Dowd, C. D., Snider, J. R. and Weingartner, E. (2005) 'The effect of physical and chemical aerosol properties on warm cloud droplet activation', *Atmospheric Chemistry and Physics Discussions*, 5(5), pp. 8507–8646. doi: 10.5194/acpd-5-8507-2005.
- Meskhidze, N., Petters, M. D., Tsigaridis, K., Bates, T., O'Dowd, C., Reid, J., Lewis, E. R., Gantt, B., Anguelova, M. D.,
- 10 Bhave, P. V., Bird, J., Callaghan, A. H., Ceburnis, D., Chang, R., Clarke, A., de Leeuw, G., Deane, G., Demott, P. J., Elliot, S., Facchini, M. C., Fairall, C. W., Hawkins, L., Hu, Y., Hudson, J. G., Johnson, M. S., Kaku, K. C., Keene, W. C., Kieber, D. J., Long, M. S., Mårtensson, M., Modini, R. L., Osburn, C. L., Prather, K. A., Pszenny, A., Rinaldi, M., Russell, L. M., Salter, M., Sayer, A. M., Smirnov, A., Suda, S. R., Toth, T. D., Worsnop, D. R., Wozniak, A. and Zorn, S. R. (2013) 'Production mechanisms, number concentration, size distribution, chemical composition, and optical properties of sea spray aerosols',
- 15 *Atmospheric Science Letters*, 14(4), pp. 207–213. doi: 10.1002/asl2.441.
- [Miles, R.E.H., Walker, J.S., Burham, D.R. and Reid, J.P. \(2012\) 'Retrieval of the complex refractive index of aerosol droplets from optical tweezers measurements', \*Physical Chemistry Chemical Physics\*, 14, pp.3037-3037](#)
- Mishra, A., Koren, I. and Ruddich, Y. (2015) 'Effect of aerosol vertical distribution on aerosol-radiation interaction: A theoretical prospect', *Atmospheric Environment*, 2, doi: 10.1016/j.heliyon.2015.e00036.
- 20 Moise, T., Flores, J. M. and Rudich, Y. (2015) 'Optical Properties of Secondary Organic Aerosols and Their Changes by Chemical Processes', *Chemical Reviews*, 115(10), pp. 4400–4439. doi: 10.1021/cr5005259.
- Moosmüller, H., Chakrabarty, R. K., Ehlers, K. M. and Arnott, W. P. (2011) 'Absorption Ångström coefficient, brown carbon, and aerosols: Basic concepts, bulk matter, and spherical particles', *Atmospheric Chemistry and Physics*, 11(3), pp. 1217–1225. doi: 10.5194/acp-11-1217-2011.
- 25 Nakayama, T., Matsumi, Y., Sato, K., Imamura, T., Yamazaki, A. and Uchiyama, A. (2010) 'Laboratory studies on optical properties of secondary organic aerosols generated during the photooxidation of toluene and the ozonolysis of  $\alpha$ -pinene', *Journal of Geophysical Research: Atmospheres*, 115(24), pp. 1–11. doi: 10.1029/2010JD014387.
- Nakayama, T., Sato, K., Matsumi, Y., Imamura, T., Yamazaki, A. and Uchiyama, A. (2013) 'Wavelength and NOx dependent complex refractive index of SOAs generated from the photooxidation of toluene', *Atmospheric Chemistry and Physics*, 13(2),
- 30 pp. 531–545. doi: 10.5194/acp-13-531-2013.
- Nakayama, T., Sato, K., Tsuge, M., Imamura, T. and Matsumi, Y. (2015) 'Complex refractive index of secondary organic aerosol generated from isoprene/NOx photooxidation in the presence and absence of SO<sub>2</sub>', *Journal of Geophysical Research: Atmospheres*, pp. 1–19. doi: 10.1002/2014JD022121. Received.
- Park, S. S. and Yu, J. (2016) 'Chemical and light absorption properties of humic-like substances from biomass burning

- emissions under controlled combustion experiments', *Atmospheric Environment*, 136, pp. 114–122. doi: 10.1016/j.atmosenv.2016.04.022.
- Petty, G. W. (2006) *A First Course in Atmospheric Radiation*. Sundog Publishing, Madison, Wisconsin.
- Pfrang, C., Sebastiani, F., Lucas, C. O. M., King, M. D., Hoare, I. D., Chang, D. and Campbell, R. A. (2014) 'Ozonolysis of methyl oleate monolayers at the air-water interface: oxidation kinetics, reaction products and atmospheric implications.', *Physical chemistry chemical physics*, 16(26), pp. 13220–8. doi: 10.1039/c4cp00775a.
- Pokhrel, R. P., Wagner, N. L., Langridge, J. M., Lack, D. A., Jayarathne, T., Stone, E. A., Stockwell, C. E., Yokelson, R. J. and Murphy, S. M. (2016) 'Parameterization of Single Scattering Albedo (SSA) and Absorption Ångström Exponent (AAE) with EC/OC for Aerosol Emissions from Biomass Burning', *Atmospheric Chemistry and Physics Discussions*, (March), pp. 1–27. doi: 10.5194/acp-2016-184.
- Pöschl, U. (2005) 'Atmospheric aerosols: Composition, transformation, climate and health effects', *Angewandte Chemie - International Edition*, 44(46), pp. 7520–7540. doi: 10.1002/anie.200501122.
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T. and Rosenfeld, D. (2001) 'Aerosols, Climate and the Hydrological Cycle', *Science*, 294(December), pp. 2119–2125. doi: 10.1126/science.1064034.
- Rathod, T., Sahu, S. K., Tiwari, M., Yousaf, A., Bhargare, R. C. and Pandit, G. G. (2016) 'Light Absorbing Properties of Brown Carbon Generated from Pyrolytic Combustion of Household Biofuels', *Aerosol and Air Quality Research*, 1, pp. 108–116. doi: 10.4209/aaqr.2015.11.0639.
- Redemann, J., Turco, R. P., Liou, K. N., Russell, P. B., Bergstrom, R. W., Schmid, B., Livingston, J. M., Hobbs, P. V., Hartley, W. S., Ismail, S., Ferrare, R. A. and Browell, E. V. (2000) 'Retrieving the vertical structure of the effective aerosol complex index of refraction from a combination of aerosol in situ and remote sensing measurements during TARFOX', *Journal of Geophysical Research: Atmospheres*, 105(D8), pp. 9949–9970. doi: 10.1029/1999JD901044.
- Reid, J. S., Koppmann, R., Eck, T. F. and Eleuterio, D. P. (2005) 'A review of biomass burning emissions part II: intensive physical properties of biomass burning particles', *Atmospheric Chemistry and Physics Discussions*, 4(5), pp. 799–825. doi: 10.5194/acpd-4-5135-2004.
- Rkiouak, L., Tang, M. J., Camp, J. C. J., McGregor, J., Watson, I. M., Cox, R. A., Kalberer, M., Ward, A. D. and Pope, F. D. (2014) 'Optical trapping and Raman spectroscopy of solid particles.', *Physical chemistry chemical physics*, 16(23), pp. 11426–34. doi: 10.1039/c4cp00994k.
- Rosenfeld, D., Lohmann, U., Raga, G. B., O'Dowd, C. D., Kulmala, M., Fuzzi, S., Reissell, A. and Andreae, M. O. (2008) 'Flood or drought: how do aerosols affect precipitation?', *Science*, 321(5894), pp. 1309–1313. doi: 10.1126/science.1160606.
- Russell, P. B., Bergstrom, R. W., Shinozuka, Y., Clarke, A. D., DeCarlo, P. F., Jimenez, J. L., Livingston, J. M., Redemann, J., Holben, B., Dubovik, O. and Strawa, A. (2009) 'Absorption Ångström Exponent in AERONET and related data as an indicator of aerosol composition', *Atmospheric Chemistry and Physics Discussions*, 9(5), pp. 21785–21817. doi: 10.5194/acpd-9-21785-2009.
- Sandradewi, J., Prévôt, A. S. H., Weingartner, E., Schmidhauser, R., Gysel, M. and Baltensperger, U. (2008) 'A study of wood

Deleted: Angstrom

Deleted: Angstrom

- burning and traffic aerosols in an Alpine valley using a multi-wavelength Aethalometer', *Atmospheric Environment*, pp. 101–112. doi: 10.1016/j.atmosenv.2007.09.034.
- Schnaiter, M., Gimmler, M., Llamas, I., Linke, C., Jäger, C. and Mutschke, H. (2006) 'Strong spectral dependence of light absorption by organic carbon particles formed by propane combustion', *Atmospheric Chemistry and Physics Discussions*, 6(2), pp. 1841–1866. doi: 10.5194/acpd-6-1841-2006.
- Schnaiter, M., Horvath, H., Möhler, O., Naumann, K. H., Saathoff, H. and Schöck, O. W. (2003) 'UV-VIS-NIR spectral optical properties of soot and soot-containing aerosols', *Journal of Aerosol Science*, 34(10), pp. 1421–1444. doi: 10.1016/S0021-8502(03)00361-6.
- Schweiger, S., (1990) 'Observation of input and output structural resonances in the Raman spectrum of a single spheroidal dielectric microparticle' *Optics Letters*, 15(3), pp. 156-158, doi: 10.1364/OL.15.000156
- Sebastiani, F., Campbell, R. A. and Pfrang, C. (2015) 'Complementarity of neutron reflectometry and ellipsometry for the study of atmospheric reactions at the air–water interface', *RSC Adv.*, 5(129), pp. 107105–107111. doi: 10.1039/C5RA22725A.
- Shen, Z., Zhang, Q., Cao, J., Zhang, L., Lei, Y., Huang, Y., Huang, R. J., Gao, J., Zhao, Z., Zhu, C., Yin, X., Zheng, C., Xu, H. and Liu, S. (2017) 'Optical properties and possible sources of brown carbon in PM<sub>2.5</sub> over Xi'an, China', *Atmospheric Environment*, 150, pp. 322–330. doi: 10.1016/j.atmosenv.2016.11.024.
- Shingler, T., Crosbie, E., Ortega, A., Shiraiwa, M., Zuend, A., Beyersdorf, A., Ziemba, L., Anderson, B., Thornhill, L., Perring, A. E., Schwarz, J. P., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., Hair, J. W., Mikoviny, T., Wisthaler, A. and Sorooshian, A. (2016) 'Airborne characterization of subsaturated aerosol hygroscopicity and dry refractive index from the surface to 6.5 km during the SEAC4RS campaign', *Journal of Geophysical Research*, pp. 4282–4303. doi: 10.1002/2014JD022963.
- Spindler, C., Abo Riziq, A. and Rudich, Y. (2007) 'Retrieval of Aerosol Complex Refractive Index by Combining Cavity Ring Down Aerosol Spectrometer Measurements with Full Size Distribution Information', *Aerosol Science and Technology*, 41(11), pp. 1011–1017. doi: 10.1080/02786820701682087.
- Stamnes, K., C. T. S., Wiscombe, W. J. and Jayaweera, K. (1988) 'Numerically stable algorithm for discrete-ordinate-method radiative transfer in multiple scattering and emitting layered media', *Applied optics*, 27, pp. 2502–2509. doi: 10.1364/AO.27.002502.
- Stocker, T. F., Qin, D. and Plattner, G. K. (2013) 'Climate Change 2013: The Physical Science Basis (Technical Summary)', *Climate Change 2013: The Physical Science Basis*, pp. 33–115. doi: 10.1017/CBO9781107415324.005.
- Tinel, L., Rossignol, S., Bianco, A., Passananti, M., Perrier, S., Wang, X., Brigante, M., Donaldson, D. J. and George, C. (2016) 'Mechanistic Insights on the Photosensitized Chemistry of a Fatty Acid at the Air/Water Interface', *Environmental Science and Technology*, 50(20), pp. 11041–11048. doi: 10.1021/acs.est.6b03165.
- Tomasi, C., Vitale, V., Lupi, A., Di Carmine, C., Campanelli, M., Herber, A., Treffeisen, R., Stone, R. S., Andrews, E., Sharma, S., Radionov, V., von Hoyningen-Huene, W., Stebel, K., Hansen, G. H., Myhre, C. L., Wehrli, C., Aaltonen, V., Lihavainen, H., Virkkula, A., Hillamo, R., Ström, J., Toledano, C., Cachorro, V. E., Ortiz, P., de Frutos, A. M., Blindheim, S., Frioud, M., Gausa, M., Zielinski, T., Petelski, T. and Yamanouchi, T. (2007) 'Aerosols in polar regions: A historical overview based on

- optical depth and in situ observations', *Journal of Geophysical Research Atmospheres*, 112(16). doi: 10.1029/2007JD008432.
- [Thompson, K.C., Rennie, A.R., King, M.D., Hardman, S.J.O., Lucas, C.O.M., Pfrang, C., Hughes, B.R. and Hughes, A.V. \(2010\) 'Reaction of a Phospholipid Monolayer with Gas-Phase Ozone at the Air-Water Interface: Measurement of Surface Excess and Surface Pressure in Real Time', \*Langmuir\*, 26 \(22\), 17295-17303. doi: 10.1021/la1022714](#)
- 5 Trainic, M., Riziq, A. A., Lavi, A., Flores, J. M. and Rudich, Y. (2011) 'and Physics The optical , physical and chemical properties of the products of glyoxal uptake on ammonium sulfate seed aerosols', pp. 9697–9707. doi: 10.5194/acp-11-9697-2011.
- Utry, N., Ajtai, T., Filep, A., Pinter, M., Torok, Z., Bozoki, Z. and Szabo, G. (2014) 'Correlations between absorption ~~Ångström~~ exponent (AAE) of wintertime ambient urban aerosol and its physical and chemical properties', *Atmospheric Environment*. Elsevier Ltd, 91, pp. 52–59. doi: 10.1016/j.atmosenv.2014.03.047.
- 10 Utry, N., Tibor, A., Filep, A., Daniel Pinter, M., Hoffer, A., Bozoki, Z. and Szabo, G. (2013) 'Mass specific optical absorption coefficient of HULIS aerosol measured by a four-wavelength photoacoustic spectrometer at NIR, VIS and UV wavelengths', *Atmospheric Environment*, 69, pp. 321–324. doi: 10.1016/j.atmosenv.2013.01.003.
- Virkkula, A., Koponen, I. K., Teinilä, K., Hillamo, R., Kerminen, V. M. and Kulmala, M. (2006) 'Effective real refractive index of dry aerosols in the Antarctic boundary layer', *Geophysical Research Letters*, 33(6), pp. 10–13. doi: 10.1029/2005GL024602.
- 15 Washenfelder, R. A., Flores, J. M., Brock, C. A., Brown, S. S. and Rudich, Y. (2013) 'Broadband measurements of aerosol extinction in the ultraviolet spectral region', *Atmospheric Measurement Techniques*, 6(4), pp. 861–877. doi: 10.5194/amt-6-861-2013.
- 20 Wex, H., Petters, M. D., Carrico, C. M., Hallbauer, E., Massling, A., McMeeking, G. R., Poulain, L., Wu, Z., Kreidenweis, S. M. and Stratmann, F. (2009) 'Towards closing the gap between hygroscopic growth and activation for secondary organic aerosol: Part 1 – Evidence from measurements', *Atmospheric Chemistry and Physics Discussions*, 9(1), pp. 955–989. doi: 10.5194/acpd-9-955-2009.
- Wild, M. (2009) 'Global dimming and brightening: A review', *Journal of Geophysical Research*, 114. doi: 25 10.1029/2008JD011470.
- Wolff, E. W. (1990) 'Signals of atmospheric pollution in polar snow and ice', *Antarctic Science*, 2(3), pp. 189–205. doi: 10.1017/S095410209000027X.
- Yamasoe, M. A., Artaxo, P., Miguel, A. H. and Allen, A. G. (2000) 'Chemical composition of aerosol particles from direct emissions of vegetation fires in the Amazon Basin: Water-soluble species and trace elements', *Atmospheric Environment*, 34(10), pp. 1641–1653. doi: 10.1016/S1352-2310(99)00329-5.
- 30 Yamasoe, M. A., Kaufman, Y. J., Dubovik, O., Remer, L. A., Holben, B. N. and Artaxo, P. (1998) 'Retrieval of the real part of the refractive index of smoke particles from Sun/sky measurements during SCAR-B', *Journal of Geophysical Research*, 103(98), p. 31893. doi: 10.1029/98JD01211.
- Yu, Y., Ezell, M. J., Zelenyuk, A., Imre, D., Alexander, L., Ortega, J., D'Anna, B., Harmon, C. W., Johnson, S. N. and

Deleted: Angstrom

Finlayson-Pitts, B. J. (2008) 'Photooxidation of alpha-pinene at high relative humidity in the presence of increasing concentrations of NO<sub>x</sub>', *Atmospheric Environment*, 42(20), pp. 5044–5060. doi: 10.1016/j.atmosenv.2008.02.026.

Zhang, X., Lin, Y. H., Surratt, J. D. and Weber, R. J. (2013) 'Sources, composition and absorption Angström exponent of light-absorbing organic components in aerosol extracts from the los angeles basin', *Environmental Science and Technology*, 47(8), pp. 3685–3693. doi: 10.1021/es305047b.

Zhao, W., Dong, M., Chen, W., Gu, X., Hu, C., Gao, X., Huang, W. and Zhang, W. (2013) 'Wavelength-resolved optical extinction measurements of aerosols using broad-band cavity-enhanced absorption spectroscopy over the spectral range of 445–480 nm', *Analytical Chemistry*, 85(4), pp. 2260–2268. doi: 10.1021/ac303174n.

Zhou, S., Gonzalez, L., Leithead, A., Finewax, Z., Thalman, R., Vlasenko, A., Vagle, S., Miller, L. A., Li, S. M., Burekul, S., Furutani, H., Uematsu, M., Volkamer, R. and Abbatt, J. (2014) 'Formation of gas-phase carbonyls from heterogeneous oxidation of polyunsaturated fatty acids at the air-water interface and of the sea surface microlayer', *Atmospheric Chemistry and Physics*, 14(3), pp. 1371–1384. doi: 10.5194/acp-14-1371-2014.






Figure 1: The typical Mie spectra as a function of wavelength obtained for the urban atmospheric aerosol extract collected over the four seasons and the remote atmospheric aerosol extract. From the Mie spectra, the refractive index and radius of the trapped droplet could be determined. Simulated and measured Mie spectra are compared.

Deleted: angstrom



Deleted:

Deleted:









Figure 2: Typical Mie spectra for (a) the wood smoke aerosol extract B and (b) the humic acid aerosol with simulated Mie spectra with and without absorption. The simulated Mie spectra when the measured absorption Ångström exponent,  $\alpha$ , was included or when the absorption Ångström exponent was held at zero are shown for wood smoke aerosol extract and humic acid aerosol. Note the absorption measured in Fig. 2 is required to provide a good fit to the intensity of the data, especially at short wavelength.

Deleted: Ångstrom

Deleted: Ångstrom



Deleted:




Figure 3: The absorbance (base 10) of the aqueous humic acid solution and wood smoke aerosol in isopropanol. The mass density of humic acid in water is  $7.00 \times 10^{-5} \text{ g cm}^{-3}$  and the mass density of the wood smoke extract in isopropanol is  $6.60 \times 10^{-5} \text{ g cm}^{-3}$ . Using mass densities of pure humic acid ( $1.52 \text{ g cm}^{-3}$ ) and wood smoke extract B ( $1.64 \text{ g cm}^{-3}$ ) allows calculation of the imaginary refractive index of the pure components as displayed in the second, lower panel. Parameters for equation (2) displayed in the figure are contained in table 2. The simulated absorbance curves in the upper panel have been displaced upwards for clarity.

**Deleted:** and imaginary refractive index

**Deleted:**

**Deleted:** is shown for wood smoke aerosol extract and aqueous humic acid samples

**Formatted:** Font: +Theme Body (Times New Roman), Font color: Text 1

**Formatted:** Font: +Theme Body (Times New Roman), Font color: Text 1





Figure 4: Refractive index dispersions for urban, remote and wood smoke atmospheric aerosol extracts and humic acid aerosol, compared to refractive index values from literature. A sample of literature studies investigated aerosols from (1) remote locations e.g. b Virkkula (2006), (2) biomass burning e.g. a: Hoffer (2005), c: Guyon (2003) i: Yamasoe (1998), n: Chakrabarty (2010) and (o) Dinar (2008) and (3) organic aerosols e.g. d: Kim and Paulson (2013), e: Nakayama (2013), h: Lambe (2013), j: Spindler (2007), k: Kim (2010), l: Yu (2008), m: Flores (2014), n: Chakrabarty (2010) and p: Trainic (2011).

Deleted: ... [2]



5 **Figure 5: The experimental Mie spectra for the urban spring aerosol extract with the simulated best fit perturbed in three different scenarios to demonstrate the sensitivity of fitting simulated Mie spectra to experimental Mie spectra. Initially the simulated fit (red solid line) is recalculated with a refractive index increased and decreased by 0.015 ( $n \pm 0.015$ ), followed by the simulated fit (red solid line) recalculated with a radius increased and decreased by 6nm ( $r \pm 6\text{nm}$ ). The final set of simulated Mie spectra consider the simulated fit (red solid line) is recalculated with a droplet radius increased and decreased by 12nm, but the refractive index re-optimized to get the best fit to the experimental fit ( $r \pm 12\text{nm}$  and RI re-optimized). A clear demonstration that the quoted uncertainties in radius ( $\pm 6\text{nm}$ ) and refractive index ( $\pm 0.015$ ) are conservative and more than adequate.**

10




Figure 6: The aerosol radiative effect for the change in top of the atmosphere albedo upon core-shell morphology formation in atmospheric aerosols. The shell material consists of the urban atmospheric aerosol extract (top), remote atmospheric aerosol extract (middle) or wood smoke atmospheric aerosol extract (bottom), whilst the core for all calculations was water. The radius of the core was varied 7 times. The left hand side of the graph corresponds to pure aqueous droplets, the right hand side to pure organic droplets.

Deleted:

5

Table 1: The number of aerosols studied, their determined Cauchy coefficients and the particle sizes. Many more particles were studied but were too small to produce Mie spectra that could be fitted with confidence (i.e. the Mie spectra lacked structure). A very limited amount of sample prevented a larger aerosol becoming optically trapped. Where more than one particle was analyzed for a single type of sample, e.g. urban spring, the average and standard deviation of the Cauchy coefficients and real refractive index were reported for the particle studied. The standard deviation does not reflect the uncertainty estimated from the fitting process (which is  $\pm 0.015$  for the real refractive index) but a spread of values obtained for the few particles studied. The range in particle sizes studied is also reported. Note the wood smoke aerosol extract was plentiful and repeated experiments could be performed.

- Deleted: Table 1 describes the
- Deleted: analysed
- Deleted: are
- Deleted: and the
- Deleted: trapped
- Deleted: shown
- Formatted Table

Sample	Aerosols Analysed	A	B (nm <sup>2</sup> )	C (nm <sup>4</sup> )	Real Refractive Index (589 nm)	Radius Range (μm)	
Urban	Spring	3	1.478±0.010	3750±3250	0	1.489	0.49-0.762
	Summer	4	1.465±0.005	4625±1200	0	1.478	0.474-1.252
	Autumn	1	1.505	6000	0	1.522	0.492
	Winter	2	1.495±0.007	4000	0	1.507	0.515
Antarctic Summer 2015	1	1.467	1000	0	1.470	0.503	
Wood Smoke	Extract A	6	1.543	15700± 750	0	1.588± 0.002	0.500-0.723
	Extract B	6	1.541± 0.003	14800± 2900	0	1.584± 0.007	0.475-0.593
Test	Humic Acid (aqueous)	1	1.449	3425	1.25×10 <sup>-8</sup>	1.460	1.307

10

**Table 1: The mass densities and absorption properties of the wood smoke and Humic acid samples used in the work described here.**

Sample	Mass density / $\text{g cm}^{-3}$	Absorption coefficient, $\beta$ , at $\lambda_0 = 460 \text{ nm}$ / $\text{cm}^{-1}$	$k_0$ ( $\lambda_0 = 460 \text{ nm}$ )	$\text{Abs}_0$ ( $\lambda_0 = 460 \text{ nm}$ )	$\alpha$
Pure Wood smoke extract B	1.64	3033	$0.0111 \pm 0.0010$	-	$9.0 \pm 0.1$
Wood smoke extract B in isopropanol (Fig. 3 – top pane)	$6.60 \times 10^{-5}$	0.122	$(4.47 \pm 0.40) \times 10^{-7}$	0.122	$9.0 \pm 0.1$
Aqueous Humic acid (Fig. 3 – top pane)	$7.00 \times 10^{-5}$	1.513	$(4.214 \pm 0.38) \times 10^{-6}$	0.499	$4.2 \pm 0.1$
Pure Humic acid	1.52	25,000	$0.092 \pm 0.046$	-	$4.2 \pm 0.1$
Optically trapped aqueous Humic acid droplet	0.016	273.2	$(1.00 \pm 0.50) \times 10^{-3}$	-	$4.2 \pm 0.1$

By substitution of Eq. (3) (Petty (2006)) into Eq. 2, it was observed that the absorption Ångstrom exponent is related to the imaginary component of the refractive index by  $\alpha-1$  (see Appendix A).

$$Abs = \frac{4k\pi\rho_1 l}{\lambda\rho_2}, \quad (3)$$

In Eq. (3),  $Abs$  is the absorbance measured by the UV-VIS spectrometer,  $\rho_1$  is the mass density of the sample in the solvent,  $\rho_2$  is the mass density of the pure sample only,  $l$  is the path length of the UV-VIS sample cell,  $k$  is the imaginary component of the refractive index and  $\lambda$  is the wavelength. Once the two mass density values were determined for each sample, the mass absorption coefficient and then the imaginary component of the refractive index were calculated. For the aqueous humic acid aerosol, the absorbance of the solvent water over the wavelength range considered was assumed to be negligible.




Figure 5